TACAN is a novel regulator of PKD2

Xiong Liu[1#], Rui Zhang[2#], Mohammad Fatehi[3#], Yifang Wang[2], Wentong Long[3], Rui Tian[2], Xiaoling Deng[2], Ziyi Weng[2], Qinyi Xu[1], Peter E Light[3], Jingfeng Tang[2], Xing-Zhen Chen[1*]

[1]Membrane Protein Disease Research Group, Department of Physiology, Faculty of Medicine and Dentistry, University of Alberta, T6G 2H7, Edmonton, AB, Canada

[2]National "111" Center for Cellular Regulation and Molecular Pharmaceutics, Hubei University of Technology, Wuhan, Hubei 430068, China

[3]Department of Pharmacology, Faculty of Medicine and Dentistry, University of Alberta, T6G 2H7, Edmonton, Alberta, Canada.

[*]Correspondence: Xing-Zhen Chen (xzchen@ualberta.ca)

[#]These authors contributed equally.

Abstract

Autosomal dominant polycystic kidney disease (ADPKD) is caused by mutations in membrane receptor PKD1 or cation channel PKD2. TACAN (also named TMEM120A), recently reported as an ion channel in neuron cells for mechano and pain sensing, is also distributed in diverse non-neuronal tissues such as kidney, heart and intestine, suggesting its involvement in other functions. In this study, we found that TACAN is in complex with PKD2 in native renal cell lines. Using the two-electrode voltage clamp in *Xenopus* oocytes we found that TACAN inhibited the channel activity of PKD2 gain-of-function mutant F604P. The first and last transmembrane domains of TACAN were found to interact with the PKD2 C- and N-terminal portions, respectively. We showed that the TACAN N-terminus acted as a blocking peptide and that TACAN inhibits the PKD2 function through the PKD2/TACAN binding. By patch clamping in mammalian cells, we found that TACAN inhibits both the single channel conductance and open probability of PKD2 and mutant F604P. Further, PKD2 co-expressed with TACAN, but not PKD2 alone, exhibited pressure sensitivity. In summary, this study revealed that TACAN acts as a PKD2 inhibitor and mediates mechano sensitivity of the PKD2/TACAN channel complex.

Keywords: ADPKD, TMEM120A, electrophysiology, *Xenopus* oocyte, mammalian cell, zebrafish, CRISPR/Cas9

Introduction

Autosomal dominant polycystic kidney disease (ADPKD), characterized by accumulation of multiple cysts in both kidneys, is one of the most common human genetic diseases (Harris and Torres, 2009). Extrarenal pathologies associated with ADPKD may include hepatic and pancreatic cysts, cerebral and intracranial aneurysms, and cardiovascular abnormalities (Luciano and Dahl, 2014). Although mutations in PKD1, PKD2 (also called polycystin-1/2) or their dosage alterations account for ADPKD, their biophysical and physiological functions are not well understood (Douguet et al., 2019). PKD1 is a receptor-like membrane protein with 11 transmembrane (TM) segments (S1-S11) and a large extracellular N-terminus while PKD2 is a Ca^{2+}-permeable cation channel belonging to the transient receptor potential (TRP) superfamily of cation channels possessing six TMs (S1-S6) and pore domain S5-loop-S6 (Bergmann et al., 2018).PKD2 forms homotetramers but can also for heterotetramers to fulfill different functions (Cheng et al., 2010). As revealed by cryo-electron microscopy (EM) PKD1/PKD2 formed heterotetramers at 1:3 stoichiometry (Su et al., 2018). Through functional studies, we found that PKD1/PKD2 has higher Ca^{2+} permeability than homomeric PKD2 (Wang et al., 2019) but mechanisms of how PKD1 contributes to the selectivity filter and pore gate in PKD1/PKD2 remains elusive. It was reported that the extracellular N-terminus of PKD1 functions as an activation ligand of the PKD1/PKD2 complex (Ha et al., 2020). Additional reports demonstrated that PKD2 interacts with other ion channels including TRPV4 (Köttgen et al., 2008), TRPC1 (Kobori et al., 2009) and Piezo1 (Peyronnet et al., 2013) to form channel complexes with distinct biophysical properties. Nevertheless, how PKD2 is regulated by interacting partners remains largely unknown and deserves further studies.

TMEM120A (transmembrane protein 120A, also called TACAN) was initially reported as a nuclear envelope transmembrane protein critical for adipocyte differentiation (Batrakou et al., 2015). It was recently shown to form a mechano-sensitive ion channel sensing the pain (Beaulieu-Laroche et al., 2020), but this was challenged by subsequent structural and functional studies (Del Rosario et al., 2021; Ke et al., 2021; Niu et al., 2021; Parpaite et al., 2021; Rong et al., 2021; Xue et al., 2021). Global knockout of TACAN led to embryo death indicating its importance for embryonic development (Beaulieu-Laroche et al., 2020). Wide distribution of TACAN in non-neuronal tissues such as kidney, heart and intestine indicate its other plausible biological functions besides pain sensation (Beaulieu-Laroche et al., 2020). Previous proteomic screen suggested a potential interaction of PKD2 with TACAN (Sharif-Naeini et al., 2009) but their interaction and functional implications has yet to be explored and characterized

In the present study, we examined how TACAN modulates the PKD2 channel function using the two-electrode voltage clamp (TEVC) electrophysiology in *Xenopus* oocytes and single-channel patch clamp electrophysiology in Chinese hamster ovary (CHO) cells. We explored the subcellular localization of endogenous PKD2 and TACAN in different renal cell lines. We also characterized the interaction between TACAN and PKD2 by means of co-immunoprecipitation (co-IP) and immunofluorescence, and explored the relationship between physical association and functional regulation. Further, we examined how TACAN regulates PKD2 deficiency-associated disorders in zebrafish models by clustered regularly interspaced short palindromic repeats/CRISPR-associated protein 9 (CRISPR/Cas9).

Results

Effects of TACAN on the PKD2 channel function

TACAN was indicated as a potential interactor of PKD2 through a proteomic screen using smooth muscle cells (Sharif-Naeini et al., 2009). To determine whether and how TACAN modulates the PKD2 channel function, we utilized *Xenopus* oocyte expression together with the two-electrode voltage clamp (TEVC). The whole-cell channel activity of wild-type (WT) PKD2 in oocytes is hardly detectable in part due to its low expression on the plasma membrane and unknown agonist, Therefore, we instead used the current mediated by PKD2 gain-of-function (GOF) mutant F604P (Arif Pavel et al., 2016) as induced large currents in oocytes, we used currents mediated by mutant F604P as a function readout, similarly as our previous studies (Zheng et al., 2018a; Zheng et al., 2018b). We found that while oocytes expressing human TACAN alone does not exhibit any significant increase in the current compared with water-injected oocytes (Fig. 1A and B). Co-expression of human PKD2 mutant F604P with TACAN is associated with significantly decreased current amplitudes compared with those with F604P expressed alone (Fig. 1A and B). Our data from whole-cell immunofluorescence (Fig. 1C) and biotinylation assays (Fig. 1D) indicated that TACAN did not significantly affect the surface expression of F604P, suggesting TACAN inhibits the F604P function. Interestingly, expression of F604P reduced both the total and surface expression of TACAN (Fig. 1E), but the underlying mechanism remains unknown.

Next, we examined whether TACAN affected the F604P ion selectivity using extracellular solutions containing 100 mM Na^+, Li^+, K^+ or non-permeable N-methyl-d-glucamine (NMDG, negative control). We found that the reversal potential shifts due to solution changes for F604P/TACAN but remain the same as those for F604P alone (Fig. 1F), indicating that TACAN does not significantly affect the cation permeability ratios (with $P_{Na} : P_{Li} : P_K = 1 : 1.65 : 2.15$ for F604P, consistent with previous reports (Shen et al., 2016) and $P_{Na} : P_{Li} : P_K = 1 : 1.66 : 2.15$ for F604P/TACAN). Extracellular Ca^{2+} is known to reduce the PKD2 conductance to monovalent cations (Arif Pavel et al., 2016; Shen et al., 2016). Here we found that Ca^{2+} exhibits a similar inhibitory effect on F604P/TACAN and F604P alone (Fig. 1G and H). Taken together, our data showed that TACAN inhibits the mutant F604P channel activity but does not affect the cation selectivity or inhibition of F604P by extracellular Ca^{2+}.

To further characterize how TACAN inhibits the PKD2 channel function, we performed single-channel patch clamp electrophysiology in CHO cells. Expression of TACAN with mutant F604P or WT PKD2 substantially reduced the single-channel amplitude and open probability while TACAN expressed alone did not induce any specific current under our experimental condition (Fig. 2A and B), consistent with whole-cell data obtained using oocytes. We also noticed that at +80 mV in the absence of TACAN the single-channel amplitude and open probability values for PKD2 are significantly lower than those for mutant F604P, which may account for its much lower whole-cell currents.

We next examined pressure sensitivity of PKD2 in CHO cells with or without co-expression of TACAN. PKD2 in complex with PKD1 or TRPV4 was reported to be involved in mechano sensing (Köttgen et al., 2008; Nauli et al., 2003) but whether PKD2, PKD1, TRPV4 or an unknown binding protein senses mechanical stimuli is not well understood. With PKD2 expressed alone in CHO cells, pressure up to -30 mm Hg had no effect on the single-channel amplitude or open probability (Fig. 2C). In contrast, in the presence of TACAN co-expression,

the open probability value significantly increased with negative pressure while the single-
channel current amplitude was insensitive to the pressure (Fig. 2D). These data showed that
channel complex PKD2/TACAN possesses mechano sensitivity mediated by TACAN, which the
single-channel open probability, but not the unitary current, stimulated by the pressure. Of note,
no channel activity was observed in cells expressing TACAN alone with pressure up to -30 mm
Hg (Fig. S1). Increasing the pressure beyond -30 mm Hg in our setup generated non-specific
leak currents, presumably due to broken cell membrane.

Besides inhibiting the F604P steady-state currents, TACAN induced significant inactivation of
the F604P-mediated currents at membrane depolarization (Fig. 3A). The steady-state current to
the peak current ratio for F604P alone was close to 1 (1.01 ± 0.01 at +100 mV, N = 7, p = 0.73),
i.e., there was no appreciable inactivation, while this ratio decreased to 0.67 ± 0.04 (N = 7, p <
0.001) with TACAN co-expression (Fig. 3B). The current inactivation at depolarizations (+50 to
+100 mV) in the presence of TACAN was voltage dependent, with time constant (τ) values in
the range of 139.23 ± 15.43 and 33.8 ± 4.80 ms (N = 7) obtained from exponential fits (Fig. 3C).
Interestingly, significant inactivation at polarized voltages was also present in PKD2 GOF gate
mutant L677G expressed alone (Fig. 3D). Unlike the F604P mutation, which locks PKD2 in an
activated configuration, the GOF of the L677G mutation is due to direct increases in the pore
hydrophilicity and size, which would infer that the L677G mutant protein may still be in a basal
configuration similar to WT PKD2 (Zheng et al., 2018b). Interestingly, although TACAN
significantly reduced the steady-state currents of L677G (Fig. 3D-F, it did not affect the
inactivating part of the L677G currents, which were associated with a similar range of
inactivation time constant compared with F604P/TACAN (Fig. 3G vs Fig. 3C). These data
together seemed to suggest that inactivation occurs when PKD2 protein is in an inhibited
(F604P+TACAN) or a basal state (L677G, with or without TACAN) but not in an activated state
(F604P alone).

Physical interaction between PKD2 and TACAN
TACAN was shown to be highly expressed in the kidney (Beaulieu-Laroche et al., 2020). We
checked the subcellular localization of PKD2 and TACAN by double immunofluorescence
assays in ciliated epithelia Madin-Darby Canine Kidney (MDCK), inner medullary collecting
duct (IMCD), and Lilly Laboratories Cell-Porcine Kidney 1 (LLC-PK1) cell lines. We found that
the endogenous PKD2 and TACAN both display enhanced staining in the primary cilia in all
three cell lines (Fig. 4A), indicating co-distribution of the two proteins. We also performed co-IP
experiments using these cell lines to assess their interaction and found that PKD2 is present in
the immunoprecipitates obtained with an anti-TMEM120A antibody, but not in those with IgG
(control) (Fig. 4B), demonstrating that PKD2 and TACAN are in the same protein complex in
these native cells.

We next characterized the physical interaction between PKD2 and TACAN interaction in
oocytes to better relate to our functional data. Indeed, TACAN co-immunoprecipitated TACAN
and mutant F604P in oocytes and reversely, mutant F604P was able to precipitate TACAN (Fig.
5A-C), indicating that PKD2 and TACAN are in the same complex in oocytes. We then used
chemical cross-linking to examine the subunit stoichiometry of the PKD2/TACAN complex.
However, the complex cannot be detected under our conditions although we successfully
detected TACAN dimers and different PKD2 oligomers (Fig. S2A). The coiled-coil domain
(amino-acid (aa) G9-L100) in the TACAN N-terminus was proposed to be important for
oligomerization (Batrakou et al., 2015). Indeed, we found aggregation of the N-terminus in our

chemical cross-linking experiments. Further, TACAN lacking the entire N-terminus still formed
the dimer (Fig. S2B), indicating that the TACAN dimeric assembly is independent of its coiled-
coil domain. Furthermore, we generated different fragments of TACAN, which possesses six
TMs (S1-S6), to narrow down the domain(s) mediating physical and functional interaction with
PKD2 (Fig. S3). By co-IP assays we found that TACAN fragments containing S1 (K135-T160)
or S6 (L296-D343), but not the N-terminus (K135X), interacts with PKD2 F604P (Fig. 5D).
While S1-containing TACAN fragments exhibited no functional effect on F604P, interestingly,
S6-containing fragments lacking the N-terminus exhibited significant inhibitory effects (Fig. 5E).
Further, this inhibition was even stronger than the one by full-length TACAN (Fig. 5E),
presumably because these fragments can't form dimers (Fig. S2B), which enhanced their
interaction with F604P.

Reversely, we tried to identify which part(s) in PKD2 is involved in the interaction with
TACAN. We performed co-IP experiments in CHO cells expressing TACAN together with the
PKD2 N-terminus (PKD2-N, M1-K215), C-terminus (PKD2-C, D682-V968), or PKD2 lacking
both the N- and C-termini (PKD2-TM, S209-K688). PKD2-TM but not PKD2-N or PKD2-C
conferred interaction with TACAN (Fig. 6A), which challenges the previous proteomic screen
study indicating that TACAN binds to the PKD2 C-terminus (Sharif-Naeini et al., 2009).

The PKD2 pore domain is formed by S5, S5-S6 loop and S6 (A594-Q693) (Shen et al., 2016).
We next examined whether TACAN participates in the pore formation in the PKD2/TACAN
channel complex. For this purpose, we tested whether TACAN interacts with truncation mutant
A594X (no pore domain) or fragment N580-L700 containing the pore domain (Fig. S3).
Surprisingly, TACAN can interact with either of them (Fig. 6B), indicating that PKD2 also
contains at least two domains interacting with TACAN. Because the TACAN S1 and S6
interacted with PKD2, we continued to narrow down the corresponding binding domains in
PKD2. We found by co-IP that PKD2 A594X interacts with TACAN S6, while PKD2 N580-
L700 interacts with TACAN S1 (Fig. 6C). Since the interaction between S6 was functionally
critical, it highly suggested that instead of acting as the pore formation subunit, TACAN mainly
binds at the peripheral domain (S1-S4) to regulate PKD2 channel function.

We wondered whether the PKD2/TACAN binding is required for inhibition of the PKD2
function by TACAN. For this, we co-expressed the TACAN N-terminus (K135X) in the
presence of F604P and TACAN, and indeed found that K135X represses the PKD2/TACAN
association (Fig. 7A), indicating that K135X acts as an effective blocking peptide, presumably
because it can compete with PKD2 for binding (dimerizing) with TACAN (see Fig. S2). Further,
K135K which had no functional effect on PKD2 (Fig. 5E) significantly represses the inhibition
of the PKD2 function by TACAN (Fig. 7B). Our data together thus demonstrated that inhibition
of the PKD2 function by TACAN is mediated by their physical association.

Regulation of the PKD2 function in zebrafish by TACAN
We next utilized zebrafish models to investigate the regulatory role *in vivo* of TACAN in PKD2.
Sufficient reduction in the PKD2 expression or function in larval zebrafish results in the presence
of tail curvature and pronephric cysts resembling renal cysts in mammals (Zheng et al., 2016).
Here, we employed the clustered regularly interspaced short palindromic repeats/CRISPR-
associated protein 9 (CRISPR/Cas9) technique and successfully knocked down the endogenous
PKD2 in larval fish at 3 days post-fertilization (dpf) through injection of 200 pg PKD2 sgRNA
and 300 pg Cas9 per embryo, which resulted in tail curling (Fig. 8A and B). We reasoned that if

we reduce the injected PKD2 sgRNA and Cas9 amounts there may be no or low occurrence of
tail curling but co-expression of TACAN may significantly aggravate the severity of the disease
if the PKD2 function is inhibited.

Indeed, while injection of 100 pg PKD2 sgRNA and 200 pg Cas9 per embryo was associated
with low percentage of fish with tail curling, co-expression of human TACAN through co-
injection on *in vitro* transcribed mRNA (100 pg/embryo) substantially increased the occurrence
of tail curling (Fig. 8C and D), consistent with the assumption that the PKD2 function is
inhibited by over-expressed TACAN, which was observed in oocytes and CHO cells. Our
Western blot experiments showed that the PKD2 expression is not significantly reduced by
TACAN over-expression (Fig. 8E), which is in agreement with our data from cellular models.
We also observed that TACAN over-expression alone did not significantly increase the
occurrence of tail curling (Fig. 8D), indicating that the functional inhibition of the endogenous
PKD2 by TACAN is in general insufficient to induce the disease. We also examined pronephric
cysts in larval fish and found that TACAN co-expression substantially increases the occurrence
of pronephric cysts while TACAN over-expression or PKD2 knockdown alone exhibits no or
low occurrence of pronephric cyst (Fig. 8F and G). Taken together, our data demonstrated that
TACAN aggravates PKD2-dependent disease severity in larval zebrafish, presumably through
repressing the PKD2 channel function.

Discussion

TACAN is a transmembrane protein that is evolutionarily conserved and was recently reported to be involved in sensing mechanical stimuli (Beaulieu-Laroche et al., 2020). Its intracellular N-terminal coiled-coil domain (aa G9-L100) was reported to be important for oligomerization (Batrakou et al., 2015), consistent with our cross-linking results showing that deletion of the N-terminus significantly reduced its dimerization (Fig. S3A). However, the remaining dimers indicate the presence of another domain important for dimerization. While this other domain involved in homo-dimerization remains unknown, our current study showed that the TACAN TM domains S1 and S6 mediate binding with PKD2. Interestingly, while the PKD2 C-terminal coiled-coil domain was reported to be involved in interaction with PKD1 (Qian et al., 1997; Tsiokas et al., 1997; Yu et al., 2009), heterotetrameric PKD1/PKD2 structure was revealed by cryo-EM in the absence of the PKD2 coiled-coil domain (Su et al., 2018), demonstrating that the domain is not essential for the PKD1/PKD2 complexing. In fact, we here showed that the PKD2 TMs but not its N- or C-terminus interacts with the TACAN S1 and S6.

Besides homotetramerization, PKD2 can heteromerize with PKD1 and TRP channels including TRPC1, -C3, -C4, -C7 and -V4 (Cheng et al., 2010; Grieben et al., 2017; Shen et al., 2016). PKD2 homo-and heterotetrameric channels have distinct biophysical properties as well as distinct subcellular localizations (Cheng et al., 2010). The PKD1/PKD2 heterotetrameric channel defective in ADPKD has a 1:3 stoichiometry (Su et al., 2018). While it is well known that PKD1 promotes the plasma membrane trafficking of PKD2, how it contributes to the channel pore formation is not well understood (Ha et al., 2020; Wang et al., 2019). PKD1/PKD2 located in renal epithelial primary cilia was previously shown to mediate mechano-dependent Ca^{2+} entry(Nauli et al., 2003) but subsequent studies found that, controversially, the Ca^{2+} entry is PKD1-independent (Liu et al., 2018). PKD2/TRPC4 in the MDCK primary cilia was found to be involved in flow sensing but the role of primary cilia and mammalian TRP channels in mechano sensing remains debatable (Delling et al., 2016; Köttgen et al., 2008; Nikolaev et al., 2019). Our current study found that in the PKD2/TACAN channel complex TACAN but not PKD2 confers mechano sensitivity (Fig. 2D). It will be interesting to investigate whether TACAN plays a role in the pathogenesis of ADPKD.

Mammalian TACAN (TMEM120A) and its homologue TMEM120B (70.5% sequence identity) are generated as a result of gene duplication. However, when they were expressed in CHO cells, only TACAN induced mechano-sensitive currents (Beaulieu-Laroche et al., 2020). TACAN, but not TMEM120B, was reported to inhibit the Piezo2 channel function (Del Rosario et al., 2021). It will be of interest to determine whether TMEM120B regulates the PKD2 function in future studies.

Zebrafish has a relatively simple renal system called pronephros that develop within 24 hours post-fertilization (hpf) and represents an established PKD model with easy visualization of organs and tissues with transparent embryos and larvae, and the ability for rapid phenotype analysis. PKD2 knockdown (KD) by injection of morpholino antisense oligonucleotides (MO) in zebrafish embryos results in tail curling and pronephric cysts at 3-5 days post-fertilization (dpf) (Zheng et al., 2016). However, because MO knockdown may have off-target effects, in the present study we knocked down PKD2 by CRISPR/Cas9, which is known to be more specific than MO knockdown. Our study using larval zebrafish demonstrated that TACAN aggravates the severity of PKD2 insufficiency-dependent diseases, presumably through repressing the PKD2

function but not the expression, consistent with its regulatory effect on PKD2 using cellular
models.

Interestingly, it was reported that TMEM33, an activator of PKD2, which forms a complex on
the endoplasmic reticulum (ER) membrane, but not in primary cilia, fails to regulate PKD2-
dependent pronephric renal cystogenesis in zebrafish (Arhatte et al., 2019), indicating that PKD2
located on the ER membrane is not involved in cystogenesis. Thus, the colocalization or
interaction of PKD2 and TACAN in the renal primary cilia may be important for TACAN to
confer its regulatory role in PKD2 and the associated disease phenotypes in zebrafish.

The report that TACAN acts as an ion channel has been challenged by recent structural and
functional studies (Beaulieu-Laroche et al., 2020; Del Rosario et al., 2021; Ke et al., 2021; Niu et
al., 2021; Parpaite et al., 2021; Rong et al., 2021; Xue et al., 2021). However, the concept that
TACAN is an ion channel is supported by a recent report showing that a point mutation at a
candidate pore gate residue significantly increases the channel activity (Chen et al., 2021) and
that, by molecular dynamic simulations, each TACAN monomer contains an ion conducting
pathway but multiple subunits might be required for TACAN to form a functional ion channel.
Also, TACAN was shown to be involved in regulation of ion transport based on its involvement
in regulation of the expression of different proteins including Ca^{2+} ATPase, ATP2a1 and
TMEM150C, a general regulator of mechano-sensitive channels such as Piezo1, Piezo2 and the
potassium channel TREK-1 (Anderson et al., 2018). Structural studies found that TACAN
structure resembles that of the long-chain fatty acid elongase 7 (Ke et al., 2021; Niu et al., 2021;
Rong et al., 2021; Xue et al., 2021), suggesting the possibility that TACAN may act as an
enzyme. In addition to our current finding, TACAN was previously reported to be an inhibitor of
Piezo2, but not Piezo1 or TREK-1, but with an unclear mechanism of inhibition (Del Rosario et
al., 2021).

In summary, our present study found that TACAN inhibits the PKD2 function through the
physical PKD2/TACAN association. It is unlikely that TACAN participates in the pore
formation in the PKD2/TACAN channel complex since TACAN-S6 that doesn't seem to be
sufficient for contributing to pore formation still exhibited an inhibitory effect. As a novel
regulator of PKD2, TACAN inhibits PKD2 *in vitro* and *in vivo*. Blocking peptides such as
K135X may eventually lead to specific peptide molecules with therapeutic potential for ADPKD.

Methods

Plasmids, reagents and antibodies

HA-tagged human PKD2 (HA-PKD2) in vector pGEMHE for *Xenopus* oocyte expression and PKD2 mutant F604P in vector pcDNA3.1 for mammalian cell expression were obtained from Dr. Yong Yu (St. John's University, NY). Human HA-TACAN in a modified pCMV vector for mammalian cell expression, a kind gift from Dr. Reza Sharif-Naeini (McGill University, QC, Canada), was subcloned into vector PGEMHE for oocyte expression, with a 5' Flag tag. Human PKD2, PKD2-N, PKD2-TM and PKD2-C in vector pEGFP for oocyte expression were constructed previously (Wang et al., 2012). All mutations were carried out using Q5 High-Fidelity 2X Master Mix from New England Biolabs (NEB, Ottawa, ON, Canada) and verified by sequencing. Antibodies against β-actin and Arl13b were from Santa Cruz Biotechnology (Santa Cruz, CA) and those against Flag, HA, TMEM120A and PKD2 were from Proteintech Group (Rosemont, IL). Secondary antibodies were purchased from GE Healthcare (Waukesha, WI). DTT was purchased from Thermo Fisher Scientific (Ottawa, ON, Canada) and disuccinimidyl tartrate (DST) from CovaChem (Loves Park, IL).

Xenopus oocyte expression

Capped RNAs encoding human PKD2, human TACAN or their mutants were synthesized by *in vitro* transcription using the T7 mMESSAGE mMACHINE kit (Invitrogen, Waltham, MA) and injected into oocytes (25 ng each), as described (Cai et al., 2020). Control oocytes were injected with the same amount of water. Oocytes were incubated at 18℃ for 2-3 days before measurements. The present study was approved by the Ethical Committee for Animal Experiments of the University of Alberta and performed in accordance with the Guidelines for Research with Experimental Animals of the University of Alberta and the Guide for the Care and Use of Laboratory Animals (NIH Guide) revised in 1996.

Culture and transfection of mammalian cells

CHO cells were cultured in Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12 (DMEM/F12) supplemented with L-glutamine, penicillin-streptomycin and 10% fetal bovine serum (FBS). MDCK, IMCD and LLC-PK1 cells were cultured in DMEM supplemented with L-glutamine, penicillin-streptomycin and 10% FBS. All cultured cells were kept at 37°C with 5% CO_2. Transfection of cDNAs was performed using Lipofectamine 3000 (Invitrogen) according to the manufacturer's protocol.

Two-electrode voltage clamp

The two-electrode voltage clamp electrophysiology experiments in *Xenopus* oocytes were performed as we described previously (Zheng et al., 2018a) Briefly, an electrode made of a capillary glass pipette (Warner Instruments, Hamden, CT) was filled with 3 M KCl and impaled an oocyte to form a tip resistance of 0.3-2 MΩ. The Geneclamp 500B amplifier and Digidata 1322A AD/DA converter (Molecular Devices, Union City, CA) were used to record the whole-cell currents and membrane potentials. The pClamp 9 software (Axon Instruments, Union City, CA) was used to acquire and analyze the data. Both signals were digitized at 200 μs/sample and filtered at 2 kHz through a Bessel filter. Data were plotted using SigmaPlot 13 (Systat Software, San Jose, CA) or GraphPad Prism 8 (GraphPad Software, San Diego, CA).

Patch clamp

Patch-clamp recordings were carried out as we described previously (Fatehi et al., 2017), under
voltage clamp at room temperature (~22°C). The recording pipette and chamber were coupled by
Ag/AgCl electrodes to an Axopatch 200B patch-clamp amplifier and Digidata 1200A BNC data-
acquisition system, controlled by pCLAMP 10 software (Axon Instruments). Both bath and
pipette solutions were composed of (in mM) 140 NaCl, 5 KCl, 2 $CaCl_2$, 2 $MgCl_2$, 10 HEPES, 10
glucose and pH 7.4. The seal resistance was no less than 5 GΩ for all cell-attached recordings.
The recordings were low-pass filtered at 1 kHz following acquisition. Linear stepwise negative
pressures (by suction) of various magnitudes were applied to the interior of the glass pipette
using a high-speed pressure clamp system (HSPC-1, ALA Scientific, Farmingdale, NY).

Western blot and surface protein biotinylation
Xenopus oocytes after 3-times washing with ice-cold PBS solution were incubated with 0.5
mg/ml sulfo-NHS-SS-Biotin (Pierce, Rockford, IL) for 30 min at room temperature. Non-reacted
biotin was quenched by 1 M NH_4Cl. Oocytes were then washed with ice-cold PBS solution and
harvested in ice-cold CelLytic M lysis buffer (Sigma-Aldrich, St. Louis, MO) supplemented with
proteinase inhibitor cocktail (Thermo Fisher Scientific). Upon addition of 100 μl streptavidin
(Pierce) lysates were incubated at 4℃ overnight with gentle shaking. The surface protein bound
to streptavidin was resuspended in SDS loading buffer and subjected to SDS-PAGE.

Chemical cross-linking
Chemical cross-linking assays were performed as previously described (Yu et al., 2009). Oocytes
expressing desired proteins were harvested in ice-cold CelLytic M lysis buffer (Sigma-Aldrich)
supplemented with proteinase inhibitor cocktail (Thermo Fisher Scientific). Fresh crosslinker
stock solutions were prepared by dissolving DST (CovaChem) into dimethylsulfoxide (Sigma-
Aldrich). Cross-linking reactions were carried out by diluting the stock solution into the cell
lysate samples, followed by incubation on ice for 4 h. SDS loading buffer was used to stop the
reactions. The samples were incubated at 37 °C for 30 min and subjected to SDS-PAGE.

Immunofluorescence
Whole-mount immunofluorescence assays using *Xenopus* oocytes were performed as described
(Zheng et al., 2018a). Briefly, oocytes were washed in PBS, fixed in 4% paraformaldehyde for
15 min, washed three times in PBS plus 50 mM NH_4Cl, and then permeabilized with 0.1%
Triton X-100 for 4 min. Oocytes were then blocked in PBS plus 3% skim milk for 30 min and
then incubated overnight with indicated primary antibodies, followed by incubation with
secondary Alexa-488-conjugated donkey anti-rabbit or Cy3-conjugated goat anti-mouse antibody
(Jackson ImmunoResearch Laboratories, West Grove, PA) for 30 min. Oocytes were then
mounted in Vectashield (Vector Labs, Burlington, ON, Canada) and examined on an AIVI
spinning disc confocal microscopy (Cell Imaging Facility, Faculty of Medicine and Dentistry,
University of Alberta).

Co- IP
Co-IP experiments were performed as we previously described (Zheng et al., 2018a). Briefly, a
group of 20-30 oocytes washed with PBS were solubilized in ice-cold CelLytic-M lysis buffer
(Sigma-Aldrich) supplemented with proteinase inhibitor cocktail. Supernatants were collected
after centrifugation at 13,200 rpm for 15 min and precleaned for 1 h with 50% protein G-
Sepharose (GE Healthcare), followed by incubation with an indicated antibody at 4℃ for 4 h.
Upon addition of 100 μl of 50% protein G-Sepharose, the mixture was incubated at 4℃

overnight with gentle shaking. The immune complexes conjugated to protein G-Sepharose were
washed three times with cold PBS solution containing 1% Nonidet P-40 and eluted by SDS
loading buffer. Precipitated proteins were subjected to Western blot analysis.

Zebrafish experiments

Zebrafish experiments were modified from one described previously (Zheng et al., 2018b).
Briefly, embryos of WT zebrafish AB line were grown at 28.5 ⁰C in water supplemented with 60
µg/ml Instant Ocean Sea salts. Single guide RNAs (sgRNAs: AGAACACCGCTCTAGTGCCG)
targeting the first coding exon of zebrafish PKD2 was designed and synthesized. The Cas9
protein was purchased from NEB. Mixture of sgRNA (100 pg) and Cas9 (200 pg) was injected
into each fertilized embryo at 1-2 hpf for PKD2 knock down. The effect of injected
CRISPR/Cas9 was confirmed by sequencing. Human TACAN mRNA was injected into
fertilized embryos at 1-2 hpf at 100 pg each. This study has been approved by the Hubei
University of Technology animal welfare regulations and maintained according to standard
protocols (http://ZFIN.org).
Histological analysis was performed as previously reported (Wang et al., 2020). Briefly, embryos
at 3 dpf were anesthetized in tricaine solution and fixed in 4% PFA at 4 ⁰C. After 3-times wash
(20 min each) in cold PBS, embryos were decalcified in EDTA solution (684 mM EDTA, pH
8.0) and then washed (2 × 20 min) with DEPC solution and dehydrated through graded alcohol.
Before paraffin processing, embryos were embedded in 1% agarose (Sigma-Aldrich) in TAE
buffer (Thermo Fisher Scientific). Subsequently, samples were sectioned transversely at 5 µm
thickness using a HM325 manual rotary microtome (Thermo Fisher Scientific). H&E staining
was processed with Varistain TM Gemini ES Automated Slide Stainer (Thermo Fisher
Scientific).

Statistical analysis

407 All statistical data in this study were represented as mean ± SEM (standard error of the mean)
408 from N measurements. Student's t-test was used for two groups comparisons and one-way
409 ANOVA for multiple groups comparisons. *, ** and *** indicate $p < 0.05$, 0.01 and 0.001,
410 respectively; ns indicates statistically not significant.

Author contributions

Conceptualization, X.L., and X.-Z.C.; Investigation, X.L., R.Z., M.F., Y.W., W.L., R.T., X.D., Z.W., Q.X.; Supervision, P.R.L., J.T., and X.-Z.C. Writing, X.L., and X.-Z.C.

Acknowledgments

We would like to thank Dr. Reza Sharif-Naeini for sharing TACAN related plasmids. This work was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), the Kidney Foundation of Canada (to X.Z.C.) and National Natural Science Foundation of China (grant #32070726, to J.T.). X.L. was a recipient of the Alberta Graduate Excellence Scholarship.

Declaration of Interests

The authors declare no competing interests.

Figure legends

Figure 1. Functional regulation of PKD2 by TACAN in *Xenopus* oocytes. **A.** Representative current traces obtained using a voltage jump protocol in oocytes expressing TACAN, PKD2 F604P (F604P for short) or F604P + TACAN, in the presence of the divalent-free Na-containing solution (in mM): 100 NaCl, 2 KCl, 10 HEPES and pH 7.5. Data from water-injected oocytes served as negative control (Ctrl). **B.** Statistical bar graphs of currents recoded at +80 mV obtained under the same experimental conditions as in **A**. Data are presented as mean ± SEM. * p < 0.05; ** p < 0.01. **C.** Representative IF images showing oocyte surface expression of TACAN or F604P. Scale bar, 50 mm. **D.** Representative Western blot data of the biotinylated (surface) and total protein of F604P and TACAN. **E.** Averaged data obtained as those from **D** (n = 3). Data are presented as mean ± SEM. ns, not significant; *** p < 0.001. **F.** Upper panel: representative I-V curves for F604P or F604P + TACAN when 100 mM of an indicated ion was used in the divalent ion-free bath solution. Lower panel: averaged reversal potentials. Data are presented as mean ± SEM. ns, not significant. **G** and **H.** Left panel: representative I-V curves for PKD2 F604P (**G**) and F604P + TACAN (**H**) in bath solution with or without 1 mM Ca^{2+}. Right panel: effect of Ca^{2+} on the inward (-80 mV) and outward (+80 mV) currents for F604P (**G**) and F604P + TACAN (**H**). Data are presented as mean ± SEM. * p < 0.05; ** p < 0.01; *** p < 0.001.

Figure 2. Effects of TACAN and pressure on PKD2 single-channel parameters in CHO cells. A. Representative single-channel recordings (at +80 mV) of CHO cells with indicated transfections. The same solution (in mM: 140 NaCl, 5 KCl, 2 $CaCl_2$, 2 $MgCl_2$, 10 HEPES, 10 glucose, and pH 7.4) was used in the pipette and bath. **B.** Quantified unitary current amplitudes and open probabilities obtained from **A**. Data are presented as mean ± SEM. ** p < 0.01; *** p < 0.001. **C** and **D.** Left panels: representative single-channel recordings (at +80 mV) in CHO cells transfected with PKD2 (**C**) or PKD2 + TACAN (**D**) showing the effect of negative pressures. Right panels: effect of the pressure on the unitary currents and normalized open probabilities of PKD2 (**C**) or PKD2 + TACAN (**D**). Data are presented as mean ± SEM. ns, not significant; ** p < 0.01.

Figure 3. Effect of TACAN on inactivation of PKD2-mediated currents in oocytes. A. Representative traces from oocytes expressing PKD2 F604P or F604P + TACAN. Traces obtained from +50 mV to +100 mV were marked blue or green, as indicated. For clarity, those at +80 mV were compared. **B.** Ratios of the steady currents (I_{steady}) to the peak currents (I_{peak}) for F604P and F604P + TACAN obtained as in **A**. **C.** Time constant (τ) of current decay for F604P + TACAN obtained through an exponential fit (N = 7). **D.** Representative traces from oocytes expressing PKD2 L677G or L677G + TACAN. Traces obtained from +50 mV to +100 mV are marked blue or green, as indicated. **E.** Left panel: Representative I-V curves from oocytes expressing L677G or L677G + TACAN. Right panel: averaged currents at +80 mV obtained from oocytes expressing L677G or L677G + TACAN. Data are presented as mean ± SEM. ** p< 0.01. **F.** Representative Western blot data of the surface (biotinylated) and total proteins of L677G and TACAN. **G.** Time constants (τ) of current decay for L677G and L677G + TACAN (N = 5).

Figure 4. Colocalization and association of the endogenous PKD2 and TACAN in renal cell lines. A. Subcellular localization of the endogenous PKD2 and TACAN in MDCK, IMCD and LLC-PK1 cells by means of immunofluorescence. Arl13b served as a primary cilia marker. Scale

bar, 5 μm. **B.** Representative co-IP data showing association between the endogenous PKD2 and TACAN in MDCK, IMCD and LLC-PK1 cells.

Figure 5. Domains of TACAN mediating association with PKD2. A and B. Representative co-IP data showing the interaction between PKD2 mutant HA-F604P and Flag-TACAN using Flag antibody (**A**) or PKD2 antibody (**B**) for precipitation. **C.** Representative co-IP data showing interaction between WT HA-PKD2 and Flag-TACAN using Flag antibody for precipitation. **D.** Representative co-IP data showing the interaction between mutant HA-F604P and an indicated TACAN truncation mutant using Flag antibody for precipitation. **E.** Representative I-V curves and averaged currents at +80 mV in oocytes expressing F604P with or without a TACAN truncation mutant. Data are presented as mean ± SEM. * p < 0.05; ** p < 0.01; *** p < 0.001.

Figure 6. Domains in PKD2 mediating association with TACAN. A. Left panels: representative co-IP data showing the interaction between GFP-PKD2 and HA-TACAN using HA antibody for precipitation. Non-transfected and GFP-transfected CHO cell lysates served as negative controls. Right panel: binding strength averaged from three independent experiments. **B.** Representative co-IP data showing the interaction between Flag-TACAN and a PKD2 truncation mutant (A594X or N580-L700) using Flag antibody for precipitation. PKD2 expressed in oocytes with or without co-expressed Flag-TACAN served as positive and negative controls, respectively. **C.** Representative co-IP data showing the interaction between PKD2 mutant A594X and TACAN mutant L296-D343, and between PKD2 mutant N580-L700 and TACAN mutant K160X using Flag antibody for precipitation. PKD2 mutants A594X and N580-L700 alone served as negative controls.

Figure 7. Effect of the TACAN N-terminus on TACAN's inhibition of the PKD2 function. A. Left panel: representative I-V curves from oocytes expressing F604P, F604P + TACAN, or F604P + TACAN + TACAN N-terminus (K135X). Right panel: averaged currents at +80 mV. Data are presented as mean ± SEM. ns, not significant, *** p < 0.001. **B.** Representative co-IP data showing the effect of K135X on the association between PKD2 and TACAN.

Figure 8. Effect of TACAN on the tail curling and pronephric cystogenesis of larval zebrafish. A. Left panel: schematic presentation of the PKD2 protein/gene domain architecture and CRISPR target sites. The gene loci are shown with exons (gray boxes) and introns (solid lines). The position of the CRISPR target sites in exon 1 in the PKD2 gene and the corresponding PKD2 protein (black box) are indicated by the dash lines, and the predicted remaining (truncated) PKD2 protein (PKD2 CRISPR) is indicated. Right panel: sequencing of PKD2 in the Ctrl and PKD2 CRISPR fish at 3 dpf. Red lines indicate CRISPR target sites. **B.** Upper panel: PKD2 protein detected by Western blot under the Ctrl (water injection) and CRISPR-directed PKD2 knock down (KD) conditions (PKD2 sgRNA 200 pg + Cas9 300 pg) at 3 dpf. Lower panels: representative pictures of embryos at 3 dpf. **C.** Representative pictures showing tails of 3 dpf zebrafish under the Ctrl, TACAN overexpression (OE) (by injection of mRNA at100 pg each), or PKD2 KD condition. **D.** Average percentages of fish displaying curled tail under different conditions. Data were from three independent experiments with the indicated total numbers of embryos. ns, not significant, ***p < 0.001. **E.** Representative Western blot data showing the expression of PKD2 and TACAN. **F.** Representative pictures of a 3-dpf embryo with PKD2 KD plus TACAN OE, showing pronephric cyst formation (red box and arrow). Ctrl fish was injected with water only. Cyst formation was also indicated by a histologic section showing

dilated pronephric tubules (asterisks). G, glomerulus; Pt, pronephric tubule. **G.** Averaged
percentages of embryos exhibiting pronephric cysts under the indicated conditions. Data are
presented as mean $\pm$ SEM. ***$p < 0.001$.

References

Anderson, E.O., Schneider, E.R., Matson, J.D., Gracheva, E.O., and Bagriantsev, S.N. (2018). TMEM150C/Tentonin3 Is a Regulator of Mechano-gated Ion Channels. Cell reports *23*, 701-708. 10.1016/j.celrep.2018.03.094.

Arhatte, M., Gunaratne, G.S., El Boustany, C., Kuo, I.Y., Moro, C., Duprat, F., Plaisant, M., Duval, H., Li, D., Picard, N., et al. (2019). TMEM33 regulates intracellular calcium homeostasis in renal tubular epithelial cells. Nature communications *10*, 2024. 10.1038/s41467-019-10045-y.

Arif Pavel, M., Lv, C., Ng, C., Yang, L., Kashyap, P., Lam, C., Valentino, V., Fung, H.Y., Campbell, T., Møller, S.G., et al. (2016). Function and regulation of TRPP2 ion channel revealed by a gain-of-function mutant. Proceedings of the National Academy of Sciences of the United States of America *113*, E2363-2372. 10.1073/pnas.1517066113.

Batrakou, D.G., de Las Heras, J.I., Czapiewski, R., Mouras, R., and Schirmer, E.C. (2015). TMEM120A and B: Nuclear Envelope Transmembrane Proteins Important for Adipocyte Differentiation. PloS one *10*, e0127712. 10.1371/journal.pone.0127712.

Beaulieu-Laroche, L., Christin, M., Donoghue, A., Agosti, F., Yousefpour, N., Petitjean, H., Davidova, A., Stanton, C., Khan, U., Dietz, C., et al. (2020). TACAN Is an Ion Channel Involved in Sensing Mechanical Pain. Cell *180*, 956-967.e917. 10.1016/j.cell.2020.01.033.

Bergmann, C., Guay-Woodford, L.M., Harris, P.C., Horie, S., Peters, D.J.M., and Torres, V.E. (2018). Polycystic kidney disease. Nature reviews. Disease primers *4*, 50. 10.1038/s41572-018-0047-y.

Cai, R., Liu, X., Zhang, R., Hofmann, L., Zheng, W., Amin, M.R., Wang, L., Hu, Q., Peng, J.B., Michalak, M., et al. (2020). Autoinhibition of TRPV6 Channel and Regulation by PIP2. iScience *23*, 101444. 10.1016/j.isci.2020.101444.

Chen, X., Wang, Y., Li, Y., Lu, X., Chen, J., Li, M., Wen, T., Liu, N., Chang, S., Zhang, X., et al. (2021). Cryo-EM structure of the human TACAN channel in a closed state. bioRxiv, 2021.2008.2023.457436. 10.1101/2021.08.23.457436.

Cheng, W., Sun, C., and Zheng, J. (2010). Heteromerization of TRP channel subunits: extending functional diversity. Protein & cell *1*, 802-810. 10.1007/s13238-010-0108-9.

Del Rosario, J.S., Gabrielle, M., Yudin, Y., and Rohacs, T. (2021). TMEM120A/TACAN inhibits mechanically activated Piezo2 channels. bioRxiv, 2021.2006.2030.450616. 10.1101/2021.06.30.450616.

Delling, M., Indzhykulian, A.A., Liu, X., Li, Y., Xie, T., Corey, D.P., and Clapham, D.E. (2016). Primary cilia are not calcium-responsive mechanosensors. Nature *531*, 656-660. 10.1038/nature17426.

Douguet, D., Patel, A., and Honoré, E. (2019). Structure and function of polycystins: insights into polycystic kidney disease. Nature reviews. Nephrology *15*, 412-422. 10.1038/s41581-019-0143-6.

Fatehi, M., Carter, C.C., Youssef, N., and Light, P.E. (2017). The mechano-sensitivity of cardiac ATP-sensitive potassium channels is mediated by intrinsic MgATPase activity. Journal of molecular and cellular cardiology *108*, 34-41. 10.1016/j.yjmcc.2017.05.004.

Grieben, M., Pike, A.C., Shintre, C.A., Venturi, E., El-Ajouz, S., Tessitore, A., Shrestha, L., Mukhopadhyay, S., Mahajan, P., Chalk, R., et al. (2017). Structure of the polycystic kidney disease TRP channel Polycystin-2 (PC2). Nature structural & molecular biology *24*, 114-122. 10.1038/nsmb.3343.

Ha, K., Nobuhara, M., Wang, Q., Walker, R.V., Qian, F., Schartner, C., Cao, E., and Delling, M. (2020). The heteromeric PC-1/PC-2 polycystin complex is activated by the PC-1 N-terminus. eLife *9*. 10.7554/eLife.60684.

Harris, P.C., and Torres, V.E. (2009). Polycystic kidney disease. Annual review of medicine *60*, 321-337. 10.1146/annurev.med.60.101707.125712.

Ke, M., Yu, Y., Zhao, C., Lai, S., Su, Q., Yuan, W., Yang, L., Deng, D., Wu, K., Zeng, W., et al. (2021). Cryo-EM structures of human TMEM120A and TMEM120B. Cell discovery *7*, 77. 10.1038/s41421-021-00319-5.

Kobori, T., Smith, G.D., Sandford, R., and Edwardson, J.M. (2009). The transient receptor potential channels TRPP2 and TRPC1 form a heterotetramer with a 2:2 stoichiometry and an alternating subunit arrangement. The Journal of biological chemistry *284*, 35507-35513. 10.1074/jbc.M109.060228.

Köttgen, M., Buchholz, B., Garcia-Gonzalez, M.A., Kotsis, F., Fu, X., Doerken, M., Boehlke, C., Steffl, D., Tauber, R., Wegierski, T., et al. (2008). TRPP2 and TRPV4 form a polymodal sensory channel complex. The Journal of cell biology *182*, 437-447. 10.1083/jcb.200805124.

Liu, X., Vien, T., Duan, J., Sheu, S.H., DeCaen, P.G., and Clapham, D.E. (2018). Polycystin-2 is an essential ion channel subunit in the primary cilium of the renal collecting duct epithelium. eLife *7*. 10.7554/eLife.33183.

Luciano, R.L., and Dahl, N.K. (2014). Extra-renal manifestations of autosomal dominant polycystic kidney disease (ADPKD): considerations for routine screening and management. Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association *29*, 247-254. 10.1093/ndt/gft437.

Nauli, S.M., Alenghat, F.J., Luo, Y., Williams, E., Vassilev, P., Li, X., Elia, A.E., Lu, W., Brown, E.M., Quinn, S.J., et al. (2003). Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells. Nature genetics *33*, 129-137. 10.1038/ng1076.

Nikolaev, Y.A., Cox, C.D., Ridone, P., Rohde, P.R., Cordero-Morales, J.F., Vasquez, V., Laver, D.R., and Martinac, B. (2019). Mammalian TRP ion channels are insensitive to membrane stretch. Journal of cell science *132*. 10.1242/jcs.238360.

Niu, Y., Tao, X., Vaisey, G., Olinares, P.D.B., Alwaseem, H., Chait, B.T., and MacKinnon, R. (2021). Analysis of the mechanosensor channel functionality of TACAN. eLife *10*. 10.7554/eLife.71188.

Parpaite, T., Brosse, L., Séjourné, N., Laur, A., Mechioukhi, Y., Delmas, P., and Coste, B. (2021). Patch-seq of mouse DRG neurons reveals candidate genes for specific mechanosensory functions. Cell reports *37*, 109914. 10.1016/j.celrep.2021.109914.

Peyronnet, R., Martins, J.R., Duprat, F., Demolombe, S., Arhatte, M., Jodar, M., Tauc, M., Duranton, C., Paulais, M., Teulon, J., et al. (2013). Piezo1-dependent stretch-activated channels are inhibited by Polycystin-2 in renal tubular epithelial cells. EMBO reports *14*, 1143-1148. 10.1038/embor.2013.170.

Qian, F., Germino, F.J., Cai, Y., Zhang, X., Somlo, S., and Germino, G.G. (1997). PKD1 interacts with PKD2 through a probable coiled-coil domain. Nature genetics *16*, 179-183. 10.1038/ng0697-179.

Rong, Y., Jiang, J., Gao, Y., Guo, J., Song, D., Liu, W., Zhang, M., Zhao, Y., Xiao, B., and Liu, Z. (2021). TMEM120A contains a specific coenzyme A-binding site and might not mediate poking- or stretch-induced channel activities in cells. eLife *10*. 10.7554/eLife.71474.

Sharif-Naeini, R., Folgering, J.H., Bichet, D., Duprat, F., Lauritzen, I., Arhatte, M., Jodar, M.,
Dedman, A., Chatelain, F.C., Schulte, U., et al. (2009). Polycystin-1 and -2 dosage regulates
pressure sensing. Cell *139*, 587-596. 10.1016/j.cell.2009.08.045.
Shen, P.S., Yang, X., DeCaen, P.G., Liu, X., Bulkley, D., Clapham, D.E., and Cao, E. (2016).
The Structure of the Polycystic Kidney Disease Channel PKD2 in Lipid Nanodiscs. Cell *167*,
763-773.e711. 10.1016/j.cell.2016.09.048.
Su, Q., Hu, F., Ge, X., Lei, J., Yu, S., Wang, T., Zhou, Q., Mei, C., and Shi, Y. (2018). Structure
of the human PKD1-PKD2 complex. Science (New York, N.Y.) *361*. 10.1126/science.aat9819.
Tsiokas, L., Kim, E., Arnould, T., Sukhatme, V.P., and Walz, G. (1997). Homo- and
heterodimeric interactions between the gene products of PKD1 and PKD2. Proceedings of the
National Academy of Sciences of the United States of America *94*, 6965-6970.
10.1073/pnas.94.13.6965.
Wang, Q., Dai, X.Q., Li, Q., Wang, Z., Cantero Mdel, R., Li, S., Shen, J., Tu, J.C., Cantiello, H.,
and Chen, X.Z. (2012). Structural interaction and functional regulation of polycystin-2 by
filamin. PloS one *7*, e40448. 10.1371/journal.pone.0040448.
Wang, X., Giusti, A., Ny, A., and de Witte, P.A. (2020). Nephrotoxic Effects in Zebrafish after
Prolonged Exposure to Aristolochic Acid. Toxins *12*. 10.3390/toxins12040217.
Wang, Z., Ng, C., Liu, X., Wang, Y., Li, B., Kashyap, P., Chaudhry, H.A., Castro, A., Kalontar,
E.M., Ilyayev, L., et al. (2019). The ion channel function of polycystin-1 in the polycystin-
1/polycystin-2 complex. EMBO reports *20*, e48336. 10.15252/embr.201948336.
Xue, J., Han, Y., Baniasadi, H., Zeng, W., Pei, J., Grishin, N.V., Wang, J., Tu, B.P., and Jiang,
Y. (2021). TMEM120A is a coenzyme A-binding membrane protein with structural similarities
to ELOVL fatty acid elongase. eLife *10*. 10.7554/eLife.71220.
Yu, Y., Ulbrich, M.H., Li, M.H., Buraei, Z., Chen, X.Z., Ong, A.C., Tong, L., Isacoff, E.Y., and
Yang, J. (2009). Structural and molecular basis of the assembly of the TRPP2/PKD1 complex.
Proceedings of the National Academy of Sciences of the United States of America *106*, 11558-
11563. 10.1073/pnas.0903684106.
Zheng, W., Cai, R., Hofmann, L., Nesin, V., Hu, Q., Long, W., Fatehi, M., Liu, X., Hussein, S.,
Kong, T., et al. (2018a). Direct Binding between Pre-S1 and TRP-like Domains in TRPP
Channels Mediates Gating and Functional Regulation by PIP2. Cell reports *22*, 1560-1573.
10.1016/j.celrep.2018.01.042.
Zheng, W., Shen, F., Hu, R., Roy, B., Yang, J., Wang, Q., Zhang, F., King, J.C., Sergi, C., Liu,
S.M., et al. (2016). Far Upstream Element-Binding Protein 1 Binds the 3' Untranslated Region of
PKD2 and Suppresses Its Translation. Journal of the American Society of Nephrology : JASN
27, 2645-2657. 10.1681/ASN.2015070836.
Zheng, W., Yang, X., Hu, R., Cai, R., Hofmann, L., Wang, Z., Hu, Q., Liu, X., Bulkley, D., Yu,
Y., et al. (2018b). Hydrophobic pore gates regulate ion permeation in polycystic kidney disease 2
and 2L1 channels. Nature communications *9*, 2302. 10.1038/s41467-018-04586-x.

Fig 1

Fig 2

Fig 3

Fig 4

A

B

IB: anti-PKD2

PKD2   Arl13b   Merge   TACAN   Arl13b   Merge

MDCK

IMCD

LLC-PK1

MDCK   IMCD   LLC-PK1

Ctrl IgG   anti-TACAN   Ctrl IgG   anti-TACAN   Ctrl IgG   anti-TACAN

100 —

IP

100 —

Input

Fig 5

Fig 6

Fig 7

Fig 8

Fig S1

Figure S1. Channel activity of TACAN in CHO cells under negative pressures, related to Figure 2. Representative single channel recordings in CHO cells transfected with none (Ctrl), GFP or GFP-TACAN, in the presence of a negative pressure, as indicated.

Fig S2

Figure S2. Oligomerization of TACAN and PKD2 F604P, related to Figure 4. A. Western blot data obtained in the presence of cross-linking, showing expression of TACAN and PKD2 F604P. Cross-linking was carried out with 1 mM disuccinimidyl tartrate (DST). Putative subunit composition of the bands is indicated. **B.** Western blot data obtained in the presence of cross-linking, showing expression of TACAN FL, K135X, and K135X-D343. Putative oligomer conditions are indicated.

Fig S3

Figure S3. Topology of human TACAN and PKD2, related to Figure 4 and 5.
Residues and their sites indicate the positions of the key truncation mutants used in the
study.