Early life exercise primes the neural epigenome to facilitate gene expression and hippocampal memory consolidation.

Raus AM*[1], Fuller TD*[2], Nelson NE*[1], Valientes DA[2], Bayat A[2], Ivy AS[1,2,3,4]

[1]Physiology/Biophysics, Anatomy/Neurobiology, University of California- Irvine School of Medicine, Irvine, CA

[2]Pediatrics, University of California- Irvine School of Medicine, Irvine, CA

[3]Neurobiology and Behavior, University of California- Irvine School of Biological Sciences, Irvine, CA

[4]Division of Neurology, Children's Hospital Orange County, Orange, CA

Correspondence to aivy@hs.uci.edu

*These authors contributed equally to the manuscript.

Abstract

Aerobic exercise promotes neuroplasticity and hippocampal memory. In the developing brain, epigenetic modifications resulting from early-life exercise (ELE) has not been fully explored. In this study we employ a line of transgenic mice expressing the NuTRAP (Nuclear tagging and Translating Ribosome Affinity Purification) cassette in Emx1 expressing cells, allowing for the simultaneous isolation of translating mRNA and nuclear chromatin from a hippocampal neuron-enriched cell population. After mice undergo ELE, we perform a novel technique of Simultaneous Isolation of Nuclei TAgged in specific Cell Types (INTACT) and Translating Ribosome Affinity Purification (TRAP) (together: "SIT") from a single hippocampus. Data from RNA- and CUT&RUN-seq were coupled to evaluate histone modifications influencing the expression of translating mRNA in neurons after ELE. We then evaluated new gene expression – histone modification relationships during memory consolidation that are likely critical for facilitated memory after ELE. Our data reveal novel candidate gene-histone modification interactions for further investigation.

Introduction

Environmental experiences engage epigenetic mechanisms to modulate gene expression and cell function in post-mitotic neurons[1,2]. Histone modifications and DNA methylation are particularly important for neuronal adaptation to environmental signals by altering transcription and synaptic function[3,4]. Behavioral outputs such as stress susceptibility, reward seeking, and long-term memory have been shown to result from changes to chromatin accessibility and gene expression in neurons[5-9]. In addition to this, the neuronal chromatin landscape undergoes waves of epigenetic modifications as a function of brain maturation itself[10-12]. Postnatal periods of heightened sensitivity to environmental stimuli can lead to lasting changes to cellular function and may result from temporally specific epigenetic mechanisms in the developing brain[13,14]. Whether gene regulatory mechanisms in postmitotic neurons are uniquely influenced by early-life experiences to inform long-term function is a question that is just beginning to be explored. Identifying epigenetic processes involved in modulating cell function, particularly during brain development, is critical for understanding how early-life experiences impact long-term behavioral outcomes.

Aerobic exercise has generally been found to enhance performance on cognitive tasks involving the hippocampus in both adult humans and animal models[15,16]. The type, timing, and duration of exercise exposure matters with regard to whether it has a persistent impact on hippocampal function[17-19]. However, in general, the effects of exercise on hippocampal memory in adulthood are transient[17-19]. Findings in adult rodents also implicate a role for histone modifying enzymes in the mechanisms of exercise-induced benefits to hippocampal memory. Both voluntary exercise or treatment with a HDAC3 inhibitor enable hippocampal memory after a subthreshold learning stimulus, increase brain-derived neurotrophic factor (BDNF), and promote acetylation of H4K8 at the BDNF promoter[20-22]. This suggests that exercise engages epigenetic regulatory mechanisms to promote plasticity. Adult exercise also opens a temporal window for persistent improvements to memory performance when a reactivating exercise exposure is introduced[23]. This suggests a "molecular memory" of the initial exercise[23]. Although the majority of these studies have been performed in adults, more recent work suggests that the effects of exercise on hippocampal memory and the associated alterations in

neurotrophic factor expression, synaptic plasticity, and neurogenesis are similar in

juvenile and adolescent periods[17,24-27]. Previous work in our lab showed that early-life

exercise (ELE) for either one week (juvenile period; postnatal days (P) 21-27) or three

weeks (juvenile-adolescence; P21-41) facilitated hippocampal long-term memory

formation in response to a learning stimulus typically insufficient for forming long-term

memory. This finding was associated with increased long-term potentiation (LTP) as well

as modulations to synaptic physiology in hippocampal CA1[17]. Notably, the hippocampal

memory effects of juvenile ELE during P21-27 persisted two weeks after exercise

cessation, which is longer than the effect of exercise on adult hippocampal function[23].

Taking the early-life and adult exercise findings together, it is possible that exercise may

"prime" hippocampal function for response to future experiences. The epigenome is a

strong consideration for this priming effect of exercise: it could serve as a platform for

the molecular memory of exercise by promoting efficient gene expression and

modulating behavioral output[28]. The mechanisms underlying sustained

electrophysiological and behavioral effects of ELE have not been assessed from the

perspective of a potential molecular memory of ELE within the epigenome.

In this study we describe a new approach to simultaneously isolate neuron-

specific chromatin and translating mRNA from a single hippocampal homogenate. Using

this approach, we discover epigenetic signatures of ELE that promote enhancements to

memory and plasticity in the developing brain. We identify unique neuronal gene

expression programs as a resulting from ELE and pair this data with ELE-induced

changes in H4K8ac or H3K27me3. By crossing an Emx1- Cre transgenic line with mice

harboring the NuTRAP (Nuclear tagging and Translating Ribosome Affinity

Purification[29]) cassette (termed "Emx1-NuTRAP" mice) we isolate whole nuclei and

translating mRNA from a single hippocampus by combining the first steps of the

INTACT (Isolation of Nuclei TAgged in specific Cell Types[30]) and TRAP (Translating

Ribosome Affinity Purification[31]) protocols. We have named this protocol "SIT"

(Simultaneous INTACT & TRAP). This approach yields DNA and RNA from the same

homogenate that is of high-quality and sufficient concentrations for our downstream

sequencing procedures RNA- and CUT&RUN-seq. Using previously undescribed

technical approaches for neuron-specific, paired transcriptomic and epigenomic bulk

sequencing, this study uncovers hippocampal gene expression programs and associated

histone modifications that are driven by ELE to promote plasticity. Importantly, these

data reveal an epigenetic signature resulting from ELE that may form the basis of

persistently enabled hippocampal function.

Results

"Emx1-NuTRAP" mouse allows for simultaneous isolation of nuclear chromatin and translating mRNA from a single population of hippocampal neurons.

Our initial goal was to obtain translating mRNA and nuclear DNA from a single population of neurons to couple transcriptional network changes with alterations in histone modifications resulting from ELE. To do this, we developed a new transgenic mouse line (Emx1-NuTRAP) and experimental protocol (SIT protocol). We crossed male NuTRAP reporter mice[29] with female Emx1-Cre[32,33] mouse lines to generate Emx1-Cre; NuTRAP transgenic progeny for this study (Fig. 1A). Emx1-expressing neurons are predominately excitatory[32] and are necessary for hippocampal neurogenesis and motor skill learning[34]. In the presence of a *loxP* site-flanked sequence, Emx1-Cre facilitates recombination in approximately 88% of neurons in the neocortex and hippocampus, and in less than 2% of GABAergic inhibitory interneurons in these regions[32]. Emx1-Cre also targets neural progenitor cells and can be expressed in mature astrocytes in striatum[35]. Of note, we exclusively used Emx1-Cre female mice in our breeding schemes given that the Cre-recombinase has been reported to be expressed in a subset of male germline cells[36].

We sought to validate neuron-specific expression of the NuTRAP cassette. First, immunohistochemistry of Emx1-NuTRAP hippocampal slices show distinct GFP and mCherry expression in CA1 and CA3 pyramidal neurons (Fig. 1B-C) as well as the granule cell layer of the dentate gyrus (DG; Fig. 1D). Furthermore, mCherry positive nuclei remained intact and bound to the magnetic beads after our modified nuclear isolation procedure (Supplementary Fig. 1A). As Emx1-expressing neural stem cells can become astrocytes[32], we sought to determine whether a significant number of mature astrocytes were obtained in the population of isolated cells. Flow cytometry experiments using THY-1 as a neuronal marker and S100β as an astrocytic marker revealed a distinct population of cells double positive for GFP and THY-1, whereas a S100β and GFP double positive cell population was absent (Fig. 1E and Supplementary Table 1).

To perform SIT, hippocampal tissue was dissected from both brain hemispheres, combined, and homogenized in one sample tube (Fig. 1A). We then performed the INTACT and TRAP methods on isolated samples by starting the protocol with the beginning steps from the INTACT procedure, modified to include cycloheximide.

Cycloheximide works rapidly to inhibit protein synthesis and is used for maintaining

crosslinks between translating mRNA and ribosomal subunits during purification in the

traditional TRAP method. Despite the presence of cycloheximide, nuclear morphology

from Emx1-expressing neurons is overall unchanged (Supplementary Fig. 1A).

Additionally, a Bioanalyzer was used to determine if there were cycloheximide-induced

double-stranded DNA breaks in our nuclear preparation that could interfere with

downstream DNA sequencing applications (such as ATAC, CUT&RUN, or CUT&Tag).

We found no evidence of DNA double strand breaks generating fragments of less than

1kb (Supplementary Fig. 2A). Following magnetic purification of biotin-labeled nuclei,

the supernatant fraction was removed and taken through TRAP, while the pelleted nuclei

were processed through the remaining steps of INTACT (see Methods). Using qPCR, we

found that TRAP isolated mRNA had significantly reduced expression of *Mog* and

Cd11b compared to total RNA, suggesting depletion of oligodendrocyte and microglial

populations in TRAP mRNA (Supplementary Fig. 1B). Combining bilateral hippocampi

from a single mouse yielded TRAP-isolated mRNA of high quality and sufficient

concentration for sequencing (RIN > 8 for all samples, average yield RNA = 14.36688

ng/ul; Supplementary Table 2 and Supplementary Fig. 2B). INTACT-isolated nuclei were

further processed using the CUT&RUN (Cleavage Under Targets and Release Using

Nuclease[37]) method to isolate antibody-specific protein-DNA interactions for

downstream DNA sequencing. The resulting DNA libraries were of high quality and

concentration when using specific antibodies (H4K8ac: average size = 1197 bp, average

concentration = 127.8 nM; H3K27me3: average size = 1056 bp, average concentration =

139.0 nM; Supplementary Table 2). In contrast, the resulting DNA libraries using the

non-specific IgG control had substantially lower concentrations (IgG: average size =

1075 bp, average concentration = 16.3 nM; Supplementary Table 2) further indicating

that nuclear DNA from both isolations was of high starting quality.

To confirm our TRAP-isolated hippocampal mRNA came primarily from

excitatory neurons, we took a bioinformatics approach to assess for neuronal gene

enrichment. Specifically, we compared RNA-seq data from whole dorsal hippocampal

tissue of wild-type mice to our TRAP-isolated RNA-seq data. We found that TRAP-

isolated mRNA had significant enrichment of several neuronal genes including *Dlg4*,

Thy1, *Eno2*, and *Syt4*, as well as a significant reduction in astrocytic, microglial,

oligodendrocyte, and inhibitory neuron genes (Fig. 1F). There was a relative expression

increase of the glial fibrillary astrocytic protein (GFAP) in our TRAP samples. This gene

can also be expressed in neural stem populations that were present in our whole

hippocampus samples, so this finding may be a reflection of the neural stem cell

population of the dentate gyrus captured as part of our TRAP isolation[38,39]. Overall, these

findings suggest that the Emx1-NuTRAP mouse model is a valid tool for neuron-enriched

isolation of sequencing-grade, translating mRNA and nuclear chromatin from the same

cell homogenate.

Figure 1. Generation and validation of the Emx1-NuTRAP mouse (A) Schematic

representation of the Emx1-NuTRAP mouse generation and workflow of the SIT

protocol. (B) Immunofluorescence imaging of the CA1 region of the hippocampus at 20x

objective and (B') 60x objective after incubation with an mCherry antibody overnight.

(C) Immunofluorescence imaging of the CA3 and CA2 region of the hippocampus at 20x

objective and (C') 60x objective after incubation with an mCherry antibody overnight.

(D) Immunofluorescence imaging of the DG region of the hippocampus at 20x objective

and (D') 60x objective after incubation with an mCherry antibody overnight. Scale bars

are set to 50μm for all images (B-D). (E) Flow cytometry for EGFP, THY-1, and S100β

(15.3% of cells THY-1+/EGFP+ and 0.13% of cells S100β+/EGFP+). (F) Heatmap of

differentially expressed neuronal and other cell type markers from RNA-seq data

comparing TRAP isolated RNA from our Emx1-NuTRAP mice and previously generated

dorsal hippocampal tissue RNA from wild type.

Performing INTACT/TRAP on the same or separate cell suspensions yields highly comparable RNA-seq and CUT&RUN-seq results.

Previous methods for performing INTACT and TRAP using NuTRAP mice have taken separate tissue homogenates for each procedure[29,40]. In this study, we performed SIT on a single tissue homogenate containing bilateral hippocampi (the products of SIT are referred to as "simultaneous isolations"). To determine if our approach of SIT compares well with separate INTACT and TRAP procedures, we also performed separate INTACT and TRAP from hippocampal tissue obtained from one brain hemisphere for each method, counterbalancing for left versus right hemispheres (we refer to this protocol as "separate isolations"). A Bioanalyzer was used to determine the amount and quality of the RNA obtained from each type of isolation (Separate isolations: average RNA concentration = 7.805 ng/ul, average RIN = 9.5; Simultaneous isolations: average RNA concentration = 14.367 ng/ul, average RIN = 9.3; Supplementary Table 2). The average RNA yield from the separate isolations (using a unilateral hippocampus) was approximately equal to half of the average yield of the simultaneous isolations (which combined bilateral hippocampi; Supplementary Table 2). Similarly, the final library concentrations for the separately isolated IgG CUT&RUN-seq libraries were also approximately half the concentration of the simultaneous isolations (average simultaneous: 24.2nM, average separate: 11nM; Supplementary Table 2). We interpret this to mean that nuclear DNA was fully intact in the simultaneous isolation because we did not obtain substantially more than double the concentration in the simultaneous vs separate isolations. Unilateral hippocampal homogenates yielded sufficient sequencing concentrations and quality to allow for library preparations from individual mice (Supplementary Table 2).

To determine if normalized RNA-seq counts were similar between the two isolation methods, we performed a Spearman's correlation between the datasets from sedentary mice. The two RNA-seq datasets were found to be highly correlated, with R values greater than 0.5 and p value less than 2.2×10^{-16} (R=1, $p<2.2 \times 10^{-16}$; Fig. 2A). CUT&RUN-seq was used to identify genomic regions interacting with either H4K8ac, an activating histone post-translational modification (PTM), or H3K27me3, a generally repressive histone PTM. We again applied Spearman's correlation to understand whether

the simultaneous vs separate INTACT isolation methods could influence CUT&RUN-seq

peak distribution. We compared normalized count data for CUT&RUN-seq peaks across

a representative chromosome (chromosome 2). We binned 100bp increments along the

entire chromosome from the simultaneous and separate isolations using datasets

generated from sedentary mice. Normalized sequencing counts, which reflected reads

assigned to binned genomic regions along chromosome 2, were highly similar between

conditions (H4K8ac: R=0.65, $p<2.2\times10^{-16}$, H3K27me3: R=0.81, $p<2.2\times10^{-16}$; Fig. 2B-C).

Taken together, these experiments suggest that translating mRNA and nuclear DNA

isolated from hippocampal homogenates using either simultaneous or separate isolation

procedures are comparable in terms of quality, concentration, functional characterization,

and normalized sequencing reads.

We next wanted to determine if the different isolation methods could bias resulting

gene expression on the basis of gene length. When plotting gene length against log fold

change of gene expression, we see a similar distribution pattern of ELE-induced

differentially expressed hippocampal genes (DEGs) between the simultaneous and

separate methods (Fig. 2D-E). We then compared the ELE-induced DEGs obtained using

either the separate or the simultaneous isolation methods. Although many of the genes

did not overlap (Fig. 2F-G and Supplementary Table 3), most Gene Ontology (GO) terms

did (14/20 GO terms for upregulated genes and 15/17 for downregulated genes; Fig. 2H-

K and Supplementary Table 3).

Figure 2. Simultaneous and separate isolations of DNA and RNA from Emx1-

NuTRAP mice are comparable. For Fig. 2A-E, n=2 mice for simultaneous isolations

and n=3 mice for separate isolations. (A) Spearman's correlation for RNA expression

between separately isolated RNA and simultaneously isolated RNA (R=1, $p<2.2\times10^{-16}$).

(B) Spearman's correlation between separate and simultaneously isolated DNA from

CUT&RUN-seq for H4K8ac with 100bp chromosome 2 position bins (R=0.65, $p<2.2\times10^{-16}$

^{16}). (C) Spearman's correlation between separate and simultaneously isolated DNA from

CUT&RUN-seq for H3K27me3 with 100bp chromosome 2 position bins (R=0.81,

$p<2.2\times10^{-16}$). (D) Gene length versus fold change distribution plot for simultaneously

isolated RNA. (E) Gene length versus fold change distribution plot for separately isolated

RNA. (F) Genes upregulated in Emx1-NuTRAP RNA-seq after ELE in either

simultaneous (n=2 sedentary mice and n=3 ELE mice) or separate isolations (n=3 mice

per group). (G) Genes downregulated in Emx1-NuTRAP RNA-seq after ELE in either

simultaneous or separate isolations. (H, I) Panther Gene Ontology: Biological Processes

Venn diagram (H) and bar graph of gene distributions (I) for genes upregulated after ELE

in either separate or simultaneous isolations. (J, K) Panther Gene Ontology: Biological

Processes Venn diagram (J) and bar graph of gene distributions (K) for genes

downregulated after ELE in either separate or simultaneous isolations.

Neuron-specific gene expression and histone PTMs are functionally comparable between left and right hippocampal hemispheres.

Many studies take advantage of the brain's structural symmetry by using tissue from each hemisphere for separate molecular processing. Prior evidence demonstrates that hippocampal lateralization can influence LTP, hemisphere-specific glutamate receptor density, and performance in certain memory tasks[41]. We wanted to determine if there were differences in transcriptional programs resulting from ELE in left vs right hippocampi. Using the separate isolation approach described above, we compared ELE-induced differential gene expression between left and right hemispheres. We found that although individual gene expression patterns were different between hippocampi originating in the left and right brain hemispheres (Fig. 3A-B and Supplementary Table 4), the Panther GO: Biological Processes terms were substantially similar with most categories overlapping (Fig. 3C-F and Supplementary Table 4). This could suggest spatial or hemispheric assignment of genes with functional similarity.

Finally, we wanted to determine if left and right hippocampal hemispheres had significant differences in DEGs and CUT&RUN-seq peak distributions at baseline (without exercise). We performed Spearman's correlation on the transcript counts from the RNA-seq (R=1, p<2.2x10^{-16}; Fig. 3G), and the CUT&RUN-seq 100 bp binned peak counts along chromosome 2 for H3K27me3 (R=0.70, p<2.2x10^{-16}; Fig. 3H) and for H4K8ac (R=0.59, p<2.2x10^{-16}; Fig. 3I). .We found that the seq peaks and transcript counts were very similar with significance considered as R values greater than 0.5 and p value less than 2.2 x 10^{-16} (Fig. 3G-I). These data suggest that in the sedentary condition, left and right hippocampal hemispheres yield similar normalized sequencing counts.

Figure 3: Histone modifications and functional categorizations of gene expression
are similar regardless of hemispheric origin of hippocampal tissue. (A-B) Venn
diagram of genes upregulated (A) or downregulated (B) by ELE identified between the
left and right hemispheres by the separate isolation protocol RNA-seq (n=3 mice per
group). (C-D) Venn diagram of Panther Gene Ontology: Biological Process terms for
genes upregulated (C) or downregulated (D) by ELE identified between the left and right
hemispheres by the separate isolation protocol RNA-seq. (E-F) Percent of genes fitting
into each GO category for the left and right hemisphere for genes upregulated (E) and
downregulated (F) by ELE by separate isolation RNA-seq. (G) Spearman's correlation
between the left and right hemispheres RNA-seq data from the separate isolation protocol
of sedentary rlog normalized expression (R=1, $p<2.2 \times 10^{-16}$). (H) Spearman's correlation
between hemispheres for CUT&RUN-seq for H4K8ac normalized count data for
chromosome 2 in 100bp bins (R=0.59, $p<2.2 \times 10^{-16}$). (I) Spearman's correlation between
hemispheres for CUT&RUN-seq for H3K27me3 normalized count data for chromosome
2 in 100bp bins (R=0.7, $p<2.2 \times 10^{-16}$).

Early-life exercise promotes expression of plasticity-related genes and is associated with transcriptional regulatory pathways implicated in hippocampal memory.

To identify the transcriptional effects of ELE in hippocampal neurons, we performed cell-type specific bulk sequencing on RNA extracted from Emx1-NuTRAP hippocampi using the SIT approach. We compared DEGs from sedentary mice vs mice that underwent voluntary wheel running between postnatal days (P) 21-41 (Supplementary Fig. 1C). Using DESeq2 to assess for DEGs (>30% expression increase with a p value < 0.05), we found that voluntary exercise during juvenile-adolescence significantly alters gene expression in Emx1 expressing neurons (297 upregulated and 338 downregulated genes; Fig. 4A and Supplementary Table 3). Many of the genes upregulated after ELE are also known to be involved in exercise or hippocampal memory mechanisms, including *Bdnf* [42] and *Nr4a1* [43]. To functionally categorize ELE-induced DEGs, we performed a Panther Gene Ontology (GO) analysis [44] focusing on the Molecular Function categorization and separated by upregulated and downregulated genes. The GO term categories with the most genes functionally assigned to them were binding, catalytic activity, molecular function regulator, transporter activity, molecular transducer activity, and structural molecule activity (Fig. 4B). Many of the genes driving these results are known to be important in neuronal function (*Kcna1, Slc24a4, Stxbp5l, Gabra2,* and *Camk2n2*), neurodevelopment (*Artn, Kdm7a, Sox21, Gap43,* and *Efna5*), and hippocampal memory (*Bdnf, Nr4a1* and *Dusp5*).

To evaluate possible transcription factors and upstream regulators implicated by ELE-induced DEGs, we applied Qiagen's Ingenuity Pathway Analysis (IPA) to our RNA-seq dataset [45]. We identified significant canonical signaling pathways implicated in ELE effects on hippocampal neuronal function (Supplementary Fig. 4A and Supplementary Table 5). Additionally, the top six upstream regulators by significance included CREB1, KMT2A, LEF1, SMAD3, EGR1, and MITF. Each of the top five transcription factors have been implicated in hippocampal development or function (Fig. 4C and Supplementary Table 5) [46-53]. The transcription factors KMT2A and LEF1 have not been previously associated with the effects of exercise on hippocampal function. Interestingly, CREB1 (through its associated CBP [54]) and KMT2A (through its methyltransferase activity [47]) both have histone modifying properties. LEF1 has been shown to regulate

neural precursor proliferation in the hippocampus[48]. SMAD3 is critical for intermediate
progenitor cell survival and negatively regulates serum insulin after exercise[49,50]. EGR1 is
an immediate early gene that recruits TET1 DNA demethylase during neural activation
and development[51]. MITF was the sixth upstream regulator identified and is a
transcription factor linked to autophagy mechanisms[52]. These specific transcription
factors were not significantly differentially expressed in our RNA-seq dataset; however,
their activity may not be linked to a change in their own expression after ELE but rather
modulated by exercise to influence programs of gene expression.

To further understand whether the transcriptional profiles resulting from ELE were
enriched for specific *a priori* assigned molecular functions, we performed Gene Set
Enrichment Analysis (GSEA)[55]. We evaluated the "Reactome" category of functional
gene sets followed by a leading-edge analysis to further determine which genes were
driving the significant categories (Fig. 4D and Supplementary Table 6). Of the DEGs
upregulated after ELE, several interesting categorizations and the genes driving them
were revealed. The genes *H3c4*, *H2bc3*, *H2bc5*, and *H2bc9* genes, which encode histone
family member proteins, were driving the categories DNA methylation, HDACs
deacetylate histones, and transcriptional regulation by smRNAs. Meiosis and meiotic
recombination also emerged in these results which was unusual; however, leading edge
analysis showed many of these genes (*Atm, Blm, Msh4*, and *Mnd1*) to be generally
involved in cell-cycling processes. Exercise is well known to increase adult neurogenesis
in hippocampal dentate gyrus and can explain cell cycle gene enrichment in our dataset[16].
Several nucleoporin complex genes (*Nup35/ 42/ 43/ 50/ 58/ 62/ 93/ 107/ 153/ 210*) were
also identified for their associations with categories such as: gene silencing by RNA,
nuclear envelope breakdown, transport of mature mRNAs derived from intronless
transcripts. *Nup* genes form a variety of nuclear pore complexes that play critical roles in
cellular processes including cell-cycle regulation, cellular differentiation, and epigenetic
control[56].

Of the downregulated pathways identified, translation-associated categories
(translation, eukaryotic translation initiation/elongation, and SRP-dependent co-
translational protein targeting to membrane) were driven by significant downregulation of
ribosomal protein gene families (*Rps*, *Rpl*, *Rplp*) and eukaryotic initiation factor 3 (*eIF3*)

subunits. Over-expression of *eIF3* subunits has been linked to Parkinson's Disease and

altered expression of *eIF3* has been associated with neurodevelopmental disorders[57].

Additionally, collagen genes (*Col1a1*, *Col1a2*, *Col2a1*, *Col3a1*, and *Col5a1*) were

downregulated after ELE, leading to the identification of categories including: MET

activates PTK2 signaling, MET promotes cell motility, and collagen degradation. *Col1a1*

and *Col1a2* have been previously identified as putative aging genes that decrease in their

expression after chronic exercise in adults[58]. By analyzing these enriched pathways and

the networks of genes driving them, we were able to identify several transcriptional

programs activated by ELE that could be candidates for further exploration.

Figure 4: ELE leads to transcriptomic changes in hippocampal neurons during

adolescence. (A) Volcano plot of differentially expressed genes reaching significance

identified by RNA-seq on ELE versus sedentary (n=2 sedentary mice and n=3 ELE mice,

absolute value $\log_2$ fold change > 0.3785 and p-value < 0.05). (B) Panther Gene

Ontology: Molecular Function top terms by most genes assigned. (C) Top 6 upstream

regulators identified by Ingenuity Pathway Analysis (IPA). (D) Representative Gene Set

Enrichment Analysis: Reactome leading-edge diagrams showing genes upregulated (D)

or downregulated (D') in ELE, and their enrichment. *Abbreviated terms in (D) transport

of mature mRNAs derived from intronless transcripts, (D') activation of the mRNA upon

binding of the cap binding complex and EIFs and subsequent binding to 43S.

Neuronal H4K8ac is enriched and H3K27me3 is reduced at a subset of plasticity genes after ELE.

371
372
The power of the Emx1-NuTRAP mouse model coupled with the SIT procedure
is the ability to directly correlate differential gene expression with candidate epigenetic
mechanisms in the same cell population. Hippocampal nuclei were obtained using
INTACT methods in our SIT procedure, followed by CUT&RUN-seq from nuclear
chromatin. We investigated ELE-induced changes in histone modifications H3K27me3
and H4K8ac across the genome. H4K8ac is a permissive histone modification that is
enriched at *Bdnf* after adult exercise, and its presence correlates with improved
hippocampal memory[22]. H3K27me3 is a repressive histone mark and a common control
in CUT&RUN-seq studies[37,59] and is decreased at the *Bdnf* promoter region after
contextual fear conditioning training[60]. Peaks were called using SEACR[61] and histone
modification enrichment peaks were overlapped with differentially expressed genes (Fig.
5A). We correlated H4K8ac and H3K27me3 peaks present with upregulated and
downregulated mRNA as a result of ELE. The loss of either H3K27me3 or H4K8ac did
not correlate with changes in mRNA expression. This may indicate that ELE-induced
additions of these two histone modifications, but not their removal, underlie differential
gene expression. 93 upregulated genes correlate with new H4K8ac peaks while 33
downregulated genes correlate with new H3K27me3 peaks (Fig. 5B and Supplementary
Table 7). However, loss of only one H4K8ac peak correlates with downregulation of
mRNA, and loss of only two H3K27me3 peaks correlate with upregulation of mRNA
(Fig. 5B and Supplementary Table 7). Unsurprisingly, many of the peaks we discovered
newly present because of ELE did not correlate with expression changes in many genes.
Notable upregulated genes with new H4K8ac after ELE are *Prox1*, *Sntb2*, and *Kptn*.
PROX1 is involved governing differentiation, maturation, and DG versus CA3 cell
identity in intermediate progenitor cells of the hippocampal DG[62], while SNTB2 is
involved in G protein-coupled receptor cell signaling[63]. KPTN encodes a protein involved
in cytoskeletal cell structure, and its mutation can cause neurodevelopmental disability
and seizures[64]. Interestingly, downregulated genes found to have new H3K27me3 after
ELE included *Col3a1*, *Efnb2*, *Epop*, and *Myoc*. COL3A1 and MYOC are structural
proteins involved in the extracellular matrix and the cytoskeleton respectively[65-67].

EFNB2 is a signaling molecule involved in cell migration and its haploinsufficiency can

cause neurodevelopmental disability[68-70]. EPOP is a known editor of the chromatin

landscape by altering H2Bub and H3K4me3 distributions[71,72]. Interestingly, many of the

histone modification changes that occurred following exercise appear to be in distal

intergenic regions for H3K27me3 and intronic regions for H4K8ac (Fig. 5C). To

investigate how ELE-induced changes to histone PTM distribution might alter gene

expression, we looked at the distribution of peaks around *Prox1* (Fig. 5D). Most apparent

changes in histone PTM peaks occurred in regions annotated as regulatory regions, rather

than the promoter region, which is concordant with our data showing that most histone

PTM changes happen outside of the promoter region. Taken together, these results

suggest that ELE promotes the new addition of H4K8ac and H3K27me3 PTMs to

regulate gene expression. Given that most identified peaks did not correlate with DEGs,

they may instead implicate regions of interest that are "primed" for facilitated gene

expression following future stimuli, such as learning.

A

H4K8ac
ELE

H4K8ac
Sedentary

RNA Up
in ELE
H3K27me3
ELE
H3K27me3
Sedentary

A'

H4K8ac
Sedentary

62
2290
5505
H4K8ac
ELE
RNA Down
in ELE
H3K27me3
ELE
H3K27me3
Sedentary

B **Presence of Histone Marks**

Up in ELE

ELE
Only

ELE &
Sedentary

Sedentary
Only

# of Genes w/ RNA Expression

Down in ELE

H3K27me3   H4K8ac

C **Relative Distribution of Annotated Genomic Regions**

% of Total Peaks

	H3K27me3 - Sedentary	H3K27me3 - ELE	H4K8ac - Sedentary	H4K8ac - ELE
top	4.92	4.15	2.77	2.39
	7.63	8.24	10.10	11.13
Distal Intergenic	41.35	46.68	32.17	28.95
Other Exon	6.87	6.94	4.84	5.49
Other Intron	10.88	12.92	22.47	24.58
Promoter (<1kb)	21.13	13.63	19.90	19.31

Legend:
- 1st Exon
- 1st Intron
- 3' UTR
- 5' UTR
- Distal Intergenic
- Downstream (<0.3kb)
- Other Exon
- Other Intron
- Promoter (<1kb)
- Promoter (1-2kb)
- Promoter (2-3kb)

D **PROX1 Gene Tracks:** chr1:189,900,750-190,843,610

Regulatory Element   Promoter Region

Figure 5: ELE alters H4K8ac and H3K27me3 occupancy. (A) Venn diagram of gene
peaks identified in CUT&RUN-seq for H4K8ac and H3K27me3 by SEACR (top 1% of
peaks FDR<0.1) overlapped with genes upregulated (A) and downregulated (A') in RNA
by ELE (n=2 sedentary mice and n=3 ELE mice). (B) CUT&RUN-seq SEACR peak calls
at genes that overlap with genes differentially expressed by ELE as determined by RNA-
seq. (C) Stacked bar of the relative distribution of peak calls at various genomic regions
identified by CUT&RUN-seq called using SEACR. (D) Representative track of
CUT&RUN-seq data around *Prox1* showing differential peak height between ELE and
sedentary conditions for H4K8ac and H3K27me3.

ELE induces changes in H4K8ac and H3K27me3 occupancy at promoters of genes involved in hippocampal memory consolidation.

Prior work from our lab and others has found that both ELE and adult exercise can facilitate hippocampal long-term memory formation in mice exposed to a typically subthreshold learning event (3 minutes of object location memory (OLM) training; typically insufficient for LTM formation in sedentary mice). These results suggest that ELE may "prime" neuronal function to facilitate hippocampal learning. Here we test the hypothesis that ELE enables memory consolidation via altering presence of the histone modifications H4K8ac and H3K27me3 in promoters of genes associated with learning and hippocampal plasticity. These changes could potentially allow for more rapid transcription of memory-related genes. Wildtype animals underwent ELE followed by object location memory training on p41 and were sacrificed 60 minutes later (during memory consolidation; Fig. 6A). Dorsal hippocampi were dissected and processed for RNA-seq, and these data were compared to the previously identified differential gene expression, H4K8ac, and H3K27me3 occupancy after ELE without learning (generated from Emx1-NuTRAP CUT&RUN-seq; Fig. 5). We categorized differentially expressed genes based on the experimental group: ELE alone, ELE+3 min learning stimulus, Sedentary+3 min learning stimulus, and Sedentary+10 min learning stimulus. To discover ELE-induced genes "primed" for regulation during memory consolidation, we focused specifically on candidate genes associated with memory after exercise (cGAME), candidate genes associated with memory after exercise or sedentary (cGAMES), and candidate genes associate with memory after sedentary (cGAMS) conditions (Fig. 6B,C). We then evaluated cGAME and cGAMES groups of genes for the new presence or absence of histone modifications H4K8ac and H3K27me3 after ELE, suggesting addition or removal of these histone marks denoting a permissive (increased gene expression) or repressive (reduced gene expression) chromatin state, respectively (Fig. 6B',C').

In the cGAMES group, there were 145 genes found to be significantly increased in the mice that underwent a threshold learning event (3 min for ELE; 10 min for sedentary mice). Of these 145 genes, 40% of them (58) had new H4K8ac following ELE. We consider this group of genes to potentially be "primed," or readied, by ELE for rapid transcription to support hippocampal memory consolidation at 3 min OLM training,

while also being involved in memory consolidation mechanisms in the threshold learning of sedentary mice (10 min OLM training). The cGAME group of genes are those which significantly increased in the ELE mice that underwent a threshold learning event but were not increased in any of the other groups (including the sedentary, threshold learning event / cGAMS group). These genes may be of unique importance for memory consolidation specifically occurring after ELE given their absence in sedentary learners. We found that the cGAME group had significant upregulation of 256 genes, with approximately 30% (76) of those genes receiving new H4K8ac following ELE alone (Fig. 6 B-C'). We also evaluated the cGAMES and cGAME that were significantly downregulated during memory consolidation. We found 304 cGAMES and 361 cGAME genes (Fig. 6 B,C). In both of these groups, a small minority of down-regulated genes had altered H3K27me3 after ELE, and this was mostly gain of H3K27me3 (cGAMES: 6.3%; cGAME: 6.4%) (Fig. 6 B',C'). The consolidation expression data is listed in in Supplementary Table 8. The lists of overlap between these groups is listed in Supplementary Table 9.

We wanted to determine how ELE "priming" directed gene expression during memory consolidation that occurs after exercise (ELE 3min OLM) relative to sedentary memory consolidation (Sed 10min OLM). We compared these to a sub-theshold learning event (Sed 3min OLM training). We compared the relative $\log_2$ fold changes (LFCs) of the genes in cGAME and cGAMES using z-scores for each group across the genes (Fig. 6D). In the vast majority of these primed genes, the ELE drove gene expression during 3 min of OLM in the same direction as 10 minutes of OLM training in sedentary animals (Fig. 6D). In some cases the gene expression changes were more exaggerated in the same direction as a sedentary threshold event (Fig. 6D). We interpret this to indicate that ELE enables a 3min consolidation event to be threshold (when 10 minutes is required for a sedentary mouse) by priming these genes for altered gene expression in the same direction as a sedentary threshold event.

We next identified upstream regulators of cGAME and cGAMES genes with ELE-induced histone modifications, as these mediators are likely involved in long-term memory formation of a subthreshold learning event following ELE. The cGAMES and cGAME lists were evaluated using Ingenuity Pathway Analysis (IPA). First, candidate

regulators were identified for the cGAME and cGAMES groups that have histone PTM

changes as a result of ELE that we measured (Fig. 6E, Supplementary Table 10).

Inhibition of MECP2 may indicate a reduction in DNA methylation. CREB1 is required

for LTM[46,73]. SNCA may play a role in neuroplasticity by modulating synaptic vesicle

transport[74]. Next, we wanted to identify gene networks that might be at play that are not

just regulated by the histone PTMs we selected. Many more histone modifications exist

and may regulate gene expression during consolidation in addition to those we selected.

We investigated the genes in cGAME and cGAMES using IPA and identified a set of

likely upstream regulators for these genes taken together (Fig. 6E'', Supplementary Table

10). Once again MECP2 inhibition and CREB1 activation appear to be indicated

validating their presence in the previous IPA. Notably new to this group are inhibition of

FMR1, KMT2D, and TCF20, along with activation of FASN, CTNNB1, and MKNK1.

FMR1 is involved in mRNA trafficking in neurons; so altered mRNA trafficking may be

an important modulator of ELE enabled consolidation. CTNNB1 is necessary for Wnt

signaling. Next, we interrogated what networks might be unique to exercise enabled

consolidation by running the cGAME list through IPA analysis (Fig. 6E', Supplementary

Table 10). Unique to this group, RTN4, an inhibitor of neurite outgrowth[75], was predicted

to trend towards inhibition. GRIN3A, a glutamatergic NMDA receptor important to

synaptic development and refinement[76], was predicted activated in this group. These

findings suggest that ELE may enable consolidation by increasing neurite and synapse

growth and development. TCF7L2 is also predicted to be activated. Given that this

transcription factor is involved in Wnt signaling, and neurogenesis[77], and activated

CTNNB1 is also predicted to be associated with threshold consolidation generally,

altered Wnt signaling may be critical for ELE enabled consolidation.

Figure 6: ELE may prime consolidation induced gene expression. (A) Experimental
design diagram. (B) Venn diagram representing upregulated genes relative to mice
without ELE or learning in either ELE, ELE and 3 min OLM training, Sed and 3 min
OLM training, and Sed and 10 min OLM training. (B') Pie chart showing the relative
percentage of genes that have new H4K8ac presence or H3K27me3 loss as a result of
ELE in specific groups identified in the Fig. 6B labeled cGAME, and cGAMES. (C) As
Fig. 6B but for downregulated genes relative to mice without ELE. (C') Pie chart of
downregulated genes in the cGAME or cGAMES groups that have new H3K27me3 or
new loss of H4K8ac as a result of ELE. (D) Z-score based heatmap of genes in the
cGAMES and cGAME groups that have new H4K8ac or H3K27me3 presence or loss as
a result of ELE. Expression values used to calculate the Z-scores were relative to mice
that did not experience OLM training or ELE. (E) Upstream regulators identified by IPA
of the genes represented by Fig. 6D. (E') Upstream regulators identified by IPA of the
genes in the cGAME (up and downregulated) group that are unique to that when only
those are included in the IPA analysis (E'') Upstream regulators identified by IPA of the
genes in the cGAME and cGAMES (up and downregulated) group when the genes in
those groups are put through IPA together. (* indicates that the activated or inhibited
indicator is registered from the IPAs based on the z-score for that regulator's expression
and not directly indicated based on software prediction).

Discussion

Identifying gene regulatory mechanisms activated by exercise during sensitive

developmental periods can aid in understanding how early-life exercise may promote a

"molecular memory" of exercise[23,60], via the epigenome, that informs long-term cell

function and behavioral outcomes. Here we present results demonstrating neuron-

specific, simultaneous characterization of translating mRNA and associated histone

modifications from a single population of hippocampal neurons obtained from ELE mice

to reveal molecular signatures of the early-life exercise experience. To our knowledge,

this is the first report describing use of the NuTRAP construct in a predominantly

neuronal population. Additionally, our unique experimental approach is a technical

advance for NuTRAP applications: by modifying the nuclear chromatin isolation

(INTACT) procedure to include cycloheximide, one can obtain both polyribosome-bound

mRNA (using TRAP) and nuclear DNA from a single lysate to directly correlate gene

expression programs with epigenetic regulatory mechanisms in the same set of cells. The

field of neuroepigenetics has been challenged by very limited approaches for coupling

cell-type specific transcriptional profiles with chromatin modifications without

compromising cell integrity (through flow sorting) or genomic coverage (through single-

cell sequencing). Therefore, the NuTRAP model offers an exciting opportunity to

overcome these challenges and perform paired transcriptomic and epigenomic analyses in

any brain cell type of interest for which a Cre line exists (cite other nutrap papers here).

Our study focuses on the impact of exercise on hippocampal gene expression and

associated epigenetic modifications specifically during juvenile – adolescent periods (the

fourth through sixth postnatal weeks of life in mice). We discover gene expression

programs, and alterations in the histone modifications H4K8ac and H3K27me3, that may

be unique to this early-life period. There have been recently published reports

investigating the effects of voluntary exercise on adolescent and adult hippocampal gene

expression and epigenetic modifications through RNA and DNA sequencing methods

from heterogenous cell populations[19,20]. In our study, several genes with either

chromatin-modifying or neurogenesis-regulating properties were either upregulated with

new H4K8ac (*Prox1*, *Sntb2*, and *Kptn*), or downregulated with new H3K27me3 (*Col3a1*,

Efnb2, *Epop*, and *Myoc*). For example, ELE could promote the expression of *Prox1* by

the addition of H4K8ac, and in turn, *Prox1* can promote neurogenic fate and cell location

assignment (such as DG granule cell fate over CA3 pyramidal cell identity[78]). ELE was

also found to alter the expression of *Kptn,* a gene encoding a protein involved in actin

dynamics and neuromorphogenesis[64]. Increased *Kptn* expression and accessibility may

possibly facilitate cytoskeletal changes occurring with altered synaptic plasticity after

exercise. Overall, the differentially expressed genes and their associated chromatin states

identified in this study provide molecular candidates involved in the effects of ELE on

hippocampal function.

Traditional epigenetic and gene expression analyses of bulk hippocampal tissue

are confounded by the cellular heterogeneity of the samples. Further, current methods for

paired analysis of the epigenome and transcriptome are restricted by the need for multiple

transgenic animal models or the use of highly expensive single-cell sequencing

technologies. However, it is critical to obtain epigenomic information in a cell-type

specific manner because there are unique epigenetic modifications distinguishing cell

subtypes of the central nervous system[10,79]. Roh et al. first developed the NuTRAP mouse

to characterize adipocyte-specific and hepatocyte-specific epigenetic and transcriptional

profiles[30]. The NuTRAP mouse has since been used to perform paired transcriptome and

DNA modification analysis in astrocytes and microglia, demonstrating its utility in CNS

cell-types[40]. The Emx1-NuTRAP mouse described here provides rapid, cost-effective

characterization of neuron-specific changes to epigenetic state and gene expression using

whole-tissue homogenates containing multiple cell types. In our technical validation

experiments, the SIT procedure produced high quality RNA and DNA of sufficient

concentration for downstream sequencing studies from a single mouse hippocampus.

This method effectively increases the total sample size used for each analysis as the tissue

sample does not need to be separated for DNA and RNA isolation. CUT&RUN and

CUT&Tag allow for low-input cell-specific chromatin analysis[61,80] further reducing the

required amount of starting material. Indeed, the DNA obtained from SIT could be used

for other applications for evaluating the chromatin landscape, including ATAC-seq.

A common approach to paired transcriptomic and epigenomic analyses is to use

separate brain tissue from one hemisphere for gene expression studies and the other side

for epigenetic analyses[81]. A previous study utilizing the NuTRAP allele in non-neuronal

brain cells used separate brain hemispheres to perform INTACT and TRAP[82]. However,

this assumes that cells of the same type from each half of the tissue will show the same

epigenetic and transcriptional profile. This is not necessarily the case for all cell types,

especially in the CNS, as structures of the brain can show high degrees of hemispheric

specialization[83-87]. To address the degree of transcriptional or epigenetic difference

between hippocampi from opposite hemispheres, we performed separate INTACT and

TRAP isolations using hippocampal tissue from one hemisphere for DNA isolation and

the contralateral hippocampus for mRNA isolation. We found that although differential

gene expression and histone modifications differed between left and right hippocampi,

functional analysis using gene ontology revealed similar biological processes between the

two sides.

Transcription is required for both hippocampal memory and LTP, a synaptic

plasticity response considered as the cellular correlate of learning and memory[88]. We

found that ELE can facilitate long-term memory formation in the hippocampus[17]. We

hypothesize that this may be due to altering the epigenomic landscape to promote a

transcriptional state supporting memory consolidation. In adult rodents, long-term

memory is formed in response to a subthreshold learning event after exercise or HDAC

inhibition and involves acetylation of neuroplasticity gene promoters[22]. Our study relates

this finding to the adolescent mouse brain and identifies new gene expression networks

and their epigenetic regulation likely involved in the hippocampal memory benefits of

ELE. We also identified a group of genes, cGAMS, that are altered in their gene

expression only in sedentary animals undergoing a threshold learning event. The

separation of this group from the cGAME and cGAMES groups indicates that ELE may

engage a separate network of genes to promote memory consolidation.

A next step of this work will be to test the necessity and sufficiency of the

identified gene networks, their histone modifications, and/or their upstream regulators

mediating the effects of ELE on hippocampal memory performance. Another unanswered

question is whether the effects of ELE on hippocampal memory persist well after

exercise cessation, and whether there is stability of ELE-induced epigenetic

modifications to allow for the persistence of ELE effects.

In summary, our work reveals ELE-induced transcriptomic and epigenomic signatures in hippocampal neurons using the Emx1-NuTRAP mouse for such comparisons. Presently, the field of neuroepigenetics is shifting to single-cell resolution for sequencing studies. Our study was performed in bulk tissue enriched for neurons. As the goal of this work was to characterize the transcriptional and epigenomic effects of early-life exercise, cell subtype identity (as is often a priority in single-cell approaches) was not necessary for tour initial hypotheses but would offer a second layer of complexity to address this study's questions. Single-cell approaches can bias against lowly-expressed genes, which presents the risk of not capturing small expression changes in rate-limiting genes that may have large changes in function as a result of ELE. A limiting factor for the use of the NuTRAP transgenic mouse is the requirement of a Cre line to exist for the experimenter's cell type of interest. In our Emx1-NuTRAP model the Cre-recombinase in Emx1 expressing cells is constitutively expressed, and this may present off-target effects when present during development. Despite this, our study uses a feasible, previously undescribed approach in the field of neuroepigenetics to pair transcriptomic and epigenomic sequencing results from a population of hippocampal neurons. Our approach revealed new molecular epigenetic candidates for further mechanistic studies to examine the effects of early-life exercise on hippocampal function. Furthermore, understanding how exercise uniquely impacts neuronal function during juvenile-adolescent developmental periods could provide insights into an early-life epigenetic memory of the ELE experience, revealing possible targets for improving memory function or mitigating memory deficits in the setting of early-life cognitive and neurodevelopmental disorders[89].

Acknowledgements

This work was supported by the National Institute of Neurological Disorders and Stroke of the National Institutes of Health, under award number K12NS098482, the Conte Center @ UCI Seed Grant, UC Irvine Institute for Clinical and Translational Sciences Pilot Award, and the Robert Wood Johnson Foundation Amos Medical Faculty Development Program (awarded to A.S.I.). We thank the UC Irvine School of Medicine, Departments of Pediatrics for support. We would also like to thank Dr. Marcelo Wood

653   for scientific discussions and critical input on the project, and Drs. Melanie Oakes and

654   Jenny Wu of the UC Irvine Genomics High Throughput Sequencing Facility for next

655   generation sequencing experiments support.

657 Author Contributions

A.M.R., T.D.F., N.E.N. and A.S.I. designed the studies and interpreted the results.

A.M.R., T.D.F., and A.S.I. performed INTACT and TRAP studies. A.M.R. built libraries

for RNA-sequencing. N.E.N. performed CUT&RUN-seq experiments and built libraries

for sequencing. N.E.N. and A.S.I. performed flow cytometry experiments. A.M.R. and

A.S.I. performed immunostainings and immunofluorescent imaging. D.A.V. and A.B.

built exercise cages and monitored running behavior. A.M.R. performed all bioinformatic

analysis of sequencing data. A.M.R. and N.E.N. analyzed sequencing data, performed

GO, GSEA and IPA analyses, and all other downstream analyses except where noted.

A.S.I. performed behavioral studies. T.D.F. performed Spearman's correlations and gene

length by fold change analysis. T.D.F. performed qPCR. N.E.N. designed the figures.

A.M.R., T.D.F., and A.S.I. wrote and edited the manuscript, with N.E.N. providing edits

to the manuscript. All authors discussed the results and edited and approved the

manuscript.

672 Competing Interests

 Methods

 Resource Availability

 Lead Contact

 Further information and requests for resources and reagents should be directed to and will

 be fulfilled by the lead contact, Dr. Autumn Ivy (aivy@uci.edu)

 Materials Availability

 • No new reagents or materials were generated for this study.

 • The mouse lines used for this study are available from Jackson Labs (Emx1-

 IRES-*Cre* knock-in mice, Jackson Laboratory Stock No: 005628, and NuTRAP

 mice, Jackson Laboratory Stock No: 029899).

 Animals

 Emx1-IRES-*Cre* knock-in mice (Jackson Laboratory Stock No: 005628),

 NuTRAP mice (Jackson Laboratory Stock No: 029899), and C57Bl6/J (Jackson

 Laboratory Stock No: 000664) wild-type mice were obtained from Jackson Laboratories.

 Female Emx1-IRES-*Cre* and male NuTRAP mice were crossed to generate the Emx1-

 NuTRAP mice used for this study. Mice were given free access to food and water, and

 lights were maintained on a standard 12-hour light/dark cycle. On postnatal day (P) 21,

 mice were weaned and pair-housed in either standard bedding cages or cages equipped

 with a running wheel. Only male mice were used for the studies performed. Experiments

 were conducted according to US National Institutes of Health guidelines for animal care

 and use and were approved by the Institutional Animal Care and Use Committee of the

 University of California, Irvine.

 Behavior

 Wild-type mice were habituated, handled and then underwent OLM training[90] for

 either three or ten minutes at P41 as described previously[17]. On postnatal day 36, mice

 were brought into a testing room with reduced room brightness. Mice were handled for

 approximately 2 min each, for a total of 5 consecutive days prior to the OLM training

 session (twice a day for the first 2 days followed by once a day for the next 3 days).

Habituation sessions were 5 min, twice per day for 3 days (P39-P41) and occurred within

chambers containing four unique spatial cues on each wall of the chamber (horizontal

lines, black X, vertical strip, and blank wall). Habituation overlapped with the last 3 days

of handling. During the training phase (P42), mice were placed into the same chambers

with two identical objects and exposed to either a subthreshold (3 min) or threshold (10

min) training period. Mice were sacrificed for consolidation experiments 1 hour later.

Exercise Paradigm

After weaning on P21, male mice from the same litters were randomly divided

and pair-housed in either standard cages without a wheel or cages equipped with a

stainless-steel running wheel similar to methods already described[17]. Animals housed

with running wheels had free access to the wheels from P21–41. Real-time data

acquisition (Vital View Software) was used to track distance ran recorded by magnetic

sensors that detect wheel revolutions. Each monitored wheel tracked running distance for

the entire cage (two mice per cage). Revolutions were quantified for every minute daily

for the duration of the exercise period and converted to distance ran per cage (km).

Exercised mice that underwent OLM training were individually tracked as described in

Valientes et al. 2021[91].

"SIT"

On P42 both ELE and sedentary control (n=4/group) mice were sacrificed by rapid

cervical dislocation and both hemispheres of the hippocampus were dissected out and

collected for tissue processing. Hippocampal tissue samples were mechanically

homogenized in 1mL of nuclear preparation buffer (NPB) (10mM HEPES (pH7.5);

1.5mM MgCl2; 10mM KCl; 250mM sucrose; 0.1% IGEPAL CA-630; 0.2mM DTT;

100μg/mL cycloheximide; 1x Complete EDTA-free protease inhibitor (Roche)) by razor

blade. Samples were then triturated with an 18G needle and syringe 50x on ice.

Homogenized tissue samples were pipetted slowly onto 100μm cell strainer caps on

5mL tubes and centrifuged at 1,000rcf for 10 minutes at 4ºC. Cell pellets were

resuspended, and samples were pipetted onto 40μm cell strainer caps on 5mL tubes and

centrifuged at 1,000rcf for 10 minutes at 4ºC. For nuclear isolation, 100μL of

Streptavidin-coated Dynabeads were washed twice with 1mL NPB and resuspended with

100μL NPB. Cell pellets were resuspended and added to washed Streptavidin-coated

Dynabeads. Samples were briefly pipetted to thoroughly suspend the beads and incubated

on ice for 20 minutes. Samples were placed on a magnetic stand to pull down nuclei

bound to the beads. The supernatant from each sample was slowly pipetted off the beads

and transferred to new 1.5mL tubes for RNA isolation (described later). Nuclei-bound

beads were washed twice with 1mL INTACT buffer (10mM HEPES (pH7.5); 1.5mM

MgCl2; 10mM KCl; 250mM sucrose; 0.1% IGEPAL CA-630; 1x Complete EDTA-free

protease inhibitor (Roche)) and resuspended in 100μL INTACT buffer. Bead-bound

nuclei were flash frozen on dry ice and stored at -80ºC until ready for CUT&RUN

chromatin digestion and DNA purification.

For RNA isolation, 100μL of concentrated TRAP isolation buffer (10mM HEPES

(pH 7.5); 117mM MgCl2; 1M KCl; 10% IGEPAL CA-630; 100μg/mL cycloheximide;

11mg/mL sodium heparin; 20mM DTT; 2.2units/μL Rnasin; 1x Complete EDTA-free

protease inhibitor (Roche)) was added to each sample (supernatant above). Samples were

mixed by gentle pipetting and centrifuged at 16,000rcf for 10 minutes at 4ºC. The

supernatant from each sample was transferred to new 1.5mL tubes and 50μL of protein-G

Dynabeads was added to each sample. (To bind anti-GFP, 100μL of protein-G

Dynabeads were washed twice with 500μL NPB and resuspended to a final 100μL

volume. 2μL anti-GFP antibody (Sigma-Aldrich, G6539, Lot:128M4867V) was added to

each tube and incubated for 1 hour on a rotating nutator at 4ºC. Excess anti-GFP was

washed off with 200μL NPB and beads were resuspended in a final 100μL volume.)

Samples were incubated for 3 hours on a rotating nutator at 4ºC. Beads were washed 3x

in 1mL high salt wash buffer (HSWB) (50mM Tris (pH 7.5); 12mM MgCl2; 300mM

KCl; 1% IGEPAL CA-630; 2mM DTT; 100μg/mL cycloheximide) at 4ºC. After the third

wash the beads were slowly resuspended in 350μL RLT buffer (from Qiagen RNeasy

Mini kit #74004) by slowly pipetting up and down and incubated on a rotating nutator for

30 minutes at room temperature. Samples were placed on a magnetic stand to pull down

the beads, and the supernatant from each were transferred to new 1.5mL tubes and stored

at -80ºC until ready for RNA purification.

Separate TRAP RNA Isolation

On P42, mice (n=3 mice/group) were sacrificed and the hippocampus was

dissected out as described above. Either the left or right brain hemisphere of the

hippocampus was processed for TRAP RNA isolation using methods similar to those

previously described[29,92] and the other hemisphere was processed for INTACT nuclear

isolation (see below). Hippocampal tissue was homogenized with a motorized pestle in

1ml of TRAP homogenization buffer (50mM Tris, (pH 7.5); 12mM $MgCl_2$; 100mM KCl;

1% NP-40; 100μg/ml cycloheximide; 1mg/ml sodium heparin; 2mM DTT; 0.2units/μl

RNasin; 1x Complete EDTA-free protease inhibitor (Roche)). Samples were centrifuged

at 16,000rcf for 10 minutes at 4ºC. The supernatant from each sample was collected and

incubated with 50μL of protein G Dynabeads (washed twice in 1mL TRAP

homogenization buffer without RNasin or protease inhibitor and resuspended in 50μL of

the same buffer). Samples were incubated for 30 minutes on a rotating nutator at 4ºC.

Following the incubation, 100μL of anti-GFP bound protein-G Dynabeads was added to

each sample. (To bind anti-GFP, 100μL of protein-G Dynabeads were washed twice with

500μL TRAP homogenization buffer without RNasin or protease inhibitor and

resuspended to a final 100μL volume. 2μL anti-GFP antibody (Sigma-Aldrich, G6539,

Lot:128M4867V) was added to each tube and incubated for 1 hour on a rotating nutator

at 4ºC. Excess anti-GFP was washed off with 200μL TRAP homogenization buffer

without RNasin or protease inhibitor and beads were resuspended in a final 100μL

volume.) Samples were incubated for 3 hours on a rotating nutator at 4ºC. Beads were

washed 3x in 1mL HSWB at 4ºC. After the final wash, the beads were resuspended in

350μL RLT buffer (from Qiagen RNeasy Mini kit #74004) and incubated on a rotating

nutator for 30 minutes at room temperature. Following the incubation, samples were

transferred to new 1.5mL tubes and stored at -80ºC until ready for RNA purification.

Separate INTACT Nuclear Isolation

The remaining hippocampal hemisphere from each mouse (not used for TRAP

RNA isolation above) was processed for INTACT DNA isolation using methods similar

to those described[29,93]. Tissue samples were quickly chopped with a razor blade, added to

1mL of INTACT buffer, and triturated with an 18G needle and syringe 50x on ice.

Samples were then added onto 100µm cell strainer caps on 5mL tubes and centrifuged at

1,000rcf for 10 minutes at 4ºC. Cell pellets were resuspended and samples were pipetted

onto 40µm cell strainer caps and centrifuged at 1,000rcf for 10 minutes at 4ºC. 100µL of

streptavidin coated Dynabeads (washed twice with 1mL INTACT buffer and resuspended

to a final 100 µL volume) were added to each sample, and samples were incubated on ice

for 20 minutes. After incubation, nuclei bound beads were washed twice with 1mL and

resuspended in 100µL INTACT buffer. Bead-bound nuclei were flash frozen on dry ice

and stored at -80ºC until ready for CUT&RUN chromatin digestion and DNA

purification.

RNA Purification and Sequencing Library Preparation

RNA was purified using the Qiagen RNeasy kit according to the manufacturer's

protocol. After purification, RNA quantity and quality were checked using the Qubit

RNA High Sensitivity assay and Agilent Bioanalyzer's Eukaryote Total RNA Pico assay,

respectively, at the University of California, Irvine Genomics High Throughput Facility

(UCI GHTF). Using an input of 15ng of total RNA, mRNA was isolated using

NEXTFLEX Poly(A) Beads 2.0, and the mRNA sequencing libraries were generated

using the NEXTFLEX Rapid Directional RNA-Seq Kit 2.0 according to the

manufacturer's instructions. Final libraries were sent to the UCI GHTF for Qubit dsDNA

High Sensitivity assay and the Agilent Bioanalyzer DNA High Sensitivity assay to

determine the quantity and quality of the RNA-seq libraries, respectively. Libraries were

sequenced at the UCI GHTF using 100bp paired end reads on a single lane of an Illumina

NovaSeq6000 to a minimum sequencing depth of 50 million reads.

Dorsal Hippocampal RNA Isolation

P42 wild-type male mice were sacrificed by cervical dislocation. Left dorsal

hippocampus was extracted by bisecting the isolated hippocampus. The tissue was

homogenized and RNA was extracted using the Qiagen RNeasy kit according to the

manufacturer's protocol. Sequencing libraries were prepared from the RNA as described

under RNA purification and sequencing library preparation above except that ERCC

ExFold RNA Spike-In Mixes (ThermoFisher cat. 4465739) were added according to the
manufacturer's instructions.
830

Cleavage Under Target and Release Using Nuclease (CUT&RUN)

For chromatin analysis, CUT&RUN was performed using the EpiCypher
CUTANA CUT&RUN protocol with slight modifications similar to what has already
been described[37]. Nuclei bound to beads were thawed and pulled down on a magnetic
stand. The supernatant was removed and beads were resuspended in 200μL wash buffer
(20mM HEPES (pH 7.5); 150mM NaCl; 0.5mM spermidine; 1x Complete EDTA-free
protease inhibitor (Roche)). Each sample was divided into three aliquots (50μL for anti-
H3K27me3 (Cell Signaling Technologies, C36B11, Lot: 16), 100μL for anti-H4K8ac
(Epicypher, 13-0047, Lot: 20202001-11), and 50μL for anti-IgG control (Rabbit IgG
Fisher Scientific, 026102)). Beads were pulled down on a magnetic stand, resuspended in
100μL wash buffer and incubated for 10 minutes at room temperature. After incubation
the beads were pulled down, the supernatant was removed, and the beads were
resuspended in 50μL ice cold antibody buffer (20mM HEPES (pH 7.5); 150mM NaCl;
0.5mM spermidine; 0.001mM digitonin; 2mM EDTA; 1x Complete EDTA-free protease
inhibitor (Roche)). 0.5μL of the appropriate antibody was added to each sample, and the
samples were mixed by gentle pipetting prior to overnight incubation at 4ºC on a nutator.

On the second day, the beads were pulled down on a magnetic stand, the
supernatant was removed and the beads were washed twice in 250μL ice cold wash
buffer with 0.001% digitonin. After the washes, beads were resuspended in 50μL ice cold
wash buffer with 0.001% digitonin, and 2.5μL of CUTANA pAG-MNase was added to
each sample and mixed by gentle pipetting. Samples were incubated for 10 minutes at
room temperature and beads were pulled down on a magnetic stand. The supernatant was
removed, and the beads were washed twice in 250μL ice cold wash buffer with 0.001%
digitonin and resuspended in 50μL of the same. 1μL of 100mM $CaCl_2$ was added to each
sample and mixed by gentle pipetting. Samples were incubated for 2 hours on a nutator at
4ºC. Following the incubation, 33μL of stop buffer (340mM NaCl; 20mM EDTA; 4mM
EGTA; 50μg/mL RNase A; 50μg/mL Glycogen) and 1.65μL of Spike-In DNA (Cell
Signaling Technology, #40366) was added to each sample. Samples were incubated in a

preheated thermocycler for 10 minutes at 37ºC. After incubation, samples were

transferred to new 1.5mL tubes and immediately processed for DNA purification.

DNA Purification and Library Preparation

DNA was purified using the Monarch PCR and DNA cleanup kit (New England

Biolabs, 13-0041) according to the manufacturer's instructions. After purification, DNA

fragments were used to generate sequencing libraries using the NEXTFLEX Rapid DNA-

Seq Kit 2.0 according to the manufacturer's instructions. Indices were diluted to a final

concentration of 1:1,000 before being added to samples, and a total of 13 PCR cycles was

used for DNA amplification of the libraries. Final libraries were sent to the UCI GHTF

for Qubit dsDNA High Sensitivity assay and the Agilent Bioanalyzer DNA High

Sensitivity assay to determine the quantity and quality of the CUT&RUN-seq libraries,

respectively. Libraries were sequenced at the UCI GHTF using 100bp paired end reads

on a single lane of an Illumina NovaSeq6000 to a minimum sequencing depth of 10

million reads.

qPCR

cDNA was generated from TRAP-isolated and total input RNA samples (n=2

samples/group) using the Transcriptor First Strand cDNA Synthesis Kit (Roche)

following the manufacturer's instructions. Relative gene expression for *Aqp4* (F: 5'-

ATCCAGCTCGATCTTTTGGA -3', R: 5'-TGAGCTCCACATCAGGACAG -3'),

Tubb3 (F: 5'-GTCTCTAGCCGCGTGAAGTC -3', R: 5'-

GCAGGTCTGAGTCCCCTACA -3'), *Cd11b* (F: 5'-CCCATGACCTTCCAAGAGAA -

3', R: 5'-ACACTGGTAGAGGGCACCTG -3'), and *Mog* (F: 5'-

AAGAGGCAGCAATGGAGTTG -3', R: 5'-GACCTGCAGGAGGATCGTAG -3') was

determined by qPCR using FastStart Essential DNA Green Master Mix (Roche,

06402712001) following the manufacturer's protocol. Cycle counts for mRNA

quantification were normalized to *Gapdh* (F: 5'-CGTCCCGTAGACAAAATGGT -3', R:

5'-GAATTTGCCGTGAGTGGAGT -3'). Quantification was performed using the Pfaffl

method [94].

Immunofluoresence

Emx1-NuTRAP mice were anesthetized intraperitoneally with 50mg/kg of

sodium pentobarbital and transfused transcardially with 4% paraformaldehyde (PFA) in

1x PBS. After perfusion, whole brains were dissected out and incubated in 4% PFA

overnight at 4ºC. After fixation, brains were incubated in 30% sucrose for 72 hours and

mounted in optimal cutting temperature (OCT) compound. 50μm coronal sections were

sliced through the hippocampus using a microtome, and every tenth section was collected

in 1mL of cryoprotectant and stored at -20ºC until ready for immunofluorescent labeling.

For immunofluorescence, sections were brought up to room temperature and washed 2x

in 1x PBS on a nutator for 5 minutes each. Sections were then washed 3x in 1x PBS with

0.3% triton-X. After washes, sections were incubated with mCherry polyclonal antibody

(ThermoFisher, PA5-34974, Lot: UG2804409F) at a 1:200 dilution in blocking solution

(1x PBS with 1% BSA) overnight at 4ºC on a nutator. The following day, sections were

washed 3x in 1x PBS. Sections were incubated for two hours at room temperature with

goat anti-rabbit Alexa Fluor 594 secondary antibody (Invitrogen, A-11012, Lot:

2119134) at a 1:200 dilution in blocking solution. After incubation, the sections were

washed twice in 1x PBS. To label nuclei, sections were incubated with DAPI at a

1:10,000 dilution in 1x PBS for 20 minutes. Sections were then washed 3x in 1x PBS and

cover slipped with Vectatshield®. Fluorescent images were acquired using a Keyence

BZ-X810 All-in-One Fluorescence microscope using the optical sectioning module.

Fluorescence Activated Cell Sorting

Fluorescence activated cell sorting (FACS) was performed to characterize

neuronal and astrocytic NuTRAP cassette expression. Whole hippocampal tissue from

Emx1-NuTRAP mice was isolated and single-cell suspensions were immunostained for

cytometric analysis, as previously described[95]. We used antibodies for THY-1 (OX7)

AlexaFluor™ 647 (Santa Cruz Biotechnology, sc-53116 AF647, Lot: F2716;

concentration: 1:50) and S100β (Abcam, ab41548, Lot:GR3326165-1; concentration:

1:200) with an AlexaFluor™ 405 goat anti-rabbit IgG (H+L) (Invitrogen, A31556, Lot:

2273716; concentration:1:800) secondary antibody. FACS analysis was performed using

a BD FACSAria™ Fusion Flow Cytometer (BD Biosciences) at the University of

California, Irvine Stem Cell Core. Samples and single-stain controls were analyzed using

the FlowJo v10.8.1 software (BD Biosciences). Samples and controls were positively

gated for live cells (SSC-A and FSC-A) and single cells (FSC-H and FSC-A), and

negatively gated for autofluorescence (Comp-Alexa Fluor 647-A (Thy1) and Comp-

BV421-A (S100β)). Fraction of GFP+ neuronal cells was determined using quartile

analysis of Thy-1 and Comp-GFP-A (GFP). Fraction of GFP+ astrocytic cells was

determined using quartile analysis of S100β and Comp-GFP-A (GFP).

Bioinformatics Analysis

Reference assembly: For all sequencing analysis that required it, the primary

reference assembly used was Genome Reference Consortium Mouse Build 38 patch

release 6 (GRCm38.p6).

RNA-seq analysis: FastQ files were quality checked for sequencing errors using

FastQC (version 0.11.9)[96]. No files were found to have sufficient quality errors to

discount their use. Files were aligned using STAR Aligner (version 2.7.3a)[97]. Duplicate

reads were removed using Picard Tools (version 1.87)[98]. SAM Tools was used to convert

BAM files to SAM files for use in downstream analysis. FastQC, alignment and duplicate

removal were all preformed on the High-Powered Compute Cluster (HPC3) operated by

The Research Cyberinfrastructure Center (RCIC) at the University of California, Irvine.

R (version 4.1.0)[99] was used for differentially expressed genes (DEG) analysis. Genomic

Alignments (version 1.28.0)[100] (summarizeOverlaps mode="IntersectionNotEmpty",

singleEnd=FALSE, ignore.strand=FALSE, fragments=TRUE), Genomic Features

(version 1.44.0) (exons by gene), and R SAM Tools (version 2.8.0)[101]

(yieldSize=100000) were used to extract a count matrix and generate a summarized

experiment object. DEseq2 (version 1.32.0)[102] was used to perform a DEG analysis.

Ensemble IDs were converted to gene symbols using BiomaRt (version 2.48.1)[103,104].

PCA plots using the top 2000 genes were generated using ggplot2 (version 3.3.4)[105].

Samples were determined as outliers if the variability between the samples heavily

weighted PC1 to those samples. This excluded 2 sedentary samples and 1 ELE sample.

Heatmaps were generated using gplots (version 3.1.1)[106] and RColorBrewer (version1.1-

2)[107]. Volcano Plots were generated using ggplot2[105]. Venn diagrams were generated

using the Venn diagram tool from Bioinformatics and Evolutionary Genomics[108]. Gene

ontology analysis was performed using Panther Classification System (version 16.0)[44].

Genes were also categorized and a leading edge heatmap was also generated using Gene

Set Enrichment Analysis (version GSEA 4.1.0)[55]. Upstream regulators were identified

from the DEGs upregulated by ELE using Qiagen's Ingenuity Pathway Analysis (IPA)

(Fall 2021 Release)[45].

CUT&RUN-seq analysis: FastQ files were quality checked for sequencing errors

using FastQC (version 0.11.9)[96]. No files were found to have sufficient quality errors to

discount their use. FastQ files were aligned using Bowtie2 (version 2.4.1)[109]. Significant

peaks were called from these aligned files using SEACR, developed in CUT&Tag for

efficient epigenomic profiling of small samples and single cells[61,109], according to

CUT&Tag Data Processing and Analysis Tutorial (updated August 12 2020):

https://www.protocols.io/view/cut-amp-tag-data-processing-and-analysis-tutorial-

bjk2kky. An FDR higher than 0.1 was considered too high for called peaks. This

threshold excluded 4 samples (3 ELE H4K8ac separate isolation samples and 1 ELE

H4K8ac SIT sample). Peaks were called using the stringent and 0.01 settings. Significant

peaks were annotated using ChIPseeker (version 1.8.6)[110] using

TxDb.Mmusculus.UCSC.mm10.knownGene[111] as a reference with the following settings:

tssRegion = c(-3000, 3000), TxDb = TxDb.Mmusculus.UCSC.mm10.knownGene, level

= "transcript", assignGenomicAnnotation = TRUE, genomicAnnotationPriority =

c("Promoter", "5UTR", "3UTR", "Exon", "Intron", "Downstream", "Intergenic"), annoDb

= NULL, addFlankGeneInfo = FALSE, flankDistance = 5000, sameStrand = FALSE,

ignoreOverlap = FALSE, ignoreUpstream = FALSE, ignoreDownstream = FALSE,

overlap = "TSS", verbose = TRUE). This approach annotated peaks to the closest gene by

distance from the promoter, except if that peak falls within a gene before the distal

intergenic region. Ensemble IDs were converted to gene symbols using BiomaRt[103,104].

Gene lists of peaks present in each condition were used to generate Venn diagrams, as

with RNA-seq analysis[108]. Further Venn diagrams were generated comparing DEGs with

peak calls. Peaks were visualized using UCSC Genome Browser and Genome Browser in

a Box[112-114].

Normalized counts (for read depth) at binned sections of the DNA as well as

RNA-seq gene expression count (normalized for read depth) were compared between

simultaneous and separate isolations as well as left and right hemispheres using

Spearman's correlations in R with the package ggpubr (version 0.4.0)[115]. Bar plots were

generated using Graph Pad 9 PRISM.

Data Availability

RNA-seq and CUT&RUN-seq data will be deposited and publicly available as of

the date of publication. Accession numbers will be listed here pending publication. Any

additional information required to reanalyze the data reported in this paper is available

from the lead contact upon request.

		References

1. Bedrosian, T. A., Quayle, C., Novaresi, N. & Gage, F. H. Early life experience drives structural variation of neural genomes in mice. *Science (New York, N.Y.)* **359**, 1395-1399, doi:10.1126/science.aah3378 (2018).

2. Stroud, H. *et al.* Early-Life Gene Expression in Neurons Modulates Lasting Epigenetic States. *Cell* **171**, 1151-1164.e1116, doi:https://doi.org/10.1016/j.cell.2017.09.047 (2017).

3. Herre, M. & Korb, E. The chromatin landscape of neuronal plasticity. *Curr Opin Neurobiol* **59**, 79-86, doi:10.1016/j.conb.2019.04.006 (2019).

4. Campbell, R. R. & Wood, M. A. How the epigenome integrates information and reshapes the synapse. *Nat Rev Neurosci* **20**, 133-147, doi:10.1038/s41583-019-0121-9 (2019).

5. Kronman, H. *et al.* Long-term behavioral and cell-type-specific molecular effects of early life stress are mediated by H3K79me2 dynamics in medium spiny neurons. *Nat Neurosci* **24**, 667-676, doi:10.1038/s41593-021-00814-8 (2021).

6. Sagarkar, S. *et al.* LSD1-BDNF activity in lateral hypothalamus-medial forebrain bundle area is essential for reward seeking behavior. *Progress in neurobiology* **202**, 102048, doi:10.1016/j.pneurobio.2021.102048 (2021).

7. Webb, W. M. *et al.* The SETD6 Methyltransferase Plays an Essential Role in Hippocampus-Dependent Memory Formation. *Biological psychiatry* **87**, 577-587, doi:10.1016/j.biopsych.2019.05.022 (2020).

8. Bolton, J. L. *et al.* Unexpected Transcriptional Programs Contribute to Hippocampal Memory Deficits and Neuronal Stunting after Early-Life Adversity. *Cell Rep* **33**, 108511, doi:10.1016/j.celrep.2020.108511 (2020).

9. Campbell, R. R. *et al.* HDAC3 Activity within the Nucleus Accumbens Regulates Cocaine-Induced Plasticity and Behavior in a Cell-Type-Specific Manner. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **41**, 2814-2827, doi:10.1523/JNEUROSCI.2829-20.2021 (2021).

10. Lister, R. *et al.* Global Epigenomic Reconfiguration During Mammalian Brain Development. *Science* **341**, 1237905, doi:10.1126/science.1237905 (2013).

11. Lavery, L. A. & Zoghbi, H. Y. The distinct methylation landscape of maturing neurons and its role in Rett syndrome pathogenesis. *Current opinion in neurobiology* **59**, 180-188, doi:10.1016/j.conb.2019.08.001 (2019).

12. Aguilar, R., Bustos, F. J., Nardocci, G., van Zundert, B. & Montecino, M. Epigenetic silencing of the osteoblast-lineage gene program during hippocampal maturation. *Journal of cellular biochemistry* **122**, 367-384, doi:10.1002/jcb.29865 (2021).

13. West, A. E. & Greenberg, M. E. Neuronal activity-regulated gene transcription in synapse development and cognitive function. *Cold Spring Harbor perspectives in biology* **3**, doi:10.1101/cshperspect.a005744 (2011).

14. Bale, T. L. *et al.* Early life programming and neurodevelopmental disorders. *Biological psychiatry* **68**, 314-319, doi:10.1016/j.biopsych.2010.05.028 (2010).

15   Cotman, C. W., Berchtold, N. C. & Christie, L.-A. Exercise builds brain health:
key roles of growth factor cascades and inflammation. *Trends in*
Neurosciences **30**, 464-472, doi:https://doi.org/10.1016/j.tins.2007.06.011
(2007).
16   Voss, M. W., Vivar, C., Kramer, A. F. & van Praag, H. Bridging animal and
human models of exercise-induced brain plasticity. *Trends in cognitive*
sciences **17**, 525-544, doi:10.1016/j.tics.2013.08.001 (2013).
17   Ivy, A. S. *et al.* A Unique Mouse Model of Early Life Exercise Enables
Hippocampal Memory and Synaptic Plasticity. *Scientific Reports* **10**, 9174,
doi:10.1038/s41598-020-66116-4 (2020).
18   Berchtold, N. C., Castello, N. & Cotman, C. W. Exercise and time-dependent
benefits to learning and memory. *Neuroscience* **167**, 588-597,
doi:10.1016/j.neuroscience.2010.02.050 (2010).
19   Urdinguio, R. G. *et al.* Physical exercise shapes the mouse brain epigenome.
Mol Metab **54**, 101398, doi:10.1016/j.molmet.2021.101398 (2021).
20   Abel, J. L. & Rissman, E. F. Running-induced epigenetic and gene expression
changes in the adolescent brain. *International journal of developmental*
neuroscience : the official journal of the International Society for
Developmental Neuroscience **31**, 382-390,
doi:10.1016/j.ijdevneu.2012.11.002 (2013).
21   Ieraci, A., Mallei, A., Musazzi, L. & Popoli, M. Physical exercise and acute
restraint stress differentially modulate hippocampal brain-derived
neurotrophic factor transcripts and epigenetic mechanisms in mice.
Hippocampus **25**, 1380-1392, doi:10.1002/hipo.22458 (2015).
22   Intlekofer, K. A. *et al.* Exercise and Sodium Butyrate Transform a
Subthreshold Learning Event into Long-Term Memory via a Brain-Derived
Neurotrophic factor-Dependent Mechanism. *Neuropsychopharmacology* **38**,
2027-2034, doi:10.1038/npp.2013.104 (2013).
23   Butler, C. W. *et al.* Exercise opens a temporal window for enhanced cognitive
improvement from subsequent physical activity. *Learn Mem* **26**, 485-492,
doi:10.1101/lm.050278.119 (2019).
24   Titterness, A. K., Wiebe, E., Kwasnica, A., Keyes, G. & Christie, B. R. Voluntary
exercise does not enhance long-term potentiation in the adolescent female
dentate gyrus. *Neuroscience* **183**, 25-31,
doi:10.1016/j.neuroscience.2011.03.050 (2011).
25   Gomes da Silva, S. & Arida, R. M. Physical activity and brain development.
Expert review of neurotherapeutics **15**, 1041-1051,
doi:10.1586/14737175.2015.1077115 (2015).
26   Gomes da Silva, S. *et al.* Early exercise promotes positive hippocampal
plasticity and improves spatial memory in the adult life of rats. *Hippocampus*
22, 347-358, doi:10.1002/hipo.20903 (2012).
27   Fernandes, J., Arida, R. M. & Gomez-Pinilla, F. Physical exercise as an
epigenetic modulator of brain plasticity and cognition. *Neuroscience and*
biobehavioral reviews **80**, 443-456, doi:10.1016/j.neubiorev.2017.06.012
(2017).

28   Intlekofer, K. A. *et al.* Exercise and sodium butyrate transform a subthreshold
1084   learning event into long-term memory via a brain-derived neurotrophic
1085   factor-dependent mechanism. *Neuropsychopharmacology* **38**, 2027-2034,
1086   doi:10.1038/npp.2013.104 (2013).
1087   29   Roh, H. C. *et al.* Simultaneous Transcriptional and Epigenomic Profiling from
1088   Specific Cell Types within Heterogeneous Tissues In Vivo. *Cell Rep* **18**, 1048-
1089   1061, doi:10.1016/j.celrep.2016.12.087 (2017).
1090   30   Deal, R. B. & Henikoff, S. The INTACT method for cell type-specific gene
1091   expression and chromatin profiling in Arabidopsis thaliana. *Nature protocols*
1092   **6**, 56-68, doi:10.1038/nprot.2010.175 (2011).
1093   31   Heiman, M. *et al.* A translational profiling approach for the molecular
1094   characterization of CNS cell types. *Cell* **135**, 738-748,
1095   doi:10.1016/j.cell.2008.10.028 (2008).
1096   32   Gorski, J. A. *et al.* Cortical excitatory neurons and glia, but not GABAergic
1097   neurons, are produced in the Emx1-expressing lineage. *J Neurosci* **22**, 6309-
1098   6314, doi:10.1523/jneurosci.22-15-06309.2002 (2002).
1099   33   Chan, C. H. *et al.* Emx1 is a marker for pyramidal neurons of the cerebral
1100   cortex. *Cerebral cortex (New York, N.Y. : 1991)* **11**, 1191-1198,
1101   doi:10.1093/cercor/11.12.1191 (2001).
1102   34   Hong, S. M. *et al.* Reduced hippocampal neurogenesis and skill reaching
1103   performance in adult Emx1 mutant mice. *Experimental neurology* **206**, 24-32,
1104   doi:10.1016/j.expneurol.2007.03.028 (2007).
1105   35   Willaime-Morawek, S. & van der Kooy, D. Cortex- and striatum- derived
1106   neural stem cells produce distinct progeny in the olfactory bulb and striatum.
1107   *The European journal of neuroscience* **27**, 2354-2362, doi:10.1111/j.1460-
1108   9568.2008.06206.x (2008).
1109   36   Luo, L. *et al.* Optimizing Nervous System-Specific Gene Targeting with Cre
1110   Driver Lines: Prevalence of Germline Recombination and Influencing Factors.
1111   *Neuron* **106**, 37-65.e35, doi:10.1016/j.neuron.2020.01.008 (2020).
1112   37   Skene, P. J., Henikoff, J. G. & Henikoff, S. Targeted in situ genome-wide
1113   profiling with high efficiency for low cell numbers. *Nature protocols* **13**,
1114   1006-1019, doi:10.1038/nprot.2018.015 (2018).
1115   38   Liu, Y. *et al.* Glial fibrillary acidic protein-expressing neural progenitors give
1116   rise to immature neurons via early intermediate progenitors expressing both
1117   glial fibrillary acidic protein and neuronal markers in the adult hippocampus.
1118   *Neuroscience* **166**, 241-251, doi:10.1016/j.neuroscience.2009.12.026 (2010).
1119   39   Garcia, A. D. R., Doan, N. B., Imura, T., Bush, T. G. & Sofroniew, M. V. GFAP-
1120   expressing progenitors are the principal source of constitutive neurogenesis
1121   in adult mouse forebrain. *Nature Neuroscience* **7**, 1233-1241,
1122   doi:10.1038/nn1340 (2004).
1123   40   Chucair-Elliott, A. J. *et al.* Inducible cell-specific mouse models for paired
1124   epigenetic and transcriptomic studies of microglia and astroglia. *bioRxiv*,
1125   862722, doi:10.1101/862722 (2019).
1126   41   Jordan, J. T. The rodent hippocampus as a bilateral structure: A review of
1127   hemispheric lateralization. *Hippocampus* **30**, 278-292,
1128   doi:https://doi.org/10.1002/hipo.23188 (2020).

42 Berchtold, N. C., Chinn, G., Chou, M., Kesslak, J. P. & Cotman, C. W. Exercise
primes a molecular memory for brain-derived neurotrophic factor protein
induction in the rat hippocampus. *Neuroscience* **133**, 853-861,
doi:10.1016/j.neuroscience.2005.03.026 (2005).
43 McNulty, S. E. *et al.* Differential roles for Nr4a1 and Nr4a2 in object location
vs. object recognition long-term memory. *Learning & memory (Cold Spring*
Harbor, N.Y.) **19**, 588-592, doi:10.1101/lm.026385.112 (2012).
44 Mi, H. & Thomas, P.   123-140 (Humana Press, 2009).
45 Krämer, A., Green, J., Pollard, J. & Tugendreich, S. Causal analysis approaches
in Ingenuity Pathway Analysis. *Bioinformatics* **30**, 523-530,
doi:10.1093/bioinformatics/btt703 (2014).
46 Kandel, E. R. The molecular biology of memory: cAMP, PKA, CRE, CREB-1,
CREB-2, and CPEB. *Molecular brain* **5**, 14, doi:10.1186/1756-6606-5-14
(2012).
47 Kerimoglu, C. *et al.* KMT2A and KMT2B Mediate Memory Function by
Affecting Distinct Genomic Regions. *Cell reports* **20**, 538-548,
doi:10.1016/j.celrep.2017.06.072 (2017).
48 Ashbrook, D. G. *et al.* Transcript co-variance with Nestin in two mouse
genetic reference populations identifies Lef1 as a novel candidate regulator
of neural precursor cell proliferation in the adult hippocampus. *Front*
Neurosci **8**, 418, doi:10.3389/fnins.2014.00418 (2014).
49 Tapia-González, S., Muñoz, M., Cuartero, M. & Sánchez-Capelo, A. Smad3 is
required for the survival of proliferative intermediate progenitor cells in the
dentate gyrus of adult mice. *Cell Communication and Signaling* **11**, 93,
doi:10.1186/1478-811x-11-93 (2013).
50 Tiano, J. P., Springer, D. A. & Rane, S. G. SMAD3 Negatively Regulates Serum
Irisin and Skeletal Muscle FNDC5 and Peroxisome Proliferator-activated
Receptor γ Coactivator 1-α (PGC-1α) during Exercise. *Journal of Biological*
Chemistry **290**, 7671-7684, doi:10.1074/jbc.m114.617399 (2015).
51 Sun, Z. *et al.* EGR1 recruits TET1 to shape the brain methylome during
development and upon neuronal activity. *Nature Communications* **10**,
doi:10.1038/s41467-019-11905-3 (2019).
52 Bouché, V. *et al.* Drosophila Mitf regulates the V-ATPase and the lysosomal-
autophagic pathway. *Autophagy* **12**, 484-498,
doi:10.1080/15548627.2015.1134081 (2016).
53 Guan, Z. *et al.* Integration of long-term-memory-related synaptic plasticity
involves bidirectional regulation of gene expression and chromatin structure.
Cell **111**, 483-493, doi:10.1016/s0092-8674(02)01074-7 (2002).
54 Korzus, E., Rosenfeld, M. G. & Mayford, M. CBP Histone Acetyltransferase
Activity Is a Critical Component of Memory Consolidation. *Neuron* **42**, 961-
972, doi:https://doi.org/10.1016/j.neuron.2004.06.002 (2004).
55 Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based
approach for interpreting genome-wide expression profiles. *Proceedings of*
the National Academy of Sciences **102**, 15545-15550,
doi:10.1073/pnas.0506580102 (2005).

56 Nofrini, V., Di Giacomo, D. & Mecucci, C. Nucleoporin genes in human
diseases. *European Journal of Human Genetics* **24**, 1388-1395,
doi:10.1038/ejhg.2016.25 (2016).
57 Gomes-Duarte, A., Lacerda, R., Menezes, J. & Romão, L. eIF3: a factor for
human health and disease. *RNA Biol* **15**, 26-34,
doi:10.1080/15476286.2017.1391437 (2018).
58 Alvarez-López, M. J. *et al.* Long-term exercise modulates hippocampal gene
expression in senescent female mice. *J Alzheimers Dis* **33**, 1177-1190,
doi:10.3233/jad-121264 (2013).
59 Butler, A. A., Sanchez, R. G., Jarome, T. J., Webb, W. M. & Lubin, F. D. O-GlcNAc
and EZH2-mediated epigenetic regulation of gene expression during
consolidation of fear memories. *Learning & memory (Cold Spring Harbor,*
N.Y.) **26**, 373-379, doi:10.1101/lm.049023.118 (2019).
60 Butler, A. A., Sanchez, R. G., Jarome, T. J., Webb, W. M. & Lubin, F. D. O-GlcNAc
and EZH2-mediated epigenetic regulation of gene expression during
consolidation of fear memories. *Learning & memory (Cold Spring Harbor,*
N.Y.) **26**, 373-379, doi:10.1101/lm.049023.118 (2019).
61 Kaya-Okur, H. S. *et al.* CUT&Tag for efficient epigenomic profiling of small
samples and single cells. *Nature Communications* **10**, doi:10.1038/s41467-
019-09982-5 (2019).
62 Stergiopoulos, A., Elkouris, M. & Politis, P. K. Prospero-related homeobox 1
(Prox1) at the crossroads of diverse pathways during adult neural fate
specification. *Front Cell Neurosci* **8**, 454, doi:10.3389/fncel.2014.00454
(2014).
63 Camp, N. D. *et al.* Individual protomers of a G protein-coupled receptor dimer
integrate distinct functional modules. *Cell Discov* **1**, 15011-,
doi:10.1038/celldisc.2015.11 (2015).
64 Baple, E. L. *et al.* Mutations in KPTN cause macrocephaly,
neurodevelopmental delay, and seizures. *Am J Hum Genet* **94**, 87-94,
doi:10.1016/j.ajhg.2013.10.001 (2014).
65 Peterson, M. D. & Titus, M. A. F-actin distribution of Dictyostelium myosin I
double mutants. *J Eukaryot Microbiol* **41**, 652-657, doi:10.1111/j.1550-
7408.1994.tb01529.x (1994).
66 Johnson Chacko, L. *et al.* Transcriptome-Wide Analysis Reveals a Role for
Extracellular Matrix and Integrin Receptor Genes in Otic Neurosensory
Differentiation from Human iPSCs. *Int J Mol Sci* **22**,
doi:10.3390/ijms221910849 (2021).
67 Truter, S., Andrikopoulos, K., Di Liberto, M., Womack, L. & Ramirez, F. Pro-
alpha 2(V) collagen gene; pairwise analysis of the amino-propeptide coding
domain, and cross-species comparison of the promoter sequence. *Connect*
Tissue Res **29**, 51-59, doi:10.3109/03008209309061966 (1993).
68 Lévy, J. *et al.* EFNB2 haploinsufficiency causes a syndromic
neurodevelopmental disorder. *Clin Genet* **93**, 1141-1147,
doi:10.1111/cge.13234 (2018).

69   Xiong, C. *et al.* Targeting Forward and Reverse EphB4/EFNB2 Signaling by a
Peptide with Dual Functions. *Sci Rep* **10**, 520, doi:10.1038/s41598-020-
57477-x (2020).
70   Zhu, F. *et al.* EFNB2 facilitates cell proliferation, migration, and invasion in
pancreatic ductal adenocarcinoma via the p53/p21 pathway and EMT.
Biomed Pharmacother **125**, 109972, doi:10.1016/j.biopha.2020.109972
(2020).
71   Ferrari, K. J., Lavarone, E. & Pasini, D. The Dual Role of EPOP and Elongin BC
in Controlling Transcriptional Activity. *Mol Cell* **64**, 637-638,
doi:10.1016/j.molcel.2016.11.009 (2016).
72   Liefke, R., Karwacki-Neisius, V. & Shi, Y. EPOP Interacts with Elongin BC and
USP7 to Modulate the Chromatin Landscape. *Mol Cell* **64**, 659-672,
doi:10.1016/j.molcel.2016.10.019 (2016).
73   Warburton, E. C. *et al.* cAMP responsive element-binding protein
phosphorylation is necessary for perirhinal long-term potentiation and
recognition memory. *J Neurosci* **25**, 6296-6303, doi:10.1523/jneurosci.0506-
05.2005 (2005).
74   Logan, T., Bendor, J., Toupin, C., Thorn, K. & Edwards, R. H. α-Synuclein
promotes dilation of the exocytotic fusion pore. *Nat Neurosci* **20**, 681-689,
doi:10.1038/nn.4529 (2017).
75   Fournier, A. E., GrandPre, T. & Strittmatter, S. M. Identification of a receptor
mediating Nogo-66 inhibition of axonal regeneration. *Nature* **409**, 341-346,
doi:10.1038/35053072 (2001).
76   Pérez-Otaño, I., Larsen, R. S. & Wesseling, J. F. Emerging roles of GluN3-
containing NMDA receptors in the CNS. *Nat Rev Neurosci* **17**, 623-635,
doi:10.1038/nrn.2016.92 (2016).
77   Chodelkova, O., Masek, J., Korinek, V., Kozmik, Z. & Machon, O. Tcf7L2 is
essential for neurogenesis in the developing mouse neocortex. *Neural Dev* **13**,
8, doi:10.1186/s13064-018-0107-8 (2018).
78   Iwano, T., Masuda, A., Kiyonari, H., Enomoto, H. & Matsuzaki, F. Prox1
postmitotically defines dentate gyrus cells by specifying granule cell identity
over CA3 pyramidal cell fate in the hippocampus. *Development (Cambridge,*
England) **139**, 3051-3062, doi:10.1242/dev.080002 (2012).
79   Luo, C. *et al.* Single-cell methylomes identify neuronal subtypes and
regulatory elements in mammalian cortex. *Science* **357**, 600-604,
doi:10.1126/science.aan3351 (2017).
80   Skene, P. J., Henikoff, J. G. & Henikoff, S. Targeted in situ genome-wide
profiling with high efficiency for low cell numbers. *Nature protocols* **13**,
1006-1019, doi:10.1038/nprot.2018.015 (2018).
81   Helbling, J.-C., Kinouchi, K., Trifilieff, P., Sassone-Corsi, P. & Moisan, M.-P.
Combined Gene Expression and Chromatin Immunoprecipitation From a
Single Mouse Hippocampus. *Current Protocols* **1**, e33,
doi:https://doi.org/10.1002/cpz1.33 (2021).
82   Chucair-Elliott, A. J. *et al.* Inducible cell-specific mouse models for paired
epigenetic and transcriptomic studies of microglia and astroglia. *Commun*
Biol **3**, 693, doi:10.1038/s42003-020-01418-x (2020).

83   Doty, R. W., Fei, R., Hu, S. & Kavcic, V. Long-term reversal of hemispheric
specialization for visual memory in a split-brain macaque. *Behavioural brain*
research **102**, 99-113, doi:10.1016/s0166-4328(99)00006-6 (1999).
84   Duboc, V., Dufourcq, P., Blader, P. & Roussigné, M. Asymmetry of the Brain:
Development and Implications. *Annual review of genetics* **49**, 647-672,
doi:10.1146/annurev-genet-112414-055322 (2015).
85   Güntürkün, O., Ströckens, F. & Ocklenburg, S. Brain Lateralization: A
Comparative Perspective. *Physiological reviews* **100**, 1019-1063,
doi:10.1152/physrev.00006.2019 (2020).
86   Hubel, D. H. & Wiesel, T. N. Receptive fields, binocular interaction and
functional architecture in the cat's visual cortex. *The Journal of Physiology*
160, 106-154, doi:https://doi.org/10.1113/jphysiol.1962.sp006837 (1962).
87   Spiers, H. J. *et al.* Unilateral temporal lobectomy patients show lateralized
topographical and episodic memory deficits in a virtual town. *Brain : a*
journal of neurology **124**, 2476-2489, doi:10.1093/brain/124.12.2476
(2001).
88   Lynch, M. A. Long-term potentiation and memory. *Physiological reviews* **84**,
87-136, doi:10.1152/physrev.00014.2003 (2004).
89   Achilly, N. P., Wang, W. & Zoghbi, H. Y. Presymptomatic training mitigates
functional deficits in a mouse model of Rett syndrome. *Nature* **592**, 596-600,
doi:10.1038/s41586-021-03369-7 (2021).
90   Kwapis, J. L. *et al.* Epigenetic regulation of the circadian gene Per1
contributes to age-related changes in hippocampal memory. *Nat Commun* **9**,
3323, doi:10.1038/s41467-018-05868-0 (2018).
91   David A Valientes, A. M. R., Autumn S Ivy. An Improved Method for Individual
Tracking of Voluntary Wheel Running in Pair-housed Juvenile Mice. *BIO-*
PROTOCOL **11**, doi:10.21769/bioprotoc.4071 (2021).
92   Long, J. Z. *et al.* A smooth muscle-like origin for beige adipocytes. *Cell*
metabolism **19**, 810-820, doi:10.1016/j.cmet.2014.03.025 (2014).
93   Deal, R. B. & Henikoff, S. A simple method for gene expression and chromatin
profiling of individual cell types within a tissue. *Developmental cell* **18**, 1030-
1040, doi:10.1016/j.devcel.2010.05.013 (2010).
94   Pfaffl, M. W. A new mathematical model for relative quantification in real-
time RT-PCR. *Nucleic Acids Res* **29**, e45, doi:10.1093/nar/29.9.e45 (2001).
95   Schwarz, J. M. Using fluorescence activated cell sorting to examine cell-type-
specific gene expression in rat brain tissue. *Journal of visualized experiments :*
JoVE, e52537, doi:10.3791/52537 (2015).
96   FastQC v. 0.11.9 (Babraham Bioinformatics, 2005).
97   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,
15-21, doi:10.1093/bioinformatics/bts635 (2013).
98   Picard v. 1.87 (Broad Institute, 2013).
99   R (R Foundation for statistical Computing, Vienna, Austria, 2010).
100 Lawrence, M. *et al.* Software for Computing and Annotating Genomic Ranges.
PLoS Computational Biology **9**, e1003118, doi:10.1371/journal.pcbi.1003118
(2013).

101 Rsamtools: Binary alignment (BAM), FASTA, variant call (BCF), and tabix file
import (Bioconductor Bioconductor 2021).
102 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**,
doi:10.1186/s13059-014-0550-8 (2014).
103 Durinck, S. *et al.* BioMart and Bioconductor: a powerful link between
biological databases and microarray data analysis. *Bioinformatics* **21**, 3439-
3440, doi:10.1093/bioinformatics/bti525 (2005).
104 Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt.
Nature protocols **4**, 1184-1191, doi:10.1038/nprot.2009.97 (2009).
105 Ggplot2: Elegant graphics for data analysis v. 3.3.4 (Springer International
Publishing, Cham, Switzerland, 2016).
106 gplots: Various R Programming Tools for Plotting Data v. 3.1.1 (Tal Galili
2020).
107 RColorBrewer: ColorBrewer Palettes (Erich Neuwirth, CRAN, 2014).
108 Venn Diagram: Calculate and draw custom Venn diagrams (BELGIUM, 2021).
109 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2.
Nature Methods **9**, 357-359, doi:10.1038/nmeth.1923 (2012).
110 Humburg, P., Helliwell, C. A., Bulger, D. & Stone, G. ChIPseqR: analysis of ChIP-
seq experiments. *BMC Bioinformatics* **12**, 39, doi:10.1186/1471-2105-12-39
(2011).
111 TxDb.Mmusculus.UCSC.mm10.knownGene: Annotation package for TxDb
object(s) v. 3.4.7 (Bioconductor 2019).
112 Kent, W. J. *et al.* The Human Genome Browser at UCSC. *Genome Research* **12**,
996-1006, doi:10.1101/gr.229102 (2002).
113 Rosenbloom, K. R. *et al.* ENCODE whole-genome data in the UCSC Genome
Browser. *Nucleic Acids Research* **38**, D620-D625, doi:10.1093/nar/gkp961
(2010).
114 Haeussler, M. *et al.* Navigating protected genomics data with UCSC Genome
Browser in a Box. *Bioinformatics* **31**, 764-766,
doi:10.1093/bioinformatics/btu712 (2014).
115 ggpubr: 'ggplot2' Based Publication Ready Plots (CRAN, 2020).
1342

Supplemental Figure 1

A
INTACT Nucleus

B
Relative Expression to GAPDH

Total RNA
TRAP RNA

TUBB3   AQP4   MOG   CD11b

C   Daily Running Distances by Cage During ELE

Running Distance (km)

Postnatal Day (P)

Separate Isolation Group
▼ Pair 1
◆ Pair 2
▲ Pair 3

Simultaneous Isolation Group
● Pair 4
■ Pair 5

Supplemental Figure 2

Supplemental Figure 3