1   **Early life exercise primes the neural epigenome to facilitate gene expression and**
2   **hippocampal memory consolidation.**
3
4

Raus AM*[1], Fuller TD*[2], Nelson NE*[1], Valientes DA[2], Bayat A[2], Ivy AS[1,2,3,4]
[1]Physiology/Biophysics, Anatomy/Neurobiology, University of California- Irvine School
of Medicine, Irvine, CA
[2]Pediatrics, University of California- Irvine School of Medicine, Irvine, CA
[3]Neurobiology and Behavior, University of California- Irvine School of Biological
Sciences, Irvine, CA
[4]Division of Neurology, Children's Hospital Orange County, Orange, CA
12
13   Correspondence to aivy@hs.uci.edu
14
15   *These authors contributed equally to the manuscript.

16   **Abstract**

Aerobic exercise is well known to promote neuroplasticity and hippocampal memory. In

the developing brain, early-life exercise (ELE) can lead to lasting improvements in

hippocampal function, yet molecular mechanisms underlying this phenomenon have not

been fully explored. In this study, adolescent transgenic mice harboring the "NuTRAP"

(Nuclear tagging and Translating Ribosome Affinity Purification) cassette in Emx1

expressing neurons ("Emx1-NuTRAP" mice) undergo ELE followed by a hippocampal

learning task, in order to determine the molecular underpinnings of exercise contributing

to improved hippocampal memory performance. We simultaneously isolate and sequence

translating mRNA and nuclear chromatin from a single hippocampus in a cell-type

specific manner (excitatory neurons), demonstrate validity of our new technical approach,

and couple multi-omics sequencing data to evaluate histone modifications H4K8ac and

H3K27me3 and their influence on gene expression after ELE. We then evaluate new gene

expression – histone modification relationships specifically during hippocampal memory

consolidation that may play a critical role in facilitated memory after ELE. Our data

reveal novel candidate gene-histone modification interactions and implicate gene

regulatory pathways involved in ELE's impact on hippocampal learning and memory.

Introduction

Environmental experiences engage epigenetic mechanisms to modulate gene expression and cell function in post-mitotic neurons[1,2]. Histone modifications and DNA methylation are particularly important for neuronal adaptation to environmental signals by altering transcription and synaptic function[3,4]. Behavioral outputs such as stress susceptibility, reward seeking, and long-term memory have been shown to result from changes to chromatin accessibility and gene expression in neurons[5-9]. In addition to this, the neuronal chromatin landscape undergoes waves of epigenetic modifications as a function of brain maturation itself[10-12]. Postnatal periods of heightened sensitivity to environmental stimuli can lead to lasting changes to cellular function and may result from temporally specific epigenetic mechanisms in the developing brain[13,14]. Whether gene regulatory mechanisms in postmitotic neurons are uniquely influenced by early-life experiences to inform long-term function is a question that is just beginning to be explored. Identifying epigenetic processes involved in modulating cell function, particularly during brain development, is critical for understanding how early-life experiences impact long-term behavioral outcomes.

Aerobic exercise enhances performance on cognitive tasks involving the hippocampus in both adult humans and animal models[15,16]. The type, timing, and duration of exercise exposure matters with regard to whether it has a persistent impact on hippocampal function[17-19]. Findings in both adolescent and adult rodents implicate a role for histone modifying enzymes in the mechanisms of exercise-induced benefits to hippocampal memory. Both voluntary exercise or treatment with a HDAC3 inhibitor enable hippocampal memory after a subthreshold learning stimulus, increase brain-derived neurotrophic factor (BDNF), and promote acetylation of H4K8 at the BDNF promoter[20-22]. This suggests that exercise engages epigenetic regulatory mechanisms to promote plasticity. Adult exercise also opens a temporal window for persistent improvements to memory performance when a reactivating exercise exposure is introduced[23], suggesting a "molecular memory" of the initial exercise[23]. Although the majority of these studies have been performed in adults, more recent work also demonstrates the effects of exercise on hippocampal memory and associated alterations in neurotrophic factor expression, synaptic plasticity, and neurogenesis are similar in

juvenile and adolescent periods[17,24-27]. Previous work in our lab showed that early-life

exercise (ELE) for either one week (juvenile period; postnatal days (P) 21-27) or three

weeks (juvenile-adolescence; P21-41) facilitated hippocampal long-term memory

formation in response to a learning stimulus typically insufficient for forming long-term

memory. This finding was associated with increased long-term potentiation (LTP) as well

as modulations to synaptic physiology in hippocampal CA1[17]. Notably, the hippocampal

memory effects of juvenile ELE persisted two weeks after exercise cessation, which is

potentially longer than the effect of exercise on adult hippocampal function[23]. Taking

these findings together, it is possible that exercise (whether in early life or adulthood)

may "prime" hippocampal function for response to future experiences (such as future

exercise bouts or hippocampal learning events). Epigenetic mechanisms are strong

candidates for the priming effects of exercise: the epigenome could harbor a "molecular

memory" of the exercise experience by readying the chromatin landscape for efficient

gene expression, thereby modulating neuronal function and behavioral output[22]. The

specific mechanisms underlying sustained electrophysiological and behavioral effects

observed in our studies of ELE have not been assessed from the perspective of a potential

molecular memory of ELE within the epigenome.

In this study we describe a new approach to simultaneously isolate neuron-

specific chromatin and translating mRNA from a single hippocampal homogenate, in

order to address our hypothesis that ELE engages the epigenome to contribute to

improved hippocampal memory and synaptic plasticity. We identify unique neuronal

gene expression programs resulting from ELE and pair this data with ELE-induced

changes in H4K8ac or H3K27me3. By crossing an Emx1-Cre transgenic line with mice

harboring the NuTRAP (Nuclear tagging and Translating Ribosome Affinity

Purification[28]) cassette (termed "Emx1-NuTRAP" mice), we isolate whole nuclei and

translating mRNA from a single hippocampus by combining the first steps of the

INTACT (Isolation of Nuclei TAgged in specific Cell Types[29]) and TRAP (Translating

Ribosome Affinity Purification[30]) published protocols. We have named our technical

approach "SIT" (Simultaneous INTACT & TRAP). This technique yields DNA and RNA

from the same homogenate that is of high-quality and sufficient concentrations for

downstream sequencing procedures in this study, TRAP-seq and CUT&RUN-seq. Using

neuron-specific, paired transcriptomic and epigenomic bulk sequencing, this study

uncovers hippocampal gene expression programs and associated histone modifications

that are driven by ELE to promote plasticity during hippocampal memory consolidation.

Importantly, these data reveal an epigenetic signature resulting from ELE that may form

the basis of persistently enabled hippocampal function.

102   **Results**

103   **"Emx1-NuTRAP" mouse allows for simultaneous isolation of nuclear chromatin**

104   **and translating mRNA from a single population of hippocampal neurons.**

105   Our initial goal was to obtain translating mRNA and nuclear DNA from a single

population of neurons to couple transcriptional network changes with alterations in

histone modifications resulting from ELE. To do this, we developed a transgenic mouse

line (Emx1-NuTRAP) and experimental protocol (SIT protocol). We crossed male

NuTRAP reporter mice[28] with female Emx1-Cre[31,32] mouse lines to generate Emx1-Cre;

NuTRAP transgenic progeny for this study ("Emx1-NuTRAP"; Fig. 1A). Emx1-

expressing neurons are predominately excitatory[31] and are necessary for hippocampal

neurogenesis and motor skill learning[33]. In the presence of a *loxP* site-flanked sequence,

Emx1-Cre facilitates recombination in approximately 88% of neurons in the neocortex

and hippocampus, and in less than 2% of GABAergic inhibitory interneurons in these

regions[31]. Emx1-Cre also targets neural progenitor cells and can be expressed in mature

astrocytes in striatum[34]. Of note, we exclusively used Emx1-Cre female mice in our

breeding schemes given that the Cre-recombinase has been reported to be expressed in a

subset of male germline cells[35].

119   We sought to validate neuron-specific expression of the NuTRAP cassette. First,

immunohistochemistry of Emx1-NuTRAP hippocampal slices show distinct GFP and

mCherry expression in CA1 and CA3 pyramidal neurons (Fig. 1B-C) as well as the

granule cell layer of the dentate gyrus (DG; Fig. 1D). Furthermore, mCherry positive

nuclei remained intact and bound to the magnetic beads after our modified nuclear

isolation procedure (Supplementary Fig. 1A). As Emx1-expressing neural stem cells can

become astrocytes[31], we sought to determine whether a significant number of mature

astrocytes were obtained in the population of isolated cells. Flow cytometry experiments

using THY-1 as a neuronal marker and S100β as an astrocytic marker revealed a distinct

population of cells double positive for GFP and THY-1, whereas a S100β and GFP

double positive cell population was absent (Fig. 1E and Supplementary Table 1).

130   To perform Simultaneous INTACT & TRAP ("SIT"), hippocampal tissue was

dissected from both brain hemispheres, combined, and homogenized in one sample tube

(Fig. 1A). We then performed the "SIT" method on isolated samples by starting the

protocol with the beginning steps from the INTACT procedure modified to include

cycloheximide. Cycloheximide works rapidly to inhibit protein synthesis and is used for

maintaining crosslinks between translating mRNA and ribosomal subunits during

purification in the traditional TRAP method. Despite the presence of cycloheximide,

nuclear morphology from Emx1-expressing neurons is overall unchanged

(Supplementary Fig. 1A). Additionally, a Bioanalyzer was used to determine if there

were cycloheximide-induced double-stranded DNA breaks in our nuclear preparation that

could interfere with downstream DNA sequencing applications (such as ATAC-,

CUT&RUN-, or CUT&Tag-seq). We found no evidence of DNA double strand breaks

generating fragments of less than 1kb (Supplementary Fig. 1B). Following magnetic

purification of biotin-labeled nuclei, the supernatant fraction was removed and taken

through TRAP, while the pelleted nuclei were processed through the remaining steps of

INTACT (see Methods). Using qPCR, we found that TRAP isolated mRNA had

significantly reduced expression of *Mog* and *Cd11b* compared to total RNA, suggesting

depletion of oligodendrocyte and microglial populations in TRAP mRNA

(Supplementary Fig. 1C). Combining bilateral hippocampi from a single mouse yielded

TRAP-isolated mRNA of high quality and sufficient concentration for sequencing (RIN >

8 for all samples, average yield RNA = 14.367 ng/ul; Supplementary Table 2 and

Supplementary Fig. 1D). INTACT-isolated nuclei were further processed using the

CUT&RUN (Cleavage Under Targets and Release Using Nuclease[36]) method to isolate

antibody-specific protein-DNA interactions for downstream DNA sequencing. The

resulting DNA libraries were of high quality and concentration when using specific

antibodies (H4K8ac: average size = 1238 bp, average concentration = 126.8 nM;

H3K27me3: average size = 1032 bp, average concentration = 146.5 nM; Supplementary

Table 2). In contrast, the resulting DNA libraries using the non-specific IgG control had

substantially lower concentrations (IgG: average size = 1035 bp, average concentration =

24.2 nM; Supplementary Table 2) further indicating that nuclear DNA from both

isolations was of high starting quality.

161   To confirm our TRAP-isolated hippocampal mRNA came primarily from

excitatory neurons, we compared RNA-seq data from whole dorsal hippocampal tissue of

wild type mice to our TRAP-isolated RNA-seq data (TRAP-seq) from Emx1-NuTRAP

mice to assess for neuronal gene enrichment. We found that TRAP-isolated mRNA had

significant enrichment of several neuronal genes, including *Dlg4*, *Thy1*, *Eno2*, and *Syt4*,

as well as a significant reduction in astrocytic, microglial, oligodendrocyte, and inhibitory

neuronal genes (Fig. 1F). There was a relative expression increase of glial fibrillary

astrocytic protein (GFAP) in our TRAP samples. This gene can also be expressed in

neural stem populations that were present in our whole hippocampus samples, so this

finding may be a reflection of the neural stem cell population of the dentate gyrus[37,38].

Overall, these findings suggest that the Emx1-NuTRAP mouse model is a valid tool for

neuron-enriched isolation of sequencing-grade, translating mRNA and nuclear chromatin

from a single brain tissue homogenate.

Figure 1. Generation and validation of the Emx1-NuTRAP mouse (A) Schematic

representation of the Emx1-NuTRAP mouse generation and workflow of the

Simultaneous INTACT and TRAP ("SIT") protocol. (B) Immunofluorescence imaging of

the CA1 region of the hippocampus at 20x objective and (B') 60x objective after

incubation with an mCherry antibody overnight. (C) Immunofluorescence imaging of the

CA3 and CA2 region of the hippocampus at 20x objective and (C') 60x objective after

incubation with an mCherry antibody overnight. (D) Immunofluorescence imaging of the

DG region of the hippocampus at 20x objective and (D') 60x objective after incubation

with an mCherry antibody overnight. Scale bars are set to 50µm for all images (B-D). (E)

Flow cytometry for EGFP, THY-1, and S100β (15.3% of cells THY-1+/EGFP+ and

0.13% of cells S100β+/EGFP+). (F) Heatmap of differentially expressed neuronal and

other cell type markers from RNA-seq data comparing TRAP-isolated RNA from our

Emx1-NuTRAP mice and previously generated dorsal hippocampal tissue RNA from

wild type mice.

Performing INTACT/TRAP on the same or separate cell suspensions yields highly

comparable TRAP-seq and CUT&RUN-seq results.

Previous methods for performing INTACT and TRAP using NuTRAP mice have

taken separate tissue homogenates for each procedure[28,39]. In this study, we performed

SIT on a single tissue homogenate containing bilateral hippocampi (the products of SIT

are referred to as "simultaneous isolations"). To determine if our approach of SIT is

comparable to separate INTACT and TRAP procedures, we also performed separate

INTACT and TRAP from hippocampal tissue obtained from one brain hemisphere for

each method, counterbalancing for left versus right hemispheres (we refer to this protocol

as "separate isolations"). A Bioanalyzer was used to determine the amount and quality of

the RNA obtained from each type of isolation (Separate isolations: average RNA

concentration = 7.805 ng/ul, average RIN = 9.5; Simultaneous isolations: average RNA

concentration = 14.367 ng/ul, average RIN = 9.3; Supplementary Table 2). The average

RNA yield from the separate isolations (using a unilateral hippocampus) was

approximately equal to half of the average yield of the simultaneous isolations (which

combined bilateral hippocampi; Supplementary Table 2). Similarly, the final library

concentrations for the separately isolated IgG CUT&RUN-seq libraries were also

approximately half the concentration of the simultaneous isolations (average

simultaneous: 24.2nM, average separate: 11nM; Supplementary Table 2). We interpret

this to mean that nuclear DNA was fully intact in the simultaneous isolation because we

did not obtain substantially more than double the concentration in the simultaneous vs

separate isolations. Unilateral hippocampal homogenates yielded sufficient sequencing

concentrations and quality to allow for library preparations from individual mice

(Supplementary Table 2).

To determine if normalized TRAP-seq counts were similar between the two isolation

methods, we performed a Spearman's correlation between the datasets from sedentary

mice. The two TRAP-seq datasets were found to be highly correlated, with R values

greater than 0.5 and p value less than 2.2×10^{-16} (R=1, $p<2.2 \times 10^{-16}$; Fig. 2A).

CUT&RUN-seq was used to identify genomic regions interacting with either H4K8ac, an

activating histone post-translational modification (PTM), or H3K27me3, a generally

repressive histone PTM. We again applied Spearman's correlation to understand whether

the simultaneous vs separate INTACT isolation methods could influence CUT&RUN-seq

peak distribution. We compared normalized count data for CUT&RUN-seq peaks across

a representative chromosome (chromosome 2). We binned 100bp increments along the

entire chromosome from the simultaneous and separate isolations using datasets

generated from sedentary mice. Normalized sequencing counts, which reflected reads

assigned to binned genomic regions along chromosome 2, were highly similar between

conditions (H4K8ac: R=0.65, p<2.2x10^{-16}; H3K27me3: R=0.81, p<2.2x10^{-16}; Fig. 2B-C).

Taken together, these experiments suggest that translating mRNA and nuclear DNA

isolated from hippocampal homogenates using either simultaneous or separate isolation

procedures are comparable in terms of quality, concentration, functional characterization,

and normalized sequencing reads.

We next wanted to determine if the different isolation methods could bias resulting

gene expression on the basis of gene length. When plotting gene length against log fold

change of gene expression, we see a similar distribution pattern of ELE-induced

differentially expressed hippocampal genes (DEGs) between the simultaneous and

separate methods (Fig. 2D-E). We then compared the biological categories of the ELE-

induced DEGs obtained using either the separate or the simultaneous isolation methods

using Panther Gene Ontology (GO). Although many of the genes did not overlap (Fig.

2F-G and Supplementary Table 3), most GO terms did (14/20 GO terms for upregulated

genes and 15/17 for downregulated genes; Fig. 2H-I and Supplementary Table 3).

Furthermore, there was high similarity between the percentage of genes found in each of

the GO categories (Fig. 2J-K). Taken together, our results demonstrate that the

simultaneous approach for isolating and sequencing mRNA and nuclear chromatin can be

performed using a single hippocampal homogenate, and can generate highly similar

results as more traditional methods using separate cell samples to pair different types of

sequencing results.

Figure 2. Simultaneous and separate isolations of DNA and RNA from Emx1-

NuTRAP mice are comparable. For Fig. 2A-E, n=2 mice for simultaneous isolations

and n=3 mice for separate isolations. (A) Spearman's correlation for RNA expression

between separately isolated RNA and simultaneously isolated RNA (R=1, p<2.2x10^{-16}).

(B) Spearman's correlation between separate and simultaneously isolated DNA from

CUT&RUN-seq for H4K8ac with 100bp chromosome 2 position bins (R=0.65, p<2.2x10^{-}

^{16}). (C) Spearman's correlation between separate and simultaneously isolated DNA from

CUT&RUN-seq for H3K27me3 with 100bp chromosome 2 position bins (R=0.81,

p<2.2x10^{-16}). (D) Gene length versus fold change distribution plot for simultaneously

isolated RNA. (E) Gene length versus fold change distribution plot for separately isolated

RNA. (F) Genes upregulated in TRAP-seq after ELE in either simultaneous (n=2

sedentary mice and n=3 ELE mice) or separate isolations (n=3 mice per group). (G)

Genes downregulated in TRAP-seq after ELE in either simultaneous or separate

isolations. (H, I) Panther Gene Ontology: Biological Processes Venn diagram for genes

upregulated (H) or downregulated (I) after ELE in either separate or simultaneous

isolations. (J, K) Panther Gene Ontology: Biological Processes bar graph of gene

distributions for genes upregulated (J) and downregulated (K) after ELE in either separate

or simultaneous isolations.

Neuron-specific gene expression and histone PTMs are functionally comparable

between left and right hippocampal hemispheres.

Many studies take advantage of the brain's structural symmetry by using tissue

from each hemisphere for separate molecular processing. Prior evidence demonstrates

that hippocampal lateralization can influence LTP, hemisphere-specific glutamate

receptor density, and performance in certain memory tasks[40]. We wanted to determine if

there were differences in transcriptional programs and biological processes resulting from

ELE in left vs right hippocampi. Using the separate isolation approach described above,

we compared ELE-induced differential gene expression between left and right

hemispheres. We found that although individual gene expression patterns were different

between hippocampi originating in the left and right brain hemispheres (Fig. 3A-B and

Supplementary Table 4), the Panther GO: Biological Processes terms were similar, with

most categories overlapping (Fig. 3C-F and Supplementary Table 4). This could suggest

spatial or hemispheric assignment of genes with functional similarity.

Finally, we wanted to determine if left and right hippocampal hemispheres had

significant differences in DEGs and CUT&RUN-seq peak distributions at baseline

(without exercise). We performed Spearman's correlation on the transcript counts from

the TRAP-seq data (R=1, $p<2.2 \times 10^{-16}$; Fig. 3G), and the CUT&RUN-seq 100 bp binned

peak counts along chromosome 2 for H4K8ac (R=0.59, $p<2.2 \times 10^{-16}$; Fig. 3H) and for

H3K27me3 (R=0.70, $p<2.2 \times 10^{-16}$; Fig. 3I). We found that the seq peaks and transcript

counts were highly similar with significance considered as R values greater than 0.5 and

p value less than 2.2×10^{-16} (Fig. 3G-I). Overall, left and right hippocampal hemispheres

did not demonstrate significant differences in normalized sequencing counts and peak

distributions in the sedentary condition, suggesting that choice of hippocampal

hemisphere is a less important factor to consider in obtaining representative data on

transcription and transcriptional regulation via histone modifications.

Figure 3: Histone modifications and functional categorizations of gene expression

are similar regardless of hemispheric origin of hippocampal tissue. (A-B) Venn

diagram of genes upregulated (A) or downregulated (B) by ELE identified between the

left and right hemispheres by the separate isolation protocol TRAP-seq (n=3 mice per

group). (C-D) Venn diagram of Panther Gene Ontology: Biological Process terms for

genes upregulated (C) or downregulated (D) by ELE identified between the left and right

hemispheres by the separate isolation protocol TRAP-seq. (E-F) Percent of genes fitting

into each GO category for the left and right hemisphere for genes upregulated (E) and

downregulated (F) by ELE by separate isolation TRAP-seq. (G) Spearman's correlation

between the left and right hemispheres TRAP-seq data from the separate isolation

protocol of sedentary rlog normalized expression (R=1, p<2.2x10^{-16}). (H) Spearman's

correlation between hemispheres for CUT&RUN-seq for H4K8ac normalized count data

for chromosome 2 in 100bp bins (R=0.59, p<2.2x10^{-16}). (I) Spearman's correlation

between hemispheres for CUT&RUN-seq for H3K27me3 normalized count data for

chromosome 2 in 100bp bins (R=0.7, p<2.2x10^{-16}).

Early-life exercise promotes expression of plasticity-related genes and is associated
with transcriptional regulatory pathways implicated in hippocampal memory.

To identify the transcriptional effects of ELE in hippocampal neurons, we performed cell-type specific bulk sequencing on TRAP-isolated mRNA extracted from Emx1-NuTRAP hippocampi using our "SIT" protocol. We compared DEGs from sedentary mice vs ELE (running distances shown in Supplementary Fig. 2). Using DESeq2 to assess for DEGs (>30% expression increase with a p value < 0.05), we found that ELE alters gene expression in Emx1 expressing neurons (297 upregulated and 338 downregulated genes; Fig. 4A and Supplementary Table 3). Many of the genes upregulated after ELE are known to be involved in exercise and/or hippocampal memory mechanisms, including *Bdnf* [41] and *Nr4a1*[42]. To functionally categorize ELE-induced DEGs, we performed a Panther Gene Ontology (GO) analysis[43] focusing on the Molecular Function categorization and separated by upregulated and downregulated genes. Regardless of gene expression directionality, GO term categories with the most genes functionally assigned to them were "binding", "catalytic activity", "molecular function regulator", "transporter activity", "molecular transducer activity", and "structural molecule activity" (Fig. 4B). Many of the upregulated genes driving these categories are known to have critical roles in neuronal function (*Kcna1, Slc24a4, Stxbp5l, Gabra2,* and *Camk2n2*), neurodevelopment (*Artn, Kdm7a, Sox21, Gap43,* and *Efna5*), and hippocampal memory (*Bdnf, Nr4a1* and *Dusp5*).

To evaluate possible transcription factors and upstream regulators implicated by ELE-induced activated gene networks, we applied Qiagen's Ingenuity Pathway Analysis (IPA) to our TRAP-seq dataset[44]. We identified significant canonical signaling pathways implicated in ELE effects on hippocampal neuronal function (Supplementary Fig. 3 and Supplementary Table 5). Additionally, the top six upstream regulators by significance included CREB1, KMT2A, LEF1, SMAD3, EGR1, and MITF. Each of the top five transcription factors have been implicated in hippocampal development or function (Fig. 4C and Supplementary Table 5)[45-52]. The transcription factors KMT2A and LEF1 have not been previously associated with the effects of exercise on hippocampal function. Interestingly, CREB1 (through its associated CBP[53]) and KMT2A (through its methyltransferase activity[46]) both have histone modifying properties. LEF1 has been

shown to regulate neural precursor proliferation in the hippocampus[47]. SMAD3 is critical

for intermediate progenitor cell survival and negatively regulates serum insulin after

exercise[48,49]. EGR1 is an immediate early gene that recruits TET1 DNA demethylase

during neural activation and development[50]. MITF was the sixth upstream regulator

identified and is a transcription factor linked to autophagy mechanisms[51]. These specific

transcription factors were not significantly differentially expressed in our TRAP-seq

dataset; however, their activity may not be linked to a change in their own expression

after ELE but rather modulated by exercise to influence programs of gene expression.

To further understand whether the transcriptional profiles resulting from ELE were

enriched for specific *a priori* assigned molecular functions, we performed Gene Set

Enrichment Analysis (GSEA)[54]. We evaluated the "Reactome" category of functional

gene sets followed by a leading-edge analysis to further determine which genes were

driving the significant categories (Fig. 4D and Supplementary Table 6). Of the DEGs

upregulated after ELE, several interesting categorizations and the genes driving them

were revealed. The genes *H3c4*, *H2bc3*, *H2bc5*, and *H2bc9* genes, which encode histone

family member proteins, were driving the categories: "DNA methylation", "HDACs

deacetylate histones", and "transcriptional regulation by smRNAs". "Meiosis" and

"meiotic recombination" also emerged in these results which was unusual; however,

leading edge analysis showed many of these genes (*Atm, Blm, Msh4*, and *Mnd1*) to be

generally involved in cell-cycling processes. Exercise is well known to increase adult

neurogenesis in hippocampal dentate gyrus and can explain cell cycle gene enrichment in

our dataset[16]. Several nucleoporin complex genes (*Nup35/ 42/ 43/ 50/ 58/ 62/ 93/ 107/*

153/ 210) were also identified for their associations with categories such as: "gene

silencing by RNA", "nuclear envelope breakdown", and "transport of mature mRNAs

derived from intronless transcripts". *Nup* genes form a variety of nuclear pore complexes

that play critical roles in cellular processes including cell-cycle regulation, cellular

differentiation, and epigenetic control[55].

Of the downregulated pathways identified, translation-associated categories

("translation", "eukaryotic translation initiation/elongation", and "SRP-dependent co-

translational protein targeting to membrane") were driven by significant downregulation

of ribosomal protein gene families (*Rps*, *Rpl*, *Rplp*) and eukaryotic initiation factor 3

(*eIF3*) subunits. Over-expression of *eIF3* subunits has been linked to neurodegeneration,

and altered expression of *eIF3* has been associated with neurodevelopmental disorders[56].

Additionally, collagen genes (*Col1a1*, *Col1a2*, *Col2a1*, *Col3a1*, and *Col5a1*) were

downregulated after ELE, leading to the identification of categories including: "MET

activates PTK2 signaling", "MET promotes cell motility", and "collagen degradation".

Col1a1 and *Col1a2* have been previously identified as putative aging genes that decrease

in their expression after chronic exercise in female adult rodents[57]. By analyzing these

enriched pathways and the networks of genes driving them, we were able to identify

several expected, as well as unexpected, transcriptional programs activated by ELE that

could be unique to exercise timing during the juvenile-adolescent period.

Figure 4: ELE leads to transcriptomic changes in hippocampal neurons during

adolescence. (A) Volcano plot of differentially expressed genes reaching significance

identified by TRAP-seq on ELE versus sedentary (n=2 sedentary mice and n=3 ELE

mice, absolute value $\log_2$ fold change > 0.3785 and p-value < 0.05). (B) Panther Gene

Ontology: Molecular Function top terms by most genes assigned. (C) Top 6 "Upstream

Regulators" identified by Ingenuity Pathway Analysis (IPA). (D) Representative Gene

Set Enrichment Analysis: Reactome leading-edge diagrams showing genes upregulated

(D) or downregulated (D') in ELE, and their categories of enrichment. *Abbreviated

terms in (D): (1) "transport of mature mRNAs derived from intronless transcripts", (D'):

(2) "activation of the mRNA upon binding of the cap binding complex and EIFs and

subsequent binding to 43S", (3) srp dependent cotranslational protein targeting to

membrane, and (4) response of eif2ak4 gcn2 to amino acid deficiency.

Neuronal H4K8ac is enriched and H3K27me3 is reduced at a subset of plasticity

genes after ELE.

The power of the Emx1-NuTRAP mouse model (coupled with the "SIT" technical

approach) is the ability to directly pair differential gene expression with governing

epigenetic mechanisms. This is achieved by obtaining both TRAP-seq and epigenomic-

sequencing data from the same cell population (Fig. 1A). To investigate ELE-induced

changes in histone PTMs across the genome, hippocampal nuclei were obtained from

ELE and sedentary mice using the INTACT method followed by CUT&RUN-seq for the

modifications H3K27me3 and H4K8ac from nuclear chromatin. We chose these two

histone PTMs for their previously described functions. H4K8ac is a permissive

modification that is enriched at *Bdnf* after adult exercise, and its presence correlates with

improved hippocampal memory[22]. H3K27me3 is a repressive histone mark and a

common control in CUT&RUN-seq studies[36,58] and is decreased at the *Bdnf* promoter

region after contextual fear conditioning training[59]. Peaks were called using SEACR[60]

and histone PTM enrichment peaks were overlapped with ELE-induced DEGs obtained

from our TRAP-seq experiments described above (Fig. 5A and A'). We evaluated for the

presence of H4K8ac and H3K27me3 peaks in union with upregulated and downregulated

mRNA as a result of ELE. 93 upregulated genes had new H4K8ac peaks, while 35

downregulated genes had new H3K27me3 peaks (Fig. 5B and Supplementary Table 7).

Notably, of those 93 upregulated genes with H4K8ac peaks, 14 also had new H3K27me3

peaks as a result of ELE. 17 genes had presence of H3K27me3 in both conditions (ELE

and sedentary; Fig. 5A). We interpret these results to mean that transcription of these 93

genes was promoted as a result of ELE-induced H4K8ac.

The loss of either H3K27me3 or H4K8ac after ELE did not substantially correlate

with changes in translating mRNA expression (Fig. 5B). This result may indicate that

ELE-induced differential gene expression is associated with the addition, rather than

removal, of these two histone modifications. We found one gene that was downregulated

and had loss of H4K8ac (*Asphd1*), and two genes downregulated with loss of H3K27me3

(*Fdxr* and *Rtl1*; Fig. 5B and Supplementary Table 7). Of particular interest, reduced

expression of paternally-inherited *Rtl1* is associated with improved hippocampal memory

performance, whereas overexpression results in memory deficits[61,62]. Unsurprisingly, the

majority of histone PTM peaks newly present as a result of ELE did not correlate with

gene expression changes. However, notable upregulated genes with new H4K8ac after

ELE include *Prox1*, *Sntb2*, and *Kptn*. PROX1 is involved governing differentiation,

maturation, and DG versus CA3 cell identity in intermediate progenitor cells of the

hippocampal DG[63], while SNTB2 is involved in G protein-coupled receptor cell

signaling[64]. KPTN encodes a protein involved in cytoskeletal cell structure, and its

mutation can cause neurodevelopmental disability and seizures[65]. Interestingly,

downregulated genes found to have new H3K27me3 after ELE included *Col3a1*, *Efnb2*,

Epop, and *Myoc*. COL3A1 and MYOC are structural proteins involved in the

extracellular matrix and the cytoskeleton respectively[66-68]. EFNB2 is a signaling molecule

involved in cell migration and its haploinsufficiency can cause neurodevelopmental

disability[69-71]. EPOP is a known editor of the chromatin landscape by altering H2Bub and

H3K4me3 distributions[72,73].

Although changes in gene expression associated with H3K27me3 peaks were less

numerous than those associated with H4K8ac, we noticed a striking difference in the

distribution of peaks across the genome (Fig. 5C). After ELE, the distribution of

H3K27me3 decreases in the promoter region and increases in the distal intergenic and

intronic regions (Fig. 5C). To investigate how ELE-induced changes to histone PTM

distribution might alter gene expression, we looked at the distribution of peaks around

Prox1 (Fig. 5D). Most apparent changes in histone PTM peaks occurred in regions

annotated as regulatory regions, rather than the promoter region, which is concordant

with our data showing that most histone PTM changes happen outside of the promoter

region. Taken together, these results suggest that ELE promotes the new addition of

H4K8ac and H3K27me3 PTMs to regulate gene expression. Given that most identified

peaks did not correlate with DEGs, they may instead implicate regions of interest that are

"primed" for facilitated gene expression following future stimuli, such as learning.

A

H4K8ac ELE
5488
H4K8ac Sedentary 63
RNA Up in ELE 121
H3K27me3 ELE 1427
H3K27me3 Sedentary

A'

H4K8ac Sedentary
62
0
2290
H4K8ac ELE 5505
RNA Down in ELE 146
H3K27me3 ELE 1425
5
5
2134
43
H3K27me3 Sedentary

B. Change in Histone Mark Presence after ELE

Up in ELE

35
2

91
1

Down in ELE

# of Genes w/ RNA Expression

Legend:
- + H3K27me3
- − H3K27me3
- + H4K8ac
- − H4K8ac

H3K27me3   H4K8ac

C. Relative Distribution of Annotated Genomic Regions

% of Total Peaks

Region	H3K27me3 - Sedentary	H3K27me3 - ELE	H4K8ac - Sedentary	H4K8ac - ELE
top	4.92	4.15	2.77	2.39
1st Intron	7.63	8.24	10.10	11.13
Distal Intergenic	41.35	46.68	32.17	28.95
Other Exon	6.87	6.94	4.84	5.49
Other Intron	10.88	12.92	22.47	24.58
Promoter (<1kb)	21.13	13.63	19.90	19.31

Legend:
- 1st Exon
- 1st Intron
- 3' UTR
- 5' UTR
- Distal Intergenic
- Downstream (<0.3kb)
- Other Exon
- Other Intron
- Promoter (<1kb)
- Promoter (1-2kb)
- Promoter (2-3kb)

D. PROX1 Gene Tracks: chr1:189,900,750-190,843,610

Sedentary
H4K8ac-Sed-Sim-NT6.306
H4K8ac-Sed-Sim-NT7.2.3

ELE
H4K8ac-Ex-Sim-NT6.304
H4K8ac-Ex-Sim-Tg7.4.1

Enhancer-gene map from ENCODE 3 (UCSD/Ren) with promoters from EPDnew
Enhancers Rep
GENCODE VM23   GENCODE VM23 Comprehensive Transcript Set (only Basic displayed by default)
Replicated associations between enhancers and gene promoters from ENCODE 3 (UCSD/Ren) and EPDnew

Sedentary
H3K27me3-Sed-Sim-NT6.306
H3K27me3-Sed-Sim-NT7.2.3

ELE
H3K27me3-Ex-Sim-NT6.304
H3K27me3-Ex-Sim-Tg7.4.1

Legend:
- Regulatory Element
- Promoter Region

Figure 5: ELE alters H4K8ac and H3K27me3 occupancy. (A) Venn diagram of gene

peaks identified in CUT&RUN-seq for H4K8ac and H3K27me3 by SEACR (top 1% of

peaks FDR<0.1) overlapped with genes upregulated (A) and downregulated (A') in RNA

by ELE (n=2 sedentary mice and n=3 ELE mice). (B) CUT&RUN-seq SEACR peak calls

at genes that overlap with genes differentially expressed by ELE as determined by TRAP-

seq. (C) Stacked bar of the relative distribution of peak calls at various genomic regions

identified by CUT&RUN-seq called using SEACR. (D) Representative track of

CUT&RUN-seq data around *Prox1* showing differential peak height between ELE and

sedentary conditions for H4K8ac (top) and H3K27me3 (bottom).

ELE alters H4K8ac and H3K27me3 occupancy at regulatory regions of genes

involved in hippocampal memory consolidation.

456   Prior work from our lab and others has found that both ELE and adult exercise

can facilitate hippocampal long-term memory formation in mice exposed to a typically

sub-threshold learning event (3 minutes of object location memory (OLM) training,

which is normally insufficient for LTM formation in sedentary mice)[17,22]. These findings

suggest that ELE may "prime" neuronal function to facilitate hippocampal learning. In

this experiment, we investigate whether ELE-enabled hippocampal memory is associated

with altered presence of histone PTMs H4K8ac and H3K27me3 in regulatory regions of

genes associated with learning and hippocampal plasticity; the prediction being that

alterations of histone PTM occupancy as a result of ELE could promote gene expression

programs necessary for facilitating memory consolidation. Wild type animals underwent

ELE followed by OLM training on P42 and were sacrificed 60 minutes later (Fig. 6A).

Dorsal hippocampi were dissected, and tissue was homogenized and processed for bulk

RNA-seq. These data were compared to the ELE-induced DEGs, H4K8ac, and

H3K27me3 occupancy occurring without learning (generated from Emx1-NuTRAP mice

TRAP-seq and CUT&RUN-seq). We categorized DEGs based on the experimental group

of wild type animals: ELE alone, ELE+3 min learning stimulus, Sedentary+3 min

learning stimulus, and Sedentary+10 min learning stimulus. To discover ELE-induced

genes "primed" for regulation during memory consolidation, we focused specifically on

<u>c</u>andidate <u>g</u>enes <u>a</u>ssociated with <u>m</u>emory after <u>e</u>xercise (cGAME), <u>c</u>andidate <u>g</u>enes

<u>a</u>ssociated with <u>m</u>emory after <u>e</u>xercise or <u>s</u>edentary (cGAMES), and <u>c</u>andidate <u>g</u>enes

<u>a</u>ssociated with <u>m</u>emory after <u>s</u>edentary (cGAMS) conditions (Fig. 6B-C and

Supplementary Table 8). We then evaluated cGAME and cGAMES groups of genes for

the new presence or absence of histone modifications H4K8ac and H3K27me3 after ELE,

suggesting addition or removal of these histone marks denoting a permissive (increased

gene expression) or repressive (reduced gene expression) chromatin state, respectively

(Fig. 6B'-C' and Supplementary Table 9).

482   In the cGAMES group, there were 145 genes found to be significantly increased

in the mice that underwent a threshold learning event (3 min for ELE; 10 min for

sedentary mice). Of these 145 genes, 40% of them (58) had new H4K8ac following ELE

(Fig. 6 B-B'). We consider this group of genes to potentially be "primed," or readied, by

ELE for rapid transcription to support hippocampal memory consolidation at 3 min OLM

training, while also being involved in memory consolidation mechanisms in the threshold

learning of sedentary mice (10 min OLM training). The cGAME group of genes are those

which significantly increased in the ELE mice that underwent a threshold learning event

but were not increased in any of the other groups (including the sedentary, threshold

learning event / cGAMS group). These genes may be of unique importance for memory

consolidation specifically occurring after ELE given their absence in sedentary learners.

We found that the cGAME group had significant upregulation of 256 genes, with

approximately 30% (76) of those genes receiving new H4K8ac following ELE alone

(Fig. 6 B-B'). We also evaluated the cGAMES and cGAME that were significantly

downregulated during memory consolidation. We found 304 cGAMES and 361 cGAME

genes (Fig. 6 C). In both of these groups, a small minority of down-regulated genes had

altered H3K27me3 after ELE, and this was mostly gain of H3K27me3 (cGAMES: 6.3%;

cGAME: 6.4%) (Fig. 6 C').

We next wanted to determine how ELE "priming" directed gene expression

during memory consolidation that occurs after exercise (ELE 3min OLM) relative to

sedentary memory consolidation (Sed 10min OLM). We compared these to the group that

experienced the sub-theshold learning event (Sed 3min OLM). We compared the relative

$\log_2$ fold changes (LFCs) of the genes in cGAME and cGAMES using z-scores for each

group across the genes (Fig. 6D). In the vast majority of these primed genes, the ELE

drove gene expression during 3 min of OLM in the same direction as 10 minutes of OLM

training in sedentary animals (Fig. 6D). In some cases, the gene expression changes were

more exaggerated in the same direction as a sedentary threshold event (Fig. 6D). We

interpret this to indicate that ELE enables a 3min consolidation event to be threshold

(when 10 minutes is required for a sedentary mouse) by priming these genes for altered

gene expression in the same direction as a sedentary threshold event.

We next identified upstream regulators of cGAME and cGAMES genes with

ELE-induced histone modifications, as these mediators are likely involved in long-term

memory formation of a subthreshold learning event following ELE. The cGAMES and

cGAME lists were evaluated using Qiagen's Ingenuity Pathway Analysis (IPA). First,

candidate regulators were identified for the cGAME and cGAMES groups that have

histone PTM changes as a result of ELE that we measured (Fig. 6E and Supplementary

Table 10). Inhibition of MECP2 may indicate a reduction in DNA methylation. CREB1 is

required for LTM[45,74]. SNCA may play a role in neuroplasticity by modulating synaptic

vesicle transport[75].

Next, we wanted to identify gene networks that might be at play that are not just

regulated by the histone PTMs we selected. Many more histone modifications exist and

may regulate gene expression during consolidation in addition to those we selected. We

investigated the genes in cGAME and cGAMES using IPA and identified a set of likely

upstream regulators for these genes taken together (Fig. 6E'' and Supplementary Table

10). Once again MECP2 inhibition and CREB1 activation appear to be indicated

validating their presence in the previous IPA. Notably new to this group are inhibition of

FMR1, KMT2D, and TCF20, along with activation of FASN, CTNNB1, and MKNK1.

Next, we interrogated what networks might be unique to exercise-enabled consolidation

by running the cGAME list through IPA analysis (Fig. 6E' and Supplementary Table 10).

Unique to this group, RTN4, an inhibitor of neurite outgrowth[76], was predicted to trend

towards inhibition. GRIN3A, a glutamatergic NMDA receptor important to synaptic

development and refinement[77], was predicted activated in this group. These findings

suggest that ELE may enable consolidation by increasing neurite and synapse growth and

development. TCF7L2 is also predicted to be activated. Given that this transcription

factor is involved in Wnt signaling, and neurogenesis[78], and activated CTNNB1 is also

predicted to be associated with threshold consolidation generally, altered Wnt signaling

may be critical for ELE enabled consolidation.

Figure 6: ELE may prime consolidation-induced gene expression. (A) Experimental

design diagram. (B) Venn diagram representing upregulated genes relative to mice

without ELE or learning in either ELE, ELE and 3 min OLM training, Sed and 3 min

OLM training, and Sed and 10 min OLM training. (B') Pie chart showing the relative

percentage of genes that have new H4K8ac presence or H3K27me3 loss as a result of

ELE in specific groups identified in the Fig. 6B labeled cGAME and cGAMES. (C) Same

as Fig. 6B but for downregulated genes relative to mice without ELE. (C') Pie chart of

downregulated genes in the cGAME or cGAMES groups that have new H3K27me3 or

new loss of H4K8ac as a result of ELE. (D) Z-score based heatmap of genes in the

cGAMES and cGAME groups that have new H4K8ac or H3K27me3 presence or loss as

a result of ELE. Expression values used to calculate the Z-scores were relative to mice

that did not experience OLM training or ELE. (E) Upstream regulators identified by IPA

of the genes represented by Fig. 6D. (E') Upstream regulators identified by IPA of the

genes in the cGAME (up and downregulated) group that are unique to that when only

those are included in the IPA analysis. (E'') Upstream regulators identified by IPA of the

genes in the cGAME and cGAMES (up and downregulated) group when the genes in

those groups are put through IPA together. (* indicates that the activated or inhibited

indicator is registered from the IPAs based on the z-score for that regulator's expression

and not directly indicated based on software prediction).

Discussion

Identifying gene regulatory mechanisms activated by exercise during sensitive

developmental periods can aid in understanding how early-life exercise may promote a

"molecular memory" of exercise[23,59], via the epigenome, that informs long-term cell

function and behavioral outcomes. Here we present results demonstrating neuron-

specific, simultaneous characterization of translating mRNA and associated histone

modifications to reveal molecular signatures of the early-life exercise experience. To our

knowledge, this is the first report describing use of the NuTRAP construct in a

predominantly neuronal population. Additionally, our unique experimental approach for

simultaneous INTACT and TRAP ("SIT") is a technical advance for NuTRAP

applications: by modifying the nuclear chromatin isolation (INTACT) procedure to

include cycloheximide, one can obtain both polyribosome-bound mRNA (using TRAP)

and nuclear DNA from a single lysate to directly correlate gene expression programs with

epigenetic regulatory mechanisms in the same set of cells. The field of neuroepigenetics

has been challenged by very limited approaches for coupling cell-type specific

transcriptional profiles with chromatin modifications without compromising cell integrity

(through flow sorting) or genomic coverage (through single-cell sequencing). Therefore,

the NuTRAP model in combination with the "SIT" protocol offers an exciting

opportunity to overcome these challenges and perform paired transcriptomic and

epigenomic analyses in any brain cell type of interest for which a Cre line exists[28,39].

This study focuses on the impact of exercise on hippocampal gene expression and

associated epigenetic modifications specifically during juvenile-adolescent periods (the

fourth through sixth postnatal weeks of life in mice). We discover gene expression

programs, and alterations in the histone modifications H4K8ac and H3K27me3, that may

be unique to this early-life period. There have been recently published reports

investigating the effects of voluntary exercise on adolescent and adult hippocampal gene

expression and epigenetic modifications through RNA and DNA sequencing methods

from heterogenous cell populations[19,20]. Our data revealed several notable genes with

either chromatin-modifying or neurogenesis-regulating properties were either upregulated

with new H4K8ac (*Prox1*, *Sntb2*, and *Kptn*), or downregulated with new H3K27me3

(*Col3a1*, *Efnb2*, *Epop*, and *Myoc*). One could consider the mechanistic implications for

these genes: for example, ELE could "prime" the expression of *Prox1* by the addition of

H4K8ac, and in turn, subsequent *Prox1* expression can promote neurogenic fate and cell

location assignment[79]. ELE was also found to alter the expression of *Kptn,* which encodes

a protein involved in actin dynamics and neuromorphogenesis[65]. Increased *Kptn*

expression could promote cytoskeletal changes occurring with altered synaptic plasticity

after exercise. Overall, the differentially expressed genes and their associated chromatin

states occurring after ELE provide molecular candidates involved in the effects of ELE

on hippocampal function.

Traditional epigenetic and gene expression sequencing approaches to bulk tissue

samples are confounded by the cellular heterogeneity. . Further, current methods for

paired analysis of the epigenome and transcriptome are restricted by the need for multiple

transgenic animal models or the use of highly expensive single-cell sequencing

technologies. It is critical to obtain epigenomic information in a cell-type specific manner

because there are unique epigenetic modifications distinguishing cell subtypes of the

central nervous system[10,80]. Roh et al. first developed the NuTRAP mouse to characterize

adipocyte-specific and hepatocyte-specific epigenetic and transcriptional profiles[29]. The

NuTRAP mouse has since been used to perform paired transcriptome and DNA

modification analysis in astrocytes and microglia, demonstrating its utility in CNS cell-

types[39]. The Emx1-NuTRAP mouse described here provides a rapid, cost-effective means

for characterizing neuron-specific changes to epigenetic state and gene expression using

whole-tissue homogenates containing multiple cell types. In our technical validation

experiments, the "SIT" technique produced high quality RNA and DNA of sufficient

concentration for downstream sequencing studies from a single mouse hippocampus.

This method effectively increases the total sample size used for each analysis as the tissue

sample does not need to be separated for DNA and RNA isolation. CUT&RUN allows

for low-input cell-specific chromatin analysis[36,60] further reducing the required amount of

starting material. Indeed, the DNA obtained from SIT could be used for other

applications for evaluating the chromatin landscape, including ATAC-seq. Presently, the

field of neuroepigenetics is shifting to single-cell resolution for sequencing studies. Our

study was performed in bulk tissue enriched for neurons. As the goal of this work was to

characterize the transcriptional and epigenomic effects of early-life exercise, cell subtype

identity (as is often a priority in single-cell approaches) was not necessary for our initial

hypotheses but would offer a second layer of complexity to address this study's

questions. Single-cell approaches can bias against lowly-expressed genes, which may

present the risk of not capturing small expression changes in rate-limiting genes that may

have large changes in function as a result of ELE.

In this study, we also assessed for hemispheric differences in DEGs and histone

modifications using our SIT protocol. A common approach to paired transcriptomic and

epigenomic analyses is to use separate brain tissue samples, taking from one hemisphere

for gene expression studies and the other side for epigenetic analyses[81]. A previous study

utilizing the NuTRAP allele in non-neuronal brain cells used separate brain hemispheres

to perform INTACT and TRAP[39]. However, this assumes that cells of the same type from

each half of the tissue will show the same epigenetic and transcriptional profile. This is

not necessarily the case for all cell types, especially in the CNS, as structures of the brain

can show high degrees of hemispheric specialization[82-86]. To address the degree of

transcriptional or epigenetic difference between hippocampi from opposite hemispheres,

we performed separate INTACT and TRAP isolations using hippocampal tissue from one

hemisphere for DNA isolation and the contralateral hippocampus for mRNA isolation.

We found that although DEGs and histone PTMs differed between left and right

hippocampi, functional analysis using Panther GO revealed similar biological processes

between the two hemispheres. We interpret these data to suggest that hemispheric origin

of hippocampal tissue is not a critical contributor to conclusions drawn from these types

of sequencing studies.

Transcription is required for both hippocampal memory and LTP. LTP is defined

as a synaptic plasticity response considered the cellular correlate of learning and

memory[87]. Given our previous finding that ELE can facilitate long-term memory and

increase LTP [17], we hypothesize that this may be due to altering the epigenomic

landscape, to ultimately promote a transcriptional state supporting hippocampal memory

consolidation. This hypothesis is supported by the finding that in adult rodents, long-term

memory is formed in response to a subthreshold learning event after exercise or HDAC3

inhibition and involves increased acetylation of neuroplasticity gene promoters[22]. This

study identifies, for the first time, novel gene expression networks and their epigenetic

regulation that may be involved in the hippocampal memory benefits of ELE. We also

identified a group of genes that are altered in their gene expression only in sedentary

animals undergoing a threshold learning event ("cGAMS" group). The separation of this

group from the cGAME and cGAMES groups indicates that ELE may engage a separate,

perhaps complementary network of genes to promote memory consolidation. A next step

of this work will be to test the necessity and sufficiency of identified genes and gene

networks, their histone modifications, and/or their upstream regulators mediating the

effects of ELE on hippocampal memory performance. Another unanswered question is

how long the memory-promoting effects of ELE last after exercise cessation, and whether

there is stability of ELE-induced epigenetic modifications to allow for the persistence of

ELE effects.

In summary, our work reveals ELE-induced transcriptomic and epigenomic

signatures in hippocampal neurons using the Emx1-NuTRAP mouse for such

comparisons. A limitation for the use of the NuTRAP transgenic mouse is the

requirement of a Cre line to exist for the experimenter's cell type of interest. In our

Emx1-NuTRAP model, the Cre-recombinase in Emx1 expressing cells is constitutively

expressed, and this may present off-target effects when present during development.

Despite this, our study uses a feasible, previously undescribed approach in the field of

neuroepigenetics to pair transcriptomic and epigenomic sequencing results from a

population of hippocampal neurons. Our approach revealed new molecular epigenetic

candidates for further mechanistic studies to examine the effects of early-life exercise on

hippocampal function. Furthermore, understanding how exercise uniquely impacts

neuronal function during juvenile-adolescent developmental periods could provide

insights into an early-life epigenetic memory of the ELE experience, revealing possible

targets for improving memory function or mitigating memory deficits in the setting of

early-life cognitive and neurodevelopmental disorders[88].

Acknowledgements

This work was supported by the National Institute of Neurological Disorders and Stroke

of the National Institutes of Health, under award number K12NS098482, the Conte

Center @ UCI Seed Grant, UC Irvine Institute for Clinical and Translational Sciences

Pilot Award, and the Robert Wood Johnson Foundation Amos Medical Faculty

Development Program (awarded to A.S.I.). We thank the UC Irvine School of Medicine,

Departments of Pediatrics for support. We would also like to thank Dr. Marcelo Wood

for scientific discussions and critical input on the project, and Drs. Melanie Oakes and

Jenny Wu of the UC Irvine Genomics High Throughput Sequencing Facility for next

generation sequencing experiments support.

Author Contributions

A.M.R., T.D.F., N.E.N. and A.S.I. designed the studies and interpreted the results.

A.M.R., T.D.F., and A.S.I. performed INTACT and TRAP studies. A.M.R. built libraries

for RNA- and TRAP-sequencing. N.E.N. performed CUT&RUN-seq experiments and

built libraries for sequencing. N.E.N. and A.S.I. performed flow cytometry experiments.

A.M.R. and A.S.I. performed immunostainings and immunofluorescent imaging. D.A.V.

and A.B. built exercise cages and monitored running behavior. A.M.R. performed all

bioinformatic analysis of sequencing data. A.M.R. and N.E.N. analyzed sequencing data,

performed GO, GSEA and IPA analyses, and all other downstream analyses except where

noted. A.S.I. performed behavioral studies. T.D.F. performed Spearman's correlations

and gene length by fold change analysis. T.D.F. performed qPCR. N.E.N. designed the

figures. A.M.R., T.D.F., and A.S.I. wrote and edited the manuscript, with N.E.N.

providing edits to the manuscript. All authors discussed the results and edited and

approved the manuscript.

Competing Interests

The authors declare no competing financial interests.

706   **Methods**

Resource Availability

Lead Contact

Further information and requests for resources and reagents should be directed to and will

be fulfilled by the lead contact, Dr. Autumn Ivy (aivy@uci.edu)

Materials Availability

•   No new reagents or materials were generated for this study.

•   The mouse lines used for this study are available from Jackson Labs (Emx1-

715   IRES-*Cre* knock-in mice, Jackson Laboratory Stock No: 005628, and NuTRAP

716   mice, Jackson Laboratory Stock No: 029899).

718   *Animals*

719   Emx1-IRES-*Cre* knock-in mice (Jackson Laboratory Stock No: 005628),

NuTRAP mice (Jackson Laboratory Stock No: 029899), and C57Bl6/J (Jackson

Laboratory Stock No: 000664) wild type mice were obtained from Jackson Laboratories.

Female Emx1-IRES-*Cre* and male NuTRAP mice were crossed to generate the Emx1-

NuTRAP mice used for this study. Mice were given free access to food and water, and

lights were maintained on a standard 12-hour light/dark cycle. On postnatal day (P) 21,

mice were weaned and pair-housed in either standard bedding cages or cages equipped

with a running wheel. Only male mice were used for the studies performed. Experiments

were conducted according to US National Institutes of Health guidelines for animal care

and use and were approved by the Institutional Animal Care and Use Committee of the

University of California, Irvine.

Behavior

732   Wild type mice were habituated, handled and then underwent OLM training[89] for

either three or ten minutes at P41 as described previously[17]. On postnatal day 36, mice

were brought into a testing room with reduced room brightness. Mice were handled for

approximately 2 min each, for a total of 5 consecutive days prior to the OLM training

session (twice a day for the first 2 days followed by once a day for the next 3 days).

Habituation sessions were 5 min, twice per day for 3 days (P39-P41) and occurred within

chambers containing four unique spatial cues on each wall of the chamber (horizontal

lines, black X, vertical strip, and blank wall). Habituation overlapped with the last 3 days

of handling. During the training phase (P42), mice were placed into the same chambers

with two identical objects and exposed to either a subthreshold (3 min) or threshold (10

min) training period. Mice were sacrificed for consolidation experiments 1 hour later.

Exercise Paradigm

After weaning on P21, male mice from the same litters were randomly divided

and pair-housed in either standard cages without a wheel or cages equipped with a

stainless-steel running wheel similar to methods already described[17]. Animals housed

with running wheels had free access to the wheels from P21–41. Real-time data

acquisition (Vital View Software) was used to track distance ran recorded by magnetic

sensors that detect wheel revolutions. Each monitored wheel tracked running distance for

the entire cage (two mice per cage). Revolutions were quantified for every minute daily

for the duration of the exercise period and converted to distance ran per cage (km).

Exercised mice that underwent OLM training were individually tracked as described in

Valientes et al. 2021[90].

Simultaneous INTACT and TRAP ("SIT")

On P42 both ELE and sedentary control (n=4/group) Emx1-NuTRAP mice were

sacrificed by rapid cervical dislocation and both hemispheres of the hippocampus were

dissected out and collected for tissue processing. Hippocampal tissue samples were

mechanically homogenized in 1mL of nuclear preparation buffer (NPB) (10mM HEPES

(pH7.5); 1.5mM $MgCl_2$; 10mM KCl; 250mM sucrose; 0.1% IGEPAL CA-630; 0.2mM

DTT; 100μg/mL cycloheximide; 1x Complete EDTA-free protease inhibitor (Roche)) by

razor blade. Samples were then triturated with an 18G needle and syringe 50x on ice.

Homogenized tissue samples were pipetted slowly onto 100μm cell strainer caps on

5mL tubes and centrifuged at 1,000rcf for 10 minutes at 4°C. Cell pellets were

resuspended, and samples were pipetted onto 40μm cell strainer caps on 5mL tubes and

centrifuged at 1,000rcf for 10 minutes at 4°C. For nuclear isolation, 100μL of

Streptavidin-coated Dynabeads were washed twice with 1mL NPB and resuspended with

100μL NPB. Cell pellets were resuspended and added to washed Streptavidin-coated

Dynabeads. Samples were briefly pipetted to thoroughly suspend the beads and incubated

on ice for 20 minutes. Samples were placed on a magnetic stand to pull down nuclei

bound to the beads. The supernatant from each sample was slowly pipetted off the beads

and transferred to new 1.5mL tubes for RNA isolation (described later). Nuclei-bound

beads were washed twice with 1mL INTACT buffer (10mM HEPES (pH7.5); 1.5mM

MgCl2; 10mM KCl; 250mM sucrose; 0.1% IGEPAL CA-630; 1x Complete EDTA-free

protease inhibitor (Roche)) and resuspended in 100μL INTACT buffer. Bead-bound

nuclei were flash frozen on dry ice and stored at -80ºC until ready for CUT&RUN

chromatin digestion and DNA purification.

For RNA isolation, 100μL of concentrated TRAP isolation buffer (10mM HEPES

(pH 7.5); 117mM MgCl2; 1M KCl; 10% IGEPAL CA-630; 100μg/mL cycloheximide;

11mg/mL sodium heparin; 20mM DTT; 2.2units/μL Rnasin; 1x Complete EDTA-free

protease inhibitor (Roche)) was added to each sample (supernatant above). Samples were

mixed by gentle pipetting and centrifuged at 16,000rcf for 10 minutes at 4ºC. The

supernatant from each sample was transferred to new 1.5mL tubes and 50μL of protein-G

Dynabeads was added to each sample. (To bind anti-GFP, 100μL of protein-G

Dynabeads were washed twice with 500μL NPB and resuspended to a final 100μL

volume. 2μL anti-GFP antibody (Sigma-Aldrich, G6539, Lot:128M4867V) was added to

each tube and incubated for 1 hour on a rotating nutator at 4ºC. Excess anti-GFP was

washed off with 200μL NPB and beads were resuspended in a final 100μL volume.)

Samples were incubated for 3 hours on a rotating nutator at 4ºC. Beads were washed 3x

in 1mL high salt wash buffer (HSWB) (50mM Tris (pH 7.5); 12mM MgCl2; 300mM

KCl; 1% IGEPAL CA-630; 2mM DTT; 100μg/mL cycloheximide) at 4ºC. After the third

wash the beads were slowly resuspended in 350μL RLT buffer (from Qiagen RNeasy

Mini kit #74004) by slowly pipetting up and down and incubated on a rotating nutator for

30 minutes at room temperature. Samples were placed on a magnetic stand to pull down

the beads, and the supernatant from each were transferred to new 1.5mL tubes and stored

at -80ºC until ready for RNA purification.

Separate TRAP RNA Isolation

On P42, mice (n=3 mice/group) were sacrificed and the hippocampus was

dissected out as described above. Either the left or right brain hemisphere of the

hippocampus was processed for TRAP RNA isolation using methods similar to those

previously described[28,91] and the other hemisphere was processed for INTACT nuclear

isolation (see below). Hippocampal tissue was homogenized with a motorized pestle in

1ml of TRAP homogenization buffer (50mM Tris, (pH 7.5); 12mM $MgCl_2$; 100mM KCl;

1% NP-40; 100μg/ml cycloheximide; 1mg/ml sodium heparin; 2mM DTT; 0.2units/μl

RNasin; 1x Complete EDTA-free protease inhibitor (Roche)). Samples were centrifuged

at 16,000rcf for 10 minutes at 4ºC. The supernatant from each sample was collected and

incubated with 50μL of protein G Dynabeads (washed twice in 1mL TRAP

homogenization buffer without RNasin or protease inhibitor and resuspended in 50μL of

the same buffer). Samples were incubated for 30 minutes on a rotating nutator at 4ºC.

Following the incubation, 100μL of anti-GFP bound protein-G Dynabeads was added to

each sample. (To bind anti-GFP, 100μL of protein-G Dynabeads were washed twice with

500μL TRAP homogenization buffer without RNasin or protease inhibitor and

resuspended to a final 100μL volume. 2μL anti-GFP antibody (Sigma-Aldrich, G6539,

Lot:128M4867V) was added to each tube and incubated for 1 hour on a rotating nutator

at 4ºC. Excess anti-GFP was washed off with 200μL TRAP homogenization buffer

without RNasin or protease inhibitor and beads were resuspended in a final 100μL

volume.) Samples were incubated for 3 hours on a rotating nutator at 4ºC. Beads were

washed 3x in 1mL HSWB at 4ºC. After the final wash, the beads were resuspended in

350μL RLT buffer (from Qiagen RNeasy Mini kit #74004) and incubated on a rotating

nutator for 30 minutes at room temperature. Following the incubation, samples were

transferred to new 1.5mL tubes and stored at -80ºC until ready for RNA purification.

Separate INTACT Nuclear Isolation

The remaining hippocampal hemisphere from each mouse (not used for TRAP

RNA isolation above) was processed for INTACT DNA isolation using methods similar

to those described[28,92]. Tissue samples were quickly chopped with a razor blade, added to

1mL of INTACT buffer, and triturated with an 18G needle and syringe 50x on ice.

Samples were then added onto 100μm cell strainer caps on 5mL tubes and centrifuged at

1,000rcf for 10 minutes at 4ºC. Cell pellets were resuspended and samples were pipetted

onto 40μm cell strainer caps and centrifuged at 1,000rcf for 10 minutes at 4ºC. 100μL of

streptavidin coated Dynabeads (washed twice with 1mL INTACT buffer and resuspended

to a final 100 μL volume) were added to each sample, and samples were incubated on ice

for 20 minutes. After incubation, nuclei bound beads were washed twice with 1mL and

resuspended in 100μL INTACT buffer. Bead-bound nuclei were flash frozen on dry ice

and stored at -80ºC until ready for CUT&RUN chromatin digestion and DNA

purification.

RNA Purification and Sequencing Library Preparation

RNA was purified using the Qiagen RNeasy kit according to the manufacturer's

protocol. After purification, RNA quantity and quality were checked using the Qubit

RNA High Sensitivity assay and Agilent Bioanalyzer's Eukaryote Total RNA Pico assay,

respectively, at the University of California, Irvine Genomics High Throughput Facility

(UCI GHTF). Using an input of 15ng of total RNA, mRNA was isolated using

NEXTFLEX Poly(A) Beads 2.0, and the mRNA sequencing libraries were generated

using the NEXTFLEX Rapid Directional RNA-Seq Kit 2.0 according to the

manufacturer's instructions. Final libraries were sent to the UCI GHTF for Qubit dsDNA

High Sensitivity assay and the Agilent Bioanalyzer DNA High Sensitivity assay to

determine the quantity and quality of the RNA-seq libraries, respectively. Libraries were

sequenced at the UCI GHTF using 100bp paired end reads on a single lane of an Illumina

NovaSeq6000 to a minimum sequencing depth of 50 million reads.

Dorsal Hippocampal RNA Isolation

P42 wild type male mice were sacrificed by cervical dislocation. Left dorsal

hippocampus was extracted by bisecting the isolated hippocampus. The tissue was

homogenized and RNA was extracted using the Qiagen RNeasy kit according to the

manufacturer's protocol. Sequencing libraries were prepared from the RNA as described

under the section "RNA purification" and the UCI GHTF performed sequencing library

preparation, except that ERCC ExFold RNA Spike-In Mixes (ThermoFisher cat.

4465739) were added according to the manufacturer's instructions.

Cleavage Under Target and Release Using Nuclease (CUT&RUN)

864   For chromatin analysis, CUT&RUN was performed using the EpiCypher

CUTANA CUT&RUN protocol with slight modifications similar to what has already

been described[36]. Nuclei bound to beads were thawed and pulled down on a magnetic

stand. The supernatant was removed and beads were resuspended in 200μL wash buffer

(20mM HEPES (pH 7.5); 150mM NaCl; 0.5mM spermidine; 1x Complete EDTA-free

protease inhibitor (Roche)). Each sample was divided into three aliquots (50μL for anti-

H3K27me3 (Cell Signaling Technologies, C36B11, Lot: 16), 100μL for anti-H4K8ac

(Epicypher, 13-0047, Lot: 20202001-11), and 50μL for anti-IgG control (Rabbit IgG

Fisher Scientific, 026102)). Beads were pulled down on a magnetic stand, resuspended in

100μL wash buffer and incubated for 10 minutes at room temperature. After incubation

the beads were pulled down, the supernatant was removed, and the beads were

resuspended in 50μL ice cold antibody buffer (20mM HEPES (pH 7.5); 150mM NaCl;

0.5mM spermidine; 0.001mM digitonin; 2mM EDTA; 1x Complete EDTA-free protease

inhibitor (Roche)). 0.5μL of the appropriate antibody was added to each sample, and the

samples were mixed by gentle pipetting prior to overnight incubation at 4ºC on a nutator.

879   On the second day, the beads were pulled down on a magnetic stand, the

supernatant was removed and the beads were washed twice in 250μL ice cold wash

buffer with 0.001% digitonin. After the washes, beads were resuspended in 50μL ice cold

wash buffer with 0.001% digitonin, and 2.5μL of CUTANA pAG-MNase was added to

each sample and mixed by gentle pipetting. Samples were incubated for 10 minutes at

room temperature and beads were pulled down on a magnetic stand. The supernatant was

removed, and the beads were washed twice in 250μL ice cold wash buffer with 0.001%

digitonin and resuspended in 50μL of the same. 1μL of 100mM $CaCl_2$ was added to each

sample and mixed by gentle pipetting. Samples were incubated for 2 hours on a nutator at

4ºC. Following the incubation, 33μL of stop buffer (340mM NaCl; 20mM EDTA; 4mM

EGTA; 50μg/mL RNase A; 50μg/mL Glycogen) and 1.65μL of Spike-In DNA (Cell

Signaling Technology, #40366) was added to each sample. Samples were incubated in a

preheated thermocycler for 10 minutes at 37ºC. After incubation, samples were

transferred to new 1.5mL tubes and immediately processed for DNA purification.

DNA Purification and Library Preparation

895   DNA was purified using the Monarch PCR and DNA cleanup kit (New England

Biolabs, 13-0041) according to the manufacturer's instructions. After purification, DNA

fragments were used to generate sequencing libraries using the NEXTFLEX Rapid DNA-

Seq Kit 2.0 according to the manufacturer's instructions. Indices were diluted to a final

concentration of 1:1,000 before being added to samples, and a total of 13 PCR cycles was

used for DNA amplification of the libraries. Final libraries were sent to the UCI GHTF

for Qubit dsDNA High Sensitivity assay and the Agilent Bioanalyzer DNA High

Sensitivity assay to determine the quantity and quality of the CUT&RUN-seq libraries,

respectively. Libraries were sequenced at the UCI GHTF using 100bp paired end reads

on a single lane of an Illumina NovaSeq6000 to a minimum sequencing depth of 10

million reads.

qPCR

908   cDNA was generated from TRAP-isolated and total input RNA samples (n=2

samples/group) using the Transcriptor First Strand cDNA Synthesis Kit (Roche)

following the manufacturer's instructions. Relative gene expression for *Aqp4* (F: 5'-

ATCCAGCTCGATCTTTTGGA -3', R: 5'-TGAGCTCCACATCAGGACAG -3'),

Tubb3 (F: 5'-GTCTCTAGCCGCGTGAAGTC -3', R: 5'-

GCAGGTCTGAGTCCCCTACA -3'), *Cd11b* (F: 5'-CCCATGACCTTCCAAGAGAA -

3', R: 5'-ACACTGGTAGAGGGCACCTG -3'), and *Mog* (F: 5'-

AAGAGGCAGCAATGGAGTTG -3', R: 5'-GACCTGCAGGAGGATCGTAG -3') was

determined by qPCR using FastStart Essential DNA Green Master Mix (Roche,

06402712001) following the manufacturer's protocol. Cycle counts for mRNA

quantification were normalized to *Gapdh* (F: 5'-CGTCCCGTAGACAAAATGGT -3', R:

5'-GAATTTGCCGTGAGTGGAGT -3'). Quantification was performed using the Pfaffl

method [93].

Immunofluoresence

Emx1-NuTRAP mice were anesthetized intraperitoneally with 50mg/kg of

sodium pentobarbital and transfused transcardially with 4% paraformaldehyde (PFA) in

1x PBS. After perfusion, whole brains were dissected out and incubated in 4% PFA

overnight at 4ºC. After fixation, brains were incubated in 30% sucrose for 72 hours and

mounted in optimal cutting temperature (OCT) compound. 50μm coronal sections were

sliced through the hippocampus using a microtome, and every tenth section was collected

in 1mL of cryoprotectant and stored at -20ºC until ready for immunofluorescent labeling.

For immunofluorescence, sections were brought up to room temperature and washed 2x

in 1x PBS on a nutator for 5 minutes each. Sections were then washed 3x in 1x PBS with

0.3% triton-X. After washes, sections were incubated with mCherry polyclonal antibody

(ThermoFisher, PA5-34974, Lot: UG2804409F) at a 1:200 dilution in blocking solution

(1x PBS with 1% BSA) overnight at 4ºC on a nutator. The following day, sections were

washed 3x in 1x PBS. Sections were incubated for two hours at room temperature with

goat anti-rabbit Alexa Fluor 594 secondary antibody (Invitrogen, A-11012, Lot:

2119134) at a 1:200 dilution in blocking solution. After incubation, the sections were

washed twice in 1x PBS. To label nuclei, sections were incubated with DAPI at a

1:10,000 dilution in 1x PBS for 20 minutes. Sections were then washed 3x in 1x PBS and

cover slipped with Vectatshield®. Fluorescent images were acquired using a Keyence

BZ-X810 All-in-One Fluorescence microscope using the optical sectioning module.

Fluorescence Activated Cell Sorting

Fluorescence activated cell sorting (FACS) was performed to characterize

neuronal and astrocytic NuTRAP cassette expression. Whole hippocampal tissue from

Emx1-NuTRAP mice was isolated and single-cell suspensions were immunostained for

cytometric analysis, as previously described[94]. We used antibodies for THY-1 (OX7)

AlexaFluor™ 647 (Santa Cruz Biotechnology, sc-53116 AF647, Lot: F2716;

concentration: 1:50) and S100β (Abcam, ab41548, Lot:GR3326165-1; concentration:

1:200) with an AlexaFluor™ 405 goat anti-rabbit IgG (H+L) (Invitrogen, A31556, Lot:

2273716; concentration:1:800) secondary antibody. FACS analysis was performed using

a BD FACSAria™ Fusion Flow Cytometer (BD Biosciences) at the University of

California, Irvine Stem Cell Core. Samples and single-stain controls were analyzed using

the FlowJo v10.8.1 software (BD Biosciences). Samples and controls were positively

gated for live cells (SSC-A and FSC-A) and single cells (FSC-H and FSC-A), and

negatively gated for autofluorescence (Comp-Alexa Fluor 647-A (Thy1) and Comp-

BV421-A (S100β)). Fraction of GFP+ neuronal cells was determined using quartile

analysis of Thy-1 and Comp-GFP-A (GFP). Fraction of GFP+ astrocytic cells was

determined using quartile analysis of S100β and Comp-GFP-A (GFP).

Bioinformatics Analysis

Reference assembly: For all sequencing analysis that required it, the primary

reference assembly used was Genome Reference Consortium Mouse Build 38 patch

release 6 (GRCm38.p6).

RNA-seq analysis: FastQ files were quality checked for sequencing errors using

FastQC (version 0.11.9)[95]. No files were found to have sufficient quality errors to

discount their use. Files were aligned using STAR Aligner (version 2.7.3a)[96]. Duplicate

reads were removed using Picard Tools (version 1.87)[97]. SAM Tools was used to convert

BAM files to SAM files for use in downstream analysis. FastQC, alignment and duplicate

removal were all preformed on the High-Powered Compute Cluster (HPC3) operated by

The Research Cyberinfrastructure Center (RCIC) at the University of California, Irvine.

R (version 4.1.0)[98] was used for differentially expressed genes (DEG) analysis. Genomic

Alignments (version 1.28.0)[99] (summarizeOverlaps mode="IntersectionNotEmpty",

singleEnd=FALSE, ignore.strand=FALSE, fragments=TRUE), Genomic Features

(version 1.44.0) (exons by gene), and R SAM Tools (version 2.8.0)[100]

(yieldSize=100000) were used to extract a count matrix and generate a summarized

experiment object. DEseq2 (version 1.32.0)[101] was used to perform a DEG analysis.

Ensemble IDs were converted to gene symbols using BiomaRt (version 2.48.1)[102,103].

PCA plots using the top 2000 genes were generated using ggplot2 (version 3.3.4)[104].

Samples were determined as outliers if the variability between the samples heavily

weighted PC1 to those samples. This excluded 2 sedentary samples and 1 ELE sample.

Heatmaps were generated using gplots (version 3.1.1)[105] and RColorBrewer (version1.1-

2)[106]. Volcano Plots were generated using ggplot2[104]. Venn diagrams were generated

using the Venn diagram tool from Bioinformatics and Evolutionary Genomics[107]. Gene

ontology analysis was performed using Panther Classification System (version 16.0)[43].

Genes were also categorized and a leading edge heatmap was also generated using Gene

Set Enrichment Analysis (version GSEA 4.1.0)[54]. Upstream regulators were identified

from the DEGs upregulated by ELE using Qiagen's Ingenuity Pathway Analysis (IPA)

(Fall 2021 Release)[44].

CUT&RUN-seq analysis: FastQ files were quality checked for sequencing errors

using FastQC (version 0.11.9)[95]. No files were found to have sufficient quality errors to

discount their use. FastQ files were aligned using Bowtie2 (version 2.4.1)[108]. Significant

peaks were called from these aligned files using SEACR, developed in CUT&Tag for

efficient epigenomic profiling of small samples and single cells[60,108], according to

CUT&Tag Data Processing and Analysis Tutorial (updated August 12 2020):

https://www.protocols.io/view/cut-amp-tag-data-processing-and-analysis-tutorial-

bjk2kky. An FDR higher than 0.1 was considered too high for called peaks. This

threshold excluded 4 samples (3 ELE H4K8ac separate isolation samples and 1 ELE

H4K8ac SIT sample). Peaks were called using the stringent and 0.01 settings. Significant

peaks were annotated using ChIPseeker (version 1.8.6)[109] using

TxDb.Mmusculus.UCSC.mm10.knownGene[110] as a reference with the following settings:

tssRegion = c(-3000, 3000), TxDb = TxDb.Mmusculus.UCSC.mm10.knownGene, level

= "transcript", assignGenomicAnnotation = TRUE, genomicAnnotationPriority =

c("Promoter", "5UTR", "3UTR", "Exon", "Intron", "Downstream", "Intergenic"), annoDb

= NULL, addFlankGeneInfo = FALSE, flankDistance = 5000, sameStrand = FALSE,

ignoreOverlap = FALSE, ignoreUpstream = FALSE, ignoreDownstream = FALSE,

overlap = "TSS", verbose = TRUE). This approach annotated peaks to the closest gene by

distance from the promoter, except if that peak falls within a gene before the distal

intergenic region. Ensemble IDs were converted to gene symbols using BiomaRt[102,103].

Gene lists of peaks present in each condition were used to generate Venn diagrams, as

with RNA-seq analysis[107]. Further Venn diagrams were generated comparing DEGs with

peak calls. Peaks were visualized using UCSC Genome Browser and Genome Browser in

a Box[111-113].

1014    Normalized counts (for read depth) at binned sections of the DNA as well as

1015 RNA-seq gene expression count (normalized for read depth) were compared between

1016 simultaneous and separate isolations as well as left and right hemispheres using

1017 Spearman's correlations in R with the package ggpubr (version 0.4.0)[114]. Bar plots were

1018 generated using Graph Pad 9 PRISM.

1019

1020 ***Data Availability***

1021    RNA-, TRAP- and CUT&RUN-seq data will be deposited and publicly available

1022 as of the date of publication. Accession numbers will be listed here pending publication.

1023 Any additional information required to reanalyze the data reported in this paper is

1024 available from the lead contact upon request.

1025   References
1026
1   Bedrosian, T. A., Quayle, C., Novaresi, N. & Gage, F. H. Early life experience
drives structural variation of neural genomes in mice. *Science (New York, N.Y.)*
359, 1395-1399, doi:10.1126/science.aah3378 (2018).
2   Stroud, H. *et al.* Early-Life Gene Expression in Neurons Modulates Lasting
Epigenetic States. *Cell* **171**, 1151-1164.e1116,
doi:https://doi.org/10.1016/j.cell.2017.09.047 (2017).
3   Herre, M. & Korb, E. The chromatin landscape of neuronal plasticity. *Curr Opin*
Neurobiol **59**, 79-86, doi:10.1016/j.conb.2019.04.006 (2019).
4   Campbell, R. R. & Wood, M. A. How the epigenome integrates information and
reshapes the synapse. *Nat Rev Neurosci* **20**, 133-147, doi:10.1038/s41583-019-
0121-9 (2019).
5   Kronman, H. *et al.* Long-term behavioral and cell-type-specific molecular effects
of early life stress are mediated by H3K79me2 dynamics in medium spiny
neurons. *Nat Neurosci* **24**, 667-676, doi:10.1038/s41593-021-00814-8 (2021).
6   Sagarkar, S. *et al.* LSD1-BDNF activity in lateral hypothalamus-medial forebrain
bundle area is essential for reward seeking behavior. *Progress in neurobiology*
202, 102048, doi:10.1016/j.pneurobio.2021.102048 (2021).
7   Webb, W. M. *et al.* The SETD6 Methyltransferase Plays an Essential Role in
Hippocampus-Dependent Memory Formation. *Biological psychiatry* **87**, 577-587,
doi:10.1016/j.biopsych.2019.05.022 (2020).
8   Bolton, J. L. *et al.* Unexpected Transcriptional Programs Contribute to
Hippocampal Memory Deficits and Neuronal Stunting after Early-Life Adversity.
Cell Rep **33**, 108511, doi:10.1016/j.celrep.2020.108511 (2020).
9   Campbell, R. R. *et al.* HDAC3 Activity within the Nucleus Accumbens Regulates
Cocaine-Induced Plasticity and Behavior in a Cell-Type-Specific Manner. *The*
Journal of neuroscience : the official journal of the Society for Neuroscience **41**,
2814-2827, doi:10.1523/JNEUROSCI.2829-20.2021 (2021).
10   Lister, R. *et al.* Global Epigenomic Reconfiguration During Mammalian Brain
Development. *Science* **341**, 1237905, doi:10.1126/science.1237905 (2013).
11   Lavery, L. A. & Zoghbi, H. Y. The distinct methylation landscape of maturing
neurons and its role in Rett syndrome pathogenesis. *Current opinion in*
neurobiology **59**, 180-188, doi:10.1016/j.conb.2019.08.001 (2019).
12   Aguilar, R., Bustos, F. J., Nardocci, G., van Zundert, B. & Montecino, M.
Epigenetic silencing of the osteoblast-lineage gene program during hippocampal
maturation. *Journal of cellular biochemistry* **122**, 367-384, doi:10.1002/jcb.29865
(2021).
13   West, A. E. & Greenberg, M. E. Neuronal activity-regulated gene transcription in
synapse development and cognitive function. *Cold Spring Harbor perspectives in*
biology **3**, doi:10.1101/cshperspect.a005744 (2011).
14   Bale, T. L. *et al.* Early life programming and neurodevelopmental disorders.
Biological psychiatry **68**, 314-319, doi:10.1016/j.biopsych.2010.05.028 (2010).
15   Cotman, C. W., Berchtold, N. C. & Christie, L.-A. Exercise builds brain health:
key roles of growth factor cascades and inflammation. *Trends in Neurosciences*
30, 464-472, doi:https://doi.org/10.1016/j.tins.2007.06.011 (2007).

16   Voss, M. W., Vivar, C., Kramer, A. F. & van Praag, H. Bridging animal and
human models of exercise-induced brain plasticity. *Trends in cognitive sciences*
17, 525-544, doi:10.1016/j.tics.2013.08.001 (2013).
17   Ivy, A. S. *et al.* A Unique Mouse Model of Early Life Exercise Enables
Hippocampal Memory and Synaptic Plasticity. *Scientific Reports* **10**, 9174,
doi:10.1038/s41598-020-66116-4 (2020).
18   Berchtold, N. C., Castello, N. & Cotman, C. W. Exercise and time-dependent
benefits to learning and memory. *Neuroscience* **167**, 588-597,
doi:10.1016/j.neuroscience.2010.02.050 (2010).
19   Urdinguio, R. G. *et al.* Physical exercise shapes the mouse brain epigenome. *Mol*
Metab **54**, 101398, doi:10.1016/j.molmet.2021.101398 (2021).
20   Abel, J. L. & Rissman, E. F. Running-induced epigenetic and gene expression
changes in the adolescent brain. *International journal of developmental*
neuroscience : the official journal of the International Society for Developmental
Neuroscience **31**, 382-390, doi:10.1016/j.ijdevneu.2012.11.002 (2013).
21   Ieraci, A., Mallei, A., Musazzi, L. & Popoli, M. Physical exercise and acute
restraint stress differentially modulate hippocampal brain-derived neurotrophic
factor transcripts and epigenetic mechanisms in mice. *Hippocampus* **25**, 1380-
1392, doi:10.1002/hipo.22458 (2015).
22   Intlekofer, K. A. *et al.* Exercise and Sodium Butyrate Transform a Subthreshold
Learning Event into Long-Term Memory via a Brain-Derived Neurotrophic
factor-Dependent Mechanism. *Neuropsychopharmacology* **38**, 2027-2034,
doi:10.1038/npp.2013.104 (2013).
23   Butler, C. W. *et al.* Exercise opens a temporal window for enhanced cognitive
improvement from subsequent physical activity. *Learn Mem* **26**, 485-492,
doi:10.1101/lm.050278.119 (2019).
24   Titterness, A. K., Wiebe, E., Kwasnica, A., Keyes, G. & Christie, B. R. Voluntary
exercise does not enhance long-term potentiation in the adolescent female dentate
gyrus. *Neuroscience* **183**, 25-31, doi:10.1016/j.neuroscience.2011.03.050 (2011).
25   Gomes da Silva, S. & Arida, R. M. Physical activity and brain development.
Expert Rev Neurother **15**, 1041-1051, doi:10.1586/14737175.2015.1077115
(2015).
26   Gomes da Silva, S. *et al.* Early exercise promotes positive hippocampal plasticity
and improves spatial memory in the adult life of rats. *Hippocampus* **22**, 347-358,
doi:10.1002/hipo.20903 (2012).
27   Fernandes, J., Arida, R. M. & Gomez-Pinilla, F. Physical exercise as an
epigenetic modulator of brain plasticity and cognition. *Neurosci Biobehav Rev* **80**,
443-456, doi:10.1016/j.neubiorev.2017.06.012 (2017).
28   Roh, H. C. *et al.* Simultaneous Transcriptional and Epigenomic Profiling from
Specific Cell Types within Heterogeneous Tissues In Vivo. *Cell reports* **18**, 1048-
1061, doi:10.1016/j.celrep.2016.12.087 (2017).
29   Deal, R. B. & Henikoff, S. The INTACT method for cell type-specific gene
expression and chromatin profiling in Arabidopsis thaliana. *Nature protocols* **6**,
56-68, doi:10.1038/nprot.2010.175 (2011).

30   Heiman, M. *et al.* A translational profiling approach for the molecular
characterization of CNS cell types. *Cell* **135**, 738-748,
doi:10.1016/j.cell.2008.10.028 (2008).
31   Gorski, J. A. *et al.* Cortical excitatory neurons and glia, but not GABAergic
neurons, are produced in the Emx1-expressing lineage. *J Neurosci* **22**, 6309-6314,
doi:10.1523/jneurosci.22-15-06309.2002 (2002).
32   Chan, C. H. *et al.* Emx1 is a marker for pyramidal neurons of the cerebral cortex.
Cerebral cortex (New York, N.Y. : 1991) **11**, 1191-1198,
doi:10.1093/cercor/11.12.1191 (2001).
33   Hong, S. M. *et al.* Reduced hippocampal neurogenesis and skill reaching
performance in adult Emx1 mutant mice. *Experimental neurology* **206**, 24-32,
doi:10.1016/j.expneurol.2007.03.028 (2007).
34   Willaime-Morawek, S. & van der Kooy, D. Cortex- and striatum- derived neural
stem cells produce distinct progeny in the olfactory bulb and striatum. *The*
European journal of neuroscience **27**, 2354-2362, doi:10.1111/j.1460-
9568.2008.06206.x (2008).
35   Luo, L. *et al.* Optimizing Nervous System-Specific Gene Targeting with Cre
Driver Lines: Prevalence of Germline Recombination and Influencing Factors.
Neuron **106**, 37-65.e35, doi:10.1016/j.neuron.2020.01.008 (2020).
36   Skene, P. J., Henikoff, J. G. & Henikoff, S. Targeted in situ genome-wide
profiling with high efficiency for low cell numbers. *Nature protocols* **13**, 1006-
1019, doi:10.1038/nprot.2018.015 (2018).
37   Liu, Y. *et al.* Glial fibrillary acidic protein-expressing neural progenitors give rise
to immature neurons via early intermediate progenitors expressing both glial
fibrillary acidic protein and neuronal markers in the adult hippocampus.
Neuroscience **166**, 241-251, doi:10.1016/j.neuroscience.2009.12.026 (2010).
38   Garcia, A. D. R., Doan, N. B., Imura, T., Bush, T. G. & Sofroniew, M. V. GFAP-
expressing progenitors are the principal source of constitutive neurogenesis in
adult mouse forebrain. *Nature Neuroscience* **7**, 1233-1241, doi:10.1038/nn1340
(2004).
39   Chucair-Elliott, A. J. *et al.* Inducible cell-specific mouse models for paired
epigenetic and transcriptomic studies of microglia and astroglia. *Commun Biol* **3**,
693, doi:10.1038/s42003-020-01418-x (2020).
40   Jordan, J. T. The rodent hippocampus as a bilateral structure: A review of
hemispheric lateralization. *Hippocampus* **30**, 278-292,
doi:https://doi.org/10.1002/hipo.23188 (2020).
41   Berchtold, N. C., Chinn, G., Chou, M., Kesslak, J. P. & Cotman, C. W. Exercise
primes a molecular memory for brain-derived neurotrophic factor protein
induction in the rat hippocampus. *Neuroscience* **133**, 853-861,
doi:10.1016/j.neuroscience.2005.03.026 (2005).
42   McNulty, S. E. *et al.* Differential roles for Nr4a1 and Nr4a2 in object location vs.
object recognition long-term memory. *Learning & memory (Cold Spring Harbor,*
N.Y.) **19**, 588-592, doi:10.1101/lm.026385.112 (2012).
43   Mi, H. & Thomas, P.   123-140 (Humana Press, 2009).

44   Krämer, A., Green, J., Pollard, J. & Tugendreich, S. Causal analysis approaches in
1160   Ingenuity Pathway Analysis. *Bioinformatics* **30**, 523-530,
1161   doi:10.1093/bioinformatics/btt703 (2014).
1162   45   Kandel, E. R. The molecular biology of memory: cAMP, PKA, CRE, CREB-1,
1163   CREB-2, and CPEB. *Molecular brain* **5**, 14, doi:10.1186/1756-6606-5-14 (2012).
1164   46   Kerimoglu, C. *et al.* KMT2A and KMT2B Mediate Memory Function by
1165   Affecting Distinct Genomic Regions. *Cell reports* **20**, 538-548,
1166   doi:10.1016/j.celrep.2017.06.072 (2017).
1167   47   Ashbrook, D. G. *et al.* Transcript co-variance with Nestin in two mouse genetic
1168   reference populations identifies Lef1 as a novel candidate regulator of neural
1169   precursor cell proliferation in the adult hippocampus. *Front Neurosci* **8**, 418,
1170   doi:10.3389/fnins.2014.00418 (2014).
1171   48   Tapia-González, S., Muñoz, M., Cuartero, M. & Sánchez-Capelo, A. Smad3 is
1172   required for the survival of proliferative intermediate progenitor cells in the
1173   dentate gyrus of adult mice. *Cell Communication and Signaling* **11**, 93,
1174   doi:10.1186/1478-811x-11-93 (2013).
1175   49   Tiano, J. P., Springer, D. A. & Rane, S. G. SMAD3 Negatively Regulates Serum
1176   Irisin and Skeletal Muscle FNDC5 and Peroxisome Proliferator-activated
1177   Receptor γ Coactivator 1-α (PGC-1α) during Exercise. *Journal of Biological*
1178   *Chemistry* **290**, 7671-7684, doi:10.1074/jbc.m114.617399 (2015).
1179   50   Sun, Z. *et al.* EGR1 recruits TET1 to shape the brain methylome during
1180   development and upon neuronal activity. *Nature Communications* **10**,
1181   doi:10.1038/s41467-019-11905-3 (2019).
1182   51   Bouché, V. *et al.* Drosophila Mitf regulates the V-ATPase and the lysosomal-
1183   autophagic pathway. *Autophagy* **12**, 484-498,
1184   doi:10.1080/15548627.2015.1134081 (2016).
1185   52   Guan, Z. *et al.* Integration of long-term-memory-related synaptic plasticity
1186   involves bidirectional regulation of gene expression and chromatin structure. *Cell*
1187   **111**, 483-493, doi:10.1016/s0092-8674(02)01074-7 (2002).
1188   53   Korzus, E., Rosenfeld, M. G. & Mayford, M. CBP Histone Acetyltransferase
1189   Activity Is a Critical Component of Memory Consolidation. *Neuron* **42**, 961-972,
1190   doi:10.1016/j.neuron.2004.06.002 (2004).
1191   54   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based
1192   approach for interpreting genome-wide expression profiles. *Proceedings of the*
1193   *National Academy of Sciences* **102**, 15545-15550, doi:10.1073/pnas.0506580102
1194   (2005).
1195   55   Nofrini, V., Di Giacomo, D. & Mecucci, C. Nucleoporin genes in human diseases.
1196   *European Journal of Human Genetics* **24**, 1388-1395, doi:10.1038/ejhg.2016.25
1197   (2016).
1198   56   Gomes-Duarte, A., Lacerda, R., Menezes, J. & Romão, L. eIF3: a factor for
1199   human health and disease. *RNA Biol* **15**, 26-34,
1200   doi:10.1080/15476286.2017.1391437 (2018).
1201   57   Alvarez-López, M. J. *et al.* Long-term exercise modulates hippocampal gene
1202   expression in senescent female mice. *J Alzheimers Dis* **33**, 1177-1190,
1203   doi:10.3233/jad-121264 (2013).

58 Butler, A. A., Sanchez, R. G., Jarome, T. J., Webb, W. M. & Lubin, F. D. O-
GlcNAc and EZH2-mediated epigenetic regulation of gene expression during
consolidation of fear memories. *Learning & memory (Cold Spring Harbor, N.Y.)*
26, 373-379, doi:10.1101/lm.049023.118 (2019).
59 Butler, A. A., Sanchez, R. G., Jarome, T. J., Webb, W. M. & Lubin, F. D. O-
GlcNAc and EZH2-mediated epigenetic regulation of gene expression during
consolidation of fear memories. *Learning & memory (Cold Spring Harbor, N.Y.)*
26, 373-379, doi:10.1101/lm.049023.118 (2019).
60 Kaya-Okur, H. S. *et al.* CUT&Tag for efficient epigenomic profiling of small
samples and single cells. *Nature Communications* **10**, doi:10.1038/s41467-019-
09982-5 (2019).
61 Chou, M.-Y. *et al.* RTL1/PEG11 imprinted in human and mouse brain mediates
anxiety-like and social behaviors and regulates neuronal excitability in the locus
coeruleus. *Human Molecular Genetics*, doi:10.1093/hmg/ddac110 (2022).
62 Kitazawa, M., Sutani, A., Kaneko-Ishino, T. & Ishino, F. The role of eutherian-
specific RTL1 in the nervous system and its implications for the Kagami-Ogata
and Temple syndromes. *Genes to Cells* **26**, 165-179,
doi:https://doi.org/10.1111/gtc.12830 (2021).
63 Stergiopoulos, A., Elkouris, M. & Politis, P. K. Prospero-related homeobox 1
(Prox1) at the crossroads of diverse pathways during adult neural fate
specification. *Front Cell Neurosci* **8**, 454, doi:10.3389/fncel.2014.00454 (2014).
64 Camp, N. D. *et al.* Individual protomers of a G protein-coupled receptor dimer
integrate distinct functional modules. *Cell Discov* **1**, 15011-,
doi:10.1038/celldisc.2015.11 (2015).
65 Baple, E. L. *et al.* Mutations in KPTN cause macrocephaly, neurodevelopmental
delay, and seizures. *Am J Hum Genet* **94**, 87-94, doi:10.1016/j.ajhg.2013.10.001
(2014).
66 Peterson, M. D. & Titus, M. A. F-actin distribution of Dictyostelium myosin I
double mutants. *J Eukaryot Microbiol* **41**, 652-657, doi:10.1111/j.1550-
7408.1994.tb01529.x (1994).
67 Johnson Chacko, L. *et al.* Transcriptome-Wide Analysis Reveals a Role for
Extracellular Matrix and Integrin Receptor Genes in Otic Neurosensory
Differentiation from Human iPSCs. *Int J Mol Sci* **22**, doi:10.3390/ijms221910849
(2021).
68 Truter, S., Andrikopoulos, K., Di Liberto, M., Womack, L. & Ramirez, F. Pro-
alpha 2(V) collagen gene; pairwise analysis of the amino-propeptide coding
domain, and cross-species comparison of the promoter sequence. *Connect Tissue*
Res **29**, 51-59, doi:10.3109/03008209309061966 (1993).
69 Lévy, J. *et al.* EFNB2 haploinsufficiency causes a syndromic neurodevelopmental
disorder. *Clin Genet* **93**, 1141-1147, doi:10.1111/cge.13234 (2018).
70 Xiong, C. *et al.* Targeting Forward and Reverse EphB4/EFNB2 Signaling by a
Peptide with Dual Functions. *Sci Rep* **10**, 520, doi:10.1038/s41598-020-57477-x
(2020).
71 Zhu, F. *et al.* EFNB2 facilitates cell proliferation, migration, and invasion in
pancreatic ductal adenocarcinoma via the p53/p21 pathway and EMT. *Biomed*
Pharmacother **125**, 109972, doi:10.1016/j.biopha.2020.109972 (2020).

72   Ferrari, K. J., Lavarone, E. & Pasini, D. The Dual Role of EPOP and Elongin BC
1251   in Controlling Transcriptional Activity. *Mol Cell* **64**, 637-638,
1252   doi:10.1016/j.molcel.2016.11.009 (2016).
1253   73   Liefke, R., Karwacki-Neisius, V. & Shi, Y. EPOP Interacts with Elongin BC and
1254   USP7 to Modulate the Chromatin Landscape. *Mol Cell* **64**, 659-672,
1255   doi:10.1016/j.molcel.2016.10.019 (2016).
1256   74   Warburton, E. C. *et al.* cAMP responsive element-binding protein
1257   phosphorylation is necessary for perirhinal long-term potentiation and recognition
1258   memory. *J Neurosci* **25**, 6296-6303, doi:10.1523/jneurosci.0506-05.2005 (2005).
1259   75   Logan, T., Bendor, J., Toupin, C., Thorn, K. & Edwards, R. H. α-Synuclein
1260   promotes dilation of the exocytotic fusion pore. *Nat Neurosci* **20**, 681-689,
1261   doi:10.1038/nn.4529 (2017).
1262   76   Fournier, A. E., GrandPre, T. & Strittmatter, S. M. Identification of a receptor
1263   mediating Nogo-66 inhibition of axonal regeneration. *Nature* **409**, 341-346,
1264   doi:10.1038/35053072 (2001).
1265   77   Pérez-Otaño, I., Larsen, R. S. & Wesseling, J. F. Emerging roles of GluN3-
1266   containing NMDA receptors in the CNS. *Nat Rev Neurosci* **17**, 623-635,
1267   doi:10.1038/nrn.2016.92 (2016).
1268   78   Chodelkova, O., Masek, J., Korinek, V., Kozmik, Z. & Machon, O. Tcf7L2 is
1269   essential for neurogenesis in the developing mouse neocortex. *Neural Dev* **13**, 8,
1270   doi:10.1186/s13064-018-0107-8 (2018).
1271   79   Iwano, T., Masuda, A., Kiyonari, H., Enomoto, H. & Matsuzaki, F. Prox1
1272   postmitotically defines dentate gyrus cells by specifying granule cell identity over
1273   CA3 pyramidal cell fate in the hippocampus. *Development (Cambridge, England)*
1274   **139**, 3051-3062, doi:10.1242/dev.080002 (2012).
1275   80   Luo, C. *et al.* Single-cell methylomes identify neuronal subtypes and regulatory
1276   elements in mammalian cortex. *Science* **357**, 600-604,
1277   doi:10.1126/science.aan3351 (2017).
1278   81   Helbling, J.-C., Kinouchi, K., Trifilieff, P., Sassone-Corsi, P. & Moisan, M.-P.
1279   Combined Gene Expression and Chromatin Immunoprecipitation From a Single
1280   Mouse Hippocampus. *Current Protocols* **1**, e33,
1281   doi:https://doi.org/10.1002/cpz1.33 (2021).
1282   82   Doty, R. W., Fei, R., Hu, S. & Kavcic, V. Long-term reversal of hemispheric
1283   specialization for visual memory in a split-brain macaque. *Behavioural brain*
1284   *research* **102**, 99-113, doi:10.1016/s0166-4328(99)00006-6 (1999).
1285   83   Duboc, V., Dufourcq, P., Blader, P. & Roussigné, M. Asymmetry of the Brain:
1286   Development and Implications. *Annual review of genetics* **49**, 647-672,
1287   doi:10.1146/annurev-genet-112414-055322 (2015).
1288   84   Güntürkün, O., Ströckens, F. & Ocklenburg, S. Brain Lateralization: A
1289   Comparative Perspective. *Physiological reviews* **100**, 1019-1063,
1290   doi:10.1152/physrev.00006.2019 (2020).
1291   85   Hubel, D. H. & Wiesel, T. N. Receptive fields, binocular interaction and
1292   functional architecture in the cat's visual cortex. *The Journal of Physiology* **160**,
1293   106-154, doi:https://doi.org/10.1113/jphysiol.1962.sp006837 (1962).

86   Spiers, H. J. *et al.* Unilateral temporal lobectomy patients show lateralized
topographical and episodic memory deficits in a virtual town. *Brain : a journal of*
neurology **124**, 2476-2489, doi:10.1093/brain/124.12.2476 (2001).
87   Lynch, M. A. Long-term potentiation and memory. *Physiological reviews* **84**, 87-
136, doi:10.1152/physrev.00014.2003 (2004).
88   Achilly, N. P., Wang, W. & Zoghbi, H. Y. Presymptomatic training mitigates
functional deficits in a mouse model of Rett syndrome. *Nature* **592**, 596-600,
doi:10.1038/s41586-021-03369-7 (2021).
89   Kwapis, J. L. *et al.* Epigenetic regulation of the circadian gene Per1 contributes to
age-related changes in hippocampal memory. *Nat Commun* **9**, 3323,
doi:10.1038/s41467-018-05868-0 (2018).
90   David A Valientes, A. M. R., Autumn S Ivy. An Improved Method for Individual
Tracking of Voluntary Wheel Running in Pair-housed Juvenile Mice. *BIO-*
PROTOCOL **11**, doi:10.21769/bioprotoc.4071 (2021).
91   Long, J. Z. *et al.* A smooth muscle-like origin for beige adipocytes. *Cell*
metabolism **19**, 810-820, doi:10.1016/j.cmet.2014.03.025 (2014).
92   Deal, R. B. & Henikoff, S. A simple method for gene expression and chromatin
profiling of individual cell types within a tissue. *Developmental cell* **18**, 1030-
1040, doi:10.1016/j.devcel.2010.05.013 (2010).
93   Pfaffl, M. W. A new mathematical model for relative quantification in real-time
RT-PCR. *Nucleic Acids Res* **29**, e45, doi:10.1093/nar/29.9.e45 (2001).
94   Schwarz, J. M. Using fluorescence activated cell sorting to examine cell-type-
specific gene expression in rat brain tissue. *Journal of visualized experiments :*
JoVE, e52537, doi:10.3791/52537 (2015).
95   FastQC v. 0.11.9 (Babraham Bioinformatics, 2005).
96   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,
15-21, doi:10.1093/bioinformatics/bts635 (2013).
97   Picard v. 1.87 (Broad Institute, 2013).
98   R (R Foundation for statistical Computing, Vienna, Austria, 2010).
99   Lawrence, M. *et al.* Software for Computing and Annotating Genomic Ranges.
PLoS Computational Biology **9**, e1003118, doi:10.1371/journal.pcbi.1003118
(2013).
100 Rsamtools: Binary alignment (BAM), FASTA, variant call (BCF), and tabix file
import (Bioconductor Bioconductor 2021).
101 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**,
doi:10.1186/s13059-014-0550-8 (2014).
102 Durinck, S. *et al.* BioMart and Bioconductor: a powerful link between biological
databases and microarray data analysis. *Bioinformatics* **21**, 3439-3440,
doi:10.1093/bioinformatics/bti525 (2005).
103 Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt.
Nature protocols **4**, 1184-1191, doi:10.1038/nprot.2009.97 (2009).
104 Ggplot2: Elegant graphics for data analysis v. 3.3.4 (Springer International
Publishing, Cham, Switzerland, 2016).
105 gplots: Various R Programming Tools for Plotting Data v. 3.1.1 (Tal Galili 2020).

106   RColorBrewer: ColorBrewer Palettes (Erich Neuwirth, CRAN, 2014).

107   Venn Diagram: Calculate and draw custom Venn diagrams (BELGIUM, 2021).

108   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2.
1343   *Nature Methods* **9**, 357-359, doi:10.1038/nmeth.1923 (2012).

1344   109   Humburg, P., Helliwell, C. A., Bulger, D. & Stone, G. ChIPseqR: analysis of
1345   ChIP-seq experiments. *BMC Bioinformatics* **12**, 39, doi:10.1186/1471-2105-12-39
1346   (2011).

1347   110   TxDb.Mmusculus.UCSC.mm10.knownGene: Annotation package for TxDb
1348   object(s) v. 3.4.7 (Bioconductor 2019).

1349   111   Kent, W. J. *et al.* The Human Genome Browser at UCSC. *Genome Research* **12**,
1350   996-1006, doi:10.1101/gr.229102 (2002).

1351   112   Rosenbloom, K. R. *et al.* ENCODE whole-genome data in the UCSC Genome
1352   Browser. *Nucleic Acids Research* **38**, D620-D625, doi:10.1093/nar/gkp961
1353   (2010).

1354   113   Haeussler, M. *et al.* Navigating protected genomics data with UCSC Genome
1355   Browser in a Box. *Bioinformatics* **31**, 764-766,
1356   doi:10.1093/bioinformatics/btu712 (2014).

1357   114   ggpubr: 'ggplot2' Based Publication Ready Plots (CRAN, 2020).

1358