Single-nucleus RNA Sequencing and Spatial Transcriptomics Reveal the

Immunological Microenvironment of Cervical Squamous Cell Carcinoma

Zhihua Ou,[1,3,13] Shitong Lin,[2,4,13] Jiaying Qiu,[1,5,13] Wencheng Ding,[2,4,13] Peidi Ren,[1,3,13]
Dongsheng Chen,[1,13] Jiaxuan Wang,[1,3,13] Yihan Tong,[1,3,6] Di Wu,[1,3,7] Ao Chen,[1,8] Yuan Deng,[9]
Mengnan Cheng,[1,5] Ting Peng,[2,4] Haorong Lu,[1,10,11] Huanming Yang,[1,12] Jian Wang,[1,12] Xin
Jin,[1] Ding Ma,[2,4] Xun Xu,[1,11,*] Yanzhou Wang,[9,*] Junhua Li,[1,3,*] Peng Wu[2,4,*]

[1] BGI-Shenzhen, Shenzhen 518083, China.

[2] Cancer Biology Research Center (Key Laboratory of the Ministry of Education), Tongji
Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan
430000, China.

[3] Shenzhen Key Laboratory of Unknown Pathogen Identification, BGI-Shenzhen, Shenzhen
518083, China.

[4] Department of Gynecologic Oncology, Tongji Hospital, Tongji Medical College, Huazhong
University of Science and Technology, Wuhan 430000, China.

[5] College of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049,
China.

[6] College of Innovation and Experiment, Northwest A&F University, Yangling 712100, China.

[7] School of Basic Medicine, Qingdao University, Qingdao 266071, China.

[8] Department of Biology, University of Copenhagen, Copenhagen DK-2200, Denmark.

[9] Department of Obstetrics and Gynecology, Southwest Hospital, Third Military Medical
University, Chongqing 400038, China.

[10] China National GeneBank, BGI-Shenzhen, Shenzhen 518120, China.

[11] Guangdong Provincial Key Laboratory of Genome Read and Write, Shenzhen 518120,
China.

[12] James D. Watson Institute of Genome Sciences, Hangzhou 310058, China.

[13] These authors contributed equally

* Correspondence: pengwu8626@tjh.tjmu.edu.cn (P.W.), lijunhua@genomics.cn (J.L.),
w.y.z@foxmail.com (Y.W.), xuxun@genomics.cn (X.X.)

Keywords: cervical cancer; tumor microenvironment; cancer-associated fibroblasts; spatial
transcriptomics; single-nucleus RNA sequencing

Abstract

Effective treatment of advanced invasive cervical cancer remains challenging nowadays.
Herein, single-nucleus RNA sequencing (snRNA-seq) and SpaTial Enhanced REsolution
Omics-sequencing (Stereo-seq) technology are used to investigate the immunological

microenvironment of cervical squamous cell carcinoma (CSCC), a major type of cervical cancers. The expression levels of most immune checkpoint genes in tumor and inflammation areas of CSCC were not significantly higher than those in the non-cancer samples except for *LGALS9* and *IDO1*. Stronger signals of CD56+ NK cells and immature dendritic cells are found in the hypermetabolic tumor areas, while more eosinophils, immature B cells, and Treg cells are found in the hypometabolic tumor areas. Moreover, a cluster of cancer-associated fibroblasts (CAFs) are identified around some tumors, which highly expressed *ACTA2, POSTN, ITGB4,* and *FAP*. The CAFs might support the growth and metastasis of tumors by inhibiting lymphocyte infiltration and remodeling the tumor extracellular matrix. Furthermore, CAFs are associated with poorer survival probability in CSCC patients and might be present in a small fraction (~20%) of advanced cancer patients. Collectively, these findings might enhance understanding of the CSCC immunological microenvironment and shed some light on the treatment of advanced CSCC.

1. Introduction

Cervical cancer is the fourth most common cancer affecting women's health globally, especially in low- and middle-income regions.[1,2] Currently, over 12 types of human papillomaviruses (HPVs) are carcinogenic.[3] Among them, HPV16 is responsible for 60%-70% of the cervical cancer cases, especially cervical squamous cell carcinoma (CSCC). Since 2018, the World Health Organization (WHO) has called for the global elimination of cervical cancer, quantifying actions in vaccination, screening, and disease treatment/management,[4] which require joint efforts from different parties for decades.

Although early stage cervical cancer receiving radical hysterectomy can achieve a favorable prognosis, the 5-year overall survival rate or disease-free survival rate of advanced cervical cancer are unsatisfactory.[5,6] At present, chemotherapy (e.g., paclitaxel, cisplatin, bevacizumab, etc.) and radiotherapy remain the main palliative treatments for metastatic or recurrent patients, with a low response rate (48%) and a short survival period (17 months).[7–10] Immunotherapy brings new hope to treating incurable cervical cancer by reversing the exhausted or suppressed immune activities. Immune-checkpoint blockade (ICB) drugs targeting programmed cell death 1 (PD1), programmed cell death ligand 1 (PD-L1), and cytotoxic T lymphocyte antigen 4 (CTLA4) are currently under trials for recurrent/metastatic cervical cancers.[11,12] Unfortunately, the overall response rates to ICB therapy were low, varying from 4% to 26%.[11,13–15] Clarifying

the immune landscape of CSCC, especially the immunosuppression status in TME, may help

us better address this phenomenon and adjust our treating strategy for cervical cancers.

Single-cell sequencing and spatial transcriptomics are state-of-the-art tools to unravel the cell

heterogeneity and microenvironment of tumors, but applications of such techniques to CSCC

investigation remain limited. In this study, we collected cervical samples from 20 individuals

and combined single-nucleus RNA sequencing (snRNA-seq) and SpaTial Enhanced REsolution

Omics-sequencing (Stereo-seq) technology to investigate the immunological profiles of

CSCC.[16] Deciphering the immunological microenvironment of CSCC would provide new

insights into the treatment of advanced CSCC, which may accelerate cervical cancer

elimination.

2. Results

2.1 snRNA-seq data revealed the cellular composition of CSCC

To fully characterize the cell composition of cervical tissues, we collected CSCC samples from

5 patients for snRNA-seq (**Figure 1A and S1, Table S1**). A total of 67,003 cells and 30,996

genes passed quality control (**Figure 1B, Table S2**), from which we identified 14 cell types

based on canonical cell markers (**Figure 1B, Table S3**), including cancer cells (6,960),

columnar epithelial cells (CECs, 22,396), endothelial cells (6,340), smooth muscle cells (4,502),

fibroblasts (9,836), B cells (689), monocytes (5,281), T cells (4,930), regulatory T (Treg) cells

(1,081), plasma cells (3,236), myeloid dendritic cells (DCs) (955), plasmacytoid DCs (272),

mast cells (384) and natural killer (NK) cells (141). The uterine cervix contains two types of

cells lining its surface, with stratified squamous epithelial cells on the ectocervix and simple

columnar epithelial cells on the endocervix and crypts. The dysplasia of squamous epithelial

cells leads to CSCC. Therefore, the cancer cells mainly expressed a known CSCC-associated

gene *SERPINB3* (Serpin Family B Member 3),[17,18] tumor gene *TP63*, *CDKN2A,* and keratin

gene *KRT15* of squamous cells (**Figure 1C&D**). Since the tissues were mainly from advanced

cancer patients (FIGO Stage IB2-IIIC1; the tissue with no staging information was collected

during cervical biopsy before chemotherapy), few normal epithelial squamous cells were

isolated. Mapping of the snRNA-seq reads against high-risk HPV reference genomes revealed

the presence of viral genes in cancer cells (**Figure 1C, Table S2**). We also identified a big

cluster of columnar epithelial cells, which highly expressed *MUC5B* and *WFDC2*. This cell

type is mainly located in the endocervix epithelia, but it can also appear at the squamocolumnar

junction in the adult uterine cervix and some glands. Smooth muscle cells, fibroblasts, and

endothelial cells are the major cell types composing the cervical stroma. The smooth muscle cells highly expressed *MYH11, MYLK, ACTG2, COL3A1, and COL1A1*, fibroblasts expressed high levels of *LAMA2* besides *COL3A1 and COL1A1*, while endothelial cells can be distinguished by high expressions of *EMCN, FLT1,* and *EGFLT* (**Figure 1C&D**). Besides the structural cells of the cervix, diverse immune cell types were also identified, with monocytes (*ITGAX, MX4A7*), T cells (*CD3E, CD247*), and plasma cells (*MZB1, IGKC*) being the most abundant (**Figure 1B&C**). In short, the snRNA-seq data revealed the structural and immune cell composition of the CSCC tissues, facilitating our downstream spatial transcriptomic analysis with specific cellular gene expression profiles.

2.2 Spatial transcriptomic characterization of CSCC

Spatial information is critical in understanding cell-cell interactions in tissues, which unfortunately was missing in the snRNA-seq data. Therefore, we utilized Stereo-seq to acquire the in situ gene expression profiles.[16] The Stereo-seq chips contained capture probe contained a 25bp coordinate identity barcode, a 10bp molecular identity barcode, and a 22bp polyT tail for mRNA hybridization. Cervical samples from 2 non-cancer and 14 CSCC patients were obtained and embedded in OCT (**Table S1**). Serial cryosections of 10 μm thickness were dissected from each OCT block for Stereo-seq, hematoxylin and eosin (H&E) staining, and immunohistochemical (IHC) staining (**Figure S1A, Table S1**). Finally, a total of 18 Stereo-seq slides were successfully obtained (**Figure 2A, Table S4**). These included 3 slides from 2 non-cancer patients and 15 slides from 14 CSCC patients. Two patients contributed more than one sample. Because the samples were all from different anatomical sites, they were all included to compensate for the small sample size. The capture spots in Stereo-seq chips were 220 nm in diameter with a center-to-center distance of 500 nm between two adjacent spots. The capture spots were grouped into bins to include sufficient genes for accurate clustering. Our preliminary analysis revealed much higher RNA abundance in tumor areas than in stroma areas. To balance the expression differences between tumor and stroma, we annotated the CSCC Stereo-seq slides at bin100 (100 x 100 spots) to fully demonstrate the tissue composition, which would cover an area of approximately 49.72 x 49.72 μm. The mean numbers of genes per bin for the CSCC Stereo-seq slides ranged from 1,767 to 4,152 (**Table S4**). Because the three Stereo-seq slides from non-cancer patients had lower gene expression intensity than CSCC slides, they were annotated at bin200 (99.72 x 99.72 μm). Uniform manifold approximation and projection (UMAP) analysis showed that bin clusters of CSCC and non-cancer tended to dissociate from each other while those of CSCC displayed some convergence (**Figure S1B**).

Considering the complex structure of cancerous tissues, we manually conducted the initial annotations of Stereo-seq slides according to professional pathological assessments (based on H&E and IHC staining results) and marker gene expression patterns (**Figure S1C&D**). Six types of tissue clusters were generally identified in our CSCC samples, including tumor, stroma (without obvious inflammation), inflammation (stroma with diffuse inflammation or focal inflammation), gland, blood vessel, and necrosis. Tumor, stroma, and inflammation clusters were widely distributed among the Stereo-seq slides of the CSCC samples, with certain samples containing necrosis, glands, and blood vessels (**Figure 2A**). Depending on the gene expression profiles, the tissue clusters may be further divided into sub-clusters with number suffixes (**Figure S1C**). It should be kept in mind that a spatial cluster was virtually a mixture of cells in a defined area but was designated by its major characteristics. For example, the tumor clusters were not purely composed of cancer cells, but may also contain other cell types though in small numbers, such as immune cells, fibroblasts, etc. Tissue-specific genes also displayed spatial patterns. Cancerous squamous cell-associated genes such *KRT5, CDKN2A,* and *SERPINB3* were mainly enriched in the Stereo-seq tumor areas, *IGKC* and *IGLC2* enriched in inflammation areas, *VIM* in stromal areas, *ADRA2A* in blood vessels, and *MUC5B* in glands (**Figure S1D and S2**). All the 14 CSCC patients donating the 15 samples were HPV-positive, with 85.7% (12/14) infected by HPV16, 7.1% (1/14) by HPV33, and 7.1% (1/14) by HPV58. The HPV reads covered 8% - 100% (about 600bp - 7905bp) of the viral genome and were mainly identified in the tumor areas (**Table S4, Figure S3 and S4**). Spatial visualization demonstrated varied capture signals of viral genes in different tumor sites of the same sample, with E5, E6, E7, and L1 frequently observed (**Figure S4**). In contrast, only marginal HPV reads were identified in the non-CSCC samples (**Table S4**). In general, higher transcriptional and translational activities, cell proliferation, oxidative phosphorylation, and immune responses were observed in the Stereo-seq tumor areas than in the other areas (**Figure 2B**). The initial manual annotations of the Stereo-seq slides were used to assist downstream analysis of the TME.

2.3 Variable immune inhibition in CSCC

Regarding the low response rates of ICB therapy to cervical cancer,[10] we decided to scrutinize the immune landscape of CSCC for clues. The expression profiles of three gene sets with different immune functions, i.e., co-stimulatory, cytotoxic/effector, and co-inhibitory/exhaustion were evaluated in both snRNA-seq (**Figure 2C**) and Stereo-seq data (**Figure 2D, Table S5**). At the single-cell level, the co-stimulatory genes were expressed in

cells of both the innate and adaptive immunity, especially in Treg, T, and NK cells (**Figure 2C**).

Treg cells were found to highly express *CD27, CD28, CD40LG, ICOS, TNFRSF18, TNFRSF4,*

and *TNFRSF9.* While these genes are necessary for the maturation and normal suppressive

function of Treg cells, overexpression of *CD27* in Treg cells may restrain the anti-tumor

immune response.[19,20] Spatially, the co-stimulatory genes tended to be enriched in tumor and

inflammation areas (**Figure 2D**). Especially, *TNFRSF18 (*also named glucocorticoid-induced

TNF receptor, *GITR)* was commonly expressed in both inflammation and tumor areas. However,

its expression was also up-regulated in the epithelia of non-cancer samples. In our snRNA-seq

data, this gene was mainly detected in Treg, NK, T cells, and mast cells. Although *TNFRSF18*

was associated with the immune suppression by Treg cells in tumors,[21,22] its high spatial

expression level may be jointly contributed by multiple types of immunocytes. The immune

cytotoxic/effector genes were mainly expressed by T and NK cells, some of which were

commonly up-regulated in the inflammation and tumor regions in the Stereo-seq slides,

including *GNLY, GZMA, GZMB,* and *NKG7* (**Figure 2D**). These genes were mainly expressed

by NK cells (**Figure 2C**), indicating their important role in the cytotoxic response against CSCC.

For the co-inhibitory/exhaustion genes, we failed to detect any prominent expression of *CTLA4*

and *PD-1* in our Stereo-seq data (**Figure 2D**), although *CTLA4* can be highly expressed by Treg

cells and *PD-1* by Treg, T, and NK cells (**Figure 2C**). *PD-L1*, which was mainly expressed by

plasmacytoid DCs, was only overexpressed in the tumor or inflammation areas of a small

fraction of Stereo-seq samples. While *CD276, ENTPD1, IDO1, LGALS9, and VSIR* were

commonly detected in the Stereo-seq samples, only *IDO1* and *LGALS9* seemed to have higher

and wider expressions in the CSCC samples compared to the non-cancer samples (**Figure 2D**).

These two genes were both expressed by DCs (**Figure 2C**) and could downregulate cytotoxic

T cell activity.[23,24] Whether IDO1 and LGALS9 could be better targets for ICB therapy against

CSCC than CTLA4 and PD-L1 remains to be explored. Moreover, when we zoomed in to check

the immune genes in the same Stereo-seq slide, their expressions can vary greatly between

different tumor areas. In sample TJH08, the tumor areas commonly expressed high *IDO1*, low

PD-L1, and very low *CTLA4* (**Figure 2E**). In contrast, only one tumor area in sample TJH37

expressed these genes (**Figure 2E and S2**). Collectively, although both our snRNA-seq and

Stereo-seq data displayed evidence of immune exhaustion in CSCC patients, the immune

microenvironment varied considerably between and within patients.

2.4 Metabolic statuses of tumors were associated with different immune responses

Metabolism can modulate the immune microenvironment of tumors, which could be putative

intervention targets for cancer therapy.[25,26] We performed gene set variation analysis (GSVA)

separately on six pathways, including hypoxia, lactic acid, glycolysis, lipid metabolism, pentose

phosphate, and oxidative phosphorylation pathways. The mean of the GSVA scores for the

above six pathways was then calculated as the metabolic score of each tumor area. Based on

the GSVA metabolic score, the Stereo-seq tumor clusters ranked top 20 were categorized as

hypermetabolic tumors, with those ranked the last 20 as hypometabolic tumors (**Figure 2F**).

Generally, the hypermetabolic tumors displayed much higher activities in the oxidative

phosphorylation, glycolysis, and lactic acid pathway, indicating active aerobic glycolysis in

proliferating cancer cells, i.e., the Warburg effect. Moreover, the hypermetabolic tumors were

also accompanied by severe hypoxia and active lipid metabolism, suggesting intense oxidative

and nutrient stress in fast-growing tumors. To further explore the relationship between

metabolism and immune response, the GSVA signature scores for different immunocytes of

each bin within the hyper- and hypometabolic tumor areas were calculated. Based on the

average signature score in each tumor area, significant differences were observed for several

cell types (**Figure 2G**). CD56+ NK cells and immature dendritic cells showed much stronger

signals in the hypermetabolic tumor areas than in the hypometabolic ones, indicating that the

hypermetabolic tumors might be more prone to be associated with innate immune response. At

the same time, there were more eosinophils, immature B cells, and Treg cells in the

hypometabolic tumor areas, which might suggest ineffective adaptive immune response. Two

of the Stereo-seq samples, TJH34 and TJH35, contained both hyper- and hypometabolic tumor

areas. Using *TOP2A* and *MS4A1* as markers for cancer cells and B lymphocytes, respectively,

the spatial association between metabolic and B cell distribution was visualized. Disseminating

MS4A1 expression was detected within and outside the hypometabolic tumor areas, consistent

with the lymphocyte distribution pattern in H&E images (**Figure 2H**) and the GSVA signature

scores (**Figure 2G**).

2.5 Identification of a cluster of cancer-associated fibroblasts (CAFs) in CSCC

When exploring the immune differences between the hyper- and hypo-metabolic tumor areas

in sample TJH34, we noticed a unique spatial cluster outside the hypermetabolic tumor regions.

This cluster was different from most stromal clusters and looked like a ribbon enclosing the

tumor. As this cluster was part of the stroma, we closely scrutinized the fibroblasts in our

snRNA-seq data. Luckily, we identified a small set of fibroblasts derived from all five samples

(**Figure 3A**), which highly expressed reported marker genes for CAFs, including *ACTA2*,

POSTN, ITGB4, and *FAP* (**Figure 3B**). The CAFs had a lower stemness score than cancer cells and fibroblasts and were in various cell cycle stages (**Figure 3C**). Function enrichments based on the hallmark gene sets (MSigDB v7.4, https://www.gsea-msigdb.org/gsea/msigdb/) showed that CAFs shared common activities with both fibroblasts and cancer cells (**Figure 3D**). CAFs were involved in similar pathways to fibroblasts including UV response down, angiogenesis, myogenesis, and epithelial-mesenchymal transition. For pathways including the p53 pathway, KRAS signaling down, estrogen response, mitotic spindle, G2/M checkpoint, and E2F targets, CAFs showed similar activities to cancer cells. At the gene level, CAFs not only highly expressed marker genes for fibroblasts, such as the collagen protein family (*COL1A1, COL3A1, COL4A1, COL5A2, COL6A3*, etc.), but also marker genes for malignant squamous cells, such as *KRT4* and *KRT13* of the keratin family (**Figure 3E, Table S6**). Though CAFs in cancers might have different origins, the epithelial characteristics indicated that the origin of the CAFs in our CSCC samples might be associated with EMT.

To locate the spatial distribution of CAFs in CSCC tissues, we adopted the multimodal intersection analysis (MIA) approach developed by Moncada et al. to integrate snRNA-seq and Stereo-seq data.[27] Briefly, this method calculated the overlapping degree of the expression levels of cell type-specific genes identified by snRNA-seq data and the area-specific genes characterized by Stereo-seq data. The smaller the resultant p-value, which was mentioned as MIA score in our later description, the stronger the correlation between a defined cell type and a spatial area. Initial MIA results showed that our Stereo-seq clustering results complied with the expected cell composition in the corresponding areas (**Figure S5A**). Unfortunately, the MIA score alone cannot fully reflect the spatial specificity of cells, especially in areas with low RNA abundance. Therefore, a high expression level of *POSTN*, which was experimentally verified to be linked with CAFs,[28–30] and a high MIA score for CAFs were simultaneously utilized to define Stereo-seq clusters of CAFs (**Figure S5B**). Results showed that CAFs were enriched around some tumor areas in 4 out of the 15 Stereo-seq slides (**Figure 3F**), including the hypermetabolic tumor areas of sample TJH34 (**Figure 2H and 3F**). The existence of CAFs in CSCC was further confirmed by IHC staining of POSTN using serial tissue sections of the same samples (**Figure S5C**). Notably, not all the tumor areas were surrounded by CAFs, making us curious about the biological differences associated with the presence of these cells.

2.6 CAFs might facilitate the growth and metastasis of CSCC from diverse aspects

To comprehensively reveal the biological functions of CAFs in CSCC, we divided the tumor

areas in the Stereo-seq slides into two types: tumor areas surrounded by CAFs (CAFs+ tumors),

and tumor areas not surrounded by CAFs (CAFs- tumors). Three Stereo-seq slides were found

to contain both CAFs+ and CAFs- tumor areas and were used for downstream analysis. We

then used the up- and down-regulated DEGs between the CAFs+ and CAFs- tumor areas

($|Log_2FC| > 1.28$, $P < 0.05$) of the 3 samples to perform GO enrichment analysis (**Figure 3G**

and 3H, Table S7). Results showed that the CAFs+ tumors were more active in energy usage,

metabolism, mitosis, and cell growth than CAFs- tumors (**Figure 3I**). Meanwhile, cellular

adhesion, apoptosis, and immune response were down-regulated in CAFs+ tumors. The above

observations coincided with the immune and metabolic heterogeneity of CSCC (**Figure 2F&G**),

especially in sample TJH34 (**Figure 2H**). These indicated that the presence of CAFs might

support tumor progression from different aspects.

Next, we calculated the gene module expression scores of 993 individual bins regarding

immune gene sets to evaluate the immune cell abundances. Results showed significantly

reduced numbers of B cells, CD4 T cells, CD8 T cells, neutrophils, DCs, NK cells, and Th1

cells in CAFs+ tumors (**Figure 4A and S5D**), indicating that CAFs might act as a physical

barrier to prevent the infiltration of pro-immunity cells into tumor areas. While more tumor-

associated macrophages (TAM) were identified in CAFs+ tumor areas (**Figure 4A),** the

distribution of M1 (tumor-suppressive) and M2 (tumor-promoting) phenotypes showed the

opposite (**Figure S5E**).[31] Due to the small sample size and the weak signal of macrophages, we

were unsure about the relationship between CAFs and TAMs of different phenotypes.

As part of the stroma, CAFs will have to interact closely with cancer cells, stromal cells, and

immune cells. Indeed, analysis of the snRNA-seq data showed potential interactions between

CAFs and the other cells regarding extracellular matrix (ECM) formation and cell-cell contact

(**Figure 4B**). CAFs highly expressed genes of the collagen family, especially *COL1A1,*

COL1A2, COL4A1, COL4A2, COL4A5, COL6A1, COL6A2, and *COL6A3*) that might interact

with *CD44* expressed by immunocytes and smooth muscle cells, which may be involved in cell

adhesion and migration. The collagens might also interact with diverse members of the integrin

family expressed by the cancer cells, immune cells, and stromal cells. Similarly, *FN1*

(fibronectin 1, a soluble glycoprotein) and laminins (*LAMA2, LAMA3, LAMA4, LAMA5,*

LAMB1, LAMB2, LAMB3, LAMC1, LAMC3) expressed by CAFs might also interact with the

other cell types through integrins. The integrins are membrane receptor proteins made up of α

and β subunits and are involved in cell adhesion and recognition. CAFs seems to use different

heterodimeric forms of integrins to contact the other cell types. They might interact with cancer

cells through integrins composed of subunits α2β1, α3β1, and αvβ8, while interact with

endothelial cells, CECs, smooth muscle cells through integrins made up of subunits α9β1, α6β1,

and α1β1, and with T cells and NK cells through integrins made up by subunits α1β1 (**Figure**

4C). Importantly, CAFs might take advantage of F11R (also called JAM1, junctional adhesion

molecule 1) to form tight junctions with cancer cells and stromal cells through F11R and JAM3,

which might prevent the infiltration of immunocytes. CAFs might also express other matrix

proteins including *THBS1* (thrombospondin 1), *THBS2,* and *TNC* (tenascin C) to communicate

with immunocytes, smooth muscle cells, cancer cells, and plasma cells through *CD44*, integrin

(α3β1), and *SDC1*. Moreover, CAFs overexpressed several tissue remodeling factors (**Figure**

4D), including *POSTN* (periostin, a secreted ECM protein), *FAP* (fibroblast activation protein,

a serine protease), *MMP1* (matrix metalloproteinase 1), *TNC* (Tenascin-C, a matrix protein)*,*

and *LOXL1* (lysyl oxidase like 1, catalyzes the cross-linking of collagen and elastin). This

evidence suggested a critical role of CAFs in shaping the tumor extracellular environment.

Besides their role in ECM construction, CAFs might also enhance the stemness and

proliferation of cancer cells through overexpressing secreted factors including *SEMA3C,*

POSTN, CXCL6 (**Figure 4B**). *SEMA3C* was reported to promote cancer stem cell maintenance,

angiogenesis, and invasion.[32,33] *POSTN* can augment cancer cell survival by activating the

Akt/PKB pathway through integrins αvβ3.[34] It may also promote cancer growth through the

PTK7-Wnt/β-Catenin signally pathway.[29] *CXCL6* (C-X-C motif chemokine ligand 6), though

mainly related to immune response, was reported to promote the growth and metastasis of

esophageal squamous cell carcinoma (**Figure 4D**).[35] Another highly expressed gene in CAFs,

SNAI2 (Slug), a snail-related zinc-finger transcription factor, may inhibit apoptosis and promote

cancer progression.[36,37] Several common growth factors such as *TGFB1* (transforming growth

factor beta 1), *EGF* (epidermal growth factor), and *VEGFA* (vascular endothelial growth factor

A) were also expressed by CAFs (**Figure 4D**). Moreover, the upregulation of *LSD1* (histone

lysine demethylase 1) in CAFs might inhibit the IFN activation to evade immune attack.[38] The

Wnt5a signaling protein produced by CAFs might also suppress the immune response to

facilitate tumor metastasis (**Figure 4B**).[39,40] In summary, CAFs might be able to potentiate the

TME, promoting the progression of tumors.

2.7 The presence of CAFs was associated with poorer clinical statuses of CSCC

To verify the pro-tumorigenic effects of CAFs, we first performed survival analyses with a

dataset from the Cancer Genome Atlas (TCGA), which contained 252 CSCC patients.[41] The

GSVA score for CAFs was calculated for each CSCC patient using the marker gene set (*ACTA2*,

POSTN, ITGB4, and *FAP*) (**Figure 3B**). Not surprisingly, higher signals of CAFs predicted

unfavorable progression-free survival (HR = 1.66, 95%CI = 1.03-2.67, *P* = 0.038) and overall

survival (HR = 1.69, 95%CI= 1.00-2.84, *P* = 0.05) for CSCC patients (**Figure 4E**). Next, we

measured the expression levels of POSTN, a biomarker of CAFs, in the stroma and tumor

regions using an independent sample set, which consisted of 65 archived formalin-fixed

paraffin-embedded (FFPE) CSCC samples (**Figure 4F**). Based on the staining intensity and

distribution pattern, we assigned POSTN staining scores to each sample (see Method). The

overall positive (weak, moderate, and strong) rate of POSTN expression in the stroma was much

higher than that in the tumor regions (**Figure 4G**). However, only 21.1% (14/65) of the FFPE

samples had moderate or strong staining of POSTN in the stroma around tumor. Chi-square test

showed that higher expression levels of POSTN in the stroma were significantly correlated with

more advanced pathological stages, poorer differentiation, larger tumor size, higher squamous

cell antigen (SCC) concentrations in peripheral blood, and older age (**Table 1**), which further

confirmed the pro-tumorigenic ability of CAFs.

Collectively, our results indicated CAFs were a crucial component of the TME of CSCC,

forming a barrier to protect the cancer cells from immune surveillance and clearance. They

might also help stimulate cell proliferation and angiogenesis, inhibit apoptosis, and reconstruct

the ECM to enhance tumor metastasis (**Figure 4H**).

3. Discussion

Although vaccines and radical hysterectomy are effective measures in preventing and treating

cervical cancers, the treatment of recurrent/metastatic cervical cancers remains a big obstacle

to achieving the goal of cervical cancer elimination. Herein, we have characterized a high-

resolution immunological landscape of CSCC combining snRNA-seq and Stereo-seq

technology, which may facilitate the management and treatment of HPV-induced cervical

cancer.

Nowadays, ICB therapies, especially those using PD-L1/PD-1 and CTLA4 inhibitors, are

among the novel methods to treat metastatic cervical cancers. Several studies reported wide

expression of PD-L1 in cervical cancers, with positive rates ranging from 34% to 96%.[42–44]

However, PD-L1 expression alone wasn't associated with the disease outcome of cervical cancer patients.[42] Indeed, the response rates to PD-1/PD-L1 and CTLA-4 inhibitors fluctuated greatly among different trials and the efficacies seemed independent of the expression status of associated checkpoint genes.[10,45] In our study, the expression levels of most immune checkpoint genes in tumor and inflammation areas of CSCC were not significantly higher than those in the non-cancer samples except for *LGALS9* and *IDO1* (**Figure 2D**). LGALS9 (i.e., galectin 9) downregulates effector T cell immunity through binding to Tim-3 on the T cell surface or inhibiting the antigen-presenting ability of DCs.[23,24] While disruption of the galectin 9 signaling pathway was shown to induce tumor regression in mice harboring pancreatic ductal adenocarcinoma, reversed effect was reported in lung metastasis mouse models.[46,47] IDO1 is mainly expressed in DCs and helps degrade tryptophan into kynurenine, which suppresses T cell functions.[48] It is found that inhibition of IDO1 could enhance the radiosensitivity of HeLa and SiHa tumorsphere cells, indicating the potential application of IDO1 inhibitors as radiosensitizers.[49] Whether targeting LGALS9 and IDO1 can improve the treatment of CSCC requires further exploration. Our study also revealed suppressive adaptive immunity in tumor areas with low metabolic activity, highlighting the critical role of metabolic modulation in TME. Indeed, recent clinical trials have combined checkpoint inhibitors with metabolic agents targeting glucose, amino acid, and nucleotide metabolism.[50] A better understanding of the crosstalk between immune response and metabolism would further benefit cancer therapies.

CAFs have been characterized in multiple types of cancers and can be classified into diverse subtypes.[51] While cell line studies have implicated a supportive role of CAFs in the proliferation of cervical cancer cells,[52,53] this is the first study to systemically describe the spatial distribution and biological properties of pro-tumorigenic CAFs in clinical samples of CSCC. Due to tissue heterogeneity, the marker genes for CAFs varied among cancers. In this study, *ACTA2*, *POSTN*, *ITGB4*, and *FAP* were adequate marker genes to identify CAFs in CSCC (**Figure 4B**). Other genes such as *KRT4*, *ITGA1*, *COL24A1*, and *COL7A1* might serve as complementary marker genes for CAFs in CSCC (**Figure 4E, Table S7**), which displayed cellular properties of both fibroblasts and cancerous squamous cells. These CAFs contributed significantly to the heterogeneity of TME, which displayed a pro-tumorigenic phenotype by facilitating tumor growth, metastasis, and immune evasion. The CAFs+ tumors were active in proliferation but lacked lymphocyte infiltration. Exposing these immune-evasive tumors to the immune system is essential to eradicate the cancer cells. However, not all the CSCC samples were positive of CAFs. Only 28.6% (4/14) of the CSCC patients in the Stereo-seq experiment showed presence

of CAFs, and only 21.1% (14/65) of the FFPE samples in the IHC experiment were positive of
POSTN, a biomarker of CAFs. As CAFs tended to be associated with more advanced
pathological status (**Table 1**), chances are that they might function in invasive cancer patients
with high variation among individuals. Researchers have tried to interfere with the activation,
the action, and the normalization processes of CAFs using antibodies or inhibitors, with several
clinical trials ongoing.[54] Since genes highly expressed by CAFs are also essential to normal
tissues, their efficacies and side effects require close monitoring.

Several limitations exist in this study. 1) Samples for snRNA-seq and Stereo-seq were not
paired. We were only able to collect paired samples from one cancer patient. Individual and
anatomical heterogeneity may hinder the comprehensive annotation of cell types. For example,
CAFs were found in 4 Stereo-seq samples, but the number of CAFs was very small in snRNA-
seq data. 2) Since all the cervical cancer cases have progressed to invasive stages, most of the
tissues collected mainly contained invasive tumors, making it impossible to compare the intra-
individual difference between normal epithelia and tumors. 3) Unlike tumor cells, the gene
expression levels of immunocytes were relatively low, hindering the spatial analysis of most
immune cells at high resolution. 4) Our results were drawn from observations of a limited
number of clinical samples. Although we have tried to incorporate public data to validate the
pro-tumorigenic phenotype of CAFs, further experimental conformation is necessary.

In conclusion, our data have demonstrated the high heterogeneity of viral gene expression,
immune response, and metabolism in CSCC, indicating that combined drugs or therapies
targeting multiple biological processes would be better practice to treat CSCC. Interventions on
CAFs and tumor metabolism may complement the current treatments of CSCC. Further
investigations into these biological aspects may facilitate the development of new drugs or
therapies against CSCC and the other HPV-induced squamous cell carcinomas.

4. Experimental Section/Methods
4.1 Patients and samples
The cervical specimens were collected from 20 patients aged 38 to 69 by the Department of
Obstetrics and Gynecology of Tongji Hospital in Wuhan and the Department of Obstetrics and
Gynecology of Southwest Hospital in Chongqing. Based on colposcopy examination, 18
patients were diagnosed with CSCC (Stage IB1 to Stage IIIC1), 2 patients were diagnosed with
benign gynecological diseases but also required surgery (**Table S1, Figure S1A**). Carcinoma

staging was conducted based on the criteria of the FIGO staging system. The freshly-taken

samples were used for single-cell RNA sequencing and Stereo-seq experiments. A total of 65

archived FFPE samples were retrospectively obtained from the Department of Obstetrics and

Gynecology of Tongji Hospital in Wuhan to verify the presence of CAFs in CSCC.

4.2 Experiments

4.2.1 Single-nucleus RNA sequencing

The collected cervical tissues were quick-frozen with liquid nitrogen for 30 minutes and then

stored in a -80°C refrigerator. Nuclei isolation and permeabilization were performed under the

guidance of Chromium Next GEM Single Cell Multiome ATAC + Gene Expression User Guide

(CG000338). snRNA-seq libraries were prepared using the Chromium Single Cell 3′ Reagent

Kits v3 (10x Genomics, USA), according to the manufacturer's instructions. Briefly, high-

quality sequencing data was obtained after a series of experimental procedures including cell

counting and quality control, gel beads-in-emulsion (GEMs) generation and barcoding, post

GEM-RT cleanup, cDNA amplification, gene expression library construction, and NovaSeq

platform (Illumina, USA) sequencing.

4.2.2 Tissue preparation for spatial transcriptomic experiment

A tissue block with an edge length of less than 1cm was dissected from the surgically removed

tissues. The tissue block was then rinsed by cold PBS, immersed in the pre-cooled tissue storage

solution (Miltenyi Biotec, Germany), and then embedded with pre-cooled OCT (Sakura, USA)

in a -30°C microtome (Thermo Fisher, USA) within 30 minutes after surgery. Three to four

serial cryosections of 10 µm thickness were cut from the OCT-embedded samples for H&E

staining, Stereo-seq library preparation, and IHC staining. Brightfield images of the H&E

samples were taken with a Motic microscope scanner (Motic, China) for histopathological

assessment.

4.2.3 Quality control of RNA obtained from OCT-embedded samples

100-200 µm thick sections were cut from each OCT-embedded sample for total RNA extraction

using the RNeasy Mini Kit (Qiagen, USA) according to the manufacturer's protocol. RNA

integrity number (RIN) was determined by a 2100 Bioanalyzer (Agilent, USA). Only samples

with RIN≥7 were qualified for the transcriptomic study. All samples used had RIN of 7-10.

4.2.4 Stereo-seq library preparation and sequencing

The spatial transcriptomic RNA library was constructed using Stereo-seq capture chips (BGI-
Shenzhen, China), which had a size of 1 cm^2. The capture spots were 220 nm in diameter with
a center-to-center distance of 500 nm between each other. Each Stereo-seq capture probe
contained a 25bp coordinate identity barcode, a 10bp molecular identity barcode, and a 22bp
polyT tail for in situ mRNA hybridization.[16] A cryosection of 10 μm thickness cut from OCT-
embedded tissue was quickly placed on the chip, incubated at 37°C for 3 minutes, and then
fixed in pre-cooled methanol at -20°C for 40 minutes. The fixed tissue section was stained with
the Qubit ssDNA dye (Thermo Fisher, USA) to check tissue integrity before fluorescent
imaging. After that, the tissue section was permeabilized using 0.1% pepsin (Sigma, USA) in
0.01 N HCl buffer, incubated at 37°C for 14 minutes, and then washed with 0.1x SSC. RNA
released from the permeabilized tissue was reverse transcribed for 1 hour at 42°C. Later, the
tissue section was digested with tissue removal buffer at 42°C for 30 min. The cDNA-
containing chip was then subjected to cDNA release enzyme treatment overnight at 55°C. The
released cDNA was further amplified with cDNA HIFI PCR mix (MGI, China). Around 20ng
cDNA was fragmented to 400-600bp, amplified for 13 cycles, and purified to generate DNA
nanoball library, which was sequenced with the single-end 50+100bp strategy on an MGI
DNBSEQ sequencer (MGI, China).

4.2.5 Immunohistochemical (IHC) staining
The IHC staining for Ki67 and POSTN were performed under the manufactures' protocol. The
frozen sections dried at room temperature were placed in a 37°C oven for 10-20 minutes, fixed
with 4% paraformaldehyde for 20 minutes, and washed thrice with PBS (pH = 7.4) for 5 minutes.
The antigens were then repaired with EDTA (pH9.0) and the endogenous peroxidase was
blocked by 3% hydrogen peroxide. The slides were further blocked with 3% BSA (G5001-100g,
Servicebio) at room temperature for 30 minutes, and then incubated with Ki67 (ab16667,
Abcam, 1:200) or POSTN (ab215199, Abcam, 1:500) at 4°C overnight. Finally, the frozen
slices were subjected to secondary antibody blocking, DAB staining, nuclear re-staining, and
dehydration. The protein expression levels of Ki67 and POSTN were evaluated by professional
pathologists under a microscope. The expression scores of POSTN in tumor and stromal regions
of 65 FFPE samples were measured according to the positivity percentage (0-5% = 0, 5-25% =
1, 26-50% = 2, 51-75% = 3, >75% = 4) and staining intensity (negative = 0, weak = 1, moderate
= 2, strong =3) (**Figure 4F and 4G**). The final score was obtained by multiplying the two scores,
ranging from 0 to 12. Negative =0, weak = 1-4, moderate = 4-8, strong = 9-12 (**Table 1**).

4.3 Bioinformatic analysis

4.3.1 Cell type characterization using snRNA-seq data

Quality control and gene expression quantification of snRNA-seq data

Raw sequencing files were first processed using CellRanger version v6.0.2 (10x Genomics, USA) to obtain gene expression matrices. After cell calling, the droplets containing no cell were excluded based on the number of filtered unique molecular identifiers (UMIs) mapped to each cell barcode. Droplets with low-quality cells or more than one cell were also removed. To obtain a gene expression matrix optimized to individual samples, the R package scCancer v2.2.1 was employed to further filter the expression matrix.[55] The filtering thresholds were determined by catching outliers from the distribution of four quality spectra, including the number of total UMIs, the number of expressed genes, the percentages of UMIs from mitochondrial genes, and the percentages of UMIs from ribosomal genes. Besides filtering cells, genes expressed in less than three cells were also excluded to avoid false-positive results. The filtering thresholds for the five samples were documented in **Table S2**.

Cell type clustering using multi-sample snRNA-seq data

Integrative analysis of the snRNA-seq data from the five patients was carried out using the IntegrateData function of Seurat v4.[56] Further analysis included normalization, log-transformation, highly variable genes identification, dimension reduction, clustering, and differential expression analysis were all conducted using default parameters of Seurat except that *dims* was set as 1:30. Initially, a total of 35 cell clusters were obtained (69,312 cells with 30,996 genes). To ensure reliable identification, we removed the cell clusters consisting of less than two samples and with less than 15 cells per sample. Finally, 14 cell clusters (67,003 cells with 30,996 genes) were determined based on reported cell type marker genes (**Table S3**).

Analysis of differentially expressed genes (DEGs)

Expression of each gene in each cluster was compared against the rest of the clusters using the Wilcoxon rank-sum test with the FindAllMarkers function of Seurat v4.[56] Significantly up- or down-regulated genes were identified using the following criteria: 1) the difference in gene expression level was >1.28 fold unless explicitly noted; 2) genes were expressed by more than 25% of the cells belonging to the target cluster. 3) the adjusted p-value was less than 0.05.

4.3.2 Processing and annotation of Stereo-seq data

Preliminary processing of Stereo-seq data

Stereo-seq raw data were automatically processed using the BGI Stereomics analytical pipeline

(http://stereomap.cngb.org/), where the reads were decoded, trimmed, deduplicated, and

mapped against the human and HPV reference genomes. The reference genomes were: Human,

GRCh38.p12; HPV16:K02718.1; HPV18: EF202147.1; HPV33: M12732.1; HPV58:

D90400.1. Data of the chip area covered by tissue was extracted based on the ssDNA and H&E

staining images using the Lasso function of the BGI Stereomics website. It's worth noting that

tumor sites usually had much higher overall mRNA levels than the other anatomical areas,

leading to a significant imbalance of transcriptomic signals between the tumor areas and the

other sites on the Stereo-seq slides. Therefore, to fully reflect the spatial transcriptomic

landscape around the tumor areas, a bin size of 100 (100 spots x 100 spots, i.e., 49.72 x 49.72

μm) was used as the analytical unit for the annotation of CSCC Stereo-seq slides, while a bin

size of 200 (200 spots x 200 spots, i.e., 99.72 x 99.72 μm) was used for the non-CSCC samples.

The downloaded data was then processed with Seurat v4.[56] We used the criteria of >200 UMIs

per bin to remove bins with low expression signals. The data were then normalized using the

SCTransform function. Dimension reduction was conducted with principal PCA. Unsupervised

clustering of bins was performed with UMAP. Sequencing and analytical details can be found

in **Table S4**.

Annotation of bin clusters in Stereo-seq slides

The bin clusters were annotated based on the in situ expression patterns of marker genes

combining H&E and IHC staining results. The spatial expression patterns of genes in Stereo-

seq slides (**Figure S1 and S2**) were conducted with the SpatialFeaturePlot function of Seurat

v4.[56] The H&E and IHC images were examined by professional pathologists to determine the

tissue types. The annotated Stereo-seq areas were confirmed to be consistent with the H&E and

IHC assessment and marker gene expression patterns.

Identification of viral RNA

The viral reads were mapped against HPV reference genomes with BWA. The genome

coverage (covered length/full length of the reference genome) and effective depth (total mapped

bases/covered length) for each type were calculated. Only samples with a viral genome

coverage of >5% and an effective depth of >50x were deemed as HPV positive (**Table S2&S4,**

Figure S3).

4.3.3 Characterization of the immunity, hypoxia, and metabolism statuses in CSCC

Signature enrichment analysis of Stereo-seq clusters

In enrichment analysis, the expression scores of signature genes were calculated for individual

bins using the AddModuleScore (on log-normalized data) of Seurat v4 with default

parameters.[56] Pathways and cell types included in the enrichment analysis, with the

corresponding reference for gene signatures, were listed as follows: hypoxia,[57] glycolysis,[58]

lipid metabolism,[59] lactic acid metabolism, oxidative phosphorylation (MSigDB,

https://www.gsea-msigdb.org/gsea/msigdb/), pentose phosphate pathway,[60] macrophages,[61]

and other immune cells (**Figure 2F and 2G**).[62]

Prediction of the spatial distribution of immunocytes

Using the gene signatures of immunocytes as input, we calculated the cell type scores of each

bin in tumor areas using the AddModuleScore of Seurat v4.[56] Combining with the spatial

coordinate of bins, the possible spatial distribution of the corresponding cell type was obtained

(**Figure S5D**).

4.3.4 Characterization of CAFs

Signature enrichment analysis of fibroblasts, CAFs, and cancer cells with snRNA-seq data

Expression scores of signature genes from MSigDB v7.4 (https://www.gsea-

msigdb.org/gsea/msigdb/) were calculated for individual cells using the AddModuleScore

function (on log-normalized data) of Seurat v4 with default parameters to assess differential

pathways in fibroblasts, CAFs, and cancer cells.[56]

Cell cycle and stemness analysis of fibroblasts, CAFs, and cancer cells with snRNA-seq data

We used the AddModuleScore function of Seurat v4 to calculate the relative average

expression of a list of G2/M and S phase markers to obtain the cell cycle scores.[63] Cell

stemness analysis was conducted using the OCLR model and the stemness signatures

embedded in the scCancer package.[55,64]

Multimodal intersection analysis (MIA)

To integrate snRNA-seq and Stereo-seq data, we calculated the overlapping degree of the

expression levels of cell type-specific genes identified by snRNA-seq data and the area-specific

genes characterized by Stereo-seq data using the MIA approach.[27] The lower the p-value, the

higher overlapping between a certain cell type and a Stereo-seq area. MIA was conducted to

confirm the consistency between cell types and Stereo-seq annotated areas and to identify the Stereo-seq areas composed of CAFs.

DEG and GO enrichment analysis of CAFs+ and CAFs- tumor areas

Expression of each gene in CAFs+ cluster was compared against that in the CAFs- clusters of the same Stereo-seq chip using the Wilcoxon rank-sum test with the FindMarkers function of Seurat v4.[56] Significantly up- or down-regulated genes (**Figure 3G**) were identified using the following criteria: 1) the difference in gene expression level was >1.28 fold unless explicitly noted; 2) genes were expressed by more than 25% of the bins belonging to the target cluster. 3) the adjusted p-value was less than 0.05. GO enrichment analysis was conducted using Metascape with default settings (**Figure 3I**).[65]

Cell-cell communication between CAFs and the other cell types in CSCC tissues

To understand the communication network between CAFs and the other cell types, we conducted cell-cell communication with CellChat with the snRNA-seq data to obtain the ligand-receptor pairs regulated by CAFs.[66] The probability of cell–cell communication was estimated by integrating gene expression with prior knowledge of the interactions between signaling ligands, receptors and their cofactors.

Prognostic analysis of CAFs with TCGA data

The gene expression profiles of CSCC were downloaded from The Cancer Genome Atlas (TCGA) (https://portal.gdc.cancer.gov/) with the latest follow-up prognostic information obtained from an integrated clinical data resource. We calculated a signature score of CAFs for each CSCC patient with GSVA using the marker genes of CAFs (*ACTA2*, *POSTN*, *ITGB4,* and *FAP*). Based on the median GSVA score, the patients were then divided into two groups: high CAFs v.s. low CAFs. The Kaplan-Meier overall survival and progression-free survival curves were generated with GraphPad Prism 6 (**Figure 4E**).

4.4 Statistical analysis and plotting

Statistical analyses, including Student's t-test, Wilcoxon's rank-sum test, Wilcoxon signed-rank test, and Chi-square test were performed in R 3.6.0. Asterisks indicate the significance levels of p-values: ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$. The schematic plots of CAFs were created with BioRender (https://biorender.com/).

Ethical statement

This study was reviewed and approved by the Medical Ethics Committee of Tongji Medical College, Huazhong University of Science and Technology (TJ-IRB20210609), Southwest Hospital, Third Military Medical University (KY2020142), and the Institutional Review Board of Beijing Genomics Institute, Shenzhen, China (BGI-IRB 21050).

Data and code availability

The data and scripts supporting the findings of this study have been deposited into CNSA (CNGB Sequence Archive) of CNGBdb (https://db.cngb.org/cnsa/, accession numbers to be updated) and are available upon reasonable request. The dataset used to verify the distinguishing ability of the DEGs between preinvasive and invasive cancerous lesions was obtained from the Gene Expression Omnibus (GEO) database (www.ncbi.nlm.nih.gov/geo) by the Accession Number of GSE63514.

Author contributions

Z. Ou, P. Wu, J. Li, W. Ding, and D. Chen designed this study. P. Wu, Z. Ou, J. Li, X. Xu, D. Ma, X. Jin, H. Yang, and J. Wang supervised this study. P. Wu, Y. Wang, W. Ding, and S. Lin coordinated sample collection. S. Lin, Y. Ding, and T. Peng conducted snRNA-seq and IHC experiments. P. Ren, Y. Tong, and D. Wu performed Stereo-seq library construction and H&E staining. A. Chen and M. Cheng provided technical support for Stereo-seq experiments. H. Lu conducted sequencing for Stereo-seq libraries. J. Qiu, J. Wang, Y. Tong, and D. Wu conducted data analysis. Z. Ou, S. Lin, P. Ren, J. Qiu, and J. Wang wrote the original manuscript. Z. Ou, S. Lin, P. Wu, J. Li, W. Ding, and D. Chen reviewed and polished the manuscript.

Declarations of interest

A. Chen and M. Cheng are applying for patents covering the chip, procedure, and applications of Stereo-seq. The other authors declare that they have no competing interests.

Acknowledgements

This work received funding from the National Key R&D Program of China (2021YFC2701201), the National Natural Science Foundation of China (82072895, 82141106, 82103134) and the Guangdong Provincial Key Laboratory of Genome Read and Write (2017B030301011). We thank China National GeneBank for providing sequencing services for this project. The authors

680 also would like to thank Miss Feiyun Xuanyuan and Mr. Geer Xuanyuan for their inspirational

681 communications.

682
683 **Tables and Figures**

684 **Table 1. Stromal POSTN expression characteristics in 65 FFPE CSCC samples.**

Characteristic	IHC score >=4	IHC score <4	p
Number of patients	24 (36.9%)	41 (63.1%)	
Pathologic stage, n (%)			**0.012**
I	5 (17.2%)	24 (82.8%)	
II	10 (50.0%)	10 (50.0%)	
III	9 (56.2%)	7 (43.8%)	
Lymph node metastasis, n (%)			0.401
N0	13 (31.0%)	29 (69.0%)	
N1	4 (50.0%)	4 (50.0%)	
N1+	6 (50.0%)	6 (50.0%)	
Differentiation, n (%)			**0.007**
Low	16 (57.1%)	12 (42.9%)	
Well/Moderate	8 (21.6%)	29 (78.4%)	
Tumor size, n (%)			**0.018**
<4 cm	10 (29.4%)	24 (70.6%)	
>=4 cm	10 (71.4%)	4 (28.6%)	
Age, n (%)			**0.042**
<=50	7 (22.6%)	24 (77.4%)	
>50	17 (50.0%)	17 (50.0%)	
SCC, n (%)			**0.030**
<=1.5 ng/ml	3 (15.0%)	17 (85.0%)	
>1.5 ng/ml	16 (48.5%)	17 (51.5%)	

685 IHC: immunohistochemistry; SCC: squamous cell carcinoma antigen.

Figure 1. The cellular composition of CSCC tissues. (A) Workflow of snRNA-seq and
Stereo-seq experiments applied to cervical tissues. n indicates the number of samples. (B)
UMAP of cells identified from the snRNA-seq data of five CSCC tissues (left). The cell number
and proportion of each cell type from each sample (right). (C) Expression of selected marker
genes and HPV genes in the major cell types of CSCC tissues. (D) Expression matrix of cell-
type marker genes in the 14 cell types isolated from CSCC tissues.

Figure 2. Transcriptomic analysis of the immune heterogeneity in CSCC. (A) Distribution of Stereo-seq clusters in the 18 cervical samples. CSCC (n=15): tumor, stroma, inflammation,

gland, blood vessel, and necrosis; non-cancer (n=3, TJH10, TJH11 and TJH15): epithelia and stroma. (B) Gene ontology (GO) enrichment of major Stereo-seq clusters. (C) Heatmap showing the expressions of gene sets associated with different immune functions (co-stimulatory, cytotoxic/effector, and co-inhibitory/exhaustion) in cell types identified by snRNA-seq data. (D) Heatmap showing the expressions of gene sets associated with different immune functions (co-stimulatory, cytotoxic/effector, and co-inhibitory/exhaustion) in Stereo-seq clusters. (E) Spatial expression of selected immune checkpoint genes (*PD-L1, CTLA4,* and *IDO1*) in two representative Stereo-seq samples. The gene tended to be enriched in tumor areas. (F) Top, metabolic scores corresponding to the tumor areas of the 15 CSCC samples. Bottom, heatmap showing the GSVA scores of hypoxia, lactic acid, glycolysis, lipid metabolism, pentose phosphate, and oxidative phosphorylation pathways for tumor areas from each Stereo-seq sample. (G) Violin plots showing abundance of CD56+ NK cells, eosinophils, immature B cells, immature dendritic cells and Treg cells in hyper- and hypo-metabolic tumors. The Y axis shows the GSVA scores for each cell type. The p-values were determined by the Wilcoxon signed-rank test: ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$. (H) Spatial expression of marker genes for proliferating cells (*TOP2A*) and B cells (*MS4A1*) in representative two Stereo-seq slides. In the HE images, the blue line indicates the approximate border between hyper- and hypometabolic tumor areas, while the red arrow indicates focal lymphocyte infiltration.

Figure 3. Identification and spatial characterization of cancer-associated fibroblasts (CAFs) in CSCC. (A) UMAP of 9,836 fibroblasts. CAFs were outlined in red. (B) Violin plot showing the expression levels of *ACTA2, POSTN, ITGB4,* and *FAP* in the 10 clusters of

fibroblasts. (C) Box plot showing the stemness and cell cycle scores of fibroblasts, CAFs, and

cancer cells. The p-values were determined by the Wilcoxon signed-rank test: ns, not

significant; *, p <0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001. (D) GSVA results of

fibroblasts, CAFs, and cancer cells. (E) DEGs identified in fibroblasts, CAFs, and cancer cells.

(F) Spatially projected CAFs in representative Stereo-seq slides. The projected area was

determined based on the MIA score of CAFs, *POSTN* expression pattern, and the IHC staining

results of POSTN (see **Figure S5B&C**). (G) Venn map showing the number of DEGs identified

in the three Stereo-seq samples with both CAFs+ and CAFs- tumors. (H) Bar plot showing the

numbers of up- and down-regulated genes in three Stereo-seq samples. The CAFs+ tumors were

compared to the CAFs- tumors. (I) Dot plot of enriched GO terms for up- and down-regulated

DEGs identified in H.

Figure 4. Functional analysis of CAFs. (A) Box plot showing the abundance of immune cells in CAFs+ and CAFs- tumors in three samples. The p-values were determined by the Student's t-test: ns, not significant; *, p <0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001. (B) Ligand-

receptor communication network between CAFs and different cervical cells predicted by
snRNA-seq data. The cell–cell communication probability was estimated by integrating gene
expression with prior knowledge of the interactions between signaling ligands, receptors and
their cofactors. Right, heatmap of the top predicted ligands expressed by CAFs. Middle,
heatmap of ligand-receptor pairs between CAFs and different cell types in CSCC. Bottom,
expression heatmap of top receptors regulated by CAFs in different cell types. CECs, columnar
epithelial cells; ECM, extracellular matrix. (C) Bar plot showing the integrin types involved in
potential communications between CAFs and the other cell types. (D) Expression heatmap of
gene sets related to functions of CAFs in tumor development in snRNA-seq data. (E) The
progression-free and overall survival probabilities of CSCC patients estimated based on the
signature scores of the marker gene set for CAFs. The analysis was conducted using the TCGA
dataset. (F) Representative IHC staining patterns of POSTN in the stroma adjacent to tumor
areas in FFPE CSCC samples. (G) Bar plot showing the IHC staining intensities of POSTN
within the tumor area or in the stroma area around tumors in 65 FFPE CSCC samples. (H)
Schematic summary of the possible functions of CAFs in CSCC.
752
753
754

Supplementary Figures

Figure S1. Experimental details and the annotation process of Stereo-seq slides. (A)
Clinical characteristics and experimental details of cervical samples from CSCC and non-
cancer patients. (B) UMAP of Stereo-seq bins from 18 cervical samples. (C) Annotation results
of 2 representative Stereo-seq slides. (D) Expression of tissue-specific genes in 2 representative
Stereo-seq slides. The outline color indicates different tissue types: purple, tumor; blue, stroma
with inflammation; green, stroma; red, blood vessel; brown, gland. Violin plots display the gene
expression levels in the Stereo-seq areas.

765 **Figure S2 (To be continued). Annotation results of the 18 Stereo-seq slides.** Non-cancer

766 samples included TJH10, TJH11, and TJH15. All the other samples were CSCC samples. Six

767 genes were selected to represent different tissue types. Cancer cells, *CDKN2A*; Endothelial cells,

768 *PECAM1*; Fibroblasts, *VIM*; Smooth muscle cells, *ADRA2A*; Plasma cells, *IGKC*; Columnar

769 epithelial cells, *MUC5B*.

Figure S2 (Continued)

Figure S3. HPV reads in the Stereo-seq sequencing data of 15 CSCC tissues. (A) Mapping of HPV reads against the corresponding reference genome. (B) Schematic plot showing the genomic arrangement of the HPV genes in a linear form.

Figure S4. Expression of HPV genes in the tumor areas of selected CSCC Stereo-seq slides.

779 The annotation result for each sample can be found in **Figure S2**.

Figure S5. Characterization of CAFs. (A) Correlation between snRNA-seq cell types and the

Stereo-seq areas defined by multimodal intersection analysis (MIA). (B) Expression of *POSTN*

in Stereo-seq clusters and the associated MIA scores for CAFs. (C) Spatial clustering of CAFs

in Stereo-seq slides and the IHC staining of POSTN in corresponding serial sections of CSCC

samples. (D) Spatial prediction of immunocytes in Stereo-seq slides. (E) The abundance of macrophages including M1 (tumor-suppressive phenotype macrophage), (tumor-promoting phenotype macrophage) M2, and TAM (tumor-associated macrophage) in CAFs+ and CAFs- tumors. Asterisks indicate the significance levels of p-values: *, p <0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001.

Supplementary Tables (see Excel files)

Table S1. Clinical characteristics and experimental details for samples.

Table S2. snRNA-seq data statistics.

Table S3. Marker genes for the annotation of cell types in CSCC.

Table S4. Stereo-seq data statistics.

Table S5. Expression matrix of immune genes in Stereo-seq areas.

Table S6. DEGs for fibroblasts, CAFs, and cancer cells.

Table S7. DEGs for CAFs+ and CAFs- tumors.

References

1. Arbyn, M. *et al.* Estimates of incidence and mortality of cervical cancer in 2018: a worldwide analysis. *The Lancet Global Health* **8**, e191–e203 (2020).

2. Sung, H. *et al.* Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA A Cancer J Clin* **71**, 209–249 (2021).

3. de Sanjose, S. *et al.* Human papillomavirus genotype attribution in invasive cervical cancer: a retrospective cross-sectional worldwide study. *The Lancet Oncology* **11**, 1048–1056 (2010).

4. WHO. A Global Strategy for elimination of cervical cancer - PAHO/WHO | Pan American Health Organization. http://www.paho.org/en/topics/cervical-cancer/global-strategy-elimination-cervical-cancer (2018).

5. Friedlander, M. & Grogan, M. Guidelines for the Treatment of Recurrent and Metastatic Cervical Cancer. *The Oncologist* **7**, 342–347 (2002).

6.   Gupta, S. *et al.* Neoadjuvant Chemotherapy Followed by Radical Surgery Versus

Concomitant Chemotherapy and Radiotherapy in Patients With Stage IB2, IIA, or IIB

Squamous Cervical Cancer: A Randomized Controlled Trial. *JCO* **36**, 1548–1555 (2018).

7.   Tewari, K. S. *et al.* Bevacizumab for advanced cervical cancer: final overall survival and

adverse event analysis of a randomised, controlled, open-label, phase 3 trial (Gynecologic

Oncology Group 240). *The Lancet* **390**, 1654–1663 (2017).

8.   Bhatla, N., Aoki, D., Sharma, D. N. & Sankaranarayanan, R. Cancer of the cervix uteri.

Int J Gynecol Obstet **143**, 22–36 (2018).

9.   Tewari, K. S., Java, J. J., Gatcliffe, T. A., Bookman, M. A. & Monk, B. J. Chemotherapy-

induced neutropenia as a biomarker of survival in advanced ovarian carcinoma: An

exploratory study of the Gynecologic Oncology Group. *Gynecologic Oncology* **133**, 439–

445 (2014).

10. Orbegoso, C., Murali, K. & Banerjee, S. The current status of immunotherapy for cervical

cancer. *Reports of Practical Oncology & Radiotherapy* **23**, 580–588 (2018).

11. Chung, H. C. *et al.* Efficacy and Safety of Pembrolizumab in Previously Treated

Advanced Cervical Cancer: Results From the Phase II KEYNOTE-158 Study. *JCO* **37**,

1470–1478 (2019).

12. Attademo, L. *et al.* Immunotherapy in cervix cancer. *Cancer Treatment Reviews* **90**,

102088 (2020).

13. Frenel, J.-S. *et al.* Safety and Efficacy of Pembrolizumab in Advanced, Programmed

Death Ligand 1–Positive Cervical Cancer: Results From the Phase Ib KEYNOTE-028

Trial. *JCO* **35**, 4035–4041 (2017).

14. Santin, A. D. *et al.* Phase II evaluation of nivolumab in the treatment of persistent or

recurrent cervical cancer (NCT02257528/NRG-GY002). *Gynecologic Oncology* **157**,

161–166 (2020).

15. Naumann, R. W. *et al.* Safety and Efficacy of Nivolumab Monotherapy in Recurrent or

Metastatic Cervical, Vaginal, or Vulvar Carcinoma: Results From the Phase I/II

CheckMate 358 Trial. *JCO* **37**, 2825–2834 (2019).

16. Chen, A. *et al.* Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA

nanoball patterned arrays. *BioRxiv* (2021) doi:10.1101/2021.01.17.427004.

17. Kato, H. & Torigoe, T. Radioimmunoassay for tumor antigen of human cervical

squamous cell carcinoma. *Cancer* **40**, 1621–1628 (1977).

18. Sun, Y., Sheshadri, N. & Zong, W.-X. SERPINB3 and B4: From biochemistry to biology.

Seminars in Cell & Developmental Biology **62**, 170–177 (2017).

19. Muth, S., Klaric, A., Radsak, M., Schild, H. & Probst, H. C. CD27 expression on Treg

cells limits immune responses against tumors. *J Mol Med* (2021) doi:10.1007/s00109-

021-02116-9.

20. Tanaka, A. & Sakaguchi, S. Regulatory T cells in cancer immunotherapy. *Cell Res* **27**,

109–118 (2017).

21. Vence, L. *et al.* Characterization and Comparison of GITR Expression in Solid Tumors.

Clin Cancer Res **25**, 6501–6510 (2019).

22. Zhu, M. M. T. *et al.* Evaluation of glucocorticoid-induced TNF receptor (GITR)

expression in breast cancer and across multiple tumor types. *Mod Pathol* **33**, 1753–1763

(2020).

23. Wang, M. *et al.* Exosomal LGALS9 in the cerebrospinal fluid of glioblastoma patients

suppressed dendritic cell antigen presentation and cytotoxic T-cell immunity. *Cell Death*

Dis **11**, 896 (2020).

24. Zhu, C. *et al.* The Tim-3 ligand galectin-9 negatively regulates T helper type 1 immunity.

Nat Immunol **6**, 1245–1252 (2005).

25. Riera-Domingo, C. *et al.* Immunity, Hypoxia, and Metabolism–the Ménage à Trois of

Cancer: Implications for Immunotherapy. *Physiological Reviews* **100**, 1–102 (2020).

26. Samec, M. *et al.* Flavonoids Targeting HIF-1: Implications on Cancer Metabolism.

Cancers **13**, 130 (2021).

27. Moncada, R. *et al.* Integrating microarray-based spatial transcriptomics and single-cell

RNA-seq reveals tissue architecture in pancreatic ductal adenocarcinomas. *Nat Biotechnol*

38, 333–342 (2020).

28. Qin, X. *et al.* TGFβ3-mediated induction of Periostin facilitates head and neck cancer

growth and is associated with metastasis. *Sci Rep* **6**, 20587 (2016).

29. Yu, B. *et al.* Periostin secreted by cancer-associated fibroblasts promotes cancer stemness

in head and neck cancer by activating protein tyrosine kinase 7. *Cell Death Dis* **9**, 1082

(2018).

30. Yue, H. *et al.* Stromal POSTN induced by TGF-β1 facilitates the migration and invasion

of ovarian cancer. *Gynecologic Oncology* **160**, 530–538 (2021).

31. Mantovani, A., Marchesi, F., Malesci, A., Laghi, L. & Allavena, P. Tumour-associated

macrophages as treatment targets in oncology. *Nat Rev Clin Oncol* **14**, 399–416 (2017).

32. Peacock, J. W. *et al.* SEMA 3C drives cancer growth by transactivating multiple receptor

tyrosine kinases via Plexin B1. *EMBO Mol Med* **10**, 219–238 (2018).

33. Zhang, X., Klamer, B., Li, J., Fernandez, S. & Li, L. A pan-cancer study of class-3

semaphorins as therapeutic targets in cancer. *BMC Med Genomics* **13**, 45 (2020).

34. Bao, S. *et al.* Periostin potently promotes metastatic growth of colon cancer by

augmenting cell survival via the Akt/PKB pathway. *Cancer Cell* **5**, 329–339 (2004).

35. Zheng, S. *et al.* CXCL6 fuels the growth and metastases of esophageal squamous cell

carcinoma cells both in vitro and in vivo through upregulation of PD-L1 via activation of

STAT3 pathway. *J Cell Physiol* **236**, 5373–5386 (2021).

36. Blanco-Gómez, A. *et al.* Stromal SNAI2 Is Required for ERBB2 Breast Cancer

Progression. *Cancer Res* **80**, 5216–5230 (2020).

37. Cobaleda, C., Pérez-Caro, M., Vicente-Dueñas, C. & Sánchez-García, I. Function of the

Zinc-Finger Transcription Factor *SNAI2* in Cancer and Development. *Annu. Rev. Genet.*

41, 41–61 (2007).

38. Sheng, W. *et al.* LSD1 Ablation Stimulates Anti-tumor Immunity and Enables

Checkpoint Blockade. *Cell* **174**, 549-563.e19 (2018).

39. Asem, M. *et al.* Host Wnt5a Potentiates Microenvironmental Regulation of Ovarian

Cancer Metastasis. *Cancer Res* **80**, 1156–1170 (2020).

40. Liu, Q. *et al.* Wnt5a-induced M2 polarization of tumor-associated macrophages via IL-10

promotes colorectal cancer progression. *Cell Commun Signal* **18**, 51 (2020).

41. Liu, J. *et al.* An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-

Quality Survival Outcome Analytics. *Cell* **173**, 400-416.e11 (2018).

42. Enwere, E. K. *et al.* Expression of PD-L1 and presence of CD8-positive T cells in pre-

treatment specimens of locally advanced cervical cancer. *Mod Pathol* **30**, 577–586

(2017).

43. Mezache, L., Paniccia, B., Nyinawabera, A. & Nuovo, G. J. Enhanced expression of PD

L1 in cervical intraepithelial neoplasia and cervical cancers. *Mod Pathol* **28**, 1594–1602

(2015).

44. Reddy, O. L., Shintaku, P. I. & Moatamed, N. A. Programmed death-ligand 1 (PD-L1) is

expressed in a significant number of the uterine cervical carcinomas. *Diagn Pathol* **12**, 45

(2017).

45. Lheureux, S. *et al.* Association of Ipilimumab With Safety and Antitumor Activity in

Women With Metastatic or Recurrent Human Papillomavirus–Related Cervical

Carcinoma. *JAMA Oncol* **4**, e173776 (2018).

46. Daley, D. *et al.* Dectin 1 activation on macrophages by galectin 9 promotes pancreatic

carcinoma and peritumoral immune tolerance. *Nat Med* **23**, 556–567 (2017).

47. Nobumoto, A. *et al.* Galectin-9 suppresses tumor metastasis by blocking adhesion to

endothelium and extracellular matrices. *Glycobiology* **18**, 735–744 (2008).

48. Munn, D. H. *et al.* GCN2 Kinase in T Cells Mediates Proliferative Arrest and Anergy

Induction in Response to Indoleamine 2,3-Dioxygenase. *Immunity* **22**, 633–642 (2005).

49. Low, H.-Y. *et al.* Reciprocal Regulation Between Indoleamine 2,3-Dioxigenase 1 and

Notch1 Involved in Radiation Response of Cervical Cancer Stem Cells. *Cancers* **12**, 1547

(2020).

50. Li, X. *et al.* Navigating metabolic pathways to enhance antitumour immunity and

immunotherapy. *Nat Rev Clin Oncol* **16**, 425–441 (2019).

51. Biffi, G. & Tuveson, D. A. Diversity and Biology of Cancer-Associated Fibroblasts.

Physiological Reviews **101**, 147–176 (2021).

52. Chu, T.-Y., Yang, J.-T., Huang, T.-H. & Liu, H.-W. Crosstalk with Cancer-Associated

Fibroblasts Increases the Growth and Radiation Survival of Cervical Cancer Cells.

Radiation Research **181**, 540–547 (2014).

53. Zhang, X. *et al.* Extracellular vesicles-encapsulated microRNA-10a-5p shed from cancer-

associated fibroblast facilitates cervical squamous cell carcinoma cell angiogenesis and

tumorigenicity via Hedgehog signaling pathway. *Cancer Gene Ther* **28**, 529–542 (2021).

54. Sahai, E. *et al.* A framework for advancing our understanding of cancer-associated

fibroblasts. *Nat Rev Cancer* **20**, 174–186 (2020).

55. Guo, W. *et al.* scCancer: a package for automated processing of single-cell RNA-seq data

in cancer. *Briefings in Bioinformatics* **22**, bbaa127 (2021).

56. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

57. Zhang, Q. *et al.* Integrative Analysis of Hypoxia-Associated Signature in Pan-Cancer.

iScience **23**, 101460 (2020).

58. Cascone, T. *et al.* Increased Tumor Glycolysis Characterizes Immune Resistance to

Adoptive T Cell Therapy. *Cell Metabolism* **27**, 977-987.e4 (2018).

59. Baenke, F., Peck, B., Miess, H. & Schulze, A. Hooked on fat: the role of lipid synthesis in

cancer metabolism and tumour development. *Dis Model Mech* **6**, 1353–1363 (2013).

60. Mues, C. *et al.* Regulation of Glucose-6-Phosphatase Gene Expression by Insulin and

Metformin. *Horm Metab Res* **41**, 730–735 (2009).

61. Heusinkveld, M. & van der Burg, S. H. Identification and manipulation of tumor

associated macrophages in human cancers. *J Transl Med* **9**, 216 (2011).

62. Charoentong, P. *et al.* Pan-cancer Immunogenomic Analyses Reveal Genotype-

Immunophenotype Relationships and Predictors of Response to Checkpoint Blockade.

Cell Reports **18**, 248–262 (2017).

63. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21

(2019).

64. Malta, T. M. *et al.* Machine Learning Identifies Stemness Features Associated with

Oncogenic Dedifferentiation. *Cell* **173**, 338-354.e15 (2018).

65. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of

systems-level datasets. *Nat Commun* **10**, 1523 (2019).

66. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat*

Commun **12**, 1088 (2021).