Potential coupling between SARS-CoV-2 replicative fitness and interactions of its nucleoprotein with human 14-3-3 proteins

Kristina V. Tugaeva [1], Andrey A. Sysoev [1], Jake L. R. Smith [2], Richard B. Cooley [3], Alfred A. Antson [2]*, Nikolai N. Sluchanko [1]*

[1] A.N. Bach Institute of Biochemistry, Federal Research Center of Biotechnology of the Russian Academy of Sciences, 119071, Moscow, Russia
[2] York Structural Biology Laboratory, Department of Chemistry, University of York, York YO10 5DD, United Kingdom
[3] Department of Biochemistry and Biophysics, Oregon State University, Corvallis, OR, 97331, USA

*Corresponding authors: nikolai.sluchanko@mail.ru (0000-0002-8608-1416),
fred.antson@york.ac.uk (0000-0002-4533-3816)

Abstract

The SARS-CoV-2 nucleocapsid protein (N) is responsible for viral genome packaging and virion assembly. Being highly abundant in the host cell, N interacts with numerous human proteins and undergoes multisite phosphorylation in vivo. When phosphorylated within its Ser/Arg-rich region, a tract highly prone to mutations as exemplified in the Omicron and Delta variants, N recruits human 14-3-3 proteins, potentially hijacking their functions. Here, we show that in addition to phosphorylated Ser197, an alternative, less conserved phosphosite at Thr205 not found in SARS-CoV N binds 14-3-3 with micromolar affinity and is in fact preferred over pS197. Fluorescence anisotropy reveals a distinctive pT205/pS197 binding selectivity towards the seven human 14-3-3 isoforms. Crystal structures explain the structural basis for the discovered selectivity towards SARS-CoV-2 N phosphopeptides, and also enable prediction for how N variants interact with 14-3-3, suggesting a link between the strength of this interaction and replicative fitness of emerging coronavirus variants.

Keywords: Host-virus interactions; RNA viruses, coronavirus, 14-3-3 protein, phosphopeptide binding; binding selectivity

Introduction

The current pandemic, caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), exposed a number of weaknesses in our approach to tackle viral outbreaks. The lack of understanding regarding host-virus interactions and the causes of virus pathogenicity at the molecular level slowed the development of novel antiviral therapies, resulting in a lack of preparedness and, consequently, major health, economic and social impacts.

One of the most abundant viral proteins, the 46-kDa nucleoprotein (N), is common for single-stranded RNA viruses, including positive-sense coronaviruses and is responsible for replication, packaging and storage of viral genomic RNA [1,2]. As one of the most abundant viral proteins, N can reach up to 1% of the total cellular protein in infected cells [3–5] and is considered to be a major factor of pathogenicity [6]. Sharing 89.1% sequence identity with SARS-CoV N, SARS-CoV-2 N consists of the structured N-terminal RNA-binding domain [7], the structured C-terminal dimerization domain [8], disordered terminal tails and the long unstructured interdomain linker (residues 176-247) (Fig. 1a) [9,10].

The latter linker domain contains a Ser/Arg-rich region (residues ~176-210) [11–13] that becomes heavily phosphorylated upon host infection [4,14–16] and is also a hotspot of viral mutations [17–19]. This interdomain linker contributes to higher-order oligomerization of N molecules [20], to N-M interactions [21], and to liquid-liquid phase separation with RNA [11,17] via the tentative amyloid-like aggregation of specific peptide stretches 179-184, 217-222 and 243-248 [22]. Phosphorylation of several Ser/Thr residues in the SR-rich region controls the flexibility, electrostatics, and protein-protein interactions of SARS-CoV-2 N [17,23,24]. Specifically, phosphorylation can regulate N interaction with 14-3-3 [25,26] - the universal and abundant phosphopeptide-binding protein-protein interaction hub represented by seven specific isoforms in human (β, γ, ε, ζ, η, σ, τ) [27,28]. In line with this, SARS-CoV N undergoes phosphorylation-dependent association with human 14-3-3 in the cytoplasm of the transfected cells in order to control nucleocytoplasmic shuttling of N and possibly hijack 14-3-3 functions via sequestration [25,26].

In previous work, we have shown that upon phosphorylation of the SR-rich region, SARS-CoV-2 N is specifically recognized by human 14-3-3 proteins, to form a 2:2 complex [26]. The use of truncated N variants narrowed the binding site to phospho-Ser197 [26]. This site is conserved between SARS-CoV and SARS-CoV-2 N proteins (Fig. 1a), as well as several animal coronaviruses, and represents a nearly canonical 14-3-3-binding motif SRNpS[197]TP [26]. Nonetheless, other phosphosites within SARS-CoV-2 N promoting 14-3-3 binding were also proposed [26]. Despite biochemical advances on SARS-CoV-2 N interaction with 14-3-3 proteins, the structural basis for such interaction remained elusive.

Production of specific phosphorylated forms of N is challenged by the dense distribution of phosphorylatable residues, resulting in heterogeneously poly-phosphorylated N forms when incubated with kinases (Fig. 1b) [26,29]. Here, we first used genetic code expansion to verify phosphoserine exclusively at position Ser197 in SARS-CoV-2 N is sufficient to enable binding to 14-3-3. Intriguingly, when poly-phosphorylated by a kinase, we found that the naturally occuring SARS-CoV-2 N variant S197L still displayed a strong phosphorylation-dependent interaction with 14-3-3. We therefore identified an adjacent phosphosite at position Thr205 that also contributes to 14-3-3 binding and is not found in SARS-CoV N. Using fluorescence anisotropy, we determined micromolar dissociation constants for the interaction of both phospho-motifs with

all seven human isoforms of 14-3-3, which revealed the characteristic affinity profiles reflecting
the affinity hierarchy of the seven 14-3-3 isoforms [28] and a remarkable selectivity of 14-3-3
towards the SARS-CoV-2 N phosphopeptides. Crystal structures of the corresponding 14-3-3
complexes helped rationalize the site-selective binding of the SARS-CoV-2 N phosphopeptides
to 14-3-3. This also captured for the first time residues 193-210 of the functional SR-rich region,
providing a structural framework for the analysis of the existing and emerging SARS-CoV-2
mutations.
87
88

89
90
Fig. 1. SARS-CoV-2 N site-specifically phosphorylated at Ser197 interacts with human 14-3-3γ.
a. Schematic representation of the SARS-CoV-2 N domain structure. NTD and CTD,
interdomain linker, Ser/Arg-rich region are labeled, Ser197 and Thr205 phosphorylation sites
are marked by red circles, and blue rectangles indicate peptide fragments prone to amyloid
aggregation [22]. b. Local alignment of the SR-regions of SARS-CoV-2 and SARS-CoV N proteins
is shown, with the phosphosites of interest marked by red font and potential 14-3-3-binding sites

by bold font. Note that SARS-CoV N has only the first potential 14-3-3-binding site. c.
Phosphorylation of SARS-CoV-2 N variants analyzed by a Phos-tag gel electrophoresis. d. The
same samples as in c were analyzed by SDS-PAGE. e,f. The interaction of human 14-3-3γ with
singly phosphorylated SARS-CoV-2 N.S197TAG (e) or the unphosphorylated SARS-CoV-2 N
counterpart (f) analyzed by SEC when 14-3-3 was added to N in excess at 220 mM NaCl in the
samples and running buffer, with SDS-PAGE of all fractions along the elution profiles. M -
molecular mass markers (indicated in kDa to the left). Positions of proteins are indicated by
arrows on the right. Note that fractions 4-6 (dashed rectangles) contained the 14-3-3:N complex
only in the case of S197TAG.
106
107
108   **Results**
109
110   **Alternative 14-3-3-binding sites within SARS-CoV-2 N**
111
The key 14-3-3-binding site of SARS-CoV-2 N was proposed to center at phospho-Ser197,
which is shared by SARS-CoV and SARS-CoV-2 N proteins (Fig. 1b) [26]. However, direct
evidence for its role was lacking and whether other phosphosites contributed to 14-3-3 binding
remained unknown. To address this, we first sought to obtain N protein containing the Ser197
phosphosite only. Due to the strong propensity of this protein to undergo multisite
phosphorylation [26], it was not possible to make monophosphorylated N enzymatically.
Therefore, we used a genetic code expansion approach to site-specifically install phosphoserine
at a genetically programmable TAG stop codon [30]. Incorporation of phosphoserine into this
protein, N.S197TAG, was confirmed by the discrete upward shift on a Phos-tag gel
electrophoresis (Fig. 1c, d). In contrast, SARS-CoV-2 N phosphorylated during co-expression
with protein kinase A (PKA) produced multiple upward shifted bands corresponding to multisite
phosphorylation (Fig. 1c, d), in agreement with earlier observations [26].
Co-elution of human 14-3-3γ with SARS-CoV-2 N.S197TAG phosphoprotein, but not with the
unphosphorylated SARS-CoV-2 N counterpart, during size-exclusion chromatography confirmed
the formation of the phosphorylation-dependent complex (Fig. 1e, f), indicating that the Ser197
phosphosite plays a role in recruiting 14-3-3.
We then asked if the S197L mutation associated with the variant of the coronavirus identified in
2020 in Europe and Brazil [31–34], which blocks phosphorylation at this position, would preclude
SARS-CoV-2 N from forming complexes with 14-3-3γ. The S197L mutant was co-expressed
with PKA for phosphorylation and then purified similar to the wild-type protein. As a control we
produced unphosphorylated N.S197L protein in the absence of PKA, and validated the
differences in phosphorylation status of the two by Phos-tag and SDS-PAGE (Fig. 1c and d).
Unexpectedly, the phosphorylated S197L mutant showed a very efficient interaction with 14-3-
3γ, as confirmed by SEC with SDS-PAGE analysis of the fractions (Fig. 2a, b) as well as MALS
analysis of the molecular mass of the complex obtained (150.8 kDa, M_w/M_n = 1.000; M_w
expected for the 2:2 complex is 148.3 kDa) (Fig. 2c). Interestingly, at lower than 220 mM NaCl,
we could detect nonspecific binding of the unphosphorylated S197L mutant of SARS-CoV-2 N
to 14-3-3γ (data not shown). This could be completely blocked by increasing salt concentration

to 300 mM, at which the specific interaction of the PKA-phosphorylated S197L mutant was still
detectable (Fig. 2d-f).
142
143

144
145
Fig. 2. The S197L mutant of SARS-CoV-2 N interacts with human 14-3-3γ in a phosphorylation-
dependent manner. a,b. The interaction of human 14-3-3γ with PKA-phosphorylated (a) or
unphosphorylated (b) SARS-CoV-2 N.S197L mutant analyzed by SEC and SDS-PAGE of all
fractions along the elution profiles. 14-3-3γ was pre-incubated in excess with N in 220 mM NaCl,
and SEC was run in the same buffer. Molecular mass markers are indicated in kDa on the right.
Positions of proteins are indicated by arrows on the left. Note that fractions 3-6 (dashed
rectangles) contained the 14-3-3:N complex only in the case of pN.S197L. c. SEC-MALS
profiles for 14-3-3γ or its mixture with phosphorylated N.S197L protein obtained using a
Superdex 200 Increase 5/150 column at 220 mM NaCl in the premixed samples and running

buffer. Average M_W values and polydispersity indexes (M_w/M_n) for the peaks of interest are
indicated. d-f. Effect of salt on the interaction. SEC profiles of 14-3-3γ, phosphorylated N.S197L
or their mixture, obtained at 200 mM NaCl (d) or 300 mM NaCl (e) in the premixed samples and
running buffer. Note that the complex is detectable even at 300 mM NaCl. f. 300 mM NaCl in the
premixed samples and SEC buffer fully blocks nonspecific interaction of 14-3-3γ with
unphosphorylated N.S197L.

161
162
163
The ability of N.S197L to efficiently interact with 14-3-3γ in a phosphorylation-dependent
manner indicated that other site(s) besides Ser197 contribute to 14-3-3 binding. In particular,
the functional SR-rich region of N contains another potential 14-3-3-binding motif around Thr205
(SRG**T**SP), which is phosphorylated *in vivo* [4,14]. It has a sequence similar to the Ser197 site
(SRN**S**TP), and since other phosphosites of the SR-rich region have more divergent sequences
from the consensus 14-3-3-binding sites [RXX(pS/pT)XP/G] [35], we focused on the interaction of
14-3-3 with these two representative phosphopeptide fragments, SSRNpS[197]TPGSS and
SSRGpT[205]SPARM. Interestingly, this Thr205 site is not conserved in SARS-CoV N, which has
only the Ser197 site (Fig. 1b) [26]. SARS-CoV N still showed a pronounced phosphorylation-
dependent interaction with human 14-3-3γ (Supplementary Fig. 1).
The results obtained using SARS-CoV-2 N.S197L, S197TAG and SARS-CoV N suggested that
the pS197 and pT205 are alternative (sufficient but not necessary) 14-3-3-binding sites within
the SR-rich region.

177
178
179
Ser197 and Thr205 phosphopeptides display nonparallel affinity profiles towards human
14-3-3 isoforms

182
Synthetic phosphopeptides corresponding to the Ser197 and Thr205 fragments, each
additionally bearing a tryptophan residue for more accurate quantification, were labeled by
fluorescein 5-isothiocyanate (FITC) and used to determine binding affinities against all seven
human 14-3-3 isoforms by fluorescence anisotropy. For this, homotypic MBP-tagged full-length
14-3-3 constructs were used to obtain highly pure protein in an efficient manner [28]. SEC-MALS
analysis of the MBP-14-3-3ζ protein confirms that the MBP fusion does not disrupt the functional
dimeric state of 14-3-3 (Supplementary Fig. 2).
We first made sure that peptide binding to 14-3-3 was saturable and specific, and that the MBP
tag does not change the K_D (Fig. 3a). The same FITC-labeled phosphopeptide failed to bind to
noncognate protein lysozyme; likewise, free FITC did not show significant binding to MBP-14-3-
3 (Fig. 3a). This is consistent with previous observations showing that FITC-labeled
phosphopeptides and their unlabeled counterparts bind similarly to 14-3-3 [28,36]. In addition, the
FITC-peptide complexes with 14-3-3 could be disrupted by titration with either the
corresponding unlabeled peptides or the canonical phosphopeptide from unrelated protein
HSPB6 (pB6) [37] (Supplementary Fig. 3). These data demonstrate the chosen N peptides bind
specifically to the common peptide-binding grooves of 14-3-3.

Both SARS-CoV-2 N phosphopeptides showed saturable interaction with all seven human 14-3-
3 isoforms, enabling quantitative analysis of the dissociation constants (Fig. 3b and c,
Supplementary Fig. 4). The average binding affinities of 14-3-3 to the SARS-CoV-2 N
phosphopeptides were somewhat lower than to the pB6 peptide, yet were in a comparable
micromolar range (Table 1 and Supplementary Fig. 3). For each SARS-CoV-2 N peptide, we
detected a remarkable peptide-affinity trend following the general peptide-affinity hierarchy of
14-3-3 isoforms reported recently (Fig. 3d and e) [28]. For the pT205 peptide, the binding affinity
to 14-3-3 decreased from gamma (γ) > eta (η) > beta (β) > epsilon (ε) > zeta (ζ) > tau (τ) >
sigma (σ); the order for the pS197 peptide was: β ~ γ ~ η > ζ ~ τ ~ ε > σ (Fig. 3d). As a result,
transformation of the K_D values into -logK_D showed that both trends correlate well with the
interactome-wide affinity hierarchy of the 14-3-3 isoforms (Pearson's r=0.87 for pS197 and
r=0.93 for pT205, Fig. 3e) [28]. To facilitate calculations of how custom peptide binding to 14-3-3
isoforms correlates with their general affinity hierarchy based on the set of seven experimentally
determined (or predicted) K_D values for individual 14-3-3 isoforms, we created a simple Excel-
based tool (Supplementary Data file 1).
The affinity profiles of 14-3-3 for the two SARS-CoV-2 N peptides were significantly shifted and
nonparallel to each other. Due to the shift, the affinity of the weakest pT205 peptide binder, 14-
3-3σ (K_D ~33 μM), was still 2 times higher than for the strongest pS197 binders, 14-3-3β/γ/η (K_D
~70 μM). Overall, the 14-3-3 family covered affinities to SARS-CoV-2 phosphopeptides differing
by an order of magnitude [~10-fold K_D ratio between the weakest (14-3-3σ/pS197) and the
strongest (14-3-3γ/pT205) pair]. The average affinity of human 14-3-3 proteins to the studied
SARS-CoV-2 phosphopeptides (K_D ~50 μM) is roughly in the same order of magnitude as the
average affinity of twenty three unrelated peptides analyzed previously (K_D ~45 μM) [28]. This
suggests viable competition of the SARS-CoV-2 N phosphopeptides with the host cell
phosphopeptides for the 14-3-3 binding, especially with the more weakly interacting portions of
the interactome.
225
226 **Selectivity trend of 14-3-3 isoforms to SARS-CoV-2 N phosphopeptides**
227
Curiously, the affinities to all 14-3-3 isoforms were consistently tighter for the pT205 peptide
than for the pSer197 phosphopeptide (4.3-fold ratio of the average K_D values) (Fig. 3f).
Moreover, we found a remarkable selectivity trend reflecting a linear correlation of the K_D ratio
for the 14-3-3 isoforms with their general affinity hierarchy (R^2=0.84, Pearson's r=0.93; Fig. 3f)
[28]. In other words, not only does 14-3-3γ turn out to be the strongest binder, it is also the most
selective isoform with respect to the SARS-CoV-2 phosphopeptides pS197 and pT205 (the $K_{D;}$
$_{197}$/$K_{D; 205}$ ratio of ~7), while 14-3-3σ is the weakest and simultaneously the least selective (Fig.
3f, Table 1). Yet, even for 14-3-3σ the affinity to the pT205 peptide was ~3 times tighter than for
the pS197 peptide. This observation was totally unexpected given the very similar sequences of
both peptides, implying that the corresponding protein-peptide complexes feature different
interfaces and chemical contacts. To understand the molecular basis of this phenomena, we
crystallized 14-3-3σ complexes with both phosphopeptides and determined structures by X-ray
crystallography at high resolution (Table 2).
241

Fig. 3. Affinity and selectivity of SARS-CoV-2 N phosphopeptide binding to the human 14-3-3 protein family. a. Fluorescence anisotropy titration data for 14-3-3γ or MBP-14-3-3γ binding to the FITC-labeled SARS-CoV-2 N pT205 peptide. The corresponding apparent K_D values (mean ± SD, n=3) are shown. Titration of the same peptide by lysozyme and titration of free FITC by MBP-14-3-3γ are shown as controls. b, c. Titration of FITC-labeled SARS-CoV-2 N phosphopeptides pT205 (b) or pS197 (c) by the seven human 14-3-3 isoforms. The data (mean ± SD, n=3) are color-coded with respect to the 14-3-3 isoforms; fitting curves and the K_D values at equilibrium are collected in Supplementary Fig. 4 and Table 1. In a-c, background anisotropy was subtracted from all values. d. Affinity profiles of the SARS-CoV-2 N phosphopeptides versus 14-3-3 isoforms shown as heatmap gradients according to the K_D scale on the right. The source K_D values are found in Table 1. e, f. Affinity profiles presented as -logK_D values (e) or selectivity coefficients (f) are correlated with the proteome-wide affinity hierarchy of the 14-3-3 isoforms represented as average $\Delta\Delta G$ values for the individual isoforms (indicated by asterisks) [28]. Coefficient of determination R^2 and Pearson's r coefficient characterizing the linear correlations are indicated.

Crystallography explains the selectivity of 14-3-3 for SARS-CoV-2 N phosphopeptides

X-ray structures of 14-3-3σ with both peptides (Table 2) had one 14-3-3 dimer in the asymmetric unit, where each protomer was occupied by the target phosphopeptide. Eight out of ten pS197 phosphopeptide residues and all ten residues for the pT205 phosphopeptide are clearly defined in electron density, revealing remarkable details of the protein/peptide interaction (Fig. 5a, b). The central six residues including the phosphosite are canonically placed facing the conserved residues of the amphipathic 14-3-3 groove, forming a developed network of hydrogen bonds (Fig. 4a, b and Supplementary Fig. 5). Residues at +1 and -1 positions with respect to the phosphosite are H-bonded to the side chains of the conserved 14-3-3σ residues Asn175 and

Asn226, while hydroxyl groups of both Thr198 and Ser206 of the peptides (position +1) are
identically involved in H-bond formation with Asn175 and the conserved Lys122 residue of 14-3-
3σ (Fig. 4a, b and Supplementary Fig. 5). An additional H-bond in the case of the pS197 peptide
between the side chain of its Asn196 residue and Asn226 of 14-3-3σ is replaced by the
additional hydrophobic contact in the case of the pT205 peptide, formed between the methyl
group of phospho-Thr205 residue and the side chain of 14-3-3σ's Val178 (Fig. 4). In contrast to
the nearly equivalent six-residue segments of both peptides (Cα RMSD=0.7 Å), crystal
structures revealed significant differences in their C-terminal regions. In the case of the pS197
peptide, we could not detect any stabilizing *in-trans* interactions after Pro199. The lack of
defined electron density for the last two Ser residues of this peptide indicate flexibility and hence
inability to form stabilizing interactions (Fig. 4a). On the contrary, the pT205 peptide was
additionally stabilized *in-trans* by i) the hydrophobic contacts of its Ala208 with Val46 of 14-3-3
and Met210 with Phe118 and Ile168 of 14-3-3, ii) the H-bonding interaction between the
peptide's carbonyl of Ala208 with the side chain of Ser45 of 14-3-3σ and iii) the salt bridge
between the peptide's Arg209 with Glu14 of 14-3-3σ (Fig. 5b). Interestingly, Glu14 is absolutely
conserved in all human isoforms of 14-3-3 (Supplementary Fig. 6). Therefore, the salt bridge
interaction involving Arg209 is likely shared by all 14-3-3 isoforms and, together with factors i)-
ii), confer the consistently higher affinity of the pT205 peptide versus pS197 peptide to 14-3-3
proteins, as observed biochemically (Fig. 3).
These peptides of the same SARS-CoV-2 N polypeptide chain cannot simultaneously occupy
both grooves of the 14-3-3 dimer, as the pS197 and pT205 phosphosites are only 7 residues
apart (the interphosphate distance is 20 Å). For bidentate binding to 14-3-3, at least 15 residues
in the extended conformation [38] are needed to sufficiently separate the phosphosites (e.g. 22 in
the 5N6N structure [39]) and cover the required ~35 Å distance between the two binding sites of
the 14-3-3 dimer (Fig. 4c). While within the same SARS-CoV-2 N molecule pS197 and pT205
are obviously alternative 14-3-3-binding sites, we cannot exclude a mixed pS197/pT205 binding
within the heterotetrameric SARS-CoV-2 N/14-3-3 complex. Our data suggest that binding of
two pT205 motifs to the 14-3-3 dimer would be thermodynamically preferred, though mutations
in the SR-rich region might affect this selectivity.
298

Fig. 4. Crystal structures reveal interfaces between 14-3-3σ and two alternative 14-3-3-binding sites within SARS-CoV-2 N. a,b. Molecular contacts formed between a 14-3-3σ subunit (gray cartoon) and the pS197 phosphopeptide (magenta sticks) (a) or the pT205 phosphopeptide (orange sticks) (b). Hydrogen bonds are indicated by yellow dashed lines, hydrophobic interactions are indicated by large semi-transparent spheres, water molecules mediating 14-3-3-peptide contacts are in aquamarine. 2Fo-Fc electron density maps for the phosphopeptides contoured at 1σ are shown as semi-transparent blue surfaces in the inserts. Sequences of the phosphopeptides are shown above the inserts. Note that two C-terminal serines in the pS197 peptide structure and a tryptophan residue, attached to each peptide from the N terminus, are not visible due to disorder. c. A model showing that the two phosphopeptides (magenta and orange sticks) of the same N chain cannot occupy both grooves of the 14-3-3σ dimer (shown as semi-transparent cartoon) simultaneously. The pT205 peptide is depicted either bound in the bottom 14-3-3 groove (semi-transparent sticks) or as a continuation of the pS197 peptide bound in the top 14-3-3 groove. Distances between two phospho-groups in one continuous peptide (black dashed line) or as separate 14-3-3-bound entities (red dashed line) are indicated.

Structure-based prediction of effects of the N mutations on 14-3-3 binding

The conformation of the 14-3-3-bound peptides allowed us to rationalize the effect of N mutations present in different coronavirus variants (Table 3) by predicting their influence on 14-

3-3 binding. First, due to the phosphorylation-dependent nature of 14-3-3 interactions with partner proteins, the S197L and T205I mutations of N would simply block the interactions at their corresponding sites (while leaving the possibility of interaction using the alternative one). Interestingly, the S197L mutation was reported to have a statistically mild outcome [32]. The T205I mutation, present in multiple variants of SARS-CoV-2, has either mild outcome or co-occurs with multiple mutations in other SARS-CoV-2 genes (https://viralzone.expasy.org/9556), making predictions about its physiological effect challenging. A similar situation applies to the P199L mutation, which should decrease the affinity to 14-3-3 at the pS197 site because of the importance of the canonical Pro residue at this position [35] (Table 3, Fig. 5a). Nevertheless, the P199L-mutated N would still retain uncompromised interactions with 14-3-3 using the pT205 site. The S194L mutation associated with severe COVID-19 [32] can lead to site-specific strengthening of the interaction with 14-3-3 due to the additional hydrophobic contacts between Leu194 and a leucine residue of 14-3-3 facing this position in the crystal structure (Fig. 5b). The additional stabilization within the 14-3-3 complex is predicted also for the widespread combined R203K/G204R mutation present in at least several variants-of-concern (e.g., alpha, gamma, omicron) [40]. As R203K/G204R mutation is expected to enhance phosphorylation of N [40], based on structural observations we anticipate this would alter 14-3-3 interactions: while the R203K mutation is expected to slightly decrease the *in-cis* contact of the side chain with the phosphate group, the Arg204 side chain is placed at an appropriate distance for making a salt bridge with the conserved Asp225 of 14-3-3 (Fig. 5c). The S202R mutation has the most profound effect on the efficiency of N to package RNA and produce new viral particles [18] and has at least two predictable effects. First, the S202R mutation is expected to enhance phosphorylation of Thr205, at least by basophilic protein kinases (by forming a canonical sequence RRXT instead of just RXT [41]). Second, the S202R mutation converts the ..**S**RGpTSP.. 14-3-3-binding site from suboptimal to canonical (..**R**RGpTSP..) [35], and thereby is expected to enhance 14-3-3 binding (Fig. 5d). The effect of the R203M mutation associated with the Delta variant is difficult to predict at the level of 14-3-3 binding, because this mutation would first of all interfere with Thr205 phosphorylation in the S**M**GT[205]SP context, at least by basophilic protein kinases.

Fig. 5. Possible effects on 14-3-3 binding of the SARS-CoV-2 N mutations improving fitness of the virus. a. Sequence of the N region (193-210) covered by crystal structures of the 14-3-

3/SARS-CoV-2 N peptide complexes aligned with the canonical 14-3-3-binding motifs [35]. The
suboptimal 14-3-3-binding sites within N are shown by boxes. Colors correspond to Fig. 4. b-d.
Structural models showing the expected interaction of the mutated N peptides with 14-3-3
based on the crystal structures determined in the present study. The mutated residues are
shown in green, the polar contacts are shown by black dashed lines, the hydrophobic
interactions are indicated by semi-transparent spheres.

Discussion

Since the beginning of the pandemic SARS-CoV-2 acquired mutations that improve the
invasion, replication, evasion of the immune system, or transmissivity have become widespread.
The benefits of mutations in the S-protein can often be rationalized via their direct effect on viral
entry, which have now been structurally characterized [42,43]. However, the effects of mutations in
N remain largely unknown. Curiously, the majority of the widespread N mutations beneficial for
the virus are mapped to a very small segment, residues 194-205 (Fig. 1a, Table 3) [18,19,33]. In
fact, all variants-of-concern defined by the World Health Organization feature at least one
mutation in the 199-205 subregion [18]. This coincides with the maximum Shannon entropy in the
SARS-CoV-2 genome centered at residue 203 and surrounding positions within N [19]. However,
this mutational hotspot SR-rich region of N is disordered, making it challenging to study the
consequences of relevant mutations at the structural level. The fact that N appears to have
many functional roles, coupled with the lack of appropriate model systems for testing the effect
of mutations on N functionality *in vivo*, further challenges our ability to deconvolute the
underlying effects.
Despite these challenges, recent work has shown that the viral-like particles can be used to
rapidly assess evolved SARS-CoV-2 variants [18]. According to this study, the large effect of the
S202R and R203M mutations within the mutational hotspot of SARS-CoV-2 N can be explained
by the vastly improved RNA packaging mechanisms and production of higher titers of infectious
virions (166- and 51-fold, respectively) [18]. The molecular and structural consequences of these
and other widespread mutations of N have yet to be elucidated.
We questioned how post-translational modifications and these mutations might affect N
interactions with host cell factors. In previous work, we showed that SARS-CoV-2 N
phosphorylated at more than 20 sites, including several ones within the SR-rich region (176-
210) that are phosphorylated *in vivo* [4,14], can be specifically recognized by human 14-3-3
proteins, presumably, using the conserved motif phosphorylated at Ser197 [26].
In the present work, we first used a unique SARS-CoV-2 N protein variant containing a single
phosphoserine at position Ser197 to verify the role of this site in recruiting 14-3-3 proteins (Fig.
1). Consistent with this, we show that SARS-CoV N, which shares the pS197 site with SARS-
CoV-2 N, also interacts with 14-3-3 in a phosphorylation-dependent manner (Supplementary
Fig. 1). Surprisingly, the S197L variant of SARS-CoV-2 N found in the natural variant circulating
in Europe in 2020 and associated with a generally mild outcome [32], also displayed a hallmark
phosphorylation-dependent interaction with 14-3-3 (Fig. 2). This indicated that N had other 14-3-
3-binding phosphorylation sites besides pS197. Since the SR-rich tract, the key determinant of

14-3-3 binding [26], also contains another similar site around Thr205, we compared the interaction
of both phosphorylated fragments, pS197 and pT205, with all human isoforms of 14-3-3.
Using high-throughput fluorescence anisotropy data (Fig. 3), we found that both SARS-CoV-2 N
phosphopeptides display the characteristic peptide-affinity profiles correlating well (r=0.87 and
0.93) with the general affinity hierarchy of the human 14-3-3 isoforms [28]. The affinity profiles for
the two SARS-CoV-2 N fragments to 14-3-3 proteins were significantly shifted and nonparallel
to each other, indicating selectivity toward the 14-3-3 isoforms (Fig. 3). Unexpectedly, the pT205
fragment interacted with all seven 14-3-3 proteins with a consistently tighter affinity than the
pS197 fragment. Moreover, the interaction of the pT205 peptide with the weakest isoform, 14-3-
3σ, was still stronger than the interaction of the pS197 peptide with the strongest isoform, 14-3-
3γ (Fig. 3). Most importantly, we identified the remarkable selectivity profile for the 14-3-3 family,
a phenomenon that, to the best of our knowledge, has not been previously recognized with
respect to the 14-3-3 proteins. This selectivity trend also correlated well (R^2=0.84; r=0.93) with
the general peptide-affinity hierarchy of the 14-3-3 isoforms [28]. According to these results, not
only the strongest peptide binder, 14-3-3γ, binds the pT205 peptide stronger, but that its
selectivity to this peptide over the pS197 peptide is more pronounced than in the case of
generally the weakest isoform, 14-3-3σ (Fig. 3f, Table 1).
To explain the established binding preferences of 14-3-3 proteins toward one of the apparently
similar phosphopeptides, we determined crystal structures of 14-3-3σ with both peptides (Fig. 4
and Table 1). The X-ray structures enabled tracing of residues 193-200 and 201-210,
respectively, and conferred the following detailed analysis. The central portions of the peptides
adopt very similar conformations forming roughly equivalent contacts with 14-3-3. By contrast,
specific stabilizing interactions beyond the central region were observed only in the case of the
pT205 peptide. The pS197 and pT205 fragments are mutually exclusive, because their
simultaneous placement within the 14-3-3 dimer is not possible due to steric reasons (Fig. 4c).
The crystal structures of the 14-3-3-bound SARS-CoV-2 N fragments 193-200 and 201-210
provide a valuable structural framework for analyzing the effect of the widespread mutations and
testing new hypotheses. Collectively, our analyses based on structural, thermodynamic and
bioinformatic data support the hypothesis [19] that beneficial SARS-CoV-2 N mutations positively
affect 14-3-3 binding (Fig. 5).
Last but not least, the 14-3-3-binding region of N, residues 193-210, is immediately flanked by
the regions 179-184 and 217-222 (Fig. 1a) that were proposed to drive amyloid aggregation of
N in the course of liquid-liquid phase separation in the presence of RNA [22]. Because this
process plays a role in viral replication [17], one can expect the influence of N phosphorylation
and 14-3-3 binding to act as another layer of regulation of these processes and also a basis for
the effect of the widespread mutations. While this warrants further investigation, we note that
interaction with host signaling proteins such as 14-3-3, mediated by structurally disordered
segments containing phosphorylation sites, is emerging as a cornerstone in virus-host
interactions [25,26,28,44–48].
436
437
438

Methods

Peptides and chemicals

SARS-CoV-2 N peptides phosphorylated at Ser197 (WSSRNpSTPGSS) or Thr205 (WSSRGpTSPARM) were synthesized by Severn Biotech (UK). The phosphopeptide of human HSPB6 (pB6; WLRRApSAPLPGLK) was obtained as previously described [37]. Expected and experimentally determined masses of the peptides are listed in Supplementary Table 1. FITC was from Sigma-Aldrich (cat no. F7250). All common chemicals were of the highest purity and quality available, and all solutions were made with milliQ grade water.

FITC labeling of the SARS-CoV-2 N phosphopeptides

SARS-CoV-2 N peptide labeling was achieved in 0.1 M carbonate buffer, pH 9.0, by mixing a 5 mg/ml solution of the peptide with 1/10 volume of the FITC stock solution (9 mg/ml) in DMSO. The pB6 phosphopeptide [37] was labeled according to the same scheme and was used for controls. The mixtures were wrapped with aluminum foil and incubated overnight at room temperature, then the reaction was quenched by the addition of Tris to a final concentration of 1 M. The FITC-labeled peptides were separated from the unreacted label by size-exclusion chromatography on a 2 ml Sephadex G10 column equilibrated and run using a 20 mM Tris-HCl buffer, pH 7.5, 150 mM NaCl at a 0.5 ml/min flow rate. The runs were operated by the Prostar 335 system (Varian Inc., Australia) following the full-spectrum absorbance. The resulting spectrum, and absorbance at 280 (A_{280}) and 493 nm (A_{493}) in particular, was used to estimate the phosphopeptide concentration ($C_{peptide}$), taking into account FITC absorbance at 280 nm using the formula [49]:

$$C_{peptide} = (A_{280}-0.3*A_{493})/\varepsilon_{peptide,\,280}$$

where $\varepsilon_{peptide,\,280}$ is the extinction coefficient of the peptide at 280 nm (for all peptides in the study 5500 M^{-1} cm^{-1}). FITC concentration was determined from the FITC-specific absorbance at 493 nm using the extinction coefficient of 68,000 M^{-1} cm^{-1}. The completion of the labeling reaction was confirmed by LC-MS on a Bruker Impact II mass-spectrometer (Supplementary Table 1 and Supplementary Fig. 7).
The labeling efficiency was 74-76% for all phosphopeptides. Labeled peptides were stored in aliquots at -80 °C.

Plasmids and mutagenesis

Cloning of the SARS-CoV-2 N sequence (Uniprot ID P0DTC9) was described in previous work [26]. The protein contained an N-terminal His$_6$-tag cleavable by rhinovirus 3C protease, which left extra residues GPA on the N terminus of the protein. The wild-type SARS-CoV-2 N plasmid was used as template to PCR amplify the S197L mutant by the megaprimer method [50] using the S197L reverse mutagenic primer (5'-CTGGAGTTAAATTTCTTGAACTG-3', the mutated codon underlined). For site-specific phosphoserine incorporation into position 197 of SARS-CoV-2 N using amber codon suppression [30], we created the S197TAG mutant, which was obtained by

the megaprimer method using the reverse mutagenic primer (5'-
CTGGAGTCTAATTTCTTGAACTG-3', the mutated codon underlined) and DNA template being
the SARS-CoV-2 N plasmid containing a C-terminal His$_6$-tag cleavable by 3C protease. The
resulting PCR product containing the S197TAG substitution and carrying the C-terminal
cleavable His$_6$-tag was treated with NdeI/HindIII restriction endonucleases and then cloned into
a pRBC_14-3-3γ/His$_6$SUMO_mBAD$_{43-204}$ vector [30] pretreated with the same endonucleases. The
C-terminal His$_6$-tag enabled the use of the phosphoserine-incorporating *E.coli* system [BL21
Δ*serB*(DE3) cells transformed by the pKW2-EFSep plasmid (chloramphenicol resistance) and
the target N plasmid] having the Release Factor 1 (responsible for translation termination at
TAG codons) [51]. Despite contamination by the truncated product due to the preliminary
translation termination at the TAG[197] codon, only the C-terminally His-tagged full-length target
protein with incorporated phosphoserine could be purified by immobilized metal-affinity
chromatography (IMAC). After cleavage by 3C protease, such protein would retain extra
residues GLEVLFQ on its C terminus. The correctness of the obtained constructs and the
presence of the desired mutations were confirmed by DNA sequencing in Evrogen, Moscow
(see exemplary chromatograms for S197L and S197TAG in Supplementary Fig. 8).
The SARS-CoV N-containing pGEX plasmid (ampicillin resistance) was kindly provided by Prof.
Fulvio Reggiori (University of Groningen, The Netherlands).
Untagged full-length human 14-3-3γ (Uniprot ID P61981) cloned into a pET21 vector (ampicillin
resistance) [52], the surface entropy reduction mutants [53] of the C-terminally truncated human 14-
3-3σ (residues 1-231), i.e. Clu1 ([159]KKE[161] → [159]AAA[161]) and Clu3 ([75]EEK[77] → [75]AAA[77]), each
containing an N-terminal 3C protease cleavable His$_6$-tag and cloned into the pET28 vector
(kanamycin resistance) have been described previously [37]. Homotypic constructs of the full-
length human 14-3-3 protein isoforms beta (β), gamma (γ), eta (η), tau (τ), zeta (ζ), epsilon (ε),
and sigma (σ), fused to the C terminus of the maltose-binding protein (MBP) and described
earlier [28], were kindly shared by the laboratory of Prof. Gilles Travé (Université de Strasbourg,
Illkirch, France). Parameters of proteins used are provided in Supplementary Table 2.
509
510   **Protein expression and purification**
511
Proteins were expressed in *E.coli* BL21(DE3) by the addition of IPTG up to 0.5 mM for 20 h at
25 °C (N proteins), or for 4-6 h at 37 °C (14-3-3 proteins). To ensure phosphorylation of SARS-
CoV-2 N, its S197L mutant, or SARS-CoV N in *E.coli* cells, target proteins were co-expressed
with the catalytically active subunit of mouse protein kinase A (PKA) encoded in a separate
pACYC-PKA plasmid (chloramphenicol resistance), essentially as described earlier [26]. The site-
specifically phosphorylated SARS-CoV-2 N protein Ser197TAG was expressed in BL21
Δ*serB*(DE3) cells tailored to co-translational phosphoserine incorporation [51].
All N proteins, and Clu1 and Clu3 mutants of 14-3-3σ used for crystallization, were purified by
subtractive immobilized metal affinity and size-exclusion chromatography, where two IMAC
steps were separated by 3C cleavage of the His$_6$-tag [37]. To remove large amounts of
contaminating nucleic acids non-specifically bound to N protein, we used a continuous washing
step (50 column volumes) by 3 M NaCl before elution with imidazole during IMAC1. Most
proteins were obtained at amounts exceeding 10 mg per 1 liter of bacterial culture. While the
yield of the SARS-CoV-2 N.Ser197TAG phosphoprotein was particularly low (~1 mg per 1 liter

of bacterial culture) and the full-length protein was expressed alongside with the truncated side-
product 1-196, the resulting protein was indeed phosphorylated according to Phos-tag SDS-
PAGE [54]. All purified N preparations used in this study showed an A_{260}/A_{280} ratio of 0.57-0.6 and
were free from nucleic acids. To minimize protein degradation and dephosphorylation, SARS-
CoV-2 N.S197TAG was purified in the presence of inhibitors of proteases (EDTA-free Pierce™
Protease Inhibitor Mini Tablets, cat. no. A32955) and phosphatases (orthovanadate,
glycerophosphate and NaF).
SARS-CoV N was expressed as a GST-fusion and purified by a combination of a GSTtrap step
1, 3C proteolysis to cut off GST, a GSTtrap step 2 and heparin-affinity chromatography.
Untagged 14-3-3γ was expressed in *E.coli* BL21(DE3) cells by IPTG induction (final
concentration of 1 mM) and purified by ammonium sulfate fractionation, anion-exchange and
size-exclusion chromatography as described earlier [26]. Expression of the MBP-14-3-3
constructs was the same, but these proteins were purified using a combination of MBPTrap
column and size-exclusion chromatography [28]. Protein concentrations were determined
spectrophotometrically at 280 nm on a NP80 nanophotometer (Implen, Germany) using
extinction coefficients listed in Supplementary Table 2.
542
543 **Analytical size-exclusion chromatography**
544
The interaction of N proteins and 14-3-3γ was analyzed using a Superdex 200 Increase 5/150
column (GE Healthcare) operated at a 0.45 ml/min flow rate on a Prostar 335 chromatographic
system (Varian, Australia). When required, the eluted fractions were collected in eppendorf
tubes and their protein content was analyzed by SDS-PAGE [55]. In specific cases, a Superdex
200 Increase 10/300 column and a 0.8 ml/min flow rate was used. The buffer, used to
equilibrate the columns, contained 20 mM Tris-HCl pH 7.6, 3 mM NaN_3 and 200-300 mM NaCl.
We noticed that free unphosphorylated SARS-CoV-2 N.S197L absorbed onto the column when
equilibrated with <220 mM NaCl. Therefore, the reported experiments contained at least this
much salt in the SEC buffer. Experiments were done at least three times with qualitatively very
similar results.
555
556 **Multi-angle light scattering (MALS)**
557
To determine absolute masses of MBP-14-3-3ζ, 14-3-3γ and its complex with phosphorylated
SARS-CoV-2 N.S197L, we used SEC coupled to MALS. To do this, a chromatographic system
was connected to the miniDAWN detector (Wyatt Technology) calibrated relative to the
scattering from toluene. The concentration signal was obtained from SEC traces recorded at
280 nm and weight extinction coefficients 1.38 for MBP-14-3-3ζ, 1.13 for 14-3-3γ, and 1.02 for
the 14-3-3γ/pN.S197L complex. The M_w distributions were calculated in Astra 8.0 software
(Wyatt Technology) using dn/dc=0.185. Protein content in the peaks was additionally analyzed
by SDS-PAGE [55].
566
567 **Fluorescence anisotropy**
568

Fluorescence anisotropy (FA) was measured at 25 °C on a Clariostar plus microplate reader
(BMG Labtech, Offenburg, Germany) using a set of bandpass filters (482±16 nm and 530±40
nm), FITC labeled phosphopeptides (100 nM, on a 20 mM HEPES-NaOH buffer, pH 7.5, 150
mM NaCl, 0.02% Tween 20) and 384-well plates (Black Nunc™ 384-Shallow Well Standard
Height Polypropylene Storage Microplates, Thermofisher scientific, cat. no. 267460). A series of
concentrations of MBP-14-3-3 proteins in a 20 mM HEPES-NaOH buffer, pH 7.5, 150 mM NaCl,
0.01% Tween 20 (33 μl each) was diluted 2-fold by a 100 nM solution of either FITC-labeled
peptide to obtain 66 μl samples containing 50 nM peptide, MBP-14-3-3 (from 0 to 110 μM) and
0.01% Tween 20. After 5 min incubation at room temperature the fluorescence anisotropy
readout did not change with time, indicating equilibrium. Then, each sample was split into three
20 μl aliquots, the plates were centrifuged for 1 min at 25 °C at 250 g to remove air bubbles,
and the FP data were recorded. The triplicate measurements were converted into mean ±
standard deviation values, which were used for the data presentation and fitting. To determine
K_D values, the binding curves were approximated using the quadratic equation [56] in Origin 9.0
(OriginLab Corporation, Northampton, MA, USA). Selectivity coefficient reflecting the preferred
14-3-3 binding to the pT205 peptide than the pS197 peptide, which by definition is an
equilibrium constant ($K_{eq} = 1/K_D$) for the corresponding displacement reaction, was determined
as the ratio of K_D for the pS197 peptide to K_D for the pT205 peptide. To determine the
correlation between selectivity or -logK_D values for the 14-3-3/phosphopeptide pairs and
average $\Delta\Delta G$ values characterizing the affinity hierarchy reported earlier [28], we used the
standard Pearson's correlation coefficient r and coefficient of determination R^2.
590
591   **Crystallization and X-ray data collection**
592
Crystallization was performed using 14-3-3σ surface entropy reduction mutants Clu1 or Clu3
(storage buffer 20 mM Tris-HCl pH 7.6, 150 mM NaCl, 0.1 mM EDTA, 2 mM DTT, 2 mM MgCl$_2$)
mixed with either pS197 or pT205 phosphopeptide at a 1:5 protein:peptide molar ratio with the
final protein concentration of 11.5 mg/ml. The best diffracting crystal for the 14-3-3σ Clu1/pT205
complex grew in condition H7 (0.2 M AmSO$_4$, 100 mM BisTris pH 5.5, 25% PEG 3350) of
JCSG+ screen (Molecular Dimensions, Sheffield, UK); the best diffracting crystal for the 14-3-3σ
Clu3/pS197 complex grew in condition G2 (0.2 M NaBr, 100 mM BisTris propane pH 7.5, 20%
PEG 3350) of PACT screen (Qiagen, Hilden, Germany). Crystals were mounted in nylon loops
directly from crystallization drops and flash frozen in liquid nitrogen without additional
cryoprotection.
Synchrotron diffraction data (Table 1) were collected at 100 K at the Diamond Light Source
(Oxfordshire, UK) and processed using DIALS [57].
605
606   **Structure determination and refinement**
607
The structures were solved using MolRep [58] and an unliganded 14-3-3σ dimer (PBD 5LU2) as a
search model. Peptide residues were manually built into difference electron density maps in
Coot [59] and the overall structures were refined using Buster 2.10.4 [60], which used NCS
information (with pruning), TLS and all-atom individual isotropic B-factor restrained refinement.

In the case of the 14-3-3σ Clu1/pT205 complex, the refinement was additionally restrained
using the target structure of an unliganded 14-3-3σ dimer (PBD 3IQU, 1.05 Å [61]). The refined
structures were validated using the comprehensive validation algorithm in Phenix 1.10 [62].
Structural illustrations were prepared using PyMOL 2.20 (Schrodinger, Inc.).
616
617 **Data availability**
618
619 The refined models and structure factors have been deposited in the Protein Data Bank under
620 the accession codes 7QIK and 7QIP (released 29 Dec 2021). All materials are available from
621 the corresponding authors upon reasonable request.
622

623 **Acknowledgements**
624
We thank Prof. Gilles Trave's laboratory for providing the MBP-14-3-3 plasmids, Prof. Fulvio
Reggiori for the SARS-CoV N plasmid, Dr. Olga Moroz for help with crystallization and Dr.
Gergo Gogl for advice concerning fluorescence anisotropy. The authors are also thankful to
Sam Smith for help with X-ray data collection. LC-MS and MALS were carried out at the
Shared-Access Equipment Centre "Industrial Biotechnology" of the Federal Research Center
"Fundamentals of Biotechnology" of the Russian Academy of Sciences. The study was
supported by the Ministry of Science and Higher education of the Russian Federation in the
framework of the Agreement no. 075-15-2021-1354 (07.10.2021) and the Wellcome Trust
(206377 award to A.A.A.). R.B.C. acknowledges support of the Medical Research Foundation at
the Oregon Health Sciences University (USA) and the Collins Medical Trust. Expression and
purification of 14-3-3 proteins were partially supported by the Russian Science Foundation (no.
19-74-10031).

637
638 **Conflict of interests**
639
640 The authors declare no conflicts of interest.
641
642 **Author Contributions**
643
644 NNS and AAA conceived studies on N interaction with 14-3-3; KVT, AAS, JLRS, NNS -
645 expressed and purified proteins; KVT - performed fluorescence anisotropy experiments; JLRS -
646 crystallized proteins; NNS - determined crystal structures; KVT, RBC, AAA, NNS - analyzed
647 data; NNS and KVT made illustrations; NNS wrote the paper with input from RBC and AAA.
648

649 **References**
650
1. Masters, P. S. Coronavirus genomic RNA packaging. *Virology* **537**, 198–207 (2019).
2. Laude, H. & Masters, P. S. The Coronavirus Nucleocapsid Protein. in *The Coronaviridae* (ed.
Siddell, S. G.) 141–163 (Springer US, 1995). doi:10.1007/978-1-4899-1531-3_7.
3. Bojkova, D. *et al.* Proteomics of SARS-CoV-2-infected host cells reveals therapy targets.
Nature **583**, 469–472 (2020).

4. Bouhaddou, M. *et al.* The Global Phosphorylation Landscape of SARS-CoV-2 Infection. *Cell*
182, 685-712 e19 (2020).
5. Sender, R. *et al.* The total number and mass of SARS-CoV-2 virions. *Proc. Natl. Acad. Sci.*
118, (2021).
6. Gussow, A. B. *et al.* Genomic determinants of pathogenicity in SARS-CoV-2 and other
human coronaviruses. *Proc. Natl. Acad. Sci.* **117**, 15193–15199 (2020).
7. Kang, S. *et al.* Crystal structure of SARS-CoV-2 nucleocapsid protein RNA binding domain
reveals potential unique drug targeting sites. *Acta Pharm Sin B* **10**, 1228–1238 (2020).
8. Zinzula, L. *et al.* High-resolution structure and biophysical characterization of the
nucleocapsid phosphoprotein dimerization domain from the Covid-19 severe acute
respiratory syndrome coronavirus 2. *Biochem Biophys Res Commun* (2020)
doi:10.1016/j.bbrc.2020.09.131.
9. Ye, Q., West, A. M. V., Silletti, S. & Corbett, K. D. Architecture and self-assembly of the
SARS-CoV-2 nucleocapsid protein. *Protein Sci* **29**, 1890–1901 (2020).
10.   Zeng, W. *et al.* Biochemical characterization of SARS-CoV-2 nucleocapsid protein.
Biochem Biophys Res Commun **527**, 618–623 (2020).
11.   Cubuk, J. *et al.* The SARS-CoV-2 nucleocapsid protein is dynamic, disordered, and
phase separates with RNA. *Nat. Commun.* **12**, 1936 (2021).
12.   Nikolakaki, E. & Giannakouros, T. SR/RS Motifs as Critical Determinants of Coronavirus
Life Cycle. *Front Mol Biosci* **7**, 219 (2020).
13.   Tylor, S. *et al.* The SR-rich motif in SARS-CoV nucleocapsid protein is important for
virus replication. *Can J Microbiol* **55**, 254–60 (2009).
14.   Davidson, A. D. *et al.* Characterisation of the transcriptome and proteome of SARS-CoV-
2 reveals a cell passage induced in-frame deletion of the furin-like cleavage site from the
spike glycoprotein. *Genome Med* **12**, 68 (2020).
15.   Liu, X. *et al.* Targeting the coronavirus nucleocapsid protein through GSK-3 inhibition.
Proc. Natl. Acad. Sci. **118**, (2021).
16.   Yaron, T. M. *et al.* The FDA-approved drug Alectinib compromises SARS-CoV-2
nucleocapsid phosphorylation and inhibits viral infection in vitro. *bioRxiv* 2020.08.14.251207
(2020) doi:10.1101/2020.08.14.251207.
17.   Savastano, A., Ibanez de Opakua, A., Rankovic, M. & Zweckstetter, M. Nucleocapsid
protein of SARS-CoV-2 phase separates into RNA-rich polymerase-containing condensates.
Nat Commun **11**, 6041 (2020).
18.   Syed, A. M. *et al.* Rapid assessment of SARS-CoV-2 evolved variants using virus-like
particles. *Science* **0**, eabl6184.
19.   Del Veliz, S., Rivera, L., Bustos, D. M. & Uhart, M. Analysis of SARS-CoV-2
nucleocapsid phosphoprotein N variations in the binding site to human 14-3-3 proteins.
Biochem. Biophys. Res. Commun. **569**, 154–160 (2021).
20.   Chang, C. K., Chen, C. M., Chiang, M. H., Hsu, Y. L. & Huang, T. H. Transient
oligomerization of the SARS-CoV N protein--implication for virus ribonucleoprotein
packaging. *PLoS One* **8**, e65045 (2013).
21.   He, R. *et al.* Characterization of protein-protein interactions between the nucleocapsid
protein and membrane protein of the SARS coronavirus. *Virus Res* **105**, 121–5 (2004).
22.   Tayeb-Fligelman, E. *et al. Inhibition of amyloid formation of the Nucleoprotein of SARS-*

CoV-2.   2021.03.05.434000   https://www.biorxiv.org/content/10.1101/2021.03.05.434000v2
(2021) doi:10.1101/2021.03.05.434000.
23.   Carlson, C. R. *et al.* Phosphoregulation of Phase Separation by the SARS-CoV-2 N
Protein Suggests a Biophysical Basis for its Dual Functions. *Mol. Cell* **80**, 1092-1103.e4
(2020).
24.   Redzic, J. S. *et al.* The Inherent Dynamics and Interaction Sites of the SARS-CoV-2
Nucleocapsid N-Terminal Region. *J. Mol. Biol.* **433**, 167108 (2021).
25.   Surjit, M. *et al.* The severe acute respiratory syndrome coronavirus nucleocapsid protein
is phosphorylated and localizes in the cytoplasm by 14-3-3-mediated translocation. *J Virol* **79**,
11476–86 (2005).
26.   Tugaeva, K. V. *et al.* The Mechanism of SARS-CoV-2 Nucleocapsid Protein Recognition
by the Human 14-3-3 Proteins. *J. Mol. Biol.* **433**, 166875 (2021).
27.   Aitken, A. 14-3-3 proteins: a historic overview. *Semin Canc Biol* **16**, 162–172 (2006).
28.   Gogl, G. *et al.* Hierarchized phosphotarget binding by the seven human 14-3-3 isoforms.
Nat. Commun. **12**, 1677 (2021).
29.   Lin, L. *et al.* Identification of phosphorylation sites in the nucleocapsid protein (N protein)
of SARS-coronavirus. *Int J Mass Spectrom* **268**, 296–303 (2007).
30.   Zhu, P., Gafken, P. R., Mehl, R. A. & Cooley, R. B. A Highly Versatile Expression
System for the Production of Multiply Phosphorylated Proteins. *ACS Chem Biol* **14**, 1564–
1572 (2019).
31.   Coppée, F., Lechien, J. R., Declèves, A.-E., Tafforeau, L. & Saussez, S. Severe acute
respiratory syndrome coronavirus 2: virus mutations in specific European populations. *New
Microbes New Infect.* **36**, 100696 (2020).
32.   Nagy, Á., Pongor, S. & Győrffy, B. Different mutations in SARS-CoV-2 associate with
severe and mild outcome. *Int. J. Antimicrob. Agents* **57**, 106272 (2021).
33.   Tsai, P.-H. *et al.* Genomic variance of Open Reading Frames (ORFs) and Spike protein
in severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *J. Chin. Med. Assoc.* **83**,
725–732 (2020).
34.   Timmers, L. F. S. M. *et al.* SARS-CoV-2 mutations in Brazil: from genomics to putative
clinical conditions. *Sci. Rep.* **11**, 11998 (2021).
35.   Yaffe, M. B. *et al.* The structural basis for 14-3-3:phosphopeptide binding specificity. *Cell*
91, 961–71 (1997).
36.   Soini, L., Leysen, S., Crabbe, T., Davis, J. & Ottmann, C. The identification and
structural analysis of potential 14-3-3 interaction sites on the bone regulator protein Schnurri-
3. *Acta Crystallogr. Sect. F Struct. Biol. Commun.* **77**, 254–261 (2021).
37.   Sluchanko, N. N. *et al.* Structural Basis for the Interaction of a Human Small Heat Shock
Protein with the 14-3-3 Universal Signaling Regulator. *Structure* **25**, 305–316 (2017).
38.   Johnson, C. *et al.* Bioinformatic and experimental survey of 14-3-3-binding sites.
Biochem J **427**, 69–78 (2010).
39.   Alblova, M. *et al.* Molecular basis of the 14-3-3 protein-dependent activation of yeast
neutral trehalase Nth1. *Proc Natl Acad Sci U A* **114**, E9811–E9820 (2017).
40.   Johnson, B. A. *et al. Nucleocapsid mutations in SARS-CoV-2 augment replication and
pathogenesis.* 2021.10.14.464390
https://www.biorxiv.org/content/10.1101/2021.10.14.464390v1   (2021)

doi:10.1101/2021.10.14.464390.

41.   Douglass, J. *et al.* Identifying protein kinase target preferences using mass
spectrometry. *Am J Physiol Cell Physiol* **303**, C715-27 (2012).

42.   Yan, R. *et al.* Structural basis for the recognition of SARS-CoV-2 by full-length human
ACE2. *Science* **367**, 1444–1448 (2020).

43.   Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. *Science* **367**, 1260–1263 (2020).

44.   Nathan, K. G. & Lal, S. K. The Multifarious Role of 14-3-3 Family of Proteins in Viral
Replication. *Viruses* **12**, (2020).

45.   Tam, E.-H. *et al.* Role of the Chaperone Protein 14-3-3ε in the Regulation of Influenza A
Virus-Activated Beta Interferon. *J. Virol.* (2021) doi:10.1128/JVI.00231-21.

46.   Liu, J. *et al.* The role of 14-3-3 proteins in cell signalling pathways and virus infection. *J.*
Cell. Mol. Med. **25**, 4173–4182 (2021).

47.   Riedl, W. *et al.* Zika Virus NS3 Mimics a Cellular 14-3-3-Binding Motif to Antagonize
RIG-I- and MDA5-Mediated Innate Immunity. *Cell Host Microbe* **26**, 493-503 e6 (2019).

48.   Edwards, M. R. *et al.* Henipavirus W Proteins Interact with 14-3-3 To Modulate Host
Gene Expression. *J Virol* **94**, (2020).

49.   Bhattacharyya, S. *et al.* Synthesis and biological evaluation of panitumumab–IRDye800
conjugate as a fluorescence imaging probe for EGFR-expressing cancers. *MedChemComm*
5, 1337–1346 (2014).

50.   Sarkar, G. & Sommer, S. S. The 'megaprimer' method of site-directed mutagenesis.
Biotechniques **8**, 404–407 (1990).

51.   Rogerson, D. T. *et al.* Efficient genetic encoding of phosphoserine and its
nonhydrolyzable analog. *Nat Chem Biol* **11**, 496–503 (2015).

52.   Chernik, I., Seit-Nebi, A., Marston, S. & Gusev, N. Small heat shock protein Hsp20
(HspB6) as a partner of 14-3-3gamma. *Mol Cell Biochem* **295**, 9–17 (2007).

53.   Goldschmidt, L., Cooper, D. R., Derewenda, Z. S. & Eisenberg, D. Toward rational
protein crystallization: A Web server for the design of crystallizable protein variants. *Protein*
Sci **16**, 1569–76 (2007).

54.   Kinoshita, E., Kinoshita-Kikuta, E., Takiyama, K. & Koike, T. Phosphate-binding Tag, a
New Tool to Visualize Phosphorylated Proteins*. *Mol. Cell. Proteomics* **5**, 749–757 (2006).

55.   Laemmli, U. K. Cleavage of structural proteins during the assembly of the head of
bacteriophage T4. *Nature* **227**, 680–685 (1970).

56.   Jarmoskaite, I., AlSadhan, I., Vaidyanathan, P. P. & Herschlag, D. How to measure and
evaluate binding affinities. *eLife* **9**, e57264 (2020).

57.   Winter, G. *et al.* DIALS: implementation and evaluation of a new integration package.
Acta Crystallogr. Sect. Struct. Biol. **74**, 85–97 (2018).

58.   Vagin, A. & Teplyakov, A. Molecular replacement with MOLREP. *Acta Crystallogr Biol*
Crystallogr **66**, 22–5 (2010).

59.   Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*
Crystallogr Biol Crystallogr **60**, 2126–32 (2004).

60.   Blanc, E. *et al.* Refinement of severely incomplete structures with maximum likelihood in
BUSTER-TNT. *Acta Crystallogr Biol Crystallogr* **60**, 2210–21 (2004).

61.   Molzan, M. *et al.* Impaired binding of 14-3-3 to C-RAF in Noonan syndrome suggests

new approaches in diseases with increased Ras signaling. *Mol Cell Biol* **30**, 4698–711
(2010).

62.   Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr Biol Crystallogr* **66**, 213–21 (2010).

63.   Arévalo, S. J. *et al. Analysis of the Dynamics and Distribution of SARS-CoV-2 Mutations*
and its Possible Structural and Functional Implications. 2020.11.13.381228
https://www.biorxiv.org/content/10.1101/2020.11.13.381228v1 (2020)
doi:10.1101/2020.11.13.381228.

64.   Bourassa, L. *et al.* A SARS-CoV-2 Nucleocapsid Variant that Affects Antigen Test
Performance. *J. Clin. Virol.* **141**, 104900 (2021).

Tables
799

800 **Table 1**. Apparent K_D values and selectivity coefficients for the interaction of seven 14-3-3
801 isoforms with two SARS-CoV-2 N phosphopeptides as determined by fluorescence anisotropy.

14-3-3 isoform	K_D ±SD, µM		Selectivity*
	pT205	pS197	
gamma	10.7 ±0.3	71.6 ±1.0	6.7
eta	13.2 ±0.3	72.0 ±1.2	5.5
beta	14.4 ±0.3	69.9 ±1.5	4.8
epsilon	17.7 ±0.4	83.7 ±1.4	4.7
zeta	19.2 ±0.4	81.4 ±1.4	4.2
tau	20.8 ±0.4	81.9 ±1.5	3.9
sigma	32.9 ±0.8	98.6 ±1.9	3.0

802 * selectivity coefficients characterizing the preferred binding of the pT205 peptide over the
803 pS197 peptide were calculated for the given isoform as the ratio of the K_D for the pS197 to the
804 K_D for the pT205 peptide.

805 **Table 2.** X-ray data collection and refinement statistics.

	14-3-3σ/pS197 structure	14-3-3σ/pT205 structure
Data collection		
Detector	Dectris EIGER2 Si 16M	Dectris EIGER2 XE 16M
Wavelength, Å	0.9795	0.9763
Resolution range, Å	57.60-2.01 (2.12 - 2.01)*	70.45 - 2.65 (2.70 - 2.65)*
Space group	$P2_12_12_1$	$P4_1$
Cell parameters : a, b, c (Å)	47.4, 111.3, 115.2	99.7, 99.7, 58.6
Total reflections	418,896 (19,731)**	200,790 (10,676)
Unique reflections	39,958 (4740)	16,912 (818)
Multiplicity	10.5 (4.2)	11.9 (13.1)
Completeness (%)	96.6 (80.8)	100 (100)
Mean I/σ(I)	11.2 (1.3)	4.8 (0.9)
R-meas	10.5 (97.6)	36.4 (225.8)
$CC_{1/2}$	0.998 (0.767)	0.989 (0.353)
Refinement		
R-work	0.192 ***	0.221 ***
R-free	0.217	0.251
Number of non-H atoms	3918	3755
Protein atoms	3580	3607
Heteroatoms	68	25
Solvent atoms	338	115
Protein residues	459	471
RMS(bonds), Å	0.014	0.014
RMS(angles), °	1.4	1.4
Ramachandran favored (%)	98.9	98.0
Ramachandran outliers (%)	0	0.2
Rotamer outliers (%)	4.0	4.3
C beta outliers	0	0

Clashscore	0.54	0.83
MolProbity overall score	1.14	1.24
Average B-factor	47.6	70.3
PDB ID	7QIK	7QIP

806
* Statistics for the outer shell are in parentheses.
** Data processing was using DIALS [57].
*** Data are from validation using Phenix [62].

Table 3. Coupling between the effect of mutations in N of SARS-CoV-2 variants and binding to
14-3-3 proteins. Bold font indicates prediction of the most pronounced stabilizing effect on 14-3-
3 binding. Underlined italics indicate that the mutation blocks 14-3-3 binding at the
corresponding site.

Mutation	Disease severity	Ref.	Predicted effect on interaction with 14-3-3 (this work)	SARS-CoV-2 name**
S194L	Severe disease	32	Can be enhanced due to hydrophobic interaction with Leu229*	B.1.1.298***
S197L	Mild disease	32	None at the S197 site	A.2.4
P199L	Uncertain/mild	63	Weakening with the S197 site, by compromising the 14-3-3 motif	B.1.526 (Iota)
S202R	More transmissible, 166-fold higher infectious virus titers	18	Significantly improved owing to rectified 14-3-3-binding motif	B.1.526 (Iota)
R203K/ G204R	Severe disease, augmented viral replication, increased N phosphorylation	40	Enhanced due to predicted Arg204:Asp225* salt bridge	B.1.1.529 (Omicron), B.1.1.28.1 (Gamma), B.1.1.7 (Alpha), B.1.1.28.3 (Theta), B.1.1.298***
R203M	More transmissible, 51-fold higher infectious virus titers	18	Uncertain: inhibited via hampered N phosphorylation, neutral due to loss of Arg203 canonical interactions but new potential hydrophobic interaction of Met203 and Leu229*	B.1.617.2 (Delta), B.1.617.1 (Kappa)
T205I	Uncertain/mild	64	None at the T205 site	B.1.351 (Beta), B.1.427 (Epsilon), B.1.429 (Epsilon), B.1.525 (Eta), B.1.214.2

814 * human 14-3-3σ residue numbering. ** for variants-of-concern and variants-of-interest, as defined by the World
815 Health Organization, Greek letters are given in parentheses. *** in B.1.1.298, co-occurrence of S194L and
816 R203K/G204R has been found with high frequency.

817
818

819 **Supplementary information**
820
821 Contents:
822
823 **Supplementary Tables**
824
825 **Supplementary Table 1**. Monoisotopic masses of the synthetic peptides and their FITC-labeled
826 derivatives used in the study.
827 **Supplementary Table 2**. Parameters of protein constructs used in this study.
828
829 **Supplementary Figures**
830
Supplementary Fig. 1. SARS-CoV N interacts with human 14-3-3γ in a phosphorylation-
dependent manner.
Supplementary Fig. 2. SEC-MALS analysis of the MBP-14-3-3ζ fusion.
Supplementary Fig. 3. SARS-CoV-2 N phosphopeptides interact with the common peptide-
binding 14-3-3 grooves.
Supplementary Fig. 4. Titration of SARS-CoV-2 N phosphopeptides by the seven human 14-3-
3 isoforms monitored by fluorescence anisotropy.
Supplementary Fig. 5. The six core residues in the two SARS-CoV-2 N phosphopeptides form
similar contacts with 14-3-3σ.
Supplementary Fig. 6. The unique salt bridge-forming glutamate residue is highly conserved in
the human 14-3-3 protein family.
Supplementary Fig. 7. MS-based validation of the peptides and their labeled derivatives.
Supplementary Fig. 8. Sequencing results for the S197L and S197TAG mutants of SARS-
CoV-2 N.

855 **Supplementary Tables**

857 **Supplementary Table 1**. Monoisotopic masses of the synthetic peptides and their FITC-labeled
858 derivatives used in the study.

peptide		M_W (expected), Da		M_W (according to LC-MS), Da	
	sequence	unlabeled	labeled by FITC	unlabeled	labeled by FITC
pS197	WSSRN**pS**TPGSS	1244.47	1633.51	1244.49	1633.51
pT205	WSSRG**pT**SPARM	1314.54	1703.58	1314.58	1703.59
pB6	WLRRA**pS**APLPGLK	1543.83	1932.86	1543.84	1932.87

861 **Supplementary Table 2**. Parameters of protein constructs used in this study.

Protein	M_W, Da	pI	Molar ext. coef. M^{-1} cm^{-1}	Ext. coef. $(mg/mL)^{-1}$ cm^{-1}
MBP-14-3-3 epsilon	69505.8	4.8	95230	1.4
MBP-14-3-3 gamma	68634.5	4.9	98210	1.4
MBP-14-3-3 zeta	68077.0	4.9	93740	1.4
MBP-14-3-3 tau	68298.4	4.9	93740	1.4
MBP-14-3-3 eta	68752.9	4.9	95230	1.4
MBP-14-3-3 sigma	68308.2	4.8	92250	1.4
MBP-14-3-3 beta	68616.5	4.9	93740	1.4
14-3-3 gamma	28302.6	4.8	31860	1.1
14-3-3 sigma (1-231 aa) (^{159}KKE^{161}-^{159}AAA^{161}, Clu1)	26225.5	4.8	25900	1.0
14-3-3 sigma (1-231 aa) (^{75}EEK^{77}-^{75}AAA^{77}, Clu3)	26224.6	4.9	25900	1.0
SARS-CoV N (no tag)	46607.9	10.1	43890	0.9
SARS-CoV-2 N S197TAG (no tag)	46412.6	10.0	43890	0.9
SARS-CoV-2 N S197L (no tag)	45877.0	10.1	43890	1.0

  Supplementary Figures

866
867
Supplementary Fig. 1. SARS-CoV N interacts with human 14-3-3γ in a phosphorylation-
dependent manner. a. Phosphorylation of SARS-CoV N analyzed by a Phos-tag gel. b. The
same samples as in a were analyzed by SDS-PAGE. c-e. The interaction of unphosphorylated
SARS-CoV N with human 14-3-3γ analyzed by SEC at a 14-3-3 excess and 220 mM NaCl in
the samples and running buffer, with SDS-PAGE of all fractions along the elution profiles. f-h.
The interaction of PKA-phosphorylated SARS-CoV N with human 14-3-3γ analyzed by SEC at a
14-3-3 excess and 220 mM NaCl in the samples and running buffer, with SDS-PAGE of all
fractions along the elution profiles. c and f correspond to SEC profiles of N or pN. d and g
correspond to SEC profiles of 14-3-3 mixtures with either N or pN. e and h correspond to SEC
profiles of 14-3-3. Molecular mass markers are indicated in kDa. Positions of proteins are
indicated by arrows. Note that only phosphorylated SARS-CoV N forms a complex with 14-3-3γ
(fractions 4-6 outlined by dashed rectangles).

881
Supplementary Fig. 2. SEC-MALS analysis of the MBP-14-3-3ζ fusion. a. SEC-MALS profile
from the Superdex 200 Increase 10/300 column followed by absorbance at 280 nm showing the
M_w distribution across the protein peak as calculated from MALS data. Polydispersity index
(M_w/M_n) and the expected M_w for the protein dimer are indicated. Flow rate was 0.8 ml/min. b.
SDS-PAGE analysis of the fractions collected from the column. Mass markers are indicated in
kDa. Protein position and its apparent mass are marked by an arrow.

888
889
Supplementary Fig. 3. SARS-CoV-2 N phosphopeptides interact with the common peptide-
binding 14-3-3 grooves. The FITC-labeled SARS-CoV-2 N pT205 phosphopeptide is
outcompeted from its complex with human 14-3-3γ either by the same unlabeled
phosphopeptide (a) or by the control unlabeled pB6 peptide (b) as monitored by fluorescence
anisotropy. c. Titration of the FITC-labeled pB6 phosphopeptide by MBP-14-3-3ε, with the
corresponding K_D. In a-c, background anisotropy was subtracted from all values. The data are
shown as mean ± standard deviation (n=3).
897

Supplementary Fig. 4. Titration of SARS-CoV-2 N phosphopeptides by the seven human 14-3-3 isoforms monitored by fluorescence anisotropy. a. Binding curves for the pT205 peptide. b. Binding curves for the pS197 peptide. Fitting of the curves was done in Origin 9.0 by standard quadratic binding equation [56]. Apparent dissociation constants (K_D) at equilibrium are shown on the panels along with the coefficient of determination R^2. The data are shown as mean ± standard deviation (n=3). Fluorescence anisotropy data throughout titrations were corrected by subtracting the background anisotropy values.

907
908
Supplementary Fig. 5. The six core residues in the two SARS-CoV-2 N phosphopeptides form
similar contacts with 14-3-3σ. a, b. Superimposition of the two crystal structures showing the
similarity of the conformation of both peptides and molecular contacts formed, looking at two
different angles. H-bonds are shown by yellow dashed lines, hydrophobic interaction between
the methyl group unique to pT205 and the side chain of Val178 of 14-3-3σ is shown by large
semi-transparent spheres, distances characterizing key polar contacts are indicated in Å. Note
that the hydrophobic contact made by the methyl group of pT205 is somewhat compensated in
the pS197 peptide by an additional H-bond from the pS197 peptide's Asn196 side chain and the
side chain of Asn226 of 14-3-3σ, making the interaction of the core part of the peptides nearly
equivalent in terms of the amount of chemical contacts.

Supplementary Fig. 6. The unique salt bridge-forming glutamate residue is highly conserved in the human 14-3-3 protein family. a. Crystal structure of 14-3-3σ subunit (grey cartoon) complexed with the pT205 phosphopeptide of SARS-CoV-2 N (orange sticks) showing the location of the unique salt bridge (red dashed lines). 2Fo-Fc electron density map for the analyzed region is contoured at 1σ. b. Close-up of the salt bridge between the side chains of Arg209 of the SARS-CoV-2 N peptide and Glu14 of 14-3-3σ showing characteristic distances in Å. 2Fo-Fc electron density map for the analyzed region is contoured at 1σ. c. Local multiple sequence alignment of the human 14-3-3 isoforms in their N-terminal region, showing the absolute conservation of the glutamate partaking in the salt bridge to Arg209 of SARS-CoV-2 N. To the best of our knowledge, this glutamate is conserved in all 14-3-3 proteins. This implies that the ability to stabilize the binding of the pT205 phosphopeptide of SARS-CoV-2 N is shared by all isoforms with no exception.

935
936
937 **Supplementary Fig. 7**. MS-based validation of the peptides and their labeled derivatives. LC-
938 MS spectra for the pS197 peptide (a) and its FITC-labeled derivative (b).