A subset of Memory B-derived antibody repertoire from 3-dose vaccinees is ultrapotent against diverse and highly transmissible SARS-CoV-2 variants, including Omicron

Authors: Kang Wang[1$], Zijing Jia[1,8$], Linlin Bao[2$], Lei Wang[1,8$], Lei Cao [1$], Hang Chi[3$], Yaling Hu[4$], Qianqian Li[5$], Yinan Jiang[6], Qianhui Zhu[1,8], Yongqiang Deng[3], Pan Liu[1], Nan Wang[1], Lin Wang[4], Min Liu[4], Yurong Li[4], Boling Zhu[1], Kaiyue Fan[1,8], Wangjun Fu[1,8], Peng Yang[1,8], Xinran Pei[1], Zhen Cui[1,8], Lili Qin[6], Pingju Ge[6], Jiajing Wu[5], Shuo Liu[5], Yiding Chen[6], Weijin Huang[5], Cheng-Feng Qin[2]*, Youchun Wang[5]*, Chuan Qin[1]*, and Xiangxi Wang[1,7, 8]*

Affiliation:

[1] CAS Key Laboratory of Infection and Immunity, National Laboratory of Macromolecules, Institute of Biophysics, Chinese Academy of Sciences, Beijing 100101, China

[2] Key Laboratory of Human Disease Comparative Medicine, Chinese Ministry of Health, Beijing Key Laboratory for Animal Models of Emerging and Remerging Infectious Diseases, Institute of Laboratory Animal Science, Chinese Academy of Medical Sciences and Comparative Medicine Center, Peking Union Medical College, Beijing, China.

[3] State Key Laboratory of Pathogen and Biosecurity, Institute of Microbiology and Epidemiology, Academy of Military Medical Sciences, Beijing, China.

[4] Sinovac Biotech Ltd, Beijing, China

[5] Division of HIV/AIDS and Sex-Transmitted Virus Vaccines, Institute for Biological Product Control, National Institutes for Food and Drug Control (NIFDC), Beijing 102629, China

[6] Acrobiosystems Inc, Beijing, China

[7] Guangzhou Laboratory, Guangzhou, Guangdong, China, 510320

[8] University of Chinese Academy of Sciences, Beijing 100049, China

*Correspondence to: X.W. (Email: xiangxi@ibp.ac.cn) or C.F.Q. (Email:qincf@bmi.ac.cn) or Y.W. (Email: wangyc@nifdc.org.cn) or C.Q. (Email: qinchuan@pumc.edu.cn)

[$] These authors contributed equally to this work.

Abstract: (~150 words)

Omicron, the most heavily mutated SARS-CoV-2 variant so far, is highly resistant to neutralizing antibodies, raising unprecedented concerns about the effectiveness of antibody therapies and vaccines. We examined whether sera from individuals who received two or three doses of inactivated vaccine, could neutralize authentic Omicron. The seroconversion rates of neutralizing antibodies were 3.3% (2/60) and 95% (57/60) for 2- and 3-dose vaccinees, respectively. For three-dose recipients, the geometric mean neutralization antibody titer (GMT) of Omicron was 15, 16.5-fold lower than that of the ancestral virus (254). We isolated 323 human monoclonal antibodies derived from memory B cells in 3-dose vaccinees, half of which recognize the receptor binding domain (RBD) and show that a subset of them (24/163) neutralize all SARS-CoV-2 variants of concern (VOCs), including Omicron, potently. Therapeutic treatments with representative broadly neutralizing mAbs individually or antibody cocktails were highly protective against SARS-CoV-2 Beta infection in mice. Atomic structures of the Omicron S in complex with three types of all five VOC-reactive antibodies defined the binding and neutralizing determinants and revealed a key antibody escape site, G446S, that confers greater resistance to one major class of antibodies bound at the right shoulder of RBD through altering local conformation at the binding interface. Our results rationalize the use of 3-dose immunization regimens and suggest that the fundamental epitopes revealed by these broadly ultrapotent antibodies are a rational target for a universal sarbecovirus vaccine.

One sentence summary

A sub-set of antibodies derived from memory B cells of volunteers vaccinated with 3 doses of an inactivated SARS-CoV-2 vaccine work individually as well as synergistically to keep variants, including Omicron, at bay.

Main Text: The ongoing evolution and emergence of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) variants raise concerns about the effectiveness of monoclonal antibodies (mAbs) therapies and vaccines [1-3], posing challenges for global pandemic control. These variants were characterized as Variant of Interest, VOI or Variant of Concern, VOC by the World Health Organization (WHO). The more recently identified Omicron variant (B.1.1.529), designated as a new VOC, has led to an unprecedented surge in COVID-19 cases in South Africa and is now spreading across the world [4]. Remarkably, Omicron is the most heavily mutated variant to emerge so far with over thirty mutations in spike (S) protein, fifteen of which occur in the receptor binding domain (RBD). In addition, there are three small deletions and one 3-residue insertion in the N-terminal domain (NTD) of S1 subunit (Fig. 1a). The pattern of some of these alterations, similar to the those noted in previous VOCs, such as Δ69-70 in Alpha, N501Y in Alpha, Beta and Gamma, P681H in Alpha and Delta, are presumably associated with enhanced transmissibility, while many substitutions, including G142D/Δ143-145, ins214EPE, K417N, T478K, E484A, Q493K and N501Y, are closely related with resistance to neutralizing antibodies and vaccine induced humoral immunity [1,3,5-9] (Figs. 1a and 1b).

Although COVID-19 vaccines continued to be effective against severe diseases and deaths, including those caused by the circulating Delta variant, waning immunity and massive breakthrough infections caused by viral diversification warrant the need for a third dose or new vaccines. To combat the current resurgence of the epidemic, the U.S. Food and Drug Administration has authorized use of a 3rd booster dose for all adults after completion of primary vaccination with approved COVID-19 vaccine [10]. This step seems essential because preliminary studies have indicated that three doses of Pfizer-BioNtech mRNA vaccine neutralize the Omicron variant with an approximate 40-fold decline, while two doses are less effective [11,12]. However, these preliminary data on the neutralization sensitivity of Omicron require further independent confirmation. The clinical impact of natural and vaccine-induced immunity with regards to protection from infection and severe disease needs urgent

95   investigation.

Authentic virus neutralization of the Omicron variant by vaccine sera

The CoronaVac, a β-propiolactone-inactivated vaccine against COVID-19, has been

approved for emergency use, and recommended for a booster dose (third) of

inactivated vaccine in older persons by WHO [13,14]. Serum specimens from two groups

of 2-dose (n=60, at month 0, 1) or 3-dose (n=60, at months 0, 1, 7) CoronaVac

vaccinee volunteers were collected for evaluating neutralization titers against the

Omicron and Delta variants using a live SARS-CoV-2. None of the volunteers

recruited for vaccination was infected by SARS-CoV-2 prior to the study. Blood

samples from vaccinees collected 4 weeks after the last vaccination were used in

this study, to compare NAb titers against circulating SARS-CoV-2 variants. An

early passage of isolated (CHK06 strain) and sequence confirmed live Omicron

virus was used for neutralization assay in this study. Among three doses of

CoronaVac recipients, the geometric mean half-maximal neutralizing titers (GMT

NT_{50}) against live wild type (WT) virus, Delta and Omicron variants were 254, 78

and 15, respectively. Compared with WT, neutralizing titers against Delta and

Omicron were, on average, 3.3-fold and 16.5-fold reduced, respectively (Fig. 1c).

Only 3 of 60 samples had a NT_{50} titer of < 8 against the Omicron with a

seroconversion rate of 95% for neutralizing antibodies (Fig. 1c). However, it's

more concerning about effectiveness for two-dose regime against Omicron

infection. Among two doses of CoronaVac recipients, NT_{50} titer against Delta was

6.3 with a 5-fold reduction when compared to WT, but none of the serum

specimens had an NT_{50} titer of >8 against Omicron (Fig. 1c). Compared to 2-dose

vaccinees, sera of the 3-dose vaccinees displayed lower reduction in neutralization

titers against Delta, which is consistent with previous observations that 3-dose

administration of inactivated vaccine leads to enhanced neutralizing breadth to

SARS-CoV-2 variants [5].

Three doses of vaccine-elicited monoclonal antibodies

125 We previously sorted immunoglobulin (IgG+) memory B cells from peripheral

blood mononuclear cells (PBMCs) of four 3-dose CoronaVac vaccinees using prefusion SARS-CoV-2 S as a bait [5,15]. In total, we sorted 1800 SARS-CoV-2 S-specific memory B cells, obtained 422 paired heavy- and light-chain antibody sequences, and selected 323 antibodies for expression. Characterization by ELISA showed that 163, 100 and 51 recognized the RBD, NTD and S2, respectively and 9 failed to bind S (Fig. 2a). Biolayer interferometry affinities (BLI) measurements showed that nearly all RBD-directed antibodies bound to WT SARS-CoV-2 at sub-nM levels and 127 of them showed neutralization activities against both authentic and pseudotyped WT SARS-CoV-2 were selected for further investigation. Of these antibodies, over 93% of these antibodies exhibited broad binding activities to most VOCs and VOIs. Notably, 85% of these antibodies cross-reacted with the Omicron RBD. Contrarily, ~80% of NTD antibodies lost their associations with Omicron. Additionally, NTD antibodies also showed relatively poor cross-reactivity to other four VOCs due to the greater diversity of the NTD (Fig. 1, a, b).

A subset of antibodies with broad neutralization against circulating variants

Results of the pseudovirus neutralization assays performed by carrying the S of WT or other VOCs [16,17] identified XX RBD targeting antibodies that were especially potent with their half-maximal inhibitory concentration (IC_{50}) ranging from 0.002 to 0.800 ng/μl against WT as well as all VOCs (Fig. 2). Among these, 28 antibodies executed their neutralization via directly blocking the interactions between the RBD and its receptor hACE2, while 3 antibodies employ other mechanisms to neutralize viral infection (Fig. 2d, Extended Data Fig. 1). Especially, a subset of RBD antibodies (13 and 24) neutralized Omicron with $IC_{50} < 0.02$ and 0.1 ng/μl, respectively. These neutralizations are as potent as those exhibited by best-in-class antibodies against WT (Fig. 2b and 2d). We obtained IC_{50} values of 0.24 and 0.28 ng/μl for well-studied therapeutic antibodies like S309 and DXP-604, respectively. These values are 10~40-fold higher than those of the subset antibodies. Concerningly, some antibody drugs, such as REGN10933, REGN10987, LY-CoV555, LY-CoV016, AZD1061 and AZD8895, almost lost their neutralization activities against Omicron (Fig. 2b)[18]. Meanwhile, specific VOC-resistant

antibodies with high neutralizing potency against WT and some other VOCs (IC_{50} <0.2 ng/μl) were identified and these comprise ~30% of the antibody repertoire, indicative of the evolution of a wide range of antibodies after 3-dose vaccination. Experiments repeated using authentic virus, including WT and all five VOCs, showed similar neutralization patterns by all these antibodies, further verifying the neutralizing potency and breadth for this subset of antibody repertoire elicited by 3-dose vaccination (Fig. 2e).

Cryo-EM structures of the Omicron Spike in complex with 3 types of all five VOCs-reactive antibodies

Antibodies targeting the RBD can be categorized into six general classes (from I to VI) based on cluster analysis on epitope from 265 available RBD-NAb complex structures [5], that are related to the four groups on the basis of competition with the hACE2 for binding to S and recognition of the up or down state of the three RBDs in S [19,20]. ELISA-based square competition matrix analysis with the aid of existing structural data revealed the presence of 3 major groups in this subset of antibody repertoire (Extended Data Fig. 2). To delineate the structural basis for antibody-mediated neutralization, we determined the cryo-EM structure of a prefusion stabilized Omicron S trimer in complex with representative Fab fragments. The two highly potent antibodies against Omicron (XGv347 and XGv289 with IC_{50} values of 0.006 and 0.016 ng/μl, respectively), one mAb (XGv282 with IC_{50} of 0.268 ng/μl) with median neutralizing activities against Omicron, but high neutralizing potency against other four VOCs, and one mAb (XGv265 with IC_{50} of 7.479 ng/μl) with >500-fold decreased neutralization against Omicron, but potent neutralization against other four VOCs were selected for structural investigations (Fig. 2b). We determined cryo-EM reconstructions of these complexes at 3.2 – 3.6 Å, and performed local refinement to further improve the densities around the binding interface between RBD and antibodies, enabling reliable analysis of the interaction details (Fig. 3, Extended Data Fig. 3, 4 and 5, Extended Data Table 1).

The XGv347-Omicron S complex structures revealed three distinct conformational

states: three XGv347 Fabs bound to a completely closed S with three down RBDs;

two XGv347 Fabs bound to either two or one up and one down RBDs on S (Fig.

3a). By contrast, each of the complex structures for XGv289, XGv282 and XGv265

showed only one configuration where three XGv289 Fabs bound to two up and one

down RBDs; three XGv282 Fabs bound to one up and two down RBDs; two XGv265

Fabs bound to S trimer with one down and one up RBD, although the XGv265-bound

up RBD conformation was weakly resolved and therefore not modeled (Fig. 3a).

Antibody XGv347 binds to an epitope at the tip of RBD, largely overlapping with the

patch targeted by ACE2 (Fig. 2d, 3b, Extended Data Fig. 1). Structural comparisons

revealed that XGv347 is very similar to A23-58.1, an ultrapotent and broadly reactive

NAb effective against 23 SARS-CoV-2 variants [21]. Furthermore, the residues of the

epitope of XGv347 match with a major subset of those targeted by S2K146, another

broadly cross-reactive sarbecovirus NAb [22,23], highlighting a plausible capability of

these NAbs to cross-neutralize Omicron, SARS-CoV-2 variants and other

sarbecoviruses through ACE2 molecular mimicry. Unexpectedly, the epitopes of

XGv347, A23-58.1 as well as their sister NAbs would be normally inaccessible for

the RBD-down conformation in the WT S, but become accessible for either up or

down RBDs in the Omicron S due to a markedly outward expansion and a ~10º

clockwise rotation of three RBDs, leading to an approximately 9 Å conformational

movement for RBM (Fig. 3b and Extended Data Fig. 6). The XGv347 paratope

constituted five complementarity determining regions (CDRs) with heavy chain and

light chain contributing 70% and 30% of the binding surface area, respectively (Fig.

3b and Extended Data Table. 2). Overall XGv289, XGv282 and XGv265 bind

patches surrounding the right shoulder of RBD with various orientations, but in a

manner similar to those observed for DH1047, BD-812 and REGN10987; antibodies

known to generally neutralize most VOCs with high potency [24-26], but showing

declined, to varying degrees, binding and neutralizing activities against Omicron due

to the presence of new N440K and G446S mutations (Fig. 2b, Extended Data Fig. 7

and Extended Data Table. 3). Notably, XGv265 and REGN10987 recognize almost

same epitopes, both nearly losing their neutralizing activities against Omicron, despite

retaining weak binding (Extended Data Fig. 7). Structural superimpositions reveal

that XGv347 and either XGv289 or XGv265 can simultaneously bind to S, informing strategies to rationally design two-antibody cocktails (Extended Data Fig. 8).

Structural basis for broad cross-neutralization activity of NAbs and immune escape

XGv347, XGv289, XGv282 and XGv265 bound Omicron with 5-40 folds lower affinity compared to their binding with WT, although the same binding modes for two orthologs were observed (Fig. 3a). For XGv347, tight binding to WT S is primarily due to extensive hydrophobic interactions contributed by F456, Y473, F486 and Y489 from WT RBD, V32, V53, W51, P100 and F111 from heavy chain, and Y33 from light chain, and 9 hydrogen bonds (Fig. 3c and Extended Data Table 3). Hydrophobic interactions between the Omicron RBD and XGv347 are largely maintained. However, substitutions of Y505H and K417N abolish three hydrogen bonds forged with K75, D31 and E104 from HCDRs, leading to conformational shifts in HCDR3 and the RBM tip (residues 470-490), which further perturb six hydrogen bonds built by Y473, A475, S477, T478, Q493 from WT RBD with T105, C107, A56, G55 and D109 from HCDRs, albeit with an extra hydrogen bond established by the mutation Q493R and G55 from HCDR2 in Omicron (Fig. 3c). Similarly, a large patch of hydrophobic interactions constructed by V445, G446, Y449, P499 from WT RBD and F33, L50, I51, Y59, W103 from HCDRs as well as extensive hydrophilic interactions facilitate tight binding between XGv289 and WT S (Fig. 3c). Substitution of G446S disrupts the hydrophobic microenvironment, substantially decreasing hydrophobic interactions between Omicron S and XGv289. Furthermore, mutations of N440K and Q498R, together with altered local conformation, also lessen hydrogen bonds formed by N439, K440, Y449, R498, T500, Q506 from Omicron RBD and D95, L98 from LCDRs as well as Y59, N62 from HCDRs that would exist in XGv289-WT S complex (Fig. 3c). Among these four representative antibodies, XGv282 showed minimal reduction in binding affinity (5-fold), but remarkable reduction in neutralization (~40-fold), versus the characterization of XGv347 with 40-fold decrease in binding, but unchanged neutralization against Omicron when compared to WT, suggesting that epitope, rather than binding affinity, might play more crucial

roles in the neutralizing potency and breadth of an antibody. Consistent with XGv289, the substitution of G446S alters the hydrophobic microenvironment generally established by RBD and a group of antibodies bound at the right shoulder, including XGv289 and XGv282, triggering a conformational shift on CDRs and disrupting antibody recognition (Fig. 3c). In addition, the mutation E484A breaks hydrogen bond-connection with R74 from XGv282 HCDR2 and losses of charge interactions between R346, K444 from WT RBD and D56, D58 of XGv265 LCDR2 due to conformational alterations, further decreasing the binding of XGv282 and XGv265 to the Omicron variant, respectively (Fig. 3c). Taken together, G446S, acting as a critical mutation site, can alter the local conformation at the binding interface, conferring greater resistance to one class of antibodies bound at the right shoulder of RBD.

The therapeutic activities of five representative antibodies against the Beta SARS-CoV-2 variant in mice

Given the excellent neutralizing breadth and potency at cell-based levels for above antibodies, we next sought to assess the correlation between *in vitro* neutralization and *in vivo* protection. A number of representative mAbs with high neutralizing potency and breadth, belonging to different classes, such as XGv347, XGv289, XGv282, XGv265 and XGv052, produced in the HEK293F cell line were selected for therapeutic evaluation in a well-established mouse model challenged with the Beta variant [27]. Upon Beta intranasal challenge, adult BALB/c showed robust viral replication in the lungs at 3-5 days post inoculation. To evaluate the protection efficacy of these mAb, BALB/c mice challenged with the Beta variant were administered a single dose of as low as 5 mg/kg of XGv347, XGv289, XGv282, XGv265 and XGv052 individually or combinations of XGv282 and XGv347 (2.5 mg/kg for each), and XGv052 and XGv289 (2.5 mg/kg for each) in therapeutic settings (Fig. 4a). Heavy viral loads with high levels of viral RNAs (> 10^9 copies/g) were detected in the lungs at 5 day post infection in the control group of mice treated with PBS. However, a single dose of XGv282 reduced the viral RNA loads by ~10,000-fold in the lungs compared to the control group (Fig. 4b). Remarkably, a

single dose of XGv289, XGv265, XGv347, XGv052 or antibody cocktails of XGv282

+ XGv347, XGv052 + XGv289 resulted in a complete clearance of viral particles in

the lungs (Fig. 4b and 4c). A potential synergistic effect was observed for combined

therapies of XGv282 + XGv347 at 2.5 mg/kg for each (Fig. 4b and 4c). In addition,

histopathological examination revealed severe interstitial pneumonia, characterized

by alveolar septal thickening, inflammatory cell infiltration and distinctive vascular

system injury developed in mice belonging to the control group at day 5 (Fig. 4d). In

contrast, no obvious lesions of alveolar epithelial cells or focal hemorrhage were

observed in the lung sections from mice that received indicated antibody treatments

(Fig. 4d). Collectively, these results suggest that some of the antibodies from the

repertoire elicited by a 3-dose vaccination regimen retain therapeutic potential

against all circulating VOCs; albeit the protective efficacy against Omicron needs

to be investigated more thoroughly.

295   Discussion

The ongoing pandemic has witnessed frequent occurrences of SARS-CoV-2 variants

that increase transmissibility and reduce potency of vaccine-induced and therapeutic

antibodies [2,11]. More recently, there has been unprecedented concern that the Omicron

variant has significantly increased antibody escape breadth due to newly occurred and

accumulated mutations in the key epitopes of most neutralizing antibodies.

Alarmingly, Omicron nearly ablates the neutralization activity of most FDA approved

antibody drugs, including LY-CoV555, LY-CoV016, REGN10933, REGN10987,

AZD8895 and AZD1061 [18]. These issues raise an urgent need to develop next-

generation antibody-based therapeutics that can broadly neutralize these variants, as

well as future variants of concern. Our previous study revealed that the regimen of 3-

dose vaccination (0, 1, 7 months) of inactivated vaccine leads to an improved

immunity response with significantly enhanced neutralizing breadth via ongoing

antibody somatic mutation and memory B cell clonal turnover [5,28]. Correlated with

this, one subset of highly potent neutralizing antibodies with broad activities (IC_{50} <

0.2 ng/μl) against all circulating VOCs, including Omicron, were present in at least

four individuals who had received three doses of inactivated ancestral SARS-CoV-2

vaccine. Some, but not limited to these of this subset antibodies fully protected
against the Beta variant infection in mice, although their *in vivo* breadth and
protective efficacy against Omicron remains unconfirmed. Furthermore, our structural
and functional analyses revealed that a newly occurred mutation, G446S, might act as
a critical antibody escape site, conferring greater resistance to one major class of
antibodies bound at the right shoulder of RBD via altering microenvironments at the
S-NAb binding interface.

In addition to evading currently available antibody therapeutics, the Omicron variant
can diminish the efficacy of all clinically approved vaccines, including the mRNA
vaccines and inactivated vaccines [11,12]. There is an ongoing debate about whether the
immune responses can be fine-turned to the Omicron variant by boosting with a
tweaked (Omicron-based) vaccine. A major hurdle for this approach is the "original
antigenic sin", a phenomenon documented in some other infectious diseases,
including flu [29]. The presence of a subset of antibodies with broad neutralizing
activities against all circulating VOCs in memory B-derived antibody repertoire from
the 3-dose vaccinees suggests a possibility that selective and expeditious recall of
humoral responses might be elicited via the Omicron/future variants infection,
conferring to a secondary protection directed by memory etched in the immune
system. Further studies are warranted to examine the advantages and disadvantages
of booster shots of an Omicron-specific vaccine or simply administration of a
booster with the original vaccines. Lastly the identification and characterization of
broadly protective antibodies against all circulating VOCs will aid in the development
of universal vaccination strategies against sarbecoviruses.

References

1   Mlcochova, P. *et al.* SARS-CoV-2 B. 1.617. 2 Delta variant replication and immune
evasion. *Nature*, 1-6 (2021).
2   Wang, G.-L. *et al.* Susceptibility of circulating SARS-CoV-2 variants to neutralization.
New England Journal of Medicine (2021).
3   Altmann, D. M., Boyton, R. J. & Beale, R. Immunity to SARS-CoV-2 variants of concern.
Science **371**, 1103-1104 (2021).
4   Karim, S. S. A. & Karim, Q. A. Omicron SARS-CoV-2 variant: a new chapter in the
COVID-19 pandemic. *The Lancet* (2021).
5   Wang, K. *et al.* A third dose of inactivated vaccine augments the potency, breadth, and

duration of anamnestic responses against SARS-CoV-2. *medRxiv* (2021).

6   Hastie, K. M. *et al.* Defining variant-resistant epitopes targeted by SARS-CoV-2
antibodies: A global consortium study. *Science* **374**, 472-478 (2021).

7   Yuan, M. *et al.* Structural and functional ramifications of antigenic drift in recent SARS-
CoV-2 variants. *Science* (2021).

8   Zhang, J. *et al.* Membrane fusion and immune evasion by the spike protein of SARS-
CoV-2 Delta variant. *Science*, eabl9463 (2021).

9   Saito, A. *et al.* Enhanced fusogenicity and pathogenicity of SARS-CoV-2 Delta P681R
mutation. *Nature*, 1-10 (2021).

10   Mbaeyi, S. *et al.* The Advisory Committee on Immunization Practices' Interim
Recommendations for Additional Primary and Booster Doses of COVID-19 Vaccines—
United States, 2021. *Morbidity and Mortality Weekly Report* **70**, 1545 (2021).

11   Wilhelm, A. *et al.* Reduced Neutralization of SARS-CoV-2 Omicron Variant by Vaccine
Sera and monoclonal antibodies. *medRxiv* (2021).

12   Gardner, B. J. & Kilpatrick, A. M. Estimates of reduced vaccine effectiveness against
hospitalization, infection, transmission and symptomatic disease of a new SARS-CoV-2
variant, Omicron (B. 1.1. 529), using neutralizing antibody titers. *medRxiv* (2021).

13   Gao, Q. *et al.* Development of an inactivated vaccine candidate for SARS-CoV-2.
Science **369**, 77-81 (2020).

14   Hasan, S. A. W. Interim statement on booster doses for COVID-19 vaccination. *Update* **4**
(2021).

15   Wang, Z. *et al.* mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating
variants. *Nature* **592**, 616-622 (2021).

16   Lv, Z. *et al.* Structural basis for neutralization of SARS-CoV-2 and SARS-CoV by a potent
therapeutic antibody. *Science* **369**, 1505-1509 (2020).

17   Yao, H. *et al.* Rational development of a human antibody cocktail that deploys multiple
functions to confer Pan-SARS-CoVs protection. *Cell research* **31**, 25-36 (2021).

18   Cao, Y. R. *et al.* B. 1.1. 529 escapes the majority of SARS-CoV-2 neutralizing antibodies
of diverse epitopes. *bioRxiv* (2021).

19   Barnes, C. O. *et al.* SARS-CoV-2 neutralizing antibody structures inform therapeutic
strategies. *Nature* **588**, 682-687 (2020).

20   Dejnirattisai, W. *et al.* The antigenic anatomy of SARS-CoV-2 receptor binding domain.
Cell **184**, 2183-2200. e2122 (2021).

21   Wang, L. *et al.* Ultrapotent antibodies against diverse and highly transmissible SARS-
CoV-2 variants. *Science* **373**, eabh1766 (2021).

22   Park, Y.-J. *et al.* Antibody-mediated broad sarbecovirus neutralization through ACE2
molecular mimicry. *Biorxiv* (2021).

23   Cameroni, E. *et al.* Broadly neutralizing antibodies overcome SARS-CoV-2 Omicron
antigenic shift. *bioRxiv* (2021).

24   Martinez, D. R. *et al.* A broadly cross-reactive antibody neutralizes and protects against
sarbecovirus challenge in mice. *Science translational medicine*, eabj7125 (2021).

25   Zhu, L. *et al.* Double lock of a potent human therapeutic monoclonal antibody against
SARS-CoV-2. *National Science Review* **8**, nwaa297 (2021).

26   Hansen, J. *et al.* Studies in humanized mice and convalescent humans yield a SARS-
CoV-2 antibody cocktail. *Science* **369**, 1010-1014 (2020).

27   Chen, Q. *et al.* Transient acquisition of cross-species infectivity during the evolution of
SARS-CoV-2. *National science review* **8**, nwab167 (2021).

28   Wang, Z. *et al.* Naturally enhanced neutralizing breadth against SARS-CoV-2 one year
after infection. *Nature* **595**, 426-431 (2021).

29   Biswas, A., Chakrabarti, A. K. & Dutta, S. Current challenges: from the path of "original
antigenic sin" towards the development of universal flu vaccines: Flu vaccine efficacy

398   encounters significant hurdles from pre-existing immunity of the host suggesting
399   assessment of host immunity before vaccination. *International reviews of immunology*
400   **39**, 21-36 (2020).

401   30   Wang, N. *et al.* Structure-based development of human antibody cocktails against
402   SARS-CoV-2. *Cell research* **31**, 101-103 (2021).

403   31   Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure
404   determination in RELION-3. *elife* **7**, e42166 (2018).

405   32   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for
406   rapid unsupervised cryo-EM structure determination. *Nature methods* **14**, 290-296
407   (2017).

408   33   Sun, Y. *et al.* Structure-based development of three-and four-antibody cocktails against
409   SARS-CoV-2 via multiple mechanisms. *Cell research* **31**, 597-600 (2021).

Materials and Methods

Facility and ethics statements

All procedures associated with SARS-CoV-2 live virus were approved by the Animal experiment Committee Laboratory Animal Center, Beijing Institute of Microbiology and Epidemiology with an approval number of IACUC-IME-2021-022 and performed in Biosafety Level 3 (BSL-3) laboratories in strict accordance with the recommendations in the Guide for Care and Use of Laboratory Animals.

Viral stock and cell lines

SARS-CoV-2 wild-type strain CN01 was originally isolated from a patient during the early phase of COVID-19 endemic in China. SARS-CoV-2 variant of concern (VOC) Beta (B.1.351 lineage) strain GDPCC was isolated in a patient from South Africa and an Omicron (B.1.1.529 lineage) strain was isolated from a patient in Hong Kong and now preserved in SinoVac Biotech Ltd. All virus strains were first purified by standard plaque assay as previously described [13] and then inoculated into Vero cells (CCL-81) grown to 95% in 10% fetal bovine serum (FBS) supplemented Dulbecco's minimal essential medium (DMEM) for amplification.

Human sera samples

The serum samples were obtained from healthy volunteers who had no history of COVID-19 and were verified by PCR and serological assay and received two doses or three doses of CoronaVac (Sinovac) inactivated vaccine specific against SARS-COV-2. All volunteers were provided informed written consent form and the whole study was conducted in accordance with the requirements of Good Clinical Practice of China.

Authentic virus neutralization assay

The serum samples were first incubated at 56 °C for 30 min for inactivation. The heat-treated samples or monoclonal antibodies (mAbs) were subject to seral dilution from

1: 4 or 50 ng/μL with DMEM in two-fold steps and mixed with a virus suspension

containing 100 $TCID_{50}$ at 36.5°C for 2h, after which, the mixtures were added to wells

seeded with confluence Vero cells and incubated at 36.5°C for another 5 days in a

humidified 5% CO_2 cell incubator. After that, the cytopathic effect (CPE) of each well

was observed under microscopes by three different individuals and the related

dilutions and concentrations were recorded and used for the titration of samples tested

by the method of Reed-Muench [13].

Pseudovirus neutralization assay

The pseudotyped viruses bearing the Spike (S) protein were generated, aliquoted and

restored as previously described [17]. Briefly, 293T cells were first transfected with the

plasmid embedded with the S gene of wild-type or VOC/VOI (Alpha, Beta, Gamma,

Delta, Lamda and Omicron) SARS-CoV-2. The transfected 293T cells were infected

with VSV G pseudotyped virus (G*ΔG-VSV) at a multiplicity of infection (MOI) of 4.

After incubation for five hours, cells were washed with PBS, and then complete

culture medium was added. After another 24 hours, the SARS-CoV-2 pseudoviruses

were produced and harvested. For the *In vitro* pseudotyped virus neutralization assay,

the plasma samples or antibodies were diluted in DMEM starting from 1:10 or 10

ng/μL with 6 additional threefold serial dilutions, each of which were mixed with the

harvested pseudovirus and incubated at 37 °C for 1h. After that, the mixtures were

added to Huh-7 cells and placed back for incubation for another 24 hours. Then, the

luciferase luminescence (RLU) of each well was measured with a luminescence

microplate reader. The neutralization percentage was calculated as following:

Inhibition (%) = [1- (sample RLU- Blank RLU) / (Positive Control RLU-Blank RLU)]

(%). Antibody neutralization titers were presented as 50% maximal inhibitory

concentration (IC_{50}).

Protein expression and purification

The sequences of VOC Omicron full-length spike (S) protein (residues 1-1208),

receptor-binding domain (RBD) (residues 319-541) and N-terminal domain (NTD)

(residues 1-304) were modified from the plasmids encoding the S, RBD and NTD of

wild-type SARS-COV-2 (GenBank: MN908947) in our lab by overlapping PCR. In

additional to the reported mutations (A67V, Δ69-70, T95I, G142D, Δ143-145, Δ211,

L212I, ins214EPE, G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N,

T478K, E484A, Q493K, G496S, Q498R, N501Y, Y505H, T547K, D614G, H655Y,

N679K, P681H, N764K, D796Y, N856K, Q954H, N969K, L981F) on Omicron, the

proline substitutions at 986 and 987, 'GSAS' substitutions at the S1/S2 furin cleavage

site (residues 682-685) and a C-terminal T4 foldon trimerization domain were also

remained in the Omicron S construct to stabilize the trimeric conformation of S

protein. For protein expression, the plasmids of these proteins were transiently

transfected into HEK293 F cells grown in suspension at 37 °C in an incubator

supplied with 8% CO_2, rotating at 130 rpm. The cell supernatants were harvested and

concentrated three days post-transfection, and further purified by affinity

chromatography using resin attached with streptavidin or Ni-NTA and size-exclusion

chromatography (SEC) using a Superdex 200 column (GE Healthcare Life Sciences)

with the buffer containing 20 mM Tris pH 8.0 and 200 mM NaCl.

Antibody expression and Fab generation

The selected 323 antibodies were subjected to gene codon optimization,

construction and expression as described previously [5]. Then the clones were

transiently transfected into mammalian HEK293F cells and incubated for 5 days in

a 5% CO_2 rotating incubator at 37°C for antibody expression, which were further

purified using protein A. The purified mAbs XGv265, XGv282, XGv289 and

XGv347 were then processed to obtain their Fab fragments using the Pierce FAB

preparation kit (Thermo Scientific) as described previously [30]. Briefly, the samples

were first applied to desalination columns to remove the salt and the flow-throughs

were collected and incubated with papain that was attached with beads to cleave Fab

fragments from the whole antibodies for 5 hours at 37°C. After that, the mixtures

were transferred into Protein A columns and the flow-throughs, i.e., the Fab fragments

were collected and dialyzed into Phosphate Buffered Saline (PBS) (ThermoFisher,

catalog #10010023).

Bio-layer interferometry

Bio-layer interferometry (BLI) experiments were run on an Octet Red 384 machine (Fortebio). To measure the binding affinities of mAbs, monoclonal antibodies were immobilized onto Protein A biosensors (Fortebio) and the threefold serial dilutions of wild-type RBD (ACROBiosystems, Cat No. SPD-C52H3), Alpha RBD (ACROBiosystems, Cat No. SPD-C52Hn), Beta RBD (ACROBiosystems, Cat No. SPD-C52Hp), Gamma RBD (ACROBiosystems, Cat No. SPD-C52Hr), Delta RBD (ACROBiosystems, Cat No. SPD-C52Hh) and Omicron RBD (ACROBiosystems, Cat No. SPD-C522e) were used as analytes. Data were then analyzed using software Octet BLI Analysis 12.2 (Fortebio) with a 1:1 fitting model. For the epitope binning by BLI, SARS-CoV-2 VOC Omicron RBD tagged with his (ACROBiosystems, Cat No. SPD-C522e) was loaded on HIS1K biosensors, which were pre-equilibrated in the buffer for at least 1 min. The loaded biosensors were immersed with the first mAb for 300 s, followed by addition of the second mAb for another 90 s. Date obtained were also analyzed by Octet BLI Analsis 12.2.

ELISA assays

To evaluate whether the given mAbs could block the interaction between human ACE2 (hACE2) and RBD, ACE2 competition ELISA was performed by using the SARS-CoV-2 (B.1.1.529) Inhibitor Screening Kit (ACROBiosystems, Cat No. EP-115) according to the recommended protocol. Briefly, each of the 10 two-fold dilution series of mAbs (starting dilution of 25 ng/μL) and 0.8 ng/μL of HRP-conjugated SARS-CoV-2 Omicron RBD were added into the ELISA plate wells which are pre-coated with hACE2 protein. After incubation at 37 ˚C for 1 hour, the plates were washed three times with PBST (0.1% Tween) and the colorimetric signals were developed by addition of 3, 3', 5, 5'-tetramethylbenzidine TMB (Thermo Fisher) for 10 min. The reaction was stopped by addition of 50 μL of 1M H_2SO_4. The absorbance was measured at 450 nm with an ELISA microplate reader. For each mAb, a blank control with no mAb was added for inhibition calculation. The area under the curve (AUC) of each mAb were determined using Prism V8.0 (GraphPad). For

competitive ELISAs to identify the domain of a given mAb, 96-well plates were first

coated with RBD (2 μg/ml) and then blocked with 2% BSA in PBS. After incubation

with the reference mAbs, the blocking antibody (15 μg/ml), the wells were followed

by directly adding the second biotinylated antibodies (0.25 μg/ml). Streptavidin-HRP

(BD Biosciences) was then added for detection. Samples with no first antibody were

used as a negative control for normalization.

Cryo-EM sample preparation, data collection

The purified S protein was mixed with each of the Fab fragments of XGv265,

XGv282, XGv289 or XGv347 with a molar ratio of 1: 1.2 for 10 s ice incubation,

and then dropped onto the pre-glow-discharged holey carbon-coated gold grid (C-

flat, 300-mesh, 1.2/1.3, Protochips In.), blotted for 7 seconds with no force in 100%

relative humidity and immediately plunged into the liquid ethane using Vitrobot

(FEI). Cryo-EM data sets of these complexes were collected at 300 kV with an FEI

Titan Krios microscope (FEI). Movies (32 frames, each 0.2 s, total dose of 60 e$^-$ Å^{-2})

were recorded using a K3 Summit direct detector with a defocus range between 1.5-

2.7 μm. Automated single particle data acquisition was carried out by SerialEM, with

a calibrated magnification of 22,500 yielding a final pixel size of 1.07 Å.

Cryo-EM data processing

A total of 3,752, 2,631, 3,955 and 5,014 micrographs of S-XGv265-complex, S-

XGv282-complex, S-XGv289-complex and S-XGv347-complex, respectively were

recorded and subjected to beam-induced motion correction using motionCorr in

Relion3.0 package [31]. The defocus value of each image was calculated by Gctf. Then,

1,302,103, 756,508, 2,332,045 and 2,320,416 particles of the S-XGv265-complex, S-

XGv282-complex, S-XGv289-complex and S-XGv347-complex, respectively were

picked and extracted for reference-free 2D alignment by cryoSPARC [32], based of

which, 422,083, 190,154, 837,832 and 614,852 particles were selected and applied for

3D classification by Relion3.0 for S-XGv265-complex, S-XGv282-complex, S-

XGv289-complex and S-XGv347-complex, respectively with no symmetry imposed

to produce the potential conformations for the complexes. Afterwards, the candidate

model for each complex was selected and processed by non-uniform auto-refinement

and postprocessing in cryoSPARC to generate the final cryo-EM density for S-

XGv265-complex, S-XGv282-complex, S-XGv289-complex and S-XGv347-

complex. To improve the resolution of the interface between RBD and mAbs, the

block-based reconstruction was performed to obtain the final resolution of the focused

interfaces which contained the interfaces of RBD and mAbs investigated here as

described previously [33]. The resolution of each structure was determined on the basis

of the gold-standard Fourier shell correlation (threshold = 0.143) and evaluated by

ResMap. All dataset processing is shown in Extended Data Fig. 3 and also

summarized in Extended Data Table 1.

Model fitting and refinement

The atomic models of the complexes were generated by first fitting the chains of the

native apo SARS-CoV-2 S trimer (PDB number of 6VYB) and Fabs (PDB number of

7LSS and 7CZW for XGv265, 5MES and 5VAG for XGv282, 6UDA and 7MEG for

XGv289 as well as 7E3K for XGv347) into the cyo-EM densities of the final S-Fab-

complexes described above by Chimera, followed by manually adjustment and

correction according to the protein sequences and densities in Coot, as well as real

space refinement using Phenix. Details of the refinement statistics of the complexes

are summarized in Extended Data Table 1.

MD simulation and ΔG estimation

Model of SARS-CoV-2 wild-type RBD in complex with XGv265, XGv282, XGv289

and XGv347 were generated in Chimera by superimposition of WT RBD and cryoEM

structure of Omicron RBD in complex with the four antibodies. Before molecular

dynamics, all models were checked by WHAT IF Web Interface

(https://swift.cmbi.umcn.nl/servers/html/index.html) to model missing sidechains and

remove atomic clashes. After that, the structure was simulated by GROMACS-2021.

Briefly, we used OPLS force field with TIP3P water model to prepare the dynamic

system and add Na+ and Cl⁻ ions to make the system electrically neutralized. Then,

the system was subjected to energy minimization using the steepest descent algorithm

until the maximum force of 1,000 kJ mol-1 has been achieved. NVT ensemb1e via the

Nose-Hoover method at 300 K and NPT ensemble at 1 bar with the Parinello-Rahman

algorithm were employed successively to make the temperature and the pressure

equilibrated, respectively. Finally, a MD production runs of 100 ns were performed

starting from random initial velocities and applying periodic boundary conditions.

The non-bonded interactions were treated using Verlet cut-off scheme, while the long-

range electrostatic interactions were treated using particle mesh Ewald (PME)

method. The short-range electrostatic and van der Waals interactions were calculated

with a cut-off of 12 Å. Average structure of the four complexes were generated using

the last 10 ns frames and ΔG between the antibodies and RBD was estimated in

ROSETTA by InterfaceAnalyzer. Atomic_burial_cutoff, sasa_calculator_probe_radius

and interfaces_cutoff values were set to 0.01, 1.4 and 8.0 respectively.

Protection against SARS-CoV-2 Beta variant strain challenge in mice

The in vivo protection efficacies of single antibody or antibody cocktail were assessed

by using a newly established mouse model based on a SARS-CoV-2 Beta variant

strain [27]. Briefly, group of 8-month-old female BALB/c mice were infected with

1×10^4 PFU of SARS-CoV-2 Beta variant strain, then infected mice were treated

intraperitoneally with a single dose of different antibodies or antibody cocktails at 1

hour after infection. The lung tissues of mice were collected at 5 dpi for viral RNA

loads assay and pathological examination.

Viral burden determination

Viral burden in lung from mice were measured as described previously [16]. Briefly,

lung tissue homogenates were clarified by centrifugation and viral RNA was extracted

using the QIAamp Viral RNA Mini Kit (Qiagen). Viral sgRNA quantification in each

tissue sample was performed by quantitative reverse transcription PCR (RT-qPCR)

targeting the S gene of SARS-CoV-2. RT-qPCR was performed using One-Step

PrimeScript RT-PCR Kit (Takara).

Histology, and RNA in situ hybridization (RNA ISH)

Lung tissues from mice were fixed with perfusion fixative (formaldehyde) for 48 h,

and embedded in paraffin according to standard histological assays. For

histopathology, lung tissues were stained with hematoxylin and eosin (H&E). Images

were captured using Olympus BX51 microscope equipped with a DP72 camera. For

RNA ISH assays were performed with an RNAscope 2.5 (Advanced Cell Diagnostics)

according to the manufacturer's instruction. Briefly, formalin-fixed paraffin-

embedded tissue sections of 5 μm were deparaffinized by incubation for 60 min at 60

ºC. Endogenous peroxidases were quenched with hydrogen peroxide for 10 min at

room temperature. Slides were then boiled for 15 min in RNAscope Target Retrieval

Reagents and incubated for 30 min in RNAscope Protease Plus before probe

hybridization. The probe targeting 2019-nCoV RNA was designed and synthesized by

Advanced Cell Diagnostics (catalog no. 848561). Tissues were counterstained with

Gill's hematoxylin and visualized with standard bright-field microscopy. Original

magnification was 10×.

Reporting summary

Further information on research design is available in the Nature Researh

Reporting Summary linked to this paper.

Data availability

The atomic coordinates of XGv347 in complex with S trimer (state I), XGv347 in

complex with S trimer (state II), XGv347 in complex with S trimer (state III),

XGv347-S have been submitted to the Protein Data Bank with accession numbers:

7WEA, 7WEC and 7WEB, respectively. Futhermore, the atomic coordinates of

XGv265, XGv282 and XGv289 have been deposited in the protein data bank under

accession code 7WE8, 7WE7 and 7WE9, respectively. Cryo-EM density maps in

this study have been deposited at the Electron Microscopy Data Bank with

accession codes EMD-32444 (state1), EMD-32446 (state2) and EMD-32445

(state3), EMD-32441 (XGv282), EMD-32442 (XGv265), and EMD-32443

(XGv289). To reveal structural details of Fab binding mechanism, the local

optimized method are used to optimized data progress and the related atomic

models and EM density maps of optimized reconstructions of Fab interaction
interface has been deposited under accession code 7WEE (XGv265), 7WED
(XGv347), 7WEF (XGv289), EMD-32447 (XGv347), EMD-32448 (XGv265),
EMD-32449 (XGv289), respectively.

Acknowledgments We thank Dr. Boling Zhu, Dr. Xujing Li and Dr. Xiaojun
Huang, Lihong Chen for cryo-EM data collection at the Center for Biological
imaging (CBI) in Institute of Biophysics for EM work. We also thank Dr. Yuanyuan
Chen, Zhenwei Yang, Bingxue Zhou for for technical support on SPR. Work was
supported by the Strategic Priority Research Program (XDB29010000,
XDB37030000), CAS (YSBR-010), National Key Research and Development
Program (2020YFA0707500, 2018YFA0900801), Emergency Key Program of
Guangzhou Laboratory, Grant No. EKPG21-09 and Beijing Municipal Science and
Technology Project (Z201100005420017). Xiangxi Wang was supported by Ten
Thousand Talent Program and the NSFS Innovative Research Group (No. 81921005).
Kang Wang was supported by the Special Research Assistant Project of the Chinese
Academy of Medical Sciences.

Author contributions X.W., K.W., C.F.Q., C.Q., and Y.W designed the whole
study; K.W., Z.J., L.B., L.C., H.C., Y.H., Q.L., Y.J., Q.Z., Y.D., L.W., M.L., Y.L.,
K.F., P.Y., X.P., Z.C., L.Q., P.G., J.W., S.L., Y.C., W.H. performed experiments;
K.W., L.W., P.L., N.W., W.F. H.C. prepared the Cryo-EM samples and solved the
structures; all authors analyzed data; X.W., K.W., C-F.Q, C.Q., and Y.W. wrote the
manuscript with input from all co-authors.

Competing interests: All authors have no competing interests.

689 **Figure legends**

Fig. 1| Evolution and neutralization characteristics of Omicron variant. **a**, , A linear representation of Omicron spike with mutations marked on. The replacements are marked in red; deletions are in grey and insertions are in purple. **b**, Distribution of mutations of Omicron on the cryo-EM structure of pre-fusion spike trimer. The mutations listed in **a** are indicated in the 'up' protomer shown as cartoon with mutated residues highlighted as spheres and colored as in **a**. The RBD, NTD, SD1 and S2 of this subunit are marked with arrow and colored in green, blue, magenta and yellow, respectively; the other two protomers in 'down' state are shown as surface in pale cyan and pale yellow, respectively. **c**, Graph shows the neutralizing antibody response against wild-type and Omicron SARS-CoV-2 authentic virus for sera from healthy vaccinees who received two doses (n=60) or three doses (n=60) of Coronavac. Neutralizing antibody titer fold decline for Delta or Omicron over wild-type for each group of sera is shown in each of the plots.

Fig. 2| Characteristics of a subset of broadly neutralizing antibodies from recipients of a booster immunization. a, Vertical slices chart shows the gross binding epitope distribution of mAbs isolated from the individuals who received three doses of inactivated vaccines. Total number of antibodies and the percentage of antibodies that recognize RBD (blue), NTD (red) and S2 domain (yellow) are indicated. **b,** Heatmap representation of 41 selected representative mAbs and another 9 mAbs approved by FDA or in clinical trials against pseudotyped viruses with wild-type or variant SARS-CoV-2 S. The color bar on the right represents the ranges of IC_{50} values for the indicated mAbs against pseudotyped viruses in **c** (yellow: 0.002-0.020 ng/µL; green: 0.020-1.000 ng/µL; red: 1.000-10.000 ng/µL). Data marked with '*' represents the data referred from the available publication in which the whole experiment system and condition of pseudovirus neutralization assay remains the

samewith this study [18]; Data marked with '-' means no related datasets here. **c**,

Heatmap with values shown in the form of AUC represents the competition ability

between the selected mAbs and hACE2. Color gradient ranging from white (1) to blue

(24) is shown on the right represents the competition ability from the weakest to the

strongest. **d**, Neutralization curves for the selected 41 antibodies on pseudotyped

viruses with the S protein of Omicron variant of concern. Data shown here are three

groups of antibodies: 1) ultrapotent antibodies against all five VOCs, 2) highly

potent antibodies against other four VOCs, but with median neutralizing activities

against Omicron, 3) highly potent antibodies against other four VOCs, but with

weak neutralizing activities against Omicron. XGv347, XGv282 and XGv265,

selected as a representative of each group are highlighted by bold curve in yellow,

green, and red, respectively, in correspondence with the color range in **e**. Heatmap

representation of the same 41 mAbs as in **b** against wild-type and variant SARS-CoV-

2 authentic viruses with color shown on the right.

Fig. 3| Structural basis of the broad and potent neutralization of representative

antibodies. a. Cryo-EM maps and the binding modes of SARS-CoV-2 omicron S

trimer in complex with XGv347 (top left), XGv289 (top right), XGv282 (bottom left)

and XGv265 (bottom right). The three states of XGv347 binding to S-Omicron RBD

are marked state I (one open and one closed RBD), state II (three closed RBDs) and

state III (two open RBDs). Cartoon representations of the structure of SARS-CoV-2

Omicron-RBD in complex with the four antibodies are zoomed in. **b**. Interactions

between the four antibodies and SARS-CoV-2 Omicron RBD. The CDRs of the four

antibodies that interact with SARS-CoV-2 Omicron RBD are displayed as cartoon

over the light green surface of the RBD. The mutation sites on RBD of Omicron are

colored red, the epitopes of antibodies are colored deep green and the overlap of them

are colored in blue. **c**. Interactions details between antibodies (XGv347, XGv289,

XGv282 and XGv265) and SARS-CoV-2 Omicron (top) and WT RBD (bottom). All

the WT structures are predicted with GROMACS. Hydrophobic patches and hydrogen

bonds are highlighted with surface and dash lines. Color scheme is the same as in **a**.

Fig. 4| Protection against SARS-CoV-2 Beta variant strain challenge in mice. a, Experimental design. Groups of BALB/c mice were infected intranasally with SARS-CoV-2 Beta variant strain, followed by a single dose of an antibody, or an antibody cocktail, or PBS as control one hour after infection as indicated in **a**. **b to d,** Examination of lung tissues of mice collected at 5 dpi for **b**, virus titer, **c**, Immunostaining and **d**, H&E. **b**, Virus RNA loads in the lungs at 5 dpi were measured by RT-qPCR and are expressed as RNA copies per gram. Data are represented as mean ± SD. Dashed line represents limit of detection. **c**, SARS-CoV-2 genome RNA ISH was performed with a SARS-CoV-2 specific probe. Brown-colored staining indicates positive results. Scale bar, 200 μm. **d**, Histopathological analysis of lung samples at 5 dpi. Scale bar: 100 μm.

Extended Data

Extended Data Fig. 1 | Fab-ACE2 Competition ELISA assay. Data shown are the curves of 31 antibodies used to compete with ACE2

		Class I				Class II		Class III		Class IV			Class V			Class VI			NC
		XGv013	XGv026	P4A1	XG017	414	H4	XGv031	P17	XGv016	S309	XG014	XGv030	XG025	FC08	XGv004	XG011	A34-2	Fc05
XGv051	1	4	3	3	4	3	9	5	6	94	89	73	71	85	49	325	117	22	101
XGv052	1	3	3	3	3	3	6	4	5	76	55	79	71	86	59	87	66	22	90
XGv053	1	3	5	3	3	3	5	4	7	75	60	76	77	74	57	84	61	22	89
XGv055	1	9	3	3	4	3	9	5	6	89	105	78	83	88	67	321	133	14	95
XGv074	1	5	4	5	5	4	8	7	19	91	61	107	77	89	87	112	80	48	97
XGv253	1	3	3	4	3	3	7	5	5	96	65	80	86	86	81	163	130	145	81
XGv261	1	5	4	3	4	3	9	8	12	91	79	78	71	75	78	272	118	25	94
XGv302	1	3	3	3	4	3	8	7	11	74	73	90	81	90	86	238	144	44	82
XGv303	1	4	4	3	4	3	7	8	13	85	78	92	86	96	89	254	148	47	69
XGv304	1	4	4	4	4	3	7	10	19	89	81	98	83	101	89	237	128	46	99
XGv387	1	6	6	5	4	5	8	15	28	80	63	75	67	60	95	95	74	23	145
XGv177	2	13	18	5	5	5	9	12	27	52	54	67	57	60	87	79	64	24	93
XGv285	2	69	38	78	84	7	10	85	94	43	71	14	72	84	96	94	81	161	107
XGv286	2	41	78	69	73	3	5	80	91	4	88	5	83	77	96	70	90	143	88
XGv287	2	115	81	78	81	15	41	118	104	69	106	32	79	96	96	164	100	133	89
XGv288	2	66	73	69	76	3	7	94	81	5	72	5	69	81	87	77	76	114	91
XGv290	2	65	73	51	76	4	11	81	82	9	60	6	54	77	94	71	69	97	101
XGv291	2	46	72	69	76	3	6	80	83	5	62	4	83	72	92	68	71	127	85
XGv292	2	69	77	62	79	3	5	74	86	7	97	5	81	77	86	75	63	107	86
XGv293	2	42	84	68	80	3	6	88	85	4	79	4	82	82	92	79	73	104	90
XGv294	2	64	78	71	81	3	6	87	88	8	61	5	76	73	87	79	72	96	88
XGv295	2	94	86	74	79	5	11	125	91	9	98	5	91	96	101	280	102	104	106
XGv297	2	57	81	75	76	3	7	101	85	4	76	4	79	72	85	73	79	131	91
XGv338	2	88	85	82	83	3	5	5	4	82	63	47	80	61	5	95	78	100	99
XGv347	2	3	3	3	3	3	7	4	5	88	84	86	96	91	85	256	149	131	97
XGv402	2	9	12	5	7	4	8	23	36	89	63	85	69	73	89	92	74	39	107
XGv420	2	131	88	88	93	3	9	120	51	25	12	5	75	46	64	318	126	57	100
XGv337	3	109	66	77	82	3	9	8	5	79	64	38	67	53	5	277	129	132	94
XGv264	4	63	79	73	84	3	6	93	86	4	5	4	86	72	79	82	84	134	86
XGv265	4	73	77	62	83	3	6	87	86	16	7	5	74	64	80	74	76	108	89
XGv266	4	73	78	87	87	3	5	91	80	12	6	4	85	64	83	86	76	127	86
XGv282	4	81	79	105	77	3	5	83	10	5	43	4	93	76	11	70	89	106	88
XGv289	4	57	78	74	72	3	7	91	77	8	71	5	87	72	82	67	86	143	87
XGv296	4	51	74	64	67	3	6	86	87	5	79	5	80	64	85	67	72	122	87
XGv345	4	88	95	87	92	3	7	97	31	22	7	4	76	12	14	248	132	121	65

Extended Data Fig. 2 | Data Sheets of ELISA assay of representative Mabs against Omicron RBD. Different Classes of Nabs (Class I-VI) are colored by yellow, green, red, blue, brown and magenta, respectively. Values are filled with black (>75), grey (50-75), silver (25-50) and white (<25).

a

b

RELION

3,955 Micrographs

Gctf
AutoPick

233,2045 Particles

cryoSPARC

2D Classification

3D Classification

RELION

163,246 particles
19.50%

66,470 particles
7.94%

401,170 particles
47.84%

118,458 particles
14.15%

88,488 particles
10.57%

cryoSPARC

NU-Refinement

3.6 Å

Local Refinement

3.8 Å

d

Extended Data Fig. 3 | Flowcharts for cryo-EM data processing. Flowcharts for

Omicron Spike protein in complex with **a**, Fab XGv347, **b**, XGv289, **c**, XGv282 and

d, XGv265 are shown.

Extended Data Fig. 4 | Resolution Estimation of the EM maps. a, The gold-standard FSC curves of overall maps of Omicron S trimer in complex with Fab XGv347, XGv289, XGv282 and XGv265 and local maps of interfaces. **b,** Local resolution assessments of cryo-EM maps using ResMap are shown.

796
797 **Extended Data Fig. 5 |Density maps and atomics models**. Cryo-EM maps of
798 Omicron S trimer in complex with Fab XGv347, XGv289, XGv282 and XGv265 and
799 their interfaces are shown. Color scheme is the same as in Fig. 3a. Residues are
800 shown as sticks with oxygen colored in red, nitrogen colored in blue and sulfurs
801 colored in yellow.
802

Extended Data Fig. 6 | Mechanism of XGv347 binding to 3 closed RBD. a,
Superimposition of Omicron to WT Spike Trimer. Omicron Spike is colored in cyan
and wild-type Spike is colored in yellow. **b,** Superimposition of Fab A23-58 onto
Omicron and WT Spike trimer and Fab XGv347 onto WT Spike trimer and are shown
as surface. Fab A23-58 is colored in magenta and XGv347 is colored in purple.

Extended Data Fig. 7 | Binding modes of XGv289, 282 and 265. Binding modes of XGv289, XGv282 and XGv 265. RBD is colored in light cyan and color scheme of XGv289, XGv282 and XGv265 is the same as in Fig. 3a. DH1047, BD-812 and REGN10987 are colored in orange, deep pink and blue, respectively.

XGv289-347   XGv282-347   XGv265-347

817 **Extended Data Fig. 8 | Potential cocktail representation.** XGv265, XGv282 and

818 XGv289 are superimposed onto XGv347 and all structure are shown as surface.

Extended Data Table. 1 | Statistics for Cryo-EM data collection, refinement, and validation

Name	XGv347 in complex with S (state 1)	XGv347 in complex with S (state 2)	XGv347 in complex with S (state 3)	XGv347-RBD-interface	XGv289 in complex with S	XGv289-RBD-interface	XGv282 in complex with S	XGv265 in complex with S	XGv2657-RBD-interface
Data collection									
Microscope	FEI Titan Krios	FEI Titan Krios	FEI Titan Krios	FEI Titan Krios	FEI Titan Krios	FEI Titan Krios	FEI Titan Krios	FEI Titan Krios	FEI Titan Krios
Camera	K3	K3	K3	K3	K3	K3	K2	K3	K3
Voltage (kV)	300	300	300	300	300	300	300	300	300
Tatal dose (e⁻/Å)	60	60	60	60	60	60	60	60	60
Symmetry imposed	C1	C3	C1	C1	C1	C1	C1	C1	C1
Data Processing									
Micrographs (Total)	5014	5014	5014	5014	3955	3955	2631	3752	3752
Micrographs (Used)	4561	4561	4561	4561	3864	3864	2598	3654	3654
Particles selected	2320416	2320416	2320416	2320416	2332045	2332045	756508	1302103	1302103
Particles included in final reconstruction	269947	85822	105455	527413	401170	401170	119800	138359	138359
Reconstruction									
Pixel size (Å)	1.07	1.07	1.07	1.07	1.07	1.07	1.04	1.07	1.07
Defocus range (μm)	-1.5--2.5	-1.5--2.5	-1.5--2.5	-1.5--2.5	-1.5--2.5	-1.5--2.5	-1.5--2.5	-1.5--2.5	-1.5--2.5
Resolution (Å) (FSC = 0.143)	3.3	3.3	3.7	3.5	3.6	3.8	3.8	3.5	4
Model Refinement									
Clashscore	9.48	9.82	11.25	6.98	11.95	8.46	13.06	9.59	8.2
Rotamer outliers (%)	0.00	0.03	0.00	0	0.03	0	0.00	0.03	0
Molprobity score	1.90	1.94	1.95	1.88	2.09	1.93	1.94	1.86	1.76
Ramachandran statistics (%)									
Most favored (%)	94.06	93.46	94.31	91.76	91.69	92.49	95.42	94.93	95.51
Allowed (%)	5.83	6.51	5.67	8.24	8.23	7.51	4.56	5.07	4.49
Outliers (%)	0.11	0.03	0.03	0	0.08	0	0.03	0	0
R.m.s.deviations									
Bond lengths (Å)	0.003	0.01	0.003	0.003	0.004	0.003	0.002	0.002	0.005
Bond angles (°)	0.714	0.666	0.567	0.7	0.62	0.624	0.557	0.576	0.72

824 **Extended Data Table. 2** |Residues of Fabs interacting with the Omicron SARS-CoV-
825 2 S trimer at the binding interface (d < 4 Å)

Omicron RBD	XGv265-Heavy chain				XGv265 Light chain		
ARG 346	Y32						
ASN 439					Y35		
ASP 442	Y32						
SER 443	I102						
LYS 444	Y54	D56	D58	W55			
VAL 445	Y54	R60	L52		T99		
SER 446	R60						
GLY 447	R60						
PRO 499					Y35	Y94	
ARG 509	Y32						

XGv265 in complex with Omicron S

Omicron RBD	XGv289 Heavy chain				XGv289 Light chain		
PHE 374					N32		
PHE 375					Y33		
ASN 439	S101				D95		
LYS 440	S102	S101			Y33		
SER 443	S101						
VAL 445	A57						
SER 446	A57	S58	G56				
PRO 499	S101						
THR 500	N62	A60	Q61		G99	S98	
GLY 502	N62				L97	S98	
VAL 503					D95	S96	L97

XGv289 in complex with Omicron S

Omicron RBD	XGv347 Heavy chain				XGv347 Light chain		
LEU 455	D31						
PHE 456	D31	V32					
TYR 473	T105						
ALA 475	S106						
GLY 476	C107						
LYS 478	D109						
GLY 485	W51						
PHE 486	P100	S108	D109	F111	Y33		
ASN 487	S108						
TYR 489	V32	S34	V53				
ARG 493	G55	T56					

XGv347 in complex with Omicron S

Omicron RBD	XGv282 Heavy chain				XGv282 Light chain	
R346	F103					
K444	S101	G102	F106			
V445					W92	
S446	R50	F106	Y107		W92	
G447	R50	F105	F106			
N448	G102	F103	F106			
Y449	I52	I59	F105	F106		
N450	G102	F103	F105	F106		
Y451	F103					
L452	I54	V55				
F490	V55					
L492	V55	K57				
R493	V55	K57				
S494	I52	K57				
R498					W92	

XGv282 in complex with Omicron S

826
827

828 **Extended Data Table. 3 | Statistics for Molecular Dynamics**

	SARS-CoV-2 Variant	KD (nM)	ΔG (kcal/mol)	ΔΔG (kcal/mol)	No.(residue$_{TOTAL}$)	No.(residue$_{RBD}$)	No.(residue$_{fab}$)	No. (HB or SB)	No.(nonpolar residue$_{RBD}$)	No.(nonpolar residue$_{fab}$)
XGv265	WT	1.475	-3.99	-0.96	21	10	11	14	4	7
	Omicron	28.52	-3.03		21	10	11	9	3	7
XGv282	WT	0.8612	-3.79	-1.82	28	15	13	13	7	8
	Omicron	4.096	-1.97		19	8	11	7	3	8
XGv289	WT	1.287	-5.94	-0.79	21	9	12	16	4	6
	Omicron	14.17	-5.15		26	11	15	12	3	4
XGv347	WT	0.1518	-5.42	-0.14	23	10	13	15	4	5
	Omicron	6.812	-5.28		26	11	15	9	4	5

829
830

831