1

Title: Optogenetic frequency scrambling of hippocampal theta oscillations dissociates working memory retrieval from hippocampal spatiotemporal codes

Authors: Guillaume Etter[1,*], Suzanne van der Veldt[1], Jisoo Choi[1], Sylvain Williams[1,*]

[1]McGill University & Douglas Mental Health University Institute, Montreal, Canada

* corresponding authors: sylvain.williams@mcgill.ca, 6875 Boulevard LaSalle, Verdun, QC H4H 1R3, Phone: 514 761-6131, Fax: 514 888-4067; etterguillaume@gmail.com

Abstract

The precise temporal coordination of activity in the brain is thought to be fundamental for

memory encoding and retrieval. Pacemaker GABAergic neurons in the medial septum (MS)

provide the largest source of innervation to the hippocampus and play a major role in

controlling hippocampal theta (~8 Hz) oscillations. While pharmacological inhibition of the

MS is known to disrupt memory, the exact role of MS inhibitory neurons and theta frequency

rhythms in hippocampal representations and memory is not fully understood. Here, we

dissociate the role of theta rhythms in spatiotemporal coding and memory using an all-optical

interrogation and recording approach in freely behaving mice. We propose a novel paradigm

to dissociate encoding of space, time and distance in freely moving mice and apply

complementary optogenetic stimulation paradigms of MS GABAergic neurons to either pace

or abolish theta altogether while recording large hippocampal cell assemblies using calcium

imaging conjointly. We first show that optogenetic frequency scrambling of MS GABAergic

neuron activity abolished theta rhythms and modulated the activity of a subpopulation of CA1

neurons. Such stimulation led to decreased memory retrieval in both a delayed non-match to

sample task, a novel place object recognition task, as well as spontaneous cue-guided linear

alternation. Strikingly, scrambled stimulations were not associated with disrupted encoding of

place, time, distance, or multiplexed information. Our study suggests that theta rhythms play a

specific and essential role in supporting working memory retrieval and maintenance while not

being necessary for hippocampal spatiotemporal codes.

Introduction

The precise temporal coordination of neuronal activity is thought to be fundamental for memory encoding and retrieval. In particular, the medial septum (MS) has been proposed to act as the main zeitgeber to downstream structures[1]. It is a neurochemically heterogeneous structure composed of GABAergic[2], mainly parvalbumin (PV)-positive neurons[3], along with cholinergic[4] and a smaller population of glutamatergic neurons[5,6]. MS PV cells project directly onto GABAergic interneurons in the hippocampus giving rise to feedforward inhibitory control of hippocampal pyramidal cells[3]. Additionally, PV interneurons within the hippocampus are essential in pacing theta (~8 Hz) rhythms[7], and optogenetic stimulation of MS PV neurons directly[8,9] or of their terminals[10] in the hippocampus are associated with frequency-specific pacing of hippocampal oscillations whereas inhibition of the medial septum (MS) *in vivo* has been associated with reduced theta oscillations power[11,12].

While complete MS optogenetic inhibition has been associated with spatial memory impairments[13], these effects could potentially be attributed to disruption of cholinergic functions[14,15], which are known to be critical for memory function. The exact role of non-cholinergic MS PV cells in memory encoding and retrieval remains an active area of research and has only been described recently[16,17]. Importantly, the hippocampus plays an essential role in encoding and retrieval of episodic[19,20] and working[21] memory. Since the MS provides the largest subcortical inputs to the hippocampus and is essential in generating and maintaining theta rhythms, disruption of MS activity is likely to impact downstream hippocampal physiology and working memory[18]. Although the exact physiological mechanisms of hippocampus-dependent memory are currently unknown, it has been proposed that hippocampal place cells[22] that encode specific locations of a given context could support episodic memory[23]. In the hippocampal subfield CA1, spatial tuning relies at least in part on contextual sensory inputs[24]. Other variables such as time and distance can also be encoded

during visually guided locomotion[25,26] but also in the absence of sensory cues[27], likely using

internal information including self-motion[28] (see Mehta[29] and McNaughton[30] for review).

Representations of time and space can be represented conjunctively in hippocampal neurons,

and such multiplexed spatiotemporal codes could be a candidate substrate for working

memory[31–33]. In addition to clamping distal visual cues using a treadmill[34–36] or using virtual

reality paradigms[25,26,37], spatiotemporal codes have also been extracted analytically using

generalized linear models that implement space, time, and distance[36] although such

approaches have not been employed extensively on recordings of neuronal activity during free

exploration.

Several studies suggest that hippocampal theta rhythms could underlie temporal codes

since theta rhythms tightly orchestrate hippocampal activity[38,39]. While time cells have also

been reported in both CA1 and CA3 of rodents performing tasks that do not require working

memory[40], pharmacological inhibition of the MS results in specific disruption of time but not

place cells and is associated with decreased working memory[35]. An important drawback of

pharmacological approaches is that they do not distinguish the relative contribution of

GABAergic versus cholinergic cells to memory function[41], and inhibition of MS cholinergic

activity was found to alter hippocampal spatial representations[42] and decrease working

memory performance[43,44]. Surprisingly, pharmacological inhibition of the MS was associated

with reduced theta oscillation power but not place fields[45], and this resilience of place cell

activity during diminished theta was not due to experience-related plasticity mechanisms[46].

Previous attempts at inhibiting MS GABAergic interneurons specifically using optogenetics

were associated with only a partial reduction, but not complete disruption of theta signals[47]. In

turn, optogenetic pacing of theta oscillation has only been associated with minor changes in

place cell characteristics, including a slight shift in firing frequency[9] and phase[48].

Additionally, while it is hypothesized that hippocampal temporal codes could be directly

controlled by MS inputs, causal evidence is still lacking. To this day, the exact role of MS-PV

neurons in orchestrating hippocampal spatiotemporal codes and memory remains unknown.

Here we controlled MS PV activity using optogenetics to pace or abolish theta

oscillations using a red-shifted excitatory opsin. We propose a novel approach to completely

abolish hippocampal theta rhythms based on optogenetic frequency scrambling stimulations

of MS neurons. Alternatively, pacing theta rhythms at their natural frequency in the same

animals provides within-subject controls. We combined optogenetic control with calcium

imaging of CA1 pyramidal cells in mice running on a linear track with sequential tones. In

these conditions, we were able to separate place, time, and distance cells using an information

theoretic approach. When performing optogenetic frequency scrambling of theta signals, both

place and time representations were preserved and only a small subset of CA1 pyramidal cells

were modulated by stimulation. We next found that ablation of theta oscillation was

associated with impaired working memory retrieval suggesting that MS PV cells play a key

role in generating hippocampal theta oscillations that are necessary for memory retrieval, but

are not involved in spatiotemporal representations.

Results

CA1 pyramidal cells encode spatiotemporal information

To examine the spatiotemporal codes in large populations of CA1 principal cells, we injected

a viral vector expressing GCaMP6$_{fast}$ under a CamKII promoter in the CA1 region of the

hippocampus, implanted a GRIN lens above the injection site, and performed calcium

imaging recordings of pyramidal neurons using open-source miniscopes[49,50] (fig. 1a,b; see

supplemental fig. 1 for detailed histology). We extracted spatial footprints of neurons (fig. 1c)

and their corresponding calcium transients (fig. 1d) using CNMFe[51]. To tease apart the spatial

and temporal tuning properties of principal cells, we developed a new task combining a linear

track with three tone cues triggered by motion sensors at both ends of the track. A new tone

was instantly triggered at the end of each run, informing the mice on its progression toward

reward delivery. Every fourth run was cued with a high-pitched, continuous tone that was

associated with the delivery of a reward at the end of the linear track (fig. 1e). The absolute

location of each mouse, along with the time elapsed and distance traveled since the departure

from the reward site were monitored (fig. 1f). Using these variables and binarized neuronal

activity, we computed probabilistic tuning curves (fig. 1g) and derived mutual information

(MI) between neuronal activity and location, time, as well as the distance for each recorded

cell. In contrast to correlation-based analyses, MI does not assume linear, monotonic

relationships between neuronal activity and behavioral variables but rather expresses the

amount of uncertainty of one variable that can be explained by the other. The significance of

MI values was tested using shuffled surrogates that underwent circular permutations (n =

1000) to preserve temporal dynamics of calcium transients. Neurons that encoded exclusively

one variable with a MI greater than shuffled surrogates 99.5 of the time ($p < 0.05$) were

labeled as either place-modulated (spatial), time-modulated (temporal), or distance-modulated

(see Methods). For the following analyses, we focused on candidate cells that only encoded

one significant variable (fig. 1h). Importantly, time-modulated cells were not systematically

active at particular locations, and place-modulated cells were not systematically active at a

given time (fig. 1i,j). While the majority of cells encoding a single variable were place-

modulated, a large portion of neurons were conjunctive neurons that encoded more than one

variable (17.98 ± 1.77 %). In contrast, cells encoding place exclusively represented 9.02 ±

1.61 % of the total recorded population, while 1.79 ± 0.68 % selectively encoded distance and

1.27 ± 0.09 % selectively encoded time (fig. 1k; additional examples of neurons tuned to time,

space, or distance along with their information content are shown in supplemental fig. 2).

Although our information theoretic approach can disentangle overlapping variables by

isolating cells that only significantly encode one variable, we further tested the relevance of

each cell type in encoding spatiotemporal variables using a naive Bayesian classifier to

decode location (fig. 1l-n), time elapsed (fig. 1o-q), and distance traveled (fig. 1r-t) on the

linear track[52]. We estimated the current state of each mouse by computing the maximum *a*

posteriori (MAP) value given neuronal activity and bootstrapped tuning curves computed

using either actual or circularly shuffled binarized activity (fig. 1l,o,r; see Methods for

detailed protocol). The quality of predictions was assessed using confusion matrices (fig.

1m,p,s) and by computing the euclidean distance between predicted state and actual state (fig.

1n,q,t). Importantly, our Bayesian decoder yielded an average error of 16.58 cm, which was

significantly lower than when decoding from shuffled surrogates (50.95 cm; t-test, $t_{98} = 127.3$,

$p < 0.0001$), and decoding using spatially modulated cells was significantly more accurate

than when using non-spatially modulated cells (t-test, $t_{98} = 29.43$, $p < 0.0001$; fig. 1n).

Similarly, the average decoding error for time elapsed was 6.45 s, which was significantly

lower than error computed using shuffled surrogates (19.12 s; t-test, $t_{98} = 101.5$, $p < 0.0001$).

Decoding using time-modulated cells yielded significantly better accuracy compared to non

time-modulated cells (t-test, $t_{98} = 10.63$, $p < 0.0001$; fig. 1q). Finally, average distance error

using our decoder was 61.40 cm, which was significantly lower than that of shuffled

surrogates (189.6 cm; t-test, $t_{98} = 141.1$, $p < 0.0001$). Decoding using distance-modulated cells

yielded significantly lower errors compared to non distance-modulated cells (t-test, $t_{98} = 9.58$,

$p < 0.0001$; fig. 1t).

Selective MS optogenetic control of theta oscillations

To examine the relative contribution of MS-generated theta signals to hippocampal

spatiotemporal codes, we transfected the red-shifted excitatory opsin ChrimsonR in the MS

(fig. 2a). In contrast to inhibitory opsins, ChrimsonR allowed us to either scramble or pace

theta signals within subjects. Additionally, ChrimsonR is more effective than the more widely

used Channelrhodopsin-2 and provides the capability to combine optogenetics with calcium

imaging[53]. 14.91 ± 3.57 % of PV cells expressed ChrimsonR which we found sufficient to

exert pervasive control over hippocampal oscillations (n = 4 mice; fig. 2b,c). In contrast, we

found virtually no expression of ChrimsonR in ChAT cells (1.05 ± 1.052 % of ChAT cells

also expressed ChrimsonR; fig. 2d,e).

We then implanted mice with fiber optics over the MS and performed 638 nm laser

stimulation while recording local field potentials (LFP) in CA1 (fig. 2f). We found that MS

scrambled and 8 Hz optogenetic stimulation could disrupt or pace theta oscillations,

respectively (fig. 2g). While baseline natural theta displays some frequency variability in the

4-12 Hz frequency band, 8 Hz stimulations led to consistent and stable hippocampal

oscillations at that frequency, whereas scrambled stimulations consistently abolished theta

rhythms (fig. 2h). We found that oscillation strength (OS) in the theta band (see Methods) was

significantly decreased by scrambled stimulations (0.45 ± 0.01 OS) compared to baseline

epochs (0.67 ± 0.01 OS, p < 0.0001) and were not significantly different from OS of white

noise control signal (0.50 ± 0.01 OS, p = 0.99). On the other hand, 8 Hz stimulations

increased theta power significantly (0.82 ± 0.01 OS, p < 0.0001; n = 60 epochs) compared to

scrambled stimulations. We also found a significant interaction between our stimulation

patterns and LFP frequency band (F_{10} = 6.467, p < 0.0001). In particular, scrambled frequency

stimulation significantly decreased theta power (0.341 ± 0.06 portion of baseline theta band

power; p = 0.0394, pairwise t-test), while 8 Hz stimulations significantly increased theta

power (3.302 ± 0.76 portion of baseline theta band power, $p = 0.0004$, pairwise t-test; fig. 2j)

leaving other frequency bands unaltered.

While our calcium imaging and electrophysiological analyses only included periods of

locomotion (see Methods), we also found that we were able to reliably abolish (supplemental

fig. 3a,b) or pace (supplemental fig. 3a,c) theta oscillations regardless of locomotor state

(including periods of restfulness). While natural theta oscillation strength is correlated to

locomotor speed (Pearson $R^2 = 0.059$, $p = 0.001$; supplemental fig. 3d), abolishing theta led to

a loss of such correlation (Pearson $R^2 = 0.008$, $p = 0.223$; supplemental fig. 3e), as did 8 Hz

stimulations (Pearson $R^2 = 0.0008$, $p = 0.714$; supplemental fig. 3f) suggesting that locomotor

states did not override the effects of optogenetic stimulations on theta oscillations.

Hippocampal sharp-wave ripples (SWRs) play an essential role in memory

consolidation[54–56] and stimulation of MS cholinergic neurons has been associated with reduced

ripple activity[57] and impaired working memory[44]. While we found virtually no expression of

ChrimsonR in MS cholinergic neurons, it was essential to measure the impact of our MS

optogenetic stimulation on ripple physiology. To this end, we recorded CA1-LFP and

performed 5s ON-OFF scrambled optogenetic stimulation in freely behaving mice exploring

an open-field (supplemental fig. 4a). We measured attributes of ripple events before and

during scrambled stimulation, and found no changes in power (t-test, $t_6 = 0.076$, $p = 0.941$;

supplemental fig. 4b, left panel), frequency of occurrence (t-test, $t_6 = -1.688$, $p = 0.142$;

supplemental fig. 4c, left panel), or width (t-test, $t_6 = 0.124$, $p = 0.905$; supplemental fig. 4d,

left panel). Similarly, applying 8 Hz optogenetic stimulation had no discernable effects on

ripple power (t-test, $t_6 = 0.378$, $p = 0.718$; supplemental fig. 4b, right panel), frequency (t-test,

$t_6 = -1.643$, $p = 0.151$; supplemental fig. 4c, right panel), and width (t-test, $t_6 = -0.138$, $p = $

0.894; supplemental fig. 4d, right panel). Together with our histological results and previous

reports that optogenetic stimulation of MS cholinergic neurons reduces the occurrence of

SWRs[57], our findings support the notion that these optogenetic stimulations are not affecting

cholinergic inputs to the hippocampus.

Combining MS optogenetic stimulation and hippocampal calcium imaging

To study the effects of theta manipulations on hippocampal spatial and temporal codes, we

combined MS optogenetic stimulation and calcium imaging in CA1. This experimental

paradigm raises two potentially important issues: GRIN lens implants involve tissue damage

that could alter the physiological state of theta oscillations, and the wavelength spectrum of

the excitation LED used to excite GCaMP6f could potentially overlap with that of the opsin

located on the terminal fibers of GABAergic MS fibers in the hippocampus.

To verify whether GRIN lens implants altered the physiology of theta, we first

implanted mice with a GRIN lens in the right hippocampus, with two electrodes in both left

and right hippocampi and found comparable theta signals in both hippocampi (fig. 3a). To

gain more statistical insights, we compared theta oscillations mice with electrodes only in the

right hippocampus, to mice with both a GRIN lens and an attached LFP electrode during open

field exploration, and found no significant differences between both groups (fig. 3b). We did

not find any significant difference between relative theta power to total LFP power of mice

implanted with a GRIN lens and LFP electrode (0.14 ± 0.007) versus an electrode only (0.154

± 0.008; t-test, $t_{54} = 1.060$, $p = 0.29$).

Although previous reports have reported that combining optogenetic stimulation of

ChrimsonR in cell bodies with calcium imaging of neurons at the vicinity of terminals is

possible with minimal crosstalk[53], we next controlled for any potential opsin activation of MS

terminals in the hippocampus by recording CA1-LFP while emitting excitation light with our

miniscope through a GRIN lens in mice injected with ChrimsonR in the MS. After calibrating

miniscope light output power (fig. 3c), we found no effect of miniscopes blue excitation light

on endogenous theta power when using excitation light power below 0.5 mW/mm^2

(1ANOVA, $F_{(4,295)}$ = 0.7729, p = 0.5435; fig. 3d). Since it has been reported that miniscope

excitation light can induce slight depolarization of terminals transfected with ChrimsonR[53],

thus hindering further optogenetic-induced depolarization, we next stimulated our control

mice while imaging with a ~0.3 mW/mm^2 miniscope LED power and were able to

significantly disrupt or pace theta using scrambled or 8 Hz stimulation respectively (Friedman

test $\chi^2 = 6, p = 0.0278$; fig. 3e).

Disruption of theta rhythms modulates a small portion of CA1 cells

We then performed phasic (5s ON, 5s OFF) optogenetic stimulations of MS while recording

CA1 pyramidal cells and as mice freely explored an open-field (fig. 4a). We found that a

portion of recorded cells were consistently excited in these conditions, while others were

inhibited (fig. 4b; see Methods). The activity of pyramidal cells during stimulation was

overall lower compared to baseline, for both scrambled stimulation during running (Pearson

correlation, R^2 = 0.567., p < 0.0001) and rest (R^2 = 0.521, p < 0.0001) periods, as well as for 8

Hz stimulation (R^2 = 0.6., p < 0.0001 for rest periods; R^2 = 0.632., p < 0.0001 for running

periods; n = 1849 cells, N = 5 mice; fig. 4c). Overall, ~6.42 ± 0.52 % of total cells were

significantly modulated by scrambled optogenetic stimulation (fig. 4d). Among those

modulated cells, 50.56 ± 6.38 % were inhibited while 49.43 ± 6.38 % were excited during

scrambled stimulations (n = 1849 cells, N = 5 mice; fig. 4e).

We next analyzed the effects of optogenetic stimulation on spatial tuning of

hippocampal neurons as mice freely explored the open field. To this end we computed activity

rate maps by using either epochs within or outside of stimulation periods (for the baseline

condition, we included epochs that followed the same 5s ON, 5s OFF pattern used for actual

stimulation; fig. 4f). Stability was then computed as the correlation between rate maps

computed for baseline and stimulation epochs. In spite of the aforementioned changes of

overall activity, rate maps displayed no change of spatial stability for either scrambled or 8 hz

stimulations (Kruskal-Wallis $H_3 = 3.5$, $p = 0.1773$; fig. 4g).

MS optogenetic stimulation alters behavior but not spatiotemporal codes

To assess the effects of theta disruption on temporal and spatial codes, we monitored CA1

pyramidal neuron activity on the 3-tone linear track (fig. 5a). Here we leverage one of the

main advantages of calcium imaging, which is the ability to register recorded cells over

several days, while performing scrambled or 8 Hz stimulations on selected days (fig. 5b; fig.

5c, bottom panel). We focused our analysis on pairs of days with the same amount of time

(48h) between testing (fig. 5c, top panel). For each condition, we assessed the portion of total

cells significantly encoding one or several variables, and found no effect of either 8 Hz or

scrambled stimulation on spatial and temporal encoding (RM-ANOVA; $F_2 = 0.807$, $p = 0.453$

for the main effect of stimulations; $F_6 = 1.283$, $p = 0.285$ for the interaction between

stimulation and encoded variable, n = 5 mice; fig. 5d).

While the portion of cells did not change under stimulation conditions, we assessed

the effect of MS stimulation on place-, time-, and distance-modulated cells' tuning curves

stability. To this end, we tracked neurons across days (fig. 5c; see Methods) and computed

stability of place and time fields as the pairwise correlation between fields across 48h.

Because CA1 is known to display prominent remapping over days[58], we also used a pair of

days with no stimulation to compute a baseline stability score to be used as a reference (fig.

5e-g). We found no change in stability during MS stimulation for place-modulated cells

(1ANOVA, $F_2 = 1.907$, $p = 0.1511$; fig. 5h), time-modulated cells (1ANOVA, $F_2 = 2.201$, $p = $

0.113; fig. 5i), and distance-modulated cells (1ANOVA, $F_2 = 0.6962$, $p = 0.5024$; fig. 5j). We

also extended the same analysis to conjunctive neurons (i.e. neurons that can encode more

than one variable; supplemental fig. 5a-f) and found no effects of optogenetic stimulations on

the stability of conjunctive spatial (1ANOVA, $F_2 = 3.731$, $p = 0.0661$; supplemental fig. 5g),

temporal (1ANOVA, $F_2 = 1.993$, $p = 0.8228$; supplemental fig. 5h), and distance cells

(1ANOVA, $F_2 = 0.469$, $p = 0.6400$; supplemental fig. 5i).

We next assessed the quality of spatiotemporal codes using random bootstrap samples

(n = 160 cells per sample, n = 50 bootstrap samples) of all recorded cells. Namely, we applied

a naive Bayesian classifier to decode location (fig. 6a,b), time (fig. 6c,d), and distance

traveled (fig. 6e,f). Decoded errors were systematically lower than shuffled surrogates,

including during 8 Hz or scrambled stimulation for location (2ANOVA, $F_{5,1100} = 5095$, $p < $

0.0001; fig. 6a), time (2ANOVA, $F_{5,1100} = 2731$, $p < 0.0001$; fig. 6c), and distance (2ANOVA,

$F_{5,1100} = 2671$, $p < 0.0001$; fig. 6e) indicating that spatiotemporal codes were preserved during

stimulation. To gain statistical insights, we z-scored decoding error by using both the actual

and shuffled results (see Methods) for a given day, for every mouse (N = 5 mice; fig. 6b,d,f).

MS optogenetic control did not significantly alter encoding of location (1ANOVA, $F_{2,11} = $

2.2332, $p = 0.1432$; effect size $\eta^2 = 0.29$; fig. 6b), time (1ANOVA, $F_{2,11} = 0.4561$, $p = 0.6452$;

effect size $\eta^2 = 0.07$; fig. 6d), or distance (1ANOVA, $F_{2,11} = 0.6102$, $p = 0.5606$; effect size η^2

= 0.09; fig. 6f).

In the behavioral paradigm used to analyze temporal modulation of neuronal activity,

mice are water-scheduled and expected to collect rewards as quickly as possible. Based on

this assumption, we quantified the number of returns to an empty reward site as errors, and

computed the percentage of correct runs to total runs as a proxy for performance (fig. 6g). In

these conditions we found a significant effect of optogenetic stimulation on performance

across days (Friedman test $\chi^2 = 6.000$, $p = 0.0278$) and in particular a significant difference in

performance between during baseline (78.70 $\pm$ 2.45 %) and scrambled stimulation (44.44 $\pm$

5.56 %; multiple comparison, $p = 0.0429$; N = 3 mice; fig. 6h). Only mice with a minimum of

12 runs were included in this analysis. Because of the limitations of this task (low cognitive

load and low number of mice tested), we next set out to assess the effect of MS optogenetic

stimulation in standardized behavioral tasks. Importantly, it was previously reported that

pacing theta oscillations could decrease locomotor speed and its variability[10]. To thoroughly

assess the specificity of MS optogenetic stimulation on memory, we performed additional

experiments to rule out the direct effects of the optogenetic stimulations on locomotor

velocity.

A subset of mice injected with ChrimsonR and implanted with fiber optics in the MS

as well as LFP electrodes in CA1 were allowed to explore an open-field freely while being

subjected to 5s ON, 5s OFF optogenetic stimulations (supplemental fig. 6a). Consistently, 8

Hz stimulation led to consistent pacing of hippocampal oscillations to that frequency

(supplemental fig. 6b). Those stimulations were not associated with any apparent change in

locomotor behavior (supplemental fig. 6c), including mean speed (t-test, $t_{116} = 0.4140$, $p =$

0.6796; supplemental fig. 6d) and speed coefficient of variation (CV; t-test, $t_{116} = 0.8296$, $p =$

0.4095, n = 59 stimulation epochs; supplemental fig. 6e). Similarly, scrambled stimulation

consistently led to abolished theta oscillations (supplemental fig. 6f) but no apparent changes

in locomotor behavior (supplemental fig. 6g), including mean speed (t-test, $t_{118} = 0.2268$, $p =$

0.8210; supplemental fig. 6h) and speed CV (t-test, $t_{118} = 1.838$, $p = 0.686$, n = 60 stimulation

epochs; supplemental fig. 6i).

While natural theta frequency and locomotor speed are correlated[59], the exact direction

of causation between these variables remains poorly understood. To answer this question, we

performed optogenetic stimulation using a 5s ON, 5s OFF paradigm and a random frequency

was selected for each stimulation epoch (supplemental fig. 6j). These stimulation frequencies

covered the entirety of the theta band spectrum (supplemental fig. 6k). As expected, we found

that natural theta oscillations frequencies are directly correlated to running speed for one

example mouse ($R^2 = 0.1114$, $p < 0.0001$; supplemental fig. 6l). Importantly, when theta

oscillation frequency results from MS optogenetic control, the correlation drops below chance

level ($R^2 = 0.0006779$, $p = 0.6244$; supplemental fig. 6m) suggesting that locomotion dictates

theta frequency, but not the opposite. We systematically replicated these results across mice

and observed a drop in the correlation between theta frequency and locomotor speed under

optogenetic stimulation control (paired t-test, $t_3 = 3.922$, $p = 0.0295$; supplemental fig. 6n).

Altogether, these results support that our optogenetic stimulation does not alter locomotor

behavior, which is highly relevant for the next behavioral assays.

Disruption of theta signals impairs spatial recognition and working memory retrieval

To test the role of theta signals in spatial memory, we prepared a dedicated group of mice

injected with ChrimsonR and implanted with a fiber optic in the MS (fig. 7a, left panel). Mice

were subjected to a novel place object recognition (NPOR) task (fig. 7a, right panel). To

assess the role of theta oscillations in memory encoding and retrieval, we performed

optogenetic stimulations specifically during the sample and test phases and computed the

recognition index (RI; see Methods). When stimulating during retrieval, memory performance

was significantly reduced for both scrambled (0.472 ± 0.048 RI, n = 6 mice) and 8 Hz

stimulations (0.435 ± 0.057 RI, n = 6 mice) groups, compared to YFP controls that displayed

a significant increase in object exploration during testing (0.61 ± 0.018 RI, n = 12 mice;

2ANOVA, $F_{4,31} = 3.283$ for the main effect of treatment, $p = 0.0097$; effect size for the main

effect of group, η^2_p = 0.306 fig. 7b). On the other hand, scrambled stimulations during

encoding did not impair memory at test time (0.60 ± 0.034 RI, p = 0.0157, n = 6 mice) but 8

Hz stimulation during encoding lowered memory performance to chance levels (0.39 ± 0.074

RI, p = 0.8740, n = 6 mice,).

To further examine the effect of optogenetic control of the MS on specific phases of

working memory function (encoding, maintenance, and retrieval), mice were transfected with

ChrimsonR and implanted with fiber optics in the MS and trained in a delayed non-match to

sample (DNMTS) task. In the sample phase, mice were forced to run to a randomly

designated arm to collect a reward. After a delay (10 s), they could either run in the opposite

arm (correct choice) to receive another reward, or run in the same, unrewarded arm (incorrect

arm; fig. 7c). The advantage of this task is to allow for repetitive testing, specific isolation of

task phases (training, delay, testing) and within-subject controls. Mice were trained in this

task with no stimulation until criterion performance of 0.8 (portion of correct trials) was

reached for at least two days. Both ChrimsonR and YFP control mice displayed significant

improvement over time (RMANOVA, F_8 = 8.737, p < 0.0001 for the main effect of training

days). Importantly, we found no learning rate differences between the two groups (pairwise

Tukey; p = 0.42; fig. 7d). Once mice had learned the rule associated with the DNMTS task,

we assessed performance while delivering either scrambled (0.792 ± 0.045) or 8 Hz (0.850 ±

0.036) optogenetic stimulation in the encoding phase (forced choice) only, and did not

observe any difference in performance compared to baseline (0.825 ± 0.030, RM-ANOVA,

F_{11} = 2.197, p = 0.547 , n = 12; fig. 7e). When stimulated only during the delay period, only

scrambled stimulations significantly decreased memory performance (0.733 ± 0.057)

compared to baseline (0.825 ± 0.030, RM-ANOVA, F_{11} = 3.483, p = 0.0495; effect size for

the main effect of treatment, η^2_p = 0.109; fig. 7f). In contrast, when stimulated during

retrieval, mice stimulated with 8 Hz displayed significantly reduced memory performance

(0.675 ± 0.049) compared to baseline (0.825 ± 0.030, RM-ANOVA, F_{11} = 3.265, p = 0.050;

fig. 7g). In contrast, YFP control mice were not affected by stimulations during encoding

(RM-ANOVA, F_2 = 0.1314, p = 0.8781), the delay period (RM-ANOVA, F_2 = 0.2020, p =

0.8197), or retrieval (RM-ANOVA, F_2 = 0.0454, p = 0.9557; effect size for the main effect of

treatment, η^2_p = 0.196) of working memory.

Surprisingly, 8 Hz optogenetic stimulations that are associated with pacing of theta

oscillations led to decreased performance when applied during either encoding or retrieval of

episodic memory in the NPOR task, as well as spatial working memory retrieval in the

DNMTS task. To further understand the effects of pacing theta on hippocampal physiology,

we analyzed phase locking of theta oscillation before and during 8 Hz optogenetic

stimulations (supplemental fig. 7a) and found that such stimulation led to a high phase

synchrony across stimulation epochs (supplemental fig. 7b). Furthermore, we analyzed the

cross-correlation of theta oscillation across dorsal CA1 (~1 mm distance between electrodes

along the septotemporal axis (supplemental fig. 7c) and found that pacing theta rhythms at 8

Hz led to increased cross correlation between those two electrodes (supplemental fig. 7d).

Discussion

Within the septohippocampal system, the exact causal relationships between (1) MS activity,

(2) hippocampal oscillations, (3) hippocampal neuron activity, and (4) behavior including

memory remain an active area of research. In particular, whether and how the MS supports

encoding of place and time in the hippocampus, as well as its specific contribution to memory

function remain unclear. Here, we leveraged more recent techniques that allowed us to record

>1000 neurons while performing optogenetic stimulation of MS neurons with a sub-second

resolution to control theta rhythms and assess its role in hippocampal physiology and

memory. Importantly, using an excitatory opsin and either scrambled or 8 Hz stimulations, we

were able to consistently and robustly abolish or pace hippocampal theta respectively.

Compared to MS inhibition using either an inhibitory opsin or pharmacological compounds

(e.g. muscimol), our approach enables a within-subject comparison of two opposed states

(paced vs abolished theta) while maintaining activity levels in MS-PV cells. We combined a

novel tone-cued linear track task that, together with information theoretic approaches, allowed

us to disentangle spatial and temporal hippocampal codes. This method alleviates the need for

arbitrary thresholds, enabling a standardized approach to calcium imaging analysis as

described previously[52]. While a large portion of CA1 pyramidal neurons expressed a mixture

of spatiotemporal codes, we focused our analyses on neurons tuned specifically to place, time,

or distance traveled.

While spatial coding has been extensively investigated in CA1, temporal and distance

codes have recently gained more interest. Temporal[27,34–36] and distance[36] codes have been

extracted by clamping visuospatial cues or extracted analytically using generalized linear

models in virtual reality paradigms[25,26,37]. Here we propose a novel approach to disentangle

spatial, temporal, and distance codes using an information theoretic approach which, together

with our novel cued-alternation task, enables the analysis of real world spatiotemporal and

multiplexed codes in freely moving animals. Strikingly, a large amount of CA1 pyramidal

neurons encode multiplexed information of location, distance, and time as previously

reported[25,26,37] in addition to self-motion signals such as acceleration, speed, and orientation[60].

Frequency scrambling and 8 Hz optogenetic stimulations drastically abolished or

paced theta rhythms, respectively, and led to an overall decreased activity in a subpopulation

of CA1 pyramidal cells while not causing significant changes in place cell activity, similar to

previous reports using either pharmacological inhibition[35,45,46], or optogenetic[9,48] pacing of MS

or septal inputs to the hippocampus. As MS neurons are known to be the primary driver of

hippocampal oscillations, we expected that MS stimulation would be associated with

disrupted time or distance cell activity in conditions of reduced theta rhythms, but did not

observe any change for both abolished and paced theta conditions. Additionally, pacing of

hippocampal oscillations to 8 Hz did not lead to changes in the quality of spatial codes as

reported previously[9], and did not alter temporal codes as observed in our behavioral paradigm.

While it was reported that time (but not place) cells could depend directly on medial

entorhinal cortex (MEC) inputs[61], other studies show that MEC lesions do not lead to any

alterations in hippocampal time cell physiology[62]. Notably, a more recent investigation found

that optogenetic pacing of the MS did not disrupt the spatial codes of grid cells in the

entorhinal cortex[63], further suggesting that MS activity is not directly involved in spatial codes

within the hippocampal formation. Together with our results, this indicates that temporal

codes could either be the result of MS-dependent computations within the entorhinal cortex[63]

(1), or be generated intrinsically within the hippocampus itself (2).

While early studies found that the MS played an essential role in maintaining time

cells and supporting working memory[35], recent experimental evidence shows that the exact

contribution of time cells to working memory performance could be less than previously

thought[62]. Since MS-PV neurons likely maintain significant activity levels during frequency

scrambling optogenetic stimulation (in contrast to inhibition approaches), our results strongly

support the role of precisely timed septal activity in supporting working memory, as

hypothesized previously[18]. Although we observed reduced memory performance when using

optogenetic stimulation during retrieval, stimulation during the encoding phase of the delayed

non-match to sample tasks was not associated with reduced memory in the DNMTS task.

Pacing theta oscillations at 8 Hz during either the encoding or retrieval phase of the NOPR

task led to impaired memory retrieval. Our electrophysiological analyses further reveal that

such stimulation caused distant regions of dorsal CA1 to be entrained at the same phase, with

non-physiological phase locking. While we did not directly investigate the causal relationship

between phase changes and memory performance, phase locking of pyramidal neurons to

theta oscillation has previously been shown to be a predictor of memory performance[64].

While we describe spatiotemporal codes in freely moving animals, one notable

limitation of our approach to establish temporal and distance tuning properties is that it

requires four times more sampling than a regular linear track which increases the chances of

photobleaching when adding stimulation conditions to the experimental timeline. While we

found that spatial codes remained unaffected by optogenetic stimulations within the same

session, more recent imaging hardware that includes sensors with increased sensitivity could

help prevent photobleaching and allow for longer recording sessions, and thus sampling of

temporal and distance cells along with stimulations. We also found that the pool of neurons

that significantly and specifically encode only one variable is small in comparison to the

number of conjunctive neurons, making sampling requirements for these highly specific

neurons much higher.

We provide experimental evidence that manipulation of MS does not alter locomotor

speed and that conversely, locomotor speed does dictate theta frequency. Even though place

and time codes were preserved during our MS stimulations, a portion (~6%) of cells were

directly modulated, and could account for memory impairments observed in our behavioral

assays. PV interneurons in the hippocampus have previously been reported to be part of a

microcircuit essential in regulating memory consolidation[65,66], and our optogenetic

manipulations could be associated with silencing of these cells to some extent. Moreover,

although we did not observe changes in frequency bands other than theta, we cannot exclude

that CA1 pyramidal neuron spike timing could be drastically altered when scrambling theta

oscillations, while 8 Hz stimulation were shown to not result in increased hippocampal

activity[17], which could explain the differential effects of those stimulation patterns on working

memory performance. The memory impairments we observed were likely non-cholinergic:

firstly, in our immunohistological experiments we found virtually no expression of

ChrimsonR in ChAT neurons of the MS. Secondly, our optogenetic stimulations were not

associated with changes in hippocampal ripples, while previous reports indicate that ChAT

neurons stimulation is associated with reduced ripple frequency[44,57]. ChrimsonR transfection

of MS VGLUT2 neurons is also unlikely, since activation of these neurons is associated with

direct increase in locomotor activity[67], which we did not observe. Although the precise

temporal arrangement of CA1 pyramidal spikes could explain, at least partially, the effects of

MS stimulation on memory, alternative mechanisms should be taken into consideration.

Notably, in addition to the hippocampus and the entorhinal cortex, MS PV neurons project to

the retrosplenial cortex[3], and could be responsible for some of the memory impairments

observed here.

In summary, using calcium imaging, optogenetics, and electrophysiology, we found

that theta rhythms can be paced or abolished using stimulation of the MS. Such stimulation

impaired episodic as well as working memory retrieval. These effects were non-cholinergic,

and did not disrupt hippocampal ripple activity. Finally, while a small fraction of hippocampal

neurons responded directly to optogenetic stimulation of the MS, place, time, and distance

cells were not disrupted by manipulations of theta oscillations. Together these results suggest

that while MS input to the hippocampus plays an essential role in memory, multiplexed codes

in CA1 pyramidal neurons might not be the direct substrate for such functions.

Methods

Animals

All procedures were approved by the McGill University Animal Care Committee and the

Canadian Council on Animal Care (protocol 2015-7650). A total of n = 41 male and female

(8-16 weeks old) parvalbumin-cre mice (Jackson Laboratory) were used in this study. Mice

were housed individually on a 12-hour light/dark cycle at 22°C and 40% humidity with food

and water ad libitum. All experiments were carried out during the light portion of the

light/dark cycle.

Adeno-associated viral vectors

Adeno-associated virus AAV5.CamKII.GCaMP6f.WPRE.SV40 (Addgene # 100834,

obtained from University of Pennsylvania Vector Core) was used in all calcium imaging

experiments. Adeno-associated virus (AAV) of serotype *dj* (hybrid capsid created from 8

different AAV serotypes) AAVdj-hSyn-ChrimsonR-tdTomato were obtained from the Vector

Core Facility at Oregon Health and Science University in Portland, Oregon. Although hSyn is

a housekeeping gene, we did not observe transfection in cholinergic cells (see Results and fig.

2b-e). An eYFP construct without the ChrimsonR sequence was used as a control (termed

YFP control in this manuscript).

Surgical procedures

Mice were anesthetized with isoflurane (5% induction, 0.5-1 % maintenance) and placed in a

stereotaxic frame (David Kopf Instruments). Body temperature was maintained with a heating

pad, eyes were hydrated with gel (Optixcare), and Carprofen (10 ml/kg) was administered

subcutaneously. The skull was completely cleared of all connective tissue and small

craniotomies were performed using a dental drill for subsequent injection or implant.

Viral injections

All viral injections were performed using a glass pipette connected to a Nanoject III

(Drummond) injector at a flow rate of 1 nL/second. 500 nl of AAVdj-hSyn-ChrimsonR-

tdTomato (or eYFP control) was delivered into the medial septum at a rate of 1 nL/s, at the

following coordinates based on reference mouse stereotaxic atlas[68] (distance from Bregma in

mm): anteroposterior (AP) 0.85, mediolateral (ML) 0, dorsoventral (DV) −4.50 using a 5º

angle in the ML plane. After surgery, animals were continuously monitored until recovery.

Fiber optic implant

Two weeks post-injection, mice were anesthetized for implantation following the same

surgical procedure. 200 μm diameter fiber optics with ceramic ferrule (Thor labs) were

implanted at the same coordinates. Implants were cemented in place using C&B-Metabond®

(Patterson dental, MN). Black nail polish was applied over the dental cement to block light

emission during optogenetic stimulation.

Electrode implants

An array of 7 tungsten microelectrodes (~1 MΩ impedance) was lowered in dorsal CA1

spanning through *stratum pyramidale* (pyr), *stratum radiatum* (rad) and *stratum lacunosum*

moleculare (lm). Screws placed in the bone above the frontal cortex and cerebellum served as

ground and reference, respectively. Following electrode, ground and reference placement,

dental cement was applied to secure the implant permanently to the skull.

Implant for calcium imaging

We injected the AAV5.CamKII.GCaMP6f virus (200 nL at 1 nl.s−1) in hippocampal CA1

using the following coordinates: anteroposterior (AP) −1.86 mm from bregma, mediolateral

(ML) 1.5 mm, dorsoventral (DV) 1.5 mm. 2 weeks following the injection, the mouse was

anesthetized with isoflurane and the skull was cleared. A ~2 mm diameter craniotomy was

performed in the skull above the injection site. An anchor screw was placed on the posterior

plate above the cerebellum. The dura was removed, and the portion of the cortex above the

injection site was aspirated using a vacuum pump, until the corpus callosum was visible.

These fiber bundles were then gently aspirated without applying pressure on the underlying

hippocampus, and a 1.8 mm diameter gradient index (GRIN; Edmund Optics) lens was

lowered at the following coordinates: AP −1.86 mm from bregma, ML 1.5 mm, DV 1.2 mm.

The GRIN lens was permanently attached to the skull using C&B-Metabond (Patterson

Dental), and Kwik-Sil (World Precision Instruments) silicone adhesive was placed on the

GRIN to protect it. 4 weeks later, the silicone cap was removed and CA1 was imaged using a

miniscope mounted with an aluminum base plate while the mouse was under light anesthesia

(~0.5 % isoflurane) to allow the visualization of cell activity. When a satisfying field of view

was found (large neuronal assembly, visible landmarks), the base plate was cemented above

the GRIN lens, the miniscope was removed, and a plastic cap was placed on the base plate to

protect the GRIN lens.

Simultaneous calcium imaging and electrophysiological recordings

To control for the effects of GRIN implants on theta physiology as well as cross-talk between

miniscope and optogenetic excitation lights, we attached tungsten micro-electrodes to GRIN

lenses. To this end, we placed GRIN lenses horizontally under a low-magnification

microscope in a dust-free environment. A tungsten microelectrode was gently placed on the

top-edge of the GRIN lens using a micromanipulator. We used the known diameter of the

GRIN lens as a reference unit to estimate the desired protrusion of the electrode (~50μm,

further assessed digitally after taking a microphotograph of the preparation) in accordance

with our planned implantation coordinates. Small amounts (~50μL) of superglue were

deposited on the top edge of the GRIN lens with the electrode in place, left to dry for ~10min,

before applying the next layer of glue. 5 thin layers were used to maintain the electrode

attached to the GRIN lens. After implantation of these GRIN-electrode assemblies using the

protocol described above, the protruding wires were gently bent and hidden under a protective

cap (1-1.5 cm high) retrieved from the pear of a manual suction pipette as a replacement for

Kwik-Sil.

***In vivo* behavioral procedures**

Habituation

Mice were habituated by being gently handled for ~5 min over the course of one week, with

progressive habituation to the plugging procedure (fiber optic, baseplate, and

electrophysiological pre-amplified tethers. Animals were then water-scheduled (2h access per

day). Miniscope recordings were performed at 30 Hz for 15 minutes, with only one recording

session per day to minimize photobleaching.

Miniscope recordings

Miniscopes (V3) were assembled using open-source instructions as described previously

(miniscope.org). Imaging data were acquired using a CMOS imaging sensor (Aptina,

MT9V032) and multiplexed through a lightweight coaxial cable. Data was acquired using a

data acquisition (DAQ) box connected via a USB host controller (Cypress, CYUSB3013).

Animal behavior was recorded using a consumer grade webcam (Logitech) mounted above

the environment. Calcium and behavioral data were recorded using miniscope.org open

source DAQ custom acquisition software. The DAQ simultaneously acquired behavioral and

cellular imaging streams at 30 Hz as uncompressed .avi files of 1000 frames, along with

timestamps to enable precise temporal alignment of behavioral and calcium imaging data.

In vivo electrophysiological recordings

Following 1 week of post-surgical recovery and one week of habituation to the tethering

setup, LFP from implanted mice was recorded. All recorded signals from implanted

electrodes were amplified by the tether pre-amplifier before being digitized at 22 KHz using a

digital recording system (Neuralynx, USA). Recordings of each channel signal were saved in

a dedicated file along with video recordings and TTL signals from the Laser Diode for

subsequent analysis.

Optogenetic stimulation

Laser stimulation was delivered through a fiber optic cord (200 μm diameter) using a laser

diode fiber light source (Doric Lenses). Light intensity was calibrated and wavelength-

corrected using the Power Meter Bundle with the PM100D Console and S130C Slim

Photodiode Sensorlight (Thorlabs). Every pacing stimulation (including 8 Hz) was performed

using 5 ms square pulses. To apply scrambled light stimulation, we used an Arduino

microcontroller to generate a white noise oscillation signal directly fed into the analogue input

of the laser diode driver. To perform randomly selected frequency stimulation, we used a

standard 5s ON, 5s OFF protocol but for each stimulation epoch, we used an Arduino

microcontroller to randomly select a stimulation frequency in the theta band. When applying

optogenetic stimulation during behavior, a loose piece of heat shrink tubing was fitted around

the junction between the patch cord and the mouse' ferrule implant to limit visible light

emission. Light intensities are expressed as nominal power, as measured at the tip of the fiber

optic implant - cord assembly, and corrected for the appropriate wavelength.

Behavioral assays

Sequential tone linear track

Mice were water scheduled and could access water only for 2h per day from 6-8 pm. To

disentangle spatial, temporal, and distance codes, we build a 134 cm long linear track using

medium gray Lego® bricks which allows easy modifications and implementations without the

need to permanently modify the structure of the maze. Pyroelectric sensors were placed in

each end of the linear track, and were connected to an Arduino microcontroller. Each

detection triggered a new tone in sequence, indicating the progress to reward delivery. The

following tones were used and delivered using a piezo speaker: 1Hz beeps at 2,000Hz, 4Hz

beeps at 3,000Hz, and a continuous tone at 4,000Hz. When the last, continuous tone was

triggered, a reward (10% sucrose in water) was delivered at the starting end of the linear track

in the cap of a 15 mL Falcon tube. Changes in running direction before triggering the next

tone were considered as errors and did not trigger reward delivery. Performance was

measured as the number of correct runs (with no error) divided by the number of total runs.

Novel object place recognition

On the first day, mice were allowed to freely explore a 45x45 cm dark gray open-field that

contained visual cues (large white horizontal and vertical gratings) on its walls for 10 min. On

the second day, two identical objects (helping hand base) were presented for 10 min. On the

third day, mice were allowed to explore the same open field while the location of one of the

two objects was displaced. To control for potential spatial preference biases, both the initial as

well as the displaced position of objects were randomized. Mice were attributed a random

testing order that was kept identical throughout the three days of testing. Behavior was

recorded with a video camera and analyzed offline. Behavioral analysis was performed blind

to the genotype and treatment. Object explorations were defined as epochs where mice have

their nose within 1 cm of an object. Recognition index (RI) was computed as the following:

642
$$RI = \frac{t_{novel}}{t_{novel} + t_{familiar}}$$
643

where t_{novel} and $t_{familiar}$ represent the time exploring novel and familiar objects, respectively.

This metric can also be computed in the sample phase for the prospectively novel (displaced)

object to assess potential pre-existing preferential biases.

Automated delayed non-match to sample task

Mice were water-scheduled and trained in a continuous T-maze to a delayed non-match to

sample task. Briefly, each trial was divided into two phases: sample and test. In the sample

phase, one arm was blocked and mice were forced to explore the opposite arm where they

received a 50 μL of 10% sucrose water reward. After completing the sample phase, mice were

subjected to a delay (10 s). Then, during the test phase, both arms could be explored, but only

the opposite (unexplored) arm is baited, so that mice have to alternate locations between

sample and test phases. Mice were subjected to 10 trials (sample + test) per day, for 10

consecutive days, and the daily success rate was calculated at the number of correct choices

divided by the total number of trials.

Post-mortem histological analyses

After completion of behavioral testing, mice were deeply anesthetized with a mixture

ketamine/xylazine/acepromazide (100, 16, 3 mg/kg, respectively, intraperitoneal injection)

and perfused transcardially with 4% paraformaldehyde (PFA) in PBS. Brains were extracted

and postfixed overnight in PFA at 4 ºC and subsequently washed in PBS for an additional 24

h at 4 ºC. Brains and sections were cryoprotected in a solution of 30 % ethylene glycol, 30 %

glycerol, and 40 % PBS until used. Each brain was then sectioned at 50 μm using a

vibratome: every section was sequentially collected in 4 different 1.5 mL tubes to allow

different analyses (electrode location, immunohistochemistry).

Immunohistology

Using one tube of collected brain sections (25% sampling) for each analysis, sections were

washed 3x5 min in PBS to remove cryoprotective solution. Sections were first incubated

overnight with PGT (0.45 % Gelatin and 0.25 % Triton in PBS) at 4 ºC. Next, slices were

incubated with primary antibodies: either 1:200 goat anti-choline-acetyl-transferase from

Millipore or 1:500 mouse anti-parvalbumin monoclonal IgG1 from Sigma-Aldrich in PGT at

room temperature for 48 h or 2 h respectively. Following washes of 10, 20, and 30 min,

sections were then incubated with secondary antibodies [1:2000 donkey anti-goat coupled

with Alexa 488 or 1:500 goat anti-mouse IgG1 coupled to Alexa 488 (Life Technologies)] in

PGT for 45 min. Following 10, 20, and 30 min washes, sections were then mounted on glass

slides and permanently coverslipped with Fluoromount mounting medium that contained

DAPI. Only mice with histologically confirmed placement of implants were included in this

study. For GRIN lenses, the surface of the lens had to be < 100μm above *stratum pyramidale*

and GCaMPf expression was confirmed using fluorescence microscopy. Electrophysiological

implants had to include at least one microelectrode in CA1 *stratum radiatum* or *stratum*

pyramidale. Finally, the tips of fiber optics had to be within 100 μm of the medial septum

region, and proper construct expression was assessed using fluorescence microscopy.

Data analysis

Except for analyses of sharp-wave ripples and explicit impact of locomotion on physiological

recordings, electrophysiological and calcium imaging analyses were performed only on

periods of locomotion (>2 cm.s^{-1} in the open field; >5 cm.s^{-1} on the linear track).

Automatized tracking of behavior

To extract information about the position, velocity, and head direction of mice, we used

DeepLabCut[69,70]. Briefly, we trained a model to detect mice ears, nose, body, and tailbase.

Head direction was estimated using the angle between either each ear or the nose and body

depending on measurement availability. Location data was interpolated to calcium imaging

sampling frequency using linear interpolation. Velocity was extracted by computing Δd/Δt

where *d* is the distance and *t* time, and subsequently smoothed the result by applying a

gaussian filter with sigma = 33 ms to remove movement artifacts. Velocity signals were used

to identify periods of locomotor activity, using a threshold of 5 cm.s^{-1}, and used selectively to

compute place, time, and distance modulation of activity.

Calcium imaging analysis

Calcium imaging data analysis was performed using MATLAB 2020a and Python 3.8.4.

Video   recordings   were   analyzed   using   the   Miniscope   Analysis   pipeline

(https://github.com/etterguillaume/MiniscopeAnalysis).   Briefly,   rigid   (rotation   and

translation) motion correction was applied using NoRMCorre[71], and videos were spatially

downsampled (3x) before concatenation. Calcium traces were extracted using CNMFe[51] using

the following parameters: gSig = 3 pixels (width of gaussian kernel), gSiz = 20 pixels

(approximate neuron diameter), background_model = 'ring', spatial_algorithm = 'hals',

min_corr = 0.8 (minimum pixel correlation threshold), min_PNR = 8 (minimum peak-to-

noise ratio threshold).

Raw calcium traces were filtered to remove high-frequency fluctuations and binarized:

briefly, neurons were considered active when normalized calcium signal amplitude > 2

standard-deviations, and first-order derivative > 0 (see Etter et al., 2020 for additional details

on the methodology[52]). To extract neurons tuning to specific variables, location, time, and

distance were binned (location: 3 cm bins; time: 1 s bins; distance: 3 cm bins). From binarized

signals, we computed the marginal likelihood of cells being active $P(A)$ that we use as a

proxy for neuronal activity. More importantly, we then derive activity likelihood, or the

'probability of being active given the state of a variable' $P(A|S_i)$ using binned variables:

721
$$P(A|S_i) = \frac{time\,active\,while \in bin\,i}{time \in state}$$

723   where A is the activity level (0 or 1) of a given neuron, when being in bin i of a given state S.

724   These values were smoothed using a Gaussian kernel with $\sigma = 5$, 10, or 15 for spatial,

725   temporal, and distance tuning curves respectively.

726   To identify place, time, and distance cells, we computed the mutual information (MI)

727   between neuronal activity and the bin i of a given state S using the following:

728
$$MI = \sum_{i=1}^{M} \sum_{j=1}^{2} P(S_i \cap A_j) \, x \log 2 \left(P \frac{(S_i \cap A_j)}{P(S_i) \, x P(A_j)} \right)$$

where M is the total number of possible behavioral states, $P(S_i \cap A_j)$ is the joint probability of

the animal being in bin i concurrently with activity level j (0 or 1). To assess the significance

of the obtained MI value, we then generated n = 1000 shuffled surrogates using random

circular permutations. We chose circular permutations because they remove the temporal

relationship between neuronal activity and behavior, while still preserving the temporal

structure of calcium transients and therefore leads to more conservative results (as opposed to

complete randomization of every data point, which inflates the significance value of results).

Because shuffled surrogates were not systematically normally distributed, we used a non-

parametric approach where the p-value (p_N) corresponds to the number of datapoints from the

shuffled distribution that are greater than the actual data for each bin, divided by the number

of permutations[52,72].

To determine the modulation of cells by optogenetic stimulations, we used a similar

approach but computed the MI between neuronal activity and the binarized stimulation

signals (thus treated as a behavioral state). The same circular shuffling procedure was then

used to extract statistical significance.

Naive Bayes decoder

To further evaluate the significance of neuronal activity at the population level, we

trained a naive Bayesian decoder with a uniform prior to predict spatiotemporal variables

from binarized neuronal activity. The posterior probability density function, which is the

probability that an animal is in a particular bin S_i, given the neural activity A is calculated

using the following equation:

751
$$P(S_i|A) = \frac{P(A|S_i) \times P(S_i)}{P(A)}$$

where P(S|A) is the posterior probability distribution of states given neuronal activity. Using

only epochs with velocity > 5 cm/s, a training dataset was generated using 90% of the data.

The remaining 10% of the data was used for testing. Decoding error was calculated using 50

bootstrapped surrogates and a pool of 160 cells using randomly chosen data points with

replacement. Every neuron is assumed to be independent of each other, which in practice is

not the case and leads to greater reconstruction errors, but decreases computational time. The

population posterior probability was derived from the following equation:

759
$$P(S|A) = \prod_{k=1}^{N} \frac{P(A_k|S) \times P(S)}{P(A_k)}$$

761 with N corresponding to the number of neurons used. Finally, the decoded data point

762 corresponds to the maximum *a posteriori* (MAP):

763
$$\hat{y} = arg\,max\,exp\left[\sum_{k=1}^{N} \log\left(1 + \frac{P(A_k|S) \times P(S)}{P(A_k)}\right) - 1\right]$$

765 The exponential-logarithm trick is used to prevent underflow when multiplying a large

766 number of small probabilities, which is a correct approximation since the posterior is a

767 smooth convex function.

768   Temporal filtering was applied to posterior probabilities using:

769
$$P(S|A_k(t)) = \prod_{k=1}^{N} \prod_{t=1}^{T} \frac{P(A_k(t)|S) \times P(S)}{P(A_k(t))}$$

where t is a given timestep, and $P(S|A_k(t))$ the *a posteriori* distribution of states at timestep t

for neuron k. The number of past timesteps used to estimate the mouse state at time t is

determined by $T,$ which we set as 2 s unless noted otherwise. On the linear track, decoding

error was assessed as:

$$decoding\, error = |decoded\, state - actual\, state|$$

In order to compare decoding scores across animals, we z-scored decoding errors

using the following equation:

$$z = \frac{\bar{x} - \bar{s}}{\sigma(x)}$$

where $\bar{x}$ is the average actual decoding error, $\bar{s}$ is the average decoding error using shuffled

surrogates, and $\sigma(x)$ is the standard deviation of the actual decoding error.

Tracking cells over multiple days

Neurons were tracked over multiple days using a probabilistic method as previously

described[73]. Briefly, spatial footprints were aligned using rigid alignment to correct for

rotations and translations. After alignment, we considered candidate sets of cells to be the

same neuron if their maximal distance was < 12 um, and used the modeled spatial correlation

threshold (usually in the range .6 - .8) to determine the identity of cell pairs across days.

Finally, we assessed the stability of the spatial representation using pairwise field correlation

(Pearson correlation of tuning curves).

Electrophysiological analysis

Electrophysiological data analysis was performed using MATLAB 2020a and both the signal

processing as well as the wavelet toolbox. Wavelet convolution was applied to LFP signals

using complex Morlet wavelets ('cmor1-1.5' in MATLAB) when both time- and frequency-

domain accuracy was required. Moving window Fourier convolution (2 s window in the theta

band, 5 s window in the gamma band, 10 ms moving steps) was used when frequency-domain

accuracy was privileged over time-domain accuracy (e.g. to plot dominant frequency when

pacing using optogenetics). Analysis of power spectral densities was performed when mice

were running at 5 cm.s^{-1} or above, unless described otherwise.

Oscillation strength

Oscillation strength was computed as the ratio of cumulative power spectral density around

the peak oscillation frequency ± 1 Hz to the cumulative band power in the theta (4 - 12 Hz)

band. This metric becomes 1 when all power spectral density falls within the peak oscillation

frequency (e.g. 8 Hz if stimulating at that frequency).

Sharp-wave ripples detection and analysis

To monitor sharp-wave ripples (SWRs), mice were allowed to freely explore an open-field for

10 min while recording. Stimulations (scrambled or 8 Hz) were performed with a 5s ON, 5s

paradigm. Only periods of quiet restfulness were considered for analysis. To this end, we

computed the z-scored ratio of theta/delta power after filtering for each frequency band and

performing a Hilbert transform, as described previously[74], and only considered periods where

the resulting value was under 0. To detect SWRs, we filtered LFP signals in the 150-250 Hz

frequency band and subsequently z-scored. Ripples were detected using the findpeaks

function in MATLAB, with the following parameters: threshold=4 sd, minpeakwidth=0 s,

minpeakdistance=0.03 s.

820   Statistics

Unless stated otherwise, all data are presented as mean ± standard error of the mean (SEM)

and statistical test details are described in the corresponding results. The distribution

normality of each group was assessed using Shapiro-Wilk normality test and parametric tests

were used only when distributions were found normal (non-parametric tests are described

where applicable). 1ANOVA: one-way ANOVA; 2ANOVA: two-way ANOVA; RM-

ANOVA: repeated measure ANOVA. $p < 0.05$ was considered statistically significant. *, $p <$

0.05; **, $p < 0.01$; ***, $p < 0.001$, ****, $p < 0.0001$. Effect size of 1ANOVA was assessed

with η^2 using sum of squares (SS):

829
$$\eta^2 = \frac{SS_{between}}{SS_{total}}$$

831   Effect size of 2ANOVA and RM-ANOVA was assessed with partial η^2 using:

832
$$\eta^2{}_p = \frac{SS_{effect}}{SS_{effect} + SS_{error}}$$

834   Author information

835   Affiliations

836   McGill University & Douglas Mental Health University Institute, Montreal, Canada

837   Corresponding authors: sylvain.williams@mcgill.ca, etterguillaume@gmail.com

838   6875 Boulevard LaSalle, Verdun, QC H4H 1R3, Phone : 514 761-6131, Fax: 514 888-4067

Contributions

GE and SW designed the study. GE performed and analyzed *in vivo* experiments. SVDV performed and analyzed both behavioral as well as immunohistological experiments. JC performed surgeries for the mice used for controlling cross-talk between calcium imaging and optogenetics. GE and SW wrote the manuscript with inputs from all collaborators.

Competing interests

The authors declare no competing interests.

Correspondence author

Correspondence should be addressed to Sylvain Williams: sylvain.williams@mcgill.ca

Acknowledgements: We thank Ke Cui for helping to maintain the colony and perfusing mice; Mark P Brandon, Alexandra Keinath, and Eric Carmichael for comments on the manuscript.

Funding

This work was supported by funding from the Canadian Institutes for Health Research (CIHR) Foundation Program FDN-148478, the Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant RGPIN-2020-06717, and a Tier 1 Canada Research Chair to SW. SVDV was supported by a Vanier Canada Graduate Scholarship and the Richard H. Tomlinson Doctoral Fellowship. JC was supported by a Healthy Brains, Healthy Lives (HBHL) Graduate Student Fellowship and a McGill Integrated Program in

Data availability

The data that support the findings of this study are available from the corresponding

author upon reasonable request.

Code availability

All source codes used in the current study are available along with instructions here:

https://github.com/etterguillaume/etter_et_al_2022

References

1. Colgin, L. L. Rhythms of the hippocampal network. *Nat. Rev. Neurosci.* **17**, 239–249 (2016).

2. Tóth, K., Freund, T. F. & Miles, R. Disinhibition of rat hippocampal pyramidal cells by

GABAergic afferents from the septum. *J. Physiol.* **500**, 463–474 (1997).

3. Unal, G., Joshi, A., Viney, T. J., Kis, V. & Somogyi, P. Synaptic targets of medial septal

projections in the hippocampus and extrahippocampal cortices of the mouse. *J. Neurosci.* **35**,

15812–15826 (2015).

4. Simon, A. P., Poindessous-Jazat, F., Dutar, P., Epelbaum, J. & Bassant, M.-H. Firing properties of

anatomically identified neurons in the medial septum of anesthetized and unanesthetized restrained

rats. *J. Neurosci.* **26**, 9038–9046 (2006).

5. Sotty, F. *et al.* Distinct electrophysiological properties of glutamatergic, cholinergic and

GABAergic rat septohippocampal neurons: novel implications for hippocampal rhythmicity. *J.*

Physiol. **551**, 927–943 (2003).

6. Manseau, F., Danik, M. & Williams, S. A functional glutamatergic neurone network in the medial

septum and diagonal band area. *J. Physiol.* **566**, 865–884 (2005).

7. Amilhon, B. *et al.* Parvalbumin interneurons of hippocampus tune population activity at theta

frequency. *Neuron* **86**, 1277–1289 (2015).

8. Etter, G. *et al.* Optogenetic gamma stimulation rescues memory impairments in an Alzheimer's

disease mouse model. *Nat. Commun.* **10**, 1–11 (2019).

9. Zutshi, I. *et al.* Hippocampal neural circuits respond to optogenetic pacing of theta frequencies by

generating accelerated oscillation frequencies. *Curr. Biol.* **28**, 1179-1188.e3 (2018).

10. Bender, F. *et al.* Theta oscillations regulate the speed of locomotion via a hippocampus to

lateral septum pathway. *Nat. Commun.* **6**, 8521 (2015).

11. Green, J. D. & Arduini, A. A. Hippocampal electrical activity in arousal. *J. Neurophysiol.* **17**,

533–557 (1954).

12. Mitchell, S. J., Rawlins, J. N., Steward, O. & Olton, D. S. Medial septal area lesions disrupt

theta rhythm and cholinergic staining in medial entorhinal cortex and produce impaired radial arm

maze behavior in rats. *J. Neurosci.* **2**, 292–302 (1982).

13. Gemzik, Z. M., Donahue, M. M. & Griffin, A. L. Optogenetic suppression of the medial septum impairs working memory maintenance. *Learn. Mem.* **28**, 361–370 (2021).

14. Knox, D. & Keller, S. M. Cholinergic neuronal lesions in the medial septum and vertical limb of the diagonal bands of Broca induce contextual fear memory generalization and impair acquisition of fear extinction. *Hippocampus* **26**, 718–726 (2016).

15. Jeong, D. U. *et al.* Improvements in memory after medial septum stimulation are associated with changes in hippocampal cholinergic activity and neurogenesis. *BioMed Res. Int.* **2014**, e568587 (2014).

16. Sans-Dublanc, A. *et al.* Septal GABAergic inputs to CA1 govern contextual memory retrieval. *Sci. Adv.* **6**, eaba5003 (2020).

17. Quirk, C. R. *et al.* Precisely timed theta oscillations are selectively required during the encoding phase of memory. *Nat. Neurosci.* **24**, 1614–1627 (2021).

18. Mehta, M. R. Role of Rhythms in Facilitating Short-Term Memory. *Neuron* **45**, 7–9 (2005).

19. Scoville, W. B. & Milner, B. Loss of recent memory after bilateral hippocampal lesions. *J Neurol Neurosurg Psychiat* **20**, 11–21 (1957).

20. Penfield, W. & Milner, B. Memory deficit produced by bilateral lesions in the hippocampal zone. *AMA Arch. Neurol. Psychiatry* **79**, 475–497 (1958).

21. Jeneson, A. & Squire, L. R. Working memory, long-term memory, and medial temporal lobe function. *Learn. Mem.* **19**, 15–25 (2012).

22. O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res.* **34**, 171–175 (1971).

23. Robinson, N. T. M. *et al.* Targeted Activation of Hippocampal Place Cells Drives Memory-Guided Spatial Behavior. *Cell* **183**, 1586-1599.e10 (2020).

24. Muller, R. U. & Kubie, J. L. The effects of changes in the environment on the spatial firing of hippocampal complex-spike cells. *J. Neurosci.* **7**, 1951–1968 (1987).

25. Ravassard, P. *et al.* Multisensory Control of Hippocampal Spatiotemporal Selectivity. *Science* **340**, 1342–1346 (2013).

26.   Aghajan, Z. M. *et al.* Impaired spatial selectivity and intact phase precession in two-

dimensional virtual reality. *Nat. Neurosci.* **18**, 121–128 (2015).

27.   Villette, V., Malvache, A., Tressard, T., Dupuy, N. & Cossart, R. Internally recurring

hippocampal sequences as a population template of spatiotemporal information. *Neuron* **88**, 357–

366 (2015).

28.   Samsonovich, a & McNaughton, B. L. Path integration and cognitive mapping in a

continuous attractor neural network model. *J. Neurosci. Off. J. Soc. Neurosci.* **17**, 5900–20 (1997).

29.   Mehta, M. R. From synaptic plasticity to spatial maps and sequence learning: Place Field

Plasticity. *Hippocampus* **25**, 756–762 (2015).

30.   McNaughton, B. L., Battaglia, F. P., Jensen, O., Moser, E. I. & Moser, M.-B. Path integration

and the neural basis of the 'cognitive map'. *Nat. Rev. Neurosci.* **7**, 663–78 (2006).

31.   Eichenbaum, H. Time cells in the hippocampus: a new dimension for mapping memories. *Nat.*

Rev. Neurosci. **15**, 732–744 (2014).

32.   Eichenbaum, H. On the Integration of Space, Time, and Memory. *Neuron* **95**, 1007–1018

(2017).

33.   Eichenbaum, H. The role of the hippocampus in navigation is memory. *J. Neurophysiol.* **117**,

1785–1796 (2017).

34.   Pastalkova, E. Internally Generated Cell Assembly Sequences in the Rat Hippocampus Eva

Pastalkova. **1322**, (2012).

35.   Wang, Y., Romani, S., Lustig, B., Leonardo, A. & Pastalkova, E. Theta sequences are

essential for internally generated hippocampal firing fields. *Nat. Neurosci.* **18**, 282–288 (2015).

36.   Kraus, B. J., Robinson, R. J., White, J. A., Eichenbaum, H. & Hasselmo, M. E. Hippocampal

'time cells': time versus path integration. *Neuron* **78**, 1090–1101 (2013).

37.   Moore, J. J., Cushman, J. D., Acharya, L., Popeney, B. & Mehta, M. R. Linking hippocampal

multiplexed tuning, Hebbian plasticity and navigation. *Nature* **599**, 442–448 (2021).

38.   O'Keefe, J. & Recce, M. L. Phase relationship between hippocampal place units and the EEG

theta rhythm. *Hippocampus* **3**, 317–330 (1993).

39.   Skaggs, W. E., McNaughton, B. L., Wilson, M. A. & Barnes, C. A. Theta phase precession in

hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* **6**, 149–172 (1996).

40.   Salz, X. D. M. *et al.* Time cells in hippocampal area ca3. **36**, 7476–7484 (2016).

41.   Givens, B. S. & Olton, D. S. Cholinergic and GABAergic modulation of medial septal area: Effect on working memory. *Behav. Neurosci.* **104**, 849–855 (1990).

42.   Ikonen, S., McMahan, R., Gallagher, M., Eichenbaum, H. & Tanila, H. Cholinergic system regulation of spatial representation by the hippocampus. *Hippocampus* **12**, 386–397 (2002).

43.   Givens, B. S., Olton, D. S. & Crawley, J. N. Galanin in the medial septal area impairs working memory. *Brain Res.* **582**, 71–77 (1992).

44.   Zhang, Y. *et al.* Cholinergic suppression of hippocampal sharp-wave ripples impairs working memory. *Proc. Natl. Acad. Sci.* **118**, (2021).

45.   Koenig, J., Linder, A. N., Leutgeb, J. K. & Leutgeb, S. The spatial periodicity of grid cells is not sustained during reduced theta oscillations. *Science* **332**, 592–595 (2011).

46.   Brandon, M. P., Koenig, J., Leutgeb, J. K. & Leutgeb, S. New and distinct hippocampal place codes are generated in a new environment during septal inactivation. *Neuron* **82**, 789–796 (2014).

47.   Boyce, R., Glasgow, S. D., Williams, S. & Adamantidis, A. Sleep research: Causal evidence for the role of REM sleep theta rhythm in contextual memory consolidation. *Science* **352**, 812–816 (2016).

48.   Mouchati, P. R., Kloc, M. L., Holmes, G. L., White, S. L. & Barry, J. M. Optogenetic "low-theta" pacing of the septohippocampal circuit is sufficient for spatial goal finding and is influenced by behavioral state and cognitive demand. *Hippocampus* (2020) doi:10.1002/hipo.23248.

49.   Aharoni, D. & Hoogland, T. M. Circuit investigations with open-source miniaturized microscopes: past, present and future. *Front. Cell. Neurosci.* **13**, (2019).

50.   Ghosh, K. K. *et al.* Miniaturized integration of a fluorescence microscope. *Nat. Methods* **8**, 871–878 (2011).

51.   Zhou, P. *et al.* Efficient and accurate extraction of in vivo calcium signals from microendoscopic video data. *eLife* **7**, e28728 (2018).

52.   Etter, G., Manseau, F. & Williams, S. A probabilistic framework for decoding behavior from

in vivo calcium imaging data. *Front. Neural Circuits* **14**, (2020).

53.   Stamatakis, A. M. *et al.* Simultaneous optogenetics and cellular resolution calcium imaging

during active behavior using a miniaturized microscope. *Front. Neurosci.* **12**, 1–16 (2018).

54.   Buzsáki, G. Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and

planning. *Hippocampus* **25**, 1073–1188 (2015).

55.   Roux, L., Hu, B., Eichler, R., Stark, E. & Buzsáki, G. Sharp wave ripples during learning

stabilize the hippocampal spatial map. *Nat. Neurosci.* **20**, 845–853 (2017).

56.   Fernández-Ruiz, A. *et al.* Long-duration hippocampal sharp wave ripples improve memory.

Science **364**, 1082–1086 (2019).

57.   Vandecasteele, M. *et al.* Optogenetic activation of septal cholinergic neurons suppresses sharp

wave ripples and enhances theta oscillations in the hippocampus. *Proc. Natl. Acad. Sci. U. S. A.*

111, (2014).

58.   Ziv, Y. *et al.* Long-term dynamics of CA1 hippocampal place codes. *Nat. Publ. Group* **16**,

264–266 (2013).

59.   Sławińska, U. & Kasicki, S. The frequency of rat's hippocampal theta rhythm is related to the

speed of locomotion. *Brain Res.* **796**, 327–331 (1998).

60.   Veldt, S. van der, Etter, G., Mosser, C.-A., Manseau, F. & Williams, S. Conjunctive spatial

and self-motion codes are topographically organized in the GABAergic cells of the lateral septum.

PLOS Biol. **19**, e3001383 (2021).

61.   Robinson, N. T. M. *et al.* Medial entorhinal cortex selectively supports temporal coding by

hippocampal neurons. *Neuron* **94**, 677-688.e6 (2017).

62.   Sabariego, M. *et al.* Time cells in the hippocampus are neither dependent on medial entorhinal

cortex inputs nor necessary for spatial working memory. *Neuron* **102**, 1235-1248.e5 (2019).

63.   Lepperød, M. E. *et al.* Optogenetic pacing of medial septum parvalbumin-positive cells

disrupts temporal but not spatial firing in grid cells. *Sci. Adv.* **7**, eabd5684.

64.   Rutishauser, U., Ross, I. B., Mamelak, A. N. & Schuman, E. M. Human memory strength is

predicted by theta-frequency phase-locking of single neurons. *Nature* **464**, 903–907 (2010).

65.   Xia, F. *et al.* Parvalbumin-positive interneurons mediate neocortical-hippocampal interactions

that are necessary for memory consolidation. *eLife* **6**, 1–25 (2017).

66.   Donato, F., Rompani, S. B. & Caroni, P. Parvalbumin-expressing basket-cell network

plasticity induced by experience regulates adult learning. *Nature* **504**, 272–276 (2013).

67.   Fuhrmann, F. *et al.* Locomotion, theta oscillations, and the speed-correlated firing of

hippocampal neurons are controlled by a medial septal glutamatergic circuit. *Neuron* **86**, 1253–

1264 (2015).

68.   Paxinos, G. & Watson, C. The rat brain in stereotaxic coordinates. *Acad. Press Inc* 439 (1997)

doi:10.1016/0166-2236(87)90017-8.

69.   Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with

deep learning. *Nat. Neurosci.* **21**, 1281–1289 (2018).

70.   Nath, T. *et al.* Using DeepLabCut for 3D markerless pose estimation across species and

behaviors. *Nat. Protoc.* **14**, 2152–2176 (2019).

71.   Pnevmatikakis, E. A. & Giovannucci, A. NoRMCorre: An online algorithm for piecewise

rigid motion correction of calcium imaging data. *J. Neurosci. Methods* **291**, 83–94 (2017).

72.   Cohen, M. X. *Analyzing neural time series data: theory and practice*. (The MIT Press, 2014).

73.   Sheintuch, L. *et al.* Tracking the same neurons across multiple days in ca2+ imaging data. *Cell*

Rep. **21**, 1102–1115 (2017).

74.   Wikenheiser, A. M. & Redish, A. D. The balance of forward and backward hippocampal

sequences shifts across behavioral states. *Hippocampus* **23**, 22–29 (2013).

1031

Figure 1. Spatial and temporal codes are present in partially overlapping populations of

CA1 neurons. a, mice were injected with GCaMP6f in hippocampal subfield CA1 and

subsequently implanted with a GRIN lens over the pyramidal cell layer for miniscope

imaging. **b**, post-mortem coronal section from one example mouse. blue, DAPI; green,

GCaMP6f. **c**, planar projection of extracted spatial footprints from one example mouse. **d**,

corresponding calcium transients from a portion of recorded cells. **e**, behavioral paradigm

used to disentangle time, place, and distance coding cells (see Methods). **f**, behavioral

outcomes for one example mouse. **g**, example pairs of neurons modulated by place (yellow),

time (green), or distance (orange) along with corresponding peak activity likelihood. **h**, 3D

scatter plot of the distribution of cells selectively encoding a single variable along with their

associated mutual information (MI) in each dimension. Cells in black either encode more than

one variable or did not encode any variable significantly more than shuffled surrogates. **i**,

individual traces from one example place-modulated cell (yellow) and one example time-

modulated cell (green). Vertical scale bars: 1ΔF/F. **j**, tuning curves for place, time, and

distance cells, sorted by the location of peak activity likelihood for each dimension (location,

time, and distance). **k**, portion of total cells (log scale) encoding for one, several, or no

variables. Bayesian decoding of location (**l-n**), time (**o-q**), and distance (**r-t**). For each

encoded variable, posterior probabilities are plotted with the corresponding actual state (**l,o,r**).

Confusion matrices between the actual vs decoded values (**m,p,s**). Decoding errors using all

cells vs shuffled surrogates (**n,q,t**, left panels), spatial cells vs non spatial cells (**n**), temporal

cells vs non temporal cells (**q**), distance vs non distance cells (**t**).

1053

Figure 2. Selective optogenetic pacing and ablation of theta oscillations. a, AAV-

ChrimsonR expressing vector was injected in the MS (blue, DAPI; red, ChrimsonR) and a

fiber optic was placed at the injection site (white arrow). Scale bar, 1mm. **b**,

immunohistological double staining for ChrimsonR and parvalbumin (PV+) neurons. **c**,

quantification of ChrimsonR+/PV+ co-expressing neurons. **d**, immunohistological double

staining for ChrimsonR and cholinergic (ChAT+) neurons. **e**, quantification of

ChrimsonR+/ChAT+ co-expressing neurons. **f**, recording configuration, with fiber optic in the

MS and electrode in CA1. **g**, wavelet spectrograms before, during, and after scrambled (left,

in red) and 8 Hz (right, in blue) stimulation patterns, with associated unfiltered LFP (white

overlay, and enlarged, black traces at the bottom). **h**, average Fourier power spectra for 60 x

5s stimulation epochs (top), and sorted power spectra for each stimulation epoch (bottom) for

baseline (left, in black), scrambled (center, in red), and 8 Hz (right, in blue) stimulations in a

representative mouse. **i**, oscillation strength for baseline vs scrambled or 8 Hz optogenetic

stimulations. Values for white noise (medium gray) and control 8 Hz sine waves are used for

comparison. **j**, group average (n = 3 mice) portion of baseline power for each stimulation

pattern and every frequency band (log scale). Frequency bands: δ, 1-4 Hz; θ, 4-12 Hz; α/β,

12-30 Hz; γslow, 30-60 Hz; γfast, 60-120 Hz; HFO (high frequency oscillations), 120-250

Hz.

1072

Figure 3. GRIN lens implant does not affect theta oscillations and enables combined

calcium imaging and optogenetics. a, diagram showing recording configuration (horizontal

plane), with LFP electrodes in CA1 of both hemispheres, and a GRIN lens only in the right

hemisphere (top). Example theta oscillations recorded in both hemispheres using wavelet

spectrograms, raw (black) and theta filtered (red) traces. **b**, relative theta power measured in

the right hemisphere of mice implanted with electrodes and GRIN lenses (blue) versus

electrodes only (green). **c**, the relationship between miniscope LED input value and measured

nominal power (purple) as well as actual light power corrected for the surface of a GRIN lens

(blue). **d**, corresponding theta power from LFP electrode attached to GRIN lens when using

different miniscope LED emitting light power values (in the absence of optogenetic

stimulations). **e**, effects of MS optogenetic stimulations on theta oscillations during miniscope

blue light emission through the lens (8AU LED power, corresponding to ~0.3mW/mm2).

1085

Figure 4. MS optogenetic stimulation modulates the activity of a subpopulation of CA1

pyramidal cells. a, recording CA1 neurons during MS optogenetic stimulation (5s ON, 5s

OFF) while simultaneously imaging CA1 neurons in mice freely exploring an open field. **b**,

peristimulus time raster plot of binarized activity for pairs of excited (left) or inhibited (right)

neurons. **c**, scatter plot for all pooled neurons (n = 1849 cells, N = 5 mice) of activity

probability during baseline vs stimulation (red, top: scrambled; blue, bottom: 8 Hz), for both

idle (left) and running (right) periods. **d**, portion of total cells significantly modulated by

stimulations (N = 5 mice). **e**, portion of modulated cells that are excited (green) or inhibited

(red). **f**, spatially tuning of example neurons before (top panels) or during (bottom panels)

stimulations. For the baseline condition, comparable 5 s epochs were used to sample spatial

tuning (baseline A vs B). Dark blue, low activity; bright yellow, high activity. Top number

indicated peak activity likelihood for each neuron. **g**, average stability of spatial tuning

between stimulated and unstimulated epochs during baseline, scrambled stimulation, and 8 Hz

stimulation sessions (n = 5 mice).

1100

Figure 5. MS optogenetic stimulation does not disrupt the stability of spatiotemporal

codes. a, we recorded mice on the 3-tone linear track to identify time-, place- and distance-

modulated cells (see fig. 1 and Methods). **b**, experimental timeline. **c**, spatial footprints of

neurons recorded over days and comparison scheme used to assess stability (top). Example

neuron (black arrow) tracked over the 5 experimental days (bottom). **d**, portion of total cells

encoding for one or several (conjunctive) variables before and during 8 Hz and frequency

scrambling optogenetic stimulations (N = 5 mice). **e**, sorted place fields for identified place

cells across day pairs. The first day is used as a baseline and mice undergo 8 Hz, scrambled,

or no stimulations (control treatment) during the other day. Stability is computed as the

pairwise correlation of fields between the two-day pairs. **f**, same for time-modulated cells. **g**,

same for distance-modulated cells. **h**, corresponding average stability for place-modulated

cells (each dot represents a cell pooled from N = 5 mice). **i**, same for time-modulated cells. **j**,

same for distance modulated cells.

1114

Figure 6. MS optogenetic stimulation alters behavior but does not disrupt

spatiotemporal codes. Using a naive Bayesian classifier, location (**a**,**b**), time elapsed (**c**,**d**),

and distance traveled (**e**,**f**) were predicted from CA1 neuronal activity. **a**,**c**,**e**, decoding error

(euclidean distance) for baseline, 8 Hz, and scrambled optogenetic stimulation conditions for

location, time, and distance, respectively (computed in a representative mouse). **b**,**d**,**f**,

decoding z-score (higher values indicate better decoding compared to shuffled surrogate) for

location, time, and distance, respectively, using N = 5 mice. **g**, example scoring of errors

(purple stars) in the tone-cued linear track task. **h**, percent of correct runs during baseline, 8

Hz, and scrambled stimulations.

1124

Figure 7. MS optogenetic stimulations disrupt maintenance and retrieval, but not

encoding of episodic and working memory. a, mice were implanted with fiber optics in the

MS after transfecting ChrimsonR (top). They were then subjected to the novel object place

recognition task (bottom, see Methods). **b**, in this task, both scrambled (red) and 8 Hz (blue)

disrupted memory performance when stimulating during the test phase. **c**, to further examine

the effect of MS stimulation on memory encoding, maintenance, and retrieval, mice were

trained in a delayed non-match to sample (DNMTS) task in an automated T-maze. **d**, mice

were trained (in absence of stimulation) to choose the correct, non-matching arm until the

performance exceeded a criterion of 0.8 portion of correct choices per day, for at least two

consecutive days. **e**,**f**,**g**, performance during stimulation at different phases of the task for

mice injected with ChrimonR (top) or YFP controls (bottom). Red shading indicates the

stimulated regions of the maze. **e**, average daily performance when performing MS

stimulations during encoding only (forced choice). **f**, average daily performance when

performing MS stimulations during the 10 s delay period only. **g**, average daily performance

when performing MS stimulations during the retrieval only (free choice).

1140

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 7