Non-neural factors influencing BOLD response magnitudes within individual subjects

Jan W. Kurzawski[1], Omer Faruk Gulban[2,3], Keith Jamison[4], Jonathan Winawer[1*], Kendrick Kay[5*]

[1]Department of Psychology, New York University, New York, NY, 10003, USA
[2]Cognitive Neuroscience Department, Faculty of Psychology and Neuroscience, Maastricht University, 62229, The Netherlands
[3]Brain Innovation, Maastricht, 62229, The Netherlands
[4]Department of Radiology, Weill Cornell Medicine, New York, NY, 10021, USA
[5]Center for Magnetic Resonance Research, Department of Radiology, University of Minnesota, Minneapolis, MN, 55455, USA

*Co-senior author

Corresponding Author: Jan W. Kurzawski, Department of Psychology, 6 Washington Place, New York University, NY, 10003 (jan.kurzawski@nyu.edu)

Keywords: BOLD signal, fMRI, vasculature, horizontal meridian, vertical meridian.

Number of pages: 23

Number of figures: 8

Number of words for: abstract | introduction | discussion: 217/250 | 700/650 | 1703/1500

Conflict of interest: The authors declare no competing financial interests.

Acknowledgments: This research was funded by the US National Eye Institute R01-EY027964, R01-MH111417 and R01-EY027401 to JW. Collection of the NSD dataset was supported by NSF CRCNS grants IIS-1822683 (to KK) and NSF IIS-1822929.

Abstract

To what extent is the size of the blood-oxygen-level-dependent (BOLD) response influenced by factors other than neural activity? In a re-analysis of three neuroimaging datasets (male and female human participants), we find large systematic inhomogeneities in the BOLD response magnitude in primary visual cortex (V1): stimulus-evoked BOLD responses, expressed in units of percent signal change, are up to 50% larger along the representation of the horizontal meridian than the vertical meridian. To assess whether this surprising effect can be interpreted as differences in local neural activity, we quantified several factors that potentially contribute to the size of the BOLD response. We find relationships between BOLD response magnitude and cortical thickness, curvature, depth and macrovasculature. These relationships are consistently found across subjects and datasets and suggest that variation in BOLD response magnitudes across cortical locations reflects, in part, differences in anatomy and vascularization. To compensate for these factors, we implement a regression-based correction method and show that after correction, BOLD responses become more homogeneous across V1. The correction reduces the horizontal/vertical difference by about half, indicating that some of the difference is likely not due to neural activity differences. We conclude that interpretation of variation in BOLD response magnitude across cortical locations should consider the influence of the potential confounding factors of thickness, curvature, depth and vascularization.

Significance statement

The magnitude of the BOLD signal is often used as a surrogate of neural activity, but the exact factors that contribute to its strength have not been studied on a voxel-wise level. Here, we examined several anatomical and measurement-related factors to assess their relationship with BOLD signal magnitude. We find that BOLD magnitude correlates with cortical anatomy, depth and macrovasculature. To remove the contribution of these factors, we propose a simple, data-driven correction method that can be used in any functional magnetic resonance imaging (fMRI) experiment. After accounting for the confounding factors, BOLD magnitude becomes more spatially homogenous. Our correction method improves the ability to make more accurate inferences about local neural activity from fMRI data.

Introduction

The blood-oxygen-level-dependent (BOLD) signal measured by fMRI is an important tool for non-invasive study of the human nervous system. However, the neural mechanisms underlying BOLD remain an active area of investigation (Herman et al., 2017). One clear conclusion is that the BOLD signal is strongly influenced by neural activity (Arthurs et al., 2000; Heeger et al., 2000; Attwell and Iadecola, 2002; Heeger and Ress, 2002; Logothetis, 2002; Lee et al., 2010; Siero et al., 2014). For a given location in the brain, and within a constrained paradigm (e.g., viewing different images and measuring the response that they elicit in visual cortex), the BOLD signal magnitude appears to be lawfully related to basic measures of neural activity. For example, as stimulus contrast increases, neural firing rates

and BOLD magnitude increase in proportion (Heeger et al., 2000). Similarly, increase in
coherence of stimulus motion boosts BOLD magnitude and firing rates in V5/MT (Britten et
al., 1993; Rees et al., 2000). When comparing different experimental paradigms or different
brain locations, however, it is less clear how to interpret differences in the magnitude of the
BOLD signal. For example, seeing a stimulus and expecting a stimulus can both elicit robust
BOLD signals in V1, but the underlying neural activity is very different in the two paradigms
(Sirotin and Das, 2009; Herman et al., 2017). It is also the case that similar BOLD signal
magnitudes in two locations may be linked to very different underlying neural activity.
These two limitations are reviewed by (Logothetis, 2008).
90
There are several reasons to believe that BOLD signal magnitudes, even within a fixed
experimental paradigm, are influenced by factors that are not directly related to neural
activity. The BOLD response, quantified in terms of percent signal change, can be especially
high in voxels containing large veins (Menon et al., 1993; Kim et al., 1994; Hoogenraad et al.,
1999; Kay et al., 2019) or unusually low, delayed, and/or displaced in voxels near cerebral
sinuses (Winawer et al., 2010; Jamison et al., 2017). The choice of MRI sequence, field
strength (van der Zwaag et al., 2009), and sequence parameters like echo time (Gorno-
Tempini et al., 2002) can also affect BOLD signal magnitude, and these effects may vary
across the brain (Herman et al., 2017). Indeed, it has been reported that BOLD may vary
across the cortex up to 40% simply due to different orientation of vasculature relative to the
direction of the static magnetic field (Gagnon et al., 2015a; Gagnon et al., 2016; Viessmann
et al., 2019). Furthermore, recent high-resolution fMRI studies have shown that BOLD signal
magnitude clearly depends on cortical depth. It is highest in the superficial depths which are
positioned near large pial veins and decreases with depth (Polimeni et al., 2010; Koopmans
et al., 2011; Zimmermann et al., 2011; Yu et al., 2014; Fracasso et al., 2016a; Fracasso et al.,
2016b; Dumoulin, 2017; Dumoulin et al., 2018; Kay et al., 2019; Self et al., 2019; van Dijk et
al., 2020).
108
In this paper, we study variations in BOLD signal magnitude within a fixed paradigm,
focusing our efforts on primary visual cortex (V1). We believe that by focusing on a single
brain region in well-controlled visual paradigms, we are in the best position to derive sound
interpretations of differences in BOLD signal magnitudes across the cortex. In three distinct
datasets, we demonstrate large differences between the meridian locations: the BOLD
magnitude in V1 is up to 50% higher along the representation of the horizontal meridian
than along the representation of the vertical meridian. We then investigate the potential
basis of these inhomogeneities by analyzing factors that are in principle distinct from neural
activity. As non-neural factors we consider cortical curvature, cortical thickness, cortical
depth, presence of macrovasculature (as indexed by bias-corrected EPI intensity), angle with
respect to B_0 magnetic field and radiofrequency (RF) coil bias. We motivate the selection of
these factors in the Methods. We find that several of these factors are systematically
related to observed variation in BOLD magnitudes across V1. To remove their influence, we
propose a simple correction method and show that the correction increases BOLD signal
homogeneity across V1, reducing the difference in response across the horizontal and
vertical meridians by about half.
125
126

Methods

Datasets

We used three publicly available visual fMRI datasets: the Human Connectome Project (HCP) 7T Retinotopy Dataset (Benson et al., 2018), the Natural Scenes Dataset (NSD) (Allen et al., 2021), and the Temporal Decomposition Method (TDM) Dataset (Kay et al., 2020). All data were acquired on 7T MR scanners using gradient-echo pulse sequences (technical details provided in Table 1). The datasets varied in stimulus properties and experimental design. HCP stimuli consisted of rings, wedges, and bars in a retinotopic mapping experiment; NSD stimuli consisted of natural scene images; and TDM stimuli consisted of high-contrast rings presented at different eccentricities. Experimental details are shown in **Figure 1**. The analyses performed in this paper start with pre-processed data from each dataset.

Temporal Decomposition Method Dataset (TDM)
- 5 subjects (2 males and 3 females)
- 0.8 x 0.8 x 0.8 mm³
- Rings varying in eccentricity
- 9 runs per subject
- TR = 2.2 s
- 3.5 s ON / 0.5 s OFF
- Maximum eccentricity = 5.5 degrees

Natural Scenes Dataset (NSD)
- 8 subjects (2 males and 6 females)
- 1.8 x 1.8 x 1.8 mm³
- Natural scene images
- 360-480 runs per subject
- TR = 1.6 s
- 3 s ON / 1 s OFF
- Maximum eccentricity = 4.2 degrees

Human Connectome Project (HCP)
- 181 subjects (72 males and 109 females)
- 1.6 x 1.6 x 1.6 mm³
- Textures viewed through spatial apertures
- 6 runs per subject
- TR = 1 s
- Continuous design
- Maximum eccentricity = 8 degrees

Figure 1 – Datasets used in this study. Stimulus images for each of the datasets are shown. For TDM, stimuli consisted of 6 rings varying in eccentricity. For NSD, stimuli consisted of natural scene images. For HCP, the experiment consisted of several retinotopic mapping runs that included expanding and contracting rings, rotating wedges, and moving bars filled with a colorful object-based texture. Additional acquisition details are provided in Table 1.

Dataset	TDM	NSD	HCP
Field strength	7T	7T	7T
TR	2200 ms	1600 ms	1000 ms
TE	22.4 ms	22.0 ms	22.2 ms
Flip angle	80	62	45
Number of slices	84	84	85
Matrix size	200×162	120×120	130×130
Field of view	160 mm $\times$ 129.6 mm	216 mm $\times$ 216 mm	208 mm $\times$ 208 mm

Nominal spatial resolution	0.8 mm	1.8 mm	1.6 mm
Multiband factor	2	3	5
iPAT factor	3	2	2
Partial Fourier	6/8	7/8	7/8

146 **Table 1 – Details on the fMRI pulse sequence used in each of the datasets.** Each column describes different
147 dataset.

Extracting BOLD magnitude

From each dataset, we extracted a measure of BOLD signal magnitude at each cortical
surface vertex. For TDM, we started with the pre-processed fMRI time-series data provided
with the dataset and analyzed the data with a GLM. Specifically, we convolved a canonical
HRF with stimulus onsets to create a regressor for each experimental condition, and then
used these regressors with GLMdenoise (Kay et al., 2013b) to estimate a beta weight for
each condition. We computed the maximum beta weight across all conditions for each voxel
as the measure of BOLD signal magnitude. These results are defined at six different depths
(equidistant from 10% to 90% of the cortical thickness) in each subject's native surface
space. (Depth assignment was achieved by a spatial interpolation of each fMRI volume at
the locations of the six depth-dependent cortical surfaces; see Kay et al. (2020) for details.)
For NSD, we took the 'meanbeta' values (1 mm data preparation, beta version 2) provided
with the dataset; these values indicate the average BOLD percent signal change observed
across all stimulus trials and all scan sessions. We then mapped these values to the 3 depth
surfaces provided in NSD (positioned at 25%, 50%, and 75% of the cortical thickness). The
HCP dataset was previously analyzed (Benson et al., 2018) with a population receptive field
(pRF) model (Dumoulin and Wandell, 2008) implemented in analyzePRF (Kay et al., 2013a).
The model includes a gain parameter that describes the amplitude of the BOLD response of
a given voxel (or vertex) to the object-based texture (covering the entire pRF) for a single
repetition time (TR = 1 s). We quantified BOLD in terms of percent signal change (%BOLD) by
dividing the gain parameter by mean signal intensity and multiplying by 100. The results are
prepared in FreeSurfer's fsaverage space.

Visual field mapping

We used retinotopic mapping to divide the primary visual cortex into a set of regions. For
HCP, we used polar angle and eccentricity estimates available from the data release. For the
TDM and NSD datasets, we mapped Benson's polar angle and eccentricity atlas using
neuropythy software (Benson and Winawer, 2018). We use the following convention for all
3 datasets: the upper vertical meridian corresponds to 0 deg, the horizontal meridian
corresponds to 90 deg, and the lower vertical meridian corresponds to 180 deg. Note that
the polar angle estimates are rescaled for the correlation and linear regression analysis (see
next section). We used Benson's definition of the extent of visual areas V1, V2, and V3 for all
3 datasets (Benson et al., 2014).

Quantification of non-neural factors

In the TDM and NSD datasets, we quantified several factors that might be related to variation in the magnitude of the BOLD signal across cortical locations. We focused on factors that can be easily extracted from either functional or anatomical data that are typically acquired in an fMRI experiment. For the purposes of the present study, we consider only within-subject factors rather than across-subject factors, with the goal of removing non-neural influences on the variation of BOLD magnitudes across voxels. We note that there are several other factors that influence variation of overall BOLD magnitude across subjects like caffeine use (Liu et al., 2004), vascular age (Tsvetanov et al., 2021), and heart rate (Chang et al., 2009). Below, we describe each of the within-subject factors that we considered in the present study.

Curvature was obtained from FreeSurfer outputs (Dale et al., 1999; Fischl and Dale, 2000), and refers to the geometry of the folding pattern of the cortical surface. Negative values correspond to gyri while positive values correspond to sulci. Curvature is quantified as $1/r$, where r is the radius of an inscribed circle measured in mm.

Thickness was also obtained from FreeSurfer outputs. It is measured in mm and corresponds to the distance between the outermost (close to cerebrospinal fluid) and innermost (close to white matter) boundaries of gray matter. Curvature and thickness are well known to vary across visual cortex. Their relationship with %BOLD remains unknown and has not been investigated in detail, especially on a voxel-by-voxel basis. We include these factors in our analysis to assess whether these anatomical factors have systematic relationships with BOLD magnitude.

Mean bias-corrected EPI was calculated as the mean signal intensity in the fMRI data divided by the estimated RF coil bias (details below). The units range from approximately 0 to 2, and indicate percentages (e.g., 0.5 means 50% of the strength of typical signal intensities). Mean bias-corrected EPI values can be viewed as high spatial frequency changes in signal intensity across space. We include this factor in the analysis as mean bias-corrected EPI was previously found to be a good predictor for venous effects (Kay et al., 2019). Proximity to veins often results in increased BOLD magnitude.

Depth was estimated by generating 6 cortical surfaces (for TDM) or 3 cortical surfaces (for NSD) equally spaced between 10% and 90% (for TDM) or 25% and 75% (for NSD) of the distance from the pial surface to the boundary between gray and white matter. These surfaces are numbered from 1 to n, where 1 is outermost and n is innermost. We include depth as a factor as it is well known that BOLD magnitude is highest in superficial depths and decreases towards the white matter (Polimeni et al., 2010).

Angle with respect to B_0 was calculated by considering the angle (*theta*) between the pial surface normal and the direction of the B_0 static magnetic field as estimated from NIFTI header information. Angle was quantified in degrees and was normalized as abs(*theta*–90) such that a final value of 0 deg indicates that the cortical surface is parallel to the magnetic field and a final value of 90 deg indicates that the cortical surface is perpendicular to the

magnetic field. We include angle with respect to B_0 in the analysis because previous reports
showed that the BOLD magnitude varies with B_0 angle (Gagnon et al., 2015b).

RF coil bias was taken to be the result of fitting a 3D polynomial to the mean signal intensity
in the fMRI data. The values are in raw scanner units and represent low spatial frequency
changes in the intensity of voxels. This estimation method has been used previously (Kay et
al., 2019). We include RF coil bias as a control in our analysis. In theory, there should not be
a systematic relationship between RF coil bias and BOLD magnitude, as we express BOLD
magnitudes at each voxel in terms of percent signal change (as is typically done in the field),
and percent signal change is sensitive to an overall scale factor on the signal.

In sum, all of these factors are known to vary across the cortical surface of V1. The exact
biophysical mechanisms that might explain their impact on %BOLD are in some cases
unknown (e.g., curvature). In other cases, we expect that some factors should not bear
systematic relationships to %BOLD (e.g., RF coil bias). In general, the work here is intended
to be a first step towards understanding the influence of potential non-neural contributions
to variations in %BOLD across individual voxels within a given subject.

Quantification of neural factors
249

Polar angle was obtained from Benson's atlas (Benson et al., 2014), representing the visual
field angle to which each cortical location is optimally tuned. For the purposes of our
analyses, we normalize polar angle such that 0 deg corresponds to the horizontal meridian
and 90 deg corresponds to the upper and lower vertical meridians. We include polar angle
as a positive control: we expect that polar angle should bear a systematic relationship with
BOLD magnitude, as this is the original observation that motivated the present study.

Definition of regions of interest
258

Using the visual field mapping results, we defined regions of interest (ROIs) corresponding
to the representation of the horizontal and vertical meridians within V1. The ROIs were
defined by limiting the eccentricity to the maximum stimulus eccentricity used in each
dataset and limiting the angle to a specific range (e.g., to create a V1 ROI for the upper
vertical meridian with a width of 20 deg, we created a mask where polar angle estimates
were higher than 0 and lower than 20).

Modelling variation in BOLD signal magnitude
267

To account for non-neural contribution to %BOLD, we used a multiple regression model. The
modeled data (Y) consisted of the %BOLD value observed at each surface vertex in visual
areas V1–V3. Although this study focuses on BOLD homogeneity in V1, we include %BOLD in
V1–V3. This is because we are attempting to establish relationships that might generalize
across different cortical regions. Furthermore, if we were to include only vertices in V1, we
would be at high risk of removing genuine neural activity differences (e.g. those that may
exist between the horizontal and vertical meridians) that correlate with the non-neural
factors.

277
The variables used to model the data included thickness, curvature, depth and mean bias-
corrected EPI intensity. (Only these four factors showed evidence of being substantially
related to BOLD magnitude; see Results.) The variables were standardized (z-scored) and,
together with a constant term, were included as predictors in the design matrix (X).
Ordinary least-squares estimates for beta weights were obtained in the following linear
model:
284
285 $$Y = X\beta + n \qquad \text{(Eq. 1)}$$
286
287 where Y is the %BOLD magnitude at each vertex, X is the 5-column design matrix, β is a set
288 of beta weights (5 per vertex), and n is a set of residuals.
289
290 **Major cortical sulci**
291
292 In several figures we show outlines of major cortical sulci. These include the calcarine sulcus
293 (CALC), parieto-occipital sulcus (POS), intraparietal sulcus (IPS), occipitotemporal sulcus
294 (OTS), and superior temporal sulcus (STS). These sulci were manually labelled on the
295 fsaverage surface and then mapped to each individual's native surface.
296
297 **Data and code availability**
298
299 The datasets used in this paper are freely available online: NSD
300 (http://naturalscenesdataset.org), HCP (https://osf.io/bw9ec/), and TDM
301 (https://osf.io/j2wsc/). Code that reproduces the main figures in this paper is available at
302 https://github.com/jk619/meridianbias/. Associated data files are available at
303 https://osf.io/2nc4x/.

Results

Stronger BOLD responses along the V1 horizontal meridian

We examined BOLD response magnitudes in three freely available datasets: the Natural
Scenes Dataset (NSD; Allen et al., 2021), the data used for the Temporal Decomposition
Method (TDM; Kay et al., 2020), and the Human Connectome Project 7T Retinotopy Dataset
(HCP; Benson et al., 2018). Each dataset contains BOLD responses to different types of visual
stimulation (see Methods). We defined one region of interest (ROI) for the horizontal
meridian (HM) and one for the vertical meridian (VM) (**Figure 2A–B**). These ROIs represent a
wedge-shaped region in the visual field centered at the horizontal meridian with a width of
40 deg (horizontal) and two wedges abutting the vertical meridian each with a width of 20
deg (vertical).
316
In each of the three datasets, we compared BOLD magnitudes expressed in percent signal
change (%BOLD) observed for the VM with BOLD magnitudes observed for the HM (**Figure
2C–E**). In each dataset, we find higher %BOLD in the HM ROIs compared to the VM ROIs. We

summarize this difference with an asymmetry ratio: (HM−VM)/mean(HM,VM). All datasets
show strong asymmetry, with an asymmetry ratio of ~30%. Positive values for the
asymmetry ratio indicate greater response for the horizontal meridian. (Note that if the
asymmetry is expressed as a percentage of the smaller vertical meridian response, the
increase reflected in the larger horizontal meridian response is up to ~50%.)
325

326
327
Figure 2 – BOLD magnitude is higher at the horizontal meridian in V1. **A)** Polar angle map of group-average
HCP subject (999999) with V1 boundary outlined in dotted black lines. **B)** Horizontal and vertical regions of
interest (ROIs) are indicated in gray and magenta, respectively. White text indicates major brain sulci (see
Methods). **C-E)** Mean BOLD magnitude for horizontal and vertical ROIs in the three datasets. Error bars
indicate standard error across subjects.

One possibility is that the horizontal and vertical V1 BOLD responses are in fact similar, but
the vertical ROIs appear to have lower signal due to mixing with signal from V2. V2 and V1
border along the vertical meridian representation, and blurring might occur either in
acquisition or in pre-processing and analysis. To further our understanding of the V1
response asymmetries, we re-computed asymmetry ratios using smaller wedges at many
locations (**Figure 3A**). Note that, because we use smaller wedges, the asymmetry at the
cardinal meridians is different from **Figure 2**. While the asymmetry is strongest at the
cardinal meridians, some horizontal/vertical asymmetry is found at least 30 deg away from
the meridians in all three datasets (**Figure 3B**). This argues against the explanation that the
asymmetry is caused by spillover from V2.
343
344

Figure 3 – BOLD asymmetries generalize to off-cardinal locations. To further understand V1 BOLD asymmetry, we manipulated the location of the wedge ROIs in 5-deg increments. **A)** The upper row shows the visual field location of wedge ROIs and the lower row shows the corresponding cortical locations (flattened left hemisphere). For clarity, we show only every other set of ROIs. **B)** Asymmetry ratio as a function of angular distance from the cardinal meridians. Error bars indicate standard error across subjects.

HM/VM asymmetry persists at inner cortical depths

The BOLD signal is strongly influenced by properties of the brain's vasculature. Uneven venous contributions across the brain can cause variation in BOLD magnitude (Menon et al., 1993; Kim et al., 1994; Hoogenraad et al., 1999; Kay et al., 2019). One possibility is that the meridian asymmetries we observe arise from non-uniformities in the vascular network. To investigate this possibility, we took advantage of the sub-millimeter resolution of the TDM dataset and examined HM/VM asymmetry as a function of depth. Because macroscopic venous effects are larger in the superficial cortex due to large pial veins (Duvernoy et al., 1981; Turner, 2002; Polimeni et al., 2010; Kay et al., 2019), by sampling BOLD responses from deeper depths, we minimize contributions from pial veins. We find that the HM/VM asymmetry is larger at the superficial depths, suggesting that part of the asymmetry may be due to differential properties in macroscopic vasculature (**Figure 4**). This depth effect is systematic: every subject shows higher asymmetry at the superficial depth than the middle depth. Nonetheless, there remains a substantial horizontal/vertical asymmetry at all depths (**Figure 4**), suggesting that macroscopic vessels near the pial surface are not the entire explanation. At the innermost depth sampled, which is least influenced by pial vessels, the

368   HM/VM asymmetry is 26% (average across subjects) and is positive in each of the 5 subjects.
369   The middle depths appear to have the least asymmetry. This could be due to a difference in
370   neural responses at intermediate depths, which generally correspond to input-related
371   cortical layers.

372
373

Figure 4 – BOLD asymmetry in V1 persists at inner depths. We exploit the high-resolution TDM dataset to
discriminate V1 BOLD responses across depth and estimate response asymmetries as a function of depth
(asymmetry is calculated in the same way as in **Figure 3**). The presence of asymmetry at the innermost depth
suggests that response asymmetries exist even with minimal contribution of large pial veins.

Assessing and modeling non-neural contributions to BOLD signal magnitude

In addition to vascular effects, other factors unrelated to neural activity evoked by the
experimental manipulation may influence variation in %BOLD across the cortical surface.
These additional factors are often neglected in fMRI analysis pipelines. Although some of
the factors are known to vary across the cortex, their influence on the BOLD signal is poorly
understood. Here, we attempt to understand how these factors may be related to BOLD
magnitude variations. To the best of our knowledge, we are unaware of any previous study
that has examined this issue in detail, especially at the level of individual voxels (or vertices)
within individual subjects.

We first identified a list of possible confounding factors (beyond cortical depth, which we
have already introduced) based on consideration of basic anatomical properties of the brain
and the nature of fMRI measurement. These factors are cortical curvature, cortical
thickness, RF coil bias, mean bias-corrected EPI signal intensity, and angle with respect to B_0.
Each of these factors can be interpreted as spatial maps, with a value at each vertex on the
cortical surface mesh. The five maps can be obtained from standard anatomical scans (T1-
weighted) or from the fMRI measurements themselves without additional MRI experiments

(see Methods for details). Example surface visualizations of these maps together with
%BOLD are shown in **Figure 5.** We hypothesize that inhomogeneities in some of these maps
might explain some of the observed inhomogeneity in %BOLD across V1.

399
400

Figure 5 – Variation in anatomical and acquisition factors across cortex. Each sphere shows data mapped on
the left hemisphere for subject S1 in the NSD dataset. Below each surface map is a histogram of the plotted
values. White outlines indicate major cortical sulci. %BOLD represents the average response to the natural
scene stimuli used in the NSD dataset. Some of the spatial variability in %BOLD might be due to variability in
the depicted non-neural factors.

To understand the potential relationships amongst these five identified factors and %BOLD,
we first performed voxel-wise correlation analyses. For these analyses, we used the TDM
dataset, as its high spatial resolution facilitates the identification of vascular effects (Kay et

al., 2019). We examined data from V1–V3 where neural activity magnitudes can be
expected to be relatively homogeneous (although biases were reported before; Liu et al.,
2006) given the simple contrast patterns used. In **Figure 6A**, we show pairwise correlations
across these five quantities, as well as retinotopic polar angle preference (rescaled between
0 and 90; see Methods) and cortical depth. We find that %BOLD correlates substantially
with four factors: curvature ($r = 0.26$), thickness ($r = -0.17$), mean bias-corrected EPI
intensity ($r = -0.25$), and depth ($r = -0.27$). We do not find a strong correlation between
%BOLD and polar angle. Although results from **Figure 2C–E**, **Figure 3C** and **Figure 4** suggest a
strong negative correlation, the previous analysis included data only from V1. Here we
analyze vertices from V1-V3 where this relationship becomes weaker ($r = -0.05$). Overall, we
can summarize as follows: %BOLD extracted from V1–V3 tends to be higher at locations that
correspond to sulci, in thinner parts of the cortex, in voxels with lower mean bias-corrected
EPI intensities, and at more superficial depths.
423
Examination of correlations amongst factors yields additional insights (**Figure 6A**). The
strongest correlation that we find is between curvature and thickness ($r = -0.28$), indicating
that sulci tend to be thin. Curvature is correlated with mean bias-corrected EPI ($r = 0.16$) and
with polar angle ($r = -0.19$), and thickness is correlated with polar angle ($r = 0.15$). Our
interpretation of these effects is that venous effects tend be stronger in gyri (consistent with
previous findings in Kay et al., 2019), and that the correlations related to polar angle simply
reflect the tendency for horizontal meridian representations to fall on sulci (e.g. the
calcarine sulcus). Overall, these complex relationships suggest that making sense of non-
neural influences on %BOLD requires a broad perspective that considers multiple factors.
433

434
435
Figure 6 – Modeling variations in BOLD signal magnitude. A) Correlation (Pearson's *r*) between a variety of
factors and %BOLD extracted from V1–V3 from the TDM dataset. Main plot shows results from data
concatenated across all subjects, while inset plots show results from individual-subject data. P-values indicate
significance of one sample *t*-test across subjects; **p* < 0.05; ***p* < 0.01; ****p* < 0.001. **B)** Regression model for

%BOLD. Based on the results of panel A, we selected curvature, thickness, depth and mean bias-corrected EPI
as the main non-neural factors that confound %BOLD. These four factors were then used in a multiple linear
regression model to predict %BOLD (top). The amount of variance explained by the model is shown in the
inset.

Correcting BOLD signal magnitude for non-neural factors

444 **Correcting BOLD signal magnitude for non-neural factors**
445
We now explore whether we can develop a statistical model to compensate for the
influence of non-neural factors on %BOLD. We operate under the assumption that any
observed correlation between the factors and %BOLD is incidental and does not reflect
genuine neural activity variation. Our model is a multiple regression model (**Figure 6B, top**)
that uses the main factors of curvature, thickness, depth and mean bias-corrected EPI
intensity as continuous variables and attempts to determine a weighted sum of these
factors that optimally accounts for variations in %BOLD across cortical locations (see
Methods for details).
454
Fitting the model, we find a strong positive contribution of curvature and negative
contributions of thickness, mean bias-corrected EPI intensity and depth (**Figure 6B, bottom**),
consistent with the earlier voxel-wise correlation analyses. Estimated beta weights are fairly
consistent across subjects, and the model on average across subjects explains 26% of the
variance in %BOLD. A multiple regression model using all 6 factors (adding RF coil bias and
angle with respect to B_0) resulted in only minimally larger explained variance, 27% vs. 26%,
consistent with the earlier correlation analyses indicating that RF coil bias and angle with
respect to B_0 bear little or no relationship with %BOLD.
463
To better understand the relationship between the identified non-neural factors and
%BOLD, we construct a 2D histogram relating the model fit (BOLD prediction based on non-
neural factors obtained by multiplying the design matrix and estimated beta weights) and
%BOLD (**Figure 7A**). This reveals a clear nonlinear relationship. To accommodate this
nonlinearity, we fit a nonlinear function relating the linear model fit and %BOLD (blue line in
Figure 7A). Finally, we remove the contribution of non-neural factors by dividing %BOLD
observed at each cortical location by the fit of the nonlinear model. We divide %BOLD by
the model fit rather than subtracting the model fit, as we believe that the influence of non-
neural factors on %BOLD might impose a type of 'gain' field on fMRI responses observed in a
given experiment. For example, if there is an excess of macrovasculature in a voxel, we
would expect the overall amplitude of the BOLD response from the voxel to be scaled. Note
that our method of rescaling BOLD magnitudes does not change the pattern of responses
across different experimental conditions within a voxel (while a subtractive approach
would). For example, if the response to condition A is 25% higher than the response to
condition B, this will continue to be the case after rescaling.
479

$$Modelfit = p(1) * (X * Beta)^{p(2)} + p(3)$$

$$Corrected\ \%BOLD = \frac{\%BOLD}{Modelfit}$$

Figure 7 – Correction of V1 BOLD inhomogeneity. A) Removal of non-neural factors. First, linear combinations
of non-neural factors are used to predict %BOLD within V1–V3 using the TDM dataset. The model is fit on data
concatenated from all 5 TDM subjects. The model is augmented with a nonlinear power-law function (blue
line), which is controlled by a gain parameter ($p(1)$), an exponent parameter ($p(2)$), and a constant term ($p(3)$).
B) Each voxel's BOLD responses are divided by the model fit, yielding the corrected %BOLD. **C)** BOLD signal
magnitude within V1 before and after the correction (TDM dataset, subject S3, most superficial depth).
Asterisk indicates the fovea and dashed lines indicate the boundary between V1 and V2. After correction,
some vertices with very high BOLD are eliminated (see white arrows). Within each plot, the color range
extends from 0 to the maximum. Each map has an associated histogram that shows all values extracted from
V1.

The result of the proposed correction procedure is shown in **Figure 7B**. We see that after
the correction procedure, the distribution of BOLD response becomes flatter, indicating the
efficacy of the procedure. (Note that what is important is the shape of the distribution of
the values, not necessarily the magnitudes of the values.) Increased homogeneity of BOLD
magnitude is also visible on the cortical surface (**Figure 7C**).

To understand whether our method generalizes across datasets, we used the same
procedure and performed correction on the NSD dataset. We summarize the effect of the
correction by showing the correlations between %BOLD and non-neural factors before and
after the correction (**Figure 8A**). The pattern of results before correction (**Figure 8A, top**) is
consistent across the TDM and NSD datasets, except for the reduced correlation with bias-

corrected EPI in NSD (see Discussion). Importantly, correlations after the correction are
substantially reduced, indicating the efficacy of the method.
To check whether accounting for non-neural factors increases the homogeneity of BOLD, we
quantified the variation of BOLD magnitudes across V1 before and after the correction.
Variation was quantified using the semi-interquartile range divided by the median (SIR).
Intuitively, if the spread of BOLD magnitudes is small (i.e., %BOLD is relatively
homogeneous), SIR will be low, whereas if the spread of BOLD magnitudes is large (i.e.,
%BOLD is relatively homogeneous), SIR will be high. We find that across subjects, the SIR
decreases from 0.42 before correction to 0.34 after correction for TDM and decreases from
0.48 to 0.42 for NSD.
512
We now return to the experimental effect that motivated this study, namely, BOLD
response asymmetries across the horizontal and vertical meridians in V1. We quantify the
asymmetry before and after correction (**Figure 8B**). We find that after accounting for the
non-neural factors, the asymmetry drops for TDM from 49.2% to 25.1% and from 40% to
18% for NSD. We thus suggest that some of the observed differences in BOLD response
magnitudes are due to non-neural factors.
519
520

521
522
Figure 8 – The effect of BOLD inhomogeneity correction. **A)** Voxel-wise correlation between the various
factors and %BOLD before and after correction. After correction, correlations are reduced, indicating that the
corrected data are less influenced by the non-neural factors. **B)** Dependence of %BOLD on polar angle in V1
before and after the correction for TDM dataset and NSD datasets. The asymmetry drops by about half.

In the results demonstrated in this paper, the correction method reduces inhomogeneities
between the horizontal and vertical meridians. But more generally, it is possible that in
other datasets, the method may reveal activity differences that are masked by non-neural
factors. For example, voxel A might have a lower neural response than voxel B, but voxel A
might reside close to a large vein which would tend to increase %BOLD. In conventional
fMRI analyses, both voxels might show similar BOLD magnitude, even though the underlying

neural activity is different. The methods proposed in this paper can be viewed as an attempt
to obtain better estimates of underlying neural activity.

Discussion

In this paper, we used three publicly available datasets to assess the degree of homogeneity
of BOLD signal magnitude in primary visual cortex. We found that stimulus-evoked BOLD
responses, expressed as percent signal change, are up to 50% stronger along the horizontal
meridian than the vertical meridian. To investigate whether these magnitude differences
can be attributed to differences in local neural activity, we systematically evaluated the
potential contribution of several non-neural factors to the observed effect. We found that
BOLD signal magnitude correlates with curvature, thickness, depth and macrovasculature
(as indexed by bias-corrected EPI intensities). Using a regression-based correction
procedure, we were able to increase the homogeneity of BOLD signal magnitude and found
that the meridian differences were reduced by half.
546
547 **Spatial variations in BOLD magnitude**
548
This study tackles the issue of the neural basis of variation in BOLD signal magnitude.
Specifically, we address variation in BOLD across cortical locations for a fixed experimental
manipulation, as opposed to variation in BOLD across experimental manipulations for a
fixed cortical location. The latter has been heavily studied (Heeger et al., 2000; Logothetis et
al., 2001; Heeger and Ress, 2002; Logothetis and Wandell, 2004; Mishra et al., 2021),
whereas the former has not yet been systematically studied to the best of our knowledge. If
there are indeed non-neural factors that influence BOLD signal variation, taking this into
account is critical when interpreting differences in fMRI responses across brain regions.
557
We acknowledge that a challenge in understanding the neural basis of the BOLD signal is
that directly comparable ground-truth measurements of neural activity are typically not
available. Moreover, the BOLD signal only indirectly measures the neural response, and its
magnitude likely depends on many aspects of neural activity. Increased BOLD signal might
be a consequence of more neurons firing, more spikes per neuron, changes in neural
correlation, changes in subthreshold activity, and/or changes in what kinds of neurons are
most active. Our approach currently does not try to distinguish amongst these causes.
565
566 In our analyses, we relied on the working assumption that the experimental paradigms of
567 the three datasets (combined with suitable averaging and analysis procedures) are expected
568 to generate relatively homogeneous patterns of neural activity in early visual cortex. Of
569 course, this may not be exactly the case.
570
571 **Non-neural factors that affect BOLD magnitude**
572
Mean bias-corrected EPI. Mean bias-corrected EPI is a convenient marker for macrovascular
contributions to the fMRI signal (Kay et al., 2019). Vertices contaminated by venous effects
show lower intensity values in mean EPI images and often result in high %BOLD magnitude.

In the TDM dataset, we found this to be the case and were able to remove, to some extent,
venous effects using the described correction method. We did not, however, find a strong
relationship between mean bias-corrected EPI and %BOLD magnitude in the NSD dataset.
We suggest that the reason for this apparent discrepancy is that effective discovery of
venous contributions requires high-resolution data where voxel size approaches the scale of
1 mm or better. Another important issue to consider is the cerebral sinuses. The sinuses are
the largest veins that drain blood from the brain and they exert major effects at certain
specific cortical locations. Complicating matters is the fact that the sinuses also produce low
EPI intensity, but instead of boosting BOLD magnitude they seem to reduce it, resulting in
low %BOLD (Winawer et al., 2010; Jamison et al., 2017). In the present study, we do not
attempt to isolate or analyze the effects of the cerebral sinuses, though preliminary analyses
indicate that the sinuses do not provide a simple explanation of the horizontal/vertical
asymmetry (data not shown).
589
Cortical anatomy. We find that curvature and thickness correlate with BOLD signal
magnitude (see **Figure 6A**). It is known that many anatomical properties vary with thickness
and with curvature (Jiang et al., 2021): (i) total neuron count is higher in gyri than it is in
sulci (Hilgetag and Barbas, 2005), (ii) gyri tend to be thicker than sulci (Welker, 1990;
Hilgetag and Barbas, 2005), (iii) venous effects (resulting in higher BOLD signal amplitude)
are more prominent in gyri than they are in sulci (Kay et al., 2019); and (iv) there may even
be intrinsic causal relationships between curvature and thickness during anatomical
development (Hilgetag and Barbas, 2005). However, the exact anatomical and biophysical
mechanisms that might link curvature and thickness to BOLD signal magnitudes are largely
unknown, to our knowledge. This is an important issue for future research. Here, we
operate under the working assumption that correlations between the BOLD signal and
curvature or thickness reflect incidental factors unrelated to local neural activity. We
therefore assume that a correction which removes their influence from the BOLD signal is
desirable.
604
Orientation of pial veins. It has been reported that regions where the cortical surface is
oriented perpendicular to the main magnetic field produce lower BOLD signal than regions
where the surface is oriented parallel (Gagnon et al., 2015a; Fracasso et al., 2018). The
proposed explanation is that this effect is caused by the orientation of pial veins, which lie
parallel to the cortical surface. Our analyses did not replicate this result and indicated little
relationship between BOLD magnitude and angle with respect to B_0 (see **Figure 6A**). One
possible explanation could be related to our pre-processing approach, in which fMRI signals
are sampled specifically in the gray matter and away from the pial veins that reside on top
of the gray matter. This may have dampened effects related to the pial veins. Nonetheless,
the prior literature would have predicted some B_0 effect even at inner cortical depths
(Viessmann et al., 2019). Alternatively, it is possible that the orientation effects depend in
some way on pulse sequence parameters, or the specific brain area being studied. A
detailed examination of different datasets would be necessary to resolve these
discrepancies.
619
RF coil effects. Due to cortical folding, gyri tend to be closer to the RF coil than sulci.
Locations that are further from the coil might have lower mean signal intensities and

therefore lower SNR (Srirangarajan et al., 2021), but this should not affect BOLD *magnitudes*
expressed in terms of percent signal change. We are not aware of any mechanism that
would alter the percent signal change in brain locations that are further away from the RF
coil. Indeed, we did not find any relationship between RF coil bias and BOLD magnitude (see
Figure 6A).
627
628
629 **Correction for the impact of non-neural factors**
630
Our results show that voxel-wise %BOLD is likely contaminated by several non-neural
factors.
To account for these factors, we developed a regression-based correction method. The goal
of this method was to introduce a simple, data-driven approach that can be applied
irrespectively of the specific experiment or brain region that is under consideration. The
underlying premise of the method is that by removing the contribution of non-neural
factors, the resulting measures would constitute a better representation of the underlying
neural activity. After application of the method, we found that %BOLD becomes more
homogenous and correlations between %BOLD and non-neural factors become significantly
reduced. Thus, our results indicate that some variation in %BOLD that might be interpreted
as change in neural activity likely reflects the variation of non-neural factors.
642
We believe the results presented in this paper constitute a first step towards developing a
cogent strategy for compensating for non-neural biases in BOLD signal magnitudes.
Suppressing the influence of non-neural factors has potential applications in pre-surgical
planning, where fMRI is routinely used to map motor, speech, and visual areas. The value of
fMRI for presurgical planning is currently limited by the accuracy of localizing neural
responses (Silva et al., 2018a). BOLD-derived maps that are a better representation of
neural activity could lead to more accurate neurosurgical interventions.
650
It remains to be seen whether the remaining asymmetry across the horizontal and vertical
meridians in V1 is a result of genuine neural activity differences, or an effect of other non-
neural factors that we were unable to quantify in the present study (which might require
additional MRI acquisition measures and/or higher resolution data). It is conceivable that
genuine neural activity differences may exist across the horizontal and vertical meridian
locations in V1. For example, there is greater cortical magnification along the horizontal
than vertical meridian (Silva et al., 2018b; Benson et al., 2021; Himmelberg et al., 2021;
Himmelberg et al., 2022), and it is plausible that this might be accompanied by differences
in the strength of neural responses.
660
Although our method is aimed towards more meaningful quantification of the BOLD signal,
it differs conceptually from quantitative BOLD (qBOLD) approaches (He and Yablonskiy,
2007; Yablonskiy et al., 2013; Cherukara et al., 2019). On the one hand, qBOLD attempts to
model the BOLD signal in terms of its underlying metabolic and hemodynamic components
(e.g., blood flow, blood volume, oxygenation extraction), and this in principle may yield
measures more closely related to neural activity. On the other hand, the approach we have
taken in this paper is to apply analytic methods to BOLD data that consider inhomogeneities

that may exist across the brain, with the goal of better estimating local neural activity. Note
that the two approaches are not mutually exclusive: one might imagine assessing whether
the magnitude of qBOLD measures co-vary with non-neural factors across the brain.
671
There are other methods that can be used to suppress the contribution of non-neural
factors to BOLD signal magnitudes. By identifying early and late components of evoked
hemodynamic responses, a temporal decomposition method can be used to estimate BOLD
response components more closely linked to the microvasculature, which presumably more
closely reflect local neural activity (Kay et al., 2020). Another analysis method focuses on
BOLD fluctuations where estimates of slow oscillations (< 0.1 Hz) are used to suppress
vascular-related effects (Kazan et al., 2016). Similarly, some methods use the amplitude of
fluctuations in resting-state data to rescale the BOLD signal (Di et al., 2013; Guidi et al.,
2020). Finally, acquisition methods, such as spin-echo pulse sequences, can be used to
suppress unwanted venous effects. Note that all these methods concern effects of the
macrovasculature, but systematic biases in BOLD signal magnitudes may in theory persist
even if BOLD responses were fully restricted to the microvasculature. Further investigation
is necessary to resolve these possibilities.
685

686 # Author Contributions

687
688 J.W.K, K.K. and J.W. conceived and designed this study. J.W.K and K.J. performed data
689 analyses, with K.K. and J.W. providing guidance. J.W.K., J.W, O.F.G and K.K. wrote and edited
690 the manuscript.
691
692
693

Allen EJ, St-Yves G, Wu Y, Breedlove JL, Prince JS, Dowdle LT, Nau M, Caron B, Pestilli F,
Charest I, Hutchinson JB, Naselaris T, Kay K (2021) A massive 7T fMRI dataset to
bridge cognitive neuroscience and artificial intelligence. Nature Neuroscience.
Arthurs OJ, Williams EJ, Carpenter TA, Pickard JD, Boniface SJ (2000) Linear coupling
between functional magnetic resonance imaging and evoked potential amplitude in
human somatosensory cortex. Neuroscience 101:803-806.
Attwell D, Iadecola C (2002) The neural basis of functional brain imaging signals. Trends
Neurosci 25:621-625.
Benson NC, Winawer J (2018) Bayesian analysis of retinotopic maps. Elife 7.
Benson NC, Butt OH, Brainard DH, Aguirre GK (2014) Correction of distortion in flattened
representations of the cortical surface allows prediction of V1-V3 functional
organization from anatomy. PLoS Comput Biol 10:e1003538.
Benson NC, Kupers ER, Barbot A, Carrasco M, Winawer J (2021) Cortical magnification in
human visual cortex parallels task performance around the visual field. Elife 10.
Benson NC, Jamison KW, Arcaro MJ, Vu AT, Glasser MF, Coalson TS, Van Essen DC,
Yacoub E, Ugurbil K, Winawer J, Kay K (2018) The Human Connectome Project 7
Tesla retinotopy dataset: Description and population receptive field analysis. J Vis
18:23.
Britten KH, Shadlen MN, Newsome WT, Movshon JA (1993) Responses of neurons in
macaque MT to stochastic motion signals. Vis Neurosci 10:1157-1169.

Chang C, Cunningham JP, Glover GH (2009) Influence of heart rate on the BOLD signal: the
cardiac response function. Neuroimage 44:857-869.
Cherukara MT, Stone AJ, Chappell MA, Blockley NP (2019) Model-based Bayesian
inference of brain oxygenation using quantitative BOLD. Neuroimage 202:116106.
Dale AM, Fischl B, Sereno MI (1999) Cortical Surface-Based Analysis: I. Segmentation and
Surface Reconstruction. NeuroImage 9:179-194.
Di X, Kannurpatti SS, Rypma B, Biswal BB (2013) Calibrating BOLD fMRI activations with
neurovascular and anatomical constraints. Cereb Cortex 23:255-263.
Dumoulin SO (2017) Layers of Neuroscience. Neuron 96:1205-1206.
Dumoulin SO, Wandell BA (2008) Population receptive field estimates in human visual
cortex. Neuroimage 39:647-660.
Dumoulin SO, Fracasso A, van der Zwaag W, Siero JCW, Petridou N (2018) Ultra-high field
MRI: Advancing systems neuroscience towards mesoscopic human brain function.
Neuroimage 168:345-357.
Duvernoy HM, Delon S, Vannson JL (1981) Cortical blood vessels of the human brain. Brain
Res Bull 7:519-579.
Fischl B, Dale AM (2000) Measuring the thickness of the human cerebral cortex from
magnetic resonance images. Proc Natl Acad Sci U S A 97:11050-11055.
Fracasso A, Petridou N, Dumoulin SO (2016a) Systematic variation of population receptive
field properties across cortical depth in human visual cortex. Neuroimage 139:427-
438.
Fracasso A, Luijten PR, Dumoulin SO, Petridou N (2018) Laminar imaging of positive and
negative BOLD in human visual cortex at 7T. Neuroimage 164:100-111.
Fracasso A, van Veluw SJ, Visser F, Luijten PR, Spliet W, Zwanenburg JJM, Dumoulin SO,
Petridou N (2016b) Lines of Baillarger in vivo and ex vivo: Myelin contrast across
lamina at 7T MRI and histology. Neuroimage 133:163-175.
Gagnon L, Sakadzic S, Lesage F, Pouliot P, Dale AM, Devor A, Buxton RB, Boas DA (2016)
Validation and optimization of hypercapnic-calibrated fMRI from oxygen-sensitive
two-photon microscopy. Philos Trans R Soc Lond B Biol Sci 371.
Gagnon L, Sakadzic S, Lesage F, Musacchia JJ, Lefebvre J, Fang Q, Yucel MA, Evans KC,
Mandeville ET, Cohen-Adad J, Polimeni JR, Yaseen MA, Lo EH, Greve DN, Buxton
RB, Dale AM, Devor A, Boas DA (2015a) Quantifying the microvascular origin of
BOLD-fMRI from first principles with two-photon microscopy and an oxygen-sensitive
nanoprobe. J Neurosci 35:3663-3675.
Gagnon L, Sakadžić S, Lesage F, Musacchia JJ, Lefebvre J, Fang Q, Yücel MA, Evans KC,
Mandeville ET, Cohen-Adad J, Polimeni JaR, Yaseen MA, Lo EH, Greve DN, Buxton
RB, Dale AM, Devor A, Boas DA (2015b) Quantifying the Microvascular Origin of
BOLD-fMRI from First Principles with Two-Photon Microscopy and an Oxygen-
Sensitive Nanoprobe. The Journal of Neuroscience 35:3663.
Gorno-Tempini ML, Hutton C, Josephs O, Deichmann R, Price C, Turner R (2002) Echo time
dependence of BOLD contrast and susceptibility artifacts. Neuroimage 15:136-142.
Guidi M, Huber L, Lampe L, Merola A, Ihle K, Moller HE (2020) Cortical laminar resting-state
signal fluctuations scale with the hypercapnic blood oxygenation level-dependent
response. Hum Brain Mapp 41:2014-2027.
He X, Yablonskiy DA (2007) Quantitative BOLD: mapping of human cerebral deoxygenated
blood volume and oxygen extraction fraction: default state. Magn Reson Med 57:115-
126.
Heeger DJ, Ress D (2002) What does fMRI tell us about neuronal activity? Nature Reviews
Neuroscience 3:142-151.
Heeger DJ, Huk AC, Geisler WS, Albrecht DG (2000) Spikes versus BOLD: what does
neuroimaging tell us about neuronal activity? Nat Neurosci 3:631-633.

Herman MC, Cardoso MMB, Lima B, Sirotin YB, Das A (2017) Simultaneously estimating the
task-related and stimulus-evoked components of hemodynamic imaging
measurements. Neurophotonics 4:031223.
Hilgetag CC, Barbas H (2005) Developmental mechanics of the primate cerebral cortex.
Anat Embryol (Berl) 210:411-417.
Himmelberg MM, Winawer J, Carrasco M (2022) Linking individual differences in human
primary visual cortex to contrast sensitivity around the visual field. Nature
Communications 13:3309.
Himmelberg MM, Kurzawski JW, Benson NC, Pelli DG, Carrasco M, Winawer J (2021)
Cross-dataset reproducibility of human retinotopic maps. Neuroimage 244:118609.
Hoogenraad FG, Hofman MB, Pouwels PJ, Reichenbach JR, Rombouts SA, Haacke EM
(1999) Sub-millimeter fMRI at 1.5 Tesla: correlation of high resolution with low
resolution measurements. J Magn Reson Imaging 9:475-482.
Jamison K, Vizioli L, Zhang R, Tao J, Winawer J, Kay K (2017) A tool for automatic
identification of cerebral sinuses and corresponding artifacts in fMRI. Journal of
Vision 17:295-295.
Jiang X, Zhang T, Zhang S, Kendrick KM, Liu T (2021) Fundamental functional differences
between gyri and sulci: implications for brain function, cognition, and behavior.
Psychoradiology 1:23-41.
Kay K, Jamison KW, Zhang RY, Ugurbil K (2020) A temporal decomposition method for
identifying venous effects in task-based fMRI. Nat Methods 17:1033-1039.
Kay K, Jamison KW, Vizioli L, Zhang R, Margalit E, Ugurbil K (2019) A critical assessment of
data quality and venous effects in sub-millimeter fMRI. Neuroimage 189:847-869.
Kay KN, Winawer J, Mezer A, Wandell BA (2013a) Compressive spatial summation in
human visual cortex. J Neurophysiol 110:481-494.
Kay KN, Rokem A, Winawer J, Dougherty RF, Wandell BA (2013b) GLMdenoise: a fast,
automated technique for denoising task-based fMRI data. Front Neurosci 7:247.
Kazan SM, Mohammadi S, Callaghan MF, Flandin G, Huber L, Leech R, Kennerley A,
Windischberger C, Weiskopf N (2016) Vascular autorescaling of fMRI (VasA fMRI)
improves sensitivity of population studies: A pilot study. Neuroimage 124:794-805.
Kim SG, Hendrich K, Hu X, Merkle H, Ugurbil K (1994) Potential pitfalls of functional MRI
using conventional gradient-recalled echo techniques. NMR Biomed 7:69-74.
Koopmans PJ, Barth M, Orzada S, Norris DG (2011) Multi-echo fMRI of the cortical laminae
in humans at 7 T. Neuroimage 56:1276-1285.
Lee JH, Durand R, Gradinaru V, Zhang F, Goshen I, Kim DS, Fenno LE, Ramakrishnan C,
Deisseroth K (2010) Global and local fMRI signals driven by neurons defined
optogenetically by type and wiring. Nature 465:788-792.
Liu T, Heeger DJ, Carrasco M (2006) Neural correlates of the visual vertical meridian
asymmetry. J Vis 6:1294-1306.
Liu TT, Behzadi Y, Restom K, Uludag K, Lu K, Buracas GT, Dubowitz DJ, Buxton RB (2004)
Caffeine alters the temporal dynamics of the visual BOLD response. Neuroimage
23:1402-1413.
Logothetis NK (2002) The neural basis of the blood-oxygen-level-dependent functional
magnetic resonance imaging signal. Philos Trans R Soc Lond B Biol Sci 357:1003-
1037.
Logothetis NK (2008) What we can do and what we cannot do with fMRI. Nature 453:869-
878.
Logothetis NK, Wandell BA (2004) Interpreting the BOLD signal. Annu Rev Physiol 66:735-
769.
Logothetis NK, Pauls J, Augath M, Trinath T, Oeltermann A (2001) Neurophysiological
investigation of the basis of the fMRI signal. Nature 412:150-157.

Menon RS, Ogawa S, Tank DW, Ugurbil K (1993) Tesla gradient recalled echo
characteristics of photic stimulation-induced signal changes in the human primary
visual cortex. Magn Reson Med 30:380-386.
Mishra A, Hall CN, Howarth C, Freeman RD (2021) Key relationships between non-invasive
functional neuroimaging and the underlying neuronal activity. Philosophical
Transactions of the Royal Society B: Biological Sciences 376:20190622.
Polimeni JR, Fischl B, Greve DN, Wald LL (2010) Laminar analysis of 7T BOLD using an
imposed spatial activation pattern in human V1. Neuroimage 52:1334-1346.
Rees G, Friston K, Koch C (2000) A direct quantitative relationship between the functional
properties of human and macaque V5. Nature Neuroscience 3:716-723.
Self MW, van Kerkoerle T, Goebel R, Roelfsema PR (2019) Benchmarking laminar fMRI:
Neuronal spiking and synaptic activity during top-down and bottom-up processing in
the different layers of cortex. Neuroimage 197:806-817.
Siero JC, Hermes D, Hoogduin H, Luijten PR, Ramsey NF, Petridou N (2014) BOLD
matches neuronal activity at the mm scale: a combined 7T fMRI and ECoG study in
human sensorimotor cortex. Neuroimage 101:177-184.
Silva MA, See AP, Essayed WI, Golby AJ, Tie Y (2018a) Challenges and techniques for
presurgical brain mapping with functional MRI. Neuroimage Clin 17:794-803.
Silva MF, Brascamp JW, Ferreira S, Castelo-Branco M, Dumoulin SO, Harvey BM (2018b)
Radial asymmetries in population receptive field size and cortical magnification factor
in early visual cortex. Neuroimage 167:41-52.
Sirotin YB, Das A (2009) Anticipatory haemodynamic signals in sensory cortex not predicted
by local neuronal activity. Nature 457:475-479.
Srirangarajan T, Mortazavi L, Bortolini T, Moll J, Knutson B (2021) Multi-band FMRI
compromises detection of mesolimbic reward responses. NeuroImage 244:118617.
Tsvetanov KA, Henson RNA, Rowe JB (2021) Separating vascular and neuronal effects of
age on fMRI BOLD signals. Philos Trans R Soc Lond B Biol Sci 376:20190631.
Turner R (2002) How much cortex can a vein drain? Downstream dilution of activation-
related cerebral blood oxygenation changes. Neuroimage 16:1062-1067.
van der Zwaag W, Francis S, Head K, Peters A, Gowland P, Morris P, Bowtell R (2009) fMRI
at 1.5, 3 and 7 T: characterising BOLD signal changes. Neuroimage 47:1425-1434.
van Dijk JA, Fracasso A, Petridou N, Dumoulin SO (2020) Linear systems analysis for
laminar fMRI: Evaluating BOLD amplitude scaling for luminance contrast
manipulations. Sci Rep 10:5462.
Viessmann O, Scheffler K, Bianciardi M, Wald LL, Polimeni JR (2019) Dependence of
resting-state fMRI fluctuation amplitudes on cerebral cortical orientation relative to
the direction of B0 and anatomical axes. Neuroimage 196:337-350.
Welker W (1990) Why Does Cerebral Cortex Fissure and Fold? In: Cerebral Cortex:
Comparative Structure and Evolution of Cerebral Cortex, Part II (Jones EG, Peters A,
eds), pp 3-136. Boston, MA: Springer US.
Winawer J, Horiguchi H, Sayres RA, Amano K, Wandell BA (2010) Mapping hV4 and ventral
occipital cortex: the venous eclipse. J Vis 10:1.
Yablonskiy DA, Sukstanskii AL, He X (2013) Blood oxygenation level-dependent (BOLD)-
based techniques for the quantification of brain hemodynamic and metabolic
properties - theoretical models and experimental approaches. NMR Biomed 26:963-
986.
Yu X, Qian C, Chen DY, Dodd SJ, Koretsky AP (2014) Deciphering laminar-specific neural
inputs with line-scanning fMRI. Nat Methods 11:55-58.
Zimmermann J, Goebel R, De Martino F, van de Moortele PF, Feinberg D, Adriany G,
Chaimow D, Shmuel A, Ugurbil K, Yacoub E (2011) Mapping the organization of axis
of motion selective features in human area MT using high-field fMRI. PLoS One
6:e28716.
868