Effector-mediated subversion of proteasome activator (PA)28αβ enhances lysoso-

mal targeting of *Legionella pneumophila* within cytokine-activated macrophages

Tshegofatso Ngwaga[1], Deepika Chauhan[1], Abigail G. Salberg[1], and Stephanie R. Shames[1]*

Division of Biology, Kansas State University, Manhattan, Kansas, 66506

*Correspondence: sshames@ksu.edu; @shames_lab

Phone: (785) 532-0110

Address: 347 Ackert Hall, 1717 Claflin Road, Manhattan, Kansas, 66506 USA

Keywords: PA28, proteasome, phagolysosomal fusion, cytokines, cell-autonomous immunity, *Le-*

gionella pneumophila, LegC4, proteostasis, oxidative stress

Abstract

Legionella pneumophila is a natural pathogen of protozoa that causes Legionnaires' Disease pneumonia via replication within host macrophages using an arsenal of hundreds of translocated virulence factors termed effector proteins. Effectors are critical for intracellular replication but can also enhance pathogen clearance by mammalian hosts via effector-triggered immunity. The effector LegC4 confers a fitness disadvantage on *L. pneumophila* in mouse models of Legionnaires' Disease and uniquely potentiates the antimicrobial activity of macrophages activated with either tumor necrosis factor (TNF) or interferon (IFN)-γ. Here, we investigated the mechanism by which LegC4 enhances macrophage restriction of *L. pneumophila* and found a central role for the induced proteasome activator (PA)28αβ. PA28αβ facilitates ubiquitin-independent proteasomal degradation of oxidant-damaged proteins to relieve oxidative stress and impaired proteasome activity results in compensatory upregulation of lysosomal degradation pathways. We found that LegC4 binds the conserved PA28α subunit and that the LegC4 restriction phenotype is abolished within PA28αβ-deficient macrophages. Moreover, LegC4 impaired the resolution of oxidative proteotoxic stress and enhanced cytokine-dependent phagolysosomal fusion with the *Legionella*-containing vacuole. PA28αβ has been traditionally associated with antigen presentation and adaptive immunity; however, our data support a model whereby suppression of PA28αβ by LegC4 culminates in the cell-autonomous lysosomal killing of *L. pneumophila* within activated macrophages. This work provides a solid foundation on which to evaluate induced proteasome regulators as mediators of innate immunity.

Significance Statement

Pro-inflammatory cytokines induce antimicrobial host defense pathways within macrophages to control intracellular pathogens. We discovered that the *Legionella pneumophila* effector protein LegC4 potentiates pathogen clearance within cytokine-activated macrophages. Here, we investigated LegC4 function and found that host proteasome activator (PA)28αβ was required for LegC4 restriction within macrophages. PA28αβ is cytokine-induced and facilitates proteasomal degradation of oxidant-damaged proteins. We found that LegC4 binds PA28α and impairs the resolution of oxidative stress. Loss of proteasome activity causes compensatory upregulation of lysosomal degradation pathways and, concomitantly, LegC4 increases phagolysosomal fusion with *Legionella*-containing vacuoles. These findings support a model whereby pathogen subversion of host proteostasis machinery triggers lysosomal pathogen targeting within activated macrophages.

Introduction

Many intracellular bacterial pathogens cause disease by replicating within host macrophages (1). Macrophages activated by T_H1 proinflammatory cytokines, such as tumor necrosis factor (TNF) and interferon (IFN)-γ, are potently and broadly microbicidal. However, activated macrophages are in a state of altered protein homeostasis (proteostasis), owing in part to robust production of reactive oxygen species (ROS), which cause collateral damage to cellular proteins. To cope with oxidative stress, cytokine signaling induces expression of proteasome activator (PA)28αβ, which engages proteolytic 20S proteasome core particles (CPs) and facilitates degradation of oxidant-damaged proteins (2). Impaired proteasome activity causes compensatory upregulation of lyso-somal degradation pathways, which are central to cell-autonomous immunity within activated macrophages (3). PA28αβ has been traditionally associated with antigen presentation and adap-tive immunity (4, 5); however, the contribution of PA28αβ and oxidative proteotoxic stress to antimicrobial host defense has not been investigated.

Legionella pneumophila is a natural pathogen of freshwater amoebae and the causative agent of Legionnaires' Disease pneumonia, which results from bacterial replication within alveolar mac-rophages. To establish its intracellular niche, the *Legionella*-containing vacuole (LCV), *L. pneu-mophila* employs hundreds of virulence factors, called effector proteins, that are translocated into infected host cells by a Dot/Icm type IV secretion system (T4SS) (6). Effectors are essential for virulence, but their activity can also trigger upregulation of pro-inflammatory cytokine production by infected macrophages via effector-triggered immunity (ETI) (7–9). We discovered that the *L. pneumophila* effector LegC4, which is a *bona fide* virulence factor in amoebae, confers a fitness disadvantage on *L. pneumophila* in a mouse model of Legionnaires' Disease (10). The *in vivo* LegC4 restriction phenotype is likely due to elevated levels of pro-inflammatory cytokines in the *L. pneumophila*-infected lung since LegC4 impairs *L. pneumophila* replication within cultured macrophages activated with either tumor necrosis factor (TNF) or interferon (IFN)-γ (10, 11).

LegC4 is the only *L. pneumophila* effector that confers quantifiable opposing host-dependent fit-

ness phenotypes and the specific augmentation of cytokine-mediated restriction represents a po-

tentially novel mechanism of ETI. However, the mechanism by which LegC4 functions and en-

hances cytokine-mediated restriction of *L. pneumophila* is unknown.

Here, we investigated how LegC4 activity suppresses *L. pneumophila* replication within activated

macrophages. We found that LegC4 interacts with host PA28α and that LegC4-mediated re-

striction was abolished within PA28αβ-deficient (*Psme1/2$^{-/-}$*) bone marrow-derived macro-

phages. We further revealed that LegC4 impairs resolution of oxidative proteotoxic stress, which

is sufficient to induce the LegC4 restriction phenotype, and that LegC4 induces phagolysosomal

fusion with intracellular *Legionella*-containing vacuoles (LCVs). In sum, these results support a

model whereby PA28αβ targeting by a translocated effector protein perturbs proteostasis and en-

hances lysosomal targeting and restriction of *L. pneumophila* within cytokine-activated macro-

phages.

Results

LegC4 potentiates redundant cytokine-mediated restriction of *L. pneumophila* We

previously reported that LegC4 confers a fitness disadvantage on *L. pneumophila* in intranasal

mouse models of Legionnaires' Disease (LD) and within cytokine-activated macrophages (10, 11).

C57Bl/6 mice are restrictive to *L. pneumophila* due to flagellin-mediated activation of the

NAIP5/NLRC4 inflammasome (12, 13). Thus, we leveraged a flagellin (FlaA)-deficient *L. pneu-*

mophila (Δ*flaA*), an established model used to study FlaA-independent host responses within

C57Bl/6 mouse models of infection (12, 13). Using competitive index experiments, we found that

L. pneumophila Δ*flaA*Δ*legC4* mutant strains outcompeted parental Δ*flaA* strains in the lungs of

C57Bl/6 wild-type (WT) mice (11). However, no LegC4 fitness phenotype was observed within

isogenic WT bone marrow-derived macrophages (BMDMs) (11). *L. pneumophila* replicates within

alveolar macrophages in the lung, but we did not observe any LegC4-dependent differences in *L.*

pneumophila replication within MH-S mouse alveolar macrophages (**Fig S1**). However, LegC4-

mediated restriction is observed within BMDMs activated with either purified recombinant

(r)TNF or rIFN-γ (11). Thus, the phenotype observed for LegC4 *in vivo* is likely due to enhanced

levels of cytokines seen in the *L. pneumophila*-infected lung compared to cultured macrophages

(14). Alternatively, a phenotype for LegC4 can be observed in non-cytokine activated (naïve)

BMDMs by overexpressing *legC4* from an inducible promoter on a complementing plasmid (**Fig**

1A) (11). The LegC4 restriction phenotype is not exclusive to mouse macrophages since LegC4

confers a comparable fitness disadvantage on *L. pneumophila* within human THP-1 macrophages

(**Fig 1B**). Further support and utility of the LegC4 overexpression model was evidenced by loss of

LegC4-mediated restriction within BMDMs unable to produce or respond to endogenous TNF

(*Myd88*$^{-/-}$ and *Tnfr1*$^{-/-}$, respectively) (**Fig 1C, 1D**) (11). IFN-γ signaling is sufficient for LegC4 re-

striction (11); however, the LegC4 restriction phenotype was retained within *Ifngr1*$^{-/-}$ BMDMs

(**Fig 1E**), indicating that the relatively low levels of IFN-γ secreted by macrophages are not nec-

essary for restriction. Furthermore, the loss of any single cytokine was not sufficient to restore *L.*

pneumophila fitness since LegC4 still conferred a fitness disadvantage on *L. pneumophila* within

the lungs of TNFR1-, MyD88- and IFNGR1-deficient mice (**Fig 1F**). These data further support

our previous work showing that LegC4 augments redundant cytokine-mediated pathogen re-

striction mechanisms (11). While overexpression of *legC4* is not necessary to observe a phenotype,

this infection model is robust and useful for our goal to define LegC4 function and mechanisms

of cytokine-mediated restriction within macrophages.

LegC4 binds host proteasome activator (PA)28α. To gain insight into the mechanism of

LegC4 function, we conducted a yeast two-hybrid (Y2H) interactomics screen using LegC4 as bait

and genomic (g)DNA from mouse splenocytes as prey. This screen revealed interactions between

LegC4 and several regulators of eukaryotic proteasomes, including proteasome activator (PA)

28α, PA28γ, and Ecm29 (**Table S1**). The highest-confidence interactor was PA28α with 28 individual clones, all of which encoded a minimal overlapping region in the PA28α C-terminal domain (amino acid residues 105-249) (**Table S1**, **Fig S2**). We initially evaluated the PA28α-LegC4 interaction by reciprocal co-immunoprecipitation from lysates of HEK 293T cells ectopically producing epitope-tagged PA28α and LegC4. We found that PA28α-Myc co-immunoprecipitated with 3xFLAG-LegC4 and that 3xFLAG-LegC4 co-immunoprecipitated with GFP-PA28α (**Fig 2A**). We also found that ectopically produced GFP-PA28α and 3xFLAG-LegC4 localize within transfected HeLa cells (**Fig 2B**). Ectopic expression of PA28α was necessary since it is normally expressed only in response to cytokine signaling (15, 16). Together, these data suggest that LegC4 binds host PA28α, which has not been previously identified as a target of bacterial effectors and provides an explanation as to why LegC4 only has a phenotype within cytokine-activated cells.

PA28αβ contributes to LegC4-dependent and -independent macrophage restriction of _L. pneumophila_. Based on the interaction between PA28α and LegC4, we tested the hypothesis that PA28α contributes to LegC4-mediated restriction of _L. pneumophila_ within BMDMs. PA28α-deficient (_Psme1-/-_) mice are not commercially available, so we evaluated the role of PA28α using BMDMs derived from PA28αβ-deficient (_Psme1/2-/-_) mice. PA28α associates with PA28β to form the PA28αβ (11S) proteasome regulator. PA28β is homologous to PA28α, but is strictly dependent on PA28α for activity and is hypothesized to enhance the interaction of PA28α with 20S proteolytic proteasome core particles (CPs) (17). We found that PA28αβ was necessary for LegC4-mediated restriction of _L. pneumophila_ since it was abolished within _Psme1/2-/-_ BMDMs (**Fig 2C**). Moreover, replication of _L. pneumophila_ Δ_flaA_ was attenuated within naïve and cytokine-activated _Psme1/2-/-_ BMDMs (**Fig 2D**), which phenocopies LegC4 and supports the hypothesis that LegC4 impairs PA28α activity. Since autocrine and paracrine TNF signaling is sufficient for the LegC4 restriction phenotype, we tested whether PA28αβ or LegC4 enhance TNF

secretion from *L. pneumophila*-infected BMDMs. However, we found no LegC4- or PA28αβ-me-

diated differences in TNF secretion from infected BMDMs (**Fig S3**). We also found that viability

of *L. pneumophila*-infected BMDMs was unaffected by either LegC4, PA28αβ, or exogenous rTNF

compared to untreated cells (**Fig S4**). These data suggest that the LegC4 restriction phenotype is

dependent on PA28αβ and occurs independently of increased cytokine secretion or cell death,

which support a novel role for PA28αβ in cell-autonomous immunity against an intracellular bac-

terial pathogen.

LegC4 impairs resolution of oxidative proteotoxic stress. In cells of the innate immune

system, PA28α is produced in response to oxidative stress, TNF, or IFN-γ and forms a heterohep-

tameric complex with PA28β to assemble the 11S (PA28αβ) proteasome regulator (2, 16, 18, 19).

PA28αβ associates with proteasome CPs to generate antigenic peptides and resolve oxidative pro-

teotoxic stress by facilitating ubiquitin-and ATP-independent proteasomal degradation of oxi-

dant-damaged (carbonylated) proteins (4, 20). Since PA28αβ was required for LegC4-mediated

L. pneumophila restriction within BMDMs (see Fig 2C), we tested the hypothesis that LegC4 im-

pairs resolution of oxidative proteotoxic stress by quantifying protein carbonyls in cell lysates.

We directly evaluated how LegC4 impacts resolution of oxidative proteotoxic stress by

quantifying protein carbonyls within RAW 264.7 cells stably producing 3xFLAG-LegC4 (RAW-

LegC4). We confirmed production of 3xFLAG-LegC4 by confocal microscopy (**Fig 3A**) and West-

ern blot (**Fig 3B**). Immunoprecipitation of 3xFLAG-LegC4 was necessary due to its low abun-

dance. However, we validated that ectopically produced LegC4 was active, since replication of *L.*

pneumophila Δ*flaA* within RAW-LegC4 cells was attenuated compared to control (RAW-Cntrl)

cells (**Fig 3C**). To test the hypothesis that LegC4 impairs resolution of oxidative proteotoxic

stress, we quantified protein carbonyls using ELISA within lysates of RAW-LegC4 and -Cntrl cells

treated with hydrogen peroxide (H_2O_2). H_2O_2 is commonly used to induce oxidative stress since

superoxide radical ($O_2^{\cdot-}$) (precursor to H_2O_2) is unstable in solution (21–23). We found that the

concentration of protein carbonyls in lysates of H_2O_2-treated RAW-LegC4 cells was significantly

greater than control cells (**Fig 3D**) and that LegC4 alone did not affect the concentration of pro-

tein carbonyls in untreated cells (**Fig 3D**). These data suggest that LegC4 is sufficient to impair

resolution of oxidative proteotoxic stress.

We subsequently evaluated how LegC4 impacts the abundance of protein carbonyls within

L. pneumophila-infected cells. To control for LegC4-mediated differences in bacterial replication

within WT BMDMs, we leveraged *Tnfr1*[-/-] BMDMs (see Fig 1). *Tnfr1*[-/-] BMDMs were infected for

4 h in the presence or absence of H_2O_2 and protein carbonyls were visualized by Western blot. We

found that the abundance of protein carbonyls was increased within H_2O_2-treated BMDMs in-

fected with *L. pneumophila* Δ*flaA*Δ*legC4* (p*legC4*) relative to Δ*flaA* and empty vector control

strains (**Fig 3E**). Within untreated BMDMs, infection with the avirulent *dotA*::Tn strain increased

the abundance of protein carbonyls (**Fig 3E**), which is likely a consequence of increased global

ROS production by macrophages infected with avirulent compared to virulent *L. pneumophila*

(24). Since no LegC4-mediated differences were observed within untreated *Tnfr1*[-/-], it is unlikely

that the increase in protein carbonyls observed is due to a LegC4-mediated upregulation global

ROS production by infected BMDMs. Together, these data support our hypothesis that LegC4

impairs resolution of oxidative proteotoxic stress within macrophages.

Oxidative stress is sufficient for LegC4-mediated restriction. Based on our data showing

that LegC4 impairs resolution of oxidative proteotoxic stress, we hypothesized that oxidative

stress is sufficient for LegC4-mediated restriction within BMDMs. To test this hypothesis, we

quantified *L. pneumophila* replication within *Tnfr1*[-/-] BMDMs, in which LegC4 is not restrictive

(see Fig 1C), in the presence or absence of a sublethal concentration of H_2O_2 to induce oxidative

stress. We found that replication of *L. pneumophila* Δ*flaA*Δ*legC4* (p*legC4*) strain was significantly

attenuated within H_2O_2-treated *Tnfr1*[-/-] BMDMs compared to *L. pneumophila* Δ*flaA* and

$\Delta flaA\Delta legC4$ empty vector (pEV) control strains (**Fig 3F**). These differences represent a 78% de-

crease in colony forming units (CFU), which is striking since experimental limitations necessi-

tated a low concentration (10 μM) of H_2O_2. It is unlikely that growth attenuation was due to direct

L. pneumophila killing by H_2O_2 since *L. pneumophila* has evolved multiple complimentary strat-

egies to protect against oxidative stress and can withstand up to 2 mM H_2O_2 *in vitro* (orders of

magnitude greater than intracellular concentrations) (22, 24–26). Thus, these data suggest that

oxidative stress is sufficient for the LegC4 restriction phenotype.

LegC4 increases phagolysosomal fusion with *Legionella*-containing vacuoles. Our

current data show that LegC4 functions through PA28αβ and confers a fitness disadvantage under

oxidative stress conditions. Impaired proteasome activity causes compensatory upregulation of

lysosomal degradation pathways to cope with proteotoxic stress (3, 27). Based on our data show-

ing that oxidative proteotoxic stress is sustained in LegC4-producing cells, we tested the hypoth-

esis that LegC4 increases lysosomal targeting of *L. pneumophila* within activated macrophages.

Virulent *L. pneumophila* establish a replicative *Legionella*-containing vacuole (LCV) by blocking

endocytic maturation of their phagosome (28, 29). However, avirulent (*dotA*::Tn) *L. pneumophila*

are unable to establish a replicative LCV and phagosomes rapidly undergo endocytic maturation,

as evidenced by localization of the lysosomal membrane marker LAMP1 to *L. pneumophila*-con-

taining phagosomes (28) (**Fig S5**). We initially tested the hypothesis that LegC4 increases LAMP1

localization to LCVs by immunofluorescence microscopy and blinded quantification of LAMP1[+]

LCVs. As expected, there were significantly fewer LAMP1[+] LCVs harboring $\Delta flaA$ and empty vec-

tor control strains compared to the avirulent control (**Fig 4A**). There was a significant increase in

the percentage of LAMP1[+] LCVs harboring $\Delta flaA\Delta legC4$ (p*legC4*) compared to $\Delta flaA$ and empty

vector control strains within both WT and *Tnfr1*[-/-] BMDMs (**Fig 4A**). However, within WT

BMDMs, LAMP1 localization to LCVs harboring $\Delta flaA\Delta legC4$ (p*legC4*) did not differ significantly

from the avirulent *dotA*::Tn strain (**Fig 4A**), which was surprising since plasmid expression of

legC4 did not impair biogenesis of a replicative LCV (≥5 bacteria/vacuole) compared to *L. pneu-*

mophila ΔflaA and *ΔflaAΔlegC4* (pEV) control strains (**Fig 4B**). As expected, the avirulent con-

trol strain was significantly attenuated for replicative LCV formation compared to the parental *L.*

pneumophila ΔflaA strain (**P<0.01, **Fig 4B**). These data correlate with our CFU data showing

that that LegC4 restriction is observed after 24 h of infection within BMDMs (Fig 1A) and chal-

lenge the dogma that LAMP1 localization and *L. pneumophila* intracellular replication are mutu-

ally exclusive. Moreover, these data support our hypothesis that LegC4 enhances phagolysosomal

fusion with LCVs.

We subsequently tested the hypothesis that phagolysosomal fusion is important for

LegC4-mediated restriction by quantifying *L. pneumophila* replication within BMDMs treated

with Bafilomycin A1 (BAF). BAF is a pharmacological inhibitor of the H^+-type vacuolar (v)-ATPase

that blocks vacuolar acidification and impairs phagolysosomal fusion (30). Thus, we evaluated

the contribution of phagolysosomal fusion to the LegC4 restriction phenotype by quantifying *L.*

pneumophila replication within BAF-treated BMDMs. LCV acidification is important for late

stages of the *L. pneumophila* lifecycle (31); however, intracellular replication is minimally im-

pacted within BMDMs treated with a low concentration (12.5 nM) of BAF (32). We found that the

LegC4 restriction phenotype was abolished within BAF-treated BMDMs (**Fig 4C**), suggesting that

phagolysosomal fusion and lysosomal acidification are important for LegC4-mediated restriction.

Together, our data support a central role for phagolysosomal fusion and vacuolar acidification in

LegC4-mediated restriction of *L. pneumophila* within BMDMs.

Discussion

This study supports a novel role for proteasome activator (PA)28αβ in cell-autonomous host de-

fense against an intracellular pathogen. We found that the *L. pneumophila* effector LegC4 en-

hances restriction of *L. pneumophila* within macrophages activated with either TNF or IFN-γ in

a PA28αβ-dependent manner. Our results suggest a model whereby LegC4 impairs resolution of

oxidative proteotoxic stress and promotes lysosomal restriction of *L. pneumophila* by subverting

PA28αβ activity (**Fig S6**). This model is supported by our data showing that LegC4 binds PA28α,

PA28αβ is required for the effects of LegC4 on host cells, LegC4 impairs resolution of oxidative

proteotoxic stress, and the contribution of both oxidative stress and phagolysosomal fusion to

LegC4-mediated restriction of *L. pneumophila*. This work is the first describe a role for PA28αβ

in innate and cell-autonomous host defense.

L. pneumophila is an accidental pathogen of humans that rarely transmits between infected indi-

viduals. Thus, an evolutionary basis for the LegC4-PA28α interaction can be explained by LegC4

targeting the PA28α homolog in amoebae. Indeed, the natural host *A. castellanii*, in which LegC4

is a *bona fide* virulence factor (10), encodes a PA28α homolog (*Ac*PA28) that shares 34% identity

with PA28α. Sequence identity between PA28α and *Ac*PA28 is primarily in the C-terminal do-

main, which is important for engaging 20S CPs and was contained in all PA28α clones identified

in our yeast two-hybrid screen (Fig 2A; Table S1). These data together with our observation that

the LegC4 phenotype is recapitulated in PA28αβ-deficient macrophages have led us to speculate

that LegC4 binding prevents PA28-CP complex formation. While the function of *Ac*PA28 has not

been described, structural studies have shown that mammalian PA28α and evolutionary distant

PA28s are similar and all associate with proteolytic proteasome core particles, suggesting that it

likely also functions as a proteasome regulator (33, 34). The constitutive 26S proteasome system

is central to *L. pneumophila*'s virulence strategy and its integrity may be preserved by sequester-

ing PA28 from CPs. We are currently investigating the molecular mechanism by which LegC4

regulates PA28 activity within mammalian cells and amoebae and how this activity is advanta-

geous to *L. pneumophila* in the natural host.

We found that inducing oxidative stress with hydrogen peroxide (H_2O_2) was sufficient for the

LegC4 restriction phenotype. H_2O_2 is routinely used to induce oxidative stress in laboratory mod-

els since both superoxide ($O_2^{\cdot-}$) and hydroxyl ($OH^\cdot$) radicals are highly reactive and unstable in

solution. Interestingly, reactive oxygen species (ROS) are important for *L. pneumophila* re-

striction by neutrophils but play a negligible role in macrophages (24, 35, 36). This difference has

been attributed to suppression of global intracellular ROS production and NADPH oxidase

(NOX2) trafficking to the LCV by virulent *L. pneumophila* within macrophages (24, 35). Direct

oxidative killing of *L. pneumophila* within activated macrophages is unlikely since *L. pneumoph-

ila* has evolved multiple complimentary strategies to protect against oxidative stress and can with-

stand up to 2 mM H_2O_2 *in vitro* (orders of magnitude greater than intracellular concentrations)

(22, 24–26). Moreover, *Tnfr1*$^{-/-}$ BMDMs treated with H_2O_2 did not suppress replication of *L.

pneumophila* control strains compared to untreated BMDMs (see Fig 3F). This suggests that the

concentration of H_2O_2 used in our study is insufficient for direct *L. pneumophila* killing. Future

studies will reveal whether LegC4 or PA28$\alpha\beta$ impact the abundance or source of ROS within *L.

pneumophila*-infected macrophages.

Phagolysosomal fusion and autophagy are cytoprotective under inflammatory conditions and

central to cytokine-mediated cell-autonomous pathogen restriction. Lysosomal pathogen re-

striction within IFN-γ-activated macrophages involves immunity-related GTPases (IRGs), which

localize to and facilitate lysosomal trafficking to pathogen-containing phagosomes (37). Although

IRGs may contribute to restriction of LegC4-producing *L. pneumophila* strains within IFN-γ-ac-

tivated macrophages (38), we found that IFNGR1 signaling is dispensable for LegC4 restriction

(Fig 1E), which suggests that IRGs are not central drivers of LegC4-mediated *L. pneumophila*

restriction. A role for TNFR1-mediated signaling in lysosomal pathogen restriction has not been

well established. Canonically, TNF signaling activates extrinsic cell death pathways; however,

TNFR1-mediated signaling induces lysosomal fusion with LCVs within 3 h of *L. pneumophila*

infection within primary mouse BMDMs, which is more rapid compared to BMDMs activated with

IFN-γ (36). The mechanism by which TNF signaling enhances phagolysosomal fusion with LCVs

is poorly understood; however, upregulation of autophagy genes is cytoprotective against TNF-

induced death in myeloid cells (39). Autophagy is also upregulated to prevent aberrant pyroptotic

cell death below a certain threshold of inflammasome agonist (40). The NLRC4 inflammasome is

dispensable for LegC4-mediated restriction (10, 11), but a role for the NLRP3 inflammasome is

possible since it is activated by proteotoxic stress (41). Inflammasome components and pro-in-

flammatory cytokines are selectively degraded below a threshold of agonist. We are currently eval-

uating the role of autophagy and the NLRP3 inflammasome in PA28αβ- and LegC4-mediated re-

striction of *L. pneumophila.*

Virulent *L. pneumophila* establish a replicative LCV by avoiding endosomal maturation of their

phagosome. The current dogma dictates that lysosomal fusion with LCVs at early time points dur-

ing infection (≤ 16 h) prevents *L. pneumophila* intracellular replication (31, 42). We found that

plasmid expression of *legC4* induced robust LAMP1 localization to LCVs within WT BMDMs, sim-

ilar to what is seen for an avirulent a Dot/Icm-deficient strain. Since we observed no attenuation

in replicative LCV formation or intracellular replication by CFU assay by 24 h post-infection, these

data suggest that *L. pneumophila* intracellular replication and LAMP1 localization to LCVs are

not mutually exclusive. It is tempting to speculate that *L. pneumophila* replication within a

LAMP1+ LCV may result from translocated effectors that block v-ATPase activity, such as SidK

(43–46). Interestingly, loss of TNFR1-mediated signaling did not fully abrogate LegC4-mediated

LAMP1 localization to LCVs. We think this may be a consequence of sustained oxidative stress

downstream of pattern recognition receptors. However, phagolysosomal fusion with LCVs is not

sufficient for LegC4 to impair replication within Tnfr1-/- BMDMs, further supporting our obser-

vation that LegC4 restriction occurs in a dose-dependent manner. The properties of a replication

permissive LAMP1[+] LCV, the contribution of additional translocated effectors, and the contribu-

tion of canonical autophagic degradation pathways are currently being investigated in our lab.

In summary, we have uncovered a new role for PA28αβ and oxidative proteotoxic stress in cell-

autonomous pathogen restriction. This work is also the first to show bacterial pathogen targeting

of PA28α and suggests that suppression of PA28αβ activity may be a pathogen-triggered host

defense mechanism. Moreover, our data suggest that inhibition of PA28αβ may enhance innate

immunity against intracellular pathogens by potentiating existing inflammatory responses. Fu-

ture studies will provide further insight into the ubiquity and mechanism(s) of pathogen re-

striction induced by proteotoxic stress and suppression of PA28αβ activity.

Materials & Methods

<u>Bacterial Strains, Plasmids, and Culture conditions</u>

Legionella pneumophila strains used in this study (**Table S2**) were cultured on supplemented

charcoal N-(2-Acetamido)-2-aminoethanesulfonic acid (ACES)-buffered yeast extract (CYE) and

grown at 37˚C as described (47). Liquid cultures were grown overnight with shaking at 37˚C in

supplemented ACES-buffered yeast extract (AYE) medium, as described (11, 48). For plasmid

maintenance, CYE was supplemented with 10 μg mL[-1] chloramphenicol and *legC4* expression was

induced with 1 mM isopropyl-β-D-1-thiogalactopyranoside (IPTG) where indicated. Plasmids

used in this study are listed in **Table S3**. Oligonucleotide primers are listed in **Table S4.**

<u>Mice and bone marrow-derived macrophages</u>

All mice were on a C57BL/6 background and purchased from the Jackson Laboratories (Bar Har-

bor, Maine) and in-house colonies were maintained in specific pathogen-free conditions at Kansas

State University. All experiments involving animals were approved by the Kansas State University

Institutional Animal Care and Use Committee (Protocols 4022, 4501, and 4386) and performed

in compliance with the Animal Welfare Act and NIH guidelines.

Bone marrow was harvested from seven- to twelve-week-old mice as described (49).

BMDMs were generated by differentiation in RPMI supplemented with 20% heat-inactivated fetal

bovine serum (HI-FBS) (Biowest, Riverside, MO) and 15% L929 cell supernatant for 6 days prior

to seeding for infection.

<u>Competitive Index</u>

Six- to twelve-week-old sex- and age-matched mice were infected as previously described. Mixed

bacterial inoculums (1:1) containing a total of 5×10^6 bacteria were diluted and plated on selective

medium (10 μg mL^{-1} chloramphenicol for plasmid selection). At 48 h post-infection, mice were

euthanized, and lung homogenates were plated on selective media as described (11). CFU were

enumerated and used to calculate CI values [(CFUcm$^R_{48h}$/CFUwt$_{48h}$)/(CFUcm$^R_{IN}$)/CFUwt$_{IN}$)].

<u>Molecular Cloning</u>

Plasmids were generated for stable and transient ectopic production in mammalian cells. For pro-

duction of 3xFLAG-LegC4, *legC4* was amplified from *L. pneumophila* genomic (g)DNA using

LegC4BamHI-F/LegC4NotI-R primer pairs and cloned as a BamHI/NotI fragment into 3xFLAG

4/TO (50). To produce GFP-PA28α, *psme1* was amplified from pCMV-3Tag-4a::*psme1* (pur-

chased from Genscript, Piscataway, New Jersey) using Psme1Sal1-F/Psme1BamHI-R primer

pairs and cloned as a Sal1/BamH1 fragment into pEGFPC1 (Clontech). Plasmid DNA was trans-

fected into mammalian cells as described below.

<u>Cell Culture and Transfections</u>

HEK 293T, HeLa cells, RAW 264.7 cells, and MH-S cells (gifts from Dr. Craig Roy, Yale Univer-

sity) were maintained at 37˚C and 5% CO_2 in Dulbecco's Modified Eagle Medium (DMEM)

supplemented with 10% HI-FBS. THP-1 cells were maintained in RPMI supplemented with 10%

HIFBS and differentiated with 100 nM phorbal 12-myristate 13-acetate (PMA) for 4 days prior to

infection with *L. pneumophila* (see below). All cell lines were used between passage 4-20.

HEK 293T cells were transfected with purified plasmid DNA (**Table S2**) using calcium

phosphate as described (51). Lysates were generated from transfected cells after 48 h and used for

immunoprecipitation and Western blot analysis (see below). HeLa cells were seeded one day prior

to transfection with jetPRIME transfection reagent according to manufacturer's guidelines. Two

hours before transfection, HeLa cells were washed and incubated in low-serum media (DMEM

4% HIFBS. Media were replaced with DMEM 10% HIFBS four hours post-transfection and as-

sayed 24 h post-transfection.

To generate RAW 264.7 cells with stable plasmid integrations, cells were transfected with

pcDNA::*3xflag-legC4* or pcDNA::*3xflag* vector using a Nucleofector 2b electroporator (Lonza,

Basel, Switzerland) and cultured over 14 days with Zeocin selection for 14 days (200 -1000 μg mL^{-}

^{1}). Stable RAW 264.7 cells were maintained in culture in the presence of 200 μg mL^{-1} Zeocin. For

L. pneumophila infections, cells were seeded one day prior to infection in the absence of Zeocin.

<u>Immunoprecipitation</u>

Transiently transfected HEK 293T cells or RAW 264.7 cells were washed with phosphate-buffered

saline and lysed in ice-cold NP-40 buffer [1% non-iodet P40 (v/v), 20 mM Tris pH 7.5, 150 mM

NaCl, 10 mM $Na_4P_2O_7$, 50 mM NaF, complete protease inhibitor (Roche)]. Lysates were clarified

and added to magnetic Protein G-conjugated Dynabeads that had been pre-incubated with either

mouse α-FLAG M2 antibody (Sigma Aldrich) or rabbit α-GFP antibody (Abcam, ab6556) accord-

ing to manufacturer's instructions. Input samples were collected from cell lysates prior to incuba-

tion with beads. Beads and input samples were resuspended in 3x Laemmli sample buffer for

Western blot analysis.

Western blot

Boiled protein samples were separated by SDS-PAGE and transferred to polyvinylidene difluoride

(PVDF) membrane using a BioRad TransBlot semidry transfer apparatus. Membranes were incu-

bated in blocking buffer [5% non-fat milk powder dissolved in Tris-buffered saline-0.1% Tween-

20 (TBST)]. Primary antibodies [rabbit a-FLAG (Sigma-Aldrich, F1804), rabbit a-Myc (Cell Sig-

naling, 2278S), rabbit a-GFP (Abcam, ab6556)] were used at 1:1,000 in blocking buffer and de-

tected with horseradish peroxidase (HRP)-conjugated secondary antibodies (1:5,000; Ther-

moFisher). Membranes were washed in TBST followed by addition of enhanced chemilumines-

cent (ECL) reagent (GE Amersham) and visualization using an Azure c300 Darkroom Replacer.

Protein carbonyl assays

418   Protein carbonyls were quantified using the OxiSelect™ Protein Carbonyl ELISA Kit (Cell

Biolabs) following manufacturer's instructions. Briefly, RAW 264.7 cells were seeded at 5×10^5 in

a 6-well plate in DMEM 10% HIFBS. The next day, media were aspirated, and cells were incubated

in DMEM/2.5% HIFBS in the presence or absence of 10 μM H_2O_2. After 24 h, cells were washed

twice in PBS and lysed in ice-cold 500 μL of detergent-free lysis buffer [25 mM HEPES pH 7.4,

150 mM NaCl, 10 mM $MgCl_2$, 1 mM EDTA, 2% glycerol (v/v)] supplemented with complete pro-

tease inhibitor cocktail (Roche). Cell lysates were scraped into a pre-chilled microcentrifuge tube

and incubated with agitation for 30 min at 4°C followed by sonication at 40% intensity three times

in 10 second intervals. Lysates were clarified by centrifugation at 4°C for 20 min at 12,000 rpm.

Supernatants were transferred to a fresh tube, snap frozen in liquid nitrogen for 2 min and stored

at -80°C until use. Protein concentrations were quantified using a Coomassie Plus (Bradford)

Protein Assay (Pierce) and diluted to 10 μg mL^{-1}. Samples and standards were adsorbed to a 96-

well protein binding plate and ELISA was performed according to manufacturer's instructions.

Absorbance at 450 nm was quantified on a BioTek Epoch2 microplate reader.

Protein carbonyls were visualized using the OxiSelect™ Protein Carbonyl Immunoblot Kit

(Cell Biolabs). *Tnrf1*[-/-] BMDMs were seeded in 24-well plates at 2.5×10^5 one day prior to infection

with indicated *L. pneumophila* strains at an MOI of 50 in the presence or absence of 10 μM H_2O_2.

One hour after infection, cells were washed 3x with PBS[-/-], media were replaced and cells were

incubated for an additional 3 h in the presence or absence of H_2O_2. Cells were washed in ice-cold

PBS-/- and lysed in 120 μL ice-cold NP-40 lysis buffer (see above). Lysates were diluted in 3x

Laemmli sample buffer and boiled for 10 min. Proteins were separated on a 4-20% gradient SDS-

PAGE gel and transferred to PVDF membrane using a BioRad TransBlot semi-dry transfer appa-

ratus. Membranes were processed according to manufacturer's instructions using 10 min wash

steps. Blots were stripped and re-probed with rabbit α-β-actin antibody (1:1000; Cell Signaling

Technology) followed by goat-α-rabbit-HRP (1:5000; ThermoFisher). Blots were visualized as de-

scribed above and densitometry was performed using ImageJ.

<u>Yeast Two-Hybrid Analysis</u>

Yeast two-hybrid screening was performed by Hybrigenics Services, S.A.S., Evry, France

(http://www.hybrigenics-services.com). The *legC4* coding sequence was PCR-amplified from

pSN85::*legC4* (11)and cloned into pB66 as a C-terminal fusion to Gal4 DNA-binding domain

(Gal4-C4). The construct was checked by sequencing and used as a bait to screen a random-

primed Mouse Spleen library constructed into pP6. pB66 derives from the original pAS2ΔΔ vector

(52) and pP6 is based on the pGADGH plasmid (53).

60 million clones (6-fold the complexity of the library) were screened using a mating ap-

proach with YHGX13 (Y187 ade2-101::loxP-kanMX-loxP, mat-α) and CG1945 (mat-a) yeast

strains as previously described (52). 303 His+ colonies were selected on a medium lacking tryp-

tophan, leucine and histidine, and supplemented with 2mM 3-aminotriazole to handle bait auto-

activation. The prey fragments of the positive clones were amplified by PCR and sequenced at

their 5' and 3' junctions. The resulting sequences were used to identify the corresponding

interacting proteins in the GenBank database (NCBI) using a fully automated procedure. A confi-

dence score (PBS, for Predicted Biological Score) was attributed to each interaction as previously

described (54).

<u>Immunofluorescence Microscopy</u>

To quantify LAMP1-LCV co-localization and the number of *L. pneumophila* per LCV, 1×10^5

BMDMs were seeded on poly-L-lysine-coated glass coverslips in 24-well plates and infected with

L. pneumophila at a multiplicity of infection (MOI) of 30. Coverslips were fixed with 4% para-

formaldehyde (PFA; ThermoFisher) and permeabilized in ice cold methanol. Coverslips were

stained with 1:1,000 rabbit α-*L. pneumophila* (Invitrogen, PA17227), rat α-LAMP1 (Developmen-

tal Studies Hybridoma Bank) primary antibodies and 1:500 Alexa 488-conjugated goat α-rabbit

and Alexa594-conjugated goat α-rat secondary antibodies (ThermoFisher) in blocking buffer

[0.1% saponin (w/v), 1% HIFBS (v/v), 0.5% bovine serum albumin (BSA; w/v)] in PBS. Nuclei

were stained with Hoechst (ThermoFisher) at 1:2,000. Coverslips were mounted on glass slides

with ProLong Gold Antifade Mountant (ThermoFisher). Slides were blinded and LAMP1 localiza-

tion to LCVs were scored on a Leica DMiL LED inverted epifluorescence microscope (n=150 cells).

To image ectopically expressed LegC4 and PA28α, 5×10^5 HeLa cells were seeded on poly-

L-lysine-coated glass coverslips in 24-well tissue culture plates and transfected for 24 h. Co-

verslips were fixed in 4% PFA and stained with mouse α-FLAG M2 (1:1,000) and rabbit α-GFP

(1:5,000) primary antibodies and 1:5,00 Alexa546-conjugated goat α-mouse and Alexa488-con-

jugated goat α-rabbit secondary antibodies, as described above. Images were acquired on a Zeiss

AxioPlan LSM-5 laser-scanning confocal microscope (KSU Division of Biology Microscopy Facil-

ity). Images were assembled using Fiji-ImageJ and Adobe Photoshop software.

<u>*L. pneumophila* intracellular growth curves</u>

To quantify *L. pneumophila* intracellular replication, macrophages were seeded in 24-well tissue

culture plates and infected at an MOI of 1 the next day. BMDMs were seeded at 2.5×10^5 in seeding

media (RPMI, 10% HIFBS, 7.5% L929 cell supernatant), THP-1 cells were seeded at 5×10^5 and

differentiated with 100 nM PMA in RPMI 10% HIFBS for three days. Cells were washed with PBS

and incubated in the absence of PMA for one day prior to infection. Stable RAW 264.7 cells were

seeded at 2.5×10^5 in DMEM 2.5%HIFBS. MH-S cells were cultured in RPMI 10% HIFBS and

seeded at 2.5×10^5. At 1 h post-infection, cells were washed 3x with PBS followed by addition of

fresh media. Macrophages were lysed in sterile water and colony forming units (CFU) were enu-

merated at the indicated time points as described (11). Where indicated, 25 ng mL^{-1} rTNF (Gibco),

5 ng mL^{-1} rIFN-γ (Gibco), 10 μM H_2O_2 (VWR), 12.5 nM bafilomycin A1 (BAF; ApexBio), or 100

nM ONX 0914 (ApexBio) were added at the time of infection and maintained throughout. DMSO

was used as a vehicle control where indicated. Fold replication was quantified by normalizing col-

ony forming units to internalized bacteria at 1h post-infection.

<u>Enzyme-linked immunosorbent assay (ELISA)</u>

To quantify TNF secretion, 2.5×10^5 BMDMs were seeded in 24-well tissue culture plates and

infected with the indicated *L. pneumophila* strains at an MOI of 10. After 1 h of infection, cells

were washed with PBS, and media were replaced. Supernatants were at 8 h post-infection and

used fresh or stored at -20˚C for up to 1 week and TNF was quantified using mouse TNF-α

ELISA MAX kit following manufacturer's instructions. Absorbance at 450 nm was quantified on

a BioTek Epoch2 microplate reader.

<u>Quantification of cell death and metabolic activity</u>

To quantify cell death, lactate dehydrogenase activity in cell supernatants was quantified. BMDMs

were seeded at 2.5×10^5 in a 24-well tissue culture plate for 24 h and infected with the indicated *L.*

pneumophila strains at an MOI of 10 in 500 μL of seeding media (RPMP/10% HIFBS/7.5% L929

cell supernatant) and incubated for 10 h. Supernatants were transferred to a 96-well plate and

centrifuged at 200 r.c.f. for 10 min. LDH was quantified using the CytoTox 96 Non-Radioactive

Cytotoxicity Assay (Promega) according to manufacturer's instructions. Absorbance at 490 nm

was quantified on a BioTek Epoch2 microplate reader and percent cytotoxicity was calculated by

normalizing absorbance values to cells treated with lysis buffer.

To quantify cellular metabolic activity, an MTT colorimetric assay was used. BMDMs were

seeded in a 96-well tissue culture plate at 1×10^5 and infected the next day with the indicated *L.*

pneumophila strains at an MOI of 10. Cells were treated with 10 ng mL-1 rTNF and/or 10 µM

staurosporine (Abcam), as indicated. Cells were assayed using the MTT Cell Proliferation Assay

Kit (Colorimetric; BioVision) at 2 h and 24 h post-infection according to manufacturer's instruc-

tions. Absorbance at 590 nm was quantified on a BioTek Epoch2 microplate reader and percent

viability was calculated by normalizing absorbance values to uninfected cells (100% viability).

<u>Statistical analysis</u>

Statistical analyses were performed with GraphPad Prism 9 software using either Mann-Whitney

U test, Students *t*-test, or two-way ANOVA as indicated with a 95% confidence interval. Unless

otherwise indicated, data are presented as mean ± standard deviation (s.d.) and statistical analy-

sis was performed on samples in triplicates.

Acknowledgements

We thank Drs. Philip Hardwidge and Kalyani Pyaram for critical reading of the manuscript and

Dr. Mary Weber for assistance with Biorender.com figure preparation. This work was funded by

NIH/NIGMS P20GM130448 (to S.R.S.); a Kansas State University Johnson Cancer Research

Center Summer Stipend Award (to T.N.); a Kansas-INBRE Postdoctoral Fellowship

(P20GM103418 to D.C.); a Kansas-INBRE Semester Scholar Award (P20GM103418 to A.G.S.);

and startup funds from Kansas State University (to S.R.S.).

REFERENCES

1. G. Mitchell, C. Chen, D. A. Portnoy, Strategies Used by Bacteria to Grow in Macrophages. *Microbiol Spectr* 4 (2016).

2. A. M. Pickering, *et al.*, The immunoproteasome, the 20S proteasome and the PA28αβ proteasome regulator are oxidative-stress-adaptive proteolytic complexes. *Biochem J* 432, 585–94 (2010).

3. I. Dikic, Proteasomal and Autophagy Degradation Systems. *Annu Rev Biochem* 86, 1–32 (2016).

4. N. de Graaf, *et al.*, PA28 and the proteasome immunosubunits play a central and independent role in the production of MHC class I-binding peptides in vivo. *Eur J Immunol* 41, 926–35 (2011).

5. T. Preckel, *et al.*, Impaired Immunoproteasome Assembly and Immune Responses in PA28−/− Mice. *Science* 286, 2162–2165 (1999).

6. D. Chauhan, S. R. Shames, Pathogenicity and Virulence of *Legionella*: Intracellular replication and host response. *Virulence* 12, 1122–1144 (2021).

7. M. F. Fontana, *et al.*, Secreted bacterial effectors that inhibit host protein synthesis are critical for induction of the innate immune response to virulent *Legionella pneumophila*. *Plos Pathog* 7, e1001289 (2011).

8. S. Asrat, A. S. Dugan, R. R. Isberg, The frustrated host response to *Legionella pneumophila* is bypassed by MyD88-dependent translation of pro-inflammatory cytokines. *Plos Pathog* 10, e1004229 (2014).

9. T. Ngwaga, D. Chauhan, S. R. Shames, Mechanisms of Effector-Mediated Immunity Revealed by the Accidental Human Pathogen *Legionella pneumophila*. *Front Cell Infect Mi* 10, 593823 (2021).

10. S. R. Shames, *et al.*, Multiple *Legionella pneumophila* effector virulence phenotypes revealed through high-throughput analysis of targeted mutant libraries. *Proc National Acad Sci* 63, 201708553 (2017).

11. T. Ngwaga, A. J. Hydock, S. Ganesan, S. R. Shames, Potentiation of Cytokine-Mediated Restriction of *Legionella* Intracellular Replication by a Dot/Icm-Translocated Effector. *J Bacteriol* 201, 13 (2019).

12. A. B. Molofsky, *et al.*, Cytosolic recognition of flagellin by mouse macrophages restricts *Legionella pneumophila* infection. *J Exp Medicine* 203, 1093 1104 (2006).

13. T. Ren, D. S. Zamboni, C. R. Roy, W. F. Dietrich, R. E. Vance, Flagellin-deficient *Legionella* mutants evade caspase-1- and Naip5-mediated macrophage immunity. *Plos Pathog* 2, e18 (2006).

14. X. Liu, M. A. Boyer, A. M. Holmgren, S. Shin, *Legionella*-Infected Macrophages Engage the
Alveolar Epithelium to Metabolically Reprogram Myeloid Cells and Promote Antibacterial In-
flammation. *Cell Host Microbe* (2020) https://doi.org/10.1016/j.chom.2020.07.019.

15. U. Seifert, *et al.*, Immunoproteasomes Preserve Protein Homeostasis upon Interferon-In-
duced Oxidative Stress. *Cell* 142, 613–624 (2010).

16. R. P. Fabunmi, W. C. Wigley, P. J. Thomas, G. N. DeMartino, Interferon gamma regulates
accumulation of the proteasome activator PA28 and immunoproteasomes at nuclear PML bod-
ies. *J Cell Sci* 114, 29–36 (2001).

17. S. Wilk, W.-E. Chen, R. P. Magnusson, Properties of the Beta Subunit of the Proteasome Ac-
tivator PA28 (11S REG). *Arch Biochem Biophys* 384, 174–180 (2000).

18. J. D. Mott, *et al.*, PA28, an activator of the 20 S proteasome, is composed of two nonidentical
but homologous subunits. *J Biological Chem* 269, 31466–71 (1994).

19. K. Hallermalm, *et al.*, Tumor necrosis factor-α induces coordinated changes in major histo-
compatibility class I presentation pathway, resulting in increased stability of class I complexes at
the cell surface. *Blood* 98, 1108–1115 (2001).

20. J. Li, S. R. Powell, X. Wang, Enhancement of proteasome function by PA28α; overexpres-
sion protects against oxidative stress. *Faseb J* 25, 883–93 (2010).

21. S. M. Rabadi, *et al.*, Antioxidant Defenses of *Francisella tularensis* Modulate Macrophage
Function and Production of Proinflammatory Cytokines. *J Biological Chem* 291, 5009–21
(2015).

22. P. Bandyopadhyay, B. Byrne, Y. Chan, M. S. Swanson, H. M. Steinman, *Legionella pneu-*
mophila Catalase-Peroxidases Are Required for Proper Trafficking and Growth in Primary Mac-
rophages. *Infect Immun* 71, 4526–4535 (2003).

23. H.-K. An, *et al.*, CASP9 (caspase 9) is essential for autophagosome maturation through regu-
lation of mitochondrial homeostasis. *Autophagy* 16, 1–20 (2019).

24. T. Harada, M. Miyake, Y. Imai, Evasion of *Legionella pneumophila* from the Bactericidal
System by Reactive Oxygen Species (ROS) in Macrophages. *Microbiol Immunol* 51, 1161–1170
(2007).

25. K. J. Flynn, M. S. Swanson, Integrative Conjugative Element ICE-βox Confers Oxidative
Stress Resistance to *Legionella pneumophila* In Vitro and in Macrophages. *Mbio* 5, e01091-14
(2014).

26. P. Bandyopadhyay, H. M. Steinman, Catalase-Peroxidases of *Legionella pneumophila*: Clon-
ing of the katA Gene and Studies of KatA Function. *J Bacteriol* 182, 6679–6686 (2000).

27. U. B. Pandey, *et al.*, HDAC6 rescues neurodegeneration and provides an essential link be-
tween autophagy and the UPS. *Nature* 447, 860–864 (2007).

28. C. R. Roy, K. H. Berger, R. R. Isberg, *Legionella pneumophila* DotA protein is required for
early phagosome trafficking decisions that occur within minutes of bacterial uptake. *Mol Micro-*
biol 28, 663–674 (1998).

29. M. S. Swanson, R. R. Isberg, Association of *Legionella pneumophila* with the macrophage
endoplasmic reticulum. *Infection and immunity* 63, 3609 3620 (1995).

30. N. Matsumoto, M. Nakanishi-Matsui, Proton pumping V-ATPase inhibitor bafilomycin A1
affects Rab7 lysosomal localization and abolishes anterograde trafficking of osteoclast secretory
lysosomes. *Biochem Bioph Res Co* 510, 421–426 (2019).

31. S. Sturgill-Koszycki, M. S. Swanson, *Legionella pneumophila* Replication Vacuoles Mature
into Acidic, Endocytic Organelles. *J Exp Medicine* 192, 1261–1272 (2000).

32. A. A. Khweek, *et al.*, A bacterial protein promotes the recognition of the *Legionella pneu-*
mophila vacuole by autophagy. *Eur J Immunol* 43, 1333 1344 (2013).

33. S. C. Xie, *et al.*, The structure of the PA28-20S proteasome complex from *Plasmodium falci-*
parum and implications for proteostasis. *Nat Microbiol* 4, 1990–2000 (2019).

34. J. Lesne, *et al.*, Conformational maps of human 20S proteasomes reveal PA28- and im-
muno-dependent inter-ring crosstalks. *Nat Commun* 11, 6140 (2020).

35. C. T. D. Price, *et al.*, Dot/Icm-Dependent Restriction of *Legionella pneumophila* within Neu-
trophils. *Mbio* 12, e01008-21 (2021).

36. P. Ziltener, T. Reinheckel, A. Oxenius, Neutrophil and Alveolar Macrophage-Mediated In-
nate Immune Control of *Legionella pneumophila* Lung Infection via TNF and ROS. *Plos Pathog*
12, e1005591 (2016).

37. J. D. MacMicking, Interferon-inducible effector mechanisms in cell-autonomous immunity.
Nat Rev Immunol 12, 367 382 (2012).

38. J. Naujoks, *et al.*, IFNs Modify the Proteome of *Legionella*-Containing Vacuoles and Restrict
Infection Via IRG1-Derived Itaconic Acid. *Plos Pathog* 12, e1005408 (2016).

39. A. Orvedahl, *et al.*, Autophagy genes in myeloid cells counteract IFNγ-induced TNF-medi-
ated cell death and fatal TNF-induced shock. *Proc National Acad Sci* 116, 16497–16506 (2019).

40. B. G. Byrne, J.-F. Dubuisson, A. D. Joshi, J. J. Persson, M. S. Swanson, Inflammasome Com-
ponents Coordinate Autophagy and Pyroptosis as Macrophage Responses to Infection. *Mbio* 4,
e00620-12 (2013).

41. J. N. Shin, E. A. Fattah, A. Bhattacharya, S. Ko, N. T. Eissa, Inflammasome Activation by Al-
tered Proteostasis*. *J Biol Chem* 288, 35886–35895 (2013).

42. A. Marra, S. J. Blander, M. A. Horwitz, H. A. Shuman, Identification of a *Legionella pneu-*
mophila locus required for intracellular multiplication in human macrophages. *Proc Natl Acad*
Sci 89, 9607–9611 (1992).

43. J. Zhao, *et al.*, Molecular basis for the binding and modulation of V-ATPase by a bacterial
effector protein. *Plos Pathog* 13, e1006394 (2017).

44. A. Choy, *et al.*, The *Legionella* effector RavZ inhibits host autophagy through irreversible
Atg8 deconjugation. *Science* 338, 1072 1076 (2012).

45. M. Rolando, *et al.*, *Legionella pneumophila* S1P-lyase targets host sphingolipid metabolism
and restrains autophagy. *Proc Natl Acad Sci* 113, 1901 1906 (2016).

46. M. Rolando, P. Escoll, C. Buchrieser, *Legionella pneumophila* restrains autophagy by modu-
lating the host's sphingolipid metabolism. *Autophagy* 12, 1053–1054 (2016).

47. J. C. Feeley, *et al.*, Charcoal-yeast extract agar: primary isolation medium for *Legionella*
pneumophila. *J Clin Microbiol* 10, 437 441 (1979).

48. A. Saito, R. D. Rolfe, P. H. Edelstein, S. M. Finegold, Comparison of liquid growth media for
Legionella pneumophila. *J Clin Microbiol* 14, 623 627 (1981).

49. C. L. Case, C. R. Roy, Analyzing caspase-1 activation during *Legionella pneumophila* infec-
tion in macrophages. *Methods Mol Biol (Clifton, NJ)* 954, 479 491 (2013).

50. A. Ingmundson, A. Delprato, D. G. Lambright, C. R. Roy, *Legionella pneumophila* proteins
that regulate Rab1 membrane cycling. *Nature* 450, 365–369 (2007).

51. S. R. Shames, *et al.*, The pathogenic *E. coli* type III effector EspZ interacts with host CD98
and facilitates host cell prosurvival signalling. *Cell Microbiol* 12, 1322 1339 (2010).

52. M. Fromont-Racine, J.-C. Rain, P. Legrain, Toward a functional analysis of the yeast genome
through exhaustive two-hybrid screens. *Nat Genet* 16, 277–282 (1997).

53. P. L. Bartel, S. Fields, [16] Analyzing protein-protein interactions using two-hybrid system.
Methods Enzymol 254, 241–263 (1995).

54. E. Formstecher, *et al.*, Protein interaction mapping: A *Drosophila* case study. *Genome Res*
15, 376–384 (2005).

Figure 1. LegC4 attenuates *L. pneumophila* replication within cytokine-activated macrophages

and the mouse lung. Growth of indicated *L. pneumophila* strains within **(A)** WT BMDMs, **(B)**

THP-1 cells, **(C)** *Myd88$^{-/-}$* BMDMs, **(D)** *Tnfr1$^{-/-}$* BMDMs, or **(E)** *Ifngr1$^{-/-}$* BMDMs over 48 h. Plas-

mid expression of *legC4* was induced with 1 mM IPTG. Macrophages were infected in triplicates

and data are presented as mean ± standard deviation (s.d.). Asterisks denote statistical signifi-

cance by Student's *t*-test (*P<0.05, **P<0.01). **(F)** Competitive index (CI) of *L. pneumophila*

ΔflaAΔlegC4 (pJB*legC4*) or *ΔflaAΔlegC4* (pJB) versus *L. pneumophila ΔflaA* in the lungs of the

indicated mice at 48 h post-infection. Each symbol represents an individual animal and asterisks

denote statistical significance by Mann-Whitney U test (**P<0.01). Data shown are representative

of at least three independent experiments.

Figure 2. LegC4 binds and functions through PA28αβ. **(A)** Western blot of co-immunoprecipi-

tation of PA28α-Myc with 3xFLAG-LegC4 on α-FLAG conjugated magnetic beads (top) or

3xFLAG-LegC4 with GFP-PA28α on α-GFP-conjugated magnetic beads (bottom) from lysates of

transfected HEK 293T cells. **(B)** Confocal microscopy to image HeLa cells co-transfected to ec-

topically produce 3xFLAG-LegC4 (red) and either GFP-PA28α or GFP (green). Scale bar repre-

sents 10 μm. **(C)** Fold replication of indicated *L. pneumophila* strains within *Psme1/2*$^{-/-}$ (left) or

WT (right) BMDMs. Plasmid expression of *legC4* was induced with 1 mM IPTG. **(D)** Fold repli-

cation of *L. pneumophila* (*L.p.*) Δ*flaA* within WT or *Psme1/2*$^{-/-}$ BMDMs at 48 h post-infection.

Where indicated BMDMs were treated with 25 ng mL^{-1} recombinant mouse (r)TNF, 5 ng mL^{-1}

rIFN-γ, or left untreated (UT). Macrophages were infected in triplicates and data are presented as

691   mean ± s.d. Asterisks denote statistical significance (**$P<0.01$) by Students' t-test. Data shown

692   are representative of at least two independent experiments.

Figure 3. LegC4 impairs resolution of oxidative stress, which is sufficient for *L. pneumophila*

restriction. **(A)** Confocal microscopy to visualize 3xFLAG-LegC4 (red) production by stable RAW

264.7 cells (RAW-LegC4). Mock transfected cells (RAW-Cntrl) were used as a control and nuclei

were imaged with Hoechst. Scale bar represents 10 μm. **(B)** Western blot analysis of 3xFLAG-

LegC4 immunoprecipitated from lysates of RAW-LegC4, RAW-Cntrl or untransfected (UT) RAW

cells using α-FLAG-conjugated beads. Arrow indicates 3xFLAG-LegC4 (~84 kDa). **(C)** Growth of

L. pneumophila Δ*flaA* within RAW-LegC4 and -Cntrl cells infected for up to 48 h. **(D)** ELISA

quantification of protein carbonyls in lysates of RAW-LegC4 and -Cntrl cells incubated for 24 h

in the presence or absence of 10 μM H_2O_2. **(E)** Western blot for protein carbonyls and actin (load-

ing control) in lysates of *Tnfr1$^{-/-}$* BMDMs infected for 4 h with the indicated strains. ΔΔ:

Δ*flaA*Δ*legC4*; plasmid expression of *legC4* was induced with 1 mM IPTG. **(F)** Fold replication of

L. pneumophila strains at 48 h post-infection within *Tnfr1$^{-/-}$* BMDMs in the presence or absence

of 10 μM H_2O_2 as indicated. Plasmid expression of *legC4* was induced with 1 mM IPTG. All quan-

titative data are presented as mean ± s.d. from samples in triplicates and are representative of at

least two independent experiments. Asterisks denote statistical significance by Students' *t*-test

(**P<0.01; n.s., not significant).

Figure 4. Role of phagolysosomal fusion in LegC4-mediated restriction of *L. pneumophila*. **(A)**
Blinded immunofluorescence scoring of LAMP1 localization to LCVs within WT or *Tnfr1*[-/-]
BMDMs after 9 h of infection. Plasmid expression of *legC4* was induced with 1 mM IPTG. Data
are representative of at least two independent experiments ($n \geq 150$ per experiment) and letters
denote statistical significance by Students' *t*-test ($P<0.05$). letters not significant) of values com-
pared to Δ*flaA*. **(B)** WT BMDMs were infected with the indicated *L. pneumophila* strains and the
number of bacteria per LCV was quantified by blinded immunofluorescence scoring. Data are
representative of at least two independent experiments ($n \geq 150$ per experiment) and asterisks
denote statistical significance by Students' *t*-test (**$P<0.01$; n.s., not significant) of values com-
pared to Δ*flaA*. Plasmid expression of *legC4* was induced with 1 mM IPTG. **(C)** Growth of

725   indicated *L. pneumophila* strains within WT BMDMs treated with 12.5 nM bafilomycin A1 (BAF)

726   or DMSO (vehicle control). Plasmid expression of *legC4* was induced with 1 mM IPTG. Data are

727   presented as mean ± s.d. of samples in triplicates and are representative of at least two independ-

728   ent experiments. Asterisks denote statistical significance by Students' *t*-test (*P<0.01).