1

Denitrification in foraminifera has ancient origin and is complemented by associated bacteria

Christian Woehle[1,5*], Alexandra-Sophie Roy[1*], Nicolaas Glock[2,6], Jan Michels[3], Tanita Wein[1,7],

Julia Weissenbach[1,8], Dennis Romero[4], Claas Hiebenthal[2], Stanislav N. Gorb[3], Joachim

Schönfeld[2], Tal Dagan[1]

[1]Institute of General Microbiology, Kiel University, Am Botanischen Garten 11, Kiel, 24118,

Germany

[2]GEOMAR Helmholtz Centre for Ocean Research Kiel, Wischhofstrasse, Kiel, 24148,

Germany

[3]Zoological Institute, Kiel University, Am Botanischen Garten 1-9, Kiel, 24118, Germany

[4]Dirección General de Investigaciones Oceanográficas y Cambio Climático, Instituto del Mar

del Perú, Callao 01, Peru 17

[5]Present address: Miltenyi Biotec B.V. & Co. KG, Friedrich-Ebert-Str. 68, Bergisch Gladbach,

51429, Germany

[6]Present address: Institute for Geology, University of Hamburg, Hamburg, 20146, Germany

[7]Present address: Department of Molecular Genetics, Weizmann Institute of Science, Rehovot

76100, Israel

[8]Present address: Center for Ecology and Evolution in Microbial model Systems-EEMiS,

Linnaeus University, Kalmar, 39182, Sweden

* Equally contributing authors

Corresponding authors: Christian Woehle woehle@gmail.com (C.W.), Tal Dagan

tdagan@ifam.uni-kiel.de (T.D.)

Keywords: eukaryotic evolution, Foraminifera, denitrification pathway, nitrogen cycle,

microbiome

Abstract

Benthic foraminifera are unicellular eukaryotes that inhabit sediments of aquatic environments. Several foraminifera of the order Rotaliida are known to store and use nitrate for denitrification, a unique energy metabolism among eukaryotes. The rotaliid *Globobulimina* spp. has been shown to encode an incomplete denitrification pathway of bacterial origin. However, the prevalence of denitrification genes in foraminifera remains unknown and the missing denitrification pathway components are elusive. Analysing transcriptomes and metagenomes of ten foraminiferal species from the Peruvian oxygen minimum zone, we show that denitrification genes are highly conserved in foraminifera. We infer of the last common ancestor of denitrifying foraminifera, which enables us to predict the ability to denitrify for additional foraminiferal species. Additionally, an examination of the foraminiferal microbiota reveals evidence for a stable interaction with *Desulfobacteracea*, which harbour genes that complement the foraminiferal denitrification pathway. Our results provide evidence that foraminiferal denitrification is complemented by the foraminifera-associated microbiome. The interaction of foraminifera with their resident bacteria is at the basis of foraminiferal adaptation to anaerobic environments that manifested in ecological success within oxygen depleted habitats.

Significance Statement

A substantial component of the global nitrogen cycle is the production of biologically inaccessible dinitrogen attributed to anaerobic denitrification by prokaryotes. Recent evidence identified a eukaryote – foraminifera – as new key players in this 'loss' of bioavailable nitrogen. The evolution of denitrification in eukaryotes constitutes a rare event and the genetic mechanisms of the denitrification pathway in foraminifera are just starting to be elucidated. We present large-scale sequencing analyses of ten denitrifying foraminiferal species, which reveals the high conservation of the foraminiferal denitrification pathway. We further find evidence for a complementation of denitrification by the foraminiferal microbiome. Together, these findings provide novel insights on the early evolution of a previously overlooked component in the marine nitrogen cycle.

Introduction

Nitrogen is an essential element for life on earth as it forms the basis for the synthesis of nucleotides and amino acids. Nonetheless, while the earth atmosphere is rich in nitrogen gas (up to 78%) (1), this gas is usually inert but can be made accessible for biological processes

by nitrogen fixation (2). Microbial organisms are important players in the global nitrogen cycle
as they facilitate the assimilation of nitrogen into bioavailable nitrogen derivatives as well as
the dissimilation of nitrogen derivatives into dinitrogen (N_2) (2). A key dissimilatory pathway is
denitrification, where nitrate (NO_3^-) is either partially or completely degraded and the final
product – N_2 – is released to the atmosphere (i.e., nitrogen loss) (2). Marine organisms are
considered major contributors to nitrogen loss from the environment with benthic organisms
being responsible for about two-thirds of the loss of reactive nitrogen in the ocean (3, 4).
Especially oxygen minimum zones (OMZs) are worth mentioning here as they are estimated
to be responsible for 20-40% of bioavailable nitrogen removal in the ocean (3, 5). The ability
to perform denitrification is abundantly found in eubacteria (6), whereas it is rare among the
eukaryotes. Partial or complete denitrification have only been reported for two species of fungi
(7) and several foraminifera of the order Rotaliida (8-11). Denitrifying foraminifera are
unicellular eukaryotes commonly found in marine sediments. Studies of foraminifera residing
in the Peruvian OMZ showed that they are found in high densities of up to 600 individuals per
cm^2, where they are estimated to contribute 20%-50% to the total benthic nitrate loss in the
OMZ (10, 12).

Rotaliid foraminifera are traditionally divided into three clades based on their phylogeny
(13). Only species in clades I and III were demonstrated to denitrify, while rotaliids classified
in clade II have been shown to lack an intracellular nitrate storage or measurable denitrification
activity (9). Class II rotaliids typically populate environments with poor-nitrate supply, e.g.
intertidal to near-shore habitats from the tropics up to boreal bioprovinces (14), hence it is
conceivable that they lack the ability to denitrify. One example is *Ammonia tepida* (class
II), that can survive episodic oxygen depletion events via dormancy (15). A study of clade III
species sampled from a hypoxic environment (Gulmarfjord, Sweden) – *Globobulimina* spp. –
showed that that their genome encodes several genes along the denitrification pathway that
are of ancient bacterial origin (11). These include the copper-containing nitrite reductase (NirK)
and nitric oxide reductase (Nor), but not the nitrate reductase (NapA/NarG) and nitrous oxide
reductase (NosZ). In addition, the *Globobulimina* genome contains a diverse gene family
encoding for nitrate transporters (Nrt) (11), a finding that is consistent with the accumulation
of an intracellular NO_3^- storage in denitrifying rotaliids (8-10, 16). More recent studies reported
the presence of those genes in at least two additional species of foraminifera and provide
alternative suggestions on the nitrate reductase homolog missing from the earlier model of
denitrification in foraminifera (17, 18). The rotaliids ability to respire both oxygen (O_2) and NO_3^-
marks them as facultative anaerobes that are able to thrive in both aerobic and anaerobic
conditions. Furthermore, while oxygen is generally considered to be preferred over NO_3^- as
electron acceptor, rotaliids from the Peruvian OMZ were reported to prefer NO_3^- over O_2 (10).

These findings suggest that genes of the denitrification pathway are widespread in rotaliids,
however their distribution remains largely unknown.

Foraminifera, like other eukaryotes (19), are populated by bacterial organisms that
reside out- and in-side their cell (i.e., test) (20-24). Microscopic observations showed that the
bacteria are localized in food vacuoles or in the cytoplasm of the cell, which led to the
suggestion that interactions of foraminifera with its microbiota may vary from prey-predator
interactions or parasitism to metabolic symbiosis (22). Further studies of the microbiota in
clade Ⅱ rotaliids identified sulphate-reducing and sulphur-oxidizing bacteria that were
suggested to participate in Sulphur cycling thus providing carbon and other nutrients to the
host (20, 21, 25). Notwithstanding, previous studies of foraminifera-associated bacteria used
microscopy observations or sequencing of marker genes (e.g., 16S rRNA) that are limited in
their resolution (20-25). It has been suggested that the microbiota of denitrifying foraminifera
include denitrifying bacteria that utilize the foraminiferal NO_3^- storage, similarly to observations
in other bacteria-protist associations [e.g., gromiids (26), allogromiids (27)]. Indeed,
metagenomics of the *Globobulimina* microbiota revealed the presence of a taxonomically
diverse species community where several members encode homologs of NapA and NosZ (11).
This finding gave rise to the hypothesis that the partial foraminiferal denitrification pathway
may be complemented by microbiota functions.

Here we investigate the evolutionary history of genes along the denitrification pathway
in foraminifera and furthermore examine the functional repertoire of the foraminiferal
microbiome. For that purpose, we studied populations of ten rotaliid species known to denitrify
(10) (except *Globobulimina pacifica* for which no information is available). Our study supplies
insights in the evolution of rotaliids and their microbiota and lays the basis for further research
of foraminiferal genome evolution.

128   Results

129   Transcriptomes of ten Peruvian rotaliids

For the purpose of our study, we sampled benthic foraminifera in the Peruvian oxygen
minimum zone. Individual foraminifers from among ten focal species were manually identified
and picked using a stereomicroscope (see sampled species in Figure 1A). Eukaryotic
transcriptomes and whole metagenomes were sequenced from the sampled foraminifera with
two biological replicates per species (ca. 160 individuals per sample). Our computational
analysis includes five additional publicly available transcriptomes of foraminifera (Suppl. Table
S1) and the genome data of one monothalamid species (*Reticulomyxa filosa*). The
assessment of transcriptome completeness showed that ~90% of the eukaryotic marker

proteins are present in the data of the ten newly sequenced species. Furthermore, we

examined the purity of the focal species in each transcriptome by testing for redundancy of

eukaryotic marker proteins that are expected as single-copy genes. As single-copy genes are

expected to exist only once in a given species, their redundancy might indicate for additional

eukaryotic species present in our samples. The highest proportion of such bystander species

were found in the *B. spissa* (78%) and *B. costata* (50%) transcriptomes (Suppl. Table S1).

The Peruvian rotaliids harbour genes of the denitrification pathway

To explore the denitrification mechanisms in the Peruvian foraminifera, we searched for

homologs to denitrification pathway genes. Our results reveal that all sampled species harbor

homologous genes to Nrt, NirK and Nor (Figure 1B; note that genes are termed by their

corresponding protein symbol). Our search for homologs in the publicly available data yielded

a putative homolog of NirK in *Rosalina* sp. and Nrt homologs in all but *R. filosa* (Fig. 1B). The

additional species we included here, especially *Ammonia sp.*, *Elphidium margaritaceum* and

Sorites sp., are considered to reside mostly in oxygenated habitats, hence they are not

expected to be able to perform denitrification. These species lack homologs to the

denitrification genes, but encode nitrate transporters (Nrt, Figure 1B). Among this group, only

A. tepida was so far experimentally tested for denitrification ability and reported as a non-

denitrifying species (9). The reconstructed phylogenies indicate that all the examined genes

are indeed encoded in the foraminiferal genomes, as all of them group with homologs

previously reported to originate from the nuclear genomes of *Globobulimina* spp. (11).

Furthermore, the phylogenies of Nrt and NirK, reveal two major sub-clades including most, but

not all, of the species sampled (Full and half circles in Fig. 1B; Suppl. Figure S2; Suppl. Table

S2). Previously we reported the presence of subclades in the Nrt and NirK phylogenies (11),

but could not clarify if those were explained by sequencing errors, gene duplication, or

speciation events. The phylogenetic trees presented here show that each subclade is

represented in most of the rotaliids sampled in the current study. Consequently, we conclude

that ancient gene duplications in the Nrt and NirK gene families led to the evolution of two or

more sub-clades that encode for different (paralogous) protein subtypes. Overall, our results

demonstrate that the denitrification pathway is highly conserved in the Peruvian rotaliids.

Figure 1. Morphological characteristics of the sampled rotaliids and denitrification gene repertoire. A) Scanning electron micrographs of the sampled species. Species with clearly distinct lateral views are shown from two sides. For more detailed views see Suppl. Figure S1. **B)** Presence/absence of homologs of denitrification proteins identified for different foraminifera. Half circles indicate species where only a single, of multiple subtypes, was found. Dashed circles illustrate homologs discarded due to low coverage and asterisks highlight corresponding species. Evidence for NO_3^- storage and/or denitrification activity is illustrated by closed circles in the last column. Question marks denote missing information. Foraminifera sampled in this study are highlighted by the location 'Peru'. Protein symbols: Nrt: nitrate/nitrite transporter; NapA: periplasmic nitrate reductase; NarG: membrane-bound nitrate reductase; NirK: copper-containing nitrite reductase; NirS: cd_1-containing nitrite reductase; Nor: nitric oxide reductase; NosZ: nitrous oxide reductase. Note that the identification of homologous genes is based not only on sequence similarity but also on transcript abundance in order to exclude bystander species in the data. The additional data filtration stage affected our findings for *Rosalina* sp., *B. plicata* and *N. auris* (Suppl. Table S2 & S3), where the presence of at least one the crucial homologs (i.e, NirK or Nor) remained in the *B. plicata* and *N. auris* metatranscriptomes.

Foraminiferal denitrification evolved in the Rotaliida ancestor

The presence of NirK and Nor homologs the tested rotaliids suggests that those genes may

have an ancient origin in this order. To study the origin of denitrification in foraminifera, we

reconstructed the phylogenetic relations among species in our data using 81 eukaryotic

marker proteins that have homologs in all species analyzed here (Suppl. Figure S3; Suppl.

Table S4). Here, we applied a phylogenomics approach where the root position is inferred

from all marker genes independent of a single species tree (28) (see methods). Our results

show that the best-supported root position (42% of the gene trees) was at the branch leading

to *R. filosa* (Suppl. Figure S3). This result is in agreement with previous studies that assumed

monothalamids to be an outgroup in foraminiferal phylogenies (29). The second most frequent

root position was found at the branch leading to the miliolid *Sorites* sp. (15% of single gene

trees), which was followed closely by a root position on the branch leading to *Rosalina* sp.

(12%). Taken together, the inferences of rooted topologies show that the Peruvian rotaliids,

together with *Globobulimina* spp. form a monophyletic group. Thus, our results reveal a shared

origin of denitrification in foraminifera and hence the evolution of denitrifying species within

the order Rotaliida.

For the inference of the denitrifying rotaliids last common ancestor (LCA; i.e., the

rotaliids root), we reconstructed phylogenetic trees from an extended set of 146 eukaryotic

marker protein coding genes considering only members of the Rotaliida (Figure 2; Suppl.

Table S4). An examination of the rooted tree topologies revealed that the branch leading to

Ammonia sp. and *E. margaritaceum* was the most frequently supported root position (14% of

single gene trees). This root inference is further supported by an alternative root position at a

branch that splits *Ammonia* sp., *E. margaritaceum* and *Rosalina* sp. from the remaining

species (10%). Overall, the different alternative root positions support with 35% a Rotaliida

LCA within clade II foraminifera (i.e., among *Ammonia sp.*, *E. margaritaceum & Rosalina* sp.;

Fig. 2). The absence of denitrification genes in clade II members (Fig. 1) further supports the

suggestion that the evolution of denitrification in rotaliids occurred by lateral gene transfer from

a prokaryotic donor rather than being inherited from the rotaliid LCA (11).

Figure 2. Phylogenetic rooting of rotaliids. A Maximum-likelihood phylogenetic reconstruction of foraminiferal species from 146 eukaryotic protein marker sequences is shown on the left. The branch with the LCA of denitrifying foraminifera is highlighted in purple, where a lighter color includes lower bound for the origin of denitrification. Nonparametric bootstrap support is 1000/1000 at all the branches. A split representation of root splits determined via the MAD approach for the 146 single gene trees is shown on the right. Each column represents a putative root branch reported as split of two groups (black and white boxes) indicating for the species found on either side of the branch. The bar graph reports the single gene tree count supporting the corresponding root split.

Ancient origin of denitrification in foraminifera

To further study the denitrifying foraminifera LCA, we examined the phylogenetic position of

the Peruvian species in the context of a foraminiferal species tree. For that purpose, we

extracted sequences of the 18S ribosomal RNA (rRNA) subunit from our transcriptome data

and used it in combination with publicly available 18S rRNA sequences to reconstruct a

foraminiferal species phylogeny (Figure 3A; Suppl. Table S5). The resulting rooted topology

was mostly in agreement with the rotaliids phylogeny, where the focal species are found next
to their closest relatives. Most species grouped well with their previously defined clades with
the exception of *Nonionella* and *Globobulimina* representatives. Furthermore, several taxa
were not monophyletic, as has been previously observed in foraminiferal 18S rRNA
phylogenies (29, 30).

Figure 3. The origin of denitrification in context of the whole foraminiferal group. A) Large-scale phylogenetic representation of Peruvian foraminifera in context of different foraminiferal taxa based on 18S sequences available in public databases. Clades I & III of the order Rotaliida appear are highly supported by bootstrap values. Questionable branching of *Nonionella* outside class III is not characteristic to data of the current study as exemplified for *N. labradorica* that as well does not group directly with clade III. *Globobulimina pacifica* forms a clade with *G. auriculata*, but within clade III. However, we consider the branching of these two species as uncertain due to the long branches and the low bootstrap support, which

we did not observe for the marker protein phylogeny (Figure 2). Lower (light purple; including *N. labradorica*) and higher (dark purple) boundaries for the origin of foraminiferal denitrification are highlighted by boxes. **B)** Extended phylogenetic representation of foraminifera including planktonic species. Lower (light purple) and higher (dark purple) boundaries for the origin of foraminiferal denitrification are highlighted by boxes. Planktonic I & III designate two to three distinct clusters comprising planktonic foraminifera. Species experimentally shown to denitrify are highlighted in blue. The only exception is *Stainforthia fusiformis*, where denitrification activity has been shown for an unspecified species of the same genus. Species also considered in Figure 1 or Suppl. Figure S3 are highlighted by asterisks. Several species labels contain as well the contig id and orientation of 18S sequences in the corresponding transcriptome assemblies. Bootstrap support values (>=70) with 1000 replicates are shown at the branches. The trees were rooted by the clade of monothalamids containing *R. filosa*. A detailed phylogeny is presented in Supplementary Figure S4.

The inference of the denitrifying foraminiferal LCA enabled us to further reconstruct
the origin of denitrification within the foraminiferal species tree. For that purpose, we examined
an extended phylogeny with additional foraminiferal groups including planktonic species
(Figure 3B; Suppl. Figure S4). While the general taxonomic relationships in the tree were
recovered as before (i.e., Figure 3A), many branches had a low statistical support. For
example, the rotaliid clades II and III were paraphyletic, in contrast to the phylogeny with less

taxa (Figure 3A). Nonetheless, an LCA of clades I and III could be inferred; the previously unassigned genera *Valvulineria* and *Cancris* (9) grouped well with members of clade III in both 18S rRNA phylogenies, hence they can be classified as clade III.

Since both clades I and III harbour several denitrifying foraminifera, it is most parsimonious to conclude that LCA of clades I and III was likely a denitrifying organism. Our inference thus suggests the presence of denitrification genes (and hence denitrification capability) in all members of clades I and III including e.g., the genera *Cibicoides* and *Virgulinella*, which were previously not considered as denitrifying species. Indeed, *Cibicidoides wullerstoerfi* has been assumed for a long time to be unable to withstand O_2 depletion (31). However, recent studies observed living *Cibicidoides spp.* thriving in environments of < 2 µmol/kg O_2 (32) and fossil specimens in the paleorecord during periods of severe O_2 depletion (33). These recent studies support our prediction that several *Cibicidoides spp.* are able to denitrify.

Composition of the foraminiferal microbiome is species-specific
To test for bacterial contribution to the foraminiferal denitrification, we examined the foraminiferal microbiome. For that purpose, we compared the composition of bacterial communities among foraminifera in our sample. In the absence of species-specific interactions, we would expect a strong impact of the environment (i.e., sampling location and water depth) on the microbiome composition. Thus, we compared the microbial community composition between foraminiferal species and their sampling depths (Figure 4A). Our results show that the individual microbiomes are clustered by the foraminiferal species rather than the sampling location, indicating the presence of species-specific bacterial communities in foraminifera. Our results reveal a similar taxonomic composition of bacterial communities associated with *Globobulimina* spp. from Sweden and *G. pacifica* from the Peruvian OMZ (Figure 4A). The similarity in microbiome composition among the *Globobulimina* species indicates the presence of genus-specific microbiome.

To identify common key players in foraminiferal microbiota we examined the relative abundance of all bacterial families in their microbiome (Figure 4B; Suppl. Table S6). Our results revealed that the most prevalent bacterial families are Desulfobacteracea followed by Planctomycetaceae. The relative abundance of these two families varied between different species and samples. Furthermore, communities with a high relative abundance of Desulfobacteracea are characterized by a low abundance of Planctomyceteaceae and vice versa (r_s=-0.77, p-value < 0.01, using Spearman correlation coefficient and t-test). Desulfobacteracea comprise sulphate-reducing bacteria and several representatives that are able to grow chemoautotrophically (34). Members of this group were previously observed in

253　association with other rotaliids like *Virgulinella fragilis* and members of clade Ⅱ (20, 25).

254　Planctomycetes are often found in association with macroalgae (35); they are characterized

255　by a diversified metabolism allowing them the of colonization of a wide range of habitats (35).

Figure 4. Comparison of microbial communities associated with different foraminifera. A) Principal coordinates analysis (PCoA) clustering of bacterial communities. Each circle represents a bacterial community; duplicates per species are connected by lines. The text next to the circles indicates for species name, replicate number or sampling depth. The clustering of most species-associated communities indicates for species-specificity. Grouping by higher taxonomy was only observed for *Globobulimina* spp. from Sweden and *Globobulimina pacifica* from the Peruvian oxygen minimum zone (OMZ). **B)** Bar graphs of the six most abundant bacterial families found in the foraminiferal bacterial communities beginning at the top. Each bar represents the proportion of reads assigned to the individual taxon relative to the total number of reads in a sample classified on family rank or lower. Note the prevalence of Desulfobacteraceae shared between the Globobulimina samples, while none of the other bacterial families showed high prevalence in the genus.

The next most abundant families were Alteromonadaceae, Vibrionaceae and

`Candidatus Brocadiaceae` that were more diverse in their relative abundances across

different species and replicates. Most reported `Candidatus Brocadiaceae` members are

autotrophic, obligately anaerobic bacteria performing anammox, which is a dissimilatory

pathway where ammonium and nitrite (NO_2^-) are metabolized into N_2 (36). The presence of

nitrate storage in foraminifera and NirK suggest that nitrite is readily available inside the

foraminiferal test. Vibrionaceae are a diverse group including multiple species that colonize

marine organisms either as symbionts (37, 38) or pathogens (39). Alteromonadaceae were

isolated from diverse marine environments including eukaryotic microbiota (40). They are

considered to be aerobe or facultative anaerobic bacteria that generally lack denitrification

capabilities (41). Finally, another abundant family, Rhodobacteracea, is worth mentioning, as

it is always found in low, but similar abundance in all samples. Marine Rhodobacteracea

species are considered ecological generalists (42-44) and have been reported to colonize

marine animals (e.g., fish larvae or sponges) (45, 46). The uniform distribution of

Rhodobacteracea in the foraminifera-microbiome suggests that members of this group are
permanent residents in the foraminiferal microbiota.
Our data thus shows that rotaliids are habitat to bacterial communities whose
composition is akin to the microbiota of other marine eukaryotes.

An ancient interaction between Desulfobacteraceae and *Globobulimina* hints to metabolic dependency

The metagenomes analysis and classification into bacterial metagenome-assembled
genomes (MAGs) resulted in a total of 263 high-quality MAGs (i.e., draft bacterial genomes)
of foraminifera-associated bacteria (Suppl. Table S1). The strongest signal for a stable core
microbiome in our data was observed in the comparison among the *Globobulimina* species,
whose microbiome is characterized by a high frequency of Desulfobacteraceae (Figure 4B).
To further explore the association between *Globobulimina* and Desulfobacteraceae we
examined the Desulfobacteraceae MAGs in our data. A total of 40 high quality MAGs were
obtained from the *G. pacifica* metagenome, including four high quality draft MAGs classified
as Desulfobacteracea. These were compared with the 26 previously published MAGs for
Globobulimina from Sweden, which include two Desulfobacteracea MAGs (11) (Suppl. Table
S7). A phylogenetic network of the Desulfobacteracea MAGs reveals two MAGs from the
Globobulimina spp. and *G. pacifica* metagenomes (Glo_11 & Gpa_30) that appear as sister
taxa (Figure 5A). The average nucleotide identity (ANI) between both MAGs was 84%, which
is within the range expected for inter-species sequence similarity (e.g., within genera) (47).
The common ancestry of these two MAGs suggests that the association between
Desulfobacteracea and *Globobulimina* has an ancient origin. Our results thus indicate that the
interaction fidelity between *Globobulimina* and Desulfobacteracea is high, similarly to
observations in other eukaryote-bacterial symbioses [e.g., oligochaete worms and sulfur
bacteria (48) or marine sponges with bacteria of the Poribacteria phylum (49)].
A central foundation in the evolution of symbiotic interaction is an exchange of
currencies (i.e., resources) between the partners (50). The versatile metabolic capabilities of
Desulfobacteraceae and the availability of intracellular NO_3^- storage in foraminifera may
suggest that the interaction between Desulfobacteracea and *Globobulimina* is based on
nutritional currencies. To further examine possible symbiotic interaction between
Desulfobacteracea and *Globobulimina*, we searched for metabolic properties of the
Desulfobacteraceae MAGs that could serve as nutritional currencies in the symbiosis. For that
purpose, we surveyed the metabolic pathways of representative Desulfobacteracea MAGs
with respect to carbon, sulfur and nitrogen metabolism (Figure 5B; Suppl. Table S8). Our
results reveal that all MAGs sampled encode the genetic repertoire required in order to
perform dissimilatory sulfate reduction as well as the Wood-Ljungdahl pathway. Anaerobic

respiration via the dissimilatory sulfate reduction to sulfide is widespread among
Desulfobacteraceae. Members of this family that encode the Wood-Ljungdahl pathway are
able to oxidize organic compounds to carbon dioxide (51). Alternatively, the Wood-Ljungdahl
pathway may function in CO_2 carbon fixation during autotrophic growth.

Figure 5. Characteristics of Desulfobacteraceae bacteria associated with Globobulimina. A) Neighbor-net network for Desulfobacteraceae associated with the genus *Globobulimina* from Sweden and Peru (*Globob-ulimina* spp. and *G. pacifica*, respectively) and additional Deltaproteobacteria. The network is based on a concatenated alignment of 33 single copy orthologous protein sequences found in all isolates shown. Red fonts highlight isolates found in association with members of the genus *Globobulimina.* The monophyly of the MAG Gpa_30 and MAG Glo_11 suggests a common ancestry. Note that the closest relatives to that pair are two uncultivated MAGs (Deltaproteobacteria bacterium B6_G6 & Desulfobacteraceae bacterium 4572_89) sampled from sediments in deep-sea hydrothermal vent sites at the Gulf of California. **B)** Presence of predicted proteins encoding metabolic pathways in representative members of Desulfobacteraceae. Three pathways are shown: Wood-Ljungdahl pathway, (dissimilatory) sulphate reduction & nitrate respiration. Asterisks highlight genomes in draft state. Circles indicate for the presence of homologs identified by the KEGG KAAS server. The thin bars link enzymes that participate in the same reaction step either by being an alternative enzyme or being part of one protein complex. The grey circles indicate for cases that were discarded following a manual inspection.

Genes along the denitrification pathway (or NO_3^- respiration) are mostly absent from
all Desulfobacteracea MAGs, except for periplasmic NO_3^- reductases (NapA) homologs that
were found in most foraminifera-associated MAGs (for a review in NapA function see ref. 52).
All MAGs lack the typical NapB protein forming a complex with NapA. However, *nap* operons

without NapB genes have been reported for other members of Deltaproteobacteria, e.g.

Desulfovibrio desulfuricans (53). Notably, we found that the genome of *Desulfobulbus*

propionicus (Accession: GCA_000186885.1), a member of the order Desulfobacterales that

has been demonstrated to grow based on NO_3^- (54), is lacking a gene for NapB. Hence NO_3^-

respiration in Desulfobacterales in the absence of NapB is possible. We found that the NapA

sequences of all MAGs (and *D. desulfuricans*) include a signal peptide of the twin-arginine

translocation pathway (reviewed in ref. 55) indicating that those proteins could be translocated

across cellular membranes. It is tenable to hypothesize that NapA is secreted by the bacteria

to *Globobulimina* intra-cellular environment. An alternative scenario is nitrate uptake by the

bacterium followed by the reduction to nitrite within the bacterial cellular compartments that

are surrounded by membranes. The nitrite is then released from the bacterial cell and can be

used by the foraminiferal host. The latter scenario may be more beneficial for the bacterium.

Overall, our results suggest that the *Globobulimina*-associated Desulfobacteraceae are able

to reduce NO_3^- to NO_2^-, and thus contribute to the foraminiferal denitrification by performing

the first reaction in the denitrification pathway for which we found no evidence in the

foraminiferal transcriptomes.

Discussion

Our results demonstrate that foraminifera are habitat to bacterial communities that may play

a role in their ability to thrive in oxygen-depleted habitats. A recent study demonstrates the

endosymbiotic contribution to denitrification within a ciliate host (56). However, previous

studies of denitrification in foraminifera argued against the possibility of bacterial contribution

to foraminiferal denitrification (8, 11). Notably, most species in the foraminiferal microbiota are

considered strict anaerobes, hence, when exposed to oxygen the foraminifera may lose their

associated microbiota. Metagenomics sequencing of samples frozen directly after sampling is

thus an important source of information on the composition and function of the foraminiferal

microbiome.

Our results indicate that Desulfobacteracea members of the foraminiferal microbiota

can utilize the NO_3^- storage accumulated by their host. Whether the NO_3^- reduction by bacteria

is beneficial to the foraminifera remains, however, an open question. The bacteria may use

the NO_3^- it for their own respiratory processes or to build up organic compounds via the

assimilation of NO_3^-. For example, it has been suggested that a proportion of NO_3^- taken up

by the foraminiferal species *Ammonia beccari* was used for amino acid synthesis, probably by

resident bacteria (21). Previously we speculated about a foraminiferal sulfite reductase

homolog performing the conversion of NO_3^- (11). Yet, considering the absence of foraminiferal

homologs to NO_3^- reductase and the presence of NapA in all sampled Desulfobacteraceae

MAGs, it is tenable to hypothesize that the reduction of NO_3^- to NO_2^- in foraminifera is
performed by resident bacteria.

An association between foraminifera and Desulfobacteracea has been previously
reported. For example, the presence of a putative Deltaproteobacterium (Desulfobacteracea)
was previously described for *Virgulinella fragilis* (20), a foraminiferal species that we here
predict to be a denitrifying species (Figure 3B). Furthermore, foraminifera from the genera
Ammonia, *Elphidium* and *Haynesina* that are not expected to denitrify were found to harbor
members of Desulfobacterales bacteria (25). Hence, the interaction between foraminifera and
Desulfobacteracea may involve alternative nutritional currencies. Previous studies refer to the
role of Desulfobacteracea in sulphur cycling and carbon/nutrient acquisition (20, 21, 25).
Especially the carbon fixation capabilities via Wood-Ljungdahl pathway would be highly
beneficial for the heterotrophic lifestyle of foraminifera, similarly to other symbioses that
involve fixed-carbon as a nutritional currency [e.g., as in deep-sea mussels (57) or sponges
(49)].

The radiation of early foraminiferal species has been estimated to occur between 690
and 1,150 million years ago (mya), while the first primitive rotaliids appeared in the late
Permian around 260 mya (58, 59). Most of the rotaliids superfamilies diverged in the mid to
late Triassic between approximately 240 and 200 mya (29, 60, 61) and the majority of extant
species diverged during the Miocene (23.8 - 5.5 mya) (62). Our phylogenetic analyses suggest
that clades I and III diversified from within clade II, while the LCA of clade I probably originated
from within clade III (Figures 2 & 3). Although most species reported to denitrify were reported
from the Peruvian OMZ and Sweden, representatives of corresponding genera are found all
over the world (Suppl. Table S5). Therefore, most extant rotaliids likely diversified from a
denitrifying LCA far back in time. The origin of foraminiferal denitrification within or after
diversification from members of clade II may coincide with the rapid increase of fossil records
at the onset of the late Cretaceous ~100 mya (60, 61). Our results therefore indicate that
eukaryotic denitrification by rotaliids emerged late in foraminiferal evolution, possibly during
the worldwide Cretaceous Ocean Anoxic Events (63, 64).

Considering the high conservation of nitrate transporters (Figure 1) and the
observation of nitrate storage in many divergent species (9), it is tenable to speculate that
foraminifera had the mechanisms for nitrate import and storage long before they evolved the
ability to denitrify. NO_3^- transporters are ancient eukaryotic enzymes that play a role in the
NO_3^- assimilation machinery and they are encoded in genomes of photoautotrophs such as
plants and saprotrophs like fungi (65, 66). In contrast to bacteria, foraminifera (as most
heterotrophs) are unable to perform NO_3^- assimilation on their own. Thus, NO_3^- accumulated
by the foraminiferal hosts could have fuelled bacterial NO_3^- metabolism of associated bacteria

in exchange for organic compounds. We propose that denitrification by symbiotic bacteria was
indeed the ancestral state of denitrification in foraminifera, similarly to gromiids (26), which
share a common evolutionary origin with foraminifera (67). The finding of bacterial-like
denitrification genes in rotaliids furthermore suggests that foraminifera may have acquired
those genes from bacteria (11). Thus, a rare gene acquisition from a foraminifera-associated,
denitrifying bacterium could have been at the origin of foraminiferal (eukaryotic) denitrification.
Notably, the rotaliids denitrification gene set is incomplete and it varies among the species
sampled here (i.e., *N. auris, B.plicata* or even *Rosalina* sp.). Species reported to release N_2O
instead of N_2 (9) further support our observation. Thus, the presence of a partial denitrification
pathway in rotaliids as well as in their resident Desulfobacteracea may suggest that the
acquisition of denitrification ability in foraminifera occurred in multiple stages and is not yet
complete.
400

Methods

Foraminifera sampling

Samples were collected off the Peruvian coast in the 2017 Austral winter (R/V Meteor M137)
as described by Glock et al. 2019 (10). Briefly, sediment samples were taken with a video-
guided sediment multiple corer (MUC) containing 6 liners along a depth transect at 12°S. The
top 1 to 3 cm of the sediment cores were sampled and immediately wet sieved with surface
water through staked sieves with mesh size of 2000µm to 63µm to retrieve benthic
foraminifera. The foraminifera were rinsed in sterile seawater obtained by filtering core-
overlying seawater with a sterile bottle top filtration system (Durapore filter, 0.2 µm) and a
vacuum pump within 40 minutes of MUC arrival on deck. Focal species were manually picked
using a stereomicroscope. These were characterised morphologically according to the
literature (68-71). Up to 160 individuals (classified into focal species) were pooled in one cryo
vial (2ml, RNAse free) and flash frozen in liquid nitrogen.

Microscopy

Individual foraminifers were dehydrated in a graded ethanol series (70 %, 80 %, 90 %, 96 %
and two times 100 %; 15 min each), air-dried at 20 °C for 12 h in a desiccator and mounted
on aluminium stubs (PLANO GmbH) using conductive and adhesive carbon pads (PLANO
GmbH). Subsequently, the preparations were sputter-coated with a 10-nm-thick gold-
palladium (80/20) layer using a high vacuum sputter coater Leica EM SCD500 (Leica
Microsystems GmbH) and visualised with a Hitachi S-4800 field emission scanning electron
microscope (Hitachi High-Technologies Corporation) at an acceleration voltage of 3 kV and
an emission current of 10 mA applying a combination of the upper detector and the lower

detector. The *B. costata* specimen was taken from previous samplings by the General

Direction of Research in Oceanography and Climate Change IMARPE (71). The

corresponding sample was air-dried at 35–38 °C, sputter-coated with gold and visualised with

an FEI Inspect S50 scanning electron microscope (Thermo Fisher Scientific Inc.) at an

acceleration voltage of 3 kV.

Nucleic acid extraction, and sequencing

Genomic DNA and total RNA from different biological samples were extracted using either the

DNeasy® Plant Mini kit (QIAGEN) for genomics (DNA) or simultaneously purified using

InnuPREP DNA/RNA Mini Kit (analytikjena) for transcriptomics (RNA). Foraminiferal cells in

each samples (ca. 160 individuals) were disrupted by pestle-crashing on ice after immersion

of the containing cryo-vial in liquid nitrogen. Samples for genomics were treated with lysozyme

(200µl of 10mg/ml TE) and proteinase K (1mg/100µl). Samples for transcriptomics samples

were treated only with lysozyme (6 ul of 20mg/ml TE) before additional crashing and 2-minute

incubation was performed. After disruption and initial lysis, manufacturer protocols for the

respective nucleic acid extraction were followed with modifications to the elution volumes (2 x

50µl for DNA and 2 x 25µl for RNA). Libraries for genomics were produced after DNA

fragmentation (Covaris target 400, intensity 5, duty cycle 5% cycles per burst 200, 55sec

treatment time) using NEBNext® Ultra II DNA Library Prep Kit for Illumina®. Whereas,

transcriptomic libraries were produced using NEBNext® Ultra RNA Library Prep Kit for

Illumina® (NEB) with mRNA isolation performed with poly-A mRNA beads. All libraries were

produced in duplicates from two different sets of pooled individuals for each species and were

produced without protocol interruption. Before sequencing, each library was quantified with a

Qubit® fluorometer (Invitrogen by Life TechnologiesTM) and qualified using a TapeStation

(Agilent technology). The libraries were sequenced paired-end (2 x 150 bp) on an Illumina®

HiSeq 4000 platform.

Sequencing datasets of foraminifera

Sequencing resulted in 4.6 billion paired-end reads covering 1.4 tera bases in total. This

includes transcriptome and metagenome datasets (BioProject accessions PRJNA494828 &

PRJNA503328). Reads were quality-checked by FastQC ver. 0.11.5

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc; Aug 2016). Filtering and trimming

of reads was performed using Trimmomatic (72) ver. 0.36 (Parameters:

ILLUMINACLIP:primers.fa:2:30:10 LEADING:5 TRAILING:5 SLIDINGWINDOW:4:5

MINLEN:21; the file 'primers.fa' contained adaptor and contaminant sequences provided by

Trimmomatic and FastQC). Processed reads from transcriptomes of the two samples per

species were assembled into transcript contigs using SPAdes (73) ver. 3.11.1 ("--rna" option).

Protein sequences were translated from transcripts as open reading frames (ORFs) using

TransDecoder (74) ver. v5.0.2 (LongOrfs; "-m 30" option). Final protein names consist of the

contig ids followed by the sequence positions covered by the CDS and an indicator for the

forwards (+) or reverse (-) strand. Transcript abundance of individual transcriptome datasets

are referring to Transcripts Per Million (TPM) determined by the Trinity pipeline (74) 2.4.0

(Trinity script 'align_and_estimate_abundance.pl') via RSEM (75) ver. 1.2.30 and Bowtie (76)

2.1.0 using paired-end reads. Additional raw sequencing reads were obtained from the Marine

Microbial Transcriptome Project (MMETSP; SRA accessions: *Rosalina* sp., SRR1296887;

Sorites sp., SRR1296734; *Ammonia* sp., SRR1300434; *Elphidium margaritaceum*,

SRR1300475) and processed as described above. We found that the assembly obtained for

Sorites sp. contained a high proportion of sequences likely originating from an algal species

(Probably *Symbiodinium* sp.), which was removed by applying a binning approach considering

only contigs with GC-content ≤38%. Finally, we included data for *Globobulimina* spp. sampled

in Sweden from our previous study (Transcriptome data: GloT15) (11) and proteins annotated

on the genome assembly of *Reticulomyxa filosa* (NCBI accession: GCA_000512085.1).

Species phylogenies and rooting

Transcriptome completeness and heterogeneity was determined by assessing genome

completeness via Benchmarking Universal Single-Copy Orthologs (77) (BUSCO v3; lineage

'eukaryota') method. Orthologous proteins determined as Complete (or Duplicated) by the

BUSCO analysis were merged into protein clusters to study phylogenetic relationships among

the species. In case of duplicated BUSCOs in metatranscriptomes, one representative was

inferred based on sequence similarity. Therefore, all orthologs retrieved for the same

metatranscriptome were compared to all members of the BUSCO protein cluster by global

pairwise alignments using needle (EMBOSS tools version 6.6.0) (78). The ortholog with the

highest median sequence similarity over all comparisons was picked as the representative

sequence. Multiple sequence alignments used in the current study were obtained using

MAFFT (79) (Version: 7; parameter: 'linsi') and the phylogenetic trees were reconstructed

using IQTREE (80) (Version: 1.5.5 or 1.6.9; default parameters; note that ModelFinder is

enabled by default). Phylogenies were reconstructed for the individual BUSCO clusters. Due

to their morphological differences from other foraminiferal groups, monothalamids like *R. filosa*

have been previously used as an outgroup for phylogenetic studies of foraminifera (29). Since

the choice of a distantly related outgroup may lead to erroneous rooted topology due to long

branch attraction (81), we further tested the robustness of the root position independent from

outgroup species. Root positions were determined using MAD (28) (Parameters: '-bsnn'). The

support in each root splits is calculated as the proportion of gene trees where the root position

was inferred as such (28). For the overall species trees, multigene phylogenies were

reconstructed using IQTREE (parameter: '-spp') considering all protein cluster alignments of
either foraminifera or Rotaliida.

Reference 18S rRNA sequences used for the large-scale phylogenies (Suppl. Table
S5) were obtained from NCBI, MMETSP, SILVA (82), the foram barcoding project
(http://forambarcoding.unige.ch/) and the Planktonic Foraminifera Ribosomal Reference
database (83) (PFR^2). These sequences were used as database sequences to identify 18S
sequences from Peruvian species transcriptomes based on BLAST (84) (version: 2.2.28+;
options: '-task blastn -evalue 1e^{-10}') searches. Among the hits retrieved we searched for a
single representative transcript contig per transcriptome assembly. The transcripts with the
highest product of sequence length and sequencing depth (i.e., coverage given by SPAdes)
were determined as representative sequences for most of the focal species. However, for *B.*
costata, *V. inflata*, *G. pacifica* and *N. auris* different representative sequences were chosen
by giving sequence information (i.e., longer sequences) a higher weight than sequencing
depth. To obtain 18S phylogenies first the reference sequences were aligned with MAFFT and
subsequently representatives 18S rRNA sequences of the Peruvian transcriptomes were
added (MAFFT options: '--addfragments --adjustdirection') to the alignment followed by the
tree reconstruction using IQTREE.

Identification of foraminiferal denitrification proteins
To identify homologs to enzymes in the denitrification pathway, we used a similar approach
as previously established in Woehle et al. 2018 (11) using the corresponding protein database
expanded by the protein sequences identified for *Globobulimina* (11). The search for
denitrification enzymes homologs in the transcriptome assemblies was performed with
BLASTP (parameter: '-max_target_seqs 1000000 e-value 1e-5'). Protein sequences of hits
with query coverage ≥ 40% and sequence identity ≥ 20% were extracted to obtain a first set
of homologs. We further applied a cutoff to discard lowly represented transcripts with TPM <
2 in at least one of the two replicates sequenced. With the resulting protein set, we reiterated
searches in the non-redundant NCBI protein (NR; version May 2018) and the RefSeq 88
database. (85) using diamond v0.9.22 applying the '--more-sensitive' option. First best hit
sequences per query were obtained and clustered with CD-HIT 4.6 (86) (option: "-c 0.98") to
reduce sequence redundancy. All obtained protein sequences of a given enzyme were aligned
with MAFFT and phylogenetic trees were reconstructed with IQTREE (parameters: '-bb 1000
-alrt 1000'). The trees were rooted using an outgroup, if available, or MAD.

Metagenomics processing
For the visualisation of metagenomic composition, trimmed paired-end reads from Peruvian
species samples and the *Globobulimina* 'Ambient' samples (11) were subsampled via BBMap

(version 36.84; https://sourceforge.net/projects/bbmap/; 'reformat.sh' script; parameters:
'samplereadstarget=10000000 addslash=t'). The resulting reads were mapped against the NR
database with ac-diamond (87) and classified using MEGAN6 (88) (Version 6.15.0; options '-
a2t   prot_acc2tax-Nov2018X1.abin   -a2eggnog   acc2eggnog-Oct2016X.abin   -a2seed
acc2seed-May2015XX.abin'). MEGAN6 was used for assessment of metagenomics
communities and visual representation (See Figure 4) that were classified up to the order rank
level with different taxon sets (e.g., all nodes or only the bacterial subtree). Metagenome
assemblies were obtained by combining all trimmed reads per focal species using MEGAHIT
(89) (Version 1.1.3; default settings). Individual bacterial genomes were obtained using the
binning approach implemented with MaxBin2 (90) (Version 2.2.4; parameter: '-
min_contig_length 500') using coverage information of the two samples per species of
foraminifera. Binning statistics were assessed using CheckM (91) (Version 1.0.11); a
threshold for completeness of at least 80% and contamination of maximum 20% was applied
to classify bacterial bins as draft genomes. The protein sequences obtained via checkM were
used for further classifications. The average nucleotide identity (ANI) was calculated using a
perl script obtained from https://github.com/chjp.
For the taxonomic assignment of genome bins, we used the diamond tool to find first
best hits for each protein of a genome bin against the NR database (option: '-k 10'; e-value ≤
1e-10) and retrieved the corresponding taxonomic assignments (as we previously described
in ref. 11). For each bin, identical taxonomic hierarchies were counted with the genus being
the lowest rank considered, sorted accordingly and stepwise searched for the lowest
taxonomic rank supporting >50% of protein bin hits starting with the most abundant taxonomy.
'Environmental samples' and 'Cellular organisms' were not considered. For the phylogenetic
reconstruction of Desulfobacteracea we determined protein families as protein clusters using
sequences of Desulfobacteracea bins associated with *Globobulimina* and additional
Desulfobacteracea genomes downloaded from NCBI. First, we determined reciprocal best
BLAST hit pairs (rBBH; parameter: '-evalue 1e-5') between all the Desulfobacteracea protein-
coding sequences (92). Then rBBH pairs were globally aligned with the needle tool and pairs
with ≥30% identical amino acids were sorted into clusters using the Markov clustering
algorithm (93) (MCL; version 12-135). The 33 resulting protein clusters that contained a single
copy for each of the Desulfobacteraceae strains (i.e., universal single-copy clusters) were
aligned with MAFFT. The resulting alignments were concatenated and a splits network was
reconstructed with SplitsTree (94) (version 4.15.1). Prediction of protein function for bacterial
genomes were obtained using the KEGG Automatic Annotation Server (95) (KAAS; BBH
method via BLAST and the following species set: hsa, dme, ath, sce, pfa, eco, sty, hin, pae,
nme, hpy, rpr, mlo, bsu, sau, lla, spn, cac, mge, mtu, ctr, bbu, syn, aae, mja, afu, pho, ape,
geo, dvu, dat, dpr, dol, dal, dak, dps, drt, dba, dao, dbr). Global pairwise identities for NapA

protein sequences were inferred with needle. Signal peptides were predicted using the
SignalP-5.0 webservice for gram-negative bacteria (96).

Data availability

Sequencing reads are deposited at the single read archive (SRA) accessions SRR8144071
to SRR8144090 and SRR7971179 to SRR7971198. The assemblies are available at the
transcriptome sequencing archive (TSA; See Suppl. Table 1 for accessions) and as whole
genome shotgun (WGS) projects (See Suppl. Table 7 for accessions). All other information
on accessing data analysed in this study is included in the manuscript or in the supplemental
information.

Acknowledgments

We thank Devani Romero Picazo and Anne Kupczok for critical comments on the manuscript.
We gratefully acknowledge the scientific party and crew of the R/V Meteor cruise M137 as
well as Asmus Petersen and Matthias Türk for their support at sea. Samples from Peru were
obtained according to Peruvian access and benefit sharing regulations. We thank the Nagoya
officer of Kiel University - Dr. Scarlett Sett - for her support of our research. We further thank
Natalia Bernabe Lopez for her assistance in collecting metadata for the 18S sequences. The
micrograph of *B. costata* was possible due to the General Direction of Research in
Oceanography and Climate Change (IMARPE). Genome sequencing was performed in the
Centre for Genome Analysis Kiel that is funded by the German Research Foundation (DFG).
The study was supported by the German Research Foundation (DFG) via the SFB 754 on
Climate–Biogeochemistry Interactions in the Tropical Ocean, the cluster of excellence The
Future Ocean, and the European Research Council (Grant No. 281357 awarded to TD). C.W.
was supported by the Kiel Life Science (KLS) Young Scientist Programme and would like to
thank the Max Planck-Genome-centre Cologne for their support in data analysis.

Author contributions

C.W., A.-S.R., N.G., T.W., J.W., D.R., J.S. & T.D. designed the research strategy and
performed sampling. C.W. analysed the transcriptomes and metagenomes. A.-S.R. performed
the experimental lab work. C.H. constructed incubation chambers. D.R. produced the
scanning electron micrograph of *B. costata*. J.M. produced all other scanning electron
micrographs, which was supported by S.N.G. C.W. and T.D. wrote the manuscript with input
from all co-authors.

Declaration of Interests

The authors declare no competing interests.

References:

1.   Cox AN (2015) *Allen's Astrophysical Quantities* ed Cox AN (Springer, New York, NY)
609   doi:10.1007/978-1-4612-1186-0.

610   2.   Thamdrup B (2012) New Pathways and Processes in the Global Nitrogen Cycle. *Annu*
611   *Rev Ecol Evol Syst* 43(1):407–428.

612   3.   Gruber N (2004) The Dynamiics of the Marine Nitrogen Cycle and its Influence on
613   Atmospheric CO_2 Variations. *The Ocean Carbon Cycle and Climate* (Springer,
614   Dordrecht, Dordrecht), pp 97–148. 2nd Ed.

615   4.   Gruber N, Galloway JN (2008) An Earth-system perspective of the global nitrogen
616   cycle. *Nature* 451(7176):293–296.

617   5.   Lam P, et al. (2009) Revising the nitrogen cycle in the Peruvian oxygen minimum
618   zone. *Proc Natl Acad Sci USA* 106(12):4752–4757.

619   6.   Philippot L (2002) Denitrifying genes in bacterial and Archaeal genomes. *Biochim*
620   *Biophys Acta* 1577(3):355–376.

621   7.   Shoun H, Kim DH, Uchiyama H, Sugiyama J (1992) Denitrification by fungi. *FEMS*
622   *Microbiology Letters* 73(3):277–281.

623   8.   Risgaard-Petersen N, et al. (2006) Evidence for complete denitrification in a benthic
624   foraminifer. *Nature* 443(7107):93–96.

625   9.   Piña-Ochoa E, et al. (2010) Widespread occurrence of nitrate storage and
626   denitrification among Foraminifera and Gromiida. *Proc Natl Acad Sci USA*
627   107(3):1148–1153.

628   10.   Glock N, et al. (2019) Metabolic preference of nitrate over oxygen as an electron
629   acceptor in foraminifera from the Peruvian oxygen minimum zone. *Proc Natl Acad Sci*
630   *USA* 116(8):2860–2865.

631   11.   Woehle C, et al. (2018) A Novel Eukaryotic Denitrification Pathway in Foraminifera.
632   *Curr Biol* 28(16):2536–2543.e5.

633   12.   Glock N, et al. (2013) The role of benthic foraminifera in the benthic nitrogen cycle of
634   the Peruvian oxygen minimum zone. *Biogeosciences* 10(7):4767–4783.

635   13.   Schweizer M, Pawlowski J, Kouwenhoven TJ, Guiard J, van der Zwaan B (2008)
636   Molecular phylogeny of Rotaliida (Foraminifera) based on complete small subunit
637   rDNA sequences. *Marine Micropaleontology* 66(3-4):233–246.

638   14.   Murray JW (2006) *Ecology and applications of benthic foraminifera* (Cambridge
639   University Press).

640   15.   LeKieffre C, et al. (2017) Surviving anoxia in marine sediments: The metabolic
641   response of ubiquitous benthic foraminifera (Ammonia tepida). *PLoS One*
642   12(5):e0177604.

16.   Bernhard JM, et al. (2012) Potential importance of physiologically diverse benthic
foraminifera in sedimentary nitrate storage and respiration. *J Geophys Res*
117(G3):G03002.

17.   Gomaa F, et al. (2021) Multiple integrated metabolic strategies allow foraminiferan
protists to thrive in anoxic marine sediments. *Sci Adv* 7(22).
doi:10.1126/sciadv.abf1586.

18.   Orsi WD, et al. (2020) Anaerobic metabolism of Foraminifera thriving below the
seafloor. *ISME J* 14(10):2580–2594.

19.   Zilber Rosenberg I, Rosenberg E (2008) Role of microorganisms in the evolution of
animals and plants: the hologenome theory of evolution. *FEMS Microbiology Reviews*
32(5):723–735.

20.   Tsuchiya M, et al. (2015) Cytologic and genetic characteristics of endobiotic bacteria
and kleptoplasts of *Virgulinella fragilis* (Foraminifera). *J Eukaryotic Microbiology*
62(4):454–469.

21.   Nomaki H, et al. (2014) Nitrate uptake by foraminifera and use in conjunction with
endobionts under anoxic conditions. *Limnology and Oceanography* 59(6):1879–1888.

22.   Bernhard JM, Martin JB, Rathburn AE (2010) Combined carbonate carbon isotopic
and cellular ultrastructural studies of individual benthic foraminifera: 2. Toward an
understanding of apparent disequilibrium in hydrocarbon seeps. *Paleoceanography*
25(4):PA4206.

23.   Bernhard JM, Habura A, Bowser SS (2006) An endobiont‐bearing allogromiid from
the Santa Barbara Basin: Implications for the early diversification of foraminifera. *J*
Geophys Res 111(G3):399.

24.   Bernhard JM (2003) Potential symbionts in bathyal foraminifera. *Science*
299(5608):861–861.

25.   Salonen IS, Chronopoulou PM, Bird C, Reichart GJ, Koho KA (2019) Enrichment of
intracellular sulphur cycle –associated bacteria in intertidal benthic foraminifera
revealed by 16S and aprA gene analysis. *Sci Rep*:1–12.

26.   Høgslund S, Cedhagen T, Bowser SS, Risgaard-Petersen N (2017) Sinks and
Sources of Intracellular Nitrate in Gromiids. *Front Microbiol* 8:617.

27.   Bernhard JM, Edgcomb VP, Casciotti KL, McIlvin MR, Beaudoin DJ (2012)
Denitrification likely catalyzed by endobionts in an allogromiid foraminifer. *ISME J*
6(5):951–960.

28.   Tria FDK, Landan G, Dagan T (2017) Phylogenetic rooting using minimal ancestor
deviation. *Nat Ecol Evol* 1:0193.

29.   Pawlowski J, et al. (2003) The evolution of early Foraminifera. *Proc Natl Acad Sci*
USA 100(20):11494–11498.

30.   Ertan KT, Hemleben V, Hemleben C (2004) Molecular evolution of some selected
benthic foraminifera as inferred from sequences of the small subunit ribosomal DNA.
Marine Micropaleontology 53(3-4):367–388.

31.   Mackensen A, Schmiedl G, Harloff J, Giese M (1995) Deep-Sea Foraminifera in the
684   South Atlantic Ocean: Ecology and Assemblage Generation. *Micropaleontology*
685   41(4):342.

686   32.   Rathbun AE, Willingham J, Ziebis W, Burkett AM, Corliss BH (2018) A New biological
687   proxy for deep-sea paleo-oxygen: Pores of epifaunal benthic foraminifera. *Sci Rep*
688   8(1):9456–8.

689   33.   Hoogakker BAA, et al. (2018) Glacial expansion of oxygen-depleted seawater in the
690   eastern tropical Pacific. *Nature* 562(7727):410–413.

691   34.   Widdel F (1987) New types of acetate-oxidizing, sulfate-reducing Desulfobacter
692   species, *D. hydrogenophilus* sp. nov., *D. latus* sp. nov., and *D. curvatus* sp. nov. *Arch*
693   *Microbiol* 148(4):286–291.

694   35.   Lage OM, Bondoso J (2014) Planctomycetes and macroalgae, a striking association.
695   *Front Microbiol* 5(5):267.

696   36.   Oshiki M, Satoh H, Okabe S (2016) Ecology and physiology of anaerobic ammonium
697   oxidizing bacteria. *Environ Microbiol* 18(9):2784–2796.

698   37.   Noguchi T, et al. (1987) *Vibrio alginolyticus*, a tetrodotoxin-producing bacterium, in the
699   intestines of the fish *Fugu vermicularis vermicularis*. *Marine Biology* 94(4):625–630.

700   38.   McFall-Ngai M (2014) Divining the essence of symbiosis: insights from the squid-
701   vibrio model. *PLoS Biol* 12(2):e1001783.

702   39.   Ben-Haim Y, Zicherman-Keren M, Rosenberg E (2003) Temperature-regulated
703   bleaching and lysis of the coral *Pocillopora damicornis* by the novel pathogen *Vibrio*
704   *coralliilyticus*. *Appl Environ Microbiol* 69(7):4236–4242.

705   40.   López-Pérez M, Rodriguez-Valera F (2014) The Family Alteromonadaceae. *The*
706   *Prokaryotes: Gammaproteobacteria*, The Prokaryotes: Gammaproteobacteria. eds
707   Rosenberg E, Delong EF, Lory S, Stackebrandt E, Thompson F (Springer Berlin
708   Heidelberg, Berlin, Heidelberg), pp 69–92.

709   41.   Ivanova EP, Flavier S, Christen R (2004) Phylogenetic relationships among marine
710   Alteromonas-like proteobacteria: emended description of the family
711   Alteromonadaceae and proposal of Pseudoalteromonadaceae fam. nov.,
712   Colwelliaceae fam. nov., Shewanellaceae fam. nov., Moritellaceae fam. nov.,
713   Ferrimonadaceae fam. nov., Idiomarinaceae fam. nov. and Psychromonadaceae fam.
714   nov. *Int J Syst Evol Microbiol* 54(Pt 5):1773–1788.

715   42.   Simon M, et al. (2017) Phylogenomics of Rhodobacteraceae reveals evolutionary
716   adaptation to marine and non-marine habitats. *ISME J* 11(6):1483–1499.

717   43.   Newton RJ, et al. (2010) Genome characteristics of a generalist marine bacterial
718   lineage. *ISME J* 4(6):784–798.

719   44.   Buchan A, Gonzalez JM, Moran MA (2005) Overview of the marine roseobacter
720   lineage. *Appl Environ Microbiol* 71(10):5665–5677.

721   45.   Hjelm M, et al. (2004) Selection and identification of autochthonous potential probiotic
722   bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. *Syst Appl Microbiol*
723   27(3):360–371.

724   46.   Webster NS, Negri AP, Munro MMHG, Battershill CN (2004) Diverse microbial
725   communities inhabit Antarctic sponges. *Environ Microbiol* 6(3):288–300.

726   47.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S (2018) High
727   throughput ANI analysis of 90K prokaryotic genomes reveals clear species
728   boundaries. *Nat Commun* 9(1):5114–8.

729   48.   Dubilier N, et al. (2001) Endosymbiotic sulphate-reducing and sulphide-oxidizing
730   bacteria in an oligochaete worm. *Nature* 411(6835):298–302.

731   49.   Kamke J, et al. (2013) Single-cell genomics reveals complex carbohydrate
732   degradation patterns in poribacterial symbionts of marine sponges. *ISME J*
733   7(12):2287–2300.

734   50.   Wein T, et al. (2019) Currency, Exchange, and Inheritance in the Evolution of
735   Symbiosis. *Trends Microbiol* 27(10):836–849.

736   51.   Strittmatter AW, et al. (2009) Genome sequence of *Desulfobacterium autotrophicum*
737   HRM2, a marine sulfate reducer oxidizing organic carbon completely to carbon
738   dioxide. *Environ Microbiol* 11(5):1038–1055.

739   52.   González PJ, Correia C, Moura I, Brondino CD, Moura JJG (2006) Bacterial nitrate
740   reductases: Molecular and biological aspects of nitrate reduction. *J Inorg Biochem*
741   100(5-6):1015–1023.

742   53.   Dias JM, et al. (1999) Crystal structure of the first dissimilatory nitrate reductase at 1.9
743   A solved by MAD methods. *Structure* 7(1):65–79.

744   54.   Widdel F, Pfennig N (1982) Studies on dissimilatory sulfate-reducing bacteria that
745   decompose fatty acids II. Incomplete oxidation of propionate by *Desulfobulbus*
746   *propionicus* gen. nov., sp. nov. *Arch Microbiol* 131(4):360–365.

747   55.   Berks BC, Ferguson SJ, Moir JW, Richardson DJ (1995) Enzymes and associated
748   electron transport systems that catalyse the respiratory reduction of nitrogen oxides
749   and oxyanions. *Biochim Biophys Acta* 1232(3):97–173.

750   56.   Graf JS, et al. (2021) Anaerobic endosymbiont generates energy for ciliate host by
751   denitrification. *Nature* 591(7850):445–450.

752   57.   Duperron S, et al. (2006) A dual symbiosis shared by two mussel species,
753   *Bathymodiolus azoricus* and *Bathymodiolus puteoserpentis* (Bivalvia: Mytilidae), from
754   hydrothermal vents along the northern Mid-Atlantic Ridge. *Environ Microbiol*
755   8(8):1441–1447.

756   58.   Maklay M (1964) Novye Rannekamennougolnye Arkhedistsidy. *Min Geol Okhrana*
757   *Nedr USSR* 1:158–159.

758   59.   Crespin I (1958) Permian Foraminifera of Australia. *Bur Miner Resour Aust Bull* 48:1–
759   207.

760   60.   Tappan H, Loeblich AR (1988) Foraminiferal evolution, diversification, and extinction.
761   *Journal of Paleontology* 62(5):695–714.

61. Loeblich AR, Tappan H (1964) *Treatise on Invertebrate Paleontology. Part C. Protista*
2. Sarcodina chiefly "thecamoebians" and Foraminiferida. (Geological Society of
America New York/University of Kansas Lawrence, New York/Lawrence).

62. Kucera M, Schönfeld J (2007) The origin of modern oceanic foraminiferal faunas and
Neogene climate change. *The Micropalaeontology Society Special Publications*:409–
425.

63. Schlanger SO, Jenkyns HC (1976) Cretaceous oceanic anoxic events: causes and
consequences. *Geologie en Mijnbouw* 55(3-4):179–184.

64. Jenkyns HC (1980) Cretaceous anoxic events: from continents to oceans. *Journal of*
the Geological Society 137(2):171–188.

65. Forde BG (2000) Nitrate transporters in plants: structure, function and regulation.
Biochim Biophys Acta 1465(1-2):219–235.

66. Pao SS, Paulsen IT, Saier MH (1998) Major facilitator superfamily. *Microbiol Mol Biol*
Rev 62(1):1–34.

67. Sierra R, et al. (2013) Deep relationships of Rhizaria revealed by phylogenomics: A
farewell to Haeckel's Radiolaria. *Mol Phylogenet Evol* 67(1):53–59.

68. Figueroa S, Marchant M, Giglio S, Ramírez M (2005) Foraminíferos bentónicos
rotalínidos del centro sur de Chile (36°S - 44°S). *Gayana* 69(2):329–363.

69. Erdem Z (2016) Reconstruction of past bottom water conditions of the Peruvian
Oxygen Minimum Zone (OMZ) for the last 22,000 years and the benthic foraminiferal
response to (de)oxygenation. *PhD thesis, Kiel University*.

70. Mallon J (2012) Benthic Foraminifera of the Peruvian and Ecuadorian Continental
Margin. *PhD thesis, Kiel University*.

71. Cardich J, et al. (2015) Calcareous benthic foraminifera from the upper central
Peruvian margin: control of the assemblage by pore water redox and sedimentary
organic matter. *Mar Ecol Prog Ser* 535:63–87.

72. Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30(15):2114–2120.

73. Nurk S, et al. (2013) Assembling single-cell genomes and mini-metagenomes from
chimeric MDA products. *J Comput Biol* 20(10):714–737.

74. Haas BJ, et al. (2013) De novo transcript sequence reconstruction from RNA-seq
using the Trinity platform for reference generation and analysis. *Nat Protoc*
8(8):1494–1512.

75. Li B, Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* 12:323.

76. Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat*
Methods 9(4):357–359.

77. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM (2015)
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics* 31(19):3210–3212.

78. Rice P, Longden I, Bleasby A (2000) EMBOSS: the European Molecular Biology
Open Software Suite. *Trends Genet* 16(6):276–277.

79. Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version
7: improvements in performance and usability. *Mol Biol Evol* 30(4):772–780.

80. Nguyen L-T, Schmidt HA, Haeseler von A, Minh BQ (2015) IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol*
Evol 32(1):268–274.

81. Felsenstein J (1978) Cases in which parsimony or compatibility methods will be
positively misleading. *Systematic Zoology* 27:401–410.

82. Yilmaz P, et al. (2014) The SILVA and "All-species Living Tree Project (LTP)"
taxonomic frameworks. *Nucleic Acids Res* 42(Database issue):D643–8.

83. Morard R, et al. (2015) PFR2: a curated database of planktonic foraminifera 18S
ribosomal DNA as a resource for studies of plankton ecology, biogeography and
evolution. *Molecular Ecology Resources* 15(6):1472–1485.

84. Altschul SF, et al. (1997) Gapped BLAST and PSI-BLAST: a new generation of
protein database search programs. *Nucleic Acids Res* 25(17):3389–3402.

85. Pruitt KD, Tatusova T, Brown GR, Maglott DR (2012) NCBI Reference Sequences
(RefSeq): current status, new features and genome annotation policy. *Nucleic Acids*
Res 40(Database issue):D130–D135.

86. Li W, Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets
of protein or nucleotide sequences. *Bioinformatics* 22(13):1658–1659.

87. Mai H, et al. (2018) AC-DIAMOND v1: accelerating large-scale DNA-protein
alignment. *Bioinformatics* 34(21):3744–3746.

88. Huson DH, et al. (2016) MEGAN Community Edition - Interactive Exploration and
Analysis of Large-Scale Microbiome Sequencing Data. *PLoS Comput Biol*
12(6):e1004957.

89. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W (2015) MEGAHIT: an ultra-fast single-
node solution for large and complex metagenomics assembly via succinct de Bruijn
graph. *Bioinformatics* 31(10):1674–1676.

90. Wu Y-W, Simmons BA, Singer SW (2016) MaxBin 2.0: an automated binning
algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics*
32(4):605–607.

91. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW (2015) CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* 25(7):1043–1055.

92. Tatusov RL, Koonin EV, Lipman DJ (1997) A genomic perspective on protein families.
Science 278(5338):631–637.

93.   Enright AJ, Van Dongen S, Ouzounis CA (2002) An efficient algorithm for large-scale
840   detection of protein families. *Nucleic Acids Res* 30(7):1575–1584.

841   94.   Huson DH (2005) Application of phylogenetic networks in evolutionary studies. *Mol
842   Biol Evol* 23(2):254–267.

843   95.   Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M (2007) KAAS: an automatic
844   genome annotation and pathway reconstruction server. *Nucleic Acids Res* 35(Web
845   Server issue):W182–5.

846   96.   Nielsen H (2017) Predicting Secretory Proteins with SignalP. *Methods Mol Biol*
847   1611(Chapter 6):59–73.

1 **Supplementary material:**

3 **Suppl. Figure S1. Scanning electron micrographs of sampled species.** Detailed views of

4 the graphs shown in Figure 1A.

B

Bacteria

Tree scale: 1 ⊢————┤

Archaea

Bacteria

Foraminifera

9	**Suppl. Figure S2. Complete phylogenies of NirK, Nor and Nrt homologs.** Phylogenetics
10	trees of the foraminiferal denitrification proteins A) NirK B) Nor and C) Nrt that survived the
11	cutoff together with homologs from public databases. Transcriptome labels contain
12	transcription values of the two replicates (tpm1 & tpm2) as well as their percentile expression
13	rank within corresponding transcriptomes. The labels contain as well the transcript id, region
14	and orientation for individual open reading frames used. Numbers of the branches represent
15	statistical support by ultrafast bootstrap and SH-like approximate likelihood ratio test results
16	(following the '/') each with 1000 replicates. Branch support values of zero are omitted. If only
17	one value is shown it represents the SH-like approximate likelihood ratio test. The background
18	colour coding indicates different taxonomic groups. *Gobobulimina* spp. homologs from Woehle
19	et al. 2018 are shown in bold. Further phylogenetic (sub-)clades, resembling those from
20	Woehle et al. 2018, are highlighted by purple bars. The Nor tree was rooted using two cbb3
21	oxidases as outgroup, while for the Nrt tree representatives of DHA14 and ACS family of MFS
22	transporters as outgroup. The NirK tree was rooted using the MAD method.
23	

24	
25	**Suppl. Figure S3. Foraminifera rooting based on single-protein trees.** A Maximum-
26	likelihood phylogenetic reconstruction of foraminifera species based on 81 eukaryotic protein
27	marker sequences is shown on the left. Parametric bootstrap support is 1000/1000 at all the
28	branches. Right of the phylogeny a split representation of root splits determined via the MAD
29	approach for the 81 single gene trees is shown. Each column represents a root. Root branches
30	are reported as separation of two groups (black and white boxes) indicating for the species
31	found on either side of the root split. The bar graph (top) reports the single gene tree count
32	supporting the corresponding root splits. These were ranked by frequency.
33	

Tree scale: 0.1

Suppl. Figure S4. Uncollapsed 18S phylogeny of foraminifera. Species shown to denitrify

experimentally are highlighted in blue. The only exception is *Stainforthia fusiformis*, where

denitrification activity has been shown for an unspecified species of the same genus. Species

also shown in Figure 2 or Figure S1 are highlighted by asterisks. Bootstrap support values

with 1000 replicates are shown at the branches. The trees were rooted by the clade of

monothalamids containing *R. filosa*. The phylogeny corresponds to the one shown in Figure

3.

Suppl. Table S1. Sequencing statistics for metagenomics & transcriptomics.

Supp. Table S2. List of foraminiferal homologs of denitrification proteins remaining

after applying cutoffs.

Supp. Table S3. List of foraminiferal homologs of denitrification proteins removed by

cutoffs.

Suppl. Table S4. List of marker proteins used for phylogenetic reconstructions.

Suppl. Table S5. 18S sequences including metadata for phylogenetic reconstructions.

Suppl. Table S6. Proportions of family rank classification of foraminifera-associated
microbial communities.
Suppl. Table S7. High quality genomic bins of foraminifera-associated species
communities.
Suppl. Table S8. Metabolic classification of Desulfobacteracea genomes.
55