ABA represses TOR and root meristem activity through nuclear exit of the SnRK1 kinase

Borja Belda-Palazón[1*], Mónica Costa[1,2,3], Tom Beeckman[2,3], Filip Rolland[4], Elena Baena-
González[1*]

[1]Instituto Gulbenkian de Ciência, 2780-156 Oeiras, Portugal and GREEN-IT Bioresources for
Sustainability, ITQB NOVA, 2780-157 Oeiras, Portugal

[2]Department of Plant Biotechnology and Bioinformatics, Ghent University, Technologiepark 71,
B-9052 Ghent, Belgium

[3]VIB-UGent Center for Plant Systems Biology, Technologiepark 71, B-9052 Ghent, Belgium

[4]Laboratory for Molecular Plant Biology, Biology Department, Katholieke Universiteit Leuven,
3001 Heverlee-Leuven, Belgium

*Corresponding authors. Instituto Gulbenkian de Ciência, Rua da Quinta Grande 6, 2780-156
Oeiras, Portugal. Tel.: +351 214464630; Fax: +351 214407970; e-mail:
bbelda@igc.gulbenkian.pt, ebaena@igc.gulbenkian.pt

Keywords: *root apical meristem, growth regulation, abscisic acid, energy signaling, SnRK1, TOR,*
Arabidopsis thaliana

ABSTRACT

The phytohormone abscisic acid (ABA) promotes plant tolerance to major stresses like drought,

partly by modulating plant growth and development. However, the underlying mechanisms are

poorly understood. Here, we show that cell proliferation in the *Arabidopsis thaliana* root meristem

is controlled by the interplay between three kinases, SNF1-RELATED KINASE 2 (SnRK2), the

main driver of ABA signaling, the SnRK1 energy sensor, and the growth-promoting TARGET OF

RAPAMYCIN (TOR) kinase. Under favorable conditions, the SnRK1α1 catalytic subunit is

enriched in the nuclei of root meristematic cells and this is accompanied by normal cell

proliferation and meristem size. Depletion of SnRK2s in a *snrk2.2 snrk2.3* double mutant causes

constitutive cytoplasmic localization of SnRK1α1 and a reduction in meristem size, suggesting

that, under non-stress conditions, SnRK2s enable growth by retaining SnRK1α1 in the nucleus. In

response to elevated ABA levels, SnRK1α1 translocates to the cytoplasm and this is accompanied

by inhibition of TOR, decreased cell proliferation and meristem size. Blocking nuclear export with

leptomycin B abrogates ABA-driven SnRK1α1 relocalization to the cytoplasm and the inhibition

of TOR. Fusion of SnRK1α1 to an SV40 nuclear localization signal leads to defective TOR

repression in response to ABA, demonstrating that SnRK1α1 nuclear exit is a premise for this

repression. Finally, the SnRK2-dependent changes in SnRK1α1 subcellular localization are

specific to the proliferation zone of the meristem, underscoring the relevance of this mechanism

for growth regulation.

The phytohormone abscisic acid (ABA) plays major roles in plant stress responses. ABA signals are transduced through a well-established pathway whose main effectors in *Arabidopsis* are SNF1-RELATED PROTEIN KINASE 2.2 (SnRK2.2), SnRK2.3, and SnRK2.6 (1). ABA promotes plant adaptation partly by modifying developmental programs, having a major impact e.g. on root architecture (2). ABA modulates primary root (PR) and lateral root (LR) growth through interactions with other hormones, ultimately affecting cell division and elongation by poorly understood mechanisms (2). We recently uncovered an intimate connection between ABA and SnRK1 signalling that is crucial for shaping root architecture in a TARGET OF RAPAMYCIN (TOR)-dependent manner (3). TOR is a protein kinase complex that promotes cell proliferation (4). It is highly enriched in the meristems and its inactivation causes reduced root meristem size and defective PR growth (4). SnRK1 is an evolutionarily conserved protein kinase complex that is activated when energy levels decline during stress, conferring protection partly by limiting growth (5). SnRK1 is also activated by ABA, enabling plants to repress growth when water is scarce (3). Under favorable conditions, the main catalytic subunit SnRK1α1 is sequestered by SnRK2-containing repressor complexes, allowing TOR to be active. In response to ABA, these complexes dissociate, releasing SnRK2 and SnRK1α1, which inhibits TOR and growth (3). Consistent with this model, the *snrk2.2 snrk2.3* mutant (*snrk2d*) shows markedly defective PR growth under favorable conditions due to aberrant repression of TOR activity (3). This defect is fully rescued by the *snrk1α1* mutation, demonstrating it is SnRK1α1-dependent (3).

To investigate further how TOR and root growth are controlled by SnRK2 and SnRK1, we examined the root meristems of Col-0 control seedlings, the *snrk2d* mutant, and its cross with *snrk1α1* (*snrk2d/1α1*) (Fig. 1*A*). As previously reported (6, 7), ABA treatment reduced the number of root meristematic cells, leading to smaller meristems in Col-0. In contrast, *snrk2d* showed a reduction in meristem size and cell number already under mock conditions, and this was fully rescued by the *snrk1α1* mutation. Consistent with the ABA hyposensitivity of *snrk2d* and *snrk2d/1α1* (3), the addition of ABA did not further decrease meristem size or cell number in these mutants (Fig. 1*A*). These meristem phenotypes correlate well with the PR length previously observed in these mutants and conditions (3) and suggest that reduced cell proliferation contributes to the reduced PR length of *snrk2d* in mock, mimicking the situation of Col-0 plants under ABA.

We next hypothesized that regulation of SnRK1 and growth by ABA and SnRK2s could
involve changes in SnRK1α1 subcellular localization. Firstly, SnRK1 and SnRK2 are highly
enriched in the nuclei of root meristematic cells (3). Secondly, SnRK2-harboring SnRK1 repressor
complexes localize to the nucleus in *Nicotiana benthamiana* epidermal cells (3). Thirdly, *in planta,*
the TOR complex subunit RAPTOR1B interacts with SnRK1α1 (3, 8) in the cytosol (9). Fourthly,
the subcellular localization of SnRK1α1 is central to its function (10). We therefore monitored
SnRK1α1 subcellular localization in mock- or ABA-treated seedlings in two independent lines
expressing SnRK1α1-GFP from its own regulatory regions (*SnRK1α1-GFP#1* and *SnRK1α1-*
GFP#2). SnRK1α1 was detected in the meristematic, elongating and differentiating zones of the
PR both in mock and ABA, displaying a known nuclear and cytoplasmic localization (11) (Fig.
1*B*). In mock, SnRK1α1 exhibited a ring-shaped pattern characteristic of nuclear proteins that are
absent from the nucleolus in meristematic cells (12) (Fig. 1*B*). In ABA, the nuclear signal appeared
to decline specifically in the meristematic zone (Fig. 1*B*). Electron micrographs confirmed that a
ring-shaped nuclear region surrounds a large-sized nucleolus in these cells (Fig. 1*C*). Closer
examination and GFP quantification in the meristematic zone revealed overall similar SnRK1α1-
GFP levels in mock- and ABA-treated roots but a reduction in the nucleus to cytoplasm ratio (N/C)
from 1.19 (line #1) and 1.2 (line #2) in mock to 0.73 (line #1) and 0.71 (line #2) in ABA (Fig. 1*D*),
suggesting that ABA induces SnRK1α1 relocalization from the nucleus to the cytoplasm.
Accordingly, the ABA effect was fully abolished by the nuclear export inhibitor Leptomycin B
(LMB; N/C ratio=1.36 and 1.23 in lines #1 and #2; Fig. 1*D*).

Given the marked reduction in meristem size (Fig. 1*A*) and PR length (3) of the *snrk2d*
mutant, we investigated the role of SnRK2s in SnRK1α1 localization using the cross of *snrk2d*
with the *SnRK1α1-GFP#2* line (Fig. 1*E*). In mock, SnRK1α1-GFP was barely present in the nuclei
of *snrk2d* meristematic cells, with most of the GFP signal being cytoplasmatic and yielding a N/C
ratio of 0.75 (Fig. 1*E*), comparable to the ABA-treated *SnRK1α1-GFP#2* control (Fig. 1*D*).
SnRK2.2-GFP suffered from overall protein degradation, but there was no change in its subcellular
localization (Fig. 1*F*). Nuclear exit of SnRK1α1 could be observed in other highly proliferating
regions of the root, such as the LR tip (Fig. 1*G*). However, in differentiated root cells, ABA
treatment and SnRK2 depletion did not alter SnRK1α1 subcellular localization (Fig. 1*H*).
Furthermore, the ABA-triggered changes in SnRK1α1 localization remained undetected in
nucleus/cytoplasm fractions of whole roots (Fig. 1*I*). Altogether, these results indicate that the

SnRK2-dependent localization of SnRK1α1 and its ABA-triggered nuclear exit is specific to
meristematic cells, consistent with reports that SnRK1α1 subcellular localization is controlled in
a cell type-specific manner (13, 14).

Given that SnRK1α1 is required for the inhibitory effect of ABA on TOR and growth (3),
we wondered whether SnRK1α1 nuclear exit is a premise for such inhibition. Indeed, pre-treatment
with LMB blocked the ABA-triggered repression of TOR activity (Fig. 2*A*), measured as
phosphorylation of ribosomal protein S6 [RPS6^{S240}; (3)]. Moreover, as compared to control plants
expressing wild-type SnRK1α1 (*control-α1*), the repression of TOR by ABA was also defective
when SnRK1α1 was fused to an SV40 nuclear localization sequence (NLS) that favors its presence
in the nucleus (*NLS-α1*; Fig. 2*B*). This demonstrates that SnRK1α1 nuclear exit is necessary for
inhibiting TOR in response to ABA. Furthermore, the fact that nuclear export is crucial for
repressing TOR (Fig. 2*A*) but only SnRK1α1 (Fig. 1*D*), and not SnRK2.2, translocates to the
cytoplasm in response to the hormone (Fig. 1*F*), suggests that the previously reported role of
SnRK2s in this process (3, 15) may be indirect *via* SnRK1α1 regulation.

We conclude that root growth is modulated by ABA through changes in SnRK1α1
subcellular localization, allowing control of TOR activity and cell proliferation in the root
meristem in accordance to *e.g.* water availability (Fig. 2*C*). When conditions are favorable,
SnRK1α1 is sequestered in the nucleus by SnRK2-containing repressor complexes. Dissociation
of these complexes in response to ABA releases SnRK1α1, which translocates to the cytoplasm
and inhibits TOR activity and growth. This mode of regulation appears to be specific to
meristematic cells, where TOR is highly enriched and where active cell proliferation takes place
(4). Future work will address whether such specificity stems from the exclusive ability of
meristematic cells to form SnRK2-SnRK1α1 complexes, to export SnRK1α1 from the nucleus or
both.

REFERENCES

1.   S. R. Cutler, P. L. Rodriguez, R. R. Finkelstein, S. R. Abrams, Abscisic Acid: Emergence
of a Core Signaling Network. *Annual Review of Plant Biology* **61**, 651–679 (2010).
2.   J. M. Harris, Abscisic Acid: Hidden Architect of Root System Structure. *Plants* **4**, 548–
572 (2015).
3.   B. Belda-Palazón, *et al.*, A dual function of SnRK2 kinases in the regulation of SnRK1
and plant growth. *Nature Plants* **6**, 1345–1353 (2020).

4.   Z. Ahmad, Z. Magyar, L. Bögre, C. Papdi, Cell cycle control by the target of rapamycin
signalling pathway in plants. *Journal of Experimental Botany* **70**, 2275–2284 (2019).
5.   M. Jamsheer K, M. Kumar, V. Srivastava, SNF1-related protein kinase 1: the many-faced
signaling hub regulating developmental plasticity in plants. *Journal of Experimental*
Botany **72**, 6042–6065 (2021).
6.   S. Promchuea, Y. Zhu, Z. Chen, J. Zhang, Z. Gong, ARF2 coordinates with PLETHORAs
and PINs to orchestrate ABA-mediated root meristem activity in Arabidopsis. *Journal of*
Integrative Plant Biology **59**, 30–43 (2017).
7.   L. Yang, *et al.*, ABA-Mediated ROS in Mitochondria Regulate Root Meristem Activity by
Controlling PLETHORA Expression in Arabidopsis. *PLOS Genetics* **10**, e1004791
(2014).
8.   J. van Leene, *et al.*, Capturing the phosphorylation and protein interaction landscape of the
plant TOR kinase. *Nature Plants* **5**, 316–327 (2019).
9.   E. Nukarinen, *et al.*, Quantitative phosphoproteomics reveals the role of the AMPK plant
ortholog SnRK1 as a metabolic master regulator under energy deprivation. *Scientific*
Reports **6**, 31697 (2016).
10. M. Ramon, *et al.*, Default activation and nuclear translocation of the plant cellular energy
sensor SnRK1 regulate metabolic stress responses and development. *Plant Cell* **31**, 1614–
1632 (2019).
11. M. Bitrián, F. Roodbarkelari, M. Horváth, C. Koncz, BAC-recombineering for studying
plant gene regulation: Developmental control and cellular localization of SnRK1 kinase
subunits. *Plant Journal* **65**, 829–842 (2011).
12. B. Desvoyes, A. Arana-Echarri, M. D. Barea, C. Gutierrez, A comprehensive fluorescent
sensor for spatiotemporal cell cycle analysis in Arabidopsis. *Nature Plants* **6**, 1330–1334
(2020).
13. C. Han, *et al.*, KIN10 promotes stomatal development through stabilization of the
SPEECHLESS transcription factor. *Nature Communications* **11**, 1–10 (2020).
14. N. E. Blanco, D. Liebsch, M. G. Díaz, Å. Strand, J. Whelan, Dual and dynamic
intracellular localization of Arabidopsis thaliana SnRK1.1. *Journal of Experimental*
Botany **70**, 2325–2338 (2019).
15. P. Wang, *et al.*, Reciprocal Regulation of the TOR Kinase and ABA Receptor Balances
Plant Growth and Stress Response. *Molecular Cell* **69**, 100-112.e6 (2018).
165

FIGURE LEGENDS

Figure 1. Impact of ABA and SnRK2s on SnRK1α1 subcellular localization and cell
proliferation in the root apical meristem. (*A*) Meristems of Col-0, *snrk2d* and *snrk2d/1α1* 7d-
old seedlings with or without ABA treatment (50 µM, 48h). Arrowheads: region for cortical cell
number and meristem length quantifications (violin plots). n=23-24; $p < 0.05$, one-way ANOVA
with Tukey's HSD test. Scale bar: 50 µm. (*B*) Root apices of 4d-old *SnRK1α1-GFP#1* seedlings
with or without ABA treatment (50 µM, 3h). Scale bar: 30 µm. Arrowheads: same position
between upper and bottom photographs. (*C*) Electron micrograph of meristematic epidermal cells

with magnification (*right panel*) showing the cellular ultrastructure. C: cytoplasm; N: nucleus; No:

nucleolus. Scale bars: left, 5 µm; right, 1 µm. (*D-E*) SnRK1α1-GFP subcellular localization in

epidermal cells of 4d-old root meristems. Scale bar: 10 µm. Dotted lines: nuclear boundary.

Quantification of SnRK1α1-GFP mean cellular fluorescence and nucleus-to-cytosol (N/C) ratios.

n=5-6 (*D*) or 8 (*E*); one-way ANOVA with Dunett´s test. (*D*) Two independent SnRK1α1-GFP

lines (#1 and #2) with or without ABA (50 µM, 3h) or LMB (2.5 µM, 1h prior to ABA addition)

treatment. (*E*) Impact of the presence (control, *SnRK1α1-GFP#2)* or absence (*SnRK1α1-GFP#2;*

snrk2d) of SnRK2s. (*F*) Expression and subcellular localization of SnRK2.2-GFP in epidermal

cells of 4d-old root meristems treated and quantified as in (*D*). Scale bar: 10 µm. *n*=5; two-tailed

Student t-test. (*G-H*) SnRK1α1-GFP subcellular localization in LRs (*G*) and the PR differentiation

zone (*H*) in seedlings (Line #1 and/or #2) with or without ABA treatment (50 µM, 3h) and in

absence (*SnRK1α1-GFP#2; snrk2d*) of SnRK2s. Scale bar: 30 µm. (*G*) Nine-day old seedlings.

(*H*) Four-day old seedlings. N/C ratios of differentiated root cells. *n*=5, two-tailed Student t-test

(line #1) or one-way ANOVA (line #2). *PI*, propidium iodide. *ns*, non-significant. (*I*)

Representative SnRK1α1 immunoblot from nuclear and cytoplasmic fractions of 14d-old Col-0

whole roots with or without ABA (50 µM ABA, 3h). H3, Histone 3 marker for nuclear fractions.

Figure 2. Impact of SnRK1α1 subcellular localization on TOR signaling and root growth. (*A-*

B) Representative immunoblots and quantification of RPS6^{S240} phosphorylation (phospho-

RPS6/total-RPS6) in Col-0 (*A*) or *control-α1* and *NLS-α1* (*B*) seedlings with or without ABA (50

µM, 3h) or LMB (2.5 µM, 1h prior to ABA addition). Images from the same gel were cropped for

showing *α1* and *NLS-α1* contiguously. (*A*) *n*=5; (*B*) *n*= 3; error bars, SEM; two-tailed Student *t*-

test. (*C*) Under favorable conditions, SnRK1α1 is sequestered in the nucleus by repressor

complexes containing SnRK2 [and a PP2C (3)], enabling TOR activity in the cytoplasm,

meristematic cell proliferation and root growth. Dissociation of these complexes in ABA by the

hormone-bound PYR/PYL/RCAR receptors releases SnRK1α1 which exits the nucleus and

inhibits TOR and growth. *TORC1*, TOR Complex 1; *N*, nucleus; *C*, cytoplasm. Created with

BioRender.com.

Materials and Methods

Experimental details are provided in SI Appendix.

All study data are included in the paper.

Acknowledgements

We thank J-K. Zhu, M. Bennett and C. Koncz for *snrk2d, SnRK2.2-GFP,* and *SnRK1α1-GFP#1*
seeds. We also thank the IGC Facilities for excellent plant care (*Plant*), sample processing and
imaging (*Electron Microscopy*) and technical advice (*Advanced Imaging*; supported by PPBI-
POCI-01-0145-FEDER-022122). This work was funded by FCT [UIDB/04551/2020, LISBOA-
01-0145-FEDER-028128, PTDC/BIA-BID/32347/2017], and by EU Horizon 2020 programme
(BBP; H2020-WF-2018-2020/H2020-WF-01-2018, Grant 867426).

Author contributions

212   BBP performed experiments and analyzed data. MC performed the root meristem assays and
213   analyzed the data. TB and FR contributed tools and expertise and supported conceptual work. EBG
214   and BBP conceived the project, designed experiments, analyzed and interpreted data and wrote the
215   manuscript.

Competing interests

217   The authors declare no competing interests.

Figure 1. Impact of ABA and SnRK2s on SnRK1α1 subcellular localization and cell proliferation in the root apical meristem. (*A*) Meristems of Col-0, *snrk2d* and *snrk2d/1α1* 7d-old seedlings with or without ABA treatment (50 μM, 48h). Arrowheads: region for cortical cell number and meristem length quantifications (violin plots). n=23-24; $p < 0.05$, one-way ANOVA with Tukey's HSD test. Scale bar: 50 μm. (*B*) Root apices of 4d-old *SnRK1α1-GFP#1* seedlings with or without ABA treatment (50 μM, 3h). Scale bar: 30 μm. Arrowheads: same position between upper and bottom photographs. (*C*) Electron micrograph of meristematic epidermal cells with magnification (*right panel*) showing the cellular ultrastructure. C: cytoplasm; N: nucleus; No: nucleolus. Scale bars: left, 5 μm; right, 1 μm. (*D-E*) SnRK1α1-GFP subcellular localization in epidermal cells of 4d-old root meristems. Scale bar: 10 μm. Dotted lines: nuclear boundary. Quantification of SnRK1α1-GFP mean cellular fluorescence and nucleus-to-cytosol (N/C) ratios. n=5-6 (*D*) or 8 (*E*); one-way ANOVA with Dunett´s test. (*D*) Two independent SnRK1α1-GFP lines (#1 and #2) with or without ABA (50 μM, 3h) or LMB (2.5 μM, 1h prior to ABA addition) treatment. (*E*) Impact of the presence (control, *SnRK1α1-GFP#2)* or absence (*SnRK1α1-GFP#2; snrk2d*) of SnRK2s. (*F*) Expression and subcellular localization of SnRK2.2-GFP in epidermal cells of 4d-old root meristems treated and quantified as in (*D*). Scale bar: 10 μm. n=5; two-tailed Student t-test. (*G-H*) SnRK1α1-GFP subcellular localization in LRs (*G*) and the PR differentiation zone (*H*) in seedlings (Line #1 and/or #2) with or without ABA treatment (50 μM, 3h) and in absence (*SnRK1α1-GFP#2; snrk2d*) of SnRK2s. Scale bar: 30 μm. (*G*) Nine-day old seedlings. (*H*) Four-day old seedlings**.** N/C ratios of differentiated root cells. n=5, two-tailed Student t-test (line #1) or one-way ANOVA (line #2). *PI*, propidium iodide. *ns*, non-significant. (*I*) Representative SnRK1α1 immunoblot from nuclear and cytoplasmic fractions of 14d-old Col-0 whole roots with or without ABA (50 μM ABA, 3h). H3, Histone 3 marker for nuclear fractions.

Figure 2. Impact of SnRK1α1 subcellular localization on TOR signaling and root growth. (*A-B*) Representative immunoblots and quantification of RPS6^{S240} phosphorylation (phospho-RPS6/total-RPS6) in Col-0 (*A*) or *control-α1* and *NLS-α1* (*B*) seedlings with or without ABA (50 μM, 3h) or LMB (2.5 μM, 1h prior to ABA addition). Images from the same gel were cropped for showing *α1* and *NLS-α1* contiguously. (*A*) *n*=5; (*B*) *n*= 3; error bars, SEM; two-tailed Student *t*-test. (*C*) Under favorable conditions, SnRK1α1 is sequestered in the nucleus by repressor complexes containing SnRK2 [and a PP2C (3)], enabling TOR activity in the cytoplasm, meristematic cell proliferation and root growth. Dissociation of these complexes in ABA by the hormone-bound PYR/PYL/RCAR receptors releases SnRK1α1 which exits the nucleus and inhibits TOR and growth. *TORC1*, TOR Complex 1; *N*, nucleus; *C*, cytoplasm. Created with BioRender.com.

247