Functional heterogeneity of cell populations increases robustness of pacemaker function in a numerical model of the sinoatrial node tissue

Alexander V. Maltsev, Michael D. Stern, Edward G Lakatta, Victor A. Maltsev*

Intramural Research Program, National Institute on Aging/NIH, Baltimore, MD, USA

*Contact information:

E-mail: **maltsevvi@mail.nih.gov**

Short title: Cell heterogeneity regulates cardiac pacemaker function

Abstract

Each heartbeat is initiated by specialized pacemaker cells operating within the sinoatrial node (SAN). While individual cells within SAN tissue exhibit substantial heterogeneity of their electrophysiological parameters and Ca cycling, the role of this heterogeneity for cardiac pacemaker function remains mainly unknown. Here we investigated the problem numerically in a 25x25 square grid of coupled-clock Maltsev-Lakatta cell models and tested the hypothesis that functional heterogeneity of cell populations increases robustness of SAN function. The tissue models were populated by cells with different degree of heterogeneity of the two key model parameters of the coupled-clock system, maximum L-type Ca current conductance (g_{CaL}) and sarcoplasmic reticulum Ca pumping rate (P_{up}). Our simulations showed that in the areas of P_{up}-g_{CaL} parametric space at the edge of the system stability where action potential (AP) firing was absent or dysrhythmic in tissues populated by identical cells, rhythmic AP generation was rescued in tissues populated by cells with random distributions of g_{CaL} or P_{up} (but keeping the same average values). This effect to increase robust AP generation was synergistic with respect to heterogeneity in both g_{CaL} and P_{up} and was further strengthened by clustering of cells with higher g_{CaL} or P_{up}. The effect of functional heterogeneity was not due to a simple summation of activity of intrinsically firing cells naturally present in SAN; rather AP firing cells locally and critically interacted with non-firing/dormant cells. When firing cells prevailed, they recruited many dormant cells to fire, strongly enhancing overall SAN function. And vice versa, prevailing dormant cells suppressed AP firing in cells with intrinsic automaticity and halted SAN automaticity.

Abbreviations and key terms

SAN – sinoatrial node

V_m – membrane potential

AP – action potential

I_{CaL} – L-type Ca current

g_{CaL} – maximum conductance of L-type Ca current

SR – sarcoplasmic reticulum

RyR - ryanodine receptor (Ca release channel)

SERCA2a - SR Ca pump

P_{up} – maximum rate of SR Ca pumping by SERCA2a

PKA - protein kinase A

CaMKII - Ca/calmodulin-dependent protein kinase II

GPU - graphics processing unit

CUDA - Compute Unified Device Architecture

****

Non-firing zone – the area of P_{up}–g_{CaL} parametric space where individual SAN cells lack automaticity

Chaotic firing zone – the area of P_{up}–g_{CaL} parametric space where individual SAN cells generate chaotic (dysrhythmic) AP firing

Rhythmic firing zone – the area of P_{up}–g_{CaL} parametric space where individual SAN cells generate rhythmic AP firing

The bifurcation line – the border line in P_{up}–g_{CaL} parametric space separating firing, non-firing, and chaotic firing zones.

Robustness – ability of SAN tissue to generate rhythmic APs in a wide range of model parameters: the larger rhythmic firing zone, the higher robustness of the tissue model.

The coupled-clock system: a coupled system of ion current oscillator in the cell membrane and intracellular Ca oscillator of the SR inside a SAN cell. According to the coupled-clock theory, the oscillators critically interact via multiple Ca- and voltage- dependent mechanisms to generate normal SAN cell automaticity.

Introduction

The sinoatrial node (SAN) of the right atrium is the heart's primary pacemaker, providing both robust and flexible automatic electrical depolarizations that capture the surrounding atrial myocardium and drive the heartbeat. These electrical depolarizations are caused by concurrent operation of pacemaker cells residing in SAN tissue. These cells are electrical oscillators that spontaneously generate rhythmic changes of their membrane potential (V_m), producing relatively periodic spontaneous action potentials (APs) (reviews (Mangoni and Nargeot, 2008;Maltsev et al., 2014)). The SAN was discovered in 1907 (Keith and Flack, 1907). However, despite enormous amount of experimental data, numerical modeling, and plausible theories, the SAN operation remains still mysterious after all these years (Weiss and Qu, 2020).

The enigma of SAN function is that the SAN is an extremely complex heterogeneous tissue regarding individual cell properties and their connections within the tissue (Boyett et al., 2000). Furthermore, at the cellular level, individual SAN cells operate via a complex coupled-clock paradigm (Maltsev and Lakatta, 2009;Lakatta et al., 2010), i.e. they possess not only an electrical oscillator within their cell surface membranes, but also an intracellular oscillator, sarcoplasmic reticulum (SR) that cycles Ca in-and-out via SR Ca pump (SERCA2a) and release channels, ryanodine receptors (RyRs). Ca releases via RyRs interact with Na/Ca exchanger and L-type Ca current (I_{CaL}), and together they drive diastolic depolarization via a positive feedback mechanism (dubbed ignition (Lyashkov et al., 2018)). The synchronization of molecular actions, in turn, is driven by enhanced enzymatic activity of Ca-activated adenylate cyclases (AC1 and AC8) (Mattick et al., 2007;Younes et al., 2008) and protein kinases (PKA and CaMKII) (Vinogradova and Lakatta, 2009;Lakatta et al., 2010) that modulate respective ion channel properties and accelerate Ca pumping.

The expression of Ca cycling proteins, and membrane ion channels vary substantially among individual SAN cells (Honjo et al., 1996;Honjo et al., 1999;Lei et al., 2001;Musa et al., 2002;Monfredi et al., 2017). For example, I_{CaL} density varies by an order of magnitude (Monfredi et al., 2017) (Fig. 1A, red squares). This results in a large degree of functional heterogeneity among individual cells, ranging from frequently and rhythmic AP firing to infrequently and dysrhythmic firing, to nonfiring cells (recently dubbed "dormant" cells) (Kim et al., 2018;Tsutsui et al., 2018;Tsutsui et al., 2021). The SAN cell population is also extremely heterogeneous with respect to the AP firing rate increase in response to β-adrenergic stimulation (Kim et al., 2021).

At the SAN tissue level, it was initially thought that a "master" pacemaker cell or a leading pacemaker center dictates the excitation rate and rhythm of other SAN pacemaker cells (Sano et al., 1978;Bleeker et al., 1980). Subsequent studies suggested that individual cells mutually entrain each other to fire APs with a common period (dubbed "democratic" process) (Jalife, 1984;Michaels et al., 1987). More recent high-resolution imaging of intact SAN tissue at a single cell level, however, demonstrated that while majority of SAN cells indeed fire synchronously with a common period, many cell fire "at will", i.e. at various rates and irregularly, or remain silent, generating only local Ca releases that do not advance into AP-induced Ca transients (Bychkov et al., 2020;Fenske et al., 2020). Thus, how structural and functional heterogeneities within and among SAN cells give rise to synchronized AP firing at the SAN exits remains a mystery at the frontier of pacemaker research (Clancy and Santana, 2020;Weiss and Qu, 2020).

Numerical modeling is a powerful approach to study complex cell interactions. Here we used a modified 2-dimensional model of SAN tissue developed by Campana (Campana, 2015),

featuring faster parallel computing via graphics processing unit (GPU) to test the hypothesis that functional heterogeneity of cells within SAN tissue increases robustness of its pacemaker function. We adapted the model of SAN tissue to include diverse cell population with respect to maximum SR Ca pumping rate (P_{up}) and I_{CaL} conductance (g_{CaL}) as observed experimentally (Honjo et al., 1996;Monfredi et al., 2017)(Fig. 1A, Table 1). We found that SAN tissue populated by heterogenous cells could generate rhythmic AP firing within the areas of P_{up} - g_{CaL} parametric space where SAN populated by identical cells lacked automaticity or generated arrhythmias. The effect was synergistic with respect to P_{up} and I_{CaL} and the robustness of heterogeneous SAN tissue could be further increased by clustering cells with similar properties.

Methods

Single-cell model of SANC

Our SAN tissue model is comprised of single cell models developed by Maltsev and Lakatta in 2009 (Maltsev and Lakatta, 2009), which encompasses a system of 29 first order differential equations (see state variables y_1-y_{29} in Table 2). This was the first SAN cell numerical model featuring coupled operation of Ca and membrane clocks. The model predicted the contribution of spontaneous Ca release during diastolic depolarization via activation of inward Na/Ca exchanger current that explained numerous experimental data. The computer code for the original model can be freely downloaded in CellML format (maltsev_2009_paper.cellml) at http://models.cellml.org/workspace/maltsev_2009 and run using the Cellular Open Resource (COR) software developed at the University of Oxford by Garny et al. (Garny et al., 2009) (for recent development of the COR see http://www.opencor.ws/).

Since 2009, the model has been tested, modified and used in numerous applications (Maltsev and Lakatta, 2010;Kurata et al., 2012;Severi et al., 2012;Maltsev and Lakatta, 2013;Campana, 2015;Sirenko et al., 2016;Kim et al., 2018;Lyashkov et al., 2018;Yang et al., 2021). According to the coupled-clock theory, a coupled system of Ca and surface membrane "clocks" can provide the robustness and flexibility required for normal pacemaker cell function. A simple numerical release-pumping-delay Ca oscillator is capable of generating any frequency between 1.3 and 6.1 Hz, but cannot generate high-amplitude signals without the assistance of membrane clocks. The coupled-clock system utilizes the greater flexibility of the SR Ca clock while simultaneously accounting for the large Ca oscillation amplitudes fueled via sarcolemmal Ca influx via I_{CaL}. A modern interpretation of the coupled-clock function is that Ca clock operates via a criticality mechanism, whereas the membrane clock operates as a limited cycle oscillator, comprising a perfect synergistic pacemaker cell system (Weiss and Qu, 2020).

Multi-cellular model of SAN

The present study examined performance of SAN tissue that was modelled as a square grid of 25 by 25 of Maltsev-Lakatta cell models (Maltsev and Lakatta, 2009), with each cell being connected to its four neighbors (except those on the border). Our model was an adapted version of the tissue model originally developed by Campana (Campana, 2015). The final formulation to compute voltage in a 2-dimensional network of SAN cells was as follows (equation 2.7 in (Campana, 2015)):

$$V_m[m] = x[n] + step \times (rad^2 \frac{\pi}{\rho C_m deltx} V_{net} - \frac{I_{ion}[n] + stim}{C_m})$$

Where $V_m[m]$ is V_m of the cell with index m within the cellular network represented by a vector (1..number of cells); x is the state variable array; for each time step, m and n denote the same cell, only in two different vectors and $x[n]$ represents the value assumed by $V_m[m]$ at the previous time step; V_{net} is the sum of voltages from the four neighboring cells (right, left, up, and down): $V_{net}=V_{net\_R}+V_{net\_L}+V_{net\_U}+V_{net\_D}$ (Fig. 2); *step* is the model integration time step of 0.005 ms; *rad* is the cell radius of 4 μm; ρ is the intracellular resistivity of 10^4 MΩ·m; C_m is the electrical membrane capacitance of 32 pF; *deltx* is the cell length of 70 μm; I_{ion} is the sum of all ion currents; *stim* is stimulus current (*stim*=0 in our study).

Simulation of functional heterogeneity in SAN tissue models

Tissue heterogeneity was introduced in the model by varying P_{up} and g_{CaL}, the two key parameters of the coupled-clock system (Maltsev and Lakatta, 2009). P_{up} determines the rate at which Ca is pumped into the SR by SERCA2a to reach a threshold of Ca load required for activation of Ca release, i.e. a central part in the timing mechanism of the coupled-clock system (Maltsev and Lakatta, 2009;Imtiaz et al., 2010;Vinogradova et al., 2010;Stern et al., 2014;Maltsev et al., 2017). Our cell model approximated Ca uptake flux (j_{up}) to the network SR compartment as a function of cytoplasmic [Ca] (Ca_i) as suggested originally by Luo and Rudy (Luo and Rudy, 1994) in ventricular cells and subsequently used in SAN cell models, e.g. (Kurata et al., 2002;Maltsev and Lakatta, 2009;Severi et al., 2012):

$$j_{up} = P_{up} /(1 + K_{up}/Ca_i)$$

where $K_{up} = 0.6 \cdot 10^{-3}$ mM is Ca_i for a half-maximal Ca uptake to the network SR. Thus, in our simulations of heterogeneous SAN tissue we only varied P_{up}, with K_{up} remaining constant. I_{CaL}

as a function of V_m, submembrane [Ca] (Ca_{sub}), and time (t) was approximated as follows (for

more details, see (Maltsev and Lakatta, 2009)) :

$$I_{CaL}=C_m \cdot g_{CaL} \cdot (V_m - E_{CaL}) \cdot d_L \cdot f_L \cdot f_{Ca}$$

$$d_{L,\infty}=1/\{1+ exp[-(V_m+13.5)/6]\}$$

$$f_{L,\infty}=1/\{1+exp[(V_m +35)/7.3]\}$$

$$\alpha_{dL} = -0.02839 \cdot (V_m + 35)/ \{exp[-(V_m+35)/2.5] - 1\} -0.0849 \cdot V_m / [exp(-V_m/4.8)- 1]$$

$$\beta_{dL} = 0.01143 \cdot (V_m - 5)/ \{exp[(V_m- 5)/2.5] -1\}$$

$$\tau_{dL}=1/(\alpha_{dL} + \beta_{dL})$$

$$\tau_{fL} = 257.1 \cdot exp\{-[(V_m+ 32.5)/13.9]^2 \}+ 44.3$$

$$f_{Ca,\infty}=K_{mfCa} / (K_{mfCa} + Ca_{sub})$$

$$\tau_{fCa} = f_{Ca,\infty}/ \alpha_{fCa}$$

$$dy_i/dt = (y_{i,\infty} - y_i)/ \tau_{yi}$$

where $y_i = d_L$, f_L, f_{Ca} are respective channel gating variables of Hodgkin-Huxley type (state

variables y_{16}, y_{17}, y_{18} in Table 2) with steady-state probabilities $d_{L,\infty}$, $f_{L,\infty}$, $f_{Ca,\infty}$ and gating time

constants τ_{dL}, τ_{fL}, τ_{fCa}. In simulations of heterogeneous SAN tissue, we varied only g_{CaL},

maintaining all other parameters unchanged. In other terms, only number of functional channels

in each cell was varied, but channel gating kinetics remained the same in all cells. We directly

linked g_{CaL} to I_{CaL} density measured experimentally by whole patch-clamp technique under

volage clamp conditions (Fig. 1, see details in Results). Once we bridged our model to

experimental data, we used a P_{up} - g_{CaL} bifurcation diagram previously reported in single cell

models (Maltsev and Lakatta, 2009) to guide our selection of P_{up} and g_{CaL} distributions for cell

populations that fell into parameter-dependent areas of rhythmic firing, chaotic firing, or no

firing, bordering at a bifurcation line (split yellow line in Fig. 3A). Since our study was focused on the system robustness, we examined the tissue operation in the areas of no firing or chaotic firing (dubbed non-firing zone and chaotic firing zone, respectively) in which individual cells lack normal automaticity but normal pacemaker function is possible in heterogeneous tissue models. The parameters for cell populations were set via a uniform random distribution within specified ranges (Table 1) and then uniformly randomly distributed over the SAN grid. For some cell populations, we also tested the effect of cell clustering within the SAN grid.

Computation, program code, data analysis and visualization

Our simulations were performed using a new computational approach suggested by Campana (Campana, 2015) based on CUDA technology (Compute Unified Device Architecture). A major advantage of this approach is its parallel processing via GPU that is critical to perform simulations of hundreds or even thousands cell models within SAN tissue within a reasonable time.

The model was originally written in CUDA C programming language for a GPU that featured 64 CUDA cores and 768MB RAM (Nvidia Quadro FX 1800). Aside from various code optimizations, we adapted the code to modern, high-performance GPUs such as TITAN RTX graphics card (NVIDIA corp. CA, USA) featuring 4608 CUDA cores and 24 GB RAM used in the present study. One specific important improvement was that we arranged CUDA blocks in a grid, rather than in a one-dimensional array in the original code. Furthermore, random numbers were properly initialized in the main function and copied to GPU memory to generate random values for g_{CaL} and P_{up} for each cell. The option for clustering random numbers in the center of the grid was also implemented in the main function.

All simulations began with the same initial conditions, near to the maximum diastolic potential (Table 2). In nearly all simulations, the system reached a steady pattern of AP firing (or no firing) in about 5 s. Thus, our standard simulations lasted 7.5 s, and we examined the system behaviors in the last 2.5 s. Some simulations were run longer, for 25 s, when the system required more time to reach its steady AP firing pattern. In all model simulation scenarios, we compared capabilities of spontaneous AP firing by SAN tissue models populated by heterogenous cells vs. control models populated by identical cells whose g_{CaL} and P_{up} were assigned to their respective mean values in heterogenous tissue models. The data analysis and visualizations (including movies of V_m in the tissue) were performed using in-house written programs in Python 3.10.1 (Python Software Foundation, www.python.org) and Delphi 10.4. (Embarcadero, Austin, TX).

Results

Matching the experimentally measured I_{CaL} densities and the model parameter g_{CaL}

To bridge realistic I_{CaL} densities (measured in pA/pF by patch clamp (Monfredi et al., 2018), Fig. 1A) and the model parameter, g_{CaL} (in nS/pF), we performed voltage-clamp simulations of our single cell model (Fig. 1B) with the voltage step protocol (see inset) similar to that used in experimental studies (Honjo et al., 1996;Monfredi et al., 2017). The I_{CaL} - V_m relationship for the peak values of the simulated I_{CaL} traces (Fig. 1C) revealed a maximum peak I_{CaL} current of 9.068 pA/pF at 0 mV with a basal state conductance of 0.464 nS/pA, bridging experimental and model data via black dash line in the dual Y scale plot in Fig. 1A. In our tissue model scenarios involving a large variety of g_{CaL} values in individual cell models, we always referred to Fig. 1A to make sure that g_{CaL} remained within the respective range I_{CaL} densities measured experimentally.

SAN models with heterogeneous g_{CaL} close to the bifurcation line

First, we tested the hypothesis that reported heterogeneity in I_{CaL} density among SAN cells (i.e. g_{CaL} in the model terms) adds robustness to AP firing within SAN tissue. We constructed scenarios in which the mean g_{CaL} values were in the non-firing zone along the bifurcation line (yellow circles in Fig. 3A), but with a substantial spread of individual g_{CaL} values (horizontal double-headed aqua arrows). Three scenarios with different g_{CaL} distributions were tested to ensure robustness of the simulation results. Our specific choice of g_{CaL} range in each scenario is given in Table 1 (scenarios 1-3). The respective simulation results are presented in Fig. 3B-D, Table 1, and visualized in Movies 1-3. The percentage of AP firing cells within the time interval from 5 to 7.5 s after simulation onset was the most important parameter, by which SAN tissue performance was evaluated (columns "% firing cells" tissue vs. separate in Table 1).

In all three scenarios, rhythmic AP firing was indeed observed in our simulations of SAN heterogeneous models (Fig.3B-D, blue lines), whereas all models populated with identical cells failed to generate APs (Fig.3B-D, red lines in middle panels). In scenarios 1 and 2, impulses originating within multiple areas in the grid spread in a wave-like fashion to other areas involving the majority of cells (Movies 1 and 2). In scenario 3, rhythmic wave-like AP propagation recruited almost all cells (83.2%), despite mean values of P_{up} and g_{CaL} fell into the chaotic firing zone (Movie 3). Thus, for scenario 3, we extended our simulation to 25 s to ensure that the control model populated with identical cells indeed generated chaotic firing (Fig. 3D, middle panel inset), rather than just ceased firing after 7.5 s in our shorter standard simulation run. Thus, scenario 3 explicitly shows that adding heterogeneity to g_{CaL} can have

antiarrhythmic effect: dysrhythmic (chaotic) firing in SAN tissue model populated by identical cells is shifted to rhythmic firing in the heterogeneous SAN model.

Histograms of AP firing length distribution calculated for all three scenarios (Fig.3B-D, left panels) exhibited sharp peaks between 400 and 500 ms with a relatively small standard deviation of about 10 ms (Table 1). Synchronization of AP firing among cells is also evident from grid-average V_m, oscillation amplitude (about 25 mV, middle panels) and the percentage of firing cells ($V_m > 0$) that oscillated with an amplitude of 15-20% (right panels).

SAN models with heterogenous P_{up}

Next, we examined 3 model scenarios (#4, 5, 6 in Table 1) in which P_{up} was uniformly randomly distributed, but g_{CaL} was fixed. The P_{up} value in a given cell mimics the functional effect of phospholamban phosphorylation to regulate SR Ca pumping via SERCA2a. The mean values of P_{up} and fixed g_{CaL} (yellow circles in Fig. 4A) were chosen to move along and close to the bifurcation line, still remaining within the non-firing zone. Here we tested the specific hypothesis that adding heterogeneity to P_{up} in cell populating SAN tissue increases its robustness to generate rhythmic AP firing.

A substantial cell-to-cell variability in phospholamban phosphorylation is expected based on a substantial variability of both basal AP firing rates and responses to β-adrenergic stimulation among isolated SAN cells (Kim et al., 2021). However, in contrast to g_{CaL}, there is no single-cell data in the literature to set up cell populations in the model within an exact P_{up} range. To get an idea of what to expect for physiologically relevant lower limit in P_{up} variations in individual cells (when testing model stability), one can refer, for example, to how low P_{up} can

descend during the physiological responses to cholinergic receptor stimulation. Within physiologic range of acetylcholine concentrations up to 1 μM, the P_{up} values can decrease on average from 12 mM/s to as low as 4 mM/s (Fig 4A in (Maltsev and Lakatta, 2010)), indicating that P_{up} can actually be very flexible in each SAN cell during its normal operation. Thus, we constructed and tested 3 model scenarios with substantial variability of P_{up} (Fig. 4; Table 1, scenarios 4-6; Movies 4-6). Our model simulations in scenarios 5 and 6 demonstrated that P_{up} heterogeneity can indeed restore normal SAN tissue operation within the non-firing zone, thus supporting our hypothesis.

In scenario 4, however, g_{CaL} (and hence cell excitability) were set so low, that none of the 625 cells in the entire grid continued to generate APs after about 4 s (Movie 4). In scenarios 5 and 6, however, despite the decreasing average of P_{up} values, the tissue generated APs (Table 1) as g_{CaL} increased (Table 1). This indicates importance of g_{CaL} and clock coupling in this mechanism of robustness. Surprisingly and in contrast to simulations involving the g_{CaL} spread in the previous section, all or almost all (100% or 98%) cells fired rhythmic APs (Table 1). However, the tissue-wide AP firing occurred in a less synchronized manner, especially in scenario 5, manifested by substantial spread in AP cycle lengths among cells (histogram in Fig. 4B, large SD value in Table 1, Movie 5). Furthermore, the percentage of firing cells at a given time was extremely noisy. A large fraction of the grid showed persistent unsynchronized AP firing that was unlike any scenario across g_{CaL} spread (#1-3) where the mean voltage of the grid exhibited a clear oscillatory pattern. Interestingly, the histogram of AP cycle length in scenario 5 appeared to be bi-modal with two peaks, indicating that cells may self-organize into two sub-populations with different AP cycle lengths.

Rescuing AP firing requires substantial heterogeneity in P_{up} or g_{CaL}

It is important to emphasize that in all above scenarios rescuing AP firing required a substantial spread of g_{CaL} values (within 0.4 nS/pF) or P_{up} values. In other terms, in the non-firing zone at each P_{up} value, the spread of g_{CaL} must reach a critical value in order to rescue AP firing. While the spread of 0.4 nS/pF is indeed substantial, it was realistic, because assigned g_{CaL} values remained within experimentally measured range of I_{CaL} as shown in our bridging dual Y plot in Fig. 1A. We tested several additional scenarios with narrower g_{CaL} distribution spreads < 0.4 nS/pF, but those non-realistic SAN tissue models were unable to converge to steady AP firing. One such failed tissue model is shown in Fig. 5A (scenario 7, no firing) by aqua double-headed arrow.

Functional synergy of P_{up} and g_{CaL} heterogeneities

To gain further insights into SAN tissue operation, we tested the hypothesis that heterogeneity of cell populations with respect to both Ca clock and membrane clock would synergistically increase robustness of the pacemaker function. When P_{up} or g_{CaL} were distributed separately in scenarios 4 and 7 (described above), the respective SAN tissue models lacked automaticity (failed to fire APs). Thus, we simulated and examined an additional special scenario (#8) that included the distributions of both g_{CaL} and P_{up} assigned in scenarios 4 and 7 (purple diagonal double-headed arrow in Fig. 5A, Table 1). Our simulations showed that such system with dual parameter distribution generated APs (Fig. 5B, Movie 7), indicating that heterogeneities in g_{CaL} and P_{up} act indeed synergistically in rescuing normal system operation within the non-firing zone. Remarkably, the percentage of firing cells in the tissue increased to 69% from 42% for the

same cell population when cells are not connected to each other, indicating ongoing recruitment of dormant cells to fire APs when cells were connected. AP firing in this scenario, however, was poorly synchronized among cells with such narrow distribution of g_{CaL} values around a low mean value of only 0.34 nS/pF.

P_{up} and g_{CaL} clustering is yet another mechanism of robust SAN function

In all previous scenarios g_{CaL} and P_{up} were uniformly randomly distributed within a specific range. Experimental studies, however, indicate that the SAN features clusters of cells with specific types of activity (Fig. 4 in (Bychkov et al., 2020)). Thus, we tested the hypothesis that the natural cell clustering provides an additional "gear" to enhance robustness of AP firing in SAN tissue. To test this hypothesis, we attempted to rescue automaticity in scenario 7 in which cells had heterogeneous g_{CaL} within an extremely narrow range from 0.3 to 0.38 nS/pF. Indeed, when cells were uniformly randomly distributed over the tissue, the modelled system of cells lacked automaticity However, the system was able to achieve rhythmic AP firing when the same heterogenous population of cells was locally redistributed with higher g_{CaL} clustering towards the grid center (Fig.5C; Table 1, scenario 9; Movie 8). Finally, we tested the hypothesis that, similar to g_{CaL}, increased heterogeneity in P_{up} can also add robustness to the system of interacting cells. Indeed, while the tissue model in scenario 4 lacked automaticity with uniformly random P_{up} distribution among cells within the tissue (Movie 4), clustering cells with higher P_{up} towards the tissue center rescued the system automaticity (scenario 10, Fig 5D, Table 1, Movie 9).

Discussion

Our new approach to the problem

Heterogeneity of SAN cells has been known and well-documented for a long time (Boyett et al., 2000), but its importance for SAN function remains unclear. Numerical model simulations performed in the present study revealed an interesting and counterintuitive aspect of the problem: While noise and heterogeneity are usually associated with undesirable disturbances or fluctuations, SAN cell heterogeneity actually increases robustness of cardiac pacemaker function. Our work took advantage of well-described properties of single isolated SAN cells in numerous previous experimental and theoretical studies. Specifically, the intrinsic behaviors of individual cells have been characterized by parametric sensitivity analyses (Maltsev and Lakatta, 2009;Kurata et al., 2012) that revealed synergistic contributions of the two key parameters P_{up} and g_{CaL} to generate normal automaticity in individual SAN cells as depicted in P_{up} - g_{CaL} diagram (panel A in Figs 3-5). We used this diagram to test specific hypotheses in different scenarios in each model simulation (Table 1) and to interpret the simulation results (discussed below). In other terms, on the one hand we investigated how the SAN tissue behaves when it is populated by a large variety of individual cells with known intrinsic properties (i.e. the systems approach). On the other hand, we limited our consideration to only two key model parameters g_{CaL} and P_{up} and their variable combinations that, in essence, determine the function of the SAN cell coupled-clock system (i.e. we combined reductionist and systems approaches).

Comparison with previous numerical studies of SAN tissue

A plethora of numerical models of SAN tissue function have been developed previously, including models in one dimension (Zhang et al., 2001;Huang et al., 2011;Glynn et al., 2014;Li

et al., 2018), two dimensions (Michaels et al., 1987;Oren and Clancy, 2010;Campana, 2015;Gratz et al., 2018;Karpaev et al., 2018), and full-scale SAN models in three dimensions (Munoz et al., 2011;Li et al., 2013;Li et al., 2014;Kharche et al., 2017;Mata et al., 2019). These models examined effects of cell-to-cell coupling (Li et al., 2013;Campana, 2015;Gratz et al., 2018;Mata et al., 2019), inter-cellular variability(Campana, 2015;Mata et al., 2019), coupling to other cell types (fibroblasts) (Oren and Clancy, 2010;Karpaev et al., 2018), and normal and abnormal impulse propagation in 3 D models (Munoz et al., 2011;Li et al., 2014;Kharche et al., 2017).

To our knowledge only two numerical studies (Campana, 2015;Mata et al., 2019) have investigated the impact of heterogeneity of cell parameters on SAN tissue function. However, in both studies cell heterogeneity was limited to **only surface membrane clock** parameters and mimicked by a simultaneous randomization of **all membrane conductances** with respect to their basal values within a fixed percentage range (dubbed sigma). Mata et. al. (Mata et al., 2019) reported time course of synchronization of AP firing vs. sigma. Campana (Campana, 2015) reported that the cell population synchronizes on a rate slightly higher than the one of the isolated cells with the basal values of conductances, but it does not equal the rate of the fastest cell. Our study, in part, confirms this result in scenarios 3 and 6 with chaotic firing in the basal state (Figs 3D and 4C), but provides further insights on specific importance of heterogeneity in I_{CaL} and P_{up} for robustness of AP firing in SAN tissue (not studied previously).

Examining Ca signals in addition and together with membrane clock parameters is vital in studies of pacemaker function because each SAN cell operates as a coupled-clock system, i.e. actions of I_{CaL} and P_{up} are intertwined in a synergistic manner in numerical models of both isolated cells (Maltsev and Lakatta, 2009;Kurata et al., 2012) and SAN tissue (in the present

study, Fig. 5A,B). Among membrane currents, we chose I_{CaL} for our analysis because it is involved not only in membrane clock function (it generates AP upstroke (Mangoni and Nargeot, 2008), but also in clock coupling by providing influx of Ca (coupled clock's oscillatory substrate) (Maltsev and Lakatta, 2009;Lakatta et al., 2010) and via positive feedback mechanisms among local Ca releases, Na/Ca exchanger, and I_{CaL} during diastolic depolarization (Torrente et al., 2016;Lyashkov et al., 2018). Distributions of I_{CaL} among individual SAN cells has been reported in previous experimental studies (Honjo et al., 1996;Honjo et al., 1999;Lei et al., 2001;Musa et al., 2002;Monfredi et al., 2017) and thus permitted realistic choices for our I_{CaL} modeling.

Recent theoretical studies have demonstrated that the presence of a Ca clock in addition to a membrane clock protects the SAN from annihilation, associated with sinus node arrest (Li et al., 2018); and that random parameter heterogeneity among oscillators can consistently rescue the system from losing synchrony (Zhang et al., 2021). These prior reports and the results of the present study support a novel SAN structure/function paradigm (resembling neuronal networks) that was proposed based on high-resolution imaging of SAN tissue (Bychkov et al., 2020): in the new paradigm, interactions of Ca signals and APs play a central role in generation of rhythmic cardiac impulses that emanate from the SAN.

The coupled-clock theory explains, in part, the effect of heterogenous cell populations to enhance robustness of SAN function

Our model simulations performed for SAN tissues operating at the edge of the system stability demonstrated that the robustness of AP firing is enhanced when either g_{CaL} or P_{up} varies among

the cells (Figs 3 and 4). This result can be explained, in part, on the basis of the coupled-clock theory (Maltsev and Lakatta, 2009). While P_{up} and g_{CaL} impact cell automaticity via different biophysical mechanisms involving Ca and membrane clocks, respectively; the clocks are coupled and the intrinsic automaticity is defined by a combined and synergistic action of the clocks as depicted in the P_{up}-g_{CaL} diagram (Maltsev and Lakatta, 2009). Perturbation of ether clock inevitably affects the other clock and the entire coupled-clock system (Yaniv et al., 2013). That is why g_{CaL} and P_{up} effectively enhanced robustness of SAN function in our simulations.

More specifically, by allowing P_{up} and/or g_{CaL} in the non-firing zone to substantially deviate from their mean values (and "penetrate" into the rhythmic firing zone indicated by double-headed arrows in panels A in Figs 3,4,5), we actually create a sub-population of cells that have intrinsic automaticity in addition to the sub-population of intrinsically non-firing cells (dubbed dormant cells (Kim et al., 2018;Tsutsui et al., 2018;Tsutsui et al., 2021). The two sub-populations of cells intimately interact, determining the ultimate performance (or failure) of the SAN tissue they comprise. Thus, the coupled-clock theory explains, in part, another important result of our study that g_{CaL} and P_{up} heterogeneities act synergistically in rescuing SAN from halt. When the SAN system featured heterogeneity in both g_{CaL} and P_{up}, almost all cells (~70%, scenario 8) generated APs in marked contrast to SAN tissue with heterogeneity in either g_{CaL} or P_{up}, in which all cells remain non-firing (scenarios 4 and 7). Indeed, the respective (purple) double-headed arrow in P_{up}-g_{CaL} diagram in Fig. 5A penetrates deeply into the rhythmic firing zone, enriching SAN tissue with cells featuring high-performance automaticity.

Furthermore, the fact that the SR Ca pumping is a key timing mechanism in the coupled-clock theory can also explain the differences in rescuing tissue automaticity via P_{up} or g_{CaL}: AP cycle length substantially varied in tissues with P_{up} heterogeneity (scenario 5), while only slight

AP cycle length variability was observed in tissues with g_{CaL} heterogeneity (scenarios 1-3). On the other hand, the fact that I_{CaL} generates AP upstroke, i.e. g_{CaL} is critical for cell excitability, can explain the presence of non-firing cells in scenarios 1-3. Indeed, as we set g_{CaL} within a substantial spread (including very low values, like in real SAN cells, Fig. 1A), the entire cell population is comprised of three sub-populations with respect to cell intrinsic properties: (i) spontaneously firing APs; (ii) excitable, but lacking automaticity; (iii) non-excitable. Thus, the intrinsically non-excitable cells do not generate APs in the SAN tissue, explaining our result in scenarios 1-3 involving g_{CaL} heterogeneity that the percentage of firing cells was always less than 100%. On the other hand, in scenarios involving P_{up} heterogeneity all or almost all (100% or 98%) cells fired rhythmic APs when g_{CaL} was chosen relatively high, thereby excluding non-excitable cells.

Emergence of SAN tissue automaticity is a critical phenomenon

While the coupled-clock theory has explained, in part, many results of the present study, its application to tissue function is limited, because it predicts only intrinsic cell properties, i.e. cell operation in isolation. Cell tissue, however, operates at the next, higher level of organization and its automaticity emerges from interactions of numerous cells with different intrinsic properties within cell communities. These interactions go beyond the coupled-clock theory (at least in its present form). One important aspect of such emergent behavior was revealed by our simulations: the percentage of firing cells in tissue in all scenarios involving either P_{up} or g_{CaL} (except scenarios 4 and 7 lacking automaticity) was much larger than that when the same cells operated in isolation (Table 1), indicating that some excitable cells lacking automaticity were

recruited to fire APs by their neighboring cells in local cell communities to increase overall performance of the SAN tissue. And vice versa, the SAN tissue surprisingly lacked automaticity in scenarios 4 or 7, despite their double-headed arrows penetrated deep into the rhythmic firing zone (Fig. 5A), indicating that a major fraction of cells (39% or 30%, respectively, Table 1) was capable of generating rhythmic APs, according to the coupled-clock theory, but did not fire APs.

Thus, our simulations at the edge of the system stability discovered strong amplification (i.e. self-organization) of either AP firing or suppression of AP firing among the cells within SAN tissue. In other terms, whether the system gains automaticity or fails, seems to be a critical phenomenon that depends on intimate interactions of intrinsically firing and non-firing cells (dormant cells). If a sub-population of firing cells decreases, these cells become surrounded by too many non-firing cells, and these non-firing cells suppress the entire system automaticity (for example in scenarios 4 and 7). And, vice versa, when intrinsically firing cells reach a critical number, they prevail: they not only fire AP themself, but also recruit a fraction of dormant cells to fire APs.

Because we distributed cells uniformly randomly, the critical phenomenon of the emerging automaticity in our tissue model can be interpreted in terms of Poisson clumping (Aldous, 1989), a phenomenon wherein random events (in space and/or time) have a tendency to occur in clusters, clumps, or bursts. In other words, forming cell clusters with high g_{CaL} and/or P_{up} could be a natural consequence of the random distribution. When these natural clusters of intrinsically firing cells are rare and small, their automaticity is suppressed by surrounding dormant cells (which, in turn, also tend to form clusters via Poisson clumping).

Cell clustering is yet another mechanism to enhance robustness of SAN function

While forming functional clusters via Poisson clumping in our model is a matter of chance (albeit predictable via statistical methods (Aldous, 1989)), the real SAN tissue, as any biological tissue, can direct cell locations to optimize its function. Indeed, high resolution imaging studies in intact SAN detected clusters of cells with various types of functional activity (Fig. 4 in (Bychkov et al., 2020)). Based on this logic, we rescued SAN tissue function in the "hopeless" scenarios 7 and 4 just by reorganizing the cell locations within the tissue models (scenarios 9 and 10, Movies 8 and 9). Dormant cells were "kicked" to SAN periphery, while intrinsically firing cells supported AP firing of each other in the center. As a result, the tissue automaticity was not only rescued, but a substantial fraction of dormant cells (at the border of the firing/non-firing cells) was recruited to fire, as evident from the increase of the percentage of firing cells (scenarios 9 and 10 in Table 1). Thus, cell clustering (e.g. with respect to g_{CaL}, P_{up}, or perhaps other parameter) may represent an additional mechanism by which SAN tissue can enhance its robust function at the edge of stability. A large number of local oscillators detected in intact SAN (Bychkov et al., 2020) may reflect cell clustering due to heterogeneity in I_{CaL} (yielding non-excitable cells, besides excitable cells, as described above, see also Fig. 1A), but the fact that the cell clusters operate at different frequencies point to (among perhaps other interpretations) cell heterogeneity in SR Ca pumping determined by various levels of phospholamban phosphorylation among cells or cell clusters (mimicked by P_{up} heterogeneity in our model). Indeed, heterogeneity in P_{up} created numerous oscillators operating at various frequencies in our tissue models (wide histograms in Figs 4B, 5B, 5D).

An apparent downside of cell clustering, however, is that the isolated clusters of automaticity lack physiological importance until their signals overcome the non-firing cell

"barriers" to interact with other clusters and/or the rest of the SAN to ultimately deliver synchronized, rhythmic (but not metronomic) APs to atria. While this appears to be a problem in our simple 2-dimensional model of SAN tissue, the real SAN tissue is a 3-dimensional, multi-layer meshwork of long and highly branched, intertwined cells (Bychkov et al., 2020). Such tight meshwork of HCN4$^+$/Conexin43$^-$ cells is structured for signal transfer within and among the functional clusters, regardless of the specific nature of cell interactions that remains unknown.

Cell heterogeneity is an antiarrhythmic mechanism

Another important result of our study is that heterogeneity in g_{CaL} or P_{up} has antiarrhythmic effect. Indeed, while the tissue models (scenarios 3 and 6) with identical cells generated chaotic firing, allowing g_{CaL} or P_{up} values to spread (but keeping same averages over the cellular network) shifted the system towards normal, rhythmic operation (Fig. 3D, Movie 3 and Fig. 4C, Movie 4). This result is remarkable and counterintuitive: when the noise of an intrinsically chaotic oscillatory system is combined with the noise of random distribution of its key parameter (g_{CaL} or P_{up}) within a SAN network, it is hard to expect that such "double-noisy" system would generate normal rhythmic automaticity; but it does.

Study limitations and future studies

SAN is an extremely heterogeneous, complex system with respect to individual cell properties, cell interactions, cell network structure, and its autonomic system modulation. The present study tested a hypothesis that heterogeneity of cell functional properties confers robustness to SAN

tissue pacemaker function. Our testing focused on two specific key parameters of the cell

coupled clock-system, g_{CaL} and P_{up} using a simple model of SAN tissue that was a square grid of

25x25 cells of only one cell model type (Maltsev-Lakatta model). Because we study robustness,

we further focused our simulations and analysis close to the edge of the system stability and did

not examine the entire parametric space of P_{up} - g_{CaL}. Thus, our study provides evidence (i.e.

Proof of Principle) supporting our specific hypothesis and does not include other numerous

important factors that determine SAN function. Future numerical studies can explore

contributions of other ion currents such as I_{Na} (Lei et al., 2004), I_{CaT} (Ono and Iijima,

2005;Tanaka et al., 2008), and funny current (for its heterogeneity data obtained in isolated cells,

see (Honjo et al., 1996;Monfredi et al., 2017)), phosphorylation of Ca cycling proteins

(phospholamban and RyRs), autonomic modulation, cell network structure, including cell

connectivity and interactions in 3 dimensions, interactions with other cell types (e.g. fibroblasts

(Camelliti et al., 2004) and telocytes (Mitrofanova et al., 2018)), mechano-sensitivity (Quinn and

Kohl, 2012), endogenous Ca buffers, pathological conditions, aging, etc.

Conclusions

Natural heterogeneity of pacemaker cells increases robustness of cardiac pacemaker function, i.e.

the SAN tissue populated by heterogeneous cells can generate normal automaticity within the

cell parameter ranges, whereas SAN populated by identical cells would exhibit either

dysrhythmic AP firing or complete lack of automaticity. This effect to enhance pacemaker

function at the edge of the system stability is synergistic with respect to heterogeneity of Ca and

membrane clocks, and it is not due to a simple summation of activity of intrinsically firing cells

that are naturally present in heterogeneous SAN. These firing cells critically interact with intrinsically non-firing cells (dormant cells) and either recruit many of them to fire granting SAN automaticity, or non-firing cells suppress firing cells and SAN tissue automaticity. Clustering cells with specific cell parameters provides an additional mechanism to enhance robustness of SAN function, but can generate isolated communities of firing cells in our simple 2-dimensional model. Therefore, additional mechanisms are required for those clusters to interact with the rest of the SAN. Such mechanisms can be explored in the future in more complex SAN tissue models. Our results provide new insights for SAN operation at the edge of stability that can be helpful not only to understanding normal SAN function especially at low rates and in HCN4[+]/Conexin43[-] cell communities (Bychkov et al., 2020;Fenske et al., 2020), but also with respect to understanding SAN function in normal aging (linked to deficient cAMP-PKA-Ca signaling (Liu et al., 2014)) and pathological conditions (such as sick sinus syndrome and SAN arrhythmias (Dobrzynski et al., 2007;John and Kumar, 2016)).

Acknowledgments

This research was supported by the Intramural Research Program of the National Institutes of Health, National Institute on Aging.

Figure legends

Fig. 1. SAN cell model simulations to match model parameter g_{CaL} to I_{CaL} density measured experimentally. A: A double Y scale plot of experimental data points of I_{CaL} densities measured by whole-cell patch clamp in a large population of SAN cells (red squares, left Y axis, replotted from (Monfredi et al., 2018)) and respective model g_{CaL} parameter values (right Y axis). B: Traces of I_{CaL} / V_m relationship simulated in Maltsev-Lakatta single cell model with the basal state value of g_{CaL} =0.464 nS/pF using voltage clamp protocol (inset) and conditions similar to those in experimental study of (Monfredi et al., 2018). The family of I_{CaL} traces was simulated by applying a series of square voltage pulses from a holding potential of -45 mV with a step of 5 mV up to 45 mV (inset). Before each stimulation, I_{CaL} activation and inactivation gating variables f_L and d_L were set to their steady-state values. To closely simulate experimental whole-cell patch clamp conditions in (Monfredi et al., 2018), 5 mM EGTA (equilibrated to 100 nM of free Ca) was added to Ca formulations of cell submembrane space and cytoplasm. C: Respective I_{CaL} / V_m relationship calculated for peak I_{CaL}. The I_{CaL} peak of 9.068 pA/pF at 0 mV was set as a bridge between experimental data and model g_{CaL} =0.464 nS/pF (black dash line in panel A).

Fig. 2. A schematic illustration of SAN tissue structure and calculation of the network voltage (V_{net}) for a cell with index n in Campana model (Campana, 2015). Pink squares denote individual SAN cells (Maltsev-Lakatta models) located at the nodes of a square grid. The cells interact via intercellular conductances placed uniformly at the grid's edges (shown by blue double-headed arrows). V_{net} is the sum of voltages from the four neighboring cells (right, left, up,

and down). Shown is only a 3x3 element the network; the tissue continues beyond the dash lines to form the entire 25x25 cell network used in majority of our simulations.

Fig 3. Heterogeneity in g_{CaL} increases robustness of AP firing in SAN tissue models close to the edge of stability. A: P_{up}- g_{CaL} bifurcation diagram previously reported for single SAN cell model (modified from (Maltsev and Lakatta, 2009)) shows the parametric space for rhythmic AP firing, no firing, and chaotic firing (white labels). Yellow circles show coordinates for fixed P_{up} and g_{CaL} mean values (all in the non-firing zone) in cell populations used in our simulations of SAN tissue function in 3 specific scenarios 1-3 (Table 1) close to the bifurcation line (yellow line). Double-headed aqua arrows show the g_{CaL} ranges for each scenario (Note: the ranges extend beyond the diagram, with the labels showing the extensions). All three tissue models fired rhythmic APs (Movies 1-3). B-D: Each panel shows the results of simulations in each scenario: histograms of AP cycle length distribution, space-average V_m vs. time, and the percentage of firing cells (V_m>0) vs. time (from 5s to 7.5 s). In scenario 3, we extended the simulation run to 25 s. Inset in panel D (middle) shows average V_m from 5 s to 25 s. Red traces show space-average V_m vs. time for the control SAN tissue model populated by identical cells with average P_{up} and g_{CaL} values of the respective heterogeneous model.

Fig 4. Heterogeneity in P_{up} increases robustness of AP firing in SAN tissue models close to the edge of stability. A: P_{up}- g_{CaL} bifurcation diagram illustrating cell populations in simulations of SAN tissue function in scenarios 4-6 (Table 1) similar to Figs 3A. Yellow circles show coordinates for mean P_{up} and fixed g_{CaL} values (in non-firing and the chaotic firing zones). Black

double-headed arrows show the exact P_{up} ranges for each scenario. The tissue model with low g_{CaL} in scenario 4 failed to generate AP firing despite a higher average P_{up} value, but scenarios 5 and 6 with higher g_{CaL} values fired rhythmic AP (Movies 4-6). B-C: the results of simulations in scenarios 5 and 6. In scenario 6, we extended the simulation run to 25 s. Inset in panel C (middle) shows average V_m from 5 s to 25 s.

Fig 5. Robustness of SAN operation is increased via combined P_{up} and g_{CaL} heterogeneities and their clustering. A: P_{up}- g_{CaL} bifurcation diagram illustrating cell populations in simulations of SAN tissue function in scenarios 7-10 (Table 1) similar to Figs 3A and 4A. Yellow circle shows coordinates for mean P_{up} and mean g_{CaL} values (in the non-firing zone). Double-headed arrows show P_{up} and g_{CaL} ranges in these scenarios. The tissue models with heterogeneity in either single parameter failed to generate rhythmic APs. However, when both parameters fluctuated, the tissue model fired rhythmic APs (Movie 7). Furthermore, when either parameter (g_{CaL} or P_{up}) was clustered, the tissue also generated APs in scenarios 9 or 10 (Movies 8, 9), respectively. B-D: the results of simulations in scenarios 8-10.

#	Distr.	P_{up}, min, mM/s	P_{up}, max, mM/s	P_{up} mean, mM/s	g_{CaL}, min, nS/pF	g_{CaL}, max, nS/pF	g_{CaL} mean, nS/pF	% firing cells, tissue	% firing cells, separate	Cycle length average, ms	Cycle length SD, ms
1	random		7		0.14	0.54	0.34	64.64	46	405.91	11.72
2	random		5		0.17	0.57	0.37	70.4	46	429.43	9.4
3	random		3		0.21	0.61	0.41	83.2	47.5	450.3	11.55
4	random	2	12	7		0.34		0	39	N.D.	N.D.
5	random	0	10	5		0.37		97.92	40	460.99	57.09
6	random	0	6	3		0.41		100	40	506.91	6.27
7	random		7		0.3	0.38	0.34	0	30	N.D.	N.D.
8	random	2	12	7	0.3	0.38	0.34	69.12	42	417.2	36.38
9	g_{CaL} cluster		7		0.3	0.38	0.34	60.8	30	457.28	8.39
10	P_{up} cluster	2	12	7		0.34		66.4	39	368.21	33.13

Table 1. Parameter values in each model simulation scenario (column #) and major results of our simulations during time interval from 5 s to 7.5 s after simulation onset, the percentage of AP firing cells in tissue, the percentage of cells firing APs in isolation, and AP cycle length average with its standard deviation (SD) calculated for all AP firing cells populating SAN model tissue model (25 x 25 square grid) in each scenario.

#	Variable	Description	Initial value
		Ca^{2+} cycling	
y_1	Ca_i	[Ca] in myoplasm, mM	0.0001
y_2	Ca_{sub}	[Ca] in submembrane space, mM	0.000223
y_3	Ca_{jSR}	[Ca] in the junctional SR (jSR), mM	0.029
y_4	Ca_{nSR}	[Ca] in the network SR (nSR), mM	1.35
y_5	f_{TC}	Fractional occupancy of the troponin-Ca site by Ca in myoplasm	0.02
y_6	f_{TMC}	Fractional occupancy of the troponin-Mg site by Ca in myoplasm	0.22
y_7	f_{TMM}	Fractional occupancy of the troponin-Mg site by Mg in myoplasm	0.69
y_8	f_{CMi}	Fractional occupancy of calmodulin by Ca in myoplasm	0.042
y_9	f_{CMs}	Fractional occupancy of calmodulin by Ca in submembrane space	0.089
y_{10}	f_{CQ}	Fractional occupancy of calsequestrin by Ca in junctional SR	0.032
y_{11}	R	RyR reactivated (closed) state	0.7499955
y_{12}	O	RyR open state	$3.4 \cdot 10^{-6}$
y_{13}	I	RyR inactivated state	$1.1 \cdot 10^{-6}$
y_{14}	RI	RyR RI state	0.25
		Electrophysiology	
y_{15}	V_m	Membrane potential, mV	-65
y_{16}	d_L	I_{CaL} activation	0
y_{17}	f_L	I_{CaL} voltage-dependent inactivation	1
y_{18}	f_{Ca}	I_{CaL} Ca dependent inactivation	1
y_{19}	p_{aF}	I_{Kr} fast activation	0
y_{20}	p_{aS}	I_{Kr} slow activation	0
y_{21}	p_i	I_{Kr} inactivation	1
y_{22}	n	I_{Ks} activation	0
y_{23}	y	I_f activation	1
y_{24}	d_T	I_{CaT} activation	0
y_{25}	f_T	I_{CaT} inactivation	1
y_{26}	q	I_{to} inactivation	1
y_{27}	r	I_{to} and I_{sus} activation	0
y_{28}	q_a	I_{st} activation	0
y_{29}	q_i	I_{st} inactivation	1

Table 2. State variables y_1-y_{29} with their descriptions and initial values assigned to all cells in our SAN tissue model simulations.

Movie legends

Each movie illustrates the result of numerical simulation of SAN tissue (25x25 cells) lasted 7.5 s in the respective model scenario described in Table 1. V_m dynamics in each cell in the grid was color-coded via red shade from -60 mV (pure black) to 20 mV (pure red). Simulation time is displayed in the upper left corner.

Movie 1. P_{up} = 7 mM/s, g_{CaL} ∈ [0.14, 0.54] nS/pF, mean = 0.34 nS/pF

Movie 2. P_{up} = 5 mM/s, g_{CaL} ∈ [0.17, 0.57] nS/pF, mean = 0.37 nS/pF

Movie 3. P_{up} = 3 mM/s, g_{CaL} ∈ [0.21, 0.61] nS/pF, mean = 0.41 nS/pF

Movie 4. P_{up} ∈ [2, 12], mean = 7 mM/s, g_{CaL} = 0.34 nS/pF

Movie 5. P_{up} ∈ [0, 10], mean = 5 mM/s, g_{CaL} = 0.37 nS/pF

Movie 6. P_{up} ∈ [0, 6], mean = 3 mM/s, g_{CaL} = 0.41 nS/pF

Movie 7. P_{up} ∈ [2, 12], mean =75 mM/s, g_{CaL} ∈ [0.3, 0.38] nS/pF, mean = 0.34 nS/pF

Movie 8. P_{up} = 7 mM/s, g_{CaL} ∈ [0.3, 0.38] nS/pF, mean = 0.34 nS/pF, cells with higher g_{CaL} were clustered to SAN center

Movie 9. P_{up} ∈ [2, 12], mean = 7 mM/s with g_{CaL} = 0.34 nS/pF, cells with higher P_{up} were clustered to SAN center

References

Aldous, D. (1989). Probability Approximations via the Poisson Clumping Heuristic. *Applied Mathematical Sciences* 7, Springer (book).

Bleeker, W.K., Mackaay, A.J., Masson-Pevet, M., Bouman, L.N., and Becker, A.E. (1980). Functional and morphological organization of the rabbit sinus node. *Circ Res* 46, 11-22.

Boyett, M.R., Honjo, H., and Kodama, I. (2000). The sinoatrial node, a heterogeneous pacemaker structure. *Cardiovasc Res* 47, 658-687.

Bychkov, R., Juhaszova, M., Tsutsui, K., Coletta, C., Stern, M.D., Maltsev, V.A., and Lakatta, E.G. (2020). Synchronized cardiac impulses emerge from multi-scale, heterogeneous local calcium signals within and among cells of heart pacemaker tissue. *JACC Clin Electrophysiol* 6, 907-931. doi: 10.1016/j.jacep.2020.06.022.

Camelliti, P., Green, C.R., Legrice, I., and Kohl, P. (2004). Fibroblast network in rabbit sinoatrial node: structural and functional identification of homogeneous and heterogeneous cell coupling. *Circ Res* 94, 828-835. doi: 10.1161/01.RES.0000122382.19400.14.

Campana, C.A. (2015). 2-Dimensional Computational Model to Analyze the Effects of Cellular Heterogeinity on Cardiac Pacemaking. *PhD thesis, Bologna, Italy: Universita di Bologna, Corso di Studio in Ingegneria Biomedical*.

Clancy, C.E., and Santana, L.F. (2020). Evolving Discovery of the Origin of the Heartbeat: A New Perspective on Sinus Rhythm. *JACC Clin Electrophysiol* 6, 932-934. doi: 10.1016/j.jacep.2020.07.002.

Dobrzynski, H., Boyett, M.R., and Anderson, R.H. (2007). New insights into pacemaker activity: promoting understanding of sick sinus syndrome. *Circulation* 115, 1921-1932.

Fenske, S., Hennis, K., Rotzer, R.D., Brox, V.F., Becirovic, E., Scharr, A., Gruner, C., Ziegler, T., Mehlfeld, V., Brennan, J., Efimov, I.R., Pauza, A.G., Moser, M., Wotjak, C.T., Kupatt, C., Gonner, R., Zhang, R., Zhang, H., Zong, X., Biel, M., and Wahl-Schott, C. (2020). cAMP-dependent regulation of HCN4 controls the tonic entrainment process in sinoatrial node pacemaker cells. *Nat Commun* 11, 5555. doi: 10.1038/s41467-020-19304-9.

Garny, A., Noble, D., Hunter, P.J., and Kohl, P. (2009). CELLULAR OPEN RESOURCE (COR): current status and future directions. *Philos Trans A Math Phys Eng Sci* 367, 1885-1905. doi: 10.1098/rsta.2008.0289.

Glynn, P., Onal, B., and Hund, T.J. (2014). Cycle length restitution in sinoatrial node cells: a theory for understanding spontaneous action potential dynamics. *PLoS One* 9, e89049. doi: 10.1371/journal.pone.0089049.

Gratz, D., Onal, B., Dalic, A., and Hund, T.J. (2018). Synchronization of pacemaking in the sinoatrial node: a mathematical modeling study. *Front Physics* 6, id.63. doi: doi.org/10.3389/fphy.2018.00063.

Honjo, H., Boyett, M.R., Kodama, I., and Toyama, J. (1996). Correlation between electrical activity and the size of rabbit sino-atrial node cells. *J Physiol* 496 (Pt 3), 795-808.

Honjo, H., Lei, M., Boyett, M.R., and Kodama, I. (1999). Heterogeneity of 4-aminopyridine-sensitive current in rabbit sinoatrial node cells. *Am J Physiol* 276, H1295-1304.

Huang, X., Mi, Y., Qian, Y., and Hu, G. (2011). Phase-locking behaviors in an ionic model of sinoatrial node cell and tissue. *Phys Rev E Stat Nonlin Soft Matter Phys* 83, 061917. doi: 10.1103/PhysRevE.83.061917.

Imtiaz, M.S., Von Der Weid, P.Y., Laver, D.R., and Van Helden, D.F. (2010). SR Ca^{2+} store refill--a key factor in cardiac pacemaking. *J Mol Cell Cardiol* 49, 412-426.

Jalife, J. (1984). Mutual entrainment and electrical coupling as mechanisms for synchronous firing of rabbit sino-atrial pace-maker cells. *J Physiol* 356, 221-243.

John, R.M., and Kumar, S. (2016). Sinus Node and Atrial Arrhythmias. *Circulation* 133, 1892-1900. doi: 10.1161/CIRCULATIONAHA.116.018011.

Karpaev, A.A., Syunyaev, R.A., and Aliev, R.R. (2018). Effects of fibroblast-myocyte coupling on the sinoatrial node activity: A computational study. *Int J Numer Method Biomed Eng* 34, e2966. doi: 10.1002/cnm.2966.

Keith, A., and Flack, M. (1907). The Form and Nature of the Muscular Connections between the Primary Divisions of the Vertebrate Heart. *J Anat Physiol* 41, 172-189.

Kharche, S.R., Vigmond, E., Efimov, I.R., and Dobrzynski, H. (2017). Computational assessment of the functional role of sinoatrial node exit pathways in the human heart. *PLoS One* 12, e0183727. doi: 10.1371/journal.pone.0183727.

Kim, M.S., Maltsev, A.V., Monfredi, O., Maltseva, L.A., Wirth, A., Florio, M.C., Tsutsui, K., Riordon, D.R., Parsons, S.P., Tagirova, S., Ziman, B.D., Stern, M.D., Lakatta, E.G., and Maltsev, V.A. (2018). Heterogeneity of calcium clock functions in dormant, dysrhythmically and rhythmically firing single pacemaker cells isolated from SA node. *Cell Calcium* 74, 168-179. doi: 10.1016/j.ceca.2018.07.002.

Kim, M.S., Monfredi, O., Maltseva, L.A., Lakatta, E.G., and Maltsev, V.A. (2021). beta-Adrenergic Stimulation Synchronizes a Broad Spectrum of Action Potential Firing Rates of Cardiac Pacemaker Cells toward a Higher Population Average. *Cells* 10. doi: 10.3390/cells10082124.

Kurata, Y., Hisatome, I., Imanishi, S., and Shibamoto, T. (2002). Dynamical description of sinoatrial node pacemaking: improved mathematical model for primary pacemaker cell. *Am J Physiol* 283, H2074-2101.

Kurata, Y., Hisatome, I., and Shibamoto, T. (2012). Roles of Sarcoplasmic Reticulum Ca^{2+} Cycling and Na^+/Ca^{2+} Exchanger in Sinoatrial Node Pacemaking: insights from bifurcation analysis of mathematical models. *Am J Physiol Heart Circ Physiol* 302, H2285-H2300.

Lakatta, E.G., Maltsev, V.A., and Vinogradova, T.M. (2010). A coupled SYSTEM of intracellular Ca^{2+} clocks and surface membrane voltage clocks controls the timekeeping mechanism of the heart's pacemaker. *Circ Res* 106, 659-673.

Lei, M., Honjo, H., Kodama, I., and Boyett, M.R. (2001). Heterogeneous expression of the delayed-rectifier K+ currents i(K,r) and i(K,s) in rabbit sinoatrial node cells. *J Physiol* 535, 703-714.

Lei, M., Jones, S.A., Liu, J., Lancaster, M.K., Fung, S.S., Dobrzynski, H., Camelliti, P., Maier, S.K., Noble, D., and Boyett, M.R. (2004). Requirement of neuronal- and cardiac-type sodium channels for murine sinoatrial node pacemaking. *J Physiol* 559, 835-848.

Li, J., Inada, S., Schneider, J.E., Zhang, H., Dobrzynski, H., and Boyett, M.R. (2014). Three-dimensional computer model of the right atrium including the sinoatrial and atrioventricular nodes predicts classical nodal behaviours. *PLoS One* 9, e112547. doi: 10.1371/journal.pone.0112547.

Li, K., Chu, Z., and Huang, X. (2018). Annihilation of the pacemaking activity in the sinoatrial node cell and tissue *AIP Advances* 8, 125319.

Li, P., Lines, G., Maleckar, M., and Tveito, A. (2013). Mathematical models of cardiac pacemaking function. *Frontiers in Physics* 1. doi: 10.3389/fphy.2013.00020.

Liu, J., Sirenko, S., Juhaszova, M., Sollott, S.J., Shukla, S., Yaniv, Y., and Lakatta, E.G. (2014). Age-associated abnormalities of intrinsic automaticity of sinoatrial nodal cells are linked to deficient cAMP-PKA-Ca^{2+} signaling. *Am J Physiol Heart Circ Physiol* 306, H1385-1397. doi: 10.1152/ajpheart.00088.2014.

Luo, C.H., and Rudy, Y. (1994). A dynamic model of the cardiac ventricular action potential. I. Simulations of ionic currents and concentration changes. *Circ Res* 74, 1071-1096.

Lyashkov, A.E., Behar, J., Lakatta, E.G., Yaniv, Y., and Maltsev, V.A. (2018). Positive Feedback Mechanisms among Local Ca Releases, NCX, and ICaL Ignite Pacemaker Action Potentials. *Biophys J* 114, 1176-1189. doi: 10.1016/j.bpj.2017.12.043.

Maltsev, A.V., Maltsev, V.A., and Stern, M.D. (2017). Stabilization of diastolic calcium signal via calcium pump regulation of complex local calcium releases and transient decay in a computational model of cardiac pacemaker cell with individual release channels. *PLoS Comput Biol* 13, e1005675. doi: 10.1371/journal.pcbi.1005675.

Maltsev, V.A., and Lakatta, E.G. (2009). Synergism of coupled subsarcolemmal Ca^{2+} clocks and sarcolemmal voltage clocks confers robust and flexible pacemaker function in a novel pacemaker cell model. *Am J Physiol Heart Circ Physiol* 296, H594-H615.

Maltsev, V.A., and Lakatta, E.G. (2010). A novel quantitative explanation for autonomic modulation of cardiac pacemaker cell automaticity via a dynamic system of sarcolemmal and intracellular proteins. *Am J Physiol Heart Circ Physiol* 298, H2010-H2023.

Maltsev, V.A., and Lakatta, E.G. (2013). Numerical models based on a minimal set of sarcolemmal electrogenic proteins and an intracellular Ca clock generate robust, flexible, and energy-efficient cardiac pacemaking. *J Mol Cell Cardiol* 59, 181-195.

Maltsev, V.A., Yaniv, Y., Maltsev, A.V., Stern, M.D., and Lakatta, E.G. (2014). Modern perspectives on numerical modeling of cardiac pacemaker cell. *J Pharmacol Sci* 125, 6-38.

Mangoni, M.E., and Nargeot, J. (2008). Genesis and regulation of the heart automaticity. *Physiol Rev* 88, 919-982.

Mata, A.N., Alonso, G.R., Garza, G.L., Fernández, J.R.G., García, M.a.C., and Ábrego, N.P.C. (2019). Parallel simulation of the synchronization of heterogeneous cells in the sinoatrial node. *Concurrency Computat Pract Exper*, e5317. doi: DOI: 10.1002/cpe.5317.

Mattick, P., Parrington, J., Odia, E., Simpson, A., Collins, T., and Terrar, D. (2007). Ca^{2+}-stimulated adenylyl cyclase isoform AC1 is preferentially expressed in guinea-pig sino-atrial node cells and modulates the I_f pacemaker current. *J Physiol* 582, 1195-1203.

Michaels, D.C., Matyas, E.P., and Jalife, J. (1987). Mechanisms of sinoatrial pacemaker synchronization: a new hypothesis. *Circ Res* 61, 704-714.

Mitrofanova, L.B., Gorshkov, A.N., Konovalov, P.V., and Krylova, J.S. (2018). Telocytes in the human sinoatrial node. *J Cell Mol Med* 22, 521-532. doi: 10.1111/jcmm.13340.

Monfredi, O., Tsutsui, K., Ziman, B., Stern, M.D., Lakatta, E.G., and Maltsev, V.A. (2018). Electrophysiological heterogeneity of pacemaker cells in the rabbit intercaval region, including the SA node: insights from recording multiple ion currents in each cell. *Am J Physiol Heart Circ Physiol* 314, H403-H414. doi: 10.1152/ajpheart.00253.2016.

Monfredi, O., Tsutsui, K., Ziman, B.D., Stern, M.D., Lakatta, E.G., and Maltsev, V.A. (2017). Electrophysiological heterogeneity of pacemaker cells in rabbit intercaval region, including SA node: insights from recording multiple ion currents in each cell. *Am J Physiol Heart Circ Physiol*, ajpheart 00253 02016. doi: 10.1152/ajpheart.00253.2016.

Munoz, M.A., Kaur, J., and Vigmond, E.J. (2011). Onset of atrial arrhythmias elicited by autonomic modulation of rabbit sinoatrial node activity: a modeling study. *Am J Physiol Heart Circ Physiol* 301, H1974-1983. doi: 10.1152/ajpheart.00059.2011.

Musa, H., Lei, M., Honjo, H., Jones, S.A., Dobrzynski, H., Lancaster, M.K., Takagishi, Y., Henderson, Z., Kodama, I., and Boyett, M.R. (2002). Heterogeneous expression of Ca^{2+} handling proteins in rabbit sinoatrial node. *J Histochem Cytochem* 50, 311-324.

Ono, K., and Iijima, T. (2005). Pathophysiological significance of T-type Ca^{2+} channels: properties and functional roles of T-type Ca^{2+} channels in cardiac pacemaking. *J Pharmacol Sci* 99, 197-204.

Oren, R.V., and Clancy, C.E. (2010). Determinants of heterogeneity, excitation and conduction in the sinoatrial node: a model study. *PLoS Comput Biol* 6, e1001041.

Quinn, T.A., and Kohl, P. (2012). Mechano-sensitivity of cardiac pacemaker function: pathophysiological relevance, experimental implications, and conceptual integration with other mechanisms of rhythmicity. *Prog Biophys Mol Biol* 110, 257-268. doi: 10.1016/j.pbiomolbio.2012.08.008.

Sano, T., Sawanobori, T., and Adaniya, H. (1978). Mechanism of rhythm determination among pacemaker cells of the mammalian sinus node. *Am J Physiol* 235, H379-384. doi: 10.1152/ajpheart.1978.235.4.H379.

Severi, S., Fantini, M., Charawi, L.A., and Difrancesco, D. (2012). An updated computational model of rabbit sinoatrial action potential to investigate the mechanisms of heart rate modulation. *J Physiol* 590, 4483-4499.

Sirenko, S.G., Maltsev, V.A., Yaniv, Y., Bychkov, R., Yaeger, D., Vinogradova, T., Spurgeon, H.A., and Lakatta, E.G. (2016). Electrochemical Na^+ and Ca^{2+} gradients drive coupled-clock regulation of automaticity of isolated rabbit sinoatrial nodal pacemaker cells. *Am J Physiol Heart Circ Physiol* 311, H251-267. doi: 10.1152/ajpheart.00667.2015.

Stern, M.D., Maltseva, L.A., Juhaszova, M., Sollott, S.J., Lakatta, E.G., and Maltsev, V.A. (2014). Hierarchical clustering of ryanodine receptors enables emergence of a calcium clock in sinoatrial node cells. *J Gen Physiol* 143, 577-604.

Tanaka, H., Komikado, C., Namekata, I., Nakamura, H., Suzuki, M., Tsuneoka, Y., Shigenobu, K., and Takahara, A. (2008). Species difference in the contribution of T-type calcium current to cardiac pacemaking as revealed by r(-)-efonidipine. *J Pharmacol Sci* 107, 99-102.

Torrente, A.G., Mesirca, P., Neco, P., Rizzetto, R., Dubel, S., Barrere, C., Sinegger-Brauns, M., Striessnig, J., Richard, S., Nargeot, J., Gomez, A.M., and Mangoni, M.E. (2016). L-type Cav1.3 channels regulate ryanodine receptor-dependent Ca^{2+} release during sino-atrial node pacemaker activity. *Cardiovasc Res* 109, 451-461. doi: 10.1093/cvr/cvw006.

Tsutsui, K., Florio, M.C., Yang, A., Wirth, A.N., Yang, D., Kim, M.S., Ziman, B.D., Bychkov, R., Monfredi, O.J., Maltsev, V.A., and Lakatta, E.G. (2021). cAMP-Dependent Signaling Restores AP Firing in Dormant SA Node Cells via Enhancement of Surface Membrane Currents and Calcium Coupling. *Front Physiol* 12, 596832. doi: 10.3389/fphys.2021.596832.

Tsutsui, K., Monfredi, O., Sirenko-Tagirova, S.G., Maltseva, L.A., Bychkov, R., Kim, M.S., Ziman, B.D., Tarasov, K.V., Tarasova, Y.S., Zhang, J., Wang, M., Maltsev, A.V., Brennan, J.A., Efimov, I.R., Stern, M.D., Maltsev, V.A., and Lakatta, E.G. (2018). A coupled-clock system drives the automaticity of human sinoatrial nodal pacemaker cells. *Sci Signal* 11, eaap7608.

Vinogradova, T.M., Brochet, D.X., Sirenko, S., Li, Y., Spurgeon, H., and Lakatta, E.G. (2010). Sarcoplasmic reticulum Ca^{2+} pumping kinetics regulates timing of local Ca^{2+} releases and spontaneous beating rate of rabbit sinoatrial node pacemaker cells. *Circ Res* 107, 767-775.

Vinogradova, T.M., and Lakatta, E.G. (2009). Regulation of basal and reserve cardiac pacemaker function by interactions of cAMP mediated PKA-dependent Ca^{2+} cycling with surface membrane channels. *Journal of Molecular and Cellular Cardiology* 47, 456-474.

Weiss, J.N., and Qu, Z. (2020). The Sinus Node: Still Mysterious After All These Years. *JACC Clin Electrophysiol* 6, 1841-1843. doi: 10.1016/j.jacep.2020.09.017.

Yang, D., Morrell, C.H., Lyashkov, A.E., Tagirova, S., Zahanich, I., Yaniv, Y., Vinogradova, T., Ziman, B., Maltsev, V.A., and Lakatta, E.G. (2021). Ca2+ and Membrane Potential Transitions During Action Potentials are Self-Similar to Each Other and to Variability of AP Firing Intervals Across the Broad Physiologic Range of AP Intervals During Autonomic Receptor Stimulation. *Front Physiol* in print. doi: DOI: 10.3389/fphys.2021.612770.

Yaniv, Y., Sirenko, S., Ziman, B.D., Spurgeon, H.A., Maltsev, V.A., and Lakatta, E.G. (2013). New evidence for coupled clock regulation of the normal automaticity of sinoatrial nodal pacemaker cells: Bradycardic effects of ivabradine are linked to suppression of intracellular Ca cycling. *J Mol Cell Cardiol* 62C, 80-89.

Younes, A., Lyashkov, A.E., Graham, D., Sheydina, A., Volkova, M.V., Mitsak, M., Vinogradova, T.M., Lukyanenko, Y.O., Li, Y., Ruknudin, A.M., Boheler, K.R., Van Eyk, J., and Lakatta, E.G. (2008). Ca^{2+}-

stimulated basal adenylyl cyclase activity localization in membrane lipid microdomains of cardiac sinoatrial nodal pacemaker cells. *J Biol Chem* 283**,** 14461-14468.

Zhang, H., Holden, A.V., and Boyett, M.R. (2001). Gradient model versus mosaic model of the sinoatrial node. *Circulation* 103**,** 584-588.

Zhang, Y., Ocampo-Espindola, J.L., Kiss, I.Z., and Motter, A.E. (2021). Random heterogeneity outperforms design in network synchronization. *Proc Natl Acad Sci U S A* 118. doi: 10.1073/pnas.2024299118.

Figure 1

$$V_{net} = V_{net\_R} + V_{net\_L} + V_{net\_U} + V_{net\_D}$$

Figure 2

Figure 3

Figure 4

A

Rate (Hz)
3.3

Scenario 4 (no firing)
Scenario 10 (P_{up} clustered)

Scenario 8 P_{up} & g_{CaL} vary

Rhythmic firing

0.3   0.34   0.38

Scenario 7 (no firing)
Scenario 9 (g_{CaL} clustered)

No firing zone

Chaotic firing

Pumping rate (P_{up}), mM/s

I_{CaL} conductance (g_{CaL}), nS/pF

B # of cells

Scenario 8 (both P_{up} & g_{CaL} vary)

AP cycle length, ms

V_m, mV

Time, s

% firing cells

Time, s

C # of cells

Scenario 9 g_{CaL} clustered

AP cycle length, ms

V_m, mV

Time, s

% firing cells

Time, s

D # of cells

Scenario 10 P_{up} clustered

AP cycle length, ms

V_m, mV

Time, s

% firing cells

Time, s

Figure 5