**Testing for genetic assimilation with phylogenetic comparative analysis: Conceptual,**

**methodological, and statistical considerations**

Alex R. Gunderson[1][*] and Liam J. Revell[2,3]

1. Department of Ecology and Evolutionary Biology, Tulane University, New Orleans, LA, USA

2. Department of Biology, University of Massachusetts Boston, Boston, MA, USA

3. Facultad de Ciencias, Universidad Católica de la Santísima Concepción, Concepción, Chile

Running header: Evolution of phenotypic plasticity

* corresponding author; email: agunderson@tulane.edu

**Abstract**

Genetic assimilation is a process that leads to reduced phenotypic plasticity during adaptation

to novel conditions, a potentially important phenomenon under global environmental change.

Null expectations when testing for genetic assimilation, however, are not always clear. For

instance, the statistical artifact of regression to the mean could bias us towards detecting

genetic assimilation when it has not occurred. Likewise, the specific mechanism underlying

plasticity expression may affect null expectations under neutral evolution. We used

macroevolutionary numerical simulations to examine both of these important issues and their

interaction, varying whether or not plasticity evolves, the evolutionary mechanism, trait

measurement error, and experimental design. We also modified an existing reaction norm

correction method to account for phylogenetic non-independence. We found: 1) regression to

the mean is pervasive and can generate spurious support for genetic assimilation; 2)

experimental design and post-hoc correction can minimize this spurious effect; and 3) neutral

evolution can produce patterns consistent with genetic assimilation without constraint or

selection, depending on the mechanism of plasticity expression. Additionally, we re-analyzed

published macroevolutionary data supporting genetic assimilation, and found that support was

lost after proper correction. Considerable caution is thus required whenever investigating

genetic assimilation and reaction norm evolution at macroevolutionary scales.

Keywords: plasticity, acclimation, global change, evolution, reaction norm

**Introduction**

Phenotypic plasticity is an important phenomenon in evolutionary biology with far-reaching

implications (Ghalambor et al. 2007; Murren et al. 2015). For example, plasticity in a given trait

can facilitate evolutionary radiations (Pfennig et al. 2010), contribute to local adaptation (Price

et al. 2003), and dampen selection on underlying genetic variation (Price et al. 2003; Crispo

2008). Interest in the expression and evolution of phenotypic plasticity has increased in recent

years due to the intense selection pressure that human activity has imposed on the natural

world (Chevin et al. 2010; Kingsolver and Buckley 2017). Taxa with greater plasticity in traits

that are important for tolerating human-induced changes to the environment are predicted to

fare better under the novel conditions created by global warming, urbanization, and species

introductions (Somero 2010; Gunderson and Stillman 2015; Seebacher et al. 2015; Gunderson

et al. 2017). For instance, plasticity in thermal physiology is predicted to decrease the negative

impact of global warming on ectotherms (Gunderson et al. 2017; Riddell et al. 2018; Rohr et al.

2018; Morley et al. 2019). Greater knowledge of the forces that shape and constrain phenotypic

plasticity is crucial both for our fundamental understanding of ecology and evolution and our

ability to predict and mitigate the consequences of global change.

Adaptation to novel conditions is sometimes associated with a reduction or loss of

plasticity in traits under selection, a process that is known as genetic assimilation (Waddington

1953; Schlichting and Pigliucci 1998). Some have argued that genetic assimilation is a key

mechanism of adaptive evolutionary change (West-Eberhard 2003; Crispo 2007). For example,

the canalization of formerly induced phenotypes can provide a means of adaptive phenotypic

divergence during local adaptation and adaptive radiation (Ehrenreich and Pfennig 2016;

Martin et al. 2016; Gunter et al. 2017; Schneider and Meyer 2017). In addition, genetic

assimilation may be an outcome of population responses to anthropogenic global change.

Lande (2009) found that the evolution of increased phenotypic plasticity followed by genetic

assimilation should result from selection due to dramatic changes in the environment.

The intersection of adaptation, genetic assimilation, and global change is highlighted by

the Baseline Tolerance/Tolerance Plasticity Trade-off Hypothesis (hereafter referred to as the

Trade-off Hypothesis). Emerging from the field of evolutionary physiology, the hypothesis

states that, as organisms evolve greater baseline tolerance to extreme temperatures (heat or

cold), plasticity in their thermal tolerance should actually decrease (van Heerwaarden and

Kellermann 2020). In other words, high levels of constitutive thermal tolerance evolve by

genetic assimilation, resulting in thermal tolerance phenotypes that are less sensitive to

environmental temperature variation (Sikkink et al. 2014).  Support for this hypothesis comes

from a variety of organisms, including marine and aquatic vertebrates and invertebrates at both

the intraspecific and interspecific levels (e.g., Stillman 2003; Esperk et al. 2016; Comte and

Olden 2017; Armstrong et al. 2019). One of several implications of the Trade-off Hypothesis is

that physiological plasticity serves as the raw material for thermal adaptation. Therefore, as

organisms evolve greater baseline tolerance in response to global change, they may lose both

the ability to evolve even greater tolerance and their capacity to track shorter-term

environmental changes via a plastic response (van Heerwaarden and Kellermann 2020).

Hypotheses that make predictions about trait change over time among individuals

and/or taxa, such as the Trade-off Hypothesis, are prone to spurious support due to the well-

known statistical phenomenon called *regression to the mean*. A consequence of this

phenomenon is that when a particular trait is measured repeatedly, the experimental units

(e.g., species, populations, or individuals) that start with particularly high or low values relative

to the population mean are also likely to record the greatest change in value over time as

subsequent measures regress towards the mean (Barnett et al. 2005). This is not a real

biological effect: rather, it is an artifact created by sampling and measurement error.

Regression to the mean is a common problem in repeated-measures analyses, and statistical

methods have been developed to overcome the spurious results that it can create. For

example, data resampling techniques have been applied to generate null distributions of trait

change (Jackson and Somers 1991), and methods to statistically remove regression to the mean

effects prior to statistical analysis have also been proposed (Kelly and Price 2005). These

approaches are most often employed at the intraspecific level (Ghalambor et al. 2015; Deery et

al. 2021), but accounting for regression to the mean is also necessary, though more often

neglected, in interspecific macroevolutionary analyses (Baker et al. 2015).

102           Motivated by current debate surrounding the Trade-off Hypothesis and increasing

interest in the evolution of phenotypic plasticity and genetic assimilation under global change

(Kelly 2019; Sasaki and Dam 2019; van Heerwaarden and Kellermann 2020; Barley et al. 2021;

Preston et al. 2021), we used simulations to explore the more general consequences of

regression to the mean for tests of genetic assimilation at the macroevolutionary level. We

conducted simulations that assumed mechanistically different models for plasticity evolution,

as well as different experimental sampling designs for measuring species phenotypic traits. In

addition, we conducted simulations with and without the assumption that phenotypes are

measured with error. We found that measurement error alone can be sufficient to recover a

macroevolutionary pattern that is consistent with genetic assimilation due entirely to the

phenomenon of regression to the mean. This problem can be avoided, however, by using an

appropriate experimental design. In addition, we found that macroevolutionary patterns

consistent with genetic assimilation are the null expectation under random Brownian motion

evolution for some models by which plasticity evolves. Finally, we applied a statistical method

designed to remove the regression to the mean effect from compromised data (Kelly and Price

2005), that we modified to take non-independence due to phylogeny into account. We found

that our approach can reduce the effect of regression to the mean in some, but not all,

circumstances. We applied this modified approach to published phylogenetic comparative data

that was thought to be consistent with the Trade-off Hypothesis, and showed that the

relationship is not significant when properly corrected for regression to the mean.

**Methods**

In our simulations, we assumed that there were two different environments that induce

different phenotypes. The difference between trait values in each environment represented the

magnitude of phenotypic plasticity and, in a two-environment system, is equivalent to the slope

of the reaction norm. In any given simulation, we first generated a random phylogenetic tree

with 50 terminal nodes using the *pbtree* function in the phytools R package (Revell 2012; R

Development Core Team 2021). Depending on the simulation, we either modeled the evolution

of plasticity across the phylogeny using the *fastBM* function or assumed no phenotypic

evolution at all (see below for details). In addition, we assumed a certain degree of sampling

error in the measurement of phenotypes in each environment. We did so by drawing

**Experimental design 1**    **Experimental design 2**    **Experimental design 3**

Figure 1. Three experimental designs that we simulated for estimating baseline phenotype and phenotypic plasticity of species. In Experimental Design 1, baseline phenotype and plasticity are estimated from a single group of individuals that have their phenotype measured twice: once after exposure to each environment. In Experimental Design 2, there are two groups of individuals that have their phenotype measured only once, after exposure to either the baseline or novel environment. In Experimental Design 3, there are also two groups of individuals. One group has their phenotype measured only after exposure to the baseline condition, and are used only to estimate the baseline phenotype. The other group of individuals have their phenotypes measured twice (as in Experimental Design 1), but they are only used to estimate phenotypic plasticity. Thermal environments are shown simply as an example.

phenotypes for individuals randomly from a Gaussian distribution with a mean value at the true

phenotypic population mean. For each simulation run, we calculated the phylogenetic

correlation coefficient for the association between the baseline phenotype and the reaction

norm slope using the *phyl.vcv* function. Starting conditions assumed a baseline phenotypic trait

value of 0 and a phenotypic value in the novel environment of 1. In simulations where evolution

occurred, the Brownian motion rate parameter ($\sigma^2$) was set at 1, 0.1, or 0.01 for respective

traits. Phenotypic measurement error was set to variance ($s^2$) of 1, 0.1, or 0.01.

We predicted that experimental design would influence whether or not regression to

the mean biased our results. As such, we modeled three different experimental designs for

measuring the phenotypes of each taxon in each environment (Figure 1). In Design 1, we

assumed that each individual has their phenotype measured twice in total: once after

acclimation to each environment. Therefore, in this case the same individuals are used to

estimate both the baseline phenotype and the reaction norm. In Design 2, we assumed that

different individuals are acclimated to each environment prior to phenotypic measurement. In

this case, individuals exposed to the baseline environment are used to estimate the baseline

phenotype, and the difference in phenotype between the two sets of individuals is used to

estimate the reaction norm. Finally, in Design 3, we assumed that one set of individuals is

acclimated only to the baseline environment prior to phenotypic trait measurement, and that a

second set of individuals have their phenotypes measured twice: once after acclimation to each

of the two environments. The former individuals are used only to estimate baseline

phenotypes, and the latter individuals are used only to estimate the slope of the reaction norm.

We also tested whether regression to the mean could emerge as a significant problem

when phenotypic plasticity evolves, and did so assuming a neutral evolutionary process (i.e.,

Brownian motion). We used two different models for the evolution of phenotypic plasticity,

that we refer to as Linked Phenotypes and Unlinked Phenotypes. In the Linked Phenotypes

model, we assumed that the slope and intercept of the reaction norms evolve directly, with the

phenotype that is expressed in each environment evolving indirectly as a byproduct. This is one

of the most common ways in which the evolution of phenotypic plasticity is conceptualized and

studied theoretically (Lande 2009; Chevin et al. 2010; Chevin et al. 2013). We refer to this as the

Linked Phenotypes model because the phenotypes that are expressed in each environment are

each determined by the slope and the intercept, linking them under a neutral evolutionary

process. In the Unlinked Phenotypes model, the phenotype that is expressed in a given

environment evolves directly and independently of the phenotype expressed in the other

environment. Therefore, the phenotypes that are expressed in each environment lack genetic

covariance (Via and Lande 1985), and the slope and intercept of the reaction norm emerge

merely as a byproduct of the trait values expressed in each environment. This model is based

on the concept that parts of a reaction norm can evolve independently of one another as

outlined in Ghalambor et al. (2007). Using each model, we conducted simulations in which the

evolving parameters diverge via Brownian motion, and then estimated the phylogenetic

correlation coefficient for an association between baseline phenotype and the reaction norm

slope, as described above.

When possible, we applied a statistical correction developed by Kelly and Price (2005) to

remove the regression to the mean effect from reaction norm values. The method was not

originally designed for use with phylogenetic comparative data, but we modified it to that

purpose here. We use the correction only with data simulated under Experimental Designs 1

and 2, as the method cannot be applied to Experimental Design 3. The approach is based on the

premise that if the plastic response is constrained by the baseline phenotype, then phenotypic

variation should be reduced under novel conditions and that phenotypes in the baseline and

novel conditions should be correlated within experimental units (Kelly and Price 2005). The

approach is implemented as follows. First, we estimate the phylogenetic variance in

phenotypes in each environment ($s_i$) using the *phyl.vcv* function, then we conduct a Pittman's

test of homogeneity of variance. If there is no difference in variance between the phenotypes,

then corrected reaction norms ($\widehat{D}^*$) are calculated using the following function:

Equation 1: $\widehat{D}^* = r(X_1 - \bar{X}_1) - (X_2 - \bar{X}_2)$

where $r$ is the phylogenetic correlation coefficient for the relationship between phenotypes in

the baseline and novel environments, $X_i$ is the vector of phenotypes in a given environment,

and $\bar{X}_i$ is the phylogenetic phenotypic mean in a given environment. If the phenotypic variance

differs between environments, then an adjusted correlation coefficient ($\hat{r}$) is used in Equation

1, calculated as follows:

Equation 2:    $\hat{r} = \dfrac{2rs_1s_2}{s_1^2 + s_2^2}$

Where $s_i^2$ is the estimated phylogenetic variance in the environment $i$. $\widehat{D}^*$ is then used as the

reaction norm value for subsequent analyses as described above.

Using our modified Kelly and Price (2005) approach, we reanalyzed previously published

data on heat tolerance plasticity across Nudibranch species (Armstrong et al. 2019). The

authors of that study used what we refer to as Experimental Design 2 (Figure 1), and originally

found support for the Trade-off Hypothesis, as Nudibranch species with greater baseline heat

tolerance had reduced heat tolerance plasticity. However, they did not adjust values for

potential regression to the mean as we do here.

**Results**

*Results without evolution but with measurement error*

We first present results of simulations in which phenotypic plasticity is present but does not

diverge among taxa. In this case, the only source of variation among taxa is measurement error.

Figure 2. Summary of phylogenetic correlation coefficients between baseline phenotype and reaction norm slope from simulations in which plasticity is present but does not diverge among taxa, and phenotypes in each environment are measured with error. Points indicate the median and black lines the central 95% of correlation coefficients from 1000 simulations. The error variance of simulations is shown above each column. Alternating shading differentiates different simulated experimental designs, which are written on the righthand side of the figure. "est." denotes estimated correlation coefficients given measurement error, while "c. est" denotes those same correlation coefficients adjusted to remove the effect of regression to the mean following our modified Kelly and Price (2005) correction. Adjusted results are not given for Experimental Design 3 because the adjustment cannot be applied to data collected with that design.

If regression to the mean is not an issue, we would expect the correlation coefficients

generated in our simulations to be centered on 0.

Instead, we found that measurement error generated a pattern consistent with genetic

assimilation (that is, a negative correlation between baseline phenotype and reaction norm

slope) due to regression to the mean, but only for Experimental Designs 1 and 2 (Figure 2). By

contrast, regression to the mean does not bias results using Experimental Design 3 (Figure 2).

This finding does not dependent on the level of plasticity organisms exhibit. For example, even

if there is no plasticity in the trait, a negative relationship between baseline phenotype and

reaction norm slope will emerge with Experimental Designs 1 and 2 if the phenotypes in each

environment are measured with error (see Supplemental Figure S1; see also Deery et al. 2001

for an individual-level example). The bias induced by Experimental Designs 1 and 2 is largely

removed, however, when the simulated data are adjusted using our phylogenetic modification

of the Kelly and Price (2005) correction (Figure 2 "corrected estimates").

*Results with neutral evolution of phenotypic plasticity*

If phenotypic plasticity evolves via the Linked Phenotype mechanism (in which reaction norm

slope and intercept evolve directly), a pattern consistent with genetic assimilation is unlikely to

evolve under neutral processes (Figure 3 "true" values). However, this assumes that there is no

measurement error when estimating phenotypes. With measurement error, regression to the

mean can produce spurious support for genetic assimilation with Experimental Designs 1 and 2,

though this effect depends on the magnitude of measurement error relative to the rate of

evolution (Figure 3 "estimates" values). The greater the measurement error, the more likely

that regression to the mean can become a problem. Adjusting data generated by Experimental

Designs 1 and 2 using our modified Kelly and Price correction removes the regression to the

mean effect (Figure 3 "corrected estimates"). These results, however, also highlight the

importance of judicious application of the Kelly and Price correction. This is because if there is a

weakly negative to positive correlation between baseline phenotype and plasticity, the Kelly

and Price correction can *create* a spurious negative relationship between the two variables (for

example, see Figure 3 results with Experimental Design 2, $\sigma^2 = 1$ and $s^2 = 0.01$).

If phenotypic plasticity evolves via the Unlinked Phenotypes mechanism (phenotypes in

each environment evolve directly and independently of one another), neutral evolution will

yield a negative relationship between baseline phenotype and reaction norm slope (Figure 4

"true" values). Furthermore, the negative relationship between these traits is maintained in the

Figure 3. Summary of phylogenetic correlation coefficients between baseline phenotype and reaction norm slope from simulations in which plasticity evolved by Brownian motion via the Linked Phenotypes mechanism. Each panel shows the result of 1000 simulations. Each row shows the results of simulations with a different experimental design (Figure 1): *Bottom*, Design 1. *Middle*: Design 2. *Top*: Design 3. Alternating shaded areas differentiate simulations with different Brownian motion rate parameters ($\sigma^2$). Each column contains simulations with a different error variance for phenotypic estimation ($s^2$). "true" denotes the true phylogenetic correlation values from the simulations. "est." denotes estimates of the true correlation coefficients given measurement error. "c. est" denotes estimated correlation coefficients corrected to remove the effect of regression to the mean following our modified Kelly and Price (2005) adjustment. Adjusted results are not given for Experimental Design 3 because the method cannot be applied to data collected with that design.

243    presence of measurement error (Figure 4 "estimated" values). In this case, however, adjusting

244    the data using the Kelly and Price correction mostly eliminates the true negative relationships

245    between the traits (Figure 4 "corrected" estimates). Conversely, if data are collected using

Figure 4. Summary of phylogenetic correlation coefficients between baseline phenotype and reaction norm slope from simulations in which plasticity evolved by Brownian motion via the Unlinked Phenotypes mechanism. Each panel shows the result of 1000 simulations. Each row shows the results of simulations with a different experimental design (Figure 1): *Bottom*, Design 1. *Middle*: Design 2. *Top*: Design 3. Alternating shaded areas differentiate simulations with different Brownian motion rate parameters ($\sigma^2$). Each column contains simulations with a different error variance for phenotypic estimation ($s^2$). "true" denotes the true phylogenetic correlation values from the simulations. "est." denotes estimates of the true correlation coefficients given measurement error. "c. est" denotes estimated correlation coefficients adjusted to remove the effect of regression to the mean following our modified Kelly and Price (2005) correction. Adjusted results are not given for Experimental Design 3 because the method cannot be applied to data collected with that design.

246    Experimental Design 3, the true negative relationship between the traits is largely maintained;

247    however, if measurement error is too high the capacity for Experimental Design 3 to yield the

248    true negative relationship is decreased (Figure 4).

Figure 5. Nudibranch heat tolerance plasticity data from Armstrong et al. (2019). Phenotypes were collected using Experimental Design 2 (Figure 1). Original data and analysis in gray. Re-analysis applying the modified Kelly and Price (2005) method to remove regression to the mean effects in black.

*Reanalysis of Nudibranch heat tolerance data*

We re-analyzed data on heat tolerance plasticity in Nudibranchs (Armstrong et al. 2019) by

applying our phylogenetic adaptation of the Kelly and Price (2005) correction. In the original

analysis, the authors found a significant negative relationship between independent contrasts

for baseline heat tolerance and heat tolerance plasticity (P < 0.005, Armstrong et al. 2019). In

our re-analysis, we found that a negative relationship remained, but was no longer significant at

the $\alpha$ = 0.05 level (P = 0.069; Figure 5).

**Discussion**

Herein, we have identified three general patterns important for testing and understanding the

evolution of genetic assimilation at the macroevolutionary level: 1) regression to the mean is a

pervasive problem; 2) experimental design and data correction can minimize the effect of

regression to the mean; and 3) neutral evolutionary processes can produce patterns consistent

with genetic assimilation, depending on the linkage between trait expression in different

environments. Below, we discuss each of these findings in greater detail.

*Regression to the mean when plasticity does not diverge*

When phenotypic plasticity was present but did not diverge among taxa, measurement error

alone generated spurious correlations between baseline phenotype and plasticity under

Experimental Designs 1 and 2 (Figure 2 "estimated" values). Furthermore, the magnitude of the

correlation coefficient was affected very little by the magnitude of measurement error

simulated (Figure 2). These results clearly demonstrate that regression to the mean can be a

significant problem when testing for genetic assimilation at the macroevolutionary level. The

choice of experimental design, however, also matters considerably. In particular, when data are

collected using Experimental Design 3, regression to the mean is not a persistent problem

(Figure 2).

Why are Experimental Designs 1 and 2 susceptible to regression to the mean while

Experimental Design 3 is not? The crux of the matter is that with Designs 1 and 2, the reaction

norm slope depends on the estimated baseline phenotype. Designs 1 and 2 have only one

estimate of baseline phenotype. That single baseline phenotype is also used to estimate

plasticity. If measurement error causes an estimate of baseline phenotype to be above the true

value, that will drive down the estimated reaction norm slope. Conversely, if error causes a

baseline estimate to be below its true value, this will in turn drive up the estimated reaction

norm slope. With Design 3, this cannot happen because the baseline phenotype and reaction

norm are estimated using separate samples.

Design 3 demands two independent estimates of phenotype under baseline conditions,

which we will call e1 and e2. One of those estimates (let's say e1) is used only as the baseline

phenotype, but does not contribute to the calculation of plasticity. The other estimate (e2)

contributes only to the calculation of plasticity. Under this design, if measurement error causes

e1 to be above or below the true value it will have no effect on the plasticity estimate, which

depends only on e2.

All hope is not lost with Experimental Designs 1 and 2, however. Using our modified

Kelly and Price correction, phenotypic values can be adjusted such that the bias towards a

spurious negative relationship is removed (Figure 2 "corrected" values). Next, we discuss the

implications of these different approaches when plasticity itself evolves.

*Regression to the mean under neutral evolution of plasticity (Linked Phenotype model)*

Under the Linked Phenotypes model, the intercept (essentially baseline phenotype) and

reaction norm slope evolve via Brownian motion. Under this model, neutral evolution does not

produce a negative relationship between baseline phenotype and plasticity under any

experimental design (Figure 3 "true" values). The introduction of measurement error, however,

can again lead to regression to the mean under Experimental Designs 1 and 2 (Figure 3

"estimated" values). The degree to which regression to the mean is a problem depends on the

magnitude of measurement error relative to the variance in true trait values among taxa as

denoted by the Brownian motion rate parameter. If measurement error is large and/or

phenotypic variation among taxa is low, regression to the mean emerges consistently for

Experimental Designs 1 and 2. By contrast, if measurement error is small and phenotypic

variation among taxa is large, regression to the mean is much less of an issue (Figure 3

"estimated" values).

Applying the Kelly and Price correction to data collected under Experimental Designs 1

and 2 can remove the effect of regression to the mean when it is present (Figure 3 "corrected"

values). Our results, however, also highlights the caveat that a Kelly and Price correction should

only be applied if the initial analysis yields a significant negative relationship. When initial

analyses yield no significant negative relationship, applying the Kelly and Price correction can

itself create a spurious negative relationship between baseline phenotype and plasticity.

Experimental design 3 again avoids the problem of regression to the mean due to the

estimation of two baseline phenotypes (see above).

*Regression to the mean under neutral evolution of plasticity (Unlinked Phenotypes)*

Under the Unlinked Phenotypes model, the phenotypes in each environment evolve

independently of one another, and the magnitude of plasticity emerges as a byproduct of the

difference between the two phenotypic values. Under these conditions, a true negative

association between baseline phenotype and plasticity consistently results under Brownian

motion evolution (Figure 4 "true" values). In other words, there are conditions in which neutral

evolution alone can produce a pattern consistent with genetic assimilation. We discuss this in

more detail below.

Unlike the Linked Phenotypes model, experimental error has little to effect on the

correlation coefficients calculated with the Unlinked Phenotypes model using Experimental

Designs 1 and 2 (Figure 4 "estimate" values). The reason for this is that phenotypic evolution via

Brownian motion in the Unlinked Phenotypes model amounts to random independent

increases or decreases in the phenotypes expressed in each environment, while experimental

error leads to spurious random increases or decreases in the phenotypes measured in each

environment, which from a statistical standpoint is the same process.

When data from the Unlinked Phenotypes model are adjusted using the Kelly and Price

correction, the negative relationship between baseline phenotype and plasticity disappears

(Figure 4, "corrected" values). In this case, however, the correction does not remove a spurious

relationship due to regression to the mean. Instead, it removes a true relationship that resulted

from a neutral evolutionary process. This occurs because of the conceptual foundation upon

which the Kelly and Price correction is predicated: it assumes that a hallmark of regression to

the mean is that phenotypic values between treatments (in this case, environments) within

experimental units are uncorrelated with one another. This, of course, is the precise pattern

that neutral evolution will produce if the phenotypes induced by plasticity are unlinked.

Correlation coefficients estimated using Experimental Design 3 can maintain the true

negative relationship between baseline phenotype and plasticity, but only if the measurement

error is not too large. As measurement error grows relative to the true phenotypic variance

among taxa, estimated correlation coefficients move away from the true values and towards 0

(Figure 4).

*Implications*

Our results make clear that regression to the mean is a serious statistical problem that can

confound tests of genetic assimilation at macroevolutionary scales. Whether or not regression

to the mean is an issue in any particular study will depend on the experimental design used.

Unfortunately, the problematic experimental designs (our Designs 1 and 2) are those most

often used to test for plasticity and genetic assimilation. This is certainly true within the field of

thermal physiology, where tests of the Trade-Off Hypothesis, which states that greater thermal

tolerance evolves via genetic assimilation (Heerwaarden and Kellerman 2020), are almost

invariably based on data collected using Experimental Designs 1 and 2 (e.g., Armstrong, others).

These studies typically find support for the Trade-off Hypothesis but do not account for

regression to the mean.

Fortunately, data collected using Experimental Designs 1 and 2 can be successfully

adjusted to remove regression to the mean using the Kelly and Price method. For example,

when plasticity is invariant among taxa, adjusting the data removes spurious relationships

between baseline phenotype and plasticity due to measurement error (Figure 3). The same

holds under neutral evolution when there is linkage between the traits expressed in each

environment (Figure 3). The caveat is that adjusting data using the Kelly and Price method can

incorrectly obfuscate a true negative relationship between baseline phenotype and plasticity if

the traits expressed in each environment are unlinked (Figure 4).

The need to adjust data can be avoided by using Experimental Design 3. This approach

does not lead to spurious negative relationships between baseline phenotype and plasticity

(Figures 3 and 4). It also recovers true negative relationships when they occur (Figure 4).

Unfortunately, however, this experimental design is significantly more labor intensive that

Designs 1 and 2. It requires that we estimate phenotypes on more groups of individuals (3

versus 2), and it's reliability depends on measurement error in the estimation of phenotypic

trait means for each environment, making large sample sizes crucial.

Overall, our results demonstrate that great care is necessary both in the design of

macroevolutionary tests for genetic assimilation, as well as in the interpretation of our results.

A lack of correlation between plasticity and baseline phenotype is not the correct null

hypothesis in many cases due to the statistical phenomenon of regression to the mean. Once

this is recognized, the potential solutions to the regression to the mean problem must be

applied with care. Additionally, the expected null hypothesis under neutral evolution depends

on the mechanism that underlies the expression of phenotypic plasticity. We hope that this

analysis will be a helpful reference for others in the design and analysis of future studies of

hypothesized genetic assimilation in phenotypically plastic traits.

**Acknowledgements**

We would like to thank Martha Muñoz, Shane Campbell-Staton, Andrew Moeller, and Kathleen

Ferris for helpful feedback on the ideas presented here.

Data availability:

Data and code to reproduce all the analyses of this study available at:

GitHub: https://nam11.safelinks.protection.outlook.com/?url=https%3A%2F%2Fgithub.com%2

Fliamrevell%2FGunderson-and-Revell-

2021&data=04%7C01%7Cagunderson%40tulane.edu%7C5ccb9b6c257243c5fe3d08d9c331

ba9a%7C9de9818325d94b139fc34de5489c1f3b%7C0%7C0%7C637755440671031894%7CUnkn

own%7CTWFpbGZsb3d8eyJWIjoiMC4wLjAwMDAiLCJQIjoiV2luMzIiLCJBTiI6Ik1haWwiLCJXVCI6M

n0%3D%7C3000&sdata=Y7HsLY01%2B5Em95N%2FkWmBCtltDGYjZMGMPyzscWSXbSk%3

D&reserved=0.

**References**

Armstrong, E. J., R. L. Tanner, and J. H. Stillman. 2019. High heat tolerance is negatively
correlated with heat tolerance plasticity in nudibranch mollusks. Physiological and
Biochemical Zoology 92:430-444.
Baker, J., A. Meade, M. Pagel, and C. Venditti. 2015. Adaptive evolution toward larger size in
mammals. Proceedings of the National Academy of Sciences 112:5093-5098.
Barley, J. M., B. S. Cheng, M. Sasaki, S. Gignoux-Wolfsohn, C. G. Hays, A. B. Putnam, S. Sheth et
al. 2021. Limited plasticity in thermally tolerant ectotherm populations: evidence for a
trade-off. Proceedings of the Royal Society B 288:20210765.
Barnett, A. G., J. C. Van Der Pols, and A. J. Dobson. 2005. Regression to the mean: what it is and
how to deal with it. International Journal of Epidemiology 34:215-220.
Chevin, L.-M., R. Gallet, R. Gomulkiewicz, R. D. Holt, and S. Fellous. 2013. Phenotypic plasticity
in evolutionary rescue experiments. Philosophical Transactions of the Royal Society B:
Biological Sciences 368:20120089.
Chevin, L.-M., R. Lande, and G. M. Mace. 2010. Adaptation, plasticity, and extinction in a
changing environment: towards a predictive theory. PLoS Biology 8:e1000357.
Comte, L., and J. D. Olden. 2017. Evolutionary and environmental determinants of freshwater
fish thermal tolerance and plasticity. Global change biology 23:728-736.
Crispo, E. 2007. The Baldwin effect and genetic assimilation: revisiting two mechanisms of
evolutionary change mediated by phenotypic plasticity. Evolution: International Journal
of Organic Evolution 61:2469-2479.
—. 2008. Modifying effects of phenotypic plasticity on interactions among natural selection,
adaptation and gene flow. Journal of Evolutionary Biology 21:1460-1469.

Deery, S. W., J. E. Rej, D. Haro, and A. R. Gunderson. 2021. Heat hardening in a pair of *Anolis*
lizards: constraints, dynamics and ecological consequences. Journal of Experimental
Biology 224:jeb240994.
Ehrenreich, I. M., and D. W. Pfennig. 2016. Genetic assimilation: a review of its potential
proximate causes and evolutionary consequences. Annals of Botany 117:769-779.
Esperk, T., A. Kjaersgaard, R. J. Walters, D. Berger, and W. U. Blanckenhorn. 2016. Plastic and
evolutionary responses to heat stress in a temperate dung fly: negative correlation
between basal and induced heat tolerance? Journal of Evolutionary Biology 29:900-915.
Ghalambor, C. K., K. L. Hoke, E. W. Ruell, E. K. Fischer, D. N. Reznick, and K. A. Hughes. 2015.
Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in
nature. Nature 525:372-375.
Ghalambor, C. K., J. K. McKay, S. P. Carroll, and D. N. Reznick. 2007. Adaptive versus non-
adaptive phenotypic plasticity and the potential for contemporary adaptation in new
environments. Functional Ecology 21:394-407.
Gunderson, A. R., M. E. Dillon, and J. H. Stillman. 2017. Estimating the benefits of plasticity in
ectotherm heat tolerance under natural thermal variability. Functional Ecology 31:1529-
1539.
Gunderson, A. R., and J. H. Stillman. 2015. Plasticity in thermal tolerance has limited potential
to buffer ectotherms from global warming. Proceedings of the Royal Society of London
B: Biological Sciences 282:20150401.
Gunter, H. M., R. F. Schneider, I. Karner, C. Sturmbauer, and A. Meyer. 2017. Molecular
investigation of genetic assimilation during the rapid adaptive radiations of East African
cichlid fishes. Molecular Ecology 26:6634-6653.
Jackson, D. A., and K. M. Somers. 1991. The spectre of 'spurious' correlations. Oecologia
86:147-151.
Kelly, C., and T. D. Price. 2005. Correcting for regression to the mean in behavior and ecology.
The American Naturalist 166:700-707.
Kelly, M. 2019. Adaptation to climate change through genetic accommodation and assimilation
of plastic phenotypes. Philosophical Transactions of the Royal Society B 374:20180176.
Kingsolver, J. G., and L. B. Buckley. 2017. Evolution of plasticity and adaptive responses to
climate change along climate gradients. Proceedings of the Royal Society B: Biological
Sciences 284:20170386.
Lande, R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic
plasticity and genetic assimilation. Journal of Evolutionary Biology 22:1435-1446.
Martin, C. H., J. E. Crawford, B. J. Turner, and L. H. Simons. 2016. Diabolical survival in Death
Valley: recent pupfish colonization, gene flow and genetic assimilation in the smallest
species range on earth. Proceedings of the Royal Society B: Biological Sciences
283:20152334.
Morley, S., L. Peck, J. Sunday, S. Heiser, and A. Bates. 2019. Physiological acclimation and
persistence of ectothermic species under extreme heat events. Global Ecology and
Biogeography 28:1018-1037.
Murren, C. J., J. R. Auld, H. Callahan, C. K. Ghalambor, C. A. Handelsman, M. A. Heskel, J.
Kingsolver et al. 2015. Constraints on the evolution of phenotypic plasticity: limits and
costs of phenotype and plasticity. Heredity 115:293-301.

Pfennig, D. W., M. A. Wund, E. C. Snell-Rood, T. Cruickshank, C. D. Schlichting, and A. P. Moczek.
467   2010. Phenotypic plasticity's impacts on diversification and speciation. Trends in Ecology
468   & Evolution 25:459-467.
469 Preston, J. C., R. Wooliver, H. Driscoll, A. Coughlin, and S. N. Sheth. 2021. Spatial variation in
470   high temperature-regulated gene expression predicts evolution of plasticity with climate
471   change in the scarlet monkeyflower. Molecular Ecology in press.
472 Price, T. D., A. Qvarnström, and D. E. Irwin. 2003. The role of phenotypic plasticity in driving
473   genetic evolution. Proceedings of the Royal Society of London. Series B: Biological
474   Sciences 270:1433-1440.
475 R Development Core Team. 2021. R: a Language and Environment for Statistical Computing.The
476   R Foundation for Statistical Computing, Vienna.
477 Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other
478   things). Methods in Ecology and Evolution 3:217-223.
479 Riddell, E. A., J. P. Odom, J. D. Damm, and M. W. Sears. 2018. Plasticity reveals hidden
480   resistance to extinction under climate change in the global hotspot of salamander
481   diversity. Science Advances 4:eaar5471.
482 Rohr, J. R., D. J. Civitello, J. M. Cohen, E. A. Roznik, B. Sinervo, and A. I. Dell. 2018. The complex
483   drivers of thermal acclimation and breadth in ectotherms. Ecology Letters 21:1425-
484   1439.
485 Sasaki, M. C., and H. G. Dam. 2019. Integrating patterns of thermal tolerance and phenotypic
486   plasticity with population genetics to improve understanding of vulnerability to warming
487   in a widespread copepod. Global Change Biology 25:4147-4164.
488 Schlichting, C. D., and M. Pigliucci. 1998, Phenotypic evolution: a reaction norm perspective,
489   Sinauer Associates Incorporated.
490 Schneider, R. F., and A. Meyer. 2017. How plasticity, genetic assimilation and cryptic genetic
491   variation may contribute to adaptive radiations. Molecular Ecology 26:330-350.
492 Seebacher, F., C. R. White, and C. E. Franklin. 2015. Physiological plasticity increases resilience
493   of ectothermic animals to climate change. Nature Climate Change 5:61-66.
494 Sikkink, K. L., R. M. Reynolds, C. M. Ituarte, W. A. Cresko, and P. C. Phillips. 2014. Rapid
495   evolution of phenotypic plasticity and shifting thresholds of genetic assimilation in the
496   nematode *Caenorhabditis remanei*. G3: Genes, Genomes, Genetics 4:1103-1112.
497 Somero, G. 2010. The physiology of climate change: how potentials for acclimatization and
498   genetic adaptation will determine 'winners' and 'losers'. The Journal of Experimental
499   Biology 213:912-920.
500 Stillman, J. H. 2003. Acclimation capacity underlies susceptibility to climate change. Science
501   301:65-65.
502 van Heerwaarden, B., and V. Kellermann. 2020. Does plasticity trade off with basal heat
503   tolerance? Trends in Ecology & Evolution 35:874-885.
504 Via, S., and R. Lande. 1985. Genotype-environment interaction and the evolution of phenotypic
505   plasticity. Evolution 39:505-522.
506 Waddington, C. H. 1953. Genetic assimilation of an acquired character. Evolution 7:118-126.
507 West-Eberhard, M. J. 2003, Developmental plasticity and evolution, Oxford University Press.
508