# Clathrin-mediated Endocytosis Facilitates Internalization of *Magnaporthe oryzae* Effectors into Rice Cells

Ely Oliveira-Garcia[1,2]*, Tej Man Tamang[1,3], Jungeun Park[3], Melinda Dalby[1], Magdalena Martin-Urdiroz[4], Clara Rodriguez Herrero[4,5], An Hong Vu[2], Sunghun Park[3], Nicholas J. Talbot[5] & Barbara Valent[1]*

[1]Department of Plant Pathology, Kansas State University, Manhattan, KS 66506, U.S.A.

[2]Department of Plant Pathology and Crop Physiology, Louisiana State University, Baton Rouge, LA 70803, U.S.A.

[3]Department of Horticulture and Natural Resources, Kansas State University, Manhattan, KS 66506, U.S.A.

[4]School of Biosciences, University of Exeter, Exeter, EX4 4QD, UK

[5]The Sainsbury Laboratory, Norwich Research Park, Norwich NR4 7UH, UK.

*Corresponding authors: E. Oliveira-Garcia; E-mail: EOliveiraGarcia@agcenter.lsu.edu and B. Valent; E-mail: bvalent@ksu.edu

## ABSTRACT

Many filamentous eukaryotic plant pathogens, such as fungi and oomycetes, deliver effectors into living plant cells to suppress defenses and control plant processes needed for infection. To date, little is known about the mechanism by which these pathogens translocate effector proteins across the plasma membrane into the plant cytoplasm. The blast fungus *Magnaporthe oryzae* secretes cytoplasmic effectors into a specialized biotrophic interfacial complex (BIC) before translocation. Here we show that cytoplasmic effectors within BICs are packaged into dynamic vesicles that can occasionally be found separated from BICs in the host cytoplasm. Live cell imaging with

fluorescently-labelled rice lines, showed that BICs are enriched in plant plasma membrane, actin, and Clathrin Light Chain-1, a marker for clathrin-mediated endocytosis (CME). We report that a novel cytoplasmic effector, Bas83, labels empty membrane vesicles surrounding BICs. Inhibition of CME using VIGS and chemical treatments results in a distinctive swollen BIC phenotype lacking effector vesicles. In contrast, fluorescent marker co-localization, VIGS and chemical inhibitor studies failed to implicate clathrin-independent endocytosis in effector vesicle formation. Taken together, this study provides evidence that cytoplasmic effector translocation is mediated by clathrin-mediated endocytosis in BICs and suggests a role for *M. oryzae* effectors in co-opting plant endocytosis.

**INTRODUCTION**

Many fungal and oomycete pathogens cause plant disease by hijacking and feeding on living plant cells, and they deliver effectors into and around host cells to promote infection (Giraldo and Valent, 2013; Lo Presti and Kahmann, 2017). This includes the Ascomycete fungus *Magnaporthe oryzae* (synonym of *Pyricularia oryzae*), which threatens global food security by causing blast diseases on rice, on several millets, and most recently on wheat (Gladieux et al., 2018; Valent et al., 2020). *M. oryzae* executes a hemibiotrophic lifestyle involving biotrophic invasion of successive plant cells by specialized intracellular invasive hyphae (IH) that grow from cell to cell (Kankanala et al., 2007). Blast IH growing in living host cells are enclosed by extensions of the plant plasma membrane termed the extra-invasive hyphal membrane (EIHM). Colonized cells die as they fill with hyphae and the EIHM and host vacuolar membranes become disrupted, while the fungus moves on to colonize neighboring cells using the same strategy (Kankanala et al., 2007; Mochizuki et al., 2015; Sakulkoo et al., 2018; Jones et al., 2021). Each successful IH undergoes hyphal morphogenesis and is associated with differentiated interfacial zones after entering the host cell lumen (Kankanala et al., 2007; Khang et al., 2010). Specifically, appressorial penetration pegs enlarge into tubular filamentous hyphae, called primary invasive hyphae, which are enclosed in EIHM with an extended dome-shaped cap, the Biotrophic Interfacial Complex (BIC). BICs accumulate cytoplasmic effectors destined for translocation into the host cytoplasm. Primary hyphae then transition to a bulbous pseudohyphal-like growth form. BICs at the tips of primary hyphae are left behind beside the first bulbous IH cell during the differentiation process but they continue to accumulate cytoplasmic effectors as the bulbous IH grow to fill the cell (Khang et al.,

2010). Mutation in an effector gene, Required-for-Focal-BIC-Formation 1 (*RBF1*), encoding a glycine-rich secreted protein, results in dispersed cytoplasmic effector localization around BIC-associated cells (the primary hyphae and first bulbous IH cell), perturbs primary hyphal structure, and decreases pathogenicity (Nishimura et al., 2016; Nishizawa et al., 2016).

In blast disease, cytoplasmic and apoplastic effectors are clearly differentiated by the fungal secretion system used and their localization pattern after secretion. Primary hyphae and the first bulbous IH focally secrete cytoplasmic effectors into BICs using a specialized, Golgi-independent secretion system that is insensitive to treatment with Brefeldin A (BFA), a chemical that inhibits conventional Golgi-mediated secretion in fungi (Giraldo et al., 2013; Van den Ackerveken, 2017; Wang et al., 2017). Fluorescent cytoplasmic effectors are routinely detected in the cytoplasm of invaded cells, and some are also detected in the cytoplasm of surrounding uninvaded host cells, apparently moving through plasmodesmata to prepare surrounding cells before invasion (Khang et al., 2010). In contrast, apoplastic effectors, such as the biotrophy-associated secreted protein 4 (Bas4), the LysM protein 1 (Slp1), and Bas113, are secreted via conventional golgi-mediated secretion, and they accumulate uniformly throughout the apoplastic compartment enclosed within the EIHM, outlining the entire IH without bleeding into the plant cell wall (Mosquera et al., 2009; Mentlak et al., 2012; Giraldo et al., 2013). Indeed, localization of secreted Bas4 fluorescent fusion proteins has become a reliable live cell imaging marker for the intactness of the EIHM around IH (Khang et al., 2010; Mochizuki et al., 2015; Jones et al., 2021). After filling the host cell, hyphal differentiation and BIC relocalization recur for each new hypha that enters a living neighbor cell (Khang et al., 2010; Sakulkoo et al., 2018).

The mechanism by which eukaryotic filamentous plant pathogens deliver effectors across the host plasma membrane inside living host cells is poorly understood and the mechanism is likely to vary in different systems (Giraldo and Valent, 2013; Petre and Kamoun, 2014; Fawke et al., 2015; Lo Presti et al., 2015; Lo Presti and Kahmann, 2017). Amino acid translocation motifs have been identified for oomycete effectors, with the RXLR motif being the most common, although cell entry motifs for fungal effectors have not been defined. The RXLR motif in oomycete effectors was reported to mediate host cell uptake through endocytosis based on binding to phosphatidylinositol-3-phosphate (PI3P) in host membrane (Kale et al., 2010), although it was recently shown that the RXLR motif from *Phytophthora infestans* effector AVR3a is cleaved off before secretion from the pathogen (Wawra et al., 2017). Therefore, the function of RXLR motif

in cell entry remains disputed (Petre and Kamoun, 2014; Trusch et al., 2018). Another amino acid

motif, YKARK, appears involved in translocation of the effector SpHtp3 from *Saprolegnia*

*parasitica*, a serious oomycete pathogen of fish, inside fish cells (Trusch et al., 2018). Specifically,

this effector appears to be internalized inside host cells via lipid-raft mediated endocytosis through

binding to a gp96-like receptor, and another effector is involved in releasing SpHtp3 from vesicles

into the fish cytoplasm. In an example from a fungal pathogen, a solvent-exposed RGD

vitronectin-like motif is required for internalization of ToxA, a host-specific proteinaceous toxin

secreted by the necrotrophic wheat pathogen *Pyrenophora tritici-repentis* (Manning and Ciuffetti,

2005). In addition to uptake by plant endocytosis, effectors might enter plant cells through

specialized translocon complexes in the plasma membrane. For example, the human malaria

parasite, *Plasmodium falciparum*, is contained within a parasitophorous vacuole inside invaded

red blood cells, and effectors are delivered across the membrane into the host cytoplasm by the

PTEX translocon (Elsworth et al., 2014). Recently, a stable protein complex comprised of five

unrelated fungal effectors and two fungal membrane proteins has been implicated in effector

translocation by the maize smut fungus *Ustilago maydis* (Ludwig et al., 2021). However, the

question of how effectors are taken up into plant cells remains largely unresolved for eukaryotic

plant pathogens.

Clathrin-mediated endocytosis (CME) is the major mechanism by which eukaryotic cells

internalize extracellular or membrane-bound cargoes, and this process is best understood in

mammalian systems and yeast (Kaksonen and Roux, 2018). Increasing understanding of CME in

plants shows both conserved and evolutionarily unique mechanistic details (Qi et al., 2018;

Reynolds et al., 2018; Johnson et al., 2020; Narasimhan et al., 2020). CME is known to play key

roles in plant-microbe interactions (Dodds and Rathjen, 2010; Beck et al., 2012; Fan et al., 2015).

CME is required for immunity mediated by pattern recognition receptor kinases, for instance,

specifically through internalization of activated extracellular pattern recognition receptors (PRRs)

for degradation in the vacuole (Mbengue et al., 2016). Examples include the pattern recognition

receptor Pep receptor 1/2 (PEPR1/2), the EF-TU receptor (EFR) recognizing the PAMP bacterial

elongation factor, and FLS2 recognizing the PAMP flagellin (flg-22) (Spallek et al., 2013;

Mbengue et al., 2016), as well as the Cf-4 receptor-like protein that recognizes the *C. fulvum* Avr4

avirulence effector (Postma et al., 2016). Trafficking of activated PRRs requires clathrin and

converges onto endosomal vesicles that are also shared by the hormone receptor Brassinosteriod

Insensitive 1 (BRI1). Additionally, it is known that pathogens can control plant CME. In addition
to its role in suppressing INF1-mediated cell death, the *P. infestans* host-translocated RXLR
effector AVR3a was recently shown to associate with the Dynamin-Related Protein 2, a plant
GTPase implicated in receptor-mediated endocytosis, and to suppress flg22-triggered defense
responses and reduce FLS2 internalization (Chaparro-Garcia et al., 2015). Therefore, AVR3a
interacts with a membrane-remodeling complex involved in immune receptor-mediated
endocytosis. Also, in host cells containing haustoria from the oomycete *P. infestans*, the late
endosomal pathway is rerouted from the vacuole to the extrahaustorial membrane (EHM)
surrounding haustoria (Bozkurt et al., 2015). Multiple effectors from this pathogen are now
implicated in rerouting the host membrane trafficking system and autophagy machinery for the
benefit of the invading pathogen (Bozkurt et al., 2015; Dagdas et al., 2018; Petre et al., 2021).

Effector translocation in the *M. oryzae*/rice pathosystem is easily assayed by live cell
imaging of fluorescently-labeled effector proteins inside host cells after secretion by IH invading
rice leaf sheath cells (Khang et al., 2010). Although some effectors naturally localize to host nuclei,
addition of a nuclear localization signal (NLS) to fluorescent effectors enhances visualization of
translocation and cell-to-cell movement. Blast cytoplasmic effectors apparently do not contain a
translocation motif as reported for oomycete effectors. Instead, preferential accumulation in the
BIC, which is a prominent feature of the blast fungus-rice cell interface, is the major predictor of
translocation into the rice cell. Previously, the Nishizawa lab reported that a cytoplasmic effector
is packaged in vesicles in the outer region of the BIC (Mochizuki et al., 2015; Nishimura et al.,
2016; Nishizawa et al., 2016). Here, we have used live-cell fluorescence imaging to characterize
the nature and dynamics of membrane-bound effector vesicles within BICs and in surrounding
host cytoplasm. We show that BICs are enriched for CME components, and we report functional
analyses using virus-induced gene silencing (VIGS) and pharmacological approaches. When
considered together, our data indicate that blast effectors are internalized through clathrin-
mediated endocytosis in the outer BIC layers, followed by escape from the vesicles into the rice
cytoplasm.

**RESULTS**
155
**BICs contain multiple vesicles labeled by cytoplasmic effectors**

BICs were previously reported as interfacial structures containing fluorescently-labelled
apoplastic effectors, such as Bas4:eGFP, in an inner base layer and a concentration of fluorescent
cytoplasmic effectors in the outer 'dome-shaped' region (Khang et al., 2010; Nishizawa et al.,
2016). We now routinely resolve the dome region of BICs by laser confocal imaging as a
concentration of vesicles containing fluorescent cytoplasmic effectors (Figure 1). These effector
vesicles are illustrated by an image of a BIC at the tip of a primary hypha (tip-BIC) expressing
fluorescent apoplastic effector Bas4:eGFP and fluorescent cytoplasmic effector Pwl2:mRFP
(Figure 1A). The Bas4:eGFP fluorescence remains evenly distributed throughout the EIHM matrix
around the growing IH, including in the inner layer of the BIC. In contrast, Pwl2:mRFP occurs as
distinct vesicles in the outer BIC region. We have observed effector vesicles in tip-BICs labeled
with additional translocated effectors Bas1 and Pwl1 fused to either eGFP or mRFP (Supplemental
Figure S1). Vesicles are also observed after tip-BICs move beside the first bulbous IH cells (side-
BICs) (Figure 1B). We evaluated the proximity of these vesicles relative to both tip- and side-
BICs. Whereas the vast majority of vesicles associated with primary IH were located in the tip-
BIC, some vesicles associated with side-BICs were found in regions of the cytoplasm at least 7
μm from the fungal cell wall at the BIC, which we define as outside the BIC (Figure 1C). Inclusion
of the apoplastic effector Bas4:eGFP in these experiments confirmed that the EIHM remained
intact at each infection site, based on its outlining of the IH and lack of spilling into the invaded
rice cell.

We identified a novel cytoplasmic effector, Bas170 (MGG_07348.6), that localized in
vesicles in the rice cytoplasm short distances from the appressorial penetration site at a stage before
primary hyphal growth is visible. The Bas170:mRFP fusion protein naturally accumulates in the
rice nucleus at this early infection stage, presumably to reprogram plant transcription essential for
*M. oryzae* infection (Supplemental Figure S2). These results suggest rapid effector translocation
from appressoria even before growth of primary IH. Bas170:mRFP also labels BIC-associated
vesicles in tip-BICs, and vesicles have been observed at a distance from the BIC in later stages.
Vesicles observed using fluorescent Bas170 fusion proteins are similar to vesicles observed with
Pwl2, Pwl1 and Bas1 fusion proteins except that these latter effectors were not observed under
appressoria at the earlier time period.

**Vesicles in BICs are dynamic through time**

To understand vesicle dynamics in BICs through time, we imaged BIC effector fluorescence during the morphogenic switch from tubular primary hyphae to bulbous IH, using a strain co-expressing two cytoplasmic effectors, Bas1:eYFP and Pwl2:mRFP (Figure 2). The vesicle content varied over the 2-hour period when the BIC was displaced from the primary hyphal tip to beside the differentiating bulbous IH cell. Tip-BICs often contained vesicles with either red or green fluorescence, rather than yellow, suggesting that effectors can be sorted into different vesicles. For this series of experiments, Pwl2:mRFP occurred in vesicles that reached outer layers of the tip-BIC compared to Bas1:eYFP. Pwl2:mRFP appeared less abundant relative to Bas1:eYFP during the switch from tip- to side-BIC, recovering again after growth of bulbous IH (Figure 2A). Both vesicle size and content varies within BICs through time (Figure 2 B-C). Specifically, BICs at primary hyphal tips contain a larger number of smaller vesicles (Figure 2D,E). Whereas BICs on primary hyphae deliver effectors in 'micro-vesicles' ranging from 41 nm to 114 nm, BICs on mature bulbous hypha contain effectors in 'macro-vesicles' ranging from 153 nm up to 2310 nm at times (Figure 2C,D). Effector vesicles beside mature bulbous IH often showed colocalization of fluorescence from different effectors compared to vesicles in tip-BICs (Figure 2A, B and F).

Cytoplasmic connections were observed between BICs and plant cytoplasm at the periphery of the invaded cell, illustrated in optical sections through a BIC in Supplemental Movie S1. This is consistent with a previous report of dynamic connections linking BICs and host cytoplasm underneath the appressorium (Khang et al., 2010). These cytoplasmic connections may facilitate delivery of effectors from cytoplasm surrounding the BIC in the cell center to the peripheral cytoplasm, in position to rapidly move through plasmodesmata into neighboring cells.

**BIC vesicles contain plant plasma membrane**

Effector-labeled vesicles in the BIC could arise through secretion of cytoplasmic effectors in exosomes by the fungus or through plant endocytosis. We previously reported that BICs contain material derived from plant plasma membrane based on infection of transgenic rice expressing a plasma membrane marker (Giraldo et al., 2013). Here we again use LTi6B:GFP-labeled rice to confirm that BICs contain abundant plant plasma membrane (Figure 3A,B). Individual effector vesicles were observed to be surrounded by labeled plant plasma membrane (Fig 3C), especially for vesicles in side-BICs on mature bulbous hyphae, where the vesicles reach diameters of 150 nm or larger.

To confirm that BIC vesicles contain plant plasma membrane, we evaluated the BIC

membrane composition by staining with the endocytosis tracker dyes FM 4-64 and FM 1-4 (Fig 3

D,E). These amphiphilic styryl dyes insert in the plasma membrane where they diffuse laterally in

the membrane and are only internalized into the cell via endocytosis (Atkinson et al., 2002; Bolte

et al., 2004). We previously showed that FM 4-64 is prevented from inserting in IH membranes,

which are sealed inside the EIHM (Kankanala et al., 2007). Rice treated with FM 4-64 and FM 1-

43 undergoing infection by *M. oryzae* strain KV177 or KV209 showed massive accumulation of

both dyes in BICs. Exclusion of these dyes from the hyphal membranes was documented by the

lack of fluorescent labeling of the fungal septa and vacuoles, which occurs if the dye inserts in the

fungal plasma membrane (Atkinson et al., 2002; Kankanala et al., 2007). Images of BICs stained

with FM 4-64 showed vesicles carrying effectors immersed in the region of intense FM 4-64

staining associated with the BIC (Figure 3E). Vesicles that carry effectors were also observed

separated from the BIC, presumably in the rice cytoplasm (Figure 3E, inset upper right), but

effector-containing vesicles have not been frequently observed at long distances from the BIC

(Supplemental Movie S2). Apparent vesicular bursting was also observed, which suggests that the

vesicles are unstable, presumably releasing the effector contents into the host cytoplasm.  Taken

together, these results indicate that BIC-associated effector vesicles are more likely derived

through plant endocytosis than from secretion of fungal exosomes.

**Bas83 is a unique cytoplasmic effector that associates with rice membrane concentrating**

**around BICs**

Since it is likely that specific blast effectors play roles in vesicle formation and effector

translocation, we were particularly interested in effectors that appear to associate with membranes

in host cells. The *BAS83* (MGG_08506.6) gene is unique to *M. oryzae*, and it is 36-fold upregulated

during biotrophic invasion (Mosquera et al., 2009). Bas83 is a cytoplasmic effector that is secreted

by the fungus using the Golgi-independent secretion pathway identified for other cytoplasmic

effectors (Giraldo et al., 2013). Bas83:mRFP is translocated into invaded host cells, but it does not

precisely colocalize with other BIC-localized cytoplasmic effectors Pwl2:eGFP and Pwl1:eGFP

(Figure 4A; Supplemental Figure S3), and it does not move ahead into uninvaded neighboring

cells as these other effectors do. In contrast, Bas83:mRFP seems to label the EIHM surrounding

the BIC-associated cells, the primary hyphae and first bulbous IH, and it labels membrane bubbles,

or vesicles, near to BICs and BIC-associated cells (Figure 4A-C). Figure 4A shows that

Bas83:mRFP-labeled vesicles do not contain the cytoplasmic effector Pwl2:eGFP, indicating they
are distinct from effector vesicles in BICs. Bas83:mRFP is also associated with vesicles observed
under appressoria (Figure 4C; Supplemental Figure S3). However, Bas83:mRFP fails to label
membrane surrounding subsequently formed bulbous IH that are associated with apoplastic
effectors and not with cytoplasmic effectors (see 2nd bulbous IH in Figure 4C). We hypothesize
that Bas83 is involved in recruiting new plant plasma membrane to form the EIHM and BICs
during biotrophic development. However, repeated attempts in two different laboratories (264
purified transformants, 4 methods and 5 experiments; KSU and Univ. of Exeter) failed to produce
knock-out mutants for further functional analysis.

## BICs contain plant actin
The actin cytoskeleton is essential for endocytic processes including plasma membrane folding,
vesiculation, and vesicle transport (Grassart et al., 2014; Narasimhan et al., 2020). Effector vesicle
formation in BICs suggests that this interfacial structure should contain abundant host actin.
Therefore, we imaged plant actin dynamics during biotrophic invasion using rice lines expressing
the actin marker LifeAct:eGFP. Infection by *M. oryzae* strains expressing chimeric protein
Pwl2:mRFP showed massive colocalization of LifeAct:eGFP in BICs (Figure 5A). Actin was
abundant in all imaged tip- and side-BICs (Figure 5B). Likewise, colocalization of fluorescent
Phalloidin conjugates and fluorescently-labeled effector confirmed that BICs are rich in plant actin
filaments (Figure 5C). We hypothesize that actin filaments are responsible for plasma membrane
manipulation, vesiculation and possibly for the transport of those vesicles carrying effectors into
the rice cell cytoplasm.

## BICs contain abundant plant clathrin, but not flotillin
Clathrin-mediated endocytosis (CME) is the major endocytosis mechanism in plants, with some
contribution by Clathrin-independent endocytosis (CIE) (Chen et al., 2011; Ewers and Helenius,
2011; Fan et al., 2015; Narasimhan et al., 2020). To assess co-localization of specific endocytosis
markers with BICs, we generated transgenic rice lines expressing a fluorescent marker for either
CME or CIE. For CME, we generated a C-terminal translational fusion of eGFP with the rice
Clathrin light chain-1 (OsCLC1), which is a component of the clathrin coat together with clathrin
heavy chain. Rice lines expressing OsCLC1:eGFP show typical clathrin foci (Konopka et al.,
2008; Narasimhan et al., 2020) in uninfected rice cells (Figure 6A). The marker for CIE was a C-

terminal translational fusion of eGFP to flotillin (Flot1), which is a component of lipid rafts (Otto

and Nichols, 2011; Li et al., 2012). Rice lines with the CME and CIE markers were used for leaf

sheath assays with an *M. oryzae* strain expressing Pwl2:mRFP to label BICs. In addition to typical

clathrin foci around the cell periphery, all tip-BICs and side-BICs contain fluorescence from

OsCLC1:eGFP co-localizing with Pwl2:mRFP (Figure 6 A,B). Clathrin fluorescence could

sometimes be observed co-localizing with effector fluorescence in BIC vesicles (Supplemental

Movie S3). In contrast, no tip-BICs colocalized with Flot1:eGFP. Some colocalization of

Flot1:eGFP and Pwl2:mRFP occurred in side-BICs besides mature bulbous hyphae (Figure 6B).

Colocalization results from several independent experiments were consistent with a major role for

CME, but not for CIE, in the internalization of cytoplasmic effectors.

**Silencing of rice endocytosis machinery blocks *M. oryzae* infection**

We assessed the impact of inhibition of CME and CIE on BICs and effector vesicles using the

brome mosaic virus system (Ding et al., 2006; Ding et al., 2007) for silencing of genes for CME

and CIE components that have been characterized by gene silencing analyses in *Arabidopsis*.

Specifically, we silenced two rice CME genes, Adaptor protein complex-2 subunit $\alpha$2 (*AP-2$\alpha$*) and

clathrin heavy chain-1 (*CHC1*) (Di Rubbo et al., 2013; Larsson et al., 2017) as well as the CIE

gene *Flotillin-1* (*Flot1*) (Li et al., 2012). Silencing of both *AP-2$\alpha$* and *CHC1* resulted in severe

reduction in pathogenicity in whole plant spray inoculation assays (Figure 7 A,B). However, as

expected, silencing of these genes caused stunting and decreased plant health (Supplemental

Figure S4), and levels of pathogenicity in blast disease are known to be decreased on unhealthy

rice plants. More significant is the phenotype of pathogen blockage at individual infection sites.

Inhibition of clathrin-mediated endocytosis led to a distinctive swollen BIC phenotype, with

enlarged, abnormally-shaped BICs that lack effector vesicles. By contrast, silencing of the CIE

marker gene *Flot1* showed minor effects on pathogenicity (Figure 7A,B), and had little impact on

BIC structure [Figure 7C,F; compare 26 hpi RNAi*Flot1* (CIE) and RNAi*AP-2$\alpha$* (CME)].

In addition to swollen BICs as sites with primary hyphae, silencing of both CME genes

decreased appressorial penetration and induced the accumulation of the cytoplasmic effector

Pwl2:mRFP under failed penetration sites where it is not normally observed (Figure 7D,E). In

contrast, silencing of Flot1 had only minor effects on appressorium penetration success and on

accumulation of Pwl2:mRFP under failed appressoria. Like our finding of early accumulation of

Bas170 under appressoria before penetration, these results again suggest that effector uptake

begins before host penetration. These results also suggest that effector uptake at the appressorial

penetration stage and from BICs is mediated by CME.

**Chemical inhibition of rice endocytosis machinery blocks *M. oryzae* infection**

Pharmacological approaches have impacted studies in diverse biological systems (Robinson et al.,

2008; Hicks and Raikhel, 2010; Grassart et al., 2014; Fan et al., 2015). To further assess potential

impact of the plant endocytic machinery on cytoplasmic effector translocation, we tested a series

of chemicals that are reported to inhibit plant CME or CIE (Table 1). Compared with the non-

treated control (Figure 8A), treatment with reported CME inhibitor cantharidin led to abnormally-

shaped, swollen BICs resembling those seen with silencing of CME components (Figure 8B).

Fluorescence intensity linescans confirm that cantharidin treatment had no impact on localization

of apoplastic effector Bas4:eGFP in the base of the BIC.

Subsequent quantification of cantharidin treatment along with treatment of 2 additional

reported CME inhibitors, chlorpromazine and triclosan (Table 1), showed a high proportion of

swollen, irregular-shaped cytoplasmic effector fluorescence patterns in BICs (Figure 9A-C). By

contrast, inhibition of CIE with filipin and methyl-β-cyclodextrin resulted in minor to no effect on

BIC effector fluorescence patterns (Figure 9A-C). Two chemical inhibitors with potential CME

association (Table 1) show different impacts on BIC effector fluorescence patterns. Wortmannin,

which inhibits phosphatidylinositol 3-kinase (PI 3-kinase) and is implicated in late endosomal

trafficking (Robinson et al., 2008) had a major impact on BIC structure, while concanamycin A,

which inhibits vacuolar-type ATPase and induces vacuolar acidification (Robinson et al., 2008),

had little impact on BIC structure (Figure 9A-C).

Moreover, treatment with CME inhibitors cantharidin, chlorpromazine and triclosan, but

not with CIE inhibitors filipin and methyl-β-cyclodextrin, resulted in reduced appressorial

penetration and induced abnormal accumulation of the cytoplasmic effector Pwl2:mRFP under

appressorial penetration sites where it is not normally observed (Figure 9A - compare 18 hpi CIE

and CME inhibitors; Figure 9D,E). Wortmannin, but not concanamycin A, also impacted

appressorial function and resulted in abnormal Pwl2:mRFP accumulation under appressoria. These

results support our findings with Bas170:mRFP (Supplementary Figure 2) in suggesting that

effector uptake begins before or at the point that host penetration occurs. Indeed, chemical

inhibition of endocytosis in rice cells undergoing *M. oryzae* infection mimics the results obtained

with VIGS, further supporting a role for CME in effector uptake into rice cells.

**DISCUSSION**

Our results, when taken together, provide strong evidence that cytoplasmic effector translocation is BIC-localized and mediated by CME. We report that diverse cytoplasmic effectors occur in dynamic vesicles in tip-BICs associated with primary hyphae that first colonize host cells and then in BICs that remain beside the first bulbous IH after hyphal differentiation. Tip-BICs contain many small effector micro-vesicles (~87 nm) and co-expression of differentially-labeled effectors Bas1:eYFP and Pwl2:mRFP showed these effectors can be sorted into different vesicles (Figure 2). Vesicle composition can be seen to vary as tip-BICs move to become side-BICs during hyphal differentiation. Specifically, side-BICs contained fewer, larger vesicles that reach >500 nm in diameter and show co-localization of Bas1:eYFP and Pwl2:mRFP. Our results confirm a previous report that cytoplasmic effectors occur in ~500 nm vesicles in the outer BIC domes beside bulbous IH (Mochizuki et al., 2015; Nishimura et al., 2016; Nishizawa et al., 2016). We also observed effector vesicles in the host cytoplasm at short distances from maturing side-BICs (Figure 2). Release of effectors from these vesicles into the host cytoplasm would require a special mechanism, perhaps mediation by an effector as suggested for release of the *S. parasitica* effector from endocytic vesicles into the fish cell cytoplasm (Trusch et al., 2018).

We report that BICs producing effector vesicles contain multiple hallmarks of active clathrin-mediated endocytosis (CEM), including focal accumulation of fluorescently-labelled Clathrin Light Chain-1 (CLC1), a key subunit of the clathrin coat in CME (Figure 6). Focal accumulation of actin in the BICs, as detected in stable transgenic plants expressing LifeAct:eGFP and by staining with rhodamine-labeled phalloidin (Figure 5), is consistent with a role in CME, although it was recently reported that actin plays a different role in CME in plants than in the well-studied systems in mammals and yeast (Qi et al., 2018; Johnson et al., 2020; Narasimhan et al., 2020). We confirm that BICs contain abundant rice plasma membrane (Giraldo et al., 2013) using transgenic plants expressing fluorescent plasma membrane fusion protein LTi6b:eGFP, and using the amphiphilic stryryl dyes FM 4-64 or FM 1-43 to track membranes that are internalized by endocytosis (Bolte et al., 2004). Indeed, we obtained occasional views of effector vesicles ringed in LTi6b:eGFP or FM 4-64 labeled membrane (Figure 3C). Interestingly, we identified a novel effector, Bas83, that labels non-effector vesicles and EIHM focused around BICs and the BIC-associated cells, which could be involved in replenishing host membranes needed for extensive

endocytotic activity in the BIC (Figure 4; Supplemental Figure S3). In contrast, flotillin (Flot1:eGFP), which is a component of lipid raft-mediated clathrin independent endocytosis, rarely co-localizes with tip-BICs where active translocation is occurring (Figure 6C). Our fluorescent marker co-localization experiments are all consistent with BICs as a focused area of CME.

For functional analyses of the role of endocytosis in effector translocation, we performed transient silencing of two rice CME components, Clathrin Heavy Chain-1 (CHC-1) and Adaptor Protein Complex subunit 2α (AP-2α), as well as silencing of the CIE component Flotillin-1. We also used 7 pharmacological agents that are reported to inhibit or be associated with CME or CIE endocytosis in plants. Both strategies have potential problems. Rice plants that received the CHC-1 and APC-2 silencing vectors were significantly reduced in rice blast symptom, compared to the empty vector control and to Flot1-silenced lines (Figure 7; Supplemental Figure S4). However, this could be due, at least in part to inevitable health costs to the plant through silencing component genes for essential cellular processes. Unhealthy rice plants are generally less susceptible to blast disease (Kato, 1974). Additionally, care must be taken in the interpretation of results in pharmacological studies based on new evidence that endocytosis inhibitors characterized in mammalian systems may not have the same targets in plants (Dejonghe et al., 2016), and pharmacological agents also have off-target effects (Hicks and Raikhel, 2010; Dejonghe and Russinova, 2017). However, both studies complement each other in that transient silencing of the two CME components and treatment with 4 different pharmacological agents associated with CME gave the same swollen BIC phenotype, in which labeled fluorescent effectors accumulate in abnormally-sized BICs lacking the discreet effector vesicles seen in control infection sites (Figure 8 and 9). In contrast, transient silencing of the CIE component Flotilin-1 and treatment with pharmacological inhibitors associated with CIE show little or no impact on BIC vesicular structure and pathogenicity (Figure 8 and 9). All results from our VIGS and chemical inhibitor experiments are consistent with our fluorescence localization experiments. Taken together, our results therefore provide evidence that BICs include a focused region of plant clathrin-mediated endocytosis for internalizing cytoplasmic effectors inside living plant cells (Figure 10).

We report two new lines of evidence that effectors are being secreted and translocated into the host cytoplasm from appressoria even before growth of primary hyphae inside host cells. First, we identified an effector, Bas170, that naturally accumulates in host nuclei before visible growth of primary hyphae, and it also localizes to vesicles underneath appressoria at this early time point.

Second, effector Pwl2:mRFP, which is not normally observed before the tip-BIC stage,
accumulates under appressoria that failed to penetrate after both VIGS silencing and chemical
inhibition of rice CME. These new findings are consistent with reports that rice cells recognize
and respond to *M. oryzae* before and independently of appressorial penetration (Xu et al., 1998).
It has been predicted that *M. oryzae* appressorial pores, a transient cell wall-less region of the
appressorium adjacent to the plant cuticle, might be involved in molecular communication between
pathogen and host before penetration (Howard and Valent, 1996). Previously, the only *M. oryzae*
effector that was a candidate for secretion through the appressorial pore was a presumed secondary
metabolite that is produced by the *AVR* gene *ACE1* encoding a hybrid polyketide synthase (PKS)
and nonribosomal peptide synthetase (NRPS) that is specifically expressed in the cytoplasm of
appressoria before and during penetration (Fudal et al., 2007; Collemare et al., 2008). Therefore,
our new effector localization results suggest similar exquisite staging of effector expression as
reported for the crucifer pathogen *Colletotrichum higginsianum*, which secretes ChEC effectors
under the appressorial pores and at additional invasion stages during the invasion of
*Arabidopsis thaliana* cells (Kleemann et al., 2012). Specifically, this fungus focally secreted
effectors Chec6 and Chec36 from the appressorium pore, suggesting these effectors may be
translocated before fungal penetration (Kleemann et al., 2012). Our results with Bas170 together
with ChECs and Ace1 protein localization patterns indicate that many effectors might be focally
secreted through appressorial pores (Fudal et al., 2007; Kleemann et al., 2012), highlighting a
potential role for fungal appressoria in effector delivery.

The effector vesicle dynamics that we report support the idea that front-loaded effector
translocation occurs during the early stages of host cell invasion. Specifically, tip-BIC vesicles are
small (~87 nm), abundant and selective in terms of sorting two cytoplasmic effectors, Bas1:eYFP
and Pwl2:mRFP, into different vesicles. In contrast, side-BICs contain fewer, larger, less-selective
vesicles that contain both Bas1:eYFP and Pwl2:mRFP. Effector vesicles observed in the cytoplasm
were almost uniformly found at the side-BIC stage, perhaps indicating a slow-down of the
mechanism for effector release from the vesicles as BICs mature. The mechanism for forming
vesicles may result in larger vesicles, or vesicles may fuse before releasing effectors to the
cytoplasm in the later BIC stages. Slowdown of the translocation system would appear consistent
with the loss of activity in BICs as host cells fill with IH. Front-loading effector translocation from
BICs would make sense because cell-to-cell movement of blast cytoplasmic effectors through

plasmodesmata must occur in the early stages of host cell invasion while the EIHM enclosing the IH remains intact and plasmodesmata in the invaded cell remain open (Sakulkoo et al., 2018; Jones et al., 2021). Active BICs are surrounded by host cytoplasm with dynamic connections to the cell periphery (Khang et al., 2010) (Supplemental Movie S2), where cytoplasmic streaming would rapidly disperse effectors to pit fields containing plasmodesmata. In contrast, nonBIC-associated IH cells invaginate the host vacuole and grow in close proximity to the host vacuolar membrane with little surrounding cytoplasm (Mochizuki et al., 2015; Jones et al., 2021). Specifically, the host vacuole remains intact, but shrinks as IH grow. Both the EIHM and vacuolar membrane become disrupted, indicating death of the invaded cell, around the time IH move into neighboring cells (Mochizuki et al., 2015; Jones et al., 2021). However, by this time, IH have established themselves in living neighboring cells.

There is evidence supporting occurrence of clathrin-mediated endocytosis at the EHM in one additional pathosystem. That is, transmission electron micrographs of the interfacial matrix between the intracellular hyphae [called monokaryotic (M)-haustoria] of the monokaryotic rust fungus *Uromyces vignae* and its host cowpea (*Vigna unguiculata*) shows clathrin-coated pits (diameter ca. 50-70nm) on the EHM at M-haustoria hyphal tips that contained dense cytoplasm and were presumably growing (Stark-Urnau and Mendgen, 1995). The coated vesicles were labeled by an antibody recognizing the heavy chain subunit of clathrin. Long tubules extending into the host cytoplasm, as well as coated vesicles in the surrounding host cytoplasm, were also labeled with the antibody, indicating that endocytosis occurs at the EHM (Stark-Urnau and Mendgen, 1995; O'Connell and Panstruga, 2006). It was suggested that these endocytic vesicles might be involved in effector uptake into host cells in addition to membrane recycling from host exocytosis. Unfortunately, lack of transformation capability and identification of effectors makes it difficult to follow up on these intriguing results with *U. vignae*. Additional TEM studies support our results that transient silencing of lipid raft-mediated endocytosis component Flotilin1, as well as filipin treatment, had little effect on BIC structure and effector vesicle formation. Freeze-fracture transmission electron microscopy (TEM) following filipin treatment revealed an absence of granular filipin-sterol complexes on the EHM of two rust fungi, *Puccinia coronata* and *Uromyces appendiculatus* (Harder and Mendgen, 1982), suggesting that the extrahaustorial membrane contains less sterol than normal plasma membranes, indicating that the EHM appears depleted in sterol-rich lipid rafts.

Differences in infection strategies among various filamentous biotrophic or hemibiotrophic

pathogens would be consistent with use of different strategies for effector translocation in different

pathosystems. A recent report indicates that a complex of seven proteins from the smut pathogen

*U. maydis* is critical for infection and implicated in translocation of cytoplasmic effectors into

maize cells (Ludwig et al., 2021). This basidiomycete fungus colonizes host tissue as intracellular

hyphae that pass through and between host cells without obviously specialized interfacial zones,

and fluorescent fusion proteins of cytoplasmic effectors have not been visualized in the host cell

cytoplasm in this pathosystem (Lo Presti et al., 2017). *U. maydis* produces galls on maize seedlings

and grows systemically as the plant grows to finally invade ovaries, which swell into dramatic

sporulating galls. In contrast, *M. oryzae* densely colonizes a localized tissue area that will become

a visible eyespot lesion that releases spores to reinitiate the infection cycle within 7 days. Different

pathogenic lifestyles could have different demands for cytoplasmic effector translocation.

Taken together, our data strongly suggest that clathrin-mediated endocytosis occurring in

BICs is a key mechanism for internalization of *M. oryzae* effectors inside living rice cells,

beginning even before appressorium penetration on the host surface. Important unknowns for

future study include the molecular mechanisms of how the endocytic machinery interacts with

effector cargos (Qi et al., 2018), including how different effectors are sorted out into different

vesicles as shown in Figure 2. It is important to understand vesicle dynamics through time and

why effector vesicles become larger and less selective as BICs mature. It is important to understand

if, and if so, how, Bas83 contributes to membrane recruitment to BICs and the EIHM surrounding

BIC-associated cells.  It is unknown how the effector vesicles are disrupted, releasing effectors

into the rice cytoplasm and how various effectors are then targeted to their destinations in host

cells including, in some cases, movement through plasmodesmata into neighboring host cells. It is

also important to understand the role for appressoria in effector secretion before penetration, and

how these early effectors, such as the host nuclear localized effector Bas170, function to promote

infection. Finally, it is critical to translate molecular mechanisms of blast biotrophic invasion into

strategies for controlling devastating diseases on rice, wheat and other food crops worldwide.

**METHODS**

**Live-cell imaging of *M. oryzae* effectors *in planta***

Fungal strains were stored in dried filter papers at -20˚C, and cultured on rice polish agar plates at

25˚C under continuous light for 2 weeks (Valent and Chumley, 1991). Rice sheath inoculations

were performed as described (Kankanala et al., 2007). The susceptible rice variety YT-16 was

used unless mentioned otherwise. Briefly, 7-cm-long sheath pieces from 3-week-old plants were

placed in a sealable Pyrex glass moist chamber. Leaf sheath sections were placed on inverted 8-

well PCR tube strips to avoid contact with wet paper and to hold epidermal cells directly above

the mid-vein horizontally flat for uniform inoculum distribution in the trimmed sheath pieces. A

spore suspension ($10^4$ spores/ml in sterile 0.25% gelatin, Cat. #G-6650, Sigma-Aldrich) was

prepared from 10-day old cultures and was injected into one end of the sheath using a 100-ml

pipette. Each segment was trimmed at 18–30 h.p.i., treated with specific dyes or inhibitors, or

imaged immediately by laser confocal microscopy. Confocal imaging was performed with a Zeiss

LSM780 confocal microscope system using two water immersion objectives, C-Apochromat

40x/1.2 WCorr. and C-Apochromat 63x/1.2WCorr. Excitation/emission wavelengths were 488

nm/505–550 nm for eGFP and FM1-43, and 543 nm/560–615 nm for mRFP, mCherry, Phalloidin

and FM4-64. Images were acquired and processed using Zeiss ZEN 2010 software. Images of

effector vesicles in Figure 3C were obtained using a Leica SP8 confocal microscope system with

a water immersion objective HC PL APO 63x/1.20 WCorr. Excitation/emission wavelengths were

488 nm/505–550 nm for eGFP and 543 nm/560–615 nm for mRFP.

**Fungal strains, DNA manipulation, and fungal transformation**

*Magnaporthe oryzae* wild type strain Guy11, a field isolate from rice in French Guiana, was

obtained from J.L. Notteghem (Centre de Cooperation Internationale en Recherche Agronomique

pour le Developpment). *M. oryzae* transformants are described in Supplemental Table S1 online.

Effector:mRFP and Effector:eGFP expression plasmids were constructed by PCR amplifying

different effector gene regions and fusing them to the N-terminus of mRFP and eGFP, respectively.

Details of construction of each plasmid are described in Supplemental Table S2, and primers are

listed in Supplemental Table S3. For all fusion constructs, transcriptional and translational fusions

were verified by DNA sequencing. See Supplemental Table S2 online for plasmids used, and

Supplemental Table S3 for PCR primers used. Plasmids were transformed into laboratory strain

Guy11 (Leung et al., 1988) using *Agrobacterium tumefaciens*–mediated transformation (Khang et

al., 2006). In several cases, two fluorescently labelled effectors were introduced by co-

transformation with separate plasmids (Sweigard et al., 1995). Because positive transformants

showed similar fluorescence patterns with varying intensities, those with strongest fluorescence

were studied (see Supplemental Table S1 online). For each construct cloned in *M. oryzae*, at least

two independent transformants showing identical phenotypes to the wild type were selected and

assayed together.

**Targeted deletion of Bas83 gene**

*Split-marker recombination method for Bas83 gene knockout via protoplast transformation*:

Targeted gene replacement mutation of the *M. oryzae Bas83* gene was attempted using the split

marker strategy as modified by Kershaw and Talbot (2009). Gene replacement was performed by

replacing the 606-bp *Bas83* locus with a hygromycin resistance selectable marker, encoding a 1.4-

kb hygromycin phosphotransferase (*HPH*) resistance cassette (Carroll et al., 1994). The two

overlapping segments of the *HPH* templates were PCR amplified using primers M13F with HY

and M13R with YG (Catlett et al., 2003) (see Supplemental Table S3 online) as described

previously (Kershaw and Talbot, 2009). One-kb DNA fragments upstream and downstream of the

*Bas83* open reading frame were generated using the primers Bas83LF-F and Bas83LFHY-R and

Bas83RFYG-F and Bas83RF-R amplified from genomic DNA of the Guy11 strain. A second-

round PCR reaction was performed to fuse the overlapping split hph marker templates with the

left and right flanking regions of the Bas83 locus. Wild-type *M. oryzae* strain Guy11 was then

transformed with the deletion cassettes (2 μg of each flank). Putative transformants were selected

in the presence of hygromycin B (200 μg/ml) and analyzed by PCR using the primers

Bas83:BASKOtest-F and Bas83:BASKOtest-R (Supplemental Table S3; Supplemental Figure

S5). This assay was repeated three times. Gene sequences on either side of *Bas83* were retrieved

from the NCBI database (https://www.ncbi.nlm.nih.gov/).

*Attempted Bas83 deletion via Agrobacterium-mediated transformation*: For gene

replacement transformation, cassettes were constructed by amplifying ∼1.0 kb of 5′- and 3′-

flanking regions for each predicted coding sequence. The *HPH* gene (Carroll et al., 1994) was

cloned between the two flanking regions using a fusion PCR strategy. The three pieces together

were cloned first into the pJET1.6 vector (Invitrogen) for sequence analysis and later into binary

vector pGKO2 (Addgene: Plasmid #63617; Khang et al., 2005) using a restriction ligation strategy.

*M. oryzae* KU70 and KU80 spores were transformed using *A. tumefaciens* (Khang et al., 2005)

strains AGL1 and EHA105 (Hellens et al., 2000). After two rounds of selection in TB3 media

containing 200 μg/mL of hygromycin, 400 independent fungal transformants were analyzed for

gene replacement events by PCR amplification using the primers Bas83:BASKOtest-F and

Bas83:BASKOtest-R (Supplemental Figure S5). This assay was repeated three times.

**558 Staining with FM 4-64, FM 1-43, and Rhodamine Phalloidin, and treatment with**

**559 pharmacological inhibitors**

To examine membrane dynamics at the BIC, rice sheaths (cv. YT16) were inoculated with *M.*

*oryzae* transformants expressing fluorescently-labeled effectors ($2x10^4$ spores/ml in 0.25% gelatin

solution) and incubated for 20-26 hours at 25˚C for subsequent FM-dye treatment. Infected rice

sheaths were treated with FM 4-64 or FM 1-43 (4 mg/ml in water). An aqueous 17mM stock

solution of FM 4-64 (Cat #13320, Invitrogen, Carlsbad, CA) and FM 1-43 (Cat # T3163,

Invitrogen, Carlsbad, CA) were prepared and stored at -20˚C as described (45). *In planta*,

inoculated leaf sheaths, 24 h.p.i., were trimmed into ultrathin sections (2-3 layers of cells; 10 x 10

mm) and incubated in a 10-mM aqueous working solution for 3–5 hours. Leaf sheaf sections were

subsequently transferred to ultrapure water and incubated for 20 min (25 ˚C) to remove excess

dyes for uniform membrane staining. For *in planta* experiments with Rhodamine Phalloidin (Cat#

R415, ThermoFisher Scientific), we prepared working solutions of 66 µM in 0.1% DMSO and

treated the inoculated tissue as described for FM dyes.

To examine the effects of endocytosis inhibitors on *in planta* effector uptake, inoculated

trimmed rice sheaths were incubated at 25˚C with Methyl-β-cyclodextrin (20 mM), or

Chlorpromazine (10 µg/mL), or Cantharidin (25 mM), or Fluazinam (20 mM), or Triclosan

(20mM), or Wortmannin (20 nM), or Concanamycin A (50 mM) (all from Sigma) solution for 1-

5 hours. Negative controls were performed with ultrapure water (Methyl-β-cyclodextrin,

Chlorpromazine, and Fluazinam) or 0.1% DMSO (Cantharidin, Filipin and Wortamanin) (see

Table 1).

**579 Generation of transgenic rice plants**

Production of transgenic plants expressing the plant plasma membrane marker low-temperature

inducible protein 6B, LTi6B:GFP was previously described (Mentlak et al., 2012; Giraldo et al.,

2013). Transformants of rice cultivar Sasanishiki expressing LifeAct:eGFP were generated in the

laboratory of NJT. Expression plasmids for CME and CIE marker genes were constructed by PCR

amplifying the marker gene regions and fusing them to the N-terminus of eGFP. The *CLC1* gene

was amplified from genomic DNA of YT16 rice using the primers KpnI_OsCLC1Prom1-F1 and

XhoI_OsCLC1-R1. The PCR product was digested with *Kpn*I and *Xho*I and integrated into

pSH1.6EGFP (Plasmid #42323, Addgene). The *Flot1* gene was amplified from genomic DNA

from YT16 rice using the primers HindIII_OsFlot1Prom1-F1 and Eco47III_OsFlot-R1. The PCR

product was digested with *Hind*III and *Eco*47III and integrated into pSH1.6EGFP. CLC1:eGFP

was amplified from pSH1.6_CLC1:eGFP using the primers CACC_OsCLC1-F1 and eGFPT2-R2,

and cloned into pENTR (pENTR™/D-TOPO™ Cloning Kit, ThermoFisher Scientific) following

the manufacturer's protocol. Flot1:eGFP was amplified form pSH1.6_Flot1:eGFP using the

primers CACC_OsFlot1-F1 and eGFPT2-R2, and cloned into pENTR following manufacturer's

protocols. Endocytosis marker:eGFP construct were cloned into the *Agrobacterium* vector

pIPKb001 (Himmelbach et al., 2007) using the Gateway LR Clonase II kit (Invitrogen) following

the manufacturer's protocol. All the fusion constructs had transcriptional and translational fusions

verified by DNA sequencing. Gene fusions for expressing the rice endocytosis markers

CLC1:eGFP (CME) and Flot1:eGFP (CIE) under their native promoters were introduced into

*Agrobacterium tumefaciens* EHA105 using the freeze-thaw method (Holsters et al., 1978). Mature

seed-derived callus from rice (*Oryza sativa L. Japonica*) cv. YT16 was used for *Agrobacterium*-

mediated transformation (Park et al., 2001). After inoculating with *A. tumefaciens*, callus was

transferred to regeneration medium for 4-10 weeks at 25˚C under a 16-h photoperiod. The

regenerated shoots were transferred to rooting medium for four more weeks, then established in

soil. Putative transformants of rice were selected on 100 μg/ml hygromycin, and expression

checked by qRT–PCR and epifluorescence microscopy.

**Silencing of rice genes**

In order to evaluate the role of plant endocytosis on disease development and effector uptake, we

performed VIGS using the Brome mosaic virus system developed by R.S. Nelson (Noble

Foundation, Ardmore, OK). We targeted *AP-2α*, *CHC1* (CME) and *Flot1* (CIE) mRNA in rice

plants (cultivar IR64) undergoing infection. The target rice sequences were cloned in the pC13/F3-

13m VIGS vector as described (Ding et al., 2006; Ding et al., 2007) and subsequently sequenced

to verify sequence accuracy. RNA transcripts were synthesized from the final VIGS vector. An

empty vector pC13/F3-13m was used as negative control. The final infectious transcripts of

pC13/F3-13m / AP-2α, pC13/F3-13m / CHC1, pC13/F3-13m /Flot1 and pC13/F3-13m (a control

virus without an insert) were inoculated with pC13/F1+2 in 7 to 10 day old IR64 rice plants

according protocols described by Ding et al (2006). Knockdown of the target genes was confirmed

by qRT-PCR at 15 days post inoculation. At this point, rice plants showing satisfactory reduction of the transcription levels were used in our standard conidial spray inoculation and leaf sheath assays with fungal strains expressing fluorescently-labeled effectors.

**Statistics**

All experiments were performed in triplicate and repeated three times. Calculations (unpaired t test, analysis of variance) were performed with the software XLSTAT version 2009.4.02 (Addinsoft).

**ACKNOWLEDGEMENTS**

We acknowledge and thank Rick Nelson (Noble Foundation, Ardmore, OK) for providing the VIGS vectors and hosting E.O.-G. in his laboratory to learn the VIGS assay. We thank the Talbot laboratory for assistance and hospitality during a working visit for E.O.-G at the University of Exeter. We thank Hiromasa Saitoh (Iwate Biotechnology Research Center, Kitakami, Iwate, Japan) and Ryohei Terauchi (Iwate Biotechnology Research Center, Kitakami, Iwate, and Kyoto University, Japan) for sharing transgenic rice expressing plant plasma membrane marker LTi6B:GFP. This project was supported by Agriculture and Food Research Initiative Competitive Grant no. #2017-67013-26525 from the USDA National Institute of Food and Agriculture Plant Biotic Interactions program. This is Contribution no. 22-053-J from the Kansas Agricultural Experiment Station.

**AUTHOR CONTRIBUTIONS**

E.O.-G. and B.V. designed experiments and analyzed data: E.O.-G. performed most of the research. T.M.T., J.P. and S.P. produced and analyzed transgenic rice lines expressing the rice CLC1:eGFP and Flot1:eGFP fluorescent fusion protein genes. M.D., M.M.-U., C.R.H. and A.H.V. assisted with microscopy and Bas83 Knockout experiments and general laboratory experiments; N.J.T. provided transgenic rice lines expressing LTi6B:GFP and LifeACT:GFP; E.O.-G., B.V. and N.J.T wrote and edited the article.

645    **Table 1. Chemical inhibitors of endocytosis**

| Chemical | Target | Mode of action | Comments | References |
|---|---|---|---|---|
| Filipin | CIE* | Binds to sterols in the membrane. | Toxic at higher concentrations. | (Rodal et al., 1999) |
| Methyl-β-cyclodextrin | CIE | Deplete cellular membranes of sterols by increasing the water solubility of the sterol. | caveolae-dependent endocytosis. | (Rodal et al., 1999) |
| Chlorpromazine | CME** | Translocates clathrin and AP2 from the cell surface to intracellular endosomes. | Also inhibits CIE in some cells. | (Wang et al., 1993) |
| Cantharidin | CME | Inhibits PP2A (protein phosphatase 2A) | Affects flg22-mediated FLS2 endocytosis. | (Serrano et al., 2007) |
| Triclosan | CME | Inhibits PP2A (protein phosphatase 2A) | Affects flg22-mediated FLS2 endocytosis. | (Serrano et al., 2007) |
| Wortmannin | nd*** | Inhibits PI3Ks, blocks late trafficking to the vacuole, multivesicular bodies and endocytosis. | Late endosomes lose their clathrin; Potential CME association. | (Bright et al., 2001; Sachse et al., 2002; Robinson et al., 2008) |
| Concanamycin A | nd | Targets V-type ATPase and blocks trafficking at trans-Golgi network, endosome acidification. | Induces acidification crucial for endocytic pathways. | (Maranda et al., 2001; Robinson et al., 2008) |

646    *CIE: clathrin-independent endocytosis; **CME: clathrin-mediated endocytosis; ***nd: not determined.

648    Figure 1

649
**Figure 1**. **BIC vesicular organization and effector vesicles in the rice cell cytoplasm**. **A.**
Cytoplasmic effector-labeled vesicles in a tip-BIC of strain KV217 expressing Pwl2:mRFP and
Bas4:eGFP at 17 hpi. The appressorium (Ap) and the first half of the primary hypha (PH) are
shown out of focus. Separate mRFP and eGFP channels (right) for the inset show the base layer
of apoplastic effector Bas4:eGFP and the outer vesicular layer labeled by cytoplasmic effector
Pwl2:mRFP. **B.** A side-BIC shown above the bulbous IH of strain KV168 expressing
Pwl2:mCherry:NLS and Bas4:eGFP at 24 hpi. Effector vesicles in are visible in the host cytoplasm
at a distance from the BIC. The Bas4:eGFP control shows that the EIHM surrounding the IH
remains intact. Addition of an artificial NLS to the fluorescent Pwl2 fusion protein facilitates
visualization of effector translocation and accumulation inside the host cell nucleus. **C.**

Quantification of the relative localization of cytoplasmic effector vesicles, assessed using strain KV217 (same as in panel A) at 98 infection sites for each BIC developmental stage; three biological replicates. Bars in C represent standard deviation. *P≤0.05. Images in A and B (merged bright field, eGFP and mRFP fluorescence) are shown as projections of confocal optical sections. Bars = 10 μm.

Figure 2

**Figure 2. Dynamics of BICs and effector vesicles during differentiation of primary hyphae**
**to first bulbous invasive hyphae. A.** Dynamics of two cytoplasmic effectors, Pwl2:mRFP and
Bas1:eYFP (Strain KV211), as a tip-BIC (top panel) becomes a side-BIC (bottom panel), imaged
every 30 min from 19 to 21 hpi. Images (left to right) are merged bright field, eYFP and mRFP
fluorescence at 5 time points; merged eYFP and mRFP at same time points in both the same and
enlarged views of the fluorescent BICs shown as projections of confocal optical sections. Bars =
10 µm. **B.** Cytoplasmic effectors Pwl2:mRFP and Bas1:eYFP (Strain KV211) co-localize in larger
vesicles in a side-BIC at 24 hpi. **C.** A rare extremely large effector vesicle (diameter 3.9 µm) in a
late stage side-BIC formed by strain KV170 expressing cytoplasmic effector Bas1:mRFP in a rice
cell at 30 hpi. **D,E.** Box and whisker plots comparing vesicle sizes (D) and  number of vesicles (E)
in tip- and side-BICs formed by strain KV211 invading rice cells. **F.** Bar chart showing
quantification of Bas1:eYFP and Pwl2:mRFP colocalization (KV211) in vesicles in tip- and side-
BICs. Bars in F are standard deviations. *$P \leq 0.05$; three times 100 BICs observed in D-F.

     Figure 3

**Figure 3. Effector vesicles are derived from plant plasma membrane. A.** Rice expressing plasma membrane marker LTi6b:eGFP in non-infected cells (left column) and in cells infected by strain KV209 expressing Pwl2:mRFP (right column). Colocalization confirms BICs as plant plasma membrane-rich structure. **B.** Quantification of the colocalization of LTi6b:eGFP with BICs at both developmental stages. Three times 90 infection sites observed. **C.** Localization assays of LTi6b:eGFP relative to Pwl2:mRFP in an infected rice cell at 22 hpi showing LTi6b:eGFP outlining vesicles in the BIC. Insets show BIC vesicles at higher magnification. Bars are 10 µm. **D.** Quantification of BICs stained by the endocytosis tracer dyes FM 4-64 and FM 1-43 at the tips of primary hyphae and beside the first bulbous IH. Three times 90 infection sites observed treatment. **E.** Colocalization of BIC effector vesicles and FM4-64-stained membranes in BICs.

Images of strain KV176 expressing Pwl2:eGFP in BICs beside the first bulbous IH cell in rice
stained with FM4-64 and observed at 19 hpi. The image is merged eGFP and FM 4-64 fluorescence
with eGFP fluorescence shown alone in the inset in the lower left. The upper right inset shows a
top-down view of a different BIC, shown below the image plane, at 22 hpi. Inset arrows indicate
Pwl2:eGFP vesicles near the BIC. Images are projections of confocal optical sections. Bar = 10
μm. See Supplemental Movie S2 for yet another side-BIC with additional examples of
colocalization of FM 4-64 fluorescence and Pwl2:eGFP.

Figure 4

**Figure 4. Bas83 is a novel effector associated membrane vesicles surrounding BICs and BIC-**
**associated cells. A.** Bas83:mRFP binds apparent membrane vesicles surrounding a side-BIC
(white arrow) labeled with Pwl2:eGFP (strain KV222). **B.** Bas83:mRFP (KV220) binds to bubble-
like structures near a primary hypha (arrow) as well as labeling a BIC and outlining BIC-associated
cells. Images (merged bright field, eGFP and mRFP fluorescence) are projections of confocal
optical sections. See additional images in **Supplemental Figure S3**. Bars = 10 μm. **C.**
Quantification of the cellular localization of Bas83:mRFP during biotrophic invasion of *M. oryzae*
(KV220). Bars are standard deviations. *$P \leq 0.05$; three times 100 infection sites observed per
experiment.

Figure 5

**Figure 5. BICs contain plant actin. A.** Rice LifeAct:eGFP localization during biotrophic invasion
by *M. oryzae*. A rice line expressing LifeAct:eGFP in non-infected cells (left column) and cells
infected by *M. oryzae* strain KV209 expressing Pwl2:mRFP in BICs (right column).
Colocalization shows BICs are rich in plant actin. **B.** Quantification of the colocalization of
LifeAct:eGFP with both tip- and side-BICs. Three times 100 infection sites observed. **C.**
Rhodamine phalloidin conjugate (red fluorescence staining actin filaments) colocalizes with
Pwl2:eGFP in rice cells infected by strain KV176, confirming the actin-rich structure of the BIC.

**Figure 6. BICs uniformly contain clathrin, but not in flotillin1. A.** Rice leaf sheaths expressing OsCLC1:eGFP (top panels) or OsFlot1:eGFP (bottom panels) were inoculated with strain KV209 expressing Pwl2:mRFP. Shown from left to right are a non-infected rice sheath cell; an infected cell containing a primary hypha (PH) with a tip-BIC at 19 hpi; and an infected cell containing a

729　bulbous IH (BH) with a BIC at 22 hpi. Arrowheads indicate endocytic foci and arrows indicate
730　BICs. Bars=10μm. **B.** Quantification of the colocalization of OsCLC1:eGFP or OsFlot1:eGFP
731　with BICs on the tip of a primary hypha and beside the first bulbous IH cell. Bars are standard
732　deviations. *$P \leq 0.05$; three times 97 BICs observed each treatment.
733

**Figure 7. Silencing of rice endocytic machinery suggests roles in pathogenicity and fungal effector translocation. A.** Disease severity in rice leaves, cultivar IR64, undergoing silencing of rice endocytic genes**.** Whereas silencing of *Flot1* gene showed minor effects on pathogenesis, the

silencing of the key clathrin-mediated endocytosis genes, *AP-2α* and CHC1 resulted in a significant reduction in pathogenicity in whole plant spray inoculation assays. **B.** Quantification of diseased area in rice expressing RNAi*AP-2α*, RNAi*CHC1*, RNAi*Flot1* or empty vector infected by *M. oryzae* strain Guy11 ($P \leq 0.05$ for all treatments; n = 9 rice plants for each treatment; mean + standard deviation, three experiments). **C.** Effects of silencing rice endocytosis genes on the localization of Pwl2:mRFP and Bas4:eGFP during biotrophic invasion of *M. oryzae* (KV217). Wild type IR64 rice and rice expressing RNAi*Flot1* or RNAi*AP-2α* were infected by KV217 expressing Pwl2:mRFP and Bas4:eGFP at 26 hpi. Note the accumulation of Pwl2:mRFP under the appressorium (Ap) and the pale Bas4:eGFP fluorescence associated with the short primary hypha (PH) in the RNAi*AP2α* rice compared to the control and RNAi*Flot1* rice. BICs formed in RNAi*AP-2α* and RNAi*CHC1* rice show similar swollen phenotype. Images in C (merged bright field, eGFP and mRFP fluorescence) are shown as projections of confocal optical sections. Bars = 10 µm. **D-F.** Quantification of the effector accumulation under appressoria (D), appressorial penetration (E), and swollen, irregular BICs (F) in rice expressing RNAi*Flot1* and RNAi*AP-2α*, RNAi*CHC1* and empty vector infected by *M. oryzae* strain KV209 expressing Pwl2:mRFP at 26 hpi. Bars are standard deviations, three experiments. (\*$P \leq 0.05$; three times 100 infection sites counted each treatment).

 Figure 8

758
**Figure 8. Treatment with CME inhibitor cantharidin impacts the localization of cytoplasmic**
**effector Pwl2, but not apoplastic effector Bas4. A.** Apoplastic effector Bas4:eGFP identifies an
inner base layer of the BIC relative to cytoplasmic Pwl2:mRFP, as shown by fluorescence intensity
linescans for GFP (green) and mRFP (red) along the path of the white arrow. This side-BIC was
formed during invasion of rice by strain KV217 expressing Pwl2:mRFP and Bas4:eGFP at 26 hpi.
**B.** After treatment with CME inhibitor cantharidin, localization of Bas4:eGFP in the BIC base
appears unaffected, compared to the non-vesicular swollen localization of Pwl2:mRFP, as shown
by fluorescence intensity linescans for GFP (green) and mRFP (red) along the path of the white
arrow. Note linescan distance is almost 3-fold greater after cantharidin treatment. This BIC was
formed by KV217 at 26 hpi, as in control panel A. Images (merged bright field, eGFP and mRFP
fluorescence) are shown as projections of confocal optical sections. Appressorium (Ap); Primary
hypha (PH). Bars = 10 μm.

773
**Figure 9. Chemical inhibition of rice CME, but not of CIE, has a major impact on cytoplasmic effector localization and appressorial penetration. A.** Effects of chemical inhibition of Clathrin-mediated endocytosis (CME) and Clathrin-independent endocytosis (CIE) in the rice leaf sheath assay with strain KV209 expressing Pwl2:mRFP at 18, 21 and 28 hpi. Whereas inhibition of CIE showed minor effects on BIC functionality and effector uptake, the chemical inhibition of CME resulted in a significant accumulation of effector fluorescence under appressoria (Ap, white arrows at 18 hpi) as well as swollen BICs on primary hyphae (PH, 21 hpi) and bulbous IH (28hpi). Images in A (merged bright field and mRFP fluorescence) are projections of confocal optical sections. Insets show close-up views of the effector fluorescence. Bars = 10

μm and bars in the insert boxes = 5 μm. **B-E.** Quantification of swollen, irregular-shaped BICs
(B), regular-shaped BICs (C), appressorial penetration (D), and effector accumulation under
appressoria (E), in rice undergoing chemical treatment and infection by strain KV209 expressing
Pwl2:mRFP at 28 hpi. Bars are standard deviations. Three times 100 infection sites observed per
treatment.

Figure 10

are secreted into BICs by a nonconventional, Brefeldin A-insensitive secretion system involving
the exocyst and SNARE protein Sso1 (red arrows). Cytoplasmic effectors also accumulate inside
the EIHM compartment around the BIC-associated cells, the PH and first differentiated bulbous
IH cell (not shown), but not in the EIHM surrounding nonBIC-associated bulbous IH cells, which
are tightly surrounded by the host vacuole. Our current study expands this model by showing that
cytoplasmic effectors are packaged in dynamic vesicles inside BICs and in the host cytoplasm
surrounding BICs, and that these effector vesicles co-localize with fluorescent plant plasma
membrane markers LTi6B:GFP and FM 4-64, as well as clathrin-mediated endocytosis (CME)
marker CLC1:eGFP. These co-localization studies, additional localization with plant actin in BICs,
and functional analyses via VIGS and pharmacological inhibitors support CME as a major
mechanism for effector translocation. We also report two novel cytoplasmic effectors, Bas83,
which labels empty membrane vesicles possibly recruiting more host membrane to the BIC and
the EIHM surrounding BIC-associated cells, and Bas170, which is delivered in vesicles to rice
cytoplasm from under the appressorium and accumulates in the host nucleus even before growth
of the primary hyphae. Cytoplasmic connections between cytoplasm surrounding BICs and
cytoplasm on the host cell periphery would facilitate movement of translocated effectors to the
host nucleus and other sites of action in the host cell, as well as movement of cytoplasmic effectors
through plasmodesmata into surrounding host cells to prepare them for invasion. Key: N, fungal
nucleus; ER, endoplasmic reticulum; G, Golgi apparatus; V, transport vesicle. **B.** Working models
for vesicle formation and effector translocation in the rice-*M. oryzae* interaction through CME or
clathrin-independent endocytosis (CIE). Although our current work supports a role for CME, some
involvement of CIE in effector translocation cannot be totally ruled out. AP2, Adaptor Protein-2
complex; EIHM, extrainvasive hyphal membrane; EIHMx, extrainvasive hyphal matrix; FPM,
fungal plasma membrane; FCW, fungal cell wall.

**REFERENCES**

Atkinson, H.A., Daniels, A., and Read, N.D. 2002. Live-cell imaging of endocytosis during

conidial germination in the rice blast fungus, *Magnaporthe grisea*. Fungal Gen. Biol.

37:233-244.

Beck, M., Heard, W., Mbengue, M., and Robatzek, S. 2012. The INs and OUTs of pattern

recognition receptors at the cell surface. Curr. Opin. Plant Biol. 15:367-374.

Bolte, S., Talbot, C., Boutte, Y., Catrice, O., Read, N.D., and Satiat-Jeunemaitre, B. 2004. FM-

dyes as experimental probes for dissecting vesicle trafficking in living plant cells. Journal

of Microscopy 214:159-173.

Bozkurt, T.O., Belhaj, K., Dagdas, Y.F., Chaparro-Garcia, A., Wu, C.-H., Cano, L.M., and

Kamoun, S. 2015. Rerouting of plant late endocytic trafficking toward a pathogen

interface. Traffic 16:204-226.

Bright, N.A., Lindsay, M.R., Stewart, A., and Luzio, J.P. 2001. The relationship between lumenal

and limiting membranes in swollen late endocytic compartments formed after wortmannin

treatment or sucrose accumulation. Traffic 2:631-642.

Carroll, A., Sweigard, J. A., and Valent, B. 1994. Improved vectors for selecting resistance to

hygromycin.  41:Article 5. https://newprairiepress.org/fgr/vol41/iss41/45/.

Catlett, N. L., Lee, B.-N., Yoder, O. C., and Turgeon, B. G. 2003. Split-marker recombination for

efficient targeted deletion of fungal genes. Fungal Genetics Newsletter 50:Article 4.

https://doi.org/10.4148/1941-4765.1150.

Chaparro-Garcia, A., Schwizer, S., Sklenar, J., Yoshida, K., Petre, B., Bos, J.I.B., Schornack, S.,

Jones, A.M.E., Bozkurt, T.O., and Kamoun, S. 2015. *Phytophthora infestans* RXLR-WY

effector AVR3a associates with dynamin-related protein 2 required for endocytosis of the

plant pattern recognition receptor FLS2. PLoS ONE 10:e0137071.

Chen, X., Irani, N.G., and Friml, J. 2011. Clathrin-mediated endocytosis: the gateway into plant

cells. Curr. Opin. Plant Biol. 14:674-682.

Collemare, J., Pianfetti, M., Houlle, A.E., Morin, D., Camborde, L., Gagey, M.J., Barbisan, C.,

Fudal, I., Lebrun, M.H., and Böhnert, H.U. 2008. *Magnaporthe grisea* avirulence gene

*ACE1* belongs to an infection-specific gene cluster involved in secondary metabolism. New

Phytol. 179:196-208.

Dagdas, Y.F., Pandey, P., Tumtas, Y., Sanguankiattichai, N., Belhaj, K., Duggan, C., Leary, A.Y.,
Segretin, M.E., Contreras, M.P., Savage, Z., Khandare, V.S., Kamoun, S., and Bozkurt,
T.O. 2018. Host autophagy machinery is diverted to the pathogen interface to mediate focal
defense responses against the Irish potato famine pathogen. eLife 7:e37476.

Dejonghe, W., and Russinova, E. 2017. Plant chemical genetics: From phenotype-based screens
to synthetic biology. Plant Physiology 174:5-20.

Dejonghe, W., Kuenen, S., Mylle, E., Vasileva, M., Keech, O., Viotti, C., Swerts, J., Fendrych,
M., Ortiz-Morea, F.A., Mishev, K., Delang, S., Scholl, S., Zarza, X., Heilmann, M.,
Kourelis, J., Kasprowicz, J., Nguyen, L.S.L., Drozdzecki, A., Van Houtte, I., Szatmári, A.-
M., Majda, M., Baisa, G., Bednarek, S.Y., Robert, S., Audenaert, D., Testerink, C.,
Munnik, T., Van Damme, D., Heilmann, I., Schumacher, K., Winne, J., Friml, J.,
Verstreken, P., and Russinova, E. 2016. Mitochondrial uncouplers inhibit clathrin-
mediated endocytosis largely through cytoplasmic acidification. Nat. Comm. 7:11710.

Di Rubbo, S., Irani, N. G., Kim, S. Y., Xu, Z.-Y., Gadeyne, A., Dejonghe, W., Vanhoutte, I.,
Persiau, G., Eeckhout, D., Simon, S., Song, K., Kleine-Vehn, J., Friml, J., De Jaeger, G.,
Van Damme, D., Hwang, I., and Russinova, E. 2013. The Clathrin Adaptor Complex AP-
2 mediates endocytosis of BRASSINOSTEROID INSENSITIVE1 in *Arabidopsis*. Plant
Cell 25:2986-2997.

Ding, X.S., Rao, C.S., and Nelson, R.S. 2007. Analysis of gene function in rice through virus-
induced gene silencing. Methods in molecular biology (Clifton, N.J.) 354:145-160.

Ding, X.S., Schneider, W.L., Chaluvadi, S.R., Mian, M.A.R., and Nelson, R.S. 2006.
Characterization of a Brome mosaic virus strain and its use as a vector for gene silencing
in monocotyledonous hosts. Mol. Plant-Microbe Interact. 19:1229-1239.

Dodds, P.N., and Rathjen, J.P. 2010. Plant immunity: towards an integrated view of plant-pathogen
interactions. Nat. Rev. Genet. 11:539-548.

Elsworth, B., Matthews, K., Nie, C.Q., Kalanon, M., Charnaud, S.C., Sanders, P.R., Chisholm,
S.A., Counihan, N.A., Shaw, P.J., Pino, P., Chan, J.A., Azevedo, M.F., Rogerson, S.J.,
Beeson, J.G., Crabb, B.S., Gilson, P.R., and de Koning-Ward, T.F. 2014. PTEX is an
essential nexus for protein export in malaria parasites. Nature 511:587-+.

Ewers, H., and Helenius, A. 2011. Lipid-mediated endocytosis. Cold Spring Harbor Perspectives
in Biology 3:10.1101/cshperspect.a004721.

Fan, L., Li, R., Pan, J., Ding, Z., and Lin, J. 2015. Endocytosis and its regulation in plants. Trends
in Plant Science 20:388-397.

Fawke, S., Doumane, M., and Schornack, S. 2015. Oomycete interactions with plants: infection
strategies and resistance principles. Microbiology and Molecular Biology Reviews 79:263-
280.

Fudal, I., Collemare, J., Böhnert, H.U., Melayah, D., and Lebrun, M.-H. 2007. Expression of
*Magnaporthe grisea* avirulence gene *ACE1* is connected to the initiation of appressorium-
mediated penetration. Eukaryotic Cell 6:546-554.

Giraldo, M.C., and Valent, B. 2013. Filamentous plant pathogen effectors in action. Nat. Rev.
Microbiol. 11:800-814.

Giraldo, M.C., Dagdas, Y.F., Gupta, Y.K., Mentlak, T.A., Yi, M., Martinez-Rocha, A.L., Saitoh,
H., Terauchi, R., Talbot, N.J., and Valent, B. 2013. Two distinct secretion systems facilitate
tissue invasion by the rice blast fungus *Magnaporthe oryzae*. Nat. Comm. 4:1996.

Gladieux, P., Condon, B., Ravel, S., Soanes, D., Maciel, J.L.N., Nhani, A., Chen, L., Terauchi, R.,
Lebrun, M.-H., Tharreau, D., Mitchell, T., Pedley, K.F., Valent, B., Talbot, N.J., Farman,
M., and Fournier, E. 2018. Gene flow between divergent cereal- and grass-specific lineages
of the rice blast fungus *Magnaporthe oryzae*. mBio 9:e01219-01217.

Grassart, A., Cheng, A.T., Hong, S.H., Zhang, F., Zenzer, N., Feng, Y., Briner, D.M., Davis, G.D.,
Malkov, D., and Drubin, D.G. 2014. Actin and dynamin2 dynamics and interplay during
clathrin-mediated endocytosis. Journal of Cell Biology 205:721-735.

Harder, D.E., and Mendgen, K. 1982. Filipin-sterol complexes in bean rust- and oat crown rust-
fungal/plant interactions: Freeze-etch electron microscopy. Protoplasma 112:46-54.

Hellens, R., Mullineaux, P., and Klee, H. 2000. Technical Focus: a guide to *Agrobacterium* binary
Ti vectors. Trends Plant Sci. 5:446-451.

Hicks, G.R., and Raikhel, N.V. 2010. Advances in dissecting endomembrane trafficking with
small molecules. Curr. Opin. Plant Biol. 13:706-713.

Himmelbach, A., Zierold, U., Hensel, G., Riechen, J., Douchkov, D., Schweizer, P., and Kumlehn,
J. 2007. A set of modular binary vectors for transformation of cereals. Plant Physiology
145:1192-1200.

Holsters, M., de Waele, D., Depicker, A., Messens, E., van Montagu, M., and Schell, J. 1978.
Transfection and transformation of *Agrobacterium tumefaciens*. Molecular and General
Genetics 163:181-187.

Howard, R.J., and Valent, B. 1996. Breaking and Entering: Host penetration by the fungal rice
blast pathogen *Magnaporthe grisea*. Annu. Rev. Microbiol. 50:491-512.

Johnson, A., Gnyliukh, N., Kaufmann, W.A., Narasimhan, M., Vert, G., Bednarek, S.Y., and
Friml, J. 2020. Experimental toolbox for quantitative evaluation of clathrin-mediated
endocytosis in the plant model *Arabidopsis*. Journal of Cell Science 133:jcs248062.

Jones, K., Zhu, J., Jenkinson, C.B., Kim, D.W., Pfeifer, M.A., and Khang, C.H. 2021. Disruption
of the interfacial membrane leads to *Magnaporthe oryzae* effector re-location and lifestyle
switch during rice blast disease. Frontiers in Cell and Developmental Biology
9:10.3389/fcell.2021.681734.

Kaksonen, M., and Roux, A. 2018. Mechanisms of clathrin-mediated endocytosis. Nat. Rev. Mol.
Cell Biol. 19:313-326.

Kale, S.D., Gu, B., Capelluto, D.G.S., Dou, D., Feldman, E., Rumore, A., Arredondo, F.D.,
Hanlon, R., Fudal, I., Rouxel, T., Lawrence, C.B., Shan, W., and Tyler, B.M. 2010.
External lipid PI3P mediates entry of eukaryotic pathogen effectors into plant and animal
host cells. Cell 142:284-295.

Kankanala, P., Czymmek, K., and Valent, B. 2007. Roles for rice membrane dynamics and
plasmodesmata during biotrophic invasion by the blast fungus. Plant Cell 19:706-724.

Kato, H. 1974. Epidemiology of rice blast disease. Review of Plant Protection Research 7:1-20.

Kershaw, M. J., and Talbot, N. J. 2009. Genome-wide functional analysis reveals that infection-
associated fungal autophagy is necessary for rice blast disease. Proceedings of the National
Academy of Sciences U.S.A. 106:15967-15972.

Khang, C.H., Park, S., Rho, H., Lee, Y., and Kang, S. 2006. *Agrobacterium tumefaciens*-mediated
transformation and mutagenesis of filamentous fungi *Magnaporthe grisea* and *Fusarium*
*oxysporum*. in: Agrobacterium Protocols, K. Wang, ed. Humana Press, Totowa.

Khang, C.H., Berruyer, R., Giraldo, M.C., Kankanala, P., Park, S.-Y., Czymmek, K., Kang, S.,
and Valent, B. 2010. Translocation of *Magnaporthe oryzae* effectors into rice cells and
their subsequent cell-to-cell movement. Plant Cell 22:1388-1403.

Kleemann, J., Rincon-Rivera, L.J., Takahara, H., Neumann, U., Ver Loren van Themaat, E., van
der Does, H.C., Hacquard, S., Stüber, K., Will, I., Schmalenbach, W., Schmelzer, E., and
O'Connell, R.J. 2012. Sequential delivery of host-induced virulence effectors by
appressoria and intracellular hyphae of the phytopathogen *Colletotrichum higginsianum*.
PLoS Path. 8:e1002643.

Konopka, C.A., Backues, S.K., and Bednarek, S.Y. 2008. Dynamics of Arabidopsis Dynamin-
Related Protein 1C and a Clathrin Light Chain at the Plasma Membrane  Plant Cell
20:1363-1380.

Larson, E. R., Van Zelm, E., Roux, C., Marion-Poll, A., and Blatt, M. R. 2017. Clathrin heavy
chain subunits coordinate endo- and exocytic traffic and affect stomatal movement. Plant
Physiol. 175:708-720.

Leung, H., Borromeo, E.S., Bernardo, M.A., and Notteghem, J.L. 1988. Genetic analysis of
virulence in the rice blast fungus *Magnaporthe grisea*. Phytopathology 78:1227-1233.

Li, R., Liu, P., Wan, Y., Chen, T., Wang, Q., Mettbach, U., Baluška, F., Samaj, J., Fang, X., Lucas,
W.J., and Lin, J. 2012. A membrane microdomain-associated protein, *Arabidopsis* Flot1,
is involved in a clathrin-independent endocytic pathway and is required for seedling
development. Plant Cell 24  2105-2122

Lo Presti, L., and Kahmann, R. 2017. How filamentous plant pathogen effectors are translocated
to host cells. Curr. Opin. Plant Biol. 38:19-24.

Lo Presti, L., Zechmann, B., Kumlehn, J., Liang, L., Lanver, D., Tanaka, S., Bock, R., and
Kahmann, R. 2017. An assay for entry of secreted fungal effectors into plant cells. New
Phytol. 213:956-964.

Lo Presti, L., Lanver, D., Schweizer, G., Tanaka, S., Liang, L., Tollot, M., Zuccaro, A., Reissmann,
S., and Kahmann, R. 2015. Fungal effectors and plant susceptibility. Annual Review of
Plant Biology 66:513-545.

Ludwig, N., Reissmann, S., Schipper, K., Gonzalez, C., Assmann, D., Glatter, T., Moretti, M., Ma,
L.-S., Rexer, K.-H., Snetselaar, K., and Kahmann, R. 2021. A cell surface-exposed protein
complex with an essential virulence function in *Ustilago maydis*. Nature Microbiology
6:722-730.

Manning, V.A., and Ciuffetti, L.M. 2005. Localization of Ptr ToxA produced by *Pyrenophora*
*tritici-repentis* reveals protein import into wheat mesophyll cells. Plant Cell 17:3203-3212.

Maranda, B., Brown, D., Bourgoin, S., Casanova, J.E., Vinay, P., Ausiello, D.A., and Marshansky,
975            V. 2001. Intra-endosomal pH-sensitive recruitment of the Arf-nucleotide exchange factor
976            ARNO and Arf6 from cytoplasm to proximal tubule endosomes. Journal of Biological
977            Chemistry 276:18540-18550.

978 Mbengue, M., Bourdais, G., Gervasi, F., Beck, M., Zhou, J., Spallek, T., Bartels, S., Boller, T.,
979            Ueda, T., Kuhn, H., and Robatzek, S. 2016. Clathrin-dependent endocytosis is required for
980            immunity mediated by pattern recognition receptor kinases. Proceedings of the National
981            Academy of Sciences 113:11034-11039.

982 Mentlak, T.A., Kombrink, A., Shinya, T., Ryder, L.S., Otomo, I., Saitoh, H., Terauchi, R.,
983            Nishizawa, Y., Shibuya, N., Thomma, B.P., and Talbot, N.J. 2012. Effector-mediated
984            suppression of chitin-triggered immunity by *Magnaporthe oryzae* is necessary for rice blast
985            disease. Plant Cell 24:322-335.

986 Mochizuki, S., Minami, E., and Nishizawa, Y. 2015. Live-cell imaging of rice cytological changes
987            reveals the importance of host vacuole maintenance for biotrophic invasion by blast
988            fungus, *Magnaporthe oryzae*. MicrobiologyOpen 4:952-966.

989 Mosquera, G., Giraldo, M.C., Khang, C.H., Coughlan, S., and Valent, B. 2009. Interaction
990            transcriptome analysis identifies *Magnaporthe oryzae* BAS1-4 as biotrophy-associated
991            secreted proteins in rice blast disease. Plant Cell 21:1273-1290.

992 Narasimhan, M., Johnson, A., Prizak, R., Kaufmann, W.A., Tan, S., Casillas-Pérez, B., and Friml,
993            J. 2020. Evolutionarily unique mechanistic framework of clathrin-mediated endocytosis in
994            plants. eLife 9:e52067.

995 Nishimura, T., Mochizuki, S., Ishii-Minami, N., Fujisawa, Y., Kawahara, Y., Yoshida, Y., Okada,
996            K., Ando, S., Matsumura, H., Terauchi, R., Minami, E., and Nishizawa, Y. 2016.
997            *Magnaporthe oryzae* glycine-rich secretion protein, Rbf1 critically participates in
998            pathogenicity through the focal formation of the biotrophic interfacial complex. PLoS Path.
999            12:e1005921.

1000 Nishizawa, Y., Mochizuki, S., Yokotani, N., Nishimura, T., and Minami, E. 2016. Molecular and
1001            cellular analysis of the biotrophic interaction between rice and *Magnaporthe oryzae* –
1002            Exploring situations in which the blast fungus controls the infection. Physiol. Mol. Plant
1003            Pathol. 95:70-76.

O'Connell, R.J., and Panstruga, R. 2006. Tête à tête inside a plant cell: establishing compatibility
between plants and biotrophic fungi and oomycetes. New Phytol. 171:699-718.

Otto, G.P., and Nichols, B.J. 2011. The roles of flotillin microdomains – endocytosis and beyond.
Journal of Cell Science 124:3933-3940.

Park, S.H., Park, J., and Smith, R.H. 2001. Herbicide and insect resistant elite transgenic rice.
Journal of Plant Physiology 158:1221-1226.

Petre, B., and Kamoun, S. 2014. How do filamentous pathogens deliver effector proteins into plant
cells? PLoS. Biol. 12:e1001801.

Petre, B., Contreras, M.P., Bozkurt, T.O., Schattat, M.H., Sklenar, J., Schornack, S., Abd-El-
Haliem, A., Castells-Graells, R., Lozano-Durán, R., Dagdas, Y.F., Menke, F.L.H., Jones,
A.M.E., Vossen, J.H., Robatzek, S., Kamoun, S., and Win, J. 2021. Host-interactor screens
of *Phytophthora infestans* RXLR proteins reveal vesicle trafficking as a major effector-
targeted process. Plant Cell 33:1447-1471.

Postma, J., Liebrand, T.W.H., Bi, G., Evrard, A., Bye, R.R., Mbengue, M., Kuhn, H., Joosten,
M.H.A.J., and Robatzek, S. 2016. Avr4 promotes Cf-4 receptor-like protein association
with the BAK1/SERK3 receptor-like kinase to initiate receptor endocytosis and plant
immunity. New Phytol. 210:627-642.

Qi, X., Pleskot, R., Irani, N.G., and Van Damme, D. 2018. Meeting report – Cellular gateways:
expanding the role of endocytosis in plant development. Journal of Cell Science
131:jcs222604.

Reynolds, G.D., Wang, C., Pan, J., and Bednarek, S.Y. 2018. Inroads into internalization: Five
years of endocytic exploration. Plant Physiology 176:208-218.

Robinson, D.G., Jiang, L.W., and Schumacher, K. 2008. The endosomal system of plants: Charting
new and familiar territories. Plant Physiology 147:1482-1492.

Rodal, S.K., Skretting, G., Garred, O., Vilhardt, F., van Deurs, B., and Sandvig, K. 1999.
Extraction of cholesterol with methyl-beta-cyclodextrin perturbs formation of clathrin-
coated endocytic vesicles. Molecular Biology of the Cell 10:961-974.

Sachse, M., Urbe, S., Oorschot, V., Strous, G.J., and Klumperman, J. 2002. Bilayered clathrin
coats on endosomal vacuoles are involved in protein sorting toward lysosomes Molecular
Biology of the Cell 13:1313-1328

Sakulkoo, W., Oses-Ruiz, M., Garcia, E.O., Soanes, D.M., Littlejohn, G.R., Hacker, C., Correia,
A., Valent, B., and Talbot, N.J. 2018. A single fungal MAP kinase controls plant cell-to-
cell invasion by the rice blast fungus. Science 359:1399-1403.

Serrano, M., Robatzek, S., Torres, M., Kombrink, E., Somssich, I.E., Robinson, M., and Schulze-
Lefert, P. 2007. Chemical interference of pathogen-associated molecular pattern-triggered
immune responses in Arabidopsis reveals a potential role for fatty-acid synthase type II
complexderived lipid signals. Journal of Biological Chemistry 282:6803-6811.

Spallek, T., Beck, M., Ben Khaled, S., Salomon, S., Bourdais, G., Schellmann, S., and Robatzek,
S. 2013. ESCRT-I mediates FLS2 endosomal sorting and plant immunity. PLoS Genetics
9:e1004035.

Stark-Urnau, M., and Mendgen, K. 1995. Sequential deposition of plant glycoproteins and
polysaccharides at the host-parasite interface of *Uromyces vignae* and *Vigna sinensis*:
Evidence for endocytosis and secretion. Protoplasma 186:1-11.

Sweigard, J. A., Carroll, A. M., Kang, S., Farrall, L., Chumley, F. G., and Valent, B. 1995.
Identification, cloning, and characterization of *PWL2*, a gene for host species specificity in
the rice blast fungus. Plant Cell 7:1221-1233.

Trusch, F., Loebach, L., Wawra, S., Durward, E., Wuensch, A., Iberahim, N.A., de Bruijn, I.,
MacKenzie, K., Willems, A., Toloczko, A., Diéguez-Uribeondo, J., Rasmussen, T.,
Schrader, T., Bayer, P., Secombes, C.J., and van West, P. 2018. Cell entry of a host-
targeting protein of oomycetes requires gp96. Nat. Comm. 9:2347.

Valent, B., and Chumley, F.G. 1991. Molecular genetic analysis of the rice blast fungus,
*Magnaporthe grisea*. Annu. Rev. Phytopathol. 29:443-467.

Valent, B., Singh, P.K., He, X., Farman, M., Tosa, Y., and Braun, H.J. 2020. Blast diseases:
Evolution and challenges of a staple food crop fungal pathogen. Pages 267-292 in:
Emerging Plant Diseases and Global Food Security, J.B. Ristaino and A. Records, eds.
American Phytopathological Society Press, St. Paul, MN.

Van den Ackerveken, G. 2017. Seeing is believing: imaging the delivery of pathogen effectors
during plant infection. New Phytol. 216:8-10.

Wang, L.H., Rothberg, K.G., and Anderson, R.G. 1993. Misassembly of clathrin lattices on
endosomes reveals a regulatory switch for coated pit formation Journal of Cell Biology
123:1107-1117.

Wang, S., Boevink, P.C., Welsh, L., Zhang, R., Whisson, S.C., and Birch, P.R.J. 2017. Delivery

1066            of cytoplasmic and apoplastic effectors from *Phytophthora infestans* haustoria by distinct

1067            secretion pathways. New Phytol. 216:205-215.

1068    Wawra, S., Trusch, F., Matena, A., Apostolakis, K., Linne, U., Zhukov, I., Stanek, J., Koźmiński,

1069            W., Davidson, I., Secombes, C.J., Bayer, P., and van West, P. 2017. The RxLR motif of

1070            the host targeting effector AVR3a of *Phytophthora infestans* is cleaved before secretion.

1071            Plant Cell 29:1184-1195.

1072    Xu, J.-R., Staiger, C.J., and Hamer, J.E. 1998. Inactivation of the mitogen-activated protein kinase

1073            Mps1 from the rice blast fungus prevents penetration of host cells but allows activation of

1074            plant defense responses. Proceedings of the National Academy of Sciences U.S.A.

1075            95:12713-12718.