# Chromatin state transition underlies the temporal changes in gene expression during cardiomyocyte maturation

Chia-Yeh Lin[1], Yao-Ming Chang[2], Hsin-Yi Tseng[1], Yen-Ling Shih[2], Hsiao-Hui Yeh[2], You-Rou Liao[1], Chia-Ling Hsu[1], Chien-Chang Chen[2], Yu-Ting Yan[2,*] and Cheng-Fu Kao[1,*]

1. Institute of Cellular and Organismic Biology, Academia Sinica, 128, Academia Road, Section 2, Nankang, Taipei, Taiwan.
2. Institute of Biomedical Sciences, Academia Sinica, 128, Academia Road, Section 2, Nankang, Taipei, Taiwan.

* Corresponding author: ckao@gate.sinica.edu.tw and yyan@ibms.sinica.edu.tw

Abstract

Congenital heart disease (CHD) is often rooted in gene expression anomalies that

occur during heart development. As cells commit to a specific lineage, chromatin

dynamics and developmental plasticity generally become more limited. However, it

remains unclear how differentiated cardiomyocytes (CMs) undergo morphological

and functional adaptations to the postnatal environment during their maturation. In

this work, we sought to identify the regulatory mechanisms controlling postnatal

cardiac gene networks. A time-series transcriptomic analysis of postnatal hearts

revealed an integrated, time-ordered transcriptional network that regulates CM

maturation. Remarkably, depletion of histone H2B ubiquitin ligase RNF20 after

formation of the four-chamber heart disrupted these highly coordinated gene networks.

Its ablation also caused early-onset cardiomyopathy, a phenotype reminiscent of CHD.

Furthermore, we found that dynamic RNF20-mediated modulation of chromatin

accessibility during CM maturation was necessary for the operative binding of cardiac

transcription factors known to drive transcriptional gene networks. Together, our

results reveal how epigenetic-mediated chromatin state transitions modulate

time-ordered gene expression during CM maturation.

Key words: RNF20, chromatin accessibility, postnatal gene networks, CM maturation

Introduction

Aberrant cellular specification and maturation during cardiogenesis may lead to congenital heart disease (CHD)[1], the most common birth defect worldwide[2]. It may also lead to heart failure, a leading cause of morbidity and mortality in adult life[3]. Since precise control of gene expression is critical for heart development and maturation[4,5], knowledge of the regulatory mechanisms controlling cardiac gene networks is key to understanding the pathogenesis of cardiovascular disease.

Gene expression is tightly controlled by transcription factors (TFs) that act in accordance with epigenetic regulatory mechanisms[6,7]. TFs bind to specific sequences of DNA adjacent to a gene promoter and control its activity[6]. However, most TFs cannot access closed chromatin, and their binding to DNA relies on the presence of a permissive chromatin environment constructed by epigenetic modulators[7,8]. Global changes in chromatin accessibility have been observed during development of several different species[9,10], and stage-specific reorganization of the chromatin landscape appears to be critical for gene expression and cell-fate control[11].

During heart development, early epigenetic marks are established at the embryonic stage[12,13], and dynamic changes in these marks are associated with cardiac lineage specification and stage transition[13-15]. An increasingly large body of clinical and experimental research indicates that aberrant epigenetic regulation can contribute to CHD[12,16]. For instance, alterations in histone modifying enzymes were shown to commonly occur as *de novo* mutations in CHD patients[16]. More recently, a critical role for histone H2B ubiquitylation (H2Bub) was demonstrated in cardiac development[17,18]. Knockdown of the *Xenopus* H2Bub E3 ligases, *rnf20* (Ring finger protein 20) and *rnf40* (Ring finger protein 40)[19], beginning at the two-cell stage, influences cardiac looping and left-right (LR) asymmetry during early embryonic development[17]. Furthermore, the results of an *in vivo* CRISPR/Cas9 genetic screen suggested that RNF20/40 is involved in cardiomyocyte (CM) maturation via H2Bub deposition[18]. Thus, RNF20 appears to be relevant to both CHD and CM maturation[17,20]. The addition of a single ubiquitin on H2B disrupts local and higher-order structures of compacted chromatin[21,22]. Through this action, the RNF20-H2Bub pathway regulates gene activities by affecting the binding of TFs or regulatory proteins at gene promoters during transcription[23,24]. H2Bub is also selectively enriched in the coding regions of highly expressed genes[25], where its tight coupling with RNA polymerase II (Pol II)[23,26] and histone chaperone FACT[27]

facilitate transcriptional elongation. Nevertheless, the molecular function of RNF20

during temporal development of mammalian heart remains uncharacterized.

Here, we identified an integrated, time-ordered transcriptional network that

regulates CM maturation, and we found that deletion of *Rnf20* in CMs disrupts this

postnatal gene network and leads to dilated cardiomyopathy. We showed that RNF20

promotes chromatin relaxation in postnatal CMs, playing a key role in the remodeling

of chromatin accessibility at this stage. Taken together, our data show how chromatin

state transitions can coordinate the temporal expression of distinct transcriptional

programs during CM maturation.

**Results**

**A time-ordered transcriptional cascade during CM maturation**

In postnatal hearts, numerous processes (e.g., cell cycle, structural, functional

and metabolic transitions) are required to drive CM maturation[4,5]. These transitional

processes are known to be coordinated by certain gene networks[5,29], but it remains

unclear whether the transcriptional programs governing each process are regulated

independently and simultaneously or by an integrated time-ordered gene network[4]. To

address this open question, we conducted time-series experiments to obtain

transcriptomic data from developing hearts (ages P2 to P28) of control mice (Figure

1A). Principal component analysis (PCA) analysis of postnatal heart transcriptomes

showed clear distinctive expression profiles among the hearts at different stages (blue

circles in Figure 1B), suggesting drastic transcriptional alterations occur during

postnatal CM maturation.

The time-ordered gene coexpression network (TO-GCN) algorithm[30] was applied

to elucidate the dynamics of developmental processes (Figure 1C and S1). Each TF

that was expressed within the time-series transcriptomes was assigned to one of nine

levels (L1 to L9), representing its expression time order over the five time-points

(high expression levels are indicated by red squares in Figure 1C and Supplementary

file 1). In the network analysis, TFs assigned to the same level are upregulated at the

same time period. Moreover, TFs assigned to consecutive levels are upregulated in

similar time orders, while TFs at lower levels are upregulated earlier than TFs at

higher levels (Figure 1C).

Next, we identified overrepresented functional pathways in heart development at

each level of TO-GCN by examining coexpressed genes (Figure 1D and Figure S1-2).

The result shows a clear time-ordered organization of transcriptional cascades that
regulate CM maturation (Left panel, Figure 1D). For example, hypoxia-inducible
factor 1 (HIF1) signaling was enriched at level 1 (L1) and quickly decreased after L2
(Left panel, Figure 1D and S2-3), reflecting the transition of hypoxia to normoxia
after birth[31]. Following the decrease in HIF1 signaling, most of the enriched pathways
from L1 to L5 (corresponding to the first week) were related to cell-cell connections,
including genes supporting adherens junctions, actin cytoskeleton, tight junction focal
adhesion, and ECM-receptor interactions (Left panel, Figure 1D and Figure S2).
Genes related to cell cycle and DNA replication pathways were activated from L1/L2
(P2) to L4 (P7), reflecting a state of hyperplasia during the first week after birth[32]
(Left panel, Figure 1D). Thyroid (T3) hormone signaling (Left panel, Figure 1D),
which is required for CM growth and structural maturation[33], was enriched from L2 to
L4. In addition to the upregulation of T3 signaling, we also observed sequential
activation of several other signaling pathways. ErbB signaling was first activated at
L2, followed by neurotrophin and PI3K-AKT signaling at L3, insulin signaling at L4
and MAPK signaling at L5 (Left panel, Figure 1D). The combination of ErbB and
insulin signaling is thought to promote sarcomere synthesis[34,35]. Notably, cross-talk
between T3 and PI3K-AKT is important for myosin heavy chain and titin isoform
transitions, and it is further critical for myocardial distensibility and
mechanosignaling[36]. Moreover, the induction of neurotrophin signaling is required for
neurodevelopment and enhances normal CM $Ca^{2+}$ cycling and contraction[37]. These
enriched signaling pathways reflect CMs undergoing structural and functional
maturation (indicated by hypertrophy, isoform switch and contraction), accompanied
by halting of the cell-cycle. The activated insulin signaling at L4 is important for the
following fatty acid metabolism and oxidative phosphorylation at L7, reflecting a
metabolism transformation in the CMs[38]. At the same time, genes involved in
mitochondrial biogenesis and organization were highly expressed (Figure S2),
consistent with idea that maturing CMs undergo a transition of primary energy source
from glucose to fatty acids[39]. Lastly, the pathway of cardiac muscle contraction was
enhanced at L8 (P21) (Left panel, Figure 1D and Figure S2), in line with enhanced
heart function at this stage. In summary, the analysis revealed an integrated, that
accurately reflects the known coordination of signaling cascades regulating CM
maturation in the first three weeks of life.
151

**Coordinated postnatal gene networks are regulated by epigenetic factor, RNF20**

To investigate how an epigenetic regulator might influence the time-ordered transcriptional network (Control, Figure 1D), *Rnf20* was conditionally ablated in CMs after four chamber formation, a transition point between cardiac morphogenesis and maturation[5,28]. This ablation was accomplished by intercrossing *Rnf20^flox/flox* with *Mck-Cre* mice (*Rnf20^flox/flox; Mck*-Cre; hereafter called *cKO*)[40] (Figure 1A and S4A). We then collected and analyzed the transcriptomic data of developing hearts from *cKO* mice (Figure 1A) and compared the results with the control data. Remarkably, we observed drastically different transcriptional regulatory cascades in *cKO* hearts (Figure 1D and S3), with disturbances observed as early as at L1 (corresponding to P2). For example, the HIF-1 signaling pathway was enriched at L1 in control hearts and L9 in *cKO* hearts. Intriguingly, the most dramatically dysregulated pathways at L1 in *cKO* hearts are related to actin cytoskeleton, tight junctions and adherens junctions, which contribute to myocyte-myocyte communication[43] and regulate cell shape (Figure 1D and Figure S2). In addition, delay of ErbB signaling and disruptions in many metabolic networks (i.e., inactivation of fructose/mannose, propanoate and 2-oxocarboxylic acid metabolism, and the early activation of fatty acid metabolism) were also observed in c*KO* (Figure 1D). These disruptions were reflected by the inactivation of genes involved in mitochondrial biogenesis (Figure S2). Thus, RNF20 is critical for the coordination of gene networks during CM maturation as early as the initial postnatal response to the hypoxia-to-normoxia shift.

**Ablation of *Rnf20* after chamber formation leads to dilated cardiomyopathy**

*Mck-Cre* ablated *Rnf20* in cardiac and skeletal muscle cells beginning at E13.5[40] (Figure S4). Notably, the conditional deletion of *Rnf20* before chamber formation by *Mesp1-Cre* (activated at embryonic day 6.5)[44] (Figure S5A) was embryonically lethal (Table S2), and the mutant embryos displayed defects in heart morphogenesis as early as E9.5 (Figure S5B). In contrast, the depletion of RNF20 by *Mck-Cre* (*cKO*) did not affect the Mendelian ratio of each genotype at postnatal day 1 (P1) (Table S1), but it caused the mice to exhibit significantly lower body weights and sudden death within 8 weeks (P56) of birth (Figure 2A and 2B). At birth, both the control (*flox/flox*) and *cKO* mice had hearts with normal structures, and as expected, the cardiac sizes increased with time (Figure 2C). However, the *cKO* hearts developed a pronounced ventricular chamber dilation after P28 (Figure 2C). Echocardiographic analysis

revealed that the left ventricle (LV) systolic function, as measured by fractional
shortening (FS) and ejection fraction (EF), was significantly impaired after P28
(Figure 2D). We also detected abnormal electrocardiogram (ECG) patterns in *cKO*
mice beginning at P14 (Figure 2E). The depolarization and repolarization of the *cKO*
ventricles were disturbed, exhibiting prolonged QRS complex and QT interval (Figure
2E). ECG also revealed that the *cKO* mice developed irregular heartbeat at P28
(Figure S6). Thus, the *cKO* mice appear to suffer from dilated cardiomyopathy[41] and
arrhythmia, which may cause their sudden death. The cardiac defects were not general
effects of myopathy since mice with muscle-specific knockout of *Rnf20* (*Rnf20^{flox/flox}*;
*HSA-Cre*[42]) exhibited identical growth curves, survival rates, heart morphologies and
cardiac functions compared to control mice at P49 (Figure S7). Taken together, these
findings led us to conclude that RNF20-mediated transcriptional network is essential
for postnatal cardiac development and function.
199

**Structural, functional and metabolic transitions in postnatal heart**

Since the results up to this point implied that lack of RNF20 may disrupt
multiple processes in CM maturation, we sought to analyze the biological outcomes
of *Rnf20 cKO* in mouse heart. We first tested whether the loss of RNF20 affects the
polarization of intercalated discs (IDs), a hallmark of structural maturation of CMs[5,29]
(Figure 3A). The major components of IDs include N-cadherin (an adherens junction
protein), connexin 43 (a gap junction protein) and desmoplakin (a cytoplasmic
desmosomal protein). These components were circumferentially distributed at P2 and
gradually became polarized (P21), migrating to the ends of the CMs at P35 in the
control hearts (Figure 3B and S8A-B). In contrast, the migration was completely
abolished in the CMs of *cKO* mice. Next, we characterized the ultrastructural changes
in IDs during CM maturation using transmission electron microscopy (TEM). IDs
were visible and adjacent to the intercellular space between myocytes in the control
CMs at P21 and P35, with clear adherent junctions (white arrow in Figure 3C) and
desmosomes (yellow arrow in Figure 3C). By contrast, the structures of IDs in mutant
hearts were fragmented and disorganized (P21). By P35, the IDs had become
degenerated with widened gaps.
The structural maturation of CMs (Figure 3D) may also be assessed by the
localization of vinculin, a membrane-cytoskeletal protein controlling cytoskeletal
mechanics and cell shape[45] (Figure 3E). In the control mouse myocardium,

cytoplasmic vinculin migrated to the cell membrane during maturation (upper panel,

Figure 3E); however, the majority of vinculin remained in the cytoplasmic region in

RNF20-depleted mutants (lower panel, Figure 3E). Furthermore, we found that the

CMs in *cKO* mice failed to adopt a rectangular shape at P35 (Figure 3E and S8C).

Since CM cell shape is intimately linked to sarcomere alignment[46], we also inspected

the myofibril organization in postnatal hearts by TEM (Figure 3F). In contrast to the

control group, the myofibril alignment in CMs of *cKO* was distorted, exhibiting loose

actin filaments (yellow arrow in Figure 3F) and fuzzy Z-lines (white arrowhead in

Figure 3F); the M-line was also difficult to distinguish (yellow arrowhead in Figure

3F).

Last, we examined the effects of *Rnf20 cKO* on CM metabolic maturation, as

indicated by the maturation of mitochondria (Figure 3G)[4,5]. Control hearts showed a

gradual morphological shift from small, tubular and round mitochondria to large,

ovoid, well-organized organelles between P7 and P35 (Figure 3H and S8D), which is

typical of developing hearts[29]. However, this morphological maturation did not occur

in mitochondria of *cKO* hearts (Figure 3H and S8D). We next analyzed the

arrangement and density of cristae by TEM. Interestingly, the arrangement of cristae

was not altered in *cKO* mutants. However, a breakdown of both inner and outer

membrane was observed at postnatal day 35 (*cKO* in Figure 3H). Furthermore,

mitochondria cristae density (an indicator of mitochondria function)[5] was low from

P21 onwards in *cKO* hearts (Figure 3H). The mitochondria DNA (mtDNA) copy

number was reduced by 20% at P35 in *cKO* mice (Figure S8E). In addition, the

protein levels of major electron transport chain components, cytochrome c oxidase

and COX VI were markedly decreased in mutant heart (Figure 3I). The observed low

enzyme activities of cytochrome c oxidase (Figure 3J) and NADH oxidase (Figure

3K)[47] were likely responsible for the drastic reduction of ATP in *cKO* myocardium

(Figure 3L). Collectively, these data suggested that RNF20 extensively regulates CM

maturation by orchestrating gene networks and their functional outcomes.

**The dynamic chromatin opening during CM maturation**

Since RNF20 is known to be an epigenetic factor that modulates chromatin

compaction via monoubiquitylating histone H2B[21], we wondered whether

Rnf20-mediated epigenetic landscaping is involved in regulating transcriptional

networks during postnatal CM development (Figure 1D). To this end, we profiled the

chromatin accessibility by ATAC-seq[48] in postnatal hearts ate early (P7) and late (P21)

stages (Figure 4A) of the postnatal transcriptional maturation program. In control

hearts, we found the number of wide-spread accessible regions (peaks) was reduced

by almost 50% when comparing P21 to P7 (Figure 4B). Most of the diminished peaks

were at promoter regions (from 13,320 to 3,714 identified peaks; reduction of 72.1%;

Figure 4B and supplementary file 2). Intriguingly, this dynamic change in chromatin

accessibility was positively correlated with the level of RNF20 in control hearts

(Figure 4C), suggesting that the chromatin in early and late stages displayed

distinctive epigenetic landscapes.

We also found that P7 *cKO* hearts had only about half the number of peaks as P7

controls. The numbers of detected peaks were decreased in all genomic regions but

decreases were most predominant in promoter regions (from 13,320 to 5,567 peaks;

reduction of 58.2%; Figure 4B). These results suggest that RNF20 facilitates

chromatin accessibility, especially during the early stage of postnatal heart

development.

**Promoter-opening is required for postnatal gene expression**

Since the loss of *Rnf20* primarily impacted accessibility at gene promoters and

nucleosome eviction at gene promoter is one of the major events during

transcriptional activation[49], we postulated that RNF20-mediated chromatin opening

may affect postnatal transcriptional regulation. To test this hypothesis, we identified

genes with ATAC-seq peaks within 3 Kb of the transcription starting sites (TSS)

(Figure 4D). A large number of peak-associated genes (11,543 genes) was identified

in the control hearts at P7, while only 6,425 genes were associated with accessible

promoters in P7 *cKO* hearts (Figure 4D). This indicated that almost half (49.5%;

5,713) of all active genes in the P7 heart may require RNF20 for promoter opening;

such genes are hereafter called "RNF20-open" genes (Figure 4D). KEGG pathway

enrichment analysis shows that "RNF20-open" genes were enriched in pathways

related to CM maturation, such as calcium signaling, cell cycle, fatty acid metabolism,

gap junction, and cardiac muscle contraction[4,5] (Figure S9). Only a small fraction of

genes (595 genes) was classified as "*cKO*-open" genes (Figure 4D), indicating their

promoters were accessible under RNF20 depletion. Notably, the "*cKO*-open" genes

showed no apparent functional links with heart development (Figure S9). These

findings revealed that RNF20-mediated chromatin opening at P7 regulates

transcriptional programs in CM maturation and thus may play a critical role in
coordinating postnatal gene networks.

At P21, the number of peak-associated genes in the control hearts was comparable
to that in the *cKO* hearts (5,288 in the control and 5,108 in *cKO*). However, the genes
in these two groups overlapped by only about 50% (Figure 4D), and there were
roughly equal numbers of "RNF20-open" (2646) and "*cKO*-open" (2466) genes. Of
note, the "RNF20-open" genes were involved in metabolism-related pathways, such
as pyruvate metabolism, glucagon signaling, regulation of lipolysis in adipocytes
(Figure S10). The RNF20-associated pathways mostly correlated with enriched
functions at later stages of CM maturation (Figure 1D and S3). In contrast, the
"*cKO*-open" genes were enriched in Wnt signaling pathway, ECM-receptor
interaction and neurotrophin signaling (Figure S10), which were highly expressed at
the neonatal phase (P2 to P7) (Figure 1D). Thus, the ablation of *Rnf20* causes the
transcriptional program of P21 heart to retain similarity with the initial postnatal stage
and impairs CM maturation.

**Chromatin remodeling facilitates dynamic binding of TFs during postnatal heart**
**development**

Previous studies reported that gene network transitions during heart development
are tightly controlled by TFs[50], so we wondered whether RNF20-mediated changes in
chromatin accessibility might affect the binding of TFs to modulate postnatal heart
development. To answer this question, we used a computational tool, TOBIAS[51] to
perform a TF footprint analysis with the ATAC-seq data (Figure 4E). This tool
assigned differential binding scores (DBSs) to 746 TFs, representing the relative
binding efficiency between the control and *cKO* groups (Figure 4F; a full list of
analyzed TFs can be found in Supplementary file 3). At P7, 199 RNF20-dependent
and 39 *cKO*-dependent TFs were respectively identified by their binding activity
($-\log_{10}$p-value > 100) in the control (DBS > 0.25) and *cKO* (DBS < -0.25) groups
(Figure 4F and supplementary file 3). Gene Ontology (GO) enrichment analysis
further showed that the RNF20-dependent TFs were involved in cardiac development
and CM maturation (Figure 4F and S11A). Remarkably, many of the
RNF20-dependent TFs at P7 turned out to be *cKO*-dependent (DBS < 0) at P21
(Figure 4F), which is in line with the chromatin opening profile in the *cKO* hearts
(p21 in Figure 4D). These results provide further explanation for the extensive

transcriptional dysregulation caused by deletion of *Rnf20* (Figure 2D). Conversely,

the binding patterns of the 39 *cKO*-dependent TFs became sparsely distributed, and

their functional categories were unrelated to heart development/function (Figure 4F

and S11B). Taken together, these observations suggest that the elevated levels of

RNF20 at the early stage (P7) of CM maturation are necessary for proper binding of

cardiac TFs that drive transcriptional gene networks, and the continual expression of

RNF20 at lower levels is required to sustain gene-network transitions (Figure 4G).

Thus, RNF20 is centrally involved in establishing a dynamic epigenetic environment

that facilitates the execution of time-ordered gene networks during postnatal heart

development (Figure 1D).

**Pioneer factors contribute to chromatin opening**

Almost 50% of genes in the control hearts of P7 (50.5%, 5830 genes) and P21

(46.44%, 2642 genes) did not require RNF20 for promoter opening (Figure 4D) and

were defined as "persistent-open" genes. The products of "persistent-open" genes

included the regulatory factors that are essential for CM maturation, such as HIF-1

signaling, ErbB signaling pathway and tight junctions (Figure 1D, S9 and S10),

suggesting there must be other factors that maintain chromatin opening for these

maturation aspects. Interestingly, many pioneer factors, including GATA families

(GATA4, GATA6) and forkhead box families (FOXA1, FOXA2) (yellow dots in

Figure 4F), did not rely on RNF20 for recognition site binding [low Dbs (x-axis) and

low p-value (y-axis); Supplementary file 3]. Pioneer factors are able to open closed

chromatin sites in order to implement cell fate programs[52]. Consistent with their

pioneering roles during embryonic heart development[53,54], expression levels of

GATA4 and GATA6 were high at early postnatal stages (L1-L2 by TO-GCN analysis,

Figure 2C). Thus, the promoter accessibilities of "persistent-open" genes are likely

attributable to the cooperation of pioneer factors via RNF20-independent mechanisms.

Intriguingly, the time-ordered expression of some "persistent-open" genes were

disrupted in the absence of RNF20 (Figure 1D, S9 and S10). As such, our data also

suggested that RNF20 may help optimize expression of "persistent-open" genes in

postnatal heart without affecting chromatin accessibility at their promoters.

**Discussion**

In this study, we found that dynamic transitions of chromatin state are required

for the time-ordered transcription of postnatal genes and CM maturation. By TO-GCN

analysis, we identified a transcriptional regulatory cascade that orchestrates the

sequential structural, cell-cycle, functional and metabolic maturation programs in the

postnatal heart (Figure 5). Evidence from our study suggests that RNF20 plays a key

role in regulating the time-ordered gene networks during postnatal CM maturation by

modulating chromatin accessibility. Mechanistically, RNF20-mediated chromatin

opening in the early postnatal heart globally resolves epigenetic barriers to allow the

maturation program to proceed. The decrease of RNF20 expression at the end-stage of

maturation then reestablishes a more compact epigenetic state with less transcriptional

flexibility, which is required for the mature cells. RNF20 is likely to modulate

chromatin accessibility through its epigenetic target, H2Bub [27], as this modification

has been shown to disrupt local and higher-order chromatin compaction *in vitro* and

*in vivo* [21,22]. However, our data do not exclude the possibility that RNF20 may

regulate other epigenetic factors to affect the postnatal chromatin landscape.

The dynamics of chromatin accessibility are modulated by the combined effects

of chromatin remodelers[56,57], TFs[58] and PTMs[59]. However, it remains uncertain how

accessibility of chromatin is modulated in response to external stimuli and

developmental cues. Overwhelming evidence supports the roles of different PTMs in

generating functional chromatin states[60], and PTMs are known to influence chromatin

accessibility via indirect mechanisms, such as altering TF binding and nucleosome

affinity of chromatin remodelers. Intriguingly, however, uncertainty remains about

how PTMs may directly contribute to accessibility remodeling of chromatin templates.

Our finding that RNF20 helps to establish a dynamic chromatin environment for CM

maturation has some intriguing implications. First, while dynamic regulation of the

RNF20-H2Bub pathway has been linked to embryonic stem cell differentiation[61-63],

the deletion of *Rnf20* after cardiac morphogenesis does not change cell fates or

identity during prenatal or neonatal heart development, Instead, it dramatically

compromises CM maturation. At the molecular level, we found that RNF20

influences chromatin opening in nearly half of the gene promoters involved in CM

maturation, suggesting that there is a division of labor for modulating chromatin

accessibility. The cardiac-specific pioneer factors most likely specialize in modulating

the chromatin states in those parts of the genome required for cardiac cell fate. In

contrast, RNF20 (possibly via H2Bub) may promote chromatin plasticity in the rest of

genome, allowing the heart cells to undergo remodeling from a fetal to an adult state

without affecting core cell identity factors. Thus, our findings suggest a novel
mechanism in which PTMs directly and extensively modulate chromatin accessibility
dynamics to orchestrate developmental processes.

The mechanisms of regulating and coordinating constituent processes during CM
maturation have received increasing attention because it is not yet possible to achieve
complete maturation of pluripotent stem cell (PSC)-induced CMs (iPSC-CMs)[4]. Our
TO-GCN analysis revealed that postnatal heart development is initiated by integrated
signals, including transient expression of hypoxia-inducible factor 1-alpha (HIF-1α),
cell-cell connection (adherens and tight junction) and structural organization
(regulation of actin cytoskeleton). Thus, the coordinated activation of these pathways
may help initiate CM maturation. In support of this idea, the synergistic application of
electrical/mechanical stimulation, extracellular matrix and non-CM co-cultures can
facilitate iPSC-CMs maturation[29,61]. Furthermore, the inhibition of HIF1α enhances
the contractile, electrophysiologic and metabolic maturation of iPSC-CMs[62,63]. The
maturation of CMs involves a complex transcriptional network coordinated by cardiac
and non-cardiac cells[29,43,64-66]. Our study provides clear staging information and
delineation of transcriptionally active maturation processes based on analyses of
whole heart transcriptomes using a comparative transcriptomic method[30]. Thus, our
approach may serve to identify the complete set of processes underlying CM
maturation. Furthermore, our mouse genetic results imply that RNF20 mutations may
cause CHD, as supported by previous observations in CHD patients and in
*Xenopus*[16,17]. The results also suggest that functional RNF20 abnormalities beginning
at late-gestation periods could be related to early-onset cardiomyopathy in children
and young adults, even without observable effects on heart morphogenesis.

In summary, our findings demonstrate that dynamic chromatin accessibility is
critical for the time-ordered expression of postnatal genes during CM maturation, and
the epigenetic factor RNF20 plays a decisive role in mediating postnatal chromatin
accessibility remodeling. We also offer a complete roadmap of gene cascades that
direct CM maturation, providing insights that may improve therapies for CHD and
facilitate the development of iPSC-CMs to treat heart disease.

**Acknowledgments**

The authors thank Dr. Ting-Fen Tsai for constructive comments during manuscript
preparation. We are grateful to the staff at the Core Facility of the Institute of Cellular
and Organismic Biology, Academia Sinica, for EM service and confocal assistance;
the Transgenic Core Facility of the Institute of Molecular Biology, Academia Sinica,
for transgenic mice production and Genomics core facility of the Institute of
Molecular Biology, Academia Sinica, for NGS service. The authors also thank the
Taiwan Mouse Clinic, Academia Sinica, and Taiwan Animal Consortium for the
technical support in electrocardiography and high-frequency ultrasound, and Dr.
Marcus Calkins for English editing. This work was supported by the intramural
funding of the Institute of Cellular and Organismic Biology, Academia Sinica, and by
grants from the Ministry of Science and Technology, Taiwan (MOST
109-2320-B-001-017-MY3).

**Author contributions**

C.-F.K. and Y.-T.Y. conceived the study. C.-F.K., Y.-T.Y. and C.-C.C. designed the
experiments. C.-Y.L., H.-Y.T., Y.-L.S., H.-H.Y. and Y.-R.L. performed the
experiments. Y.-M.C. designed and performed the bioinformatic analysis. C.-Y.L. and
C.-L.H. analyzed the data. Y.-L.S. and H.-H.Y. provided study materials and animals.
C.-Y.L. and C.-F.K. wrote the manuscript.

**Declaration of interests**

The authors declare no competing interests.

**Supplementary information**

Experimental designs, materials, additional information and references are available
in the supplemented files.
445

**References**

1     Epstein, J. A. Franklin H. Epstein Lecture. Cardiac development and
implications for heart disease. *N Engl J Med* 363, 1638-1647,
doi:10.1056/NEJMra1003941 (2010).
2     Hoffman, J. I. & Kaplan, S. The incidence of congenital heart disease. *J Am
Coll Cardiol* 39, 1890-1900, doi:10.1016/s0735-1097(02)01886-7 (2002).
3     McMurray, J. J. Clinical practice. Systolic heart failure. *N Engl J Med* 362,
228-238, doi:10.1056/NEJMcp0909392 (2010).
4     Kannan, S. & Kwon, C. Regulation of cardiomyocyte maturation during

| 455 | | critical perinatal window. *J Physiol* 598, 2941-2956, doi:10.1113/JP276754 |
| 456 | | (2020). |
| 457 | 5 | Guo, Y. & Pu, W. T. Cardiomyocyte Maturation: New Phase in Development. |
| 458 | | *Circ Res* 126, 1086-1106, doi:10.1161/CIRCRESAHA.119.315862 (2020). |
| 459 | 6 | Latchman, D. S. Transcription factors: an overview. *Int J Biochem Cell Biol* 29, |
| 460 | | 1305-1312, doi:10.1016/s1357-2725(97)00085-x (1997). |
| 461 | 7 | Liu, L., Jin, G. & Zhou, X. Modeling the relationship of epigenetic |
| 462 | | modifications to transcription factor binding. *Nucleic Acids Res* 43, 3873-3885, |
| 463 | | doi:10.1093/nar/gkv255 (2015). |
| 464 | 8 | Bannister, A. J. & Kouzarides, T. Regulation of chromatin by histone |
| 465 | | modifications. *Cell Res* 21, 381-395, doi:10.1038/cr.2011.22 (2011). |
| 466 | 9 | Liu, L. *et al.* An integrated chromatin accessibility and transcriptome |
| 467 | | landscape of human pre-implantation embryos. *Nat Commun* 10, 364, |
| 468 | | doi:10.1038/s41467-018-08244-0 (2019). |
| 469 | 10 | Thomas, S. *et al.* Dynamic reprogramming of chromatin accessibility during |
| 470 | | Drosophila embryo development. *Genome Biol* 12, R43, |
| 471 | | doi:10.1186/gb-2011-12-5-r43 (2011). |
| 472 | 11 | Li, D., Shu, X., Zhu, P. & Pei, D. Chromatin accessibility dynamics during cell |
| 473 | | fate reprogramming. *EMBO Rep* 22, e51644, doi:10.15252/embr.202051644 |
| 474 | | (2021). |
| 475 | 12 | Jarrell, D. K., Lennon, M. L. & Jacot, J. G. Epigenetics and Mechanobiology |
| 476 | | in Heart Development and Congenital Heart Disease. *Diseases* 7, |
| 477 | | doi:10.3390/diseases7030052 (2019). |
| 478 | 13 | Paige, S. L. *et al.* A temporal chromatin signature in human embryonic stem |
| 479 | | cells identifies regulators of cardiac development. *Cell* 151, 221-232, |
| 480 | | doi:10.1016/j.cell.2012.08.027 (2012). |
| 481 | 14 | Wamstad, J. A. *et al.* Dynamic and coordinated epigenetic regulation of |
| 482 | | developmental transitions in the cardiac lineage. *Cell* 151, 206-220, |
| 483 | | doi:10.1016/j.cell.2012.07.035 (2012). |
| 484 | 15 | Gilsbach, R. *et al.* Dynamic DNA methylation orchestrates cardiomyocyte |
| 485 | | development, maturation and disease. *Nat Commun* 5, 5288, |
| 486 | | doi:10.1038/ncomms6288 (2014). |
| 487 | 16 | Zaidi, S. *et al.* De novo mutations in histone-modifying genes in congenital |
| 488 | | heart disease. *Nature* 498, 220-+, doi:10.1038/nature12141 (2013). |
| 489 | 17 | Robson, A. *et al.* Histone H2B monoubiquitination regulates heart |
| 490 | | development via epigenetic control of cilia motility. *Proc Natl Acad Sci U S A* |
| 491 | | 116, 14049-14054, doi:10.1073/pnas.1808341116 (2019). |
| 492 | 18 | VanDusen, N. J. *et al.* Massively parallel in vivo CRISPR screening identifies |

RNF20/40 as epigenetic regulators of cardiomyocyte maturation. *Nat Commun*
12, 4442, doi:10.1038/s41467-021-24743-z (2021).

19    Fuchs, G. & Oren, M. Writing and reading H2B monoubiquitylation. *Biochim*
*Biophys Acta* 1839, 694-701, doi:10.1016/j.bbagrm.2014.01.002 (2014).

20    VanDusen, N. J. *et al.* Massively parallel in vivo CRISPR screening identifies
RNF20/40 as epigenetic regulators of cardiomyocyte maturation. *Nat Commun*
12, 4442, doi:10.1038/s41467-021-24743-z (2021).

21    Fierz, B. *et al.* Histone H2B ubiquitylation disrupts local and higher-order
chromatin compaction. *Nat Chem Biol* 7, 113-119, doi:10.1038/nchembio.501
(2011).

22    Moyal, L. *et al.* Requirement of ATM-dependent monoubiquitylation of
histone H2B for timely repair of DNA double-strand breaks. *Mol Cell* 41,
529-542, doi:10.1016/j.molcel.2011.02.015 (2011).

23    Zhang, F. & Yu, X. WAC, a functional partner of RNF20/40, regulates histone
H2B ubiquitination and gene transcription. *Mol Cell* 41, 384-397,
doi:10.1016/j.molcel.2011.01.024 (2011).

24    Kim, J., Hake, S. B. & Roeder, R. G. The human homolog of yeast BRE1
functions as a transcriptional coactivator through direct activator interactions.
*Mol Cell* 20, 759-770, doi:10.1016/j.molcel.2005.11.012 (2005).

25    Minsky, N. *et al.* Monoubiquitinated H2B is associated with the transcribed
region of highly expressed genes in human cells. *Nat Cell Biol* 10, 483-488,
doi:10.1038/ncb1712 (2008).

26    Xiao, T. *et al.* Histone H2B ubiquitylation is associated with elongating RNA
polymerase II. *Mol Cell Biol* 25, 637-651,
doi:10.1128/MCB.25.2.637-651.2005 (2005).

27    Pavri, R. *et al.* Histone H2B monoubiquitination functions cooperatively with
FACT to regulate elongation by RNA polymerase II. *Cell* 125, 703-717,
doi:10.1016/j.cell.2006.04.029 (2006).

28    Krishnan, A. *et al.* A detailed comparison of mouse and human cardiac
development. *Pediatr Res* 76, 500-507, doi:10.1038/pr.2014.128 (2014).

29    Karbassi, E. *et al.* Cardiomyocyte maturation: advances in knowledge and
implications for regenerative medicine. *Nat Rev Cardiol* 17, 341-359,
doi:10.1038/s41569-019-0331-x (2020).

30    Chang, Y. M. *et al.* Comparative transcriptomics method to infer gene
coexpression networks and its applications to maize and rice leaf
transcriptomes. *Proceedings of the National Academy of Sciences of the*
*United States of America* 116, 3091-3099, doi:10.1073/pnas.1817621116
(2019).

31    Semenza, G. L. Hydroxylation of HIF-1: oxygen sensing at the molecular
level. *Physiology (Bethesda)* 19, 176-182, doi:10.1152/physiol.00001.2004
(2004).

32    Porrello, E. R. *et al.* Transient regenerative potential of the neonatal mouse
heart. *Science* 331, 1078-1080, doi:10.1126/science.1200708 (2011).

33    Grais, I. M. & Sowers, J. R. Thyroid and the heart. *Am J Med* 127, 691-698,
doi:10.1016/j.amjmed.2014.03.009 (2014).

34    Sanchez-Soria, P. & Camenisch, T. D. ErbB signaling in cardiac development
and disease. *Semin Cell Dev Biol* 21, 929-935,
doi:10.1016/j.semcdb.2010.09.011 (2010).

35    Bertrand, L., Horman, S., Beauloye, C. & Vanoverschelde, J. L. Insulin
signalling in the heart. *Cardiovasc Res* 79, 238-248, doi:10.1093/cvr/cvn093
(2008).

36    Kruger, M. *et al.* Thyroid hormone regulates developmental titin isoform
transitions via the phosphatidylinositol-3-kinase/ AKT pathway. *Circ Res* 102,
439-447, doi:10.1161/CIRCRESAHA.107.162719 (2008).

37    Feng, N. *et al.* Constitutive BDNF/TrkB signaling is required for normal
cardiac contraction and relaxation. *Proc Natl Acad Sci U S A* 112, 1880-1885,
doi:10.1073/pnas.1417949112 (2015).

38    Riehle, C. & Abel, E. D. Insulin Signaling and Heart Failure. *Circ Res* 118,
1151-1169, doi:10.1161/CIRCRESAHA.116.306206 (2016).

39    Lopaschuk, G. D. & Jaswal, J. S. Energy metabolic phenotype of the
cardiomyocyte during development, differentiation, and postnatal maturation.
*J Cardiovasc Pharmacol* 56, 130-140, doi:10.1097/FJC.0b013e3181e74a14
(2010).

40    Johnson, J. E., Wold, B. J. & Hauschka, S. D. Muscle Creatine-Kinase
Sequence Elements Regulating Skeletal and Cardiac-Muscle Expression in
Transgenic Mice. *Molecular and Cellular Biology* 9, 3393-3399, doi:Doi
10.1128/Mcb.9.8.3393 (1989).

41    Japp, A. G., Gulati, A., Cook, S. A., Cowie, M. R. & Prasad, S. K. The
Diagnosis and Evaluation of Dilated Cardiomyopathy. *J Am Coll Cardiol* 67,
2996-3010, doi:10.1016/j.jacc.2016.03.590 (2016).

42    Miniou, P. *et al.* Gene targeting restricted to mouse striated muscle lineage.
*Nucleic Acids Res* 27, e27, doi:10.1093/nar/27.19.e27 (1999).

43    Tirziu, D., Giordano, F. J. & Simons, M. Cell communications in the heart.
*Circulation* 122, 928-937, doi:10.1161/CIRCULATIONAHA.108.847731
(2010).

44    Saga, Y. *et al.* MesP1: a novel basic helix-loop-helix protein expressed in the

nascent mesodermal cells during mouse gastrulation. *Development* 122,
2769-2778 (1996).

45   Goldmann, W. H. & Ingber, D. E. Intact vinculin protein is required for control
of cell shape, cell mechanics, and rac-dependent lamellipodia formation.
*Biochem Biophys Res Commun* 290, 749-755, doi:10.1006/bbrc.2001.6243
(2002).

46   Bray, M. A., Sheehy, S. P. & Parker, K. K. Sarcomere alignment is regulated
by myocyte shape. *Cell Motil Cytoskeleton* 65, 641-651,
doi:10.1002/cm.20290 (2008).

47   Anraku, Y. Bacterial electron transport chains. *Annu Rev Biochem* 57, 101-132,
doi:10.1146/annurev.bi.57.070188.000533 (1988).

48   Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and
enables interrogation of frozen tissues. *Nat Methods* 14, 959-962,
doi:10.1038/nmeth.4396 (2017).

49   John, S. *et al.* Chromatin accessibility pre-determines glucocorticoid receptor
binding patterns. *Nat Genet* 43, 264-268, doi:10.1038/ng.759 (2011).

50   Uosaki, H. *et al.* Transcriptional Landscape of Cardiomyocyte Maturation.
*Cell Rep* 13, 1705-1716, doi:10.1016/j.celrep.2015.10.032 (2015).

51   Bentsen, M. *et al.* ATAC-seq footprinting unravels kinetics of transcription
factor binding during zygotic genome activation. *Nat Commun* 11, 4267,
doi:10.1038/s41467-020-18035-1 (2020).

52   Iwafuchi-Doi, M. & Zaret, K. S. Pioneer transcription factors in cell
reprogramming. *Genes Dev* 28, 2679-2692, doi:10.1101/gad.253443.114
(2014).

53   Sharma, A. *et al.* GATA6 mutations in hiPSCs inform mechanisms for
maldevelopment of the heart, pancreas, and diaphragm. *Elife* 9,
doi:10.7554/eLife.53278 (2020).

54   Tremblay, M., Sanchez-Ferras, O. & Bouchard, M. GATA transcription factors
in development and disease. *Development* 145, doi:10.1242/dev.164384
(2018).

55   Janes, J. *et al.* Chromatin accessibility dynamics across C. elegans
development and ageing. *Elife* 7, doi:10.7554/eLife.37344 (2018).

56   Iurlaro, M. *et al.* Mammalian SWI/SNF continuously restores local
accessibility to chromatin. *Nat Genet* 53, 279-287,
doi:10.1038/s41588-020-00768-w (2021).

57   Schick, S. *et al.* Acute BAF perturbation causes immediate changes in
chromatin accessibility. *Nat Genet* 53, 269-278,
doi:10.1038/s41588-021-00777-3 (2021).

58    Klemm, S. L., Shipony, Z. & Greenleaf, W. J. Chromatin accessibility and the
regulatory epigenome. *Nat Rev Genet* 20, 207-220,
doi:10.1038/s41576-018-0089-8 (2019).

59    Allis, C. D. & Jenuwein, T. The molecular hallmarks of epigenetic control. *Nat*
*Rev Genet* 17, 487-500, doi:10.1038/nrg.2016.59 (2016).

60    Ernst, J. *et al.* Mapping and analysis of chromatin state dynamics in nine
human cell types. *Nature* 473, 43-49, doi:10.1038/nature09906 (2011).

61    Scuderi, G. J. & Butcher, J. Naturally Engineered Maturation of
Cardiomyocytes. *Front Cell Dev Biol* 5, 50, doi:10.3389/fcell.2017.00050
(2017).

62    Hu, D. *et al.* Metabolic Maturation of Human Pluripotent Stem Cell-Derived
Cardiomyocytes by Inhibition of HIF1alpha and LDHA. *Circ Res* 123,
1066-1079, doi:10.1161/CIRCRESAHA.118.313249 (2018).

63    Gentillon, C. *et al.* Targeting HIF-1alpha in combination with PPARalpha
activation and postnatal factors promotes the metabolic maturation of human
induced pluripotent stem cell-derived cardiomyocytes. *J Mol Cell Cardiol* 132,
120-135, doi:10.1016/j.yjmcc.2019.05.003 (2019).

64    Pinto, A. R. *et al.* Revisiting Cardiac Cellular Composition. *Circ Res* 118,
400-409, doi:10.1161/CIRCRESAHA.115.307778 (2016).

65    Ieda, M. *et al.* Cardiac fibroblasts regulate myocardial proliferation through
beta1 integrin signaling. *Dev Cell* 16, 233-244,
doi:10.1016/j.devcel.2008.12.007 (2009).

66    Brutsaert, D. L. Cardiac endothelial-myocardial signaling: its role in cardiac
growth, contractile performance, and rhythmicity. *Physiol Rev* 83, 59-115,
doi:10.1152/physrev.00017.2002 (2003).

67    Liu, R. *et al.* Tead1 is required for perinatal cardiomyocyte proliferation. *PLoS*
*One* 14, e0212017, doi:10.1371/journal.pone.0212017 (2019).

68    Pikkarainen, S. *et al.* GATA-4 is a nuclear mediator of mechanical
stretch-activated hypertrophic program. *J Biol Chem* 278, 23807-23816,
doi:10.1074/jbc.M302719200 (2003).

69    Xiang, F. L., Guo, M. & Yutzey, K. E. Overexpression of Tbx20 in Adult
Cardiomyocytes Promotes Proliferation and Improves Cardiac Function After
Myocardial Infarction. *Circulation* 133, 1081-1092,
doi:10.1161/CIRCULATIONAHA.115.019357 (2016).

70    Wang, J., Liu, S., Heallen, T. & Martin, J. F. The Hippo pathway in the heart:
pivotal roles in development, disease, and regeneration. *Nat Rev Cardiol* 15,
672-684, doi:10.1038/s41569-018-0063-3 (2018).

71    Kolodziejczyk, S. M. *et al.* MEF2 is upregulated during cardiac hypertrophy

and is required for normal post-natal growth of the myocardium. *Curr Biol* 9,
1203-1206, doi:10.1016/S0960-9822(00)80027-5 (1999).
72   Ferdous, A. *et al.* FoxO1-Dio2 signaling axis governs cardiomyocyte thyroid
hormone metabolism and hypertrophic growth. *Nat Commun* 11, 2551,
doi:10.1038/s41467-020-16345-y (2020).
73   Martin, O. J. *et al.* A role for peroxisome proliferator-activated receptor
gamma coactivator-1 in the control of mitochondrial dynamics during
postnatal cardiac growth. *Circ Res* 114, 626-636,
doi:10.1161/CIRCRESAHA.114.302562 (2014).

Figure Legends
Figure 1. Comparative analysis of time-series transcriptomes from mouse hearts.
(A) Schematic of sample collection strategy from control and *cKO* mice for RNA-seq
and time-ordered gene co-expression network (TO-GCN) analysis. (B) PCA plot of
RNA-seq results. n=2 per group. (C) The TO-GCN structure with TF genes as nodes
(blue circles) and the heatmap of average normalized RPKMs (z scores) at each
time-point; TF genes were evaluated at nine levels in the control heart. Examples of
TFs gene lists from each level in the control mice heart are shown. TFs at L1 and L2
were associated with cardiac development, e.g., Gata4, Nkx2.5, Tbx5, Tbx20 and
Tead1, in line with their known contributions to neonatal cell proliferation and
maturation[67-69] (Supplementary file 1). The YAP co-factors (Smad3, Runx1, Smad4/5)
were present at L3 and L4 (P7), consistent with their roles in the cardiac homeostasis
and cell cycle control at postnatal stages[70]. The postnatal activation of Mef2 family
(shown at L5 and L7) is required for normal postnatal growth of the myocardium[71],
and Foxo1 (shown at L6) is necessary for hypertrophic growth[72]. The PPARs shown
at L8 are responsible for mitochondrial biogenesis and metabolic switching of CMs[73].
(D) KEGG analysis for co-expressed genes in the control heart are displayed in order
(by TO-GCN level); the same list from *cKO* heart is shown for comparison.
677

Figure 2. Depletion of RNF20 after chamber formation leads to dilated
cardiomyopathy.

(A) Growth curve and (B) Kaplan-Meier survival curves of *Rnf20 cKO*
(*Rnf20$^{flox/flox}$;Mck-Cre*) mice and control (*Rnf20$^{flox/flox}$, Rnf20$^{flox/+}$, Mck-Cre Rnf20$^{flox/+}$*)
littermates. (C) Hematoxylin/eosin-stained sagittal (left) and cross (right) sections of
control (*Rnf20$^{flox/flox}$*) and *Rnf20 cKO* hearts at P2, P14, P28 and P42. (D) (Left panel)
Representative 2D and M-mode echocardiographic images of the *RNF20 cKO* and
control hearts. (Right panel) Echocardiographic analysis in the control and *cKO* mice
at different stages. Left ventricular ejection fraction (EF) and fractional shortening
(FS) are shown. Data are presented as mean ± SD; n=6 per group. * $p<0.05$; ** $p<$
$0.01$; *** $p< 0.001$. (E) Representative telemetric electrocardiogram (ECG) patterns
of control and *cKO* mice at P14. Quantification of the ECG changes in *Rnf20 cKO*
and control mice from P14 to P42 are shown; data are presented as mean ± SD. n=5
per group. * $p<0.05$; ** $p< 0.01$; *** $p< 0.001$.

Figure 3. Postnatal maturation defects in *Rnf20 cKO* mice heart.

(A) Model of the dynamic formation of intercalated discs (IDs; composed of

desmosome, gap junction and adherens junction) during CM maturation. (B) Confocal

micrographs of cross sections of the cardiac muscle of control and cKO mice at P2,

P21 and P35. Specific antibodies were used to identify the distributions of ID

component, N-Cadherin, and the costamere marker, Vinculin. Nuclei were visualized

by Hoechst 33342 staining; n=3 per group. Scale bar: 10 μm. (C) Transmission

electron microscopy images (TEMs) of ventricular myocardium from *Rnf20 cKO* and

control mice at P21 and P35. The insets show representative ID morphologies in

control and *cKO* mice. Abnormal structures were observed in mutant hearts, with

widened gaps in the IDs. White arrowheads: adherent junctions; yellow arrowheads:

desmosomes. Scale bar: 0.5 μm. (D) Model of the morphological changes in

immature CMs; increased size and altered organization of the contractile cytoskeleton.

(E) Staining for Vinculin (costamere marker) in ventricular sections from *Rnf20 cKO*

and control mice at P2, P21 and P35. Scale bar: 50 μm. (F) The ultrastructure of the

sarcomeres in CMs of control and *Rnf20 cKO* mice. Abnormal Z-line (white

arrowheads) and loose, distorted myofibrils (yellow arrowheads) were observed in

*cKO* mice at P21 and P35. Scale bar: 0.5 μm. (G) Schematic depicts the maturation of

mitochondria during CM maturation. (H) TEMs of mitochondria in ventricular CMs

(Scale bar: 0.5 μm). Quantification of number of cristae per μm suggested a

significantly less dense cristae structure in *cKO* mice heart than in control at P35

(*n* = 3 for each group). * $p<0.05$; ** $p< 0.01$; *** $p< 0.001$. (I) Immunoblotting for

mitochondrial proteins in control and *Rnf20 cKO* hearts (n=3). COX IV: cytochrome c

oxidase IV; VDAC: voltage-dependent anion channel; PHB1: prohibitins; PDH:

pyruvate dehydrogenase; SDHA: succinate dehydrogenase A; HSP60: heat shock

protein 60. (J-L) The mitochondrial enzyme activities for (J) cytochrome c oxidase,

(K) NADH oxidase and (L) ATP production were measured in P35 hearts (*n* = 3

for each, normalized to mitochondrial protein). * $p<0.05$; ** $p< 0.01$; *** $p< 0.001$.

Figure 4. Chromatin accessibility landscape in postnatal heart.

(A) Experimental design for sample collection from control and *cKO* mice. (B)

Numbers of peak calls from the P7 and P21 control or mutant hearts are shown

individually. Color indicates the type of genomic region overlapped by the peak. UTR,

untranslated region. (C) Immunoblot analysis of RNF20 in control heart. (D) Venn

diagram of peak-associated genes in the control and *Rnf20 cKO* heart at P7 (Left) and

P21 (Right), respectively. Middle, model of peak associated genes from ATAC-seq (3

kb ± transcription start site). (E) Schematic illustrates the dynamics of TF binding

(blue), Tn5 insertion (green), and the concept of footprint analysis using ATAC-seq

data. (F) Pairwise comparisons of TF activities in control and *cKO* mice. The volcano

plots show differential binding activity versus -$\log_{10}$ p-value (both calculated by

TOBIAS) for all investigated TF motifs; each dot represents one motif. The

control-specific TFs (RNF20-dependent) are labeled in red, and the *cKO*-specific TFs

(*cKO*-dependent) are labeled in blue. Gene ontology analyses of RNF20-dependent

and *cKO*-dependent TFs are shown. (G) Illustration of RNF20-mediated chromatin

remodeling in postnatal heart. P7: RNF20 is upregulated, chromatin accessibility at

regulatory elements (REs) of genes are upregulated, TF binding is upregulated. P21:

RNF20 is downregulated, chromatin accessibility at REs of genes are downregulated,

TF binding is downregulated.

Figure 5. Working model.

Dynamic RNF20 expression after birth mediates remodeling of chromatin

accessibility to allow the induction of a time-ordered postnatal transcription program.

RNF20 is highly expressed at the early postnatal stage, leading to a global chromatin

decompaction that facilitates the binding of transcription factors (TFs) to early

postnatal genes. RNF20 level decreases at the later stage of heart maturation, leading

to chromatin compaction and a restriction of TF binding that establishes and

maintains the mature status of the heart.

**Figure 1**

**Figure 2**

**Figure 3**

**Figure 4**

**Figure 5**