Ret loss-of-function decreases neural crest progenitor proliferation and restricts developmental fate potential during enteric nervous system development

Elizabeth Vincent [1,*], Sumantra Chatterjee [2,3,*], Gabrielle H. Cannon [1], Dallas Auer [1], Holly Ross [1], Aravinda Chakravarti [2,3,#], Loyal A. Goff [1,4,5,#]

Affiliations
1. McKusick-Nathans Department of Genetic Medicine, Johns Hopkins University School of Medicine
2. Center for Human Genetics and Genomics, NYU Grossman School of Medicine
3. Department of Neuroscience and Physiology, NYU Grossman School of Medicine
4. Solomon H. Snyder Department of Neuroscience, Johns Hopkins University School of Medicine
5. Kavli Neurodiscovery Institute, Johns Hopkins University
* Authors contributed equally
# Co-corresponding authors

Abstract

The receptor tyrosine kinase gene *RET* plays a critical role in the fate specification of enteric neural crest-derived cells (ENCDCs) during enteric nervous system (ENS) development. Pathogenic *RET* loss of function (LoF) alleles are associated with Hirschsprung disease (HSCR), which is marked by aganglionosis of the gastrointestinal (GI) tract. ENCDCs invade the developing GI tract, proliferate, migrate caudally, and differentiate into all of the major ENS cell types. Although the major phenotypic consequences and the underlying transcriptional changes from Ret LoF in the developing ENS have been described, its cell type and state-specific effects are unknown. Consequently, we performed single-cell RNA sequencing (scRNA-seq) on an enriched population of ENCDCs isolated from the developing GI tract of *Ret* null heterozygous and homozygous mouse embryos at embryonic day (E)12.5 and E14.5. We demonstrate four significant findings: (1) *Ret*-expressing ENCDCs are a heterogeneous population composed of ENS progenitors as well as glial and neuronal committed cells; (2) neurons committed to a predominantly inhibitory motor neuron developmental trajectory are not produced under Ret LoF, leaving behind a mostly excitatory motor neuron developmental program; (3) HSCR-associated and *Ret* gene regulatory network genes exhibit distinct expression patterns across *Ret*-expressing ENCDC with their expression impacted by *Ret* LoF; and (4) *Ret* deficiency leads to precocious differentiation and reduction in the number of proliferating ENS precursors. Our results support a model in which Ret contributes to multiple distinct cellular phenotypes associated with the proper development of the ENS, including the

specification of inhibitory neuron subtypes, cell cycle dynamics of ENS progenitors, and the

developmental timing of neuronal and glial commitment.

Summary Statement

Ret LoF affects proper development of the mouse ENS through multiple distinct cellular

phenotypes including restriction of neuronal fate potential, disruption of ENCDC migration, and

modulation of progenitor proliferation rate.

Introduction

The enteric nervous system (ENS) is primarily derived from migrating vagal and sacral neural

crest cells (Kulkarni et al. 2020; Le Douarin and Smith 1988; Kulkarni et al. 2018; Durbec et al. 1996;

Kapur et al. 1992) that colonize the developing gut tube between embryonic day (E)9.5 to E14.5 in mice

(Young et al. 1998) and Carnegie stage (CS) 14 to CS22 in humans (Goldstein et al. 2013), leading to

the neuronal networks of the myenteric and submucosal plexuses of the gut. The ENS has essential roles

in propulsive and mixing motility, absorption, immunity, and vascular function (Furness 2012) and

disruption of proper ENS development or organization underlies several disorders of the gastrointestinal

system (Heuckeroth 2018). Previous studies of ENS development have demonstrated that invading

enteric neural crest-derived cells (ENCDCs) are predominantly neuronal precursors early in mouse

development (E10.5), but as development proceeds they differentiate into either glia, expressing classical

glial markers including *Gfap, Plp1,* and *S100b* (Hao and Young 2009; Kang et al. 2021), or neurons (Hao

and Young 2009). Single-cell transcriptional profiling of mouse ENCDCs during mid and late (E12.5,

E15.5 and E18.5) developmental time points reflects this heterogeneity and temporal change in cell fate

(Lasrado et al. 2017; Morarach et al. 2021, 2017; Elmentaite et al. 2021).

One of the key factors involved in the migration of ENCDCs into the developing gut, and their

differentiation into enteric neurons and glia, is the receptor tyrosine kinase RET (Natarajan et al. 2002;

Lasrado et al. 2017). RET expression in ENCDCs promotes proliferation and cell survival via the MAPK

and PI3K/AKT pathways (Mulligan 2014). Consequently, gain-of-function (GoF) mutations in *RET* have

been associated with various cancers including medullary thyroid carcinoma and multiple endocrine

neoplasias type 2A and 2B (Chatterjee et al. 2016b). Conversely, *RET* LoF due to both coding and non-

coding regulatory variants leads to Hirschsprung disease (HSCR; congenital intestinal aganglionosis), an

uncommon (~1/5,000 live births) developmental disorder of the ENS characterized by absence of enteric

ganglia along variable lengths of the distal bowel (Tilghman et al. 2019; Chatterjee et al. 2016a; Kapoor

et al. 2015; Chatterjee et al. 2021; Alves et al. 2013). In the developing mouse gut, *Ret* deficiency leads

to significant transcriptional changes in transcription factors, signaling molecules, and genes involved in

specific transport and biosynthesis processes, leading to aganglionosis (Heanue and Pachnis 2006;

Chatterjee et al. 2019). Tissue-level bulk transcriptomic studies in *Ret*-deficient developing mouse gut

(Heanue and Pachnis 2006; Chatterjee et al. 2019)) have uncovered significant expression changes in

multiple transcription factors critical to neural crest migration and ENS differentiation including *Sox10,*

Pax3 and *Phox2b* (Southard-Smith et al. 1998; Lang et al. 2000; Pattyn et al. 1999), as well as genes

harboring mutations in HSCR patients such as *Ednrb* and *Gdnf* (Heanue and Pachnis 2006; Chatterjee

et al. 2019).

Development and fate specification in the ENS occurs in conjunction with a prominent migration

along the forming gut tube (Kang et al. 2021; Nagy and Goldstein 2017). As such, temporal and spatial

influences affect ENCDC competence and can modulate fate decisions during development (Kang et al.

2021; Obermayr et al. 2013). While *Ret* is well known to play a significant role in the proper establishment

of the ENS, the cell-type-specific effects of *Ret* LoF on normal ENS development have not yet been fully

elucidated and could contribute to our understanding of normal ENS development.

To identify the cell-type-specific contributions of *Ret* to the proper organization and development

of the ENS, we examined the cellular heterogeneity of the ENS during mouse gut development in the

presence and absence of *Ret*. We profiled the transcriptome of 1,003 enteric neural crest-derived single

cells in heterozygous *Ret* null and prenatally lethal homozygous *Ret* null mice at E12.5 and E14.5. The

Ret heterozygous mice do not exhibit aganglionosis in the developing gut or any other known phenotypes

associated with *Ret* LOF and hence serves as an suitable control for our study which focusses on the

cellular dysregulation of enteric neurons and glia; the primary cell affected cell type due to *Ret* LoF.

ENCDCs and their progeny were isolated using a cyan fluorescent protein (CFP) reporter driven by the

Ret promoter for enrichment (Uesaka et al. 2008). Using a combination of differential gene expression

analysis, gene co-expression pattern analysis, RNA velocity, and cell cycle state predictions, we identify

several key developmental phenotypes associated with the loss of *Ret.* We find the specific effects of

Ret LoF vary across different cellular states in ENCDCs including effects on progenitor proliferation,

migration, differentiation, and fate specification. We confirm a previously described bifurcation in enteric

neuronal developmental fate specification (Morarach et al. 2021; Elmentaite et al. 2021) at an even earlier

time point. We further demonstrate that *Ret* expression is required for commitment to one of these two

neuronal fates and subsequently, the establishment of cell types from a predominantly inhibitory neuron

developmental trajectory. Additionally, we show that homozygous *Ret* LoF leads to precocious

differentiation of ENCDCs and a significant reduction in ENCDC progenitor proliferation rate. Each of

these observed phenotypes may independently contribute to the deficits in proper ENS cell numbers,

development, organization, and function observed in HSCR. These results highlight the diverse cell-type-

specific roles for *Ret*, a key developmental gene, in the proper establishment of the ENS with implications

for understanding HSCR-related phenotypes, and provide insights into the unique nature of ENS

development.

Results

To assess cell-type-specific roles for *Ret* in the developing ENS, we used a previously described

mouse line harboring a *Ret*-null fluorescent reporter allele (Ret^{CFP}) (Uesaka et al. 2008). All ENCDCs

express *Ret* during early development (Natarajan et al. 2002) allowing for their isolation via this reporter

system. We sorted *Ret* expressing ENCDC via FACS to decipher the cell autonomous effect of *Ret* LoF.

Ret-null ($Ret^{CFP/CFP}$) mouse embryos exhibit total colonic aganglionosis and kidney agenesis, and are

perinatal lethal (Uesaka et al. 2008). Importantly, no defects in ENS formation have been observed in

Ret-heterozygous ($Ret^{CFP/+}$) mice (Uesaka et al. 2008) nor any significant difference in the genome wide

expression observed when compared to wildtype GI tract at the same developmental times (Kapoor et

al. 2017). Comparison of $Ret^{CFP/CFP}$ to $Ret^{CFP/+}$ mice can therefore elucidate the mechanisms by which

Ret LoF affects the proper formation of the ENS.

To investigate transcriptional changes of ENCDCs, we performed single-cell RNA-sequencing

(scRNA-seq) on FACS-enriched populations of ENCDCs in the developing murine gut. Briefly, CFP-

positive single cells were collected from the dissociated stomach and intestine of male and female E12.5

and E14.5 $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ embryos and subjected to a modified Smart-seq2 scRNA-Seq library

preparation (Picelli et al. 2014; Chevée et al. 2018) (see Methods). Libraries from 95 cells for each of two

biological replicates were sequenced from all age, sex, and genotype combinations, with the exception

of E14.5 $Ret^{CFP/CFP}$ embryos. Consistent with a previously described reduction in the number of CFP[+]

cells seen in a *Ret* hypomorphic mouse model (Uesaka et al. 2008), we observed reduction in the cell

capture rate in the E14.5 $Ret^{CFP/CFP}$ mice as compared to other ages and genotypes. We also observed

a pronounced sex bias in the number of CFP[+] cells recovered from E14.5 $Ret^{CFP/CFP}$ embryos, with fewer

viable cells collected from males, consistent with the established sex bias observed in HSCR (Amiel et

al. 2008) and described in other mouse models of HSCR (Dang et al. 2011; Uesaka et al. 2008). This

reduced number of $Ret^{CFP/CFP}$ cells as a function of developmental age and sex, highlights a specific

challenge of this mouse model as the number of viable CFP[+] cells rapidly diminishes in these conditions.

As such, only 22 cells from one male E14.5 $Ret^{CFP/CFP}$ embryo and 190 cells from four female E14.5

$Ret^{CFP/CFP}$ embryos were recovered (Table 1). Full-length cDNA scRNA-Seq libraries from 1,341

individual cells were sequenced to an average depth of 9.91×10^5 (standard deviation 7.43×10^5) paired-

end reads per cell. Low quality cells were removed using standard metrics (see Methods) leaving 1,003

cells meeting all quality control parameters (Figure 1A; Table 1) with an average read depth of 1.16×10^6

(standard deviation 6.83×10^5) paired-end reads per cell. Relative FPKM values were converted to

estimated RNA copies per cell (CPC) using the CENSUS algorithm (Qiu et al. 2017). We identified a set

of 14,319 expressed genes with greater than zero CPC in at least 20 cells and a standard deviation

greater than zero across all cells. To visualize the transcriptional relationships between cells, we identified

a set of 419 high-variance genes to use as input to the uniform manifold approximation and projection

(UMAP) dimensionality reduction tool to establish a biologically relevant two-dimensional embedding of

the cells (Figure 1B, Figure S1).

Annotation of cell types and states of the developing ENS

To quantify cell type and cell state differences with respect to genotype, we first used a Gaussian-

based clustering algorithm (Scrucca et al. 2016) to identify 9 clusters (Figure 1C) and calculated gene-

wise specificity scores to identify gene expression profiles for each cluster (Figure S2A). High specificity

scores relative to mean expression identified clusters of actively cycling progenitors (cluster 2, Hist1h

family genes; cluster 3, *Aurka/b*, *Cdk1*), a neurogenic commitment window (cluster 4, *Neurod4*; cluster

5, *Pcp2, Hes5*, cluster 6, *Hes6*, *Btg2*), maturing neurons (cluster 7, *Etv1*, *Vip*, *Gal, Cartpt*; cluster 8,

Slc18a3 (ChAT), Npy, Snap25, Gfra2; cluster 9, *Th, Dbh, Ddc*), as well as a population of maturing enteric

glia (cluster 1, *Plp1*, *Myl9*) (Figure S2B, Table 2). The neural progenitor cell markers p75 (*Ngfr*) and the

pan neuronal marker TuJ1 (*Tubb3*), which have traditionally been used to label mature enteric neurons

along with *Ret* (Hao and Young 2009; Young et al. 1999), have a distinct, largely non-overlapping

expression profile in *Ret*-expressing cells at these time points (Figure S2C). Consistent with this, we

observe *Ngfr* and *Tubb3* expression to be restricted to non-overlapping populations of cells at either end

of our learned manifold.

To annotate individual cells as either progenitors, developing glia, or developing neurons we next

evaluated the use of canonical marker gene expression for mature ENS cell types. However, during

development, the majority of cells may not yet express markers of mature, terminally differentiated cell

types. To address this limitation, we chose to assign cell types using a combination of transfer learning

and marker gene expression, where available, to predict the most likely annotation probabilities based

on an existing cellular atlas of adult mouse enteric neurons and glia (Zeisel et al. 2018). To obtain

transcriptional signatures of mature ENS cell types, we learned patterns of gene co-expression on the

annotated set of enteric neurons and glia from the public dataset using single-cell Coordinated Gene

Activity in Pattern Sets (scCoGAPS) (Stein-O'Brien et al. 2019) and trained a random forest (RF) model

on those learned patterns to predict cell type (see Methods). We next projected our single-cell data into

the latent spaces defined by these patterns using projectR (Stein-O'Brien et al. 2019; Sharma et al. 2019)

(Figure S3) and employed the RF model to classify cells in our dataset based on their projected pattern

weights, thereby agnostically annotating the cells in our data as glia or neuron (Figure S4A-C). We next

calculated cell type proportions by cluster (Figure S4D) and for clusters in which at least 90% of cells

shared an annotation, we transferred this majority annotation to all members of a given cluster. As a

result, clusters 1-5 (1 = 99% glia, 2 = 98%, 3 = 92%, 4 = 95%, 5 = 94%) were annotated as glia and

clusters 7 and 8 (7 = 98% neuron, 8 = 90%) were annotated as neurons. Annotations for clusters 6 (58%

glia) and 9 (63% neuron) were not altered.

Our single-cell gene expression profiling is a cross-sectional snapshot in developmental

progression for a population of cells that have not yet fully committed to mature cell type identities. To

infer the direction of state transitions for our ENCDC population, we next calculated RNA velocity

estimates (La Manno et al. 2018) and observed that the learned velocity fields comprise diverging

trajectories from a common origin (Figure 1F). Cells with high projection weights in patterns associated

with proliferating ENCDCs (pattern 40: *Brca1/2, Cdc6, Pola1/2*; pattern 41: *Ccna1/b1, Cenpe, Kif24*;

pattern 42: *Aurka/b, Bub1, Top2a*; Figure S3, Tables 3, 4, and 5) exhibit an ellipsoid pattern of velocities,

consistent with transition through the cell cycle. Extending from this progenitor population we observed

an population of cells expressing canonical markers consistent with progression towards a neuronal fate

(cluster 6: *Btg2*, *Hes6* Figure S2B) that further branches into two distinct trajectories of neuronal

specification (Figure 1F), consistent with previous observations of postmitotic branching of cellular fate

specification for neural crest-derived neurons in later gut development (Morarach et al. 2021). We infer

through shared marker gene expression that cluster 7 in our data corresponds to a previously described

developmental branch (branch A) (*Etv1*[+], *Nos1*[+], *Vip*[+], *Gal*[+]) and cluster 8 corresponds to branch B

(*Bnc2*[+], *Dlx5*[+], *Mgat4c*[+], *Ndufa4l2*[+]) (Figure S6A). These data highlight that *Ret*-expressing cells isolated

from the GI tract encompass two broad fates during development (glial-commitment and neuronal-

commitment) and confirm that neuronally-committed ENCDCs diverge post-mitotically into two distinct

developmental trajectories, outlining the diversity of cellular fate trajectories accessible to ENCDCs within

the developing mouse gut.

Ret LoF induces distinct transcriptional responses in different ENS cell states

We first asked whether loss of *Ret* changes the proportion of ENCDC subtypes in our dataset. To

assess this, we tested for genotype bias in cluster composition by comparing the proportion of Ret$^{CFP/CFP}$

cells against the naïve (overall) proportion of 34%. Using a chi-squared test for equal proportions, we

observed significant deviations in seven of the nine identified clusters after Bonferroni correction for

multiple tests: clusters 1, 8, and 9 were enriched for *Ret$^{CFP/CFP}$* cells (cluster 1: 83%, Bonferroni adjusted

p=9.86x10^{-18}; cluster 8: 57%, p=5.64x10^{-4}; cluster 9: 97%, p=1.24x10^{-11}), clusters 4-7 were depleted for

Ret$^{CFP/CFP}$ cells (cluster 4: 9%, Bonferroni adjusted p=3.48x10^{-5}; cluster 5: 20%, p=1.00x10^{-3}; cluster 6:

33%, p=3.44x10^{-3}; cluster 7: 2%, p=6.33x10^{-5}), while clusters 2 and 3 were relatively invariant (cluster 2:

42% *Ret$^{CFP/CFP}$*, Bonferroni adjusted p=0.196; cluster 3: 35%, p=1.0) (Figure 1D). Clusters 6, 7, and 8 are

neuronal-committed clusters and differ significantly from the naive proportion. Clusters 6 and 7 are

enriched for *Ret$^{CFP/+}$* cells and 8 is enriched for *Ret$^{CFP/CFP}$* cells, indicating strong genotypic effects

associated with neuronal fate specification. Three glial clusters, clusters 1, 4, and 5, differ significantly

from the naive proportion as well, with cluster 1 enriched and clusters 4 and 5 depleted for *Ret$^{CFP/CFP}$*

cells. Cluster-specific enrichment of genotype within the neuronal and glial populations suggests that

genotypic effects of *Ret* LoF modulates cell fate commitment with divergent results across ENCDC cell

states.

To identify the transcriptional effects of *Ret* LoF on the developing ENS, we tested 14,319 genes

and identified 520 as differentially expressed with respect to genotype across all cells (monocle likelihood

ratio test; Benjamini-Hochberg corrected q-value < 0.05, Figure 2A, Table 6A). Hierarchical clustering

based on expression of these 520 genes revealed that cells cluster first by cell type then by genotype,

again suggesting that the transcriptional changes observed are largely cell-type-specific. Gene ontology

analysis revealed enrichment for several relevant processes, including gliogenesis, cell fate commitment,

and axon development (hypergeometric test q-value < 0.05, Figure 2B, Table 7A). Significantly

upregulated genes expressed in $Ret^{CFP/CFP}$ progenitor/glia includes several associated with ENS glial

maturation (*Fabp7*, *Plp1*, *Myl9*), indicating that these progenitor/glia cells may be on average more

mature than their $Ret^{CFP/+}$ counterparts. A second cluster of genes is more highly expressed in $Ret^{CFP/+}$

neurons and includes genes with specific expression in the neurogenic branch A (*Cartpt, Vip, Nos1,*

Etv1), consistent with our observation that $Ret^{CFP/+}$ cells are significantly overrepresented in branch A

(cluster 7).

We next sought to map the transcriptional signatures learned from our ENCDC dataset to specific

cellular features by using coordinated sets of co-regulated genes. To this end, we applied scCoGAPS to

learn 25 patterns of gene co-expression from our ENCDC scRNA-Seq dataset (Figure S5, Table 8). We

calculated the Pearson correlation for pattern weights by cell against both known and learned cellular

annotations (Figure 2C, Table 9). This revealed patterns that correlated with technical (read depth and

alignment rates), experimental (genotype, age, and sex), as well as biological (cell type and cluster)

variables. We identified patterns correlated with the acquisition of different neuronal identities (Pattern

20: *Elavl4* (*HuD*)*, Nrxn3, Tubb3, Nnat, Syt1*; Pattern 21: *Nos1, Tubb2a, Mapt, Sncg, Ntng1*; Pattern 22:

Onecut2, Gabrg2, Gabra3, Slc18a3 (*ChAT*)*, Gfra2, Elavl2* (*HuB*)*, Elavl3* (*HuC*)) (Figure S5; Table 10), a

transient neurogenic window (Pattern 19: *Pou4f1, Gadd45g, Neurod4, Hes6, Btg1/2, Sox11, Phox2b*),

as well as glial identity (Pattern 12: *S100b, Plp1, Fabp7*). In addition, we also identified patterns well

correlated with specific genotype-biased clusters of cells including Pattern 18 (*Ret, Gal, Ass1, Etv1*) and

Pattern 21 specifically used by Branch A neurons in cluster 7, and Pattern 12, specifically used by cells

in cluster 1 undergoing glial maturation and significantly enriched for $Ret^{CFP/CFP}$ cells.

Previous in vivo studies in mice have shown a severe migratory delay beginning as early as E9.5

in the absence of *Ret (Natarajan et al. 2002; Uesaka et al. 2008, 2015)*. Consistent with this observation,

we identified four gene expression patterns (Patterns 2, 18, 19, and 25), well correlated with genotype,

where *Hox* gene family members were identified as pattern markers (Stein-O'Brien et al. 2017) (Table

10). Pattern 25 markers include the more distal *Hoxa/c9* and *Hoxa/c10*, and the separation of these cells

from the main manifold of ENCDCs, combined with the specific expression of marker genes such as

Maob and *Pcdh10*, suggests that the handful of cells specifically using Pattern 25 are likely caudal

Schwann cell precursors. The enrichment for Hox genes within the remaining three pattern marker gene

lists with strong correlation to genotype, suggests that Hox gene expression in ENCDCs may be a

discriminating feature between the two genotypes. This was confirmed by the identification of seven *Hox*

genes, (*Hoxa2*, *b2*, *d3*, and the more caudally-expressed *Hoxa4*, *a5*, *b5*, and *b6*), as well as several

anterior-posterior pattern specification genes (*Dll1, Gpc3, Hes1, Lef1, Nrarp, Sema3c,* and *Tbx3*),

significantly differentially expressed (monocle likelihood ratio test; q<0.05) with respect to genotype

(Figure 2A-B). With the exception of *Gpc3, Lef1,* and *Sema3c*, all of the differentially expressed genes

associated with anterior-posterior positioning show increased expression in $Ret^{CFP/+}$ cells (Figure 2E).

Consistent with this and other previous work at E12.5 (Hirst et al. 2018), we observe that ENCDCs in the

absence of *Ret* have not left the extrinsically innervating nerve fibers of the stomach by E12.5 nor by

E14.5 (Figure 2D), in stark contrast to $Ret^{CFP/+}$ cells which have formed a dense network throughout the

stomach and all but the end of the hindgut by E12.5. As the *Hox* gene family is well-known to contribute

to axial patterning, this differential expression of *Hox* genes may therefore be a consequence of

significant rostro-caudal positional differences in ENCDCs in the presence and absence of *Ret*.

Ret is required for commitment to the branch A neuronal developmental trajectory

Recent work in E15.5 mouse gut shows that after neuronal commitment, cells of the developing

ENS branch into two main trajectories, referred here as branch A (*Etv1+, Gal+, Nos1+*, *Vip*+) and branch

B (*Bnc2*+, *Dlx5*+, *Mgat4c*+, *Ndufa4l2*+) (Morarach et al. 2021). Our UMAP embedding reveals this

branching behavior can be observed as early as E12.5 and that the choice of branch is *Ret*-dependent

(Figure 3A, Figure S6A). As observed above, cluster 7, corresponding to branch A, is almost exclusively

composed of $Ret^{CFP/+}$ cells. Conversely, while cluster 8 (branch B) is significantly enriched for $Ret^{CFP/CFP}$

cells (Figure 1D), both genotypes are represented along the branch B trajectory. This suggests that there

is a significant bias in the choice of developmental trajectory mediated in part by *Ret* expression in

ENCDCs.

We next sought to identify branch-specific differences in gene expression with respect to

genotype. When we tested cluster 8 (branch B), we identified 12 genes differentially expressed with

respect to genotype (monocle likelihood ratio test, q<0.05, Figure 3B, Table 6B), four of which are more

highly expressed in $Ret^{CFP/+}$ and eight more highly expressed in $Ret^{CFP/CFP}$. Gene ontology analysis

indicates that genes upregulated in $Ret^{CFP/CFP}$ are enriched (hypergeometric test, q<0.05, Table 7B) for

dopamine biosynthetic processes (q<4.52x10^{-6}, *Agtr2*, *Ddc*, *Th*), catecholamine biosynthetic processes

(q<6.44x10^{-6}, *Agtr2*, *Ddc*, *Th*), and regulation of corticosteroid hormone secretion (q<3.35x10^{-4}, *Agtr2*,

Tac1). This suggests that, while both genotypes exhibit generally similar transcriptional profiles within

branch B, there may still be functional consequences of *Ret* LoF within this trajectory.

To examine differences between branch A and B trajectories and identify branch-specific markers,

we tested for differential expression irrespective of genotype to identify 111 genes (monocle likelihood

ratio test, q<0.05, Figure 3C, Table 6C). Gene ontology enrichment analysis (Table 7C) reveals that

genes with higher expression in cluster 7 are enriched for regulation of neurological system processes

(hypergeometric test, q=1.91x10^{-5}, *Cartpt, Ednrb, Fgfr1, Nos1, Nrg1, Vip*) and axonogenesis

(hypergeometric test, q=6.11x10^{-5}, *Ablim1, Auts2, Cdh11, Etv1, Lrrc4c, Lrrn3, Nrg1, Ntng1, Ret*), while

genes more highly expressed in cluster 8 are enriched for catecholamine biosynthetic processes

(hypergeometric test, q=1.15x10^{-8}, *Agtr2, Dbh, Ddc, Gata3, Gch1, Th*), regulation of neurotransmitter

levels (hypergeometric test, q=2.00x10^{-6}, *Chrna7, Dbh*, *Ddc, Gch1, Nrxn1, Prima1, Slc18a1, Sytl4, Th*),

and dopamine biosynthetic processes (hypergeometric test, q=2.36x10^{-6}, *Agtr2, Dbh*, *Ddc, Gch1, Th*).

The increased expression of genes in this latter geneset in $Ret^{CFP/CFP}$ cells in cluster 8 relative to $Ret^{CFP/+}$

cells (Figure 3B) suggests that differentiation and/or maturation rate of cells in branch B may be affected

by loss of *Ret*. Additionally, several established markers of branches A and B (Morarach et al. 2021) are

differentially expressed in accordance with expectations (*Etv1, Gal, Nos1,* and *Vip are* higher in cluster

7/branch A; *Bnc2, Dlx5,* and *Ndufa4l2* are higher in cluster 8/branch B).

Both Ret and CFP are downregulated along the branch B trajectory in addition to Gfra1 (Figure

3D), which encodes the canonical RET co-receptor, and Ednrb, which is known to be epistatic with Ret

(McCallion et al. 2003; Chatterjee and Chakravarti 2019; Carrasquillo et al. 2002). Interestingly,

expression of these same genes is maintained in Ret$^{CFP/+}$ cells along the branch A trajectory, and each

exhibits a biased expression for cells along this trajectory (Figure 3D). The exclusion of Ret$^{CFP/CFP}$ cells,

and the branch-A-specific persistence of Ret and its canonical co-receptor indicates that commitment to

the branch A developmental trajectory, and subsequently, establishment of the enteric neuronal subtypes

derived from cells adopting this trajectory, requires continued Ret expression. The downregulation of both

Ret and CFP expression along the branch B trajectory, which contains numerous cells of both genotypes

(Figure 3A&D), suggests that this trajectory may not require persistent Ret expression. While prior

immunohistochemical studies found Ret protein expression is down-regulated in emerging glia and

maintained in emerging neurons (Young et al., 1999), our scRNA-Seq data indicate that Ret exhibits

differential expression between the two branches of differentiating neurons with Ret expression

maintained in branch A and down-regulated in branch B, consistent with other recent single cell studies

(Elmentaite et al., 2021; Morarach et al., 2021). In aggregate, these results suggest that in the Ret$^{CFP/CFP}$

developing ENS, since branch A is an inaccessible developmental trajectory, and a greater portion of

neuronally committed ENCDCs may engage the branch B trajectory.

To validate Branch A inaccessibility in *Ret$^{CFP/CFP}$* ENCDCs, we selected *Cartpt* (branch A/cluster

7) and *Gfra2* (branch B/cluster 8) from the list of differentially expressed genes as markers for *in vivo*

validations due to their high expression and high specificity in each branch (Figure 3D). We performed

whole-mount immunohistochemistry (IHC) on isolated gut (esophagus, stomach, and intestine) of E12.5

and E14.5, *Ret$^{CFP/+}$* and *Ret$^{CFP/CFP}$* embryos to identify and localize cells from each branch. We probed

for CART, the product of *Cartpt*, and CFP to localize branch A cells. In the *Ret$^{CFP/+}$* samples at E12.5 we

observe co-expression of CART and CFP in the stomach (Figure 3E) but not in the esophagus (data not

shown). By E14.5, we observe co-expression of CART and CFP extending all the way to the hindgut of

Ret$^{CFP/+}$ samples (Figure 3F). Consistent with our scRNA-seq data, we do not observe any CART

expression in the stomach of $Ret^{CFP/CFP}$ samples at either age (Figure 3E-F) nor in the esophagus (data

not shown). We further validated the lack of branch A cells within the intestine of $Ret^{CFP/CFP}$ cells using

single-molecule fluorescent in situ hybridization (RNAscope) to assess mRNA expression of the branch

A marker genes *Cartpt* and *Gal* (Figure S6B-C). These results demonstrate that cells associated with the

branch A developmental trajectory are not produced within the GI tract in the absence of *Ret*.

To determine the spatial distribution of branch B cells we exploited the specific expression of

Gfra2, which encodes a non-canonical RET co-receptor, in branch B neurons, which is correlated with a

decrease in expression of *Gfra1* (Figure 3D). When we probed for GFRA2 and CFP via IHC we did not

observe GFRA2 expression in the E12.5 stomach of $Ret^{CFP/+}$ samples (Figure 3G). We did, however,

observe co-expression of GFRA2 and CFP restricted to cells along the extrinsically innervating nerve

fibers of the stomach of $Ret^{CFP/CFP}$ mice at this age (Figure 3G), confirming that branch B neuronal

development trajectory is indeed accessible in the absence of *Ret*. At E12.5 we did not observe GFRA2

outside the extrinsically innervating nerve fibers of the $Ret^{CFP/CFP}$ stomach, nor in the intestines of either

genotype (data not shown), consistent with the previously described migratory defect and failure to

effectively colonize the ENS in the $Ret^{CFP/CFP}$ mice. We were able to detect CFP$^+$/GFRA2$^+$ cells (putative

branch B neurons) in the esophagus of both genotypes, as well as CFP$^+$/GFRA2$^-$ (non-neuronal

ENCDCs) and CFP$^-$/GFRA2$^+$ (maturing branch B neurons) cells (Figure S6D), indicating that branch B

cells are spatially restricted to the esophagus in E12.5 $Ret^{CFP/+}$ samples and to the esophagus and

extrinsic nerve fibers along the stomach in E12.5 $Ret^{CFP/CFP}$. By E14.5, CFP$^+$/GFRA2$^+$ branch B maturing

neurons are observed throughout the intestine in the $Ret^{CFP/+}$; however, CFP$^+$/GFRA2$^+$ cells remain

restricted to the extrinsically innervating nerve fibers along the stomach in the $Ret^{CFP/CFP}$ mice (Figure

3H). Consistent with this difference, at E12.5 we see proportionally more $Ret^{CFP/CFP}$ cells in cluster 8

(branch B) than $Ret^{CFP/+}$ within our scRNA-Seq dataset (9/37 $Ret^{CFP/+}$, 28/37 $Ret^{CFP/CFP}$). By E14.5

however, $Ret^{CFP/+}$ are the predominant cells in cluster 8 (21/33 $Ret^{CFP/+}$, 12/33 $Ret^{CFP/CFP}$). This could be

related to an overall reduced number of E14.5 $Ret^{CFP/CFP}$ cells in the intestine, or reduced capture

efficiency as fluorescence decreases over biological time along this trajectory due to the downregulation

of *Ret (*and by proxy *CFP*) expression as cells progress along branch B (Figure 3D).

These data indicate that branch A and branch B represent two transcriptionally distinct

developmental trajectories of maturing ENS neuronal subtypes that colonize the length of the intestines

in the $Ret^{CFP/+}$ mice. Branch A neurons are never produced in the $Ret^{CFP/CFP}$ mice. Furthermore, Branch

B neurons are produced, at a potentially faster developmental rate, but remain restricted to the

extrinsically innervating nerve fibers throughout development in the $Ret^{CFP/CFP}$ mouse. As *Ret* is required

for cells to migrate into the developing ENS (Figure 2D) (Hirst et al. 2018) it follows that branch A, which

appears to require *Ret*, would be inaccessible while branch B, which is *Ret*-independent, remains

accessible in the absence of *Ret.*

HSCR-associated genes are expressed across different cellular states in the developing ENS

While coding and enhancer LoF variants in *RET* contribute to ~50% of cases of Hirschsprung

disease, HSCR is also associated with rare coding pathogenic alleles (PAs) in 23 additional genes

(*GDNF, GFRA1, NRTN, SOX10, EDNRB, EDN3, ECE1, ZEB2, PHOX2B, KIFBP, TCF4, L1CAM,*

IKBKAP, SEMA3C, SEMA3D, NRG1, ACSS2, FAM213A, ADAMTS17, ENO3, SH3PXD2A, SLC27A4

and *UBR4*) (Tilghman et al. 2019; Chatterjee et al. 2016a; Kapoor et al. 2015; Chatterjee et al. 2021).

Many of these genes are a part of the *RET-EDNRB* gene regulatory network (GRN) (Chatterjee et al.

2016a). To assess the distribution of HSCR-associated genes, we investigated the expression pattern of

all 24 genes in our data to determine their co-expression patterns and their specific effects due to *Ret*

LoF (Figure 4A). Three HSCR-associated genes (*Gdnf, Edn3, and Nrtn*) are not expressed in our ENCDC

dataset as they are predominantly expressed from the surrounding gut mesenchyme. Of the remaining

21 expressed genes (Figure S7A-C), six are significantly differentially expressed (monocle likelihood ratio

test, q<0.1, Figure 4B-C, Table 6A) between $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ cells (*Gfra1, Nrg1, Phox2b, Ednrb,*

Sema3c, and *Sox10)*. To identify cell-type-specific differential effects from *Ret* LoF we further tested the

combinatorial effect of genotype and cell type. Three HSCR-associated genes (*Gfra1, Sema3d* and

Ednrb) exhibit significant cell-type-specific differential expression (monocle likelihood ratio test, q<0.05,

Figure 4B-C, Table 6D). Both *Gfra1 and Ednrb* reach statistical significance in both tests, suggesting a

cell autonomous effect of *Ret* LoF on the expression of these two genes. Alternatively, four HSCR-

associated genes (*Nrg1, Phox2b, Sema3c*, and *Sox10*) are globally differentially expressed but are not

differentially expressed with respect to genotype:cell type interactions (monocle likelihood ratio test,

q<0.05; Table 6A,D).

We have previously shown that the proportion of cells by genotype deviates significantly from the

naïve proportion in seven of the nine clusters in our data (Figure 1D), we also observe that genotype

proportions deviate significantly from expectation in E14.5 neurons and approach statistical significance

in E14.5 glia (chi-squared test for equal proportions; neurons/p=0.0158, glia/p=0.0702) (Figure 4D).

Several of the HSCR-associated genes (*Ednrb, Gfra1, Elp1, Sox10, Zeb2*) are expressed predominantly

in progenitor/glia cells (Figure S7A,C). *Phox2b, Tcf4,* and *Ubr4* are the three within this set that are

ubiquitously expressed across all clusters in our embedding (Figure S7A,C). *L1cam* is enriched in both

maturing neurons and maturing glia, but exhibits reduced expression within the cycling progenitor

population of cells (Figure S7A,C). *Ece1* is predominantly expressed in the population of maturing glial

cells. These results suggest that some of the observed differential expression of HSCR-associated genes

when observed at a global level, but not when assessed within discrete cell states, may be primarily due

to differences in cell type proportions across genotypes. Finally, similar to the observed persistence of

Ret expression along Branch A, a subset of HSCR-associated genes are differentially expressed

(monocle likelihood ratio test; q<0.05, Table 6C) between the branches, with higher expression in branch

A relative to branch B (*Ednrb, Gfra1, Nrg1*, and *Ret*), suggesting their potential role in the establishment

of Branch A identity in conjunction with *Ret*.

Despite significant genetic heterogeneity, ~67% of HSCR patients have mutations in the genes

of the *RET-EDNRB* GRN (Tilghman et al. 2019). This GRN includes bidirectional transcriptional feedback

between *RET and EDNRB,* and the transcription factors *SOX10, GATA2, RARB,* and *NKX2-5*, as well

as *GFRA1* and its physiological ligand, *GDNF*, and the ubiquitin ligase *CBL*, which breaks down

phosphorylated RET (Chatterjee and Chakravarti 2019; Chatterjee et al. 2016a, 2019). To ascertain

whether this feedback is cell autonomous or not we looked at the expression of these genes in our

dataset. *Nkx2-5* and *Gdnf* are not expressed in our ENCDC dataset. Of the remaining six genes, four

(*Gata2, Gfra1, Rarb, and Sox10*) are significantly differentially expressed between $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$

ENCDCs (Table 6A), with *Gfra1* and *Rarb* more highly expressed in $Ret^{CFP/+}$ cells and *Sox10 and Gata2*

more highly expressed in $Ret^{CFP/CFP}$ cells*,* although these moderate differences in expression may also

reflect a disproportionate distribution of ENS cell types within our dataset with respect to genotype (Figure

4C-D, Figure S7C). Finally, with the exception of *Sox10*, which rapidly decreases expression during

neuronal commitment, and *Gata2*, which is specific to cluster 9, *Ednrb, Rarb, Gfra1*, and *Cbl* all exhibit

persistent expression along the branch A neuronal developmental trajectory (Figure 4B, Figure S7C),

consistent with a role for the *RET-EDNRB* GRN in establishing branch A identity. The expression of

Sox10 exclusively within the progenitor population explains a previous observation that *Sox10*

haploinsufficiency only affects maintenance of progenitor cells and not neurons (Paratore et al. 2002).

These results highlight the diverse cellular states in which HSCR-associated genes are

expressed, suggesting that their roles in HSCR disease etiopathology may involve modulation of different

aspects of ENCDC proliferation and development. Collectively however, expression of *RET-EDNRB*

GRN genes is affected by the loss of Ret signalling and is persistent along the Branch A trajectory, which

is inaccessible in the $Ret^{CFP/CFP}$ mouse.

Ret LoF reduces the number of actively proliferating ENCDCs in the developing gut

Reduction in proliferation capacity and precocious differentiation as a result of LoF of other genes

within the developing ENS have been proposed as significant drivers of colonic aganglionosis (Fujiwara

et al. 2018; Bergeron et al. 2016). Given the previously established role for *RET* as a proto-oncogene

(Grieco et al. 1990; Mulligan 2014) and its association with increased proliferation in other context (Park

and Bolton 2017; Perea et al. 2017; Gattelli et al. 2013), we asked whether *Ret* LoF was associated with

significant differences in the proliferation rate of ENCDC progenitors. Differential gene expression

analysis with respect to both genotype and cell type identified 331 genes (monocle likelihood ratio test,

q<0.05; Figure S8A, Table 6D), consisting of several genes associated with the mitotic cell cycle (*Cdkn1c,*

Cdk1, Top2a, Ccnd1, Cenpk, Cenpe, Cenpf, Ccna2, Ccnb1, Ccnb2), DNA replication (*Pcna, Mcm3,*

Mcm5, Mcm6, Mcm10, Gmnn), and cell cycle regulation (*Rgcc, Mad2l1, Chek2, Rad51ap1*). Gene set

enrichment analysis of this differential gene list identified several cell cycle-associated gene sets

including mitotic cell cycle processes, cell division, DNA replication, regulation of cell cycle as top scoring

Gene Ontology Biological Processes (GO:BP) (hypergeometric test, q<0.05; Table 7D). To focus on cells

that are potentially engaging the cell cycle, we next performed a differential gene expression analysis on

just the subset of ENCDCs identified as progenitors or glia within the contiguous manifold (i.e. excluding

ENCDCs committed to a postmitotic neuronal fate and excluding cluster 9). We identified 364 genes with

significant differential expression with respect to genotype (monocle likelihood ratio test, q<0.05; Figure

5A, Table 6E), and again this list was enriched for gene sets associated with cell cycle regulation, in

addition to gliogenesis, cell fate commitment, and neural crest cell differentiation (hypergeometric test,

q<0.05; Figure 5B, Table 7E). Thus, both cell cycle occupancy and differentiation state, two processes

that are intrinsically linked during developmental fate specification, are disrupted in the *Ret$^{CFP/CFP}$*

progenitor population relative to *Ret$^{CFP/+}$*.

We next asked whether we could identify any significant differences in cell cycle state

assignments for ENCDCs in the absence of Ret. We used the tricycle package (Zheng et al. 2021) to

project the 1,003 single ENCDCs into the default reference cell cycle space provided (Figure 5C). This

analysis allows us to estimate the continuous position of each cell along the cell cycle (Θ) (Table 11),

comparisons of which by genotype and age (Figure 5D) suggests that at E12.5, there are minimal

differences in cell cycle occupancy between genotypes (Kolmogorov-Smirnov test; p-value = 0.660), but

by E14.5 there are consistently fewer *Ret$^{CFP/CFP}$* cells engaging the cell cycle than *Ret$^{CFP/+}$*(Kolmogorov-

Smirnov test; p-value = 8.28x10^{-3}).

To further assess whether *Ret$^{CFP/CFP}$* ENCDCs exhibit differential cell cycle occupancy relative to

Ret$^{CFP/+}$ we next sought to infer cell cycle stage annotations for cells in our dataset using a projection-

based approach with an independent, biologically relevant data source. Using the scCoGAPS patterns

previously learned from publicly available data, and on which we trained the random forest classification

model, we observed high usage of three patterns within a population of cycling glia. Gene ontology

analysis of pattern markers revealed these patterns correspond to different proliferative phases of the

cell cycle (hypergeometric test, q<0.05, Table 4A&B). Pattern 40 was associated with DNA replication,

corresponding to S phase; pattern 41 associated with cilium assembly, organization, and segregation,

and attachment of spindle microtubules to kinetochores, corresponding to prophase or metaphase (early

M phase); and pattern 42 associated with chromatid segregation and nuclear division, corresponding to

anaphase or telophase (late M phase). To predict the cell cycle stage for individual cells in our ENCDC

data, we scaled projected pattern weights linearly within each pattern to range between 0 and 1, and

used this as a measure of cell cycle stage occupancy probability (Figure S8C&D). We tested all six

possible comparisons of the four interaction conditions of genotype and age per pattern for differences

in probabilities using two-sample Kolmogorov-Smirnov tests on the empirical cumulative density functions

(Figure S8E), and used the Bonferroni method for multiple testing correction. Across the three patterns,

E14.5 cells have significantly lower proliferative cell cycle stage occupancy probabilities than E12.5 cells

of the same genotype (S phase: $Ret^{CFP/+}$ Bonferroni adjusted p≤1.32x10^{-15}, $Ret^{CFP/CFP}$ p≤2.00x10^{-15}; early

M: $Ret^{CFP/+}$ p≤1.32x10^{-15}, $Ret^{CFP/CFP}$ p≤3.84x10^{-13}; late M: $Ret^{CFP/+}$ p≤1.38x10^{-8}, $Ret^{CFP/CFP}$ p≤1.94x10^{-}

^{9}), consistent with the expectation that as progenitor cells differentiate, the proportion of actively cycling

cells will decrease over biological time. Again, cell cycle stage occupancy probabilities for ENCDCs did

not differ significantly with respect to genotype at E12.5. By E14.5 however, $Ret^{CFP/CFP}$ cells have

significantly lower occupancy scores across all three patterns (S phase: Bonferroni adjusted p≤9.66x10^{-}

^{5}, early M: p≤1.97x10^{-4}; late M: p≤0.0193) suggesting overall fewer cycling ENCDCs in the $Ret^{CFP/CFP}$.

This may be attributed to an increased differentiation rate, and consequently decreased proliferation rate,

in $Ret^{CFP/CFP}$ cells compared to $Ret^{CFP/+}$ cells between E12.5 and E14.5.

To validate this predicted decrease in cell cycle occupancy in $Ret^{CFP/CFP}$ ENCDCs, we next used

IHC to assess the number of CFP$^+$ ENCDC co-expressing the mitotic marker phosphohistone H3 (pH3)

within the developing gut at E14.5 (Figure 5E). We confirmed a significant (Student's t-test, $p<1.0\times10^{-4}$)

overall reduction in the number of CFP$^+$ cells in the $Ret^{CFP/CFP}$ mouse model at E14.5 (Figure S8G)

consistent with the reduced number of CFP$^+$ cells observed in our FACS-collected population (Figure 1A)

and previously published studies (Uesaka et al. 2008). We also identified a slight difference in the mean

number of pH3$^+$ cells at E14.5 (Student's T-test, $p<0.05$); however, modest enough to suggest that the

overall number of mitotic cells within the developing gut remains unaffected (Figure S8H). This is

consistent with a cell-autonomous effect of Ret on cell cycle as ENCDCs make up a small portion of all

proliferating cells in the developing gut and would therefore not contribute significantly to the overall

proliferative capacity of the gut as a whole. The overall number of double positive pH3$^+$/CFP$^+$ cells was

significantly reduced in $Ret^{CFP/CFP}$ animals (Figure 5F & S7I) (Student's T-test, $p=0.029$), and while not

statistically significant, we observed a difference in the mean percentage of pH3$^+$ cells within the subset

of CFP$^+$ ENCDCs in the $Ret^{CFP/CFP}$ ENS at E14.5 indicating that a smaller fraction of developing ENCDCs

are actively engaging the cell cycle in the absence of Ret (Figure 5G & Figure S8J) (Student's t-test,

$p=0.11$). In aggregate, these results are consistent with the previously established role of Ret in

enhancing proliferation and indicate that a loss of Ret in the developing ENS may contribute to an early

reduction in the progenitor pool, possibly through induction of precocious differentiation, leading to a

failure to colonize the more distal region of the developing gut.

Discussion

The enteric nervous system (ENS), initially derived entirely from the caudally migrating vagal and

cranially migrating sacral neural crest cells, contains more than 100 million neurons of a variety of

functionally distinct subtypes, highlighting the breadth of the functions it performs (Furness 2008; Rao

and Gershon 2016). In contrast to the rest of the peripheral nervous system, the ENS is endowed with

the ability to integrate multiple neuronal signals to control gastrointestinal (GI) behavior independently

from the brain and spinal cord (Furness 2008; Gershon 2010). One of the critical genes that enables this

transition of ENCDCs to diverse types of neurons is *RET*, which encodes a cell surface receptor tyrosine

kinase. *RET* LoF leads to colonic aganglionosis and is a known factor in the etiology of Hirschsprung

disease.

In spite of the critical role *Ret* plays in establishing the normal ENS, there is a limited understanding

of the diversity of *Ret*-expressing cells in the developing GI tract and the precise nature of the cell-type-

specific effects mediated by *Ret* LoF. By performing scRNA-seq on an enriched population of *Ret*-

expressing cells during ENS development we have generated a *Ret*-dependent fate transition map of

ENCDCs as they generate diverse classes of neurons and glia in the mouse GI tract to decipher the

specific cellular phenotypes mediated by *Ret* LoF.

Ret LoF affects the establishment of a specific developmental neuronal trajectory in the

developing GI tract

Our cellular reconstruction of ENCDC differentiation demonstrates that *Ret*-expressing cells

broadly acquire neuronal and glial fates early during normal development of the GI tract and helps explain

why *Ret* LoF leads to total aganglionosis caused by impaired migration, defects in proliferation, and loss

of immature ENCDCs (Schuchardt et al. 1994). Our data highlight that *Ret* is required for the induction of

an entire neuronal specification trajectory within the developing ENS which has previously been

associated with the specific development of at least two distinct subtypes of inhibitory motor neurons: an

interneuron subtype, and a population of intrinsic primary afferent neurons (IPANs) (Morarach et al. 2021).

Though previous studies have posited that branch A corresponds to a predominantly inhibitory motor

neuron (MN) trajectory and branch B corresponds to a predominantly excitatory MN trajectory (Morarach

et al. 2021), our data are derived from embryonic timepoints which are too early to directly support this

functional distinction between the branches. At E12.5, we observe putative branch B cells (GFRA2[+]/CFP[+])

in the esophagus of the *Ret*^CFP/+, and in the esophagus and stomach of the *Ret*^CFP/CFP mice (Figure 3F,

Figure S6C). We also observe putative branch A cells (CART⁺/CFP⁺) in the stomach of *Ret*^CFP/+ mice but

not the esophagus, and we do not observe putative branch A cells anywhere in the *Ret*^CFP/CFP GI tract at

E12.5. By E14.5 we observe putative branch B cells in the stomach of the *Ret*^CFP/CFP, and throughout the

gut up to and including the wave front of migration in *Ret*^CFP/+ mice. The earlier establishment of branch B

cells within the stomach of the *Ret*^CFP/CFP is consistent with a model of precocious differentiation we identify

in this study. The specific failure to produce cells along branch A in the *Ret*^CFP/CFP suggests that the subset

of mature ENS neuronal subtypes derived from this developmental trajectory may be the most acutely

affected in disorders such as HSCR and should be the focus of future studies of other *RET*-associated

pathologies in the gut.

We provide evidence that aganglionosis observed in HSCR may arise from a combination of early

exit from the cell cycle for ENS progenitors, leading to precocious differentiation, and a failure to produce

the full repertoire of mature ENS cell types. We do not detect any transcriptional signatures of cell death,

suggesting that at this early time point apoptosis or necrosis may not be a significant contributing factor

in *Ret*-deficient aganglionosis. The presence of glial cells in our mouse models, independent of genotype,

is consistent with previous observations in HSCR patients that immature glial cells can be detected within

the aganglionic portion of the colon (Tani et al. 2017). Our data suggest that *Ret* expression is not required

for the establishment of a normal ENS glial identity; however, the observed effects on proliferation and

differentiation potential suggest that enteric glia may be a preferred fate in *Ret* LoF models.

Neuronal fates that use non-canonical RET co-receptors remain accessible in the absence of Ret

A previous scRNA-seq study of the developing ENS at E15.5, when the migration of ENCDC

through the length of the gastrointestinal tract is near complete, reports a divergence in maturing neurons

that results in two branches of progenitors that produce different cell fates (Morarach et al. 2021). We

observe this divergence of the neuronal trajectories as early as E12.5, when ENCDCs are still migrating

and undergoing neuronal maturation. While we observe a near complete absence of *Ret*^CFP/CFP cells in

branch A (cluster 7), it is interesting to note that branch B (cluster 8) contains maturing neurons from both

genotypes and is not grossly affected by the loss of *Ret*. As cells progress along branch B there is a

general reduction in the expression of *Ret*, *CFP*, and several *RET-EDNRB* GRN genes, including *Gfra1*.

Concomitant with this, cells in branch B subsequently upregulate *Gfra2*, which encodes a non-canonical

RET co-receptor. Though GFRA1 and GFRA2 are in the same family of receptors, GFRA1 preferentially

binds GDNF as its ligand, while GFRA2 preferentially binds the ligand NRTN. It is possible that GFRA2

is binding to residual RET after downregulation of transcription; however, it raises the possibility that

GFRA2 complexes with another co-receptor to activate an alternate signaling pathway responding to a

different ligand in this *Ret*-independent trajectory. Differential expression of *Gfra* family genes within the

aganglionic segment of HSCR patients has previously been observed (Lui et al. 2002), including a

specific lack of *GFRA1* expression in neuronal cells within the associated hypertrophied nerve fibers.

Additionally, *Ntrn* and *Gfra2* null mice exhibit a reduction in neuron fiber density and neuronal size in the

ENS, and in each case, result in the specific loss of cholinergic excitatory myenteric innervation

(presumably branch B-derived) within the mouse gut, leaving the nitrergic neuronal population (branch

A-derived) largely unaffected (Rossi et al. 1999; Heuckeroth et al. 1999), consistent with the segregation

of fates and corresponding gene expression differences described here. In aggregate, these results

suggest that the canonical signaling pathway of RET, GDNF and GFRA1 is responsible for the proper

establishment of ENS subtypes derived from the branch A developmental trajectory, and that the activity

of NRTN-GFRA2 signaling may play a greater role in the establishment of ENS neuron identities along

branch B, potentially independent from *Ret*.

Expression patterns of HSCR-associated genes highlight necessity of disease-relevant cellular

context

The specific expression patterns of many genes that harbor pathogenic alleles in HSCR patients

within the invading and developing ENCDCs of the ENS should lead us to reassess the nature of

aganglionosis in HSCR. Our data highlight the need to better understand the specific cellular states and

contexts in which these genes and GRNs operate, as well as the cell autonomous versus cell non-

autonomous effects in HSCR. Genes such as *Acss2, Eno2,* and *Adamts17* express very lowly in the ENS

population (Figure 4), yet are robustly expressed in both the mouse and human developing GI tract

(Tilghman et al. 2019; Chatterjee et al. 2019). Furthermore, while the nature of our study, using only

sorted *Ret*-expressing cells is by definition only able to capture cell-autonomous effects, additional non-

cell-autonomous effects have been observed in *Ret* LoF studies as well (Bogni et al. 2008). Hence

patients carrying pathogenic alleles in the genes expressed in other, non-ENCDC, may contribute to

other GI anomalies, and aganglionosis may then be a secondary effect rather than primary. Though

verification of this would require detailed patient phenotyping, the shift between neuronal trajectories we

have identified suggests developmental mechanisms that may guide future analysis of patient samples.

Ret LoF leads to proliferation defects in ENS progenitors

During formation of the ENS, the gut itself is increasing dramatically in both size and length,

creating a huge proliferative burden on ENCDCs, the progenitors of the ENS. Our in silico estimates of

cell cycle stage occupancy in conjunction with an *in vivo* assay of proliferation suggest a decrease in

proliferation of ENCDCs in the absence of *Ret*. Even a small reduction in proliferative capacity could

produce a significant decrease in population size over the course of the several days it takes to colonize

the ENS, such as the absence of CFP+ cells we observe in the $Ret^{CFP/CFP}$ mouse intestines.

Ret deletion in the mouse leads to significant changes in fate acquisition, progenitor proliferation,

and ENCDC migration within the developing ENS. It is likely that *RET* deficiency alleles in humans lead

to similar phenotypes in distinct subsets of ENS cell types and states in HSCR patients, and that each of

these phenotypes may play some combinatorial role in establishing the aganglionosis that is the hallmark

of this disease. Furthermore, different *RET*-deficiency alleles and epistatic interactions may affect these

phenotypes in different ways, potentially explaining some of the observed variable expressivity and

incomplete penetrance of HSCR. Our study highlights the specific roles that *Ret* plays within different

cellular contexts of the developing enteric nervous system, and demonstrates that appropriate cellular

context is critical for the interpretation of multiple disease-related cellular phenotypes that may emerge

from a single causal insult.

Methods

Mouse strains:

Mice heterozygous for the conditional *Ret* allele (*Ret*^{fl/+})(Gould et al. 2008; Uesaka et al. 2008) were obtained from David Ginty (Harvard University, Boston, MA). Briefly, the floxed *Ret* allele (*Ret*^{fl}) was generated by inserting a gene cassette, comprising the floxed human *RET9* cDNA with SV40 intron polyA sequence followed by a cyan fluorescent protein (CFP) cDNA with polyA sequence, into exon 1 of the mouse *Ret* gene. *Ret CFP* knock-in mice (*Ret*^{CFP/+}; MGI:3777556) were generated by crossing *Ret*^{fl/+} mice to β-actin-Cre mice to remove the *RET9* cDNA This mouse was backcrossed >20 generation to C57BL/6 to maintain the line and all experiments were conducted in this isogenic strain. All mouse experiments were conducted in accordance with the National Institutes of Health Guidance for the Care and Use of Laboratory Animals. All procedures were approved by the Johns Hopkins Animal Care and Use Committee (Protocol number: MO21M56). All mice embryos used were either E12.5 or E14.5 and both male and female mice were used.

Mouse breeding and genotyping:

Homozygous *Ret* null mice are perinatal lethal (P0), therefore we intercrossed *Ret*^{CFP/+} mice to generate all possible genotypes. Embryos were genotyped from yolk sac genomic DNA to allow for genotype- and sex-specific analysis. Routine mouse genotyping was performed by PCR with primer pairs and amplicons as follows: wild-type *Ret* allele forward primer 5'-CAGCGCAGGTCTCTCATCAGTACCGCA-3' and reverse primer 5'-CAGCTAGCCGCAGCGACCCGGTTC-3' resulting in a 449 bp PCR product in *Ret*^{+/+} and *Ret*^{CFP/+} embryos (modified from (Jain and Varadarajan 2014)), CFP knock-in allele forward primer 5' ATGGTGAGCAAGGGCGAGGAGCTGTT-3' and reverse primer 5'-CTGGGTGCTCAGGTAGTGGTTGTC-3' resulting in a 615 bp PCR product from *Ret*^{CFP/+} and *Ret*^{CFP/CFP} embryos. Embryo sex was determined by PCR using yolk sac genomic DNA with the forward primer 5'-CTGAAGCTTTTGGCTTTGAG-3' and reverse primer 5'-CCGCTGCCAAATTCTTTGG-3,' mapping to

exons 9 and 10 of the *Kdm5c/d* genes resulting in two 331 bp X chromosome-specific amplicons in

females and an additional 301 bp Y chromosome-specific amplicon in males (Clapcote and Roder 2005).

Dissection, Dissociation, Flow Cytometry and Sorting.

Whole gut tissue (stomach, fore and hind gut) from $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ male and female

embryos at E12.5 and E14.5 were dissociated into single-cell suspensions using Accumax (Sigma, USA).

The cells were filtered serially through a 100 μm and 40 μm cell strainer, and centrifuged at 2000 rpm for

5 m. The cell pellet was resuspended in 5% FBS, 4 mM EDTA in Leibovitz L-15 medium for cell sorting

using MoFlo XDP (BD Biosciences) with a blue-violet excitation laser for CFP. Single cells were sorted

directly into 96 well plates containing 1.9 μL Triton X-100 (Sigma-Aldrich, 0.2% v/v, #T9284), 1 μL dNTP

mix (Amresco, 10 mM each, #N557), 1 μL 5' biotinylated oligo-dT$_{30}$VN primer (IDT, 10 μM), and 0.1 μL

RNasin Plus (Promega, 40 U/μL, #N2615) per well. As a no-template negative control, one well per plate

did not receive a cell. Immediately after collection, sorted plates were centrifuged (2500 rpm for ~5 s),

frozen on dry ice, and stored at -80°C until library preparation.

Single-cell RNA-Seq library Preparation

Libraries were prepared from 2 embryos for each of the 8 conditions ($Ret^{CFP/+}$ and $Ret^{CFP/CFP}$,

female and male, E12.5 and E14.5) with two exceptions: libraries were prepared from 1 $Ret^{CFP/CFP}$ male

E14.5 embryo and 4 $Ret^{CFP/CFP}$ female E14.5 embryos owing to differential viability. Single-cell RNA-seq

libraries were prepared following a modified Smart-seq2 protocol (Picelli et al. 2014; Chevée et al. 2018).

Reverse transcription and preamplification (20 cycles for E12.5 cells, 22 cycles for E14.5 cells) followed

the original Smart-seq2 protocol, with the exception of using 5' biotinylated NGG template-switching

oligonucleotides. Sequencing libraries were generated following the Nextera XT protocol (Illumina, #FC-

131-1096) on a quartered scale, using approximately 200 pg input DNA per sample and 14 enrichment

PCR cycles. Nextera XT v2 Index Sets (Illumina, #FC-131-2004) were diluted 1:5 in water prior to use.

Libraries were sequenced (paired end, 50 bp read length) on an Illumina HiSeq 2500 to an average depth

of 9.91×10^5 (standard deviation 7.43×10^5) paired-end reads per cell.

Data preprocessing

Single-cell RNA-seq libraries were aligned with hisat2 version 2.0.1-beta to a mouse mm10 hisat2

index with the inserted CFP cDNA sequence included in the index. Using samtools 1.2, sam files were

compressed to bam files then sorted and indexed. Using Cufflinks v2.2.1 indexed bam files were

quantified against the Gencode mouse vM10 assembly and normalized across all 1,369 samples (1,351

cells, 18 negative control wells). 51 samples, both in the bottom 5th percentile for total number of reads

and for which less than 50% of reads were mapped, were removed. 22 PCA outliers were removed

iteratively. The remaining 1,296 samples (1,280 cells and 16 negative control wells) were normalized

again, as before. 261 cells for which less than 1,000 genes were detected at non-zero fragments per

kilobase of transcript per million mapped reads (FPKM) were removed, and 4 more cells were

subsequently removed as PCA outliers. 28 cells were then removed due to a high ratio of mitochondrial

to nuclear gene expression, indicating poor library quality. 1,003 cells and 0 negative control wells passed

these quality control measures. The age, sex, and genotype of these 1,003 cells is as follows: 183

embryonic day (E)12.5 female *Ret*$^{CFP/+}$, 164 E12.5 male *Ret*$^{CFP/+}$, 173 E12.5 female *Ret*$^{CFP/CFP}$, 125 E12.5

male *Ret*$^{CFP/CFP}$, 161 E14.5 female *Ret*$^{CFP/+}$, 156 E14.5 male *Ret*$^{CFP/+}$, 37 E14.5 female *Ret*$^{CFP/CFP}$, and 4

E14.5 male *Ret*$^{CFP/CFP}$. FPKM values were transformed to copies per cell (CPC) using the Census

algorithm (Qiu et al. 2017). Raw and processed data have been deposited in the NCBI Gene Expression

Omnibus (GEO) under accession number GSE192676.

scRNA-Seq analysis

To reduce the computational burden, analyses of gene expression were limited to expressed

genes defined as genes with non-zero CPC values in at least 20 cells, and greater than 1×10^{-20} standard

deviation. Differential gene expression tests were performed using the Monocle R package (Trapnell et

al. 2014). To reduce noise and batch effects in visualizations a set of 419 genes with high dispersion was

defined (Chevée et al. 2018). To determine these genes, we normalized expression by read depth across

all expressed genes, then subset cells into five sets according to the pool in which each cell was

sequenced. Estimated dispersions were calculated for normalized expression of expressed genes in

each set. Overdispersed genes were defined as those with greater than 1.4 times the dispersion fit across

all five sets. Uniform manifold approximation and projection (UMAP) was performed on these 419 high-

dispersion genes using the Umapr R package (McInnes et al. 2018; [CSL STYLE ERROR: reference with

no printed form.]). RNAVelocity estimates were obtained using the Velocyto R package(La Manno et al.

2018). Cluster assignment was performed using the mClust package (Scrucca et al. 2016). Specificity

scores for each gene were calculated using the cummeRbund package (Goff et al. 2013). Briefly, Jensen-

Shannon (JS) distance was calculated between a vector of mean expression level for each cluster to a

unit vector of exclusive expression in each cluster, and specificity is defined as one minus the JS distance.

Pattern discovery with scCoGAPS

The Coordinated Gene Activity in Pattern Sets (CoGAPS) algorithm employs non-negative matrix

factorization to determine patterns of gene expression across cells and returns an amplitude (A) matrix

of gene weights by pattern and a pattern (P) matrix of cell weights by pattern. The optimal number of

patterns to learn was determined by running several iterations and selecting the maximum number of

patterns before observing redundancies and noise. Loom files for publicly available scRNA-seq data from

the enteric nervous system of adolescent mice were downloaded for analysis (mousebrain.org). 70

patterns were learned on log-transformed unique molecular identifier (UMI) counts from 11,624 libraries

prepared through the 10X Genomics platform using single-cell CoGAPS (scCoGAPS)(Stein-O'Brien et

al. 2019). We projected our data into these patterns using projectR (Sharma et al. 2019; Stein-O'Brien et

al. 2019), which returned projection weights for each pattern in each cell and each gene. For our data,

25 patterns were learned on a log-transformed, normalized expression matrix of CPC limited to

expressed genes, defined above. We projected the publicly available data into these patterns learned on

our data again using projectR.

Random Forest Model

While gene expression patterns were learned on the complete public dataset, which consists of 10,519

glia and 1,105 neurons, we trained the random forest model on all neurons and a subset of 1,660 glia to

create a 60:40 glia:neurons training dataset to avoid biasing the model to classifying glial cells. The glia

were subsampled such that the distribution of glial cells across clusters in the subset matched the

distribution across clusters in the whole glial population. The model was trained on 90% of cells to predict

cluster assignment from 70 pattern weights with 10-fold cross-validation resulting in an accuracy of

89.8%. To keep model input consistent, and because pattern weights are constrained to [0,1], projected

pattern weights were normalized to [0,1] before being used as input to predict cluster assignment. The

parent cell type of each cluster was taken as the predicted cell type for our data.

Immunofluorescence assay on whole mouse gut

Whole gut tissue (stomach, fore and hind gut) at E12.5 and E14.5 was dissected and washed in ice-cold

PBS followed by fixation in 4% PFA at 4°C for 1 h. The fixed tissues were permeabilized and blocked in

5% normal donkey serum + 0.3% Triton-X for 1 h at room temperature (RT). The tissues were incubated

with appropriate primary antibodies at 4°C overnight, washed with PBS, then incubated in appropriate

secondary antibodies at 4°C overnight (see Table 12 for antibodies and dilutions). Tissues were washed

in PBS and and incubated with 0.2 µg/mL of DAPI (4′,6-diamidino-2-phenylindole) for 10 m at RT,

followed by incubation for 20 m at RT with CUBIC-1 (Susaki et al. 2014) diluted 1:10 with PBS. Samples

were washed with PBS and mounted on Fisherbrand Superfrost Plus microscope slides with ProLong

Gold Antifade mounting media and imaged using a Nikon Eclipse Ti confocal microscope.

RNAscope

Whole gut tissue (stomach, fore and hind gut) at E12.5 was dissected and washed in ice-cold PBS

followed by fixation in 1% PFA at room temperature (RT) for several hours. Tissue was dehydrated

through an ethanol gradient and cleared with xylenes before embedding in paraffin. Tissue blocks were

sectioned on a microtome at 10 µm thickness and mounted on Fisherbrand Superfrost Plus microscope

slides. Slides were baked for 1 h at 60°C, allowed to cool to RT, then deparaffinized with xylenes.

Fluorescence in situ hybridization was performed following the manufacturer's protocol for the RNAscope

Fluorescent Multiplex V2 Assay using the RNAscope Multiplex Fluorescent Reagent Kit v2 (ACDBio

#323100). Briefly, deparaffinized sections were incubated in hydrogen peroxide for 10 m at RT and

incubated in 1X Target Retrieval for 15 m at RT, then incubated in Protease Plus for 30 m at 40°C.

Sections were incubated in a 50:1:1 mixture of RNAscope Probes CFP (ACDBio #529381), Mm-Cartpt-

C2 (ACDBio #432001-C2), and Mm-Gal-C3 (ACDBio #400961-C3) for 2 h at 40°C. Sections were

incubated with Amp 1, 2, and 3 for 30 m, 30 m, and 15 m at 40°C, respectively. Next, for each of the

three channels sections were incubated in the appropriate HRP for 15 m at 40°C, the appropriate

fluorophore for 30 m at 40°C, then HRP blocker for 15 m at 40°C. The fluorophores used were 1:750

dilutions of TSA Plus fluorescein (PerkinElmer #NEL741E001KT), cyanine 3 (PerkinElmer

#NEL744E001KT), and cyanine 5 (PerkinElmer #NEL745E001KT). Sections were then incubated in 1X

Trueblack Lipofuscin Autofluorescence Quencher (VWR #10119-144) for 1 m at RT to reduce red blood

cell autofluorescence. Sections were then incubated in RNAscope DAPI for 30 s at RT and mounted with

ProLong Gold Antifade mounting media and imaged using a Nikon Eclipse Ti confocal microscope.

Image acquisition

Tissues were visualized using a Nikon Eclipse Ti confocal microscope and processed using Fiji.

Confocal stacks were taken using a 10X air objective (Figure 2D), 20X air objective (Figure S6B-C), 40X

oil objective with Type A immersion oil (Figure 5E and proliferation quantification), or 60X oil objective

with Type A immersion oil (Figure 3E-G, Figure S6D). For image presentation, stacks were z-projected

to single images using the maximum value for each pixel. For Figure S6B, two fields of view were stitched

together (Preibisch et al. 2009).

Proliferation assay and quantification

To measure cell cycle occupancy we stained replicate $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ whole GI tracts from both

male and female mice at E14.5 for phosphohistone H3 (Anti-H3S10p; Millipore Sigma #06-570), an

immunomarker specific to cells undergoing mitosis (Skaland et al. 2009), and CFP protein (Anti-GFP;

Aves labs #GFP-1010) using the immunofluorescence assay described above. Confocal stack images

were acquired on the Nikon Eclipse Ti confocal microscope with a 40X oil objective and Type A immersion

oil and quantified using a custom CellProfiler (Carpenter et al. 2006; McQuin et al. 2018) workflow. Briefly,

primary objects were identified from each input channel (pH3, CFP) using the Otsu Adaptive threshold

strategy with a smoothing scale of 1.34888 and a threshold correction factor upper bound limit of 0.99.

The adaptive window size was set at 40. The typical diameter size ranged between 30-120 pixels

depending on whether nuclei (pH3) or cell bodies were being identified (CFP). Objects were related and

filtered using standard parameters and cells were quantified as either CFP^+, $pH3^+$, or $CFP^+/pH3^+$ double

positive. The proportion of dividing ENS precursors for each genotype was determined by estimating the

mean % of pH3+ cells within the subset of CFP+ cells for each animal. To directly test the two genotypes,

we performed Student's T-Test on the percentage of $pH3^+$ CFP^+ cells between $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$

aggregated to each replicate.

Acknowledgements

We would like to thank Hao Zhang at the Bloomberg School of Public Health, Johns Hopkins University

Cell Sorting facility for help with single cell sorting. E.V. is supported in part by the Johns Hopkins

Predoctoral Training Grant in Human Genetics (T32-GM007814-39). L.A.G. and E.V. are supported by

an award from the National Institute of Aging (R01AG066768). L.A.G. is additionally supported by the

National Institute of Aging (R01AG072305), the National Science Foundation (IOS-1665692), and a

Johns Hopkins Catalyst award. SC is supported by a NYU Grossman School of Medicine Startup Grant

(59-D-37802-30609-NYUPRJ) and AC is supported by a National Institutes of Health (Eunice Kennedy

Shriver National Institute of Child Health and Human Development) R01 award HD028088.

Competing interests

The authors declare no competing interests

Author Contributions

Elizabeth Vincent: Investigation, Data Analysis, Writing, Original draft preparation, Review and editing, Validation, Visualization; Sumantra Chatterjee: Conceptualization, Investigation, Writing, Original draft preparation, Review and editing, Supervision. Gabrielle Cannon: Investigation; Dallas Auer: Investigation; Holly Ross: Investigation; Aravinda Chakravarti: Conceptualization, Writing, Review and editing, Supervision, Funding acquisition; Loyal A. Goff: Conceptualization, Data Analysis, Writing, Original draft preparation, Review and editing, Supervision, Funding acquisition

Figures

Figure 1: A cellular map of Ret expressing enteric nervous system cells.

(A) Barplot of post quality control (QC) count of sorted $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ expressing cells from E12.5

& E14.5 embryonic gut in male and female mice. (B) Uniform Manifold Approximation and Projection

(UMAP) of 1003 cells colored by genotype. (C) UMAP embedding colored by 9 learned clusters (D)

Stacked barplot of the genotype proportions for each cluster. Asterisk denotes significant (Chi-squared

test for equal proportions, p<0.05 after multiple testing correction) deviation from expected naive

proportion. (E) UMAP embedding of _Ret_-expressing cells classified as glia/progenitor or neuron (F) RNA

velocity vector field along the UMAP embedding demonstrates direction of cellular differentiation. Dotted

line indicates neurogenic commitment, solid line indicates gliogenic commitment.

Figure 2: Cellular and molecular changes driven by Ret loss of function

(A) Heatmap of differentially expressed genes between $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ cells. (B) Significant

(hypergeometric test, p<0.05) gene ontology (GO) gene sets include neuron differentiation, gliogenesis

and cell fate commitment. C) Correlation plot of scCoGAPS patterns vs. annotated cellular features

including developmental age, sex and genotype. (D) Immunofluorescence staining of mouse GI tract at

E12.5 and E14.5 showing the organization of CFP-expressing cells. The first two columns show stomach

and the third column shows hindgut. 10X magnification, scale bar is 200 µm. (E) Violin plots of

significantly differentially expressed genes associated with the anterior-posterior patterning GO term.

Figure 3: RET expression is required for access to the Branch A developmental trajectory

(A) UMAP embedding of the neuronal cells comprising both developmental branches colored by

genotype (top) or by cluster assignment (bottom). Branch A (cluster 7) is almost entirely $Ret^{CFP/+}$ cells

while both genotypes are represented in branch B (cluster 8). (B) Heatmap of significant differentially

expressed genes with respect to genotype for cells in branch B. (C) Heatmap of significant differentially

expressed genes with respect to developmental lineage (branch) across all neurons from clusters 7 and

8. Row cluster segmentation represents the top six divisions of the hierarchical clustering of gene

expression values. (D) UMAP embeddings highlighting the persistent expression of *Ret, CFP,* and *Gfra1*

along the branch A developmental trajectory, the specific induction of the non-canonical Ret co-receptor

Gfra2 along cells in branch B, and the specific expression of the branch A marker gene *Cartpt*. (E,F)

Immunofluorescence assay of branch A marker CART at E12.5 in $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ stomach (E)

and E14.5 in $Ret^{CFP/+}$ hindgut and $Ret^{CFP/CFP}$ stomach (F) demonstrates presence of CART positive cells

only in the $Ret^{CFP/+}$ mice. (G,H) Immunofluorescence assay of branch B marker GFRA2 at E12.5 in

$Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ stomach (G) and E14.5 in $Ret^{CFP/+}$ hindgut and $Ret^{CFP/CFP}$ stomach (H). All IF assay

images are 60X magnification and scale bars are 40 μm.

Figure 4: Cellular diversity of Hirschsprung disease (HSCR) genes

(A) Heatmap of expression levels for 24 HSCR-associated genes across all isolated ENCDCs. Significant

differentially expressed genes are highlighted in red. (B) UMAP embeddings showing the expression of

11 genes that harbor pathogenic alleles in HSCR patients or are part of the *RET-EDNRB* GRN and are

differentially expressed (monocle likelihood ratio test q<0.05) with respect to genotype or with respect to

genotype-cell type interactions. Unannotated genes are HSCR-associated but not part of the *RET-*

EDNRB GRN, # indicates genes that are both associated with HSCR and are part of the *RET-EDNRB*

GRN, and ¶ indicates genes that are part of the *RET-EDNRB* GRN but have not been associated with

HSCR. (C) Violin plots of these 11 genes show expression with respect to genotype, with mean

expression value indicated by a dot. (D) Barplot showing the proportion of neurons and progenitors/glia

by genotype at E12.5 and E14.5. Dotted lines indicate the naive genotype proportion at E12.5 and E14.5.

Asterisk indicates a significant deviance from the naive proportion (Chi-squared test for equal proportions,

p<0.05).

Vincent_Fig5

Figure 5: **Ret** *LoF results in reduced cell cycle occupancy and significant reduction in the*

number of actively proliferating ENCDC progenitors.

(A) Heatmap of significant differentially expressed genes between $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ within the non-

neuronal subset of isolated ENCDCs. (B) Barplots of significant biological process gene ontology terms

from gene ontology enrichment analysis of genes with significant differential expression between

genotypes in non-neuronal ENCDCs. (C) Tricycle embedding of ENCDCs to estimate cell cycle position.

(D) Kernel density estimates for cell cycle position (θ) for each genotype at each age. Proliferative phase

of the cell cycle (non-G0/G1) is between 0.25π and 1.75π. (E) Representative immunofluorescence

staining of the proliferation marker pH3 and CFP in the stomach of E12.5 $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ mice.

Yellow arrowheads indicate $CFP^+/pH3^+$ cells. 40X magnification, scale bars are 20 μm. (F) Box &

whiskers plot showing the quantification of $CFP^+/pH3^+$ double positive cells per embryo, normalized by

the total fields of view analyzed per embryo. (G) Box & whiskers plot showing the quantification of the %

of CFP^+ cells with $pH3^+$ staining in the GI tract of E14.5 $Ret^{CFP/+}$ or $Ret^{CFP/CFP}$ mice.

Supplemental Figures

Figure S1: Uniform Manifold Approximation and Projection (UMAP) embedding of 1,003 cells colored by (A) age, (B) sex, and (C) sequencing batch.

Figure S2: Cluster-specific gene expression profiles

(A) Scatter plot of mean gene expression per cluster vs. cluster specificity scores for each of the 9 learned clusters of enriched ENCDCs. (B) UMAP embeddings highlighting the gene expression of top specific cluster markers for each cluster. (C) UMAP embedding showing the expression levels for the ENS glial marker *Ngfr* and the neuronal marker *Tubb3*.

Figure S3: Projection analysis of adult ENS cell types

Each panel describes the usage of a single scCoGAPS pattern learned from (Zeisel et al. 2018). For each of the 70 patterns, the left panel is a reduced dimensionality visualization of the glial cells and the middle panel is for the neuronal cells from (Zeisel et al. 2018). Cells are colored based on their usage of the learned pattern. The right panel is a UMAP embedding of the ENCDCs isolated in this study and cells are colored by the corresponding projected pattern weight, highlighting the shared use of specific biological features across both datasets.

Figure S4: Cell type annotation via transfer learning

(A) Reduced dimensionality visualization of mature ENS cells from (Zeisel et al. 2018) colored by previously annotated cluster designations. (B) UMAP embedding of sorted ENCDCs colored by the random forest model predictions for corresponding clusters from Zeisel 2018. (C) UMAP embedding of sorted ENCDCs colored by top-level ENS cell type designation. (D) Stacked barplot of cell type proportions for each of the nine learned clusters of ENCDCs. (E) UMAP embedding of Zeisel 2018 mature ENS cells by previously annotated cell type designations.

Figure S5: Learned gene expression patterns across all cells

25 single-cell coordinated gene activity in pattern sets (scCoGAPS) patterns learned from sorted *Ret-*expressing ENCDCs. Each UMAP embedding is colored by the usage of a specific pattern of gene co-expression learned through non-negative matrix factorization.

Figure S6: RNAscope single molecule fluorescence in situ hybridization validation of branch A and branch B cells

(A) UMAP embedding of ENCDC cells highlighting previously identified (Morarach et al. 2021) marker gene expression for neuron developmental branches (branch A: *Nos1, Vip, Gal, Etv1*; branch B: *Bnc2, Dlx5, Mgat4c, Ndufa4l2*). (B,C) RNAscope single molecule fluorescence *in situ* hybridization (smFISH) validation of *Gal/Cartpt* in cross section of E12.5 stomach in both *Ret^{CFP/+}* and *Ret^{CFP/CFP}* mice. The yellow box in (B) indicates the portion isolated in (C). 10X magnification, scale bars are 200 μm (B) and 50 μm (C). (D) Whole mount immunofluorescence validation of branch B cells (GFRA2⁺/CFP⁺) in E12.5 esophagus of both *Ret^{CFP/+}* and *Ret^{CFP/CFP}* mice. 60X magnification, scale bars are 20 μm.

Figure S7: Expression profiles for Hirschsprung-associated and RET-EDNRB gene regulatory network genes

(A) UMAP embeddings showing normalized expression of 21 expressed genes associated with Hirschsprung disease (HSCR). (B) Violin plots showing expression of these same 21 genes by genotype. (C) Violin plots showing expression of these same 21 genes by cluster. (D) UMAP embeddings showing normalized expression of 3 genes in the *RET-EDNRB* GRN that have not been associated with Hirschsprung disease. (E) Violin plots showing expression of these same 3 genes by genotype. (C) Violin plots showing expression of these same 3 genes by cluster. Genes in red are significantly differentially expressed (monocle likelihood ratio test q<0.05) either with respect to genotype or with respect to

genotype-cell type interactions. # indicates genes that are shared across the HSCR-associated and *RET-*

EDNRB GRN sets.

Figure S8: Cell cycle is modulated by Ret LoF

(A) Heatmap of significant differentially expressed genes with respect to the combinatorial effect of both

genotype and celltype (i.e. genotype:celltype effect). (B) Barplot of significantly enriched gene ontology

terms for the list of genes identified as differentially expressed in A. Ontology coloring matches the row

color annotations for each gene in A. (C) UMAP embeddings showing the projected pattern weights for

the cell cycle stage-specific patterns (40-42) learned from (Zeisel et al. 2018). (D) UMAP embedding

colored by the maximum projection value across each projected cell cycle stage-specific pattern. (E)

Empirical cumulative density functions for projection values of cell cycle stage-specific patterns for each

of the interaction terms between age and genotype. Significant differences in distributions were

determined using the Kolmogorov-Smirnov test (* = $p<0.05$; ** = $p<0.01$ after multiple testing correction.

(F) Empirical cumulative density function (eCDF) curve across all projected (Zeisel et al. 2018)

scCoGAPS patterns. One curve for each of the interaction terms between age and genotype are

presented. (G-J) Box & whisker plots showing the distribution of quantifications for pH3 and CFP

immunofluorescence staining in E14.5 $Ret^{CFP/+}$ and $Ret^{CFP/CFP}$ mouse GI tracts. Each point represents a

separate sampled field of view from one of four replicate mice from each condition. Points are colored by

replicate embryos. G) Count of CFP^+ cells, H) Count of $pH3^+$ cells, I) Count of double positive $CFP^+/pH3^+$

cells, and J) Percent of CFP^+ cells that are $pH3^+$.

Tables

Table 1: Distribution of cells before and after QC
The age, sex, and *Ret* genotype of each embryo, as well as the number of libraries that were prepared,
sequenced, and passed all quality control filters.
915
Table 2: Gene specificity scores by cluster

Specificity scores for each gene (row) in each cluster (column). Specificity scores range from [0, 1], with
a higher score indicating greater specificity.

Table 3: Gut atlas scCoGAPS A and P weights
Sheet "A-Pmatrix" contains the Pattern (P) matrix of pattern weight values for each cell (row) in each
pattern (column) across 70 patterns learned on gut scRNAseq atlas data from Zeisel et al, 2018. Sheet
"B-Amatrix" contains the Amplitude (A) matrix of pattern weight values for each gene (row) in each
pattern (column).

Table 4: Gut atlas pattern markers
Sheet "A-patternMarkers" lists genes that are pattern markers for the 70 patterns in Table 3. Sheet "B-
GO_terms" lists the corresponding gene ontology:biological process (GO:BP) terms for which the
pattern marker lists are enriched. Sheet "C-GO_term_genes" lists the specific genes associated with
each enriched GO:BP term. An entry of "not run" on sheet B indicates the pattern marker gene list for
that particular pattern was not long enough (at least 10 genes) after converting to Entrez IDs, therefore
gene ontology enrichment was not run on that list. An empty column under a pattern name on sheet B
indicates no GO:BP terms reached significance (q<.05) for enrichment in pattern markers of that
pattern.

Table 5: ENCDC data projected into gut atlas
Projected pattern weights (sheet "projection") and p-values (sheet "pval") for the 70 patterns (row)
learned on the gut atlas into each cell (column) from the Ret mutant dataset.

Table 6: Differentially expressed gene lists
Each sheet presents results from a single differential expression test. At the top of each sheet we report
which cells were tested ("cells"), the parameter tested ("test"), the full model formula ("full_model"), and
the reduced model formula ("reduced_model"). Gene lists are subset to significantly differentially
expressed (q < 0.05) genes.

Table 7: Differentially expressed gene GO lists
Each sheet presents results from a single gene ontology (GO) enrichment test, including the number of
differentially expressed genes associated with each term ("Count") and the official gene symbol for
those genes ("geneID") . The sheets in Table 7 report the GO enrichment results for the corresponding
differentially expressed gene lists in Table 6. GO enrichment lists are subset to significantly enriched (q
< 0.05) terms.

Table 8: ENCDC scCoGAPS A and P weights
Sheet "A-Pmatrix" contains the Pattern (P) matrix of pattern weight values for each cell (row) in each
pattern (column) across 25 patterns learned on the Ret mutant dataset. Sheet "B-Amatrix" contains the
Amplitude (A) matrix of pattern weight values for each gene (row) in each pattern (column).

Table 9: ENCDC patterns correlated with metadata

Correlation (sheet "A-correlation") and the p-value for correlation (sheet "B-correlation-pvalue") of
various cell metadata parameters (row) with pattern weights for 25 patterns learned on the Ret mutant
data (column).
962
963   Table 10: ENCDC data pattern markers
964   A list of marker genes each of the 25 patterns (column) learned on the Ret mutant data.
965
966   Table 11: Tricycle theta estimates
967   Theta estimates from tricycle ("tricyclePosition") as well as relevant metadata for each cell (row).
968
969   Table 12: Antibodies and Concentrations Used
970   A list of all antibodies used in this study.

References

Alves MM, Sribudiani Y, Brouwer RWW, Amiel J, Antiñolo G, Borrego S, Ceccherini I, Chakravarti A, Fernández RM, Garcia-Barcelo M-M, et al. 2013. Contribution of rare and common variants determine complex diseases-Hirschsprung disease as a model. *Dev Biol* **382**: 320–329. http://dx.doi.org/10.1016/j.ydbio.2013.05.019.

Amiel J, Sproat-Emison E, Garcia-Barcelo M, Lantieri F, Burzynski G, Borrego S, Pelet A, Arnold S, Miao X, Griseri P, et al. 2008. Hirschsprung disease, associated syndromes and genetics: a review. *J Med Genet* **45**: 1–14. http://jmg.bmj.com/content/45/1/1.short.

Bergeron K-F, Nguyen CMA, Cardinal T, Charrier B, Silversides DW, Pilon N. 2016. Upregulation of the Nr2f1-A830082K12Rik gene pair in murine neural crest cells results in a complex phenotype reminiscent of Waardenburg syndrome type 4. *Dis Model Mech* **9**: 1283–1293. http://dx.doi.org/10.1242/dmm.026773.

Bogni S, Trainor P, Natarajan D, Krumlauf R, Pachnis V. 2008. Non-cell-autonomous effects of Ret deletion in early enteric neurogenesis. *Development* **135**: 3007–3011. http://dx.doi.org/10.1242/dev.025163.

Carpenter AE, Jones TR, Lamprecht MR, Clarke C, Kang IH, Friman O, Guertin DA, Chang JH, Lindquist RA, Moffat J, et al. 2006. CellProfiler: image analysis software for identifying and quantifying cell phenotypes. *Genome Biol* **7**: R100. http://dx.doi.org/10.1186/gb-2006-7-10-r100.

Carrasquillo MM, McCallion AS, Puffenberger EG, Kashuk CS, Nouri N, Chakravarti A. 2002. Genome-wide association study and mouse model identify interaction between RET and EDNRB pathways in Hirschsprung disease. *Nat Genet* **32**: 237–244. http://dx.doi.org/10.1038/ng998.

Chatterjee S, Chakravarti A. 2019. A gene regulatory network explains RET-EDNRB epistasis in Hirschsprung disease. *Hum Mol Genet* **28**: 3137–3147. http://dx.doi.org/10.1093/hmg/ddz149.

Chatterjee S, Kapoor A, Akiyama JA, Auer DR, Lee D, Gabriel S, Berrios C, Pennacchio LA, Chakravarti A. 2016a. Enhancer Variants Synergistically Drive Dysfunction of a Gene Regulatory Network In Hirschsprung Disease. *Cell* **167**: 355–368.e10. http://dx.doi.org/10.1016/j.cell.2016.09.005.

Chatterjee S, Karasaki KM, Fries LE, Kapoor A, Chakravarti A. 2021. A multi-enhancer RET regulatory code is disrupted in Hirschsprung disease. *Genome Res* **31**: 2199–2208. http://genome.cshlp.org/cgi/pmidlookup?view=long&pmid=34782358 (Accessed July 19, 2021).

Chatterjee S, Nandakumar P, Auer DR, Gabriel SB, Chakravarti A. 2019. Gene- and tissue-level interactions in normal gastrointestinal development and Hirschsprung disease. *Proceedings of the National Academy of Sciences* **116**: 26697–26708. http://dx.doi.org/10.1073/pnas.1908756116.

Chatterjee S, Stine ZE, Mccallion AS, Chakravarti A. 2016b. RET Mutation and Function in HSCR, MEN2, and Other Cancers. In *Epstein's Inborn Errors of Development* (eds. R.P. Erickson and A.J. Wynshaw-Boris), pp. 517–523, Oxford University Press http://oxfordmedicine.com/view/10.1093/med/9780199934522.001.0001/med-9780199934522-chapter-69 (Accessed August 24, 2021).

Chevée M, Robertson JDJ, Cannon GH, Brown SP, Goff LA. 2018. Variation in Activity State, Axonal

Projection, and Position Define the Transcriptional Identity of Individual Neocortical Projection
Neurons. *Cell Rep* **22**: 441–455. http://dx.doi.org/10.1016/j.celrep.2017.12.046.

Clapcote SJ, Roder JC. 2005. Simplex PCR assay for sex determination in mice. *Biotechniques* **38**:
702, 704, 706. http://dx.doi.org/10.2144/05385BM05.

Dang R, Torigoe D, Suzuki S, Kikkawa Y, Moritoh K, Sasaki N, Agui T. 2011. Genetic Background
Strongly Modifies the Severity of Symptoms of Hirschsprung Disease, but Not Hearing Loss in
Rats Carrying Ednrbsl Mutations. *PLoS ONE* **6**: e24086.
http://dx.doi.org/10.1371/journal.pone.0024086.

Durbec PL, Larsson-Blomberg LB, Schuchardt A, Costantini F, Pachnis V. 1996. Common origin and
developmental dependence on c-ret of subsets of enteric and sympathetic neuroblasts.
Development **122**: 349–358. http://dx.doi.org/10.1242/dev.122.1.349.

Elmentaite R, Kumasaka N, Roberts K, Fleming A, Dann E, King HW, Kleshchevnikov V, Dabrowska
M, Pritchard S, Bolt L, et al. 2021. Cells of the human intestinal tract mapped across space and
time. *Nature* **597**: 250–255. http://dx.doi.org/10.1038/s41586-021-03852-1.

Fujiwara N, Nakazawa-Tanaka N, Miyahara K, Arikawa-Hirasawa E, Akazawa C, Yamataka A. 2018.
Altered expression of laminin alpha1 in aganglionic colon of endothelin receptor-B null mouse
model of Hirschsprung's disease. *Pediatr Surg Int* **34**: 137–141. http://dx.doi.org/10.1007/s00383-
017-4180-6.

Furness JB. 2012. The enteric nervous system and neurogastroenterology. *Nat Rev Gastroenterol*
Hepatol **9**: 286–294. http://dx.doi.org/10.1038/nrgastro.2012.32.

Furness JB. 2008. The enteric nervous system: normal functions and enteric neuropathies.
Neurogastroenterol Motil **20 Suppl 1**: 32–38. http://dx.doi.org/10.1111/j.1365-2982.2008.01094.x.

Gattelli A, Nalvarte I, Boulay A, Roloff TC, Schreiber M, Carragher N, Macleod KK, Schlederer M,
Lienhard S, Kenner L, et al. 2013. Ret inhibition decreases growth and metastatic potential of
estrogen receptor positive breast cancer cells. *EMBO Mol Med* **5**: 1335–1350.
http://dx.doi.org/10.1002/emmm.201302625.

Gershon MD. 2010. Developmental determinants of the independence and complexity of the enteric
nervous system. *Trends Neurosci* **33**: 446–456. http://dx.doi.org/10.1016/j.tins.2010.06.002.

Goff L, Trapnell C, Kelley D. 2013. cummeRbund: Analysis, exploration, manipulation, and visualization
of Cufflinks high-throughput sequencing data. *R package version* **2**.

Goldstein AM, Hofstra RMW, Burns AJ. 2013. Building a brain in the gut: development of the enteric
nervous system. *Clin Genet* **83**: 307–316. http://dx.doi.org/10.1111/cge.12054.

Gould TW, Yonemura S, Oppenheim RW, Ohmori S, Enomoto H. 2008. The neurotrophic effects of
glial cell line-derived neurotrophic factor on spinal motoneurons are restricted to fusimotor
subtypes. *J Neurosci* **28**: 2131–2146. http://dx.doi.org/10.1523/JNEUROSCI.5185-07.2008.

Grieco M, Santoro M, Berlingieri MT, Melillo RM, Donghi R, Bongarzone I, Pierotti MA, Della Porta G,
Fusco A, Vecchio G. 1990. PTC is a novel rearranged form of the ret proto-oncogene and is
frequently detected in vivo in human thyroid papillary carcinomas. *Cell* **60**: 557–563.
http://dx.doi.org/10.1016/0092-8674(90)90659-3.

Hao MM, Young HM. 2009. Development of enteric neuron diversity. *J Cell Mol Med* **13**: 1193–1210.
http://dx.doi.org/10.1111/j.1582-4934.2009.00813.x.

Heanue TA, Pachnis V. 2006. Expression profiling the developing mammalian enteric nervous system
identifies marker and candidate Hirschsprung disease genes. *Proceedings of the National
Academy of Sciences* **103**: 6919–6924. http://dx.doi.org/10.1073/pnas.0602152103.

Heuckeroth RO. 2018. Hirschsprung disease - integrating basic science and clinical medicine to
improve outcomes. *Nat Rev Gastroenterol Hepatol* **15**: 152–167.
http://dx.doi.org/10.1038/nrgastro.2017.149.

Heuckeroth RO, Enomoto H, Grider JR, Golden JP, Hanke JA, Jackman A, Molliver DC, Bardgett ME,
Snider WD, Johnson EM Jr, et al. 1999. Gene targeting reveals a critical role for neurturin in the
development and maintenance of enteric, sensory, and parasympathetic neurons. *Neuron* **22**:
253–263. http://dx.doi.org/10.1016/s0896-6273(00)81087-9.

Hirst CS, Stamp LA, Bergner AJ, Hao MM, Tran MX, Morgan JM, Dutschmann M, Allen AM, Paxinos G,
Furlong TM, et al. 2018. Publisher Correction: Kif1bp loss in mice leads to defects in the peripheral
and central nervous system and perinatal death. *Sci Rep* **8**: 9085.
http://dx.doi.org/10.1038/s41598-018-26661-5.

Jain PC, Varadarajan R. 2014. A rapid, efficient, and economical inverse polymerase chain reaction-
based method for generating a site saturation mutant library. *Anal Biochem* **449**: 90–98.
http://dx.doi.org/10.1016/j.ab.2013.12.002.

Kang Y-N, Fung C, Vanden Berghe P. 2021. Gut innervation and enteric nervous system development:
a spatial, temporal and molecular tour de force. *Development* **148**.
http://dx.doi.org/10.1242/dev.182543.

Kapoor A, Auer DR, Lee D, Chatterjee S, Chakravarti A. 2017. Testing the Ret and Sema3d genetic
interaction in mouse enteric nervous system development. *Hum Mol Genet* **26**: 1811–1820.
http://dx.doi.org/10.1093/hmg/ddx084.

Kapoor A, Jiang Q, Chatterjee S, Chakraborty P, Sosa MX, Berrios C, Chakravarti A. 2015. Population
variation in total genetic risk of Hirschsprung disease from common RET, SEMA3 and NRG1
susceptibility polymorphisms. *Hum Mol Genet* **24**: 2997–3003.
http://dx.doi.org/10.1093/hmg/ddv051.

Kapur RP, Yost C, Palmiter RD. 1992. A transgenic model for studying development of the enteric
nervous system in normal and aganglionic mice. *Development* **116**: 167–175.
http://dx.doi.org/10.1242/dev.116.Supplement.167.

Kulkarni S, Ganz J, Bayrer J, Becker L, Bogunovic M, Rao M. 2018. Advances in Enteric Neurobiology:
The "Brain" in the Gut in Health and Disease. *J Neurosci* **38**: 9346–9354.
http://dx.doi.org/10.1523/JNEUROSCI.1663-18.2018.

Kulkarni S, Saha M, Becker L, Wang Z, Liu G, Leser J, Kumar M, Bakhshi S, Anderson M, Lewandoski
M, et al. 2020. Neural crest-derived neurons are replaced by a newly identified mesodermal
lineage in the post-natal and aging enteric nervous system. *bioRxiv* 2020.08.25.262832.
https://www.biorxiv.org/content/10.1101/2020.08.25.262832v1 (Accessed July 19, 2021).

La Manno G, Soldatov R, Zeisel A, Braun E, Hochgerner H, Petukhov V, Lidschreiber K, Kastriti ME,

Lönnerberg P, Furlan A, et al. 2018. RNA velocity of single cells. *Nature* **560**: 494–498.
http://dx.doi.org/10.1038/s41586-018-0414-6.

Lang D, Chen F, Milewski R, Li J, Lu MM, Epstein JA. 2000. Pax3 is required for enteric ganglia
formation and functions with Sox10 to modulate expression of c-ret. *J Clin Invest* **106**: 963–971.
http://dx.doi.org/10.1172/JCI10828.

Lasrado R, Boesmans W, Kleinjung J, Pin C, Bell D, Bhaw L, McCallum S, Zong H, Luo L, Clevers H,
et al. 2017. Lineage-dependent spatial and functional organization of the mammalian enteric
nervous system. *Science* **356**: 722–726. http://dx.doi.org/10.1126/science.aam7511.

Le Douarin NM, Smith J. 1988. Development of the peripheral nervous system from the neural crest.
Annu Rev Cell Biol **4**: 375–404. http://dx.doi.org/10.1146/annurev.cb.04.110188.002111.

Lui VCH, Samy ET, Sham MH, Mulligan LM, Tam PKH. 2002. Glial Cell Line–Derived Neurotrophic
Factor Family Receptors Are Abnormally Expressed in Aganglionic Bowel of a Subpopulation of
Patients with Hirschsprung's Disease. *Laboratory Investigation* **82**: 703–712.
http://dx.doi.org/10.1097/01.lab.0000017364.13014.ae.

McCallion AS, Stames E, Conlon RA, Chakravarti A. 2003. Phenotype variation in two-locus mouse
models of Hirschsprung disease: tissue-specific interaction between Ret and Ednrb. *Proc Natl*
Acad Sci U S A **100**: 1826–1831. http://dx.doi.org/10.1073/pnas.0337540100.

McInnes L, Healy J, Saul N, Großberger L. 2018. UMAP: Uniform Manifold Approximation and
Projection. *J Open Source Softw* **3**: 861. http://joss.theoj.org/papers/10.21105/joss.00861.

McQuin C, Goodman A, Chernyshev V, Kamentsky L, Cimini BA, Karhohs KW, Doan M, Ding L,
Rafelski SM, Thirstrup D, et al. 2018. CellProfiler 3.0: Next-generation image processing for
biology. *PLoS Biol* **16**: e2005970. http://dx.doi.org/10.1371/journal.pbio.2005970.

Morarach K, Memic F, Zeisel A, Hochgerner H, Marklund U, Linnarsson S. 2017. Single-cell molecular
interrogation of enteric nervous system development. *Mech Dev* S67.
https://www.infona.pl/resource/bwmeta1.element.elsevier-65aec147-c938-33e9-beeb-
d4eb732d66ef.

Morarach K, Mikhailova A, Knoflach V, Memic F, Kumar R, Li W, Ernfors P, Marklund U. 2021.
Diversification of molecularly defined myenteric neuron classes revealed by single-cell RNA
sequencing. *Nat Neurosci* **24**: 34–46. http://dx.doi.org/10.1038/s41593-020-00736-x.

Mulligan LM. 2014. RET revisited: expanding the oncogenic portfolio. *Nat Rev Cancer* **14**: 173–186.
http://dx.doi.org/10.1038/nrc3680.

Nagy N, Goldstein AM. 2017. Enteric nervous system development: A crest cell's journey from neural
tube to colon. *Semin Cell Dev Biol* **66**: 94–106. http://dx.doi.org/10.1016/j.semcdb.2017.01.006.

Natarajan D, Marcos-Gutierrez C, Pachnis V, de Graaff E. 2002. Requirement of signalling by receptor
tyrosine kinase RET for the directed migration of enteric nervous system progenitor cells during
mammalian embryogenesis. *Development* **129**: 5151–5160.
https://www.ncbi.nlm.nih.gov/pubmed/12399307.

Obermayr F, Hotta R, Enomoto H, Young HM. 2013. Development and developmental disorders of the
enteric nervous system. *Nat Rev Gastroenterol Hepatol* **10**: 43–57.

http://dx.doi.org/10.1038/nrgastro.2012.234.

Paratore C, Eichenberger C, Suter U, Sommer L. 2002. Sox10 haploinsufficiency affects maintenance
of progenitor cells in a mouse model of Hirschsprung disease. *Hum Mol Genet* **11**: 3075–3085.
http://dx.doi.org/10.1093/hmg/11.24.3075.

Park H-J, Bolton EC. 2017. RET-mediated glial cell line-derived neurotrophic factor signaling inhibits
mouse prostate development. *Development* **144**: 2282–2293.
http://dx.doi.org/10.1242/dev.145086.

Pattyn A, Morin X, Cremer H, Goridis C, Brunet JF. 1999. The homeobox gene Phox2b is essential for
the development of autonomic neural crest derivatives. *Nature* **399**: 366–370.
http://dx.doi.org/10.1038/20700.

Perea D, Guiu J, Hudry B, Konstantinidou C, Milona A, Hadjieconomou D, Carroll T, Hoyer N,
Natarajan D, Kallijärvi J, et al. 2017. Ret receptor tyrosine kinase sustains proliferation and tissue
maturation in intestinal epithelia. *EMBO J* **36**: 3029–3045.
http://dx.doi.org/10.15252/embj.201696247.

Picelli S, Faridani OR, Björklund AK, Winberg G, Sagasser S, Sandberg R. 2014. Full-length RNA-seq
from single cells using Smart-seq2. *Nat Protoc* **9**: 171–181.
http://dx.doi.org/10.1038/nprot.2014.006.

Preibisch S, Saalfeld S, Tomancak P. 2009. Globally optimal stitching of tiled 3D microscopic image
acquisitions. *Bioinformatics* **25**: 1463–1465. http://dx.doi.org/10.1093/bioinformatics/btp184.

Qiu X, Hill A, Packer J, Lin D, Ma Y-A, Trapnell C. 2017. Single-cell mRNA quantification and
differential analysis with Census. *Nat Methods* **14**: 309–315. http://dx.doi.org/10.1038/nmeth.4150.

Rao M, Gershon MD. 2016. The bowel and beyond: the enteric nervous system in neurological
disorders. *Nat Rev Gastroenterol Hepatol* **13**: 517–528.
http://dx.doi.org/10.1038/nrgastro.2016.107.

Rossi J, Luukko K, Poteryaev D, Laurikainen A, Sun YF, Laakso T, Eerikäinen S, Tuominen R, Lakso
M, Rauvala H, et al. 1999. Retarded Growth and Deficits in the Enteric and Parasympathetic
Nervous System in Mice Lacking GFRα2, a Functional Neurturin Receptor. *Neuron* **22**: 243–252.
https://www.sciencedirect.com/science/article/pii/S0896627300810867.

Schuchardt A, D'Agati V, Larsson-Blomberg L, Costantini F, Pachnis V. 1994. Defects in the kidney and
enteric nervous system of mice lacking the tyrosine kinase receptor Ret. *Nature* **367**: 380–383.
http://dx.doi.org/10.1038/367380a0.

Scrucca L, Fop M, Murphy TB, Raftery AE. 2016. *mclust 5: Clustering, Classification and density*
estimation using Gaussian finite mixture models. R J. 8: 289--317. doi: 10.32614. RJ-2016-021.

Sharma G, Colantuoni C, Goff LA, Fertig EJ, Stein-O'Brien G. 2019. projectR: An R/Bioconductor
package for transfer learning via PCA, NMF, correlation, and clustering. *bioRxiv* 726547.
https://www.biorxiv.org/content/10.1101/726547v1.abstract (Accessed August 26, 2019).

Skaland I, Janssen EAM, Gudlaugsson E, Hui Ru Guo L, Baak J. 2009. The prognostic value of the
proliferation marker phosphohistone H3 (PPH3) in luminal, basal-like and triple negative phenotype
invasive lymph node-negative breast cancer. *Anal Cell Pathol* **31**: 261–271.

https://content.iospress.com/articles/analytical-cellular-pathology/clo464.

Southard-Smith EM, Kos L, Pavan WJ. 1998. Sox10 mutation disrupts neural crest development in
Dom Hirschsprung mouse model. *Nat Genet* **18**: 60–64. http://dx.doi.org/10.1038/ng0198-60.

Stein-O'Brien GL, Carey JL, Lee WS, Considine M, Favorov AV, Flam E, Guo T, Li S, Marchionni L,
Sherman T, et al. 2017. PatternMarkers & GWCoGAPS for novel data-driven biomarkers via whole
transcriptome NMF. *Bioinformatics* **33**: 1892–1894. http://dx.doi.org/10.1093/bioinformatics/btx058.

Stein-O'Brien GL, Clark BS, Sherman T, Zibetti C, Hu Q, Sealfon R, Liu S, Qian J, Colantuoni C,
Blackshaw S, et al. 2019. Decomposing Cell Identity for Transfer Learning across Cellular
Measurements, Platforms, Tissues, and Species. *Cell Syst* **8**: 395–411.e8.
http://dx.doi.org/10.1016/j.cels.2019.04.004.

Susaki EA, Tainaka K, Perrin D, Kishino F, Tawara T, Watanabe TM, Yokoyama C, Onoe H, Eguchi M,
Yamaguchi S, et al. 2014. Whole-brain imaging with single-cell resolution using chemical cocktails
and computational analysis. *Cell* **157**: 726–739. http://dx.doi.org/10.1016/j.cell.2014.03.042.

Tani G, Tomuschat C, O'Donnell AM, Coyle D, Puri P. 2017. Increased population of immature enteric
glial cells in the resected proximal ganglionic bowel of Hirschsprung's disease patients. *J Surg Res*
218: 150–155. http://dx.doi.org/10.1016/j.jss.2017.05.062.

Tilghman JM, Ling AY, Turner TN, Sosa MX, Krumm N, Chatterjee S, Kapoor A, Coe BP, Nguyen K-
DH, Gupta N, et al. 2019. Molecular Genetic Anatomy and Risk Profile of Hirschsprung's Disease.
N Engl J Med **380**: 1421–1432. http://dx.doi.org/10.1056/NEJMoa1706594.

Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ, Mikkelsen TS,
Rinn JL. 2014. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal
ordering of single cells. *Nat Biotechnol* **32**: 381–386. http://dx.doi.org/10.1038/nbt.2859.

Uesaka T, Nagashimada M, Enomoto H. 2015. Neuronal Differentiation in Schwann Cell Lineage
Underlies Postnatal Neurogenesis in the Enteric Nervous System. *J Neurosci* **35**: 9879–9888.
http://dx.doi.org/10.1523/JNEUROSCI.1239-15.2015.

Uesaka T, Nagashimada M, Yonemura S, Enomoto H. 2008. Diminished Ret expression compromises
neuronal survival in the colon and causes intestinal aganglionosis in mice. *J Clin Invest* **118**: 1890–
1898. http://dx.doi.org/10.1172/JCI34425.

Young HM, Ciampoli D, Hsuan J, Canty AJ. 1999. Expression of Ret-, p75(NTR)-, Phox2a-, Phox2b-,
and tyrosine hydroxylase-immunoreactivity by undifferentiated neural crest-derived cells and
different classes of enteric neurons in the embryonic mouse gut. *Dev Dyn* **216**: 137–152.
https://onlinelibrary.wiley.com/doi/10.1002/(SICI)1097-0177(199910)216:2%3C137::AID-
DVDY5%3E3.0.CO;2-6.

Young HM, Hearn CJ, Ciampoli D, Southwell BR, Brunet JF, Newgreen DF. 1998. A single rostrocaudal
colonization of the rodent intestine by enteric neuron precursors is revealed by the expression of
Phox2b, Ret, and p75 and by explants grown under the kidney capsule or in organ culture. *Dev*
Biol **202**: 67–84. http://dx.doi.org/10.1006/dbio.1998.8987.

Zeisel A, Hochgerner H, Lönnerberg P, Johnsson A, Memic F, van der Zwan J, Häring M, Braun E,
Borm LE, La Manno G, et al. 2018. Molecular Architecture of the Mouse Nervous System. *Cell*
174: 999–1014.e22. http://dx.doi.org/10.1016/j.cell.2018.06.021.

Zheng S, Stein-O'Brien G, Augustin J, Slosberg J, Carosso GA, Winer B, Shin G, Bjornsson HT, Goff
LA, Hansen KD. 2021. Universal prediction of cell cycle position using transfer learning. *bioRxiv*.
https://www.biorxiv.org/content/10.1101/2021.04.06.438463v2.abstract.

umapr: ARCHIVED UMAP dimensionality reduction in R. Github https://github.com/ropensci-
archive/umapr (Accessed October 22, 2021).

212