Title Page:

AtERF60, **negatively regulates** *ABR1* **to modulate basal resistance response in**

Arabidopsis

Running title: *AtERF60* **regulates biotic stress response**

Authors: Ashutosh Joshi[1#], Ujjal J Phukan[1#], Neeti Singh[1,2#], Gajendra Singh

Jeena[1†],Sunil Kumar[1†],Durgesh Kumar Pandey[1], Alok Pandey[2], Vineeta Tripathi[2,4],

Rakesh Kumar Shukla*[1,2]

*Corresponding Author

#- Authors contributed equally

†- Authors contributed equally.

Contact details:

1. Plant Biotechnology Division, CSIR-Central Institute of Medicinal and Aromatic Plants,

P.O. CIMAP, Lucknow-226015, India.

2. Academy of Scientific and Innovative Research (AcSIR), CSIR-Central Institute of

Medicinal and Aromatic Plants (CSIR-CIMAP)

3. Microbial Technology Department, CSIR-Central Institute of Medicinal and Aromatic

Plants, P.O. CIMAP, Lucknow-226015, India.

4. Ethanobotany Division, CSIR- Central Drug Research Institute, Jankipuram Extension,

Sitapur Road, Lucknow-226031, India.

Corresponding Author addresses and email id: Plant Biotechnology Division, CSIR-Central

Institute of Medicinal and Aromatic Plants, P.O. CIMAP, Lucknow-226015, India.

email: rk.shukla@cimap.res.in

Total word count: 4182

Colour figure: 5

Supplementary Figure 5

Supplementary Table: 3

Abstract

APETALA2 (AP2)/ERF family transcription factors (TFs) contribute an important function

against various external cues. Our study reports *AtERF60*, an AP2/ERF TF, which plays a

key role in regulating the *ABR1* gene, leading to an altered basal resistance response in

Arabidopsis. *AtERF60* is induced in response to drought, salt, abscisic acid (ABA), salicylic

acid (SA), and bacterial pathogen *Pst*DC3000 infection. *AtERF60* interacts with

DEHYDRATION RESPONSE ELEMENTS and GCC box, indicating its ability to regulate

multiple responses. The overexpressing lines of *AtERF60* demonstrated increased resistance

to *Pst*DC3000 infection, whereas *erf60* mutant lines showed increased susceptibility.

Complementation of the *erf60* mutant background exhibits no significant difference towards

*Pst*DC3000 infection compared with the WT Col-0. Microarray and qRT-PCR analysis of

overexpression and mutant lines indicated that *AtERF60* regulates stress-inducible genes. The

induction of these differentially expressed transcripts was significantly increased in *erf60*

mutant lines, whereas it was reversed in wild-type lines when *AtERF60* was complemented.

ABR1 was one of the differentially expressed transcripts, and we discovered that *AtERF60*

interacts with the DRE *cis*-elements in the *ABR1* promoter. Further, qRT-PCR expression

analysis after infection with *Pst*DC3000 in *AtERF60- OX*, *erf60* and its complementation

background suggest that the mutation in *AtERF60* upregulates *ABR1* activity, leading to the

enhanced susceptibility towards *Pst*DC3000. Conversely, *AtERF60* overexpression

suppresses *ABR1* activity, strengthening the basal resistance responses of Arabidopsis.

Keywords

AtERF60, AP2/ERF, Abiotic stress, *Pst*DC3000, DRE, Overexpression

Introduction

Plants face multiple environmental stresses, including biotic and abiotic stresses, which

disturb their normal growth and development, leading to massive agricultural losses (He *et*

al., 2018). To adapt to these environmental conditions, plants have developed various

evolutionary mechanisms governed by a complex regulatory network. Transcription factors

regulate different interconnected and diverse signalling cascades by interacting with *cis*-

elements present in their promoters (Liu *et al*., 2014). AP2/ERFs are one of the most

important transcription factor families that appeared as a key regulator of a large cluster of

downstream target genes involved in various stress responses (Phukan *et al*., 2017). The

AP2/ETHYLENE RESPONSIVE ELEMENT BINDING FACTOR (EREB) domain of

AP2/ERFs is made up of conserved 40–70 amino acids that act as a DNA binding domain

(Nakano *et al*., 2006). Based on the DNA binding domain, AP2/ERFs are classified into three

groups: first is ERF (single ERF domain, contains subgroup I to X, VI-L and Xb-L), second

is AP2 (tandem copies of two AP2 domains and few of them are with single AP2 domain),

and third is RAV (ERF domain associated with a B3 DNA-binding domain). Soloist is a

protein that contains an ERF domain; however, its sequence and gene structure strongly

diverge from ERF TFs (Shigyo and Ito, 2004, Nakano *et al*., 2006, Swaminathan *et al*., 2008,

Zhuang *et al*., 2008; Licausi *et al*., 2010, Licausi *et al*., 2013). A member of AP2/DREB type

TF, RAP2.4 was induced in response to salt and drought stress (Feng *et al*., 2005). WOUND

INDUCED DEDIFFERENTIATION 1–4 (WIND1-4), a member of RAP2.2, are induced on

wounding and play important roles in callus formation at wounded sites (Iwase *et al*., 2011).

These proteins integrate multiple phytohormone signals and regulate abiotic and biotic stress

responses (Gutterson and Reuber, 2004). ERF binds with the ETHYLENE-RESPONSE

ELEMENT (ERE) or GCC-box to provide biotic stress resistance (Franco-Zorrilla *et al*.,

2014).

Different properties of AP2/ERFs, including the DNA binding properties and induction upon

various stresses, implement these TFs to coordinate multiple responses and participate in

regulatory processes (Abiri *et al*., 2017). It is evident that AP2/ERFs and various

phytohormones such as ethylene, Methyl jasmonate (MeJA), ABA, SA act in concert to

regulate various plant processes (Licausi *et al*., 2013). The coordinated action of ABA and

different abiotic stresses leads to the activation of several stress-inducible genes and DREB2s

(Lee *et al*., 2016). AP2/ERFs such as ABA-INSENSITIVE 4 (ABI4) and CBFA (CCAAT

binding factor A) are ABA-responsive and help to activate the ABA-dependent/independent

stress-responsive genes (Zhang *et al*., 2013). AP2/ERF mutants with altered hormone

sensitivity and abiotic stress responses have been identified and studied, mentioning this

family of TFs as an important candidate for studying the interactions between plant hormones

and abiotic stresses.

The ERF subfamily protein is usually conscious of pathogen attack and contributes to plant

immunity. ERF subfamily protein's overexpression is generally related to altered disease

resistance phenotypes in plants (Onate-Sanchez *et al*., 2007; Zhang *et al*., 2016; Zheng *et al*.,

2019). Overexpression of rice *OsEREBP1* activates ABA and JA signalling pathways,

eventually resulting in enhanced tolerance under biotic and abiotic conditions (Jisha *et al*.,

2015). ABA regulates various important agronomical traits of plant development and

physiology, such as seed dormancy and maturation, along with responses to several
environmental stress factors, such as salinity, drought, and cold stress mediated through
different components of ABA signalling (Himmelbach *et al*., 2003; Sah *et al*., 2016). During
the investigation of gene expression regulated with ABA, an AP2-domain containing a
protein known as *ABR1* that regulates ABA responses was identified (Pandey *et al*., 2005).
Later, it was studied that *ABR1* (group X of AP2/ERF) acts as a transcriptional activator and
is involved in the wounding response (Baumler *et al*., 2019). *ABR1* is an important molecule
thought to play a functional role in the host-pathogen interface and acts as a susceptibility
core that binds with different effector molecules of *Pseudomonas syringae* (Schreiber *et al*.,
2021). Despite the importance of *ABR1* in regulating biotic stress responses, its
transcriptional regulator is unknown. In our study, we have identified and characterized
AtERF60 (homologue of MaRAP2 from *Mentha arvensis*) (Phukan *et al*., 2018). We
investigated that *AtERF60* regulates the *ABR1* gene by binding with their promoter and
modulates biotic stress response in Arabidopsis.

Materials and Methods

Plant growth conditions and stress treatment

Arabidopsis Col-0 ecotype was utilized in this study. The plants were grown in the Conviron
growth chamber model A-1000 as described in Phukan *et al*. (2018). The T-DNA insertion
mutant lines were obtained from the Arabidopsis Biological Resource Centre (ABRC) and
screened for homozygous lines. For growing, the Arabidopsis seeds were surface sterilized
(3% sodium hypochlorite solution), stratified at 4^0C for 96 hours, and then transferred to the
growth chamber under controlled conditions with photoperiod (16:8 h light-dark) at 22°C and
60% relative humidity. MeJA treatment was provided by spraying a 100 μM solution made in
dimethyl sulfoxide (DMSO) and Triton-X. Control plants were sprayed with a solution
containing only DMSO and Triton- X. For SA and ABA treatment, plants were sprayed with
100 μM of SA and ABA, while control plants were sprayed with water. The aerial part,
including the stem and leaves, was punctured with a needle to avoid major injuries. To avoid
contamination, treated samples were obtained at various periods and cleaned thoroughly with
sterile water.

Expression analysis and phylogenetic analysis

Total RNA was isolated from plant samples utilizing RNeasy Plant Mini Kit (Qiagen), and

cDNA was prepared utilizing a high-capacity cDNA reverse transcription kit (Applied

Biosystems). Total cDNA was checked for quality by PCR using the control primers

provided in the kit. The qRT-PCR (Applied biosystems 7900-HT Fast Real-Time PCR) was

performed to determine the expression level of transcripts utilizing the SYBR Green PCR

master mix kit (Takara). The unique region of transcripts was selected for designing qRT-

PCR primers and mentioned in Table S3. Actin and ubiquitin were used as endogenous

controls for normalizing gene expression. The $2^{-\Delta\Delta Ct}$ method was utilized to calculate the

relative expression of genes. The protein sequences of all the AP2/ERF family members

related to Arabidopsis were downloaded from the TAIR database. The sequence alignment

was performed by using clustal omega online program using default parameters

(https://www.ebi.ac.uk/Tools/msa/clustalo/), and the phylogenetic tree was constructed using

MEGAXI using the neighbour-joining method, with 1000 bootstrap replicates.

Cloning experiments, generation of transgenic lines, and mutant screening

The *AtERF60* was amplified using gene-specific primers and cloned in the PTZ57R/T vector.

To analyze the in vitro interaction of protein and DNA, *AtERF60* was cloned in pGEX-4-T2

expression vector fused with GST. The positive clones were confirmed by PCR and digestion

with primer-specific restriction enzymes (*BamHI* and *Xho*I) (Fig. S2a). The resulting positive

construct was transformed in Lon and OmpT protease deficient *E. coli BL21* (DE3) strain for

expression in the bacterial system. To study the functional role of *AtERF60* in planta, we

cloned it in the pBI121 binary vector downstream of the CaMV35S promoter. The positive

clones were confirmed by PCR, sequencing and final digestion utilizing the specific

restriction enzymes (*BamHI* and *Xba*I) (Fig. S3a). The *AtERF60* transgenic lines in

Arabidopsis were raised using the *Agrobacterium*-mediated gene transformation (Fig. S3b).

Four independent transgenic lines containing *AtERF60* were generated, and two lines

showing similar responses were used in the study. Transgenic lines in Arabidopsis were

confirmed by using genomic DNA PCR using pBI-121 nptII (Kan[R]) specific primers. The

PCR amplification ensures the successful integration of the desired gene (Fig. S3d). To

determine the functional role of the *AtERF60*, we used T-DNA mutant lines of *AtERF60*

(SALK_138492). The *erf60* mutants were screened using genomic DNA PCR with specific

primers designed from the left border of T-DNA (Fig. S4).

Electrophoretic mobility shift assay

To study the in vitro protein-DNA interaction, *AtERF60* was cloned in the bacterial expression vector pGEX-4-T2. The open reading frame was continued with GST without disturbing the amino acid sequences. The cloned construct was then transformed in Lon and OmpT protease deficient *E. coli BL21* (DE3) strain and induced with 0.4 mM IPTG at 37°C for 5 h. The recombinant fusion protein was purified with Glutathione Sepharose beads (Sigma). Definite probes were designed for different *cis*-elements and their mutated versions (Table S3). EMSA followed the manufacturer's protocol mentioned in the 2nd generation DIG gel shift kit (Roche).

Microarray analysis

The complete RNA was extracted from the plant samples (WT, *ERF60*-OX, and *erf60* mutant) under controlled conditions utilizing the RNeasy Plant Mini Kit (Qiagen) following the manufacturer's instructions. The Arabidopsis GXP 4x44K AMADID slide was used to hybridize the RNA, which was further aided and labelled with Cy3-CTP. The microarray was performed and scanned at 535 nm, and the images were analyzed using Agilent Feature Extraction software (v10.7) to calculate signal and background strength. The images obtained from the microarray were cleaned and are uniformly intense, with very little background noise. Gene-Spring GX 12.6 software was utilized for statistical significance and normalization. The fold induction values obtained from two different lines were normalized to a single fold induction. The student's t-test was used to correct P-values for the downregulation and upregulation of genes in experimental and biological replicates. The P-value cut-off for gene up- and downregulation was set to <0.05.

Bacterial pathogen assays with *Pseudomonas syringae* pv. tomato DC3000 (*Pst*DC3000)

Bacterial pathogen *Pst*DC3000 was grown at 28°C overnight on LB media containing 50μg/ml rifampicin. For infiltration, bacteria were re-suspended in 10 mM MgCl$_2$ to obtain OD$_{600}$ = 0.2. The leaves of 5-week-old WT, *AtERF60*-OX, *erf60* mutant, and *AtERF60*-complementation Arabidopsis plants were syringe infiltrated with *Pst*DC3000. Disease symptoms were observed and photographed at regular intervals. Bacterial populations [Log (CFU/cm^2)] in leaf tissues of WT, *AtERF60*-OX, *erf60* and *AtERF60*-complementation plants at 0, 2, and 4 dpi, were determined according to Katagiri *et al*. (2002). Four leaf discs of

equal area (0.5 cm2) were taken from each WT, *AtERF60*-OX, *erf60* and *AtERF60*-
complementation plants inoculated with *Pst*DC3000 at 4 dpi for electrolyte leakage assay.
These leaf discs were then agitated in a tube containing 10 ml milli-Q water for 3 h. The
conductivity was measured using the electrical conductivity meter for each sample (HORIBA
Scientific, F74BW). Afterwards, leaf discs for each sample were autoclaved to release the
total ions, and the conductivity corresponding to total ions was measured. Electrolyte leakage
values (conductivity value at 3 h) were presented as the percentage relative to total ions.

Pathogen colonization in Arabidopsis plants was visualized using a Zeiss confocal laser-
scanning microscope (LSM-880). We used GFP-tagged *Pst*DC3000 for this assay. Confocal
images of leaf samples of WT, *AtERF60*-OX, *erf60* and *AtERF60*-complementation plants
inoculated with GFP-tagged *Pst*DC3000 were taken at 2 dpi. GFP acquisition was performed
at 488 nm excitation with emission collection at 493-598 nm. Image processing was
performed using Zeiss application software. Leaf samples from Arabidopsis plants were
collected at 6 hpi (hours post infiltration) to examine the induction of selected differentially
expressed genes in response to the *Pst*DC3000 challenge using the qRT-PCR assay. The
leaves infiltrated with 10 mM $MgCl_2$ were taken as control. Actin and ubiquitin were used as
endogenous controls for gene normalization.

Statistical analysis and R packages used

The experiments in the study were performed in two independent biological replicates, each
with three technical repeats. The data shown in the study are mean $\pm$ SD. Student t-test was
performed using instat.exe version 3.0 software to measure the degree of significance with a
P-value $\leq$ 0.05 and denoted by an asterisk above the bar graph in the figures. P-values $\leq$
0.001 are designated with three asterisks while p values $\leq$ 0.0001are designated with four
asterisks. For GSEA Go enrichment analysis, R cluster Profiler and enrich plot package were
used. For the ridge plot, the r ggridges package was used. The Tidyverse package was used to
make a volcano, dot plot, and bar plots. For heatmap, the pheatmap package was used.

Results

AtERF60 **is induced under salt, dehydration, ABA and SA treatments**

The *AtERF60* CDS is 819 bp long and encodes a 273 amino acid polypeptide. The alignment
of complete ORF with its amino acid sequence has shown that it has 58 amino acids

conserved AP2/ERF DNA binding domain (DBD) (Fig. S1). The phylogenetic homology of

AtERF60 was investigated using MEGAXI software with the neighbour-joining method and

1000 bootstrap repetitions, and it was discovered that it is closely related to Arabidopsis

ERF056. (Fig. 1a). Chromosomal location and mutant information used in the study are

depicted in (Fig.1 b). To study *AtERF60* under various conditions, we checked the induction

of *AtERF60* in response to abscisic acid (ABA), salt, dehydration, (SA) salicylic acid,

wounding and methyl jasmonate (MeJA) treatments in *Arabidopsis thaliana* Col-0 plants. We

observed that *AtERF60* is induced early after 1 hour of ABA treatment. *AtERF60* showed

maximum induction after 3 hours of salt and dehydration treatment. Transcripts of *AtERF60*

showed a similar level of expression after 1, 3 and 5 hours of SA treatment (Fig. 1c).

Additionally, *AtERF60* is not induced in response to MeJA and wounding treatment as

compared to the untreated control WT Col-0 (Fig. 1c).

Recombinant AtERF60 interacts with DRE/GCC box *cis*-elements

The recombinant GST-tagged-AtERF60 was induced and purified in the bacterial expression

system (Fig. S2b, c). An expected molecular mass of 56.0 kDa along with GST-tagged-

AtERF60 was observed and separated using SDS-PAGE (Fig. S2b). To study the DNA

binding property of *AtERF60*, we performed EMSA of recombinant AtERF60 with DRE-1

and GCC-box *cis*-elements. Specific probes were designed for DRE-1 (GCCGAC) and GCC-

box (AGCCGCC) *cis*-element along with their mutated forms containing a single nucleotide

substitution (Fig. 1d). We found that the recombinant AtERF60 interacts with DRE-1 and

GCC-box *cis*-elements, while it does not bind with its mutated probe carrying a change of

single nucleotide (Fig. 1e). The EMSA demonstrate that *AtERF60* TF interacts specifically

with DRE-1 and GCC-box *cis*-elements and might be involved in the regulation of

downstream target genes.

***AtERF60* regulates basal resistance response in Arabidopsis**

ABR1   is   a   functionally   important   molecule   that   interacts   with

different *Pseudomonas syringae* effectors (Schreiber *et al*., 2021). We sought to test our idea

that *AtERF60* regulates the central key component *ABR1* and modulates plant response to

biotic stress. We generated independent homozygous lines by constitutively overexpressing

AtERF60 under *CaMV 35S* to investigate its role in response to biotic stress (Fig S3 a, b, c d)

and validated the overexpression lines (Fig 2a). Additionally, separate homozygous lines were

generated by complementing the *erf60* mutant lines (Fig 2a). The *AtERF60*-OX, *erf60* mutant,

and *AtERF60* complemented lines were infected with *Pseudomonas syringae* pv. *tomato* (*Pst*)

strain DC3000 to examine the bacterial pathogen response. The *AtERF60*-OX lines exhibited

reduced chlorosis and necrosis at 4 days of post-inoculation (dpi), whereas *erf60* mutant lines

displayed increased disease symptoms as compared to the Col-0 plants. Furthermore,

the *erf60* mutant lines supported a significantly higher population of bacterial *Pst*DC3000

than the Col-0 plants. The bacterial population in the *AtERF60*- OX lines was reduced by 0.93

log (CFU/cm^2) at 2 dpi while it was reduced by 1.34 log (CFU/cm^2), after 4 dpi (Fig. 2c). The

erf60 lines, on the other hand, have a 0.61 log (CFU/cm2) and 0.70 log (CFU/cm2) increase in

bacterial population at 2 and 4 dpi, respectively (Fig. 2c). A significant reduction in

electrolyte leakage was observed in *AtERF60*-OX lines (~16.04 %) compared to the Col-0

plants at 4 dpi with *Pst*DC3000 (Fig. 2d). Contrary to that, electrolyte leakage was found to be

significantly higher in *erf60* lines (~ 29.43 %) compared to the Col-0 (Fig. 2c) plants. No

significant difference in the bacterial population and electrolyte leakage was observed in the

AtERF60 complemented lines compared to Col-0. Further, we used confocal microscopy to

observe the colonization of *Pst*DC3000 in Arabidopsis plants using the GFP-

tagged *Pst*DC3000 strain. In vivo imaging revealed the formation of the distinct

microcolonies of GFP-tagged *Pst*DC3000 that were evenly dispersed in the intercellular

spaces of leaves of Col-0 plants. We observed reduced colonization of GFP-

tagged *Pst*DC3000 in *AtERF60*- OX lines while *erf60* lines displayed enhanced colonization

of bacterial pathogen (Fig. 2b). Together, these results suggest that mutation in

the *AtERF60* gene causes an increase in susceptibility of Arabidopsis towards bacterial

pathogen, *Pst*DC3000.

Loss of *AtERF60* function affects multiple gene expression

To understand the regulatory role of *AtERF60* in Arabidopsis, we performed a microarray

analysis of *AtERF60*-OX and *erf60* mutant lines under control conditions in two independent

biological replicates (Table S1). We observed a significant differential expression gene

(DEG) change in mutants and OX lines. There are 1527 DEGs upregulated and 1050 DEGs

downregulated in OX lines with p-value >0.05 and 2-fold change increase/decrease, while

1226 DEGs are upregulated and 773 are down-regulated in mutant lines. (Fig. 3a, b). Further,

these DEGs are categorised into function GO categories. The categories are differentiated

based on whether they are activated or suppressed in biological processes, cellular

components and molecular function. The top 10 enriched categories in OX and mutant lines

are shown (Fig. 3c, e). Photosynthesis-associated pathways are seen enriched in, and protein

quality control pathways are suppressed in OX lines while Sugar pathways are enriched in

mutant and transport pathways are suppressed. To interpret up/down-regulated pathways

grouped by gene set, density plots are generated using the frequency of fold change values

per gene within each set of top enriched categories in OX and mutant lines (Fig. 3d, f). We

further validated the expression of 13- above identified genes using qRT-PCR analysis in

both *AtERF60*-OX *erf60* and its complemented lines as compared to the Col-0 (Fig. 4). We

found that most of the genes encoding COMPROMIZED RECOGNITION OF TCV1

(CRT1), *ABR1*, Peroxidase superfamily protein, Aspartyl protease family protein, O-glucosyl

hydrolases family protein, Alcohol dehydrogenase 1 (ADH1), Acyl-CoA synthetase 5

(ACS5), and EIDI like 3, were significantly upregulated in the *erf60* mutant background as

compared to the Col-0. Complementation of *AtERF60* in a mutant background restored the

increased expression of these downregulated transcripts to their wild type levels, indicating

that *AtERF60* suppresses the expression of these genes (Fig 4). The microarray and qRT-PCR

analysis revealed that *AtERF60* might be involved in downregulating these target genes

directly or indirectly by binding with their promoter *cis*-elements. Therefore, we screened out

the individual promoters and searched for probable *cis*-elements (Supplementary file 1).

***AtERF60* interacts with *ABR1* promoter**

Sequence analysis revealed the presence of dehydration (A/GCCGAC) and ABA-responsive

cis-elements (ACGTC/G) in the *ABR1* promoter. Based on the presence of these core *cis*-

elements in the *ABR1* promoter, we studied the interaction of *AtERF60* with the *ABR1*

promoter using EMSA. Probes specific to DRE and ABRE *cis*-elements were designed along

with their mutated counterpart with a change in a single nucleotide (Fig. 5a). It was observed

that *AtERF60* interacts with both DRE-1 and DRE-2 *cis*-elements (Fig. 5b). At the same time,

it is not able to produce gel shift with the ABRE *cis*-elements and its counterpart with a

single nucleotide mutation (Data not shown). This showed that the binding of *AtERF60* is

specific to A/GCCGAC *cis*-elements and is completely abolished when the sequence-specific

nucleotide is mutated.

We examined the expression of *AtERF60* and *ABR1* genes in Arabidopsis plants following
infection with *Pst*DC3000. The expression of the *AtERF60* and *ABR1* genes are induced in
Col-0 in response to *Pst*DC3000 infection as observed at 6 hours post-inoculation (hpi) (Fig
5c, d). On the other hand, *ADH1, ACS, CHS*, and *ABR1* genes show induced expression in the
erf60 mutant plants, followed by a *Pst*DC3000 infection of 6 hours. The expressions of these
transcripts were downregulated in the *AtERF60*-OX lines after 6 hours of *Pst*DC300 infection
(Fig 5d-g). When *AtERF60* is complemented into *erf60* mutant lines, it causes a reversal in
the expression of these transcripts to the Col-0 level. Moreover, after being challenged with 6
hours of *Pst*DC3000, the *erf60* mutant plants show a considerable increase in *ABR1*
expression, while the *AtERF60*-OX plants show a significant downregulation. (Fig. 5c-g).
Thus, our data suggest a negative regulation of *ABR1* by *AtERF60* under biotic stress
response.

Discussion

AP2/ERFs regulate various downstream signalling cascades by interacting with different *cis*-
elements of target gene promoters and regulating multiple responses (Phukan *et al*., 2018). In
this study, we have identified a transcription factor, *AtERF60*, induced in response to drought
conditions, salt stress, ABA/SA treatment, and bacterial pathogen *Pst*DC3000 infection.
Induction of the *AtERF60* in response to ABA and SA suggests that the *AtERF60* might
regulate   downstream   responses in   an   ABA/SA-dependent
manner. In Arabidopsis, *ERF53* (Hsieh *et al*., 2013), *RAP2.6* (Zhu *et al*., 2010), and *ERF-*
VIIs (Yao *et al*., 2017) are induced in response to ABA to upregulate the genes containing
DRE/ABRE *cis*-elements. There are limited numbers of AP2/ERFs reported to be involved in
response to SA treatment (Xie *et al*., 2019). The potential of AP2/ERFs to counter different
signals and regulate multiple stresses facilitates them to make a highly complex stress
regulatory network. Some AP2/ERFs are induced at earlier stages, while others are induced
upon prolonged stress, which suggests that their function might be influenced mutually (Van
den Broeck *et al*., 2017).

The diversity of AP2/ERFs in response to various stresses depends on the flexibility of the
AP2 domain that implements the binding to different *cis*-elements such as DRE/CRT and
GCC-box elements (Huang *et al*., 2008, Cheng *et al*. 2013, Franco-Zorrilla *et al*., 2014,
Catinot *et al*., 2015). Our previous study conclusively demonstrated that AP2/ERF TFs such

as *PsAP2* and *MaRAP2-4* from the *Opium poppy* and *Mentha arvensis*, respectively, bind

with both GCC and DRE box *cis*-elements to modulate biotic and abiotic stress response in

transgenic plants (Mishra *et al.*, 2015; Phukan *et al.*, 2018). We observed that it interacts with

the core *cis*-element of DRE (GCCGAC), and the binding was abolished when we used the

mutated version by changing the 4th nucleotide G to T (GCC**T**AC). We also observed that

AtERF60 interacts with the core GCC box element GCCGCC, and mutation of the probe into

GCCACC abolished its interaction. These results suggest that CCGCC and CCGAC are the

core sequences preferred for *AtERF60* to interact. If we mutate G of CCGCC to CCACC, the

binding was abolished. We find that the mutation in the last two CC to AC for the DRE

sequence does not affect its binding ability. Some ERFs, including *ERF71*, *ERF4*, and *ERF1*,

are also reported to interact with both DRE and GCC elements in Arabidopsis (Lee *et al.*,

2015; Xie *et al.*, 2019). The ability of *AtERF60* to interact specifically with DRE and GCC

further supports its probable involvement in stress response.

Additional insight into *AtERF60* function in biotic stress response was gained by processing

and analysing the microarray and expression data in both the overexpression and mutant

background. Microarray analysis, promoter binding activity, and phenotypic studies suggest

the regulatory role of *AtERF60* in Arabidopsis. The fold expressions of the majority of

differentially expressed transcripts are significantly higher in the *erf60* mutant background

under control. Additionally, we identified the upregulation of target genes in the *erf60* mutant

plants following *Pst*DC3000 inoculation, including *ABR1, chalcone synthase gene*, *Acyl CoA*

synthase 5, and *ADH1*.

In this study we discovered that, the mutation in *AtERF60* enhanced susceptibility against

bacterial pathogen *Pst*DC3000. In contrast, *AtERF60*-OX plants exhibited reduced

susceptibility to *Pst*DC3000. This work identifies *AtERF60* as the pathogen-responsive gene

which downregulates *ABR1,* which contributes to plant immunity by interacting with

Pseudomonas syringae effector molecule HopZ1a and other different effectors (Schreiber *et*

al., 2021). Increased expression of *ABR1* in the *erf60* mutant plants in response to the

pathogen infection suggests that *AtERF60* downregultes the *ABR1* gene expression during

biotic stress. Under similar conditions, *AtERF60*-OX plants show a suppression of *ABR1*

expression. Our results suggest that *AtERF60* plays an important role in plant bacterial-

pathogen interaction via downregulating *ABR1* expression under the control conditions. The

AtERF60-OX lines exhibited reduced susceptibility, and its functional mutant showed an

enhanced susceptibility response. Our results conclude that mutation in *AtERF60* makes
susceptibility hub *ABR1* constitutively active, leading to susceptible phenotype towards
bacterial pathogen *Pst*DC3000.

Acknowledgements

The authors acknowledge the Director, CSIR-CIMAP, for providing the necessary facilities.
Authors acknowledge ABRC for providing the mutant seeds of *AtERF60*. We also
acknowledge Genotypic for performing the microarray analysis of overexpression and mutant
background of *AtERF60*.

Funding

RKS acknowledge CSIR-CIMAP and SERB for financial support to his lab. GSJ, UJP, AJ,
NS, DP and SK acknowledge CSIR-UGC for fellowship. AP is grateful to the DST -
INSPIRE Faculty Award (IFA13-LSPA-20).

Author contributions

AJ has generated the complementation lines, contributed to the bacterial pathogen infiltration
experiments and CFU analysis, and performed electrolyte leakage; UJP has cloned the
AtERF60 gene made the over-expression lines and screened the *erf60* mutant lines, giving
samples for microarray analysis and contributed to analysis of figure 3. NS has contributed to
the bacterial pathogen infiltration experiments and CFU analysis and electrolyte leakage,
maintained the seed stock and plating of seeds. GSJ has performed EMSA experiments. SK
has completed validation of q-RT-PCR of *AtERF60-OX*, *erf60* and its complementation lines
under the control condition and after bacterial infiltration. DP has contributed to bacterial
CFU analysis. VT has contributed to planning experiments related to mutant background and
procuring of the mutant background of *AtERF60*. AP has supervised the bacterial colony
infiltration. AJ, SK, NS, and RKS has analyzed the result, RKS has supervised the work,
planned the experiments and edited the manuscript. AJ, SK, NS, and RKS have finalized the
draft manuscript.

Data Availability Statement

ERF60- AT4G39780, *ADH 1-* AT1G77120, *ACOS5-* AT1G62940, *Chalcone and stilbene*
synthase- AT4G34850, *CYP450 family protein-* AT1G69500, *Aspartyl protease family*
protein- AT5G24820, *Actin 1-* AT2G37620, *Acyl-transferase family protein-* AT1G03390,
EID1-like 3- AT3G63060, *ABR1-* AT5G64750, *CRT1-* AT4G36290, *alpha/beta-Hydrolases*
superfamily protein- AT1G73480, *O-Glycosyl hydrolases family 17 protein-* AT4G14080,
Myb-like HTH transcriptional regulator family protein- AT1G18960, *UDP-*
Glycosyltransferase superfamily protein- AT1G10400, *RING/U-box superfamily protein-*
AT1G49220, Microarray data deposited with the accession number GSE179600.

Conflict of interest

The authors declare no conflict of interest.

References

Abiri R, Shaharuddin NA, Maziah M, Yusof ZNB, Atabaki N, Sahebi M, *et al*. **2017**
Role of ethylene and the APETALA 2/ethylene response factor superfamily in rice under
various abiotic and biotic stress conditions. *Environ. Exp. Bot* 134: **33**–44.
Baumler J, Riber W, Klecker M, Müller L, Dissmeyer N, Weig AR, *et al*. **2019**
AtERF111/*ABR1* is a transcriptional activator involved in the wounding response. *Plant J*
100 (5): 969-990.
Catinot J, Huang JB, Huang PY, Tseng MY, Chen YL, Gu SY, *et al*. **2015** ETHYLENE
RESPONSE FACTOR 96 positively regulates Arabidopsis resistance to necrotrophic
pathogens by direct binding to GCC elements of jasmonate and ethylene responsive defence
genes. *Plant, cell & environment* **38** (12): 2721–2734.
Causier B, Ashworth M, Guo W, Davies B 2012 The TOPLESS interactome: a framework
for gene repression in Arabidopsis. *Plant Physiol* **158**: 423–438.
Cheng MC, Liao PM, Kuo WW, Lin TP 2013 The Arabidopsis ETHYLENE RESPONSE
FACTOR1 regulates abiotic stress-responsive gene expression by binding to different cis-
acting elements in response to different stress signals. *Plant Physiology* **162:** 1566–1582.
Feng JX, Liu D, Pan Y, Gong W, Ma LG, Luo JC, *et al*. **2005.** An annotation update via
cDNA sequence analysis and comprehensive profiling of developmental, hormonal or
environmental responsiveness of the *Arabidopsis* AP2/EREBP transcription factor gene
family. *Plant Mol. Biol.* **59**: 853–868.

Franco-Zorrilla JM, Lopez-Vidriero I, Carrasco JL, Godoy M, Vera P, Solano R 2014
DNA-binding specificities of plant transcription factors and their potential to define target
genes. *Proc. Natl. Acad. Sci. U.S.A.* **111**: 2367–2372.

Gutterson N, Reuber TL 2004 Regulation of disease resistance pathways by AP2/ERF-
transcription factors. *Curr. Opin. Plant Biol* **7**: 465–471.

He M, He CQ, Ding NZ 2018 Abiotic Stresses: General Defenses of Land Plants and
Chances for Engineering Multistress Tolerance. *Frontiers in plant science* **9**: 1771.

Himmelbach A, Yang Y, Grill E 2003 Relay and control of abscisic acid signaling. ***Curr***
Opin Plant Biol **6**: 470–479.

Huang D, Wu W, Abrams SR, Cutler AJ 2008 The relationship of drought-related gene
expression in Arabidopsis thaliana to hormonal and environmental factors. *J. Exp. Bot* **59**:
2991–3007.

Hsieh EJ, Cheng MC, Lin TP 2013 Functional characterization of an abiotic stress-
inducible transcription factor AtERF53 in Arabidopsis thaliana. *Plant Mol. Biol* **82**: 223–237.

Iwase A, Harashima H, Ikeuchi M, Rymen B, Ohnuma M, Komaki S, *et al*. 2017 WIND1
promotes shoot regeneration through transcrip- tional activation of ENHANCER OF SHOOT
REGENERATION1 in Arabidopsis. *Plant Cell* **29**: 54–69.

Jisha V, Dampanaboina L, Vadassery J, Mithofer A, Kappara S, Ramanan R 2015
Overexpression of an AP2/ERF Type transcription factor OsEREBP1 confers biotic and
abiotic stress tolerance in rice. *PLoS ONE 10*: e0127831.

Kagale S, Rozwadowski K 2011 EAR motif-mediated transcriptional repression in plants:
an underlying mechanism for epigenetic regulation of gene expression. *Epigenetics 6*: 141–
146.

Katagari F, Thlimony R, He SY 2002 The Arabidopsis *thaliana- Pseudomonas*
syringae interaction. Arabidopsis Book 1: e0039

Lee DK, Jung H, Jang G, Jeong JS, Kim YS, Ha SH, *et al*. 2016 Overexpression of the
OsERF71 transcription factor alters rice root structure and drought resistance. *Plant Physiol*
172(1): 575–588.

Lee SY, Boon NJ, Webb AA, Tanaka RJ 2016 Synergistic Activation of RD29A via
integration of salinity stress and abscisic acid in Arabidopsis thaliana. *Plant Cell Physiol* **57:**
2147–2160.

Lee SY, Hwang EY, Seok HY, Tarte VN, Jeong MS, Jang SB, *et al*. 2015 Arabidopsis
AtERF71/HRE2 functions as transcriptional activator via cis-acting GCC box or DRE/CRT

element and is involved in root development through regulation of root cell expansion. *Plant Cell Rep* **34**: 223–231.

Licausi F, Giorgi F, Zenoni S, Osti F, Pezzotti M, Perata P 2010 Genomic and transcriptomic analysis of the AP2/ERF superfamily in Vitis vinifera. *BMC Genomics* **11**: 719.

Licausi F, Ohme-Takagi M, Perata P 2013 APETALA2/Ethylene Responsive Factor (AP2/ERF) transcription factors: mediators of stress responses and developmental programs. *New Phytol* **199**: 639–649.

Liu JH, Peng T Dai W 2014 Critical cis-Acting Elements and Interacting Transcription Factors: Key Players Associated with Abiotic Stress Responses in Plants. *Plant Mol Biol Rep* **32**: 303–317.

Mishra S, Phukan UJ, Tripathi V, Singh DK, Luqman S, Shukla RK 2015 PsAP2 an AP2/ERF family transcription factor from papaver somniferum enhances abiotic and biotic stress tolerance in transgenic tobacco. *Plant Mol. Biol* **89**: 173–186.

Nakano T, Suzuki K, Fujimura T, Shinshi H 2006 Genome-wide analysis of the ERF gene family in Arabidopsis and rice. *Plant Physiol* **140**: 411–432.

Oñate-Sánchez L, Anderson JP, Young J, Singh KB 2007 AtERF14, a member of the ERF family of transcription factors, plays a nonredundant role in plant defense. *Plant physiology* **143**(1): 400–409.

Pandey GK, Grant JJ, Cheong YH, Kim BG, Li L, Luan S 2005 *ABR1*, an APETALA2-domain transcription factor that functions as a repressor of ABA response in Arabidopsis. *Plant Physiol* **139**(3): 1185-93.

Phukan UJ, Jeena GS, Tripathi V, Shukla RK 2018 MaRAP2-4, a waterlogging responsive ERF from Mentha, regulates bidirectional sugar transporter AtSWEET10 to modulate stress response in Arabidopsis. *Plant Biotechnol J* **16**(1): 221-233.

Phukan UJ, Jeena GS, Triptahi V, Shukla RK 2017 Regulation of Apetala2/Ethylene Response *Frontiers in Plant Science* **8**: 150.

Sah SK, Reddy KR, Li J 2016 Abscisic Acid and Abiotic Stress Tolerance in Crop Plants. *Frontiers in plant science* **7**: 571.

Schreiber KJ, Hassan JA, Lewis JD 2021 Arabidopsis Abscisic Acid Repressor 1 is a susceptibility hub that interacts with multiple Pseudomonas syringae effectors. *The Plant journal*: for cell and molecular biology **105**(5): 1274–1292.

Shigyo M, Ito M 2004 Analysis of gymnosperm two-AP2-domain-containinggenes.
Development Genes and Evolution **214**: 105–114.

Song CP, Agarwal M, Ohta M, Guo Y, Halfter U, Wang P, *et al.* **2005.** Role of an
Arabidopsis AP2/EREBP-type transcriptional repressor in abscisic acid and drought stress
responses. *Plant Cell* **17**: 2384–2396.

Swaminathan K, Peterson K, Jack T 2008 The plant B3 superfamily. *Trends in Plant*
Science **13**: 647–655.

Van den Broeck L, Dubois M, Vermeersch M, Storme V, Matsui M, Inze D 2017 From
network to phenotype: the dynamic wiring of an Arabidopsis transcriptional network induced
by osmotic stress. *Mol. Syst. Biol* **13**: 961.

Xie Z, Nolan T, Jiang H, *et al.* **2019** The AP2/ERF Transcription factor TINY modulates
brassinosteroid-regulated plant growth and drought responses in Arabidopsis. *Plant Cell*
31(8):1788–1806.

Xie Z, Nolan TM, Jiang H, Yin Y 2019 AP2/ERF Transcription Factor Regulatory
Networks in Hormone and Abiotic Stress Responses in Arabidopsis. *Frontiers in plant*
science **10**: 228.

Yao Y, Chen X, Wu AM 2017 ERF-VII members exhibit synergistic and separate roles in
Arabidopsis. *Plant Signal. Behav* **12**: e1329073.

Zhang H, Hong Y, Huang L, Li D, Song F 2016 Arabidopsis AtERF014 acts as a dual
regulator that differentially modulates immunity against Pseudomonas syringae pv. tomato
and Botrytis cinerea. *Sci. Rep* **6**: 1–15.

Zhang P, Berardini TZ, Ebert D, Li Q, Mi H, Muruganujan A, Prithvi T, Reiser L,
Sawant S, Thomas PD, Huala E 2020 PhyloGenes: An online phylogenetics and functional
genomics resource for plant gene function inference. *Plant Direct.* **4**(12):e00293.

Zhang ZW, Feng LY, Cheng J, Tang H, Xu F, Zhu F, *et al.* **2013** The roles of two
transcription factors, ABI4 and CBFA, in ABA and plastid signalling and stress responses.
Plant Mol. Biol **83**: 445–458.

Zheng X, Xing J, Zhang K, Pang X, Zhao Y, Wang G, *et al.* **2019** Ethylene response
factor ERF11 activates BT4 transcription to regulate immunity to Pseudomonas syringae.
Plant Physiol **180**: 1132–1151.

Zhu Q, Zhang J, Gao X, Tong J, Xiao L, Li W, *et al.* **2010** The Arabidopsis AP2/ERF
transcription factor RAP2.6 participates in ABA, salt and osmotic stress responses. *Gene* **457**:
1–12.

Zhuang J, Cai B, Peng RH, Zhu B, Jin XF, Xue Y, *et al*. 2008 Genome-wide analysis of

the AP2/ERF gene family in Populus trichocarpa. *Biochem Biophys Res Commun* **371**(3):

468–474.

Figure legends

Figure 1. Phylogeny, expression analysis, and Electrophoretic mobility shift assay

(EMSA) of *AtERF60*. (a) The protein sequences of 122 AP2/ERF family members from

Arabidopsis thaliana were retrieved from the TAIR database. The amino acid sequences were

aligned by clustalW and the phylogenetic tree was constructed using MEGAXI by neighbour-

joining (NJ) method and 1000 bootstrap replicates. **(b)** Gene structure of *AtERF60***,** its

chromosome location and T-DNA insertion line**. (c)** Relative expression of *AtERF60*

transcript in response to ABA, salt, dehydration, SA, wounding, and MeJA treatment after 1,

3, and 5 h. Relative expression of transcripts was calculated by taking untreated plant samples

as a control (WT). Actin and ubiquitin were used as endogenous controls for gene

normalisation. Error bars indicate mean ± SD. Student's t-test, **, $P < 0.01$. **(d)** Probes

containing DRE (GCCGAC) and GCC-box (AGCCGCC) *cis*-elements were designed to

study the DNA-protein interaction. The desired *cis*-elements are red, while the binding site

carrying mutations are marked blue. **(e)** EMSA of *AtERF60* showed that it interacts

specifically with DRE-1 and GCC regulatory *cis*-elements (M-mutated/substituted and L-

DIG-labelled). Error bars indicate mean ± SD. Student's t-test, **, $P < 0.01$.

Figure 2. Evaluation of *Pst*DC3000 colonization in *AtERF60* overexpressing, mutant

and complemented lines. (a) Relative expression of *AtERF60* transcripts in overexpression

lines (OX), *erf60* mutant lines and *AtERF60* complementation lines compared to Col-0

plants. Actin and ubiquitin were used as endogenous controls for gene normalization. The

error-bars represents SE, *, $P < 0.05$; **, $P < 0.01$. **(b)** Differential interference contrast

(DIC) and confocal images of leaves of WT, *AtERF60-OX* and *erf60* mutant lines with GFP-

tagged *Pst*DC3000 at 2 dpi. GFP fluorescence (green) was acquired using 488 nm excitation

and 493-598 nm emission. Images of DIC and confocal microscopy (scale bar, 100 µm).

(c) Bacterial growth estimation in leaves of WT, *AtERF60*-OX, *erf60* mutant and *AtERF60*

complemented lines inoculated with *Pst*DC3000 at 0, 2, and 4 dpi. **(d)** Electrolyte leakage

from leaves of WT, *AtERF60*-OE *erf60* mutant and *AtERF60* complemented lines inoculated

with *Pst*DC3000 at 4 dpi. Electrolyte leakage values are given as the percentage of total ions.

Figure 3. Microarray analysis of *AtERF60*-OX and *erf60* mutant plants under controlled
conditions. (a) For the *AtERF60* overexpression lines (OX), univariate analysis of DEGs in
volcano plot with negative logarithm of the p value on the y-axis and logarithm of the fold
change on the x-axis. Red dots show upregulated genes with fold change ≥ 2, and blue dots
show down-regulated genes with fold change ≤ 2. The red threshold line shows a p-value of
0.05. The grey dots are non-significant DEGs. The subplot shows the number of up-and-
down-regulated genes. **(b)** For the mutant line, volcano plot and bar plot of up-and down-
regulated DEGs. **(c)** Enrichment plot (dot plot) of OX lines with GO categories up and down-
regulated in biological processes (BP), cellular component (CC) and molecular function (MF).
(d) Ridge plot (density plot) of OX lines summarizing the "core enriched" genes for each gene
set via (log2) fold-change. The colour describes the adjusted p-value listed in the colour bar,
notifying whether an enriched term is significant. **(e)** Enrichment dot plot and **(f)** ridge plot of
mutant lines showing GO categories and core enriched genes.

Figure 4. Validation of differentially expressed target genes between *AtERF60*-OX and
***erf60* mutant through qRT-PCR.** Relative expression of differential target genes as
determined by qRT-PCR in the *AtERF60*-OX, *erf60* mutant and *AtERF60* complemented
lines compared to the WT. 13 differentially expressed genes were selected for validation
through qRT-PCR. Actin and ubiquitin were used as endogenous controls for gene
normalization. Error bars indicate mean $\pm$ SD. Student's t-test, *, $P < 0.05$; **, $P < 0.01$.

Figure 5. EMSA analysis to determine binding ability of *AtERF60*. (a) Probes containing
DRE1/2 (A/GCCGAC) and ABRE1/2 (ACGTC/G) *cis*-elements were designed to study the
DNA-protein interaction. The desired *cis*-elements are red, while the binding site containing
mutations is blue. **(b)** EMSA of *AtERF60* showed that it interacts with the regulatory DRE1
and DRE2 cis-elements in the *ABR1* promoter. In contrast, it does not interact with the
ABRE1/2 *cis*-elements (M-mutated/substituted and L-DIG-labelled). **Validation of**
differentially expressed target genes between *AtERF60*-OX and *erf60* mutant under
biotic stress. qRT-PCR analysis of (c) *ERF60,* (d) *ABR1,* (e) *chalcone synthase,* (f) *ADH1*
and (g) *ACS* genes in Col-0, OX and mutant lines following infection with *Pst* DC3000 after
6 hours post-inoculation (hpi) relative to Col-0 infiltrated with 10mM $MgCl_2$ (mock).

Supplementary data

Figure S1. Nucleotide and amino acid alignment of *AtERF60*. Complete cDNA (1224 bp)
carrying 819 bp of ORF. Red represents the amino acid sequence, and blue represents the
AP2/ERF DNA binding domain. *AtERF60* has a 145 bp 5' UTR and 259 bp of 3' UTR. It
encodes a protein of 273 amino acids having a molecular weight of 30 kDa.

Figure S2. Protein induction and purification of *AtERF60*. (a) Confirmation of positive
clones of *AtERF60* by PCR and restriction digestion in bacterial expression vector pGEX-4-
T2. **(b)** The recombinant AtERF60 protein was induced with different concentrations (mM)
of IPTG at 37^0C. **(c)** The AtERF60-GST fusion protein was purified from the *E. coli* BL21
(DE3) strain following induction for 5 hours at 37^0C with 0.3 mM IPTG. An affinity-purified
recombinant protein with a molecular weight of 56.0 kDa was segregated on SDS-PAGE.

Figure S3. Cloning and validation of transgenic lines of *AtERF60*. (a) Confirmation of
positive clones of *AtERF60* by PCR and restriction digestion in plant expression vector
pBI121. **(b)** Selection of transgenic lines on kanamycin-supplemented half MS media. **(c)**
PCR confirmation of transgenic lines. Actin was used as an internal control. **(d)** Genomic
DNA PCR of transgenic lines with NPT II (KanR) primers.

Figure S4. Mutant screening of *AtERF60* lines. Genomic DNA PCR of mutant *AtERF60*
Salk lines. LB- left border, RP- right primer, LP- left primer. The resulting amplification
using primers specific LB and RP showed mutant homozygous lines.

Supplementary file 1. List of selected Arabidopsis promoter sequences used in the study.

Table S1. Microarray data showing the list of up and downregulated genes obtained.

Table S2. List of genes identified from microarray analysis with significant p-value (<0.05).

Table S3. List of primers used in the study.

Fig. 1

Fig. 2

(a)

(b)

(c)

(d)

Fig. 3

Fig. 4

Fig. 5

(a)

DRE-1: AGA**GCCGAC**AGAT**GCCGAC**TG**GCCGAC**CT

DRE-1 (M): AGA**GCCTAC**AGAT**GCCTAC**TG**GCCTAC**CT

DRE-2: AGA**ACCGAC**AGAT**ACCGAC**TG**GCCGCCC**T

DRE-2 (M): AGA**ACCTAC**AGAT**ACCTAC**TG**ACCTAC**CT

(b)

(c)

(d)

(e)

(f)

(g)