Age-associated B cells are long-lasting effectors that restrain reactivation of latent

γHV68

Isobel C. Mouat[1,2], Iryna Shanina[2], Marc S. Horwitz[2]

[1] Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK

[2] Department of Microbiology and Immunology, The University of British Columbia, Vancouver,

British Columbia, Canada

Corresponding author:

Marc S. Horwitz

Room 3551, Life Sciences Centre

2350 Health Sciences Mall

University of British Columbia

Vancouver, B. C. Canada V6T 1Z3

604-822-6298

mhorwitz@mail.ubc.ca

Abstract

Age-associated B cells (ABCs; $CD19^+CD11c^+T-bet^+$) are a unique population that are increased in

an array of viral infections, though their role during latent infection is largely unexplored. Here,

we use murine gammaherpesvirus 68 (γHV68) to demonstrate that ABCs remain elevated long-

term during latent infection and express IFNγ and TNF. Using a strain of γHV68 that is cleared

following acute infection, we show that ABCs persist in the absence of latent virus, though their

expression of IFNγ and TNF is decreased. With a fluorescent virus we demonstrate that ABCs

are infected with γHV68 at similar rates to other previously activated B cells. We find that mice

without ABCs display defects in anti-viral IgG2a/c antibodies and are more susceptible to γHV68

reactivation when challenged with heterologous infection. Together, these results indicate that

ABCs are a persistent effector subset during latent viral infection that restrains γHV68

reactivation.

Conflict of interest statement

The authors have declared that no conflict of interest exists.

Introduction

Humans are infected with an array of herpesviruses that persist within us throughout our lives and require continuous surveillance by the host immune system. Gammaherpesvirus-68 (γHV68), like other herpesviruses, deploys distinct transcriptional programs during lytic and latent infection, each of which requires a distinct immune response[1]. Age-associated B cells (ABCs) are a unique B cell population implicated in viral infection, autoimmunity, and aging[2–5]. ABCs (CD19+CD11c+T-bet+) are induced following γHV68 infection[6], though their function throughout the course of infection is not clear. In this study we examine the response of and role for ABCs throughout γHV68 infection, from acute infection through long-term latency.

ABCs were identified in 2011 in the context of female aging and autoimmunity[2,3] and have since been shown to be increased in an array of viral infections including lymphocytic choriomeningitis virus (LCMV), murine cytomegalovirus, γHV68, vaccinia, human immunodeficiency virus, rhinovirus, SARS-CoV2, and influenza[6–11]. ABCs are elevated in the spleen and circulation during active viral infections and persist primarily in the spleen during chronic infection or upon infection resolution[8,9]. ABCs display multiple functional capacities including the secretion of antibodies and anti-viral cytokines and activation of T cells[12,13]. T-bet expression in B cells is required for IgG2a/c class switching[14] and lack of ABCs exacerbates LCMV chronic infection in mice[15]. We predict that ABCs could be playing a role throughout γHV68 infection due to their long-term persistence, activation of T cells, and continuous cytokine and antibody production.

Lytic γHV68 infection is primarily cleared by CD8[+] T cells[16,17], following which γHV68 persists in a

latent state mainly in previously activated B cells[18]. During latency the immune response

persists even though there is minimal production of viral genes and limited production of new

virions[19]. Both humoral and cellular responses contribute to controlling γHV68 infection long-

term[20,21]. Mice lacking either antibodies or T cells are both able to control γHV68 latency,

though mice lacking both antibodies and T cells display elevated γHV68 reactivation[20,22]. IFNγ is

critical for controlling γHV68 replication and reactivation from latency[23,24].

B cells are also known to be important during latent γHV68 infection. B cells act as the primary

latent viral reservoir, are required for movement from the lung to the spleen at the onset of

latency, and mice deficient in B cells are unable to develop latency following intranasal

infection[18,25–27]. B cells secrete anti-viral antibodies long term[28], but also may inhibit

reactivation via secretion of anti-viral cytokines or help to virus-specific T cells. The precise B

cell subsets, mechanisms, and factors that facilitate the maintenance of latency and prevent

reactivation are not fully understood.

The role of ABCs during acute and latent herpesvirus infection is unknown. Herein, using

various tools, we investigate the relationship between γHV68 and ABCs throughout lytic and

latent infection. We find that ABCs expand during acute γHV68 infection and persist in the

spleen during latency. Well after the initial establishment of latency, ABCs continuously secrete

anti-viral cytokines and antibodies and in the face of heterologous challenge, ABCs are required

to restrain latent γHV68 reactivation. Thus, our findings highlight a novel role for ABCs as

effector cells during γHV68 latency.

Results

ABCs are induced and maintained following γHV68 infection in a sex-biased manner

To examine the relative proportion of ABCs during acute and latent γHV68 infection, C57BL/6(J)

mice were mock-infected with media (naïve) or infected with γHV68 for 6, 35, and 150 days,

blood and spleen were collected and analyzed by flow cytometry to measure ABCs. A gating

scheme, described in Figure 1A, was used to identify CD11c$^+$T-bet$^+$ as a proportion of previously

activated B cells (CD19$^+$IgD$^-$). We observed the relative proportion of ABCs increased in

circulation during acute γHV68 infection (6 days p.i.) compared to naïve mice (Figure 1B-C).

Once latency was well established (day 35and 150 p.i.), the proportion of circulating ABCs

decreased to pre-infection levels (Figure 1B-C). This indicates that ABCs circulate in response to

the lytic γHV68 infection but do not remain elevated in circulation during latency. In the spleen,

the relative proportion and total number of ABCs was increased during acute γHV68 infection

(day 6 p.i.) relative to naïve mice and remained elevated during latent γHV68 infection at 35

days p.i. and 150 days p.i. (Figure 1D-E, Figure S1). These results support previous findings that

the spleen is the major site for ABCs following the clearance of active viral infection[9].

Intriguingly, we observed that the proportion of ABCs is increased in females as compared to

males in both the blood and spleen throughout infection (Figure 1F-M). Previously, we

demonstrated that ABCs display a sex bias during experimental autoimmune encephalomyelitis

and at 35 days post-γHV68 infection in the spleen[29]. Here, we have extended this finding and

shown that ABCs display female sex bias in both the circulation and spleen during lytic and

latent γHV68 infection. While ABCs increase in both females and males following γHV68

infection, there is a sex bias leading to greater expansion in females. These results demonstrate

that ABCs are increased in circulation during acute γHV68 infection and continue to endure in

the spleen long-term during latent infection.

ABCs express anti-viral cytokines in response to latent γHV68 infection

To begin investigating the role of ABCs in the anti-viral response, we measured ABC expression

of anti-viral cytokines interferon-γ (IFNγ) and tumor necrosis factor-α (TNF) in the spleen at 6,

35, and 150 days p.i.. We found that during acute γHV68 infection (6 days p.i.), roughly 40% of

ABCs in the spleen expressed IFNγ and TNF, as compared to about 10% of ABCs in naïve mice

(Figure 2A-B). During latent infection (35 days and 150 days p.i.), a significantly increased

proportion of ABCs continued to express IFNγ and TNF compared to naïve mice, though the

proportion of ABCs expressing IFNγ and TNF was decreased compared to acute infection (Figure

2A-B). Throughout the acute and latent infection, we observed a downregulation of IL-17A

expression on ABCs (Figure 2C).

To determine if the level of cytokine expression was distinct from other B cells, we compared

the proportion of ABCs expressing IFNγ and TNF to non-ABC B cells. Specifically, we examined

expression on ABCs compared to CD11c⁻Tbet⁺ and CD11c⁻Tbet⁻ B cells 35 days post-γHV68

infection (Figure 2D). A significantly increased proportion of ABCs expressed IFNγ and TNF

compared to non-ABC B cells, both CD11c⁻Tbet⁺ and CD11c⁻Tbet⁻ B cells, in the spleen (Figure

2E). The mean fluorescent intensity (MFI) of IFNγ and TNF, of IFNγ or TNF positive cells, was

significantly increased on ABCs compared to both non-ABC populations (Figure S2A, B). CD11c⁻

Tbet⁺ B cells expressed an intermediate level of IFNγ and TNF between ABCs and CD11c⁻Tbet⁻ B

cells (Figure 2E, Figure S2A, B). Sex differences in ABC cytokine expression were not observed

(Figure 2A-C, Figure S3 A-C). The high level of IFNγ and TNF cytokine expression indicates that

ABCs may be functioning in a unique anti-viral capacity during latent infection.

To explore the relationship between latent γHV68 and the ABC population, we infected mice

with ACRTA-γHV68, a recombinant strain of γHV68 in which the genes responsible for latency

are deleted and a lytic gene, RTA, is constitutively expressed[30]. As a result, γHV68 infection is

cleared following acute infection without ever establishing latency. We found that, at 35 days

p.i., a time point at which ACRTA-γHV68 is cleared, ABCs are increased to the same level in the

spleens of ACRTA-γHV68 infected mice as compared to spleens in mice infected with WT γHV68

(Figure 2F). Thus, ABCs persist in the absence of latent virus, indicating that ABCs act similarly to

memory cells that remain following acute infection.

While the proportion of ABCs remained elevated in the absence of latent virus, cytokine

expression was altered. The proportion of ABCs in ACRTA-γHV68-infected mice expressing IFNγ

and TNF at 35 days p.i. was significantly reduced compared to those infected with WT γHV68

while remaining significantly elevated compared to naïve mice (Figure 2G-H). Additionally, the

IFNγ$^+$ ABCs in mice infected with ACRTA-γHV68 displayed a lower MFI than those from γHV68-

infected mice, though the TNF MFI was not significantly different between the two groups

(Figure S2C, D). Without the presence of the latent virus, we observe a decrease specifically in

the proportion of ABCs that doubly express both IFNγ and TNF as well as an increase in ABCs

that are negative for the expression of either IFNγ and TNF as compared to ABCs from WT

γHV68-infected mice (Figure 2I). This finding indicates that ABCs respond to the latent virus by

producing both IFNγ and TNF.

Together, these results show that ABCs are a predominant B cell subset expressing anti-viral

cytokines IFNγ and TNF during latent γHV68. Further, our data establish that high levels of

expression of these cytokines are dependent on the presence of the latent virus. These findings

suggest that ABCs recognize the presence of the quiescent latent virus and respond by

expressing anti-viral cytokines.

ABCs are susceptible to γHV68 infection but are not a major viral reservoir

γHV68 is known to infect B cells, in particular germinal centre and memory B cells[31–34]. To

determine if ABCs are directly infected by γHV68, we used a previously developed fluorescent

strain of γHV68, γHV68.H2bYFP[35] where fluorescence is easily detectable during acute infection,

but over time falls off during latent infection. Mice were infected with γHV68.H2bYFP and 8

days p.i. spleens were collected, and flow cytometry was performed. We found that, during

acute infection, a small proportion of ABCs were infected with γHV68. The same proportion of

ABCs were positive for the fluorescent virus (1.2 ± 0.3%) as observed in previously activated B

cells (1.1 ± 0.2%, Figure 3A-B), demonstrating that ABCs are not preferably targeted for

infection. To our knowledge this is the first evidence of a virus directly infecting ABCs. We next

asked what proportion of the γHV68 reservoir is made up of ABCs. The primary reservoir for

γHV68 is previously activated B cells which aligns with our results that 73% of γHV68-infected

cells are IgD⁻ B cells (Figure 3C). We found that ABCs make up only 6% of γHV68-infected cells

(Figure 3C). Other infected cell populations included innate cells such as macrophages and DCs,

in alignment with previous findings[31], as well as a small proportion of NK and T cells (Figure 3C).

We also examined the relationship between infection of ABCs and expression of IFNγ. We

observed that during acute infection the proportion of ABCs expressing IFNγ was significantly

lower in ABCs infected with γHV68 (4.3±1.3 %) as compared to uninfected ABCs (32.2±1.0%,

Figure 3D). These results indicate that direct infection of ABCs was not driving IFNγ production

and that ABCs are susceptible to γHV68-driven downregulation of IFNγ. These findings

demonstrate that although a portion of ABCs are infected with γHV68, ABCs are not the major

target for γHV68.

ABCs are dispensable for the control of acute infection and establishment of latency

To further examine the role(s) of ABCs in γHV68 infection, mice with a floxed B cell specific T-

bet deletion were followed post-infection. Specifically, $Tbx21^{fl/fl}Cd19^{cre/+}$ (KO) and littermate

$Tbx21^{fl/fl}Cd19^{+/+}$ (Ctrl) mice were infected with γHV68 for 6 or 35 days and the spleen was

collected to examine the quantity of γHV68 by qPCR and the immune cell composition (Figure

4A). Mice were genotyped by PCR and loss of ABCs in KO mice was confirmed by flow cytometry

(Figure 4B). No differences in clinical symptoms or weight changes were observed during γHV68

infection between Ctrl or KO mice (Figure S4A).

To determine if knocking out ABCs alters immune cell composition, Ctrl and KO mice were

infected or mock-infected with γHV68 and spleens analyzed at 6 or 35 days by flow cytometry

to examine various T cell, B cell, and innate immune cell populations. No difference in the total

number of splenocytes was observed (Figure S4B). The composition of the splenic immune

profile was similar between Ctrl and KO mice with no significant differences observed in the

relative proportions of B cells, CD8 T cells, DCs, neutrophils, NK cells, or macrophages between

mock-infected mice or those infected with γHV68 for 6 or 35 days, though KO mice had greater

proportions of CD4 T cells than Ctrl mice 35 days post-infection (Figure S5).

We then measured the quantity of γHV68 in the spleens of Ctrl and KO mice infected for 6 and

35 days by qPCR. We found no difference in that the quantity of γHV68 at 6 and 35 days p.i.

between Ctrl and KO mice (Figure 4C-D) and the quantity of γHV68 did not differ between male

and female mice (Figure S4C-D). To confirm that KO mice effectively control the lytic infection

and develop latency, we infected Ctrl and KO mice with latency-deficient ACRTA-γHV68 for 35

days. We found that KO mice, like Ctrl mice, displayed similar viral loads that were near or

below the limit of detection when infected with ACRTA-γHV68 (Figure 4E). As no virus was

detected at day 35, mice lacking ABCs were effectively controlling the lytic infection.

Furthermore, no differences were observed in the relative expression of viral genes associated

with lytic infection (*Orf50*, *Orf68*) or latent infection (*Orf73*) between Ctrl and KO mice (Figure

4F). *Orf50* encodes the replication and transcription activator protein, which initiates viral lytic

gene expression[36]. *Orf68* encodes a packaging protein that assists in moving the newly

replicated viral genomes to the packaging motor, where they are loaded into capsids[37]. *Orf73*

encodes for the latency-associated nuclear antigen that is required for the establishment and

maintenance of latent infection[38,39]. As such, equal expression of lytic and latency-associated

genes *Orf50*, *Orf68,* and *Orf73* between Ctrl and KO mice suggests that Ctrl and KO mice harbor

similar levels of lytic and latent γHV68. Together, these results demonstrate that ABCs are not

required for the clearance of acute infection and establishment of latency in steady-state

conditions.

Lack of ABCs results in a dysregulated γHV68 antibody response but does not alter the

viral reservoir or anti-viral T cell response

ABCs are known to secrete anti-viral IgG2a/c[6], and transfer of serum into mice without ABCs

during LCMV infection was partially able to restore control of the infection[15]. To examine the

antibody response in Ctrl versus KO mice, mice were infected with γHV68 for 35 days and sera

was collected. The levels of anti-γHV68 IgG were the identical between Ctrl and KO mice,

though Th1-associated IgG2c was significantly decreased in KO mice, while Th2-associated IgG1

was significantly elevated, when compared to Ctrl mice (Figure 5A-C). These results suggest that

ABCs are the primary secreters of anti-γHV68 IgG2, though in their absence there is

compensation by other B cell subsets resulting in increased anti-γHV68 IgG1 antibodies.

We next asked if the γHV68 reservoir was altered in mice without ABCs, as a portion of γHV68-

infected cells were ABCs. In particular, we posited that an altered viral reservoir could impact

the ability of KO mice to control γHV68 latency; previous findings indicate that various cell types

infected by γHV68 may display differential susceptibility to reactivation[23,40]. To compare the

viral reservoir in Ctrl and KO mice, we infected mice with a fluorescent strain, γHV68.H2bYFP,

and, at 8 days p.i., examined immune cell populations that comprise the γHV68-infected

population. We observed no difference in the proportion of infected cell populations, including

T cells, DCs, macrophages, NK cells, and naïve and previously activated B cells (Figure 5D). This

suggests that the cell populations infected with γHV68 are not substantially altered in KO mice

compared to Ctrl mice, making it unlikely that changes to the reservoir impacted susceptibility

to viral reactivation.

To ask whether ABCs influence anti-viral immune cells in the spleen, we next examined three

cell populations previously shown to be important for control of latent γHV68: IFNγ-producing

cells, γHV68-specific CD8$^+$ T cells, and Vβ4$^+$CD8$^+$ T cells. IFNγ is present at low levels during

latency[41] and is critical for controlling γHV68 reactivation from latency[23,24,42]. In particular, IFNγ-

producing T cells are known to block γHV68 reactivation[23,43–45]. CD8$^+$ T cells that harbor the Vβ4

TCR were also examined. CD8$^+$Vβ4$^+$ T cells expand following γHV68 infection and reach their

highest levels during latency, wherein they persist throughout infection without taking on an

exhausted phenotype[46–48]. γHV68-specific CD8$^+$ T cells were also measured with tetramers to

p79, an immunodominant γHV68 epitope for which CD8-specific memory T cells remain

throughout latent infection[49]. We observed that the proportion of CD4$^+$ and CD8$^+$ T cells that

express IFNγ were unchanged between mice with and without ABCs (Figure 5E-F). Further, no

differences were observed in the proportion of either Vβ4$^+$CD8$^+$ T cells or γHV68 p79-specific

CD8[+] T cells between mice with and without ABC circulating populations (Figure 5G-H).

Additionally, we found that the proportion of NK cells and B cells that express IFNγ, and the MFI

of IFNγ on these populations, was not changed between Ctrl and KO mice whether mock-

infected or infected for 6 or 35 days (Figure S6). That a similar proportion of B cells expressed

IFNγ indicates that another B cell subset likely compensates for the loss of IFNγ-expressing

ABCs. Collectively, these findings indicate that ABCs are not acting to stimulate anti-viral

immune cell populations in the spleen.

Together, these data indicate that knocking out ABCs leads to dysregulation of the γHV68

antibody response, without altering the γHV68 reservoir or T cell populations responding to the

latent virus.

Susceptibility to γHV68 reactivation increases in the absence of ABCs

We posited that the persistence of ABCs long-term during γHV68 and their secretion of IFNγ,

TNF, and anti-viral antibodies suggest that ABCs have an important role in the suppression of

γHV68 reactivation. To examine this role, ex vivo reactivation assays were performed on

splenocytes from Ctrl and KO mice at 35 days p.i.. Splenocytes from γHV68-infected KO mice

reactivated more frequently in culture than those from Ctrl mice (Figure 6A). This finding

indicates that there is an increased propensity for γHV68 reactivation in the absence of ABCs,

although the assay does not disentangle whether infected cells have an altered cell-intrinsic

susceptibility to reactivation or if there is a cell-extrinsic influence on reactivation.

To further interrogate the role of ABCs in suppressing γHV68 reactivation, we next asked if mice

without ABCs are more susceptible to γHV68 reactivation following infectious challenge.

Heterologous viral infection was performed as a physiological mimic to reactivate latent γHV68.

Importantly, we chose viruses that do not typically reactivate γHV68,[50] LCMV and coxsackievirus

B4 (CVB4). Specifically, KO and Ctrl mice were infected with γHV68 for 35 days and then

challenged with LCMV or CVB4 (Figure 6B). No clinical symptoms were observed during either

of the challenges in Ctrl or KO mice and there was no difference in the relative quantity of

LCMV and CVB4 in the spleens of Ctrl versus KO mice (Figure 6C-D). Following challenge with

LCMV or CVB4, mice lacking ABCs had elevated quantities of γHV68 in the spleen compared to

mice with ABCs (Figure 6E-F). We reasoned that the increased γHV68 load following challenge

may be due to increased reactivation and, in support, observed increased relative expression of

two lytic-associated γHV68 genes, *Orf50* and *Orf68*, in KO mice compared to Ctrl mice following

challenge with the heterologous viruses (Figure 6G-H). Alternatively, we did not observe a

difference in expression of the latency-associated gene *Orf73*, between Ctrl and KO mice

(Figure 6I). However, it remains possible that the increased quantity of γHV68 may result from

expansion of infected B cells in response to the challenge, rather than reactivation. Collectively,

these findings indicate that mice lacking ABCs may be more susceptible to γHV68 reactivation in

the face of heterologous challenge.

Discussion

Here we have shown that ABCs are increased during acute γHV68 infection and persist during

latency for at least 150 days. Previous studies have shown that ABCs persist following clearance

of viral infections[9] and during chronic infection[15], though their contribution during a latent

infection was unexplored. Our results demonstrate a novel role for ABCs: they continuously

respond in an effector manner to latent viral infection by secreting anti-viral cytokines and

antibodies. Further, our results indicate that ABCs may play an important role in suppressing

γHV68 reactivation during subsequent heterologous infectious challenges.

Our results show that ABCs continuously expresses anti-viral cytokines during latency and the

presence of the latent virus is required for the sustained anti-viral cytokine expression by ABCs.

Compared to other B cells, ABCs display uniquely high expression of anti-viral cytokines IFNγ

and TNF during γHV68. In addition to expressing cytokines, ABCs also produce anti-viral

antibodies, as deficiency of ABCs results in a significant loss of γHV68-specific IgG2c antibodies.

We hypothesized that the production of anti-viral cytokines and antibodies by ABCs could be a

way in which ABCs restrain γHV68 reactivation. Using mice deficient in ABCs, we showed that

ABCs are important for the suppression of γHV68 reactivation in the face of heterologous viral

challenge.

A surprising finding of our study is that ABCs are increased more so in female than male mice

throughout γHV68 infection, in both the blood and spleen. Sex-differences are well-

documented in anti-viral immune responses[51] and ABCs have previously been shown to display

a female sex bias in contexts of aging and autoimmunity[3,52]. It has recently been shown that the

ABC female sex bias in lupus mice is abolished following the duplication of *Tlr7* in male mice[52].

While we show that there is no difference in the quantity of γHV68 between female and male

mice, TLR7 stimulation is known to be important for ABC differentiation[2,3] and TLR7 is critical

for the control of lytic γHV68 infection and maintenance of latency[53]. While increased

frequencies of ABCs were observed in female mice compared to males, we found that a similar

proportion of the ABCs express anti-viral cytokines in both males and female mice. No

differences between male and female mice were observed in viral load or reactivation.

Using a parallel approach to knockout T-bet in B cells with bone marrow chimeras, it was

previously shown that mice lacking ABCs display an increased quantity of γHV68 at day 14 p.i.

compared to controls[6]. This inconsistency with our results is most easily explained by a clear

difference in the inoculating dose (lower herein) and the method of ABC knockout. Additionally,

the two studies measure mice at different timepoints (day 14 p.i., the establishment of latency

versus day 35 p.i., steady-state latency) and this likely hints at potential differences in the

kinetics of latency establishment in mice lacking ABCs.

ABCs are known to possess an array of functional capacities and the precise mechanism(s) of

ABC contribution to the restraint of γHV68 reactivation will be the focus of future studies. Here

we have shown that only the antibody response is altered in mice deficient in ABCs and

unexpectedly, changes are not observed in the anti-viral T cell response or viral reservoir. The

ability of ABCs to continuously secrete anti-viral cytokines during latent infection is an

additional mechanism that likely contributes to the suppression of γHV68 reactivation.

Understanding the precise localization of ABCs in relation to γHV68-infected B cells will aid in

our understanding of the development of splenic structures, in particular germinal centres, as

ABCs have been shown to play a role in their development[54].

Whether ABCs play a role in suppressing the reactivation of human gamma-herpesvirus

infections such as Epstein-Barr virus (EBV) and Kaposi sarcoma-associated herpesvirus (KSHV), is

not known. Understanding this role would have important relevance in diseases associated with

these human gammaherpesviruses, including various malignancies, myalgic

encephalomyelitis/chronic fatigue syndrome, and chronic-active EBV[55–57].

This work demonstrates that ABCs are a long-lasting effector population during γHV68 latency

and suggests that they are playing a unique role in the suppression of viral reactivation.

Methods

Mice

Tbx21[fl/fl]*Cd19*[cre/+] mice were generated by crossing *Tbx21*[fl/fl]*Cd19*[cre/+] and *Tbx21*[fl/fl]*Cd19*[+/+] mice.

Tbx21[fl/fl] and *Cd19*[cre/+] mice were provided by Dr. Pippa Marrack[54]. C57BL/6(J) mice were

originally purchased from The Jackson Laboratory. All animals were bred and maintained in the

animal facility at the University of British Columbia. All animal work was performed per

regulations of the Canadian Council for Animal Care (Protocols A17- 0105, A17-0184).

γHV68, ACRTA-γHV68, and γHV68H2B.YFP infection

γHV68 WUMS strain (ATCC), ACRTA-γHV68 (developed by Dr. Ting-Ting Wu, gift of Dr. Marcia A.

Blackman)[30], and γHV68H2B.YFP (developed and provided by Dr. Samuel H. Speck) were

propagated in Baby Hamster Kidney cells (BHK, ATCC). Viruses were diluted in Minimum

Essential Media (MEM) prior to infection and maintained on ice. 6- to 8-week-old mice were

infected i.p. with 10^4 PFU of γHV68, ACRTA-γHV68, γHV68H2B.YFP, or mock-infected with MEM.

Clinical symptoms were not observed during γHV68, ACRTA-γHV68, or γHV68H2B.YFP infections

in C57BL/6(J) nor *Tbx21*[fl/fl]*Cd19*[cre/+] or *Tbx21*[fl/fl]*Cd19*[+/+] mice.

LCMV infection

LCMV Armstrong strain 53b (originally acquired from Dr. M.B. Oldstone) was propagated on

BHK cells. Prior to infection, virus was diluted in RPMI-1640 media (Gibco) and maintained on

ice. Mice (11- to 13-week-old) were infected i.p. with 2 x 10^5 PFU LCMV or mock infected with

media. No clinical symptoms were observed from LCMV infection in $Tbx21^{fl/fl}Cd19^{cre/+}$ nor

$Tbx21^{fl/fl}Cd19^{+/+}$ mice.

CVB4 infection

CVB4 Edward strain was propagated on HeLa cells and titred by plaque assay. Virus was diluted

in Dulbecco's Modified Eagle Medium (DMEM, Gibco) and maintained on ice prior to infection.

Mice (11- to -13-week-old) were infected i.p. with 100 PFU CBV4. No clinical symptoms were

observed from CVB4 infection in $Tbx21^{fl/fl}Cd19^{cre/+}$ or $Tbx21^{fl/fl}Cd19^{+/+}$ mice.

Tissue harvesting and processing for flow cytometry

Mice were anesthetised with isoflurane and euthanized by cardiac puncture. For flow

cytometry, blood was collected by cardiac puncture into 100 ul 0.5 M

Ethylenediaminetetraacetic acid (EDTA) to prevent clotting and placed on ice until processing.

Spleen extracted and placed into 2 ml PBS and kept on ice until processing. Blood was

incubated in 10 ml warmed (37°C) ACK lysis buffer for 15 minutes at room temperature to

remove red blood cells and washed twice with FACS buffer. Spleens were mashed through a 70

μm cell strainer with a 3 ml syringe insert to make a single cell suspension for each sample.

Splenocytes were incubated in 4 ml warmed (37°C) ACK lysis buffer for 10 minutes on ice to

lyse red blood cells and remaining cells were resuspended in FACS buffer and kept on ice until

further use.

Flow cytometry analysis of cell-type specific surface antigens and intracellular cytokines

For analysis of extracellular and intracellular antigens, 2 million cells per spleen sample or all

cells collected per blood sample were stained with appropriate antibodies. Prior to staining,

samples were incubated at 4°C covered from light for 30 minutes with 2 μl/ml Fixable Viability

Dye eFluor506 (Thermo Fisher) while in PBS and then resuspended in rat anti-mouse CD16/32

(Fc block, BD Biosciences) antibody for 10 minutes. Then, samples were stained with

fluorochrome labeled antibodies (Table 1) against cell surface antigens for 30 minutes covered

from light at 4°C, washed, and resuspended in Fix/Perm buffer (Thermo Fisher) for 30 minutes-

12 hours while maintained covered from light at 4°C. Samples were then washed twice with

perm buffer and incubated 40 minutes with antibodies for intracellular antigens (Table 1) in

Perm buffer at covered from light at RT. Cells were then washed and resuspended in FACS

buffer with 2 mM EDTA. For analysis of cytokine production, 4 million cells per sample were

stimulated ex vivo for 3 hours at 5% CO_2 at 37°C in Minimum Essential Media (Gibco)

containing 10% fetal bovine serum (FBS, Sigma-Aldrich), 1 μl/ml GolgiPlug (BD Biosciences), 10

ng/ml PMA (Sigma-Aldrich) and 500 ng/ml ionomycin (Thermo Fisher) and washed prior to

staining. Samples were collected on an Attune NxT Flow Cytometer (Thermo Fisher) and

analyzed with FlowJo software v10 (FlowJo LLC).

Antigen	Fluor	Clone	Dilution	Company
CD3e	eFluor 450	500A2	1:200	Thermo Fisher Scientific
CD19	PE	1D3	1:200	Thermo Fisher Scientific
CD19	Pe-Cy7	1D3	1:200	Thermo Fisher Scientific
CD19	SB780	1D3	1:100	Thermo Fisher Scientific
IgD	Pe-Cy7	11-26	1:200	Thermo Fisher Scientific
CD11c	Alexafluor700	N418	1:100	Thermo Fisher Scientific
CD11c	APC	N418	1:100	Biolegend
T-bet	PerCpCy5.5	4B10	1:100	Thermo Fisher Scientific
IFNγ	APC	XMG1.2	1:100	Thermo Fisher Scientific

TNF	Pe-Cy7	MP6-XT22	1:100	Biolegend
IL-17A	PEDazzle594	TC11-18H10.1	1:100	Biolegend
CD45	Alexafluor700	30-F11	1:200	Thermo Fisher Scientific
CD45	APC-eFluor 780	30-F11	1:200	Thermo Fisher Scientific
CD4	SB645	RM4-5	1:100	Thermo Fisher Scientific
CD8	APC-eFluor 780	53-6.7	1:200	Thermo Fisher Scientific
CD8	PEDazzle594	53-6.7	1:100	Biolegend
CD11b	Pe-Cy7	M1/70	1:100	Thermo Fisher Scientific
Ly6G	PerCpCy5.5	RB6-8C5	1:100	Thermo Fisher Scientific
CD335	FitC	29A1.4	1:100	Thermo Fisher Scientific
NK1.1	PE	PK136	1:100	Thermo Fisher Scientific
F4/80	APC-eFluor 780	BM8	1:100	Thermo Fisher Scientific
CD62L	PerCpCy5.5	MEL-14	1:100	Thermo Fisher Scientific
CD44	Alexafluor700	IM7	1:200	Thermo Fisher Scientific
PD1	PerCpeFluor 710	RMP1-30	1:100	Thermo Fisher Scientific
CXCR5	PE	SPRCL5	1:100	Thermo Fisher Scientific
FoxP3	Alexafluor700	FJK-16s	1:100	Thermo Fisher Scientific
CD138	PEDazzle594	281-2	1:100	Biolegend
CD95	FitC	SA367H8	1:100	Biolegend
IgM	APCCy7	RMM-1	1:100	Biolegend
TCR Vβ4	FitC	KT4	1:100	BD Bioscience

Table 1 Flow cytometry antibodies

Tetramer staining

γHV68-specific CD8 T cells were identified by staining with a tetramer acquired from the NIH

Tetramer Facility. 2 million cells per well were incubated for 1 hour at RT with p79

(K^b/ORF61$_{524-531}$ TSINFVKI, diluted 1:400) following viability staining and Fc receptor block, as

described above, but before extracellular antigen staining. Uninfected mice stained with

tetramers used as negative gating controls.

γHV68 qPCR

DNA was isolated from 4 x 10^6 splenocytes using PureLink™ Genomic DNA Mini Kit (Thermo

Fisher) and quantified with a spectrophotometer. qPCR was performed using 2x QuantiNova

Probe Mastermix (Qiagen, USA) on the Bio-Rad CFX96 Touch™ Real Time PCR Detection system.

Copies of γHV68 were quantified in duplicate wells with 150 ng DNA per reaction using primers

and probes specific to γHV68 *Orf50* and mouse *Ptger2* (Table 2). Standard curves were obtained

by serial dilutions of *Orf50* and *Ptger2* gBlocks (*Orf50*: $2x10^6 - 2x10^1$; *Ptger2*: $5x10^7$-$5x10^2$) that

were amplified in parallel.

433
434

Gene	Sequence
Ptger2 Forward Primer	5⦸-TACCTTCAGCTGTACGCCAC-3⦸
Ptger2 Reverse Primer	5⦸-GCCAGGAGAATGAGGTGGTC-3⦸
Ptger2 Probe	5⦸-/56-FAM/CCTGCTGCT/ZEN/TATCGTGGCTG/3IABkFQ/-3⦸
Orf50 Forward Primer	5⦸-TGGACTTTGACAGCCCAGTA-3⦸
Orf50 Reverse Primer	5⦸- TCCCTTGAGGCAAATGATTC-3⦸
Orf50 Probe	5⦸-/56-FAM/TGACAGTGC/ZEN/CTATGGCCAAGTCTTG/3 IABkFQ/-3⦸

Table 2 γHV68 qPCR primers and probe sequences

RT-qPCR for γHV68 lytic and latency-associated genes and relative quantity of LCMV and CVB4

Portion of spleen stabilized in RNAlater immediately following collection and stored at -80°C for

up to 12 weeks. RNA extracted with RNeasy mini kit (Qiagen, cat no. 74104) and cDNA

immediately synthesized with High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher, cat

no. 4368814). Reaction performed with 500ng per reaction in duplicate wells. cDNA quantified

with a spectrophotometer. qPCR was performed using iQTM SYBR® Green supermix (Bio-Rad)

on the Bio-Rad CFX96 Touch™ Real Time PCR Detection system. Transcript specific primers have

been previously described for *Orf50*, *Orf73*, and *Orf68*[58], LCMV[59], and CVB4[60]. Primers were

ordered from Integrated DNA Technologies (Table 3). Normalized to the ribosomal

housekeeping gene *18s* and expression determined relative to control group.

Gene	Sequence
Orf50 forward	5'-GGCCGCAGACATTTAATGAC-3'
Orf50 reverse	5'-GCCTCAACTTCTCTGGATATGCC-3'
Orf73 forward	5'-AAGGGTTGTCTTGGCCTAC TGTG-3'
Orf73 reverse	5'-AGAGATGCTGTGGGACCATGTTG-3'
Orf68 forward	5'-CTCAAATACACTGGCCGCCATC-3'
Orf68 reverse	5'-CGTGCTTGAGATATGAGTGAGT-3'
CVB4 forward	5'-CCCACAGGACGCTCTAATA-3'
CVB4 reverse	5'-CAGAGTTACCCGTTACGACA -3'
LCMV GP forward	5'-TGCCTGACCAAATGGATGATT-3'
LCMV GP reverse	5'-CTGCTGTGTTCCCGAAACACT-3'
18s forward	5'-GTAACCCGTTGAACCCCATT-3'
18s reverse	5'-CCATCCAATCGGTAGTAGCG-3'

Table 3 qPCR for lytic and latent γHV68 genes primer sequences

Ex vivo limiting dilution reactivation assay

A single cell suspension following RBC lysis with ACK lysis buffer was prepared from freshly

collected spleens. To analyze ex vivo reactivation, limiting dilution reactivation assay was

performed as previously described[17]. Two-fold serial dilutions of splenocytes starting at 10^5

cells per well were plated onto monolayer of mouse embryonic fibroblast (MEF, C57BL/6) cells

(ATCC) in 96-well flat-bottom polystyrene tissue culture plates. Twelve wells were plated per

dilution. Plates were incubated at 37°C 5% CO_2 for 10 days and scored for microscopically for

viral cytopathic effects (CPE). Data were plotted as sigmoidal dose curves and interpolation was

used to determine the cell density per sample at which 63.2% of wells were positive for CPE.

Anti-γHV68 antibodies

Serum was collected via cardiac puncture and maintained at RT to allow for clotting. The sera

were isolated by centrifugation 2000 × g for 10 min, aliquoted, and stored for up to 12 months

at -80°C prior to running the ELISA. Anti-γHV68 antibodies were quantified by standard indirect

ELISA. Briefly, γHV68 virions were inactivated in 4% paraformaldehyde (PFA) for 20 minutes at

RT. Then, ELISA plates (NUNC, Thermo Fisher) were coated with 5 μg/ml γHV68 in coating

buffer (0.05 M NaHCO$_3$, pH 9.6) with 1% PFA overnight at 4°C. The plate was then washed 4x

with wash buffer (PBS-0.05% Tween-20), blocked in blocking buffer (5% NBCS in PBS) for 1 hour

at 37°C, and incubated with serial dilutions (1:20, 1:40, 1:80, 1:160, 1:320, 1:640, 1:1280,

1:2560) of test sera diluted in blocking buffer for 2 hours at 37°C. The plate was washed 4x with

wash buffer and bound antibody was incubated with HRP-conjugated goat anti-mouse IgG

(Thermo Fisher), rat anti-mouse IgG1 (BD Biosciences), or goat anti-mouse IgG2c (Thermo

Fisher), all diluted 1:500 in blocking buffer, for 1 hour at 37°C, washed 4x with wash buffer, and

detected by TMB substrate (BD Biosciences). Absorbance was read at 450 nm on a VarioSkan

Plate Reader (Thermo Fisher).

Statistical analyses

Data analysis and presentation as well as statistical tests were performed using GraphPad Prism

software 8.4.2 (GraphPad Software Inc.). Results are presented as mean ± SEM. Statistical

tests, significance (p-value), sample size (n, number of mice per group) and number of

experimental replicates are stated in the figure legends. Statistical analyses included: two-way

ANOVA with Geisser-Greenhouse's correction, Mann-Whitney test, and one-way ANOVA. P-

values indicated by asterisks as follows: ****$p<0.0001$, *** $p<0.001$, ** $p<0.01$, * $p<0.05$.

Study approval

All work was approved by the Animal Care Committee (ACC) of the University of British

Columbia (Protocols A17- 0105, A17-0184).

Acknowledgements

The following reagent was obtained through the NIH Tetramer Core Facility: class I p79

tetramer. The authors thank Martin Richer for the helpful feedback on the manuscript.

References

1. Barton E, Mandal P, Speck SH. Pathogenesis and host control of gammaherpesviruses: lessons from the mouse. *Annu Rev Immunol*. 2011;29:351-397. doi:10.1146/annurev-immunol-072710-081639

2. Hao Y, O'Neill P, Naradikian MS, Scholz JL, Cancro MP. A B-cell subset uniquely responsive to innate stimuli accumulates in aged mice. *Blood*. 2011;118(5):1294-1304. doi:10.1182/blood-2011-01-330530

3. Rubtsov AV, Rubtsova K, Fischer A, et al. Toll-like receptor 7 (TLR7)–driven accumulation of a novel CD11c+ B-cell population is important for the development of autoimmunity. *Blood*. 2011;118(5):1305-1315. doi:10.1182/blood-2011-01-331462

4. Cancro MP. Age-Associated B Cells. *Annu Rev Immunol*. 2020;38(1):315-340. doi:10.1146/annurev-immunol-092419-031130

5. Mouat IC, Horwitz MS. Age-associated B cells in viral infection. *PLOS Pathog*. 2022;18(3):e1010297. doi:https://doi.org/10.1371/journal. ppat.1010297

6. Rubtsova K, Rubtsov AV, Dyk LF van, Kappler JW, Marrack P. T-box transcription factor T-bet, a key player in a unique type of B-cell activation essential for effective viral clearance. *Proc Natl Acad Sci*. 2013;110(34):E3216-E3224. doi:10.1073/pnas.1312348110

7. Rubtsova K, Rubtsov AV, Halemano K, et al. T Cell Production of IFNγ in Response to TLR7/IL-12 Stimulates Optimal B Cell Responses to Viruses. *PloS One*. 2016;11(11):e0166322. doi:10.1371/journal.pone.0166322

8. Knox JJ, Buggert M, Kardava L, et al. T-bet+ B cells are induced by human viral infections and dominate the HIV gp140 response. *JCI Insight*. 2017;2(8). doi:10.1172/jci.insight.92943

9. Johnson JL, Rosenthal RL, Knox JJ, et al. The Transcription Factor T-bet Resolves Memory B Cell Subsets with Distinct Tissue Distributions and Antibody Specificities in Mice and Humans. *Immunity*. April 2020. doi:10.1016/j.immuni.2020.03.020

10. Eccles JD, Turner RB, Kirk NA, et al. T-bet+ Memory B Cells Link to Local Cross-Reactive IgG upon Human Rhinovirus Infection. *Cell Rep*. 2020;30(2):351-366.e7. doi:10.1016/j.celrep.2019.12.027

11. Woodruff MC, Ramonell RP, Nguyen DC, et al. Extrafollicular B cell responses correlate with neutralizing antibodies and morbidity in COVID-19. *Nat Immunol*. 2020;21(12):1506-1516. doi:10.1038/s41590-020-00814-z

12. Rubtsov AV, Rubtsova K, Kappler JW, Jacobelli J, Friedman RS, Marrack P. CD11c-Expressing B Cells Are Located at the T Cell/B Cell Border in Spleen and Are Potent APCs. *J Immunol Baltim Md 1950*. 2015;195(1):71-79. doi:10.4049/jimmunol.1500055

13. Mendoza A, Yewdell WT, Hoyos B, et al. Assembly of a spatial circuit of T-bet–expressing
T and B lymphocytes is required for antiviral humoral immunity. *Sci Immunol*. 2021;6(60).
doi:10.1126/sciimmunol.abi4710

14. Peng SL, Szabo SJ, Glimcher LH. T-bet regulates IgG class switching and pathogenic
autoantibody production. *Proc Natl Acad Sci U S A*. 2002;99(8):5545-5550.
doi:10.1073/pnas.082114899

15. Barnett BE, Staupe RP, Odorizzi PM, et al. B cell intrinsic T-bet expression is required to
control chronic viral infection. *J Immunol Baltim Md 1950*. 2016;197(4):1017-1022.
doi:10.4049/jimmunol.1500368

16. Ehtisham S, Sunil-Chandra NP, Nash AA. Pathogenesis of murine gammaherpesvirus
infection in mice deficient in CD4 and CD8 T cells. *J Virol*. 1993;67(9):5247-5252.
doi:10.1128/JVI.67.9.5247-5252.1993

17. Weck KE, Barkon ML, Yoo LI, Speck SH, Virgin HW IV. Mature B cells are required for
acute splenic infection, but not for establishment of latency, by murine gammaherpesvirus
68. *J Virol*. 1996;70(10):6775-6780. doi:10.1128/JVI.70.10.6775-6780.1996

18. Willer DO, Speck SH. Long-Term Latent Murine Gammaherpesvirus 68 Infection Is
Preferentially Found within the Surface Immunoglobulin D-Negative Subset of Splenic B
Cells In Vivo. *J Virol*. 2003;77(15):8310-8321. doi:10.1128/JVI.77.15.8310-8321.2003

19. Simas JP, Swann D, Bowden R, Efstathiou S. Analysis of murine gammaherpesvirus-68
transcription during lytic and latent infection. *J Gen Virol*. 1999;80 (Pt 1):75-82.
doi:10.1099/0022-1317-80-1-75

20. Kim IJ, Flaño E, Woodland DL, Blackman MA. Antibody-mediated control of persistent
gamma-herpesvirus infection. *J Immunol Baltim Md 1950*. 2002;168(8):3958-3964.
doi:10.4049/jimmunol.168.8.3958

21. Freeman ML, Burkum CE, Woodland DL, Sun R, Wu TT, Blackman MA. Importance of
antibody in virus infection and vaccine-mediated protection by a latency-deficient
recombinant murine γ-herpesvirus-68. *J Immunol Baltim Md 1950*. 2012;188(3):1049-1056.
doi:10.4049/jimmunol.1102621

22. Stevenson PG, Cardin RD, Christensen JP, Doherty PC. Immunological control of a murine
gammaherpesvirus independent of CD8+ T cells. *J Gen Virol*. 1999;80 (Pt 2):477-483.
doi:10.1099/0022-1317-80-2-477

23. Steed A, Buch T, Waisman A, Virgin HW. Gamma Interferon Blocks Gammaherpesvirus
Reactivation from Latency in a Cell Type-Specific Manner. *J Virol*. 2007;81(11):6134-
6140. doi:10.1128/JVI.00108-07

24. Goodwin MM, Canny S, Steed A, Virgin HW. Murine gammaherpesvirus 68 has evolved
gamma interferon and stat1-repressible promoters for the lytic switch gene 50. *J Virol*.
2010;84(7):3711-3717. doi:10.1128/JVI.02099-09

25.   Stewart JP, Usherwood EJ, Ross A, Dyson H, Nash T. Lung Epithelial Cells Are a Major
Site of Murine Gammaherpesvirus Persistence. *J Exp Med*. 1998;187(12):1941-1951.

26.   Usherwood EJ, Stewart JP, Robertson K, Allen DJ, Nash AA. Absence of splenic latency in
murine gammaherpesvirus 68-infected B cell-deficient mice. *J Gen Virol*. 1996;77 (Pt
11):2819-2825. doi:10.1099/0022-1317-77-11-2819

27.   Weck KE, Kim SS, Virgin HW, Speck SH. B Cells Regulate Murine Gammaherpesvirus 68
Latency. *J Virol*. 1999;73(6):4651-4661.

28.   Siegel AM, Rangaswamy US, Napier RJ, Speck SH. Blimp-1-Dependent Plasma Cell
Differentiation Is Required for Efficient Maintenance of Murine Gammaherpesvirus
Latency and Antiviral Antibody Responses. *J Virol*. 2010;84(2):674-685.
doi:10.1128/JVI.01306-09

29.   Mouat IC, Allanach JR, Fan V, et al. *Gammaherpesvirus Infection Licenses Age-Associated
B Cells for Pathogenicity in MS and EAE*.; 2021:2021.07.22.453263.
doi:10.1101/2021.07.22.453263

30.   Jia Q, Freeman ML, Yager EJ, et al. Induction of Protective Immunity against Murine
Gammaherpesvirus 68 Infection in the Absence of Viral Latency. *J Virol*. 2010;84(5):2453-
2465. doi:10.1128/JVI.01543-09

31.   Flaño E, Husain SM, Sample JT, Woodland DL, Blackman MA. Latent murine gamma-
herpesvirus infection is established in activated B cells, dendritic cells, and macrophages. *J
Immunol Baltim Md 1950*. 2000;165(2):1074-1081. doi:10.4049/jimmunol.165.2.1074

32.   Sunil-Chandra NP, Efstathiou S, Nash AA. Murine gammaherpesvirus 68 establishes a
latent infection in mouse B lymphocytes in vivo. *J Gen Virol*. 1992;73 (Pt 12):3275-3279.
doi:10.1099/0022-1317-73-12-3275

33.   Marques S, Efstathiou S, Smith KG, Haury M, Simas JP. Selective Gene Expression of
Latent Murine Gammaherpesvirus 68 in B Lymphocytes. *J Virol*. 2003;77(13):7308-7318.
doi:10.1128/JVI.77.13.7308-7318.2003

34.   Collins CM, Boss JM, Speck SH. Identification of Infected B-Cell Populations by Using a
Recombinant Murine Gammaherpesvirus 68 Expressing a Fluorescent Protein. *J Virol*.
2009;83(13):6484-6493. doi:10.1128/JVI.00297-09

35.   Collins CM, Speck SH. Tracking murine gammaherpesvirus 68 infection of germinal center
B cells in vivo. *PloS One*. 2012;7(3):e33230. doi:10.1371/journal.pone.0033230

36.   Staudt MR, Dittmer DP. The Rta/Orf50 transactivator proteins of the gamma-herpesviridae.
Curr Top Microbiol Immunol. 2007;312:71-100. doi:10.1007/978-3-540-34344-8_3

37.   Didychuk AL, Gates SN, Gardner MR, Strong LM, Martin A, Glaunsinger BA. A
pentameric protein ring with novel architecture is required for herpesviral packaging. Carter
AP, Wolberger C, Graham SC, eds. *eLife*. 2021;10:e62261. doi:10.7554/eLife.62261

38. Fowler P, Marques S, Simas JP, Efstathiou S. ORF73 of murine herpesvirus-68 is critical
for the establishment and maintenance of latency. *J Gen Virol*. 2003;84(Pt 12):3405-3416.
doi:10.1099/vir.0.19594-0

39. Moorman NJ, Willer DO, Speck SH. The gammaherpesvirus 68 latency-associated nuclear
antigen homolog is critical for the establishment of splenic latency. *J Virol*.
2003;77(19):10295-10303. doi:10.1128/jvi.77.19.10295-10303.2003

40. Wang G, Zarek C, Chang T, Tao L, Lowe A, Reese TA. Th2 Cytokine Modulates
Herpesvirus Reactivation in a Cell Type Specific Manner. *J Virol*. February
2021:JVI.01946-20. doi:10.1128/JVI.01946-20

41. Barton ES, White DW, Cathelyn JS, et al. Herpesvirus latency confers symbiotic protection
from bacterial infection. *Nature*. 2007;447(7142):326-329. doi:10.1038/nature05762

42. Tibbetts SA, van Dyk LF, Speck SH, Virgin HW. Immune Control of the Number and
Reactivation Phenotype of Cells Latently Infected with a Gammaherpesvirus. *J Virol*.
2002;76(14):7125-7132. doi:10.1128/JVI.76.14.7125-7132.2002

43. Gredmark-Russ S, Cheung EJ, Isaacson MK, Ploegh HL, Grotenbreg GM. The CD8 T-Cell
Response against Murine Gammaherpesvirus 68 Is Directed toward a Broad Repertoire of
Epitopes from both Early and Late Antigens. *J Virol*. 2008;82(24):12205-12212.
doi:10.1128/JVI.01463-08

44. Steed AL, Barton ES, Tibbetts SA, et al. Gamma Interferon Blocks Gammaherpesvirus
Reactivation from Latency. *J Virol*. 2006;80(1):192-200. doi:10.1128/JVI.80.1.192-
200.2006

45. Dutia BM, Clarke CJ, Allen DJ, Nash AA. Pathological changes in the spleens of gamma
interferon receptor-deficient mice infected with murine gammaherpesvirus: a role for CD8
T cells. *J Virol*. 1997;71(6):4278-4283.

46. Tripp RA, Hamilton-Easton AM, Cardin RD, et al. Pathogenesis of an Infectious
Mononucleosis-like Disease Induced by a Murine γ-Herpesvirus: Role for a Viral
Superantigen? *J Exp Med*. 1997;185(9):1641-1650.

47. Hardy CL, Flaño E, Cardin RD, et al. Factors controlling levels of CD8+ T-cell
lymphocytosis associated with murine gamma-herpesvirus infection. *Viral Immunol*.
2001;14(4):391-402. doi:10.1089/08828240152716637

48. Evans AG, Moser JM, Krug LT, Pozharskaya V, Mora AL, Speck SH. A
gammaherpesvirus-secreted activator of Vbeta4+ CD8+ T cells regulates chronic infection
and immunopathology. *J Exp Med*. 2008;205(3):669-684. doi:10.1084/jem.20071135

49. Stevenson PG, Belz GT, Altman JD, Doherty PC. Changing patterns of dominance in the
CD8+ T cell response during acute and persistent murine gamma-herpesvirus infection. *Eur
J Immunol*. 1999;29(4):1059-1067. doi:10.1002/(SICI)1521-
4141(199904)29:04<1059::AID-IMMU1059>3.0.CO;2-L

50.   Barton ES, Rajkarnikar S, Langston PK, Price MJ, Grayson JM. Gammaherpesvirus latency
638   differentially impacts the generation of primary versus secondary memory CD8+ T cells
639   during subsequent infection. *J Virol*. 2014;88(21):12740-12751. doi:10.1128/JVI.02106-14

640   51.   Jacobsen H, Klein SL. Sex Differences in Immunity to Viral Infections. *Front Immunol*.
641   2021;12:3483. doi:10.3389/fimmu.2021.720952

642   52.   Ricker E, Manni M, Flores-Castro D, et al. Altered function and differentiation of age-
643   associated B cells contribute to the female bias in lupus mice. *Nat Commun*.
644   2021;12(1):4813. doi:10.1038/s41467-021-25102-8

645   53.   Bussey KA, Murthy S, Reimer E, et al. Endosomal Toll-Like Receptors 7 and 9 Cooperate
646   in Detection of Murine Gammaherpesvirus 68 Infection. *J Virol*. 2019;93(3):e01173-18.
647   doi:10.1128/JVI.01173-18

648   54.   Rubtsova K, Rubtsov AV, Thurman JM, Mennona JM, Kappler JW, Marrack P. B cells
649   expressing the transcription factor T-bet drive lupus-like autoimmunity. *J Clin Invest*.
650   2017;127(4):1392-1404. doi:10.1172/JCI91250

651   55.   Vockerodt M, Yap LF, Shannon-Lowe C, et al. The Epstein-Barr virus and the pathogenesis
652   of lymphoma. *J Pathol*. 2015;235(2):312-322. doi:10.1002/path.4459

653   56.   Shikova E, Reshkova V, Kumanova A, et al. Cytomegalovirus, Epstein-Barr virus, and
654   human herpesvirus-6 infections in patients with myalgic encephalomyelitis/chronic fatigue
655   syndrome. *J Med Virol*. March 2020. doi:10.1002/jmv.25744

656   57.   Tsuge I, Morishima T, Kimura H, Kuzushima K, Matsuoka H. Impaired cytotoxic T
657   lymphocyte response to Epstein-Barr virus-infected NK cells in patients with severe
658   chronic active EBV infection. *J Med Virol*. 2001;64(2):141-148. doi:10.1002/jmv.1029

659   58.   Abernathy E, Clyde K, Yeasmin R, et al. Gammaherpesviral gene expression and virion
660   composition are broadly controlled by accelerated mRNA degradation. *PLoS Pathog*.
661   2014;10(1):e1003882. doi:10.1371/journal.ppat.1003882

662   59.   Welsh RM, Seedhom MO. LCMV: Propagation, quantitation, and storage. *Curr Protoc
663   Microbiol*. 2008;CHAPTER:Unit-15A.1. doi:10.1002/9780471729259.mc15a01s8

664   60.   Walter DL, Benner SE, Oaks RJ, et al. Coxsackievirus B4 Exposure Results in Variable
665   Pattern Recognition Response in the Kidneys of Female Non-Obese Diabetic Mice Before
666   Establishment of Diabetes. *Viral Immunol*. 2020;33(7):494-506.
667   doi:10.1089/vim.2019.0188

668
669

Figure 1

Figure 1 ABCs expand in the blood and spleen throughout γHV68 infection

Blood and spleen collected from C57BL/6(J) mice (6–8-week-old at infection) mock-infected

with media (naïve, black circles) or infected i.p. with γHV68 for 6 (red circles), 35 (grey circles),

or 150 (blue circles) days and processed for flow cytometry. **(A)** Representative gating strategy

of ABCs. IgD, CD11c, and T-bet gating used fluorescence-minus-one (FMO) controls. **(B-E)**

Proportion of ABCs (CD11c$^+$T-bet$^+$) of previously activated B cells (CD19$^+$IgD$^-$) in the blood and

spleen of male and female mice. **(F-I)** Proportion of ABCs (CD11c$^+$T-bet$^+$) of previously activated

B cells (CD19$^+$IgD$^-$) in the blood or spleen of male and female mice. Same data as presented in

panels B-E. n=10-15 mice per group, data compiled from 4 experiments. Each data point

represents an individual mouse. Data presented as mean ± SEM. Analyzed by **(B-E)** one-way

ANOVA or **(F-M)** Mann-Whitney test. P-values indicated as asterixis as follows: *** p<0.001, **

p<0.01, * p<0.05.

Figure 2

Figure 2 Cytokine production by ABCs in the spleen during γHV68 and ACRTA-γHV68 infection

(**A-C**) C57BL/6(J) mice (6–8-week-old at infection) mock-infected with media (naïve) or infected

i.p. with γHV68 for 6, 35, or 150 days. Spleens collected and processed for flow cytometry.

Proportion of ABCs in the spleen expressing (**A**) IFNγ, (**B**) TNF, or (**C**) IL-17A, with females

presented as open circles and males as filled circles. (**D**) Gating of ABCs (CD11c$^+$T-bet$^+$) and non-

ABC B cells (CD11c$^-$T-bet$^+$ and CD11c$^-$T-bet$^-$), previously gated on CD19$^+$IgD$^-$ cells in the spleen.

(**E**) Representative histograms and dot plots of the proportion of ABCs and non-ABC B cells

(CD11c$^-$T-bet$^+$ and CD11c$^-$T-bet$^-$) expressing IFNγ and TNF from mice infected with γHV68 for 35

days. (**F-H**) C57BL/6(J) mice mock-infected with media (Naïve, black circles) or infected i.p. with

γHV68 (grey circles) or ACRTA-γHV68 (green diamonds) for 35 days and spleens processed for

flow cytometry. (**F**) Proportion of ABCs (CD11c$^+$T-bet$^+$) of previously activated B cells (CD19$^+$IgD$^-$)

in the spleen. Proportion of ABCs (CD19$^+$CD11c$^+$T-bet$^+$) in the spleen expressing (**G**) IFNγ or (**H**)

TNF. (**I**) Representative plots and quantification of the proportion of ABCs expressing IFNγ and

TNF. Equivalent numbers of females and per group in all panels. (**A-C**) n=6-10 mice per group,

data combined from two experiments. (**E-I**) n=6-8 mice per group, representative of two

experiments. Each data point represents an individual mouse. Data presented as mean ± SEM.

Analyzed by one-way ANOVA. P-values indicated as asterixis as follows: ****p<0.0001, ***

p<0.001, ** p<0.01, * p<0.05.

Figure 3

A

Previously activated

| | T cells CD3+ | B cells CD19+IgD- | ABCs CD19+CD11c+T-bet+ |

γHV68 WT

- T cells: 0.06 ± 0.03
- B cells: 0.07 ± 0.03
- ABCs: 0.10 ± 0.04

γHV68 H2bYFP

- T cells: 0.05 ± 0.02
- B cells: 1.2 ± 0.3
- ABCs: 1.1 ± 0.2

SSC / YFP

B

%YFP+

- ● γHV68
- △ γHV68 H2bYFP

p=0.94

T cells, Innate cells, Naive B cells, Previously activated B cells, ABCs

C

- ☐ 73% Previously activated B cells (CD3-CD19+IgD-)
- ☐ 6%   ABCs (CD3-CD19+CD11c+T-bet+)
- ☐ 6%   Naive B cells (CD3-CD19+IgD+)
- ☐ 3%   NK cells (CD3-CD19-NK1.1+)
- ☐ 2%   Macrophages (CD3-CD19-F4/80+)
- ☐ 3%   DCs (CD3-CD19-CD11c+)
- ☐ 8%   T cells (CD3+CD19-)

708
709

D

ABCs

- 0.5 ± 0.07
- 0.06 ± 0.1
- 66.6 ± 6.1
- 31.9 ± 6.0

γHV68 / IFNγ

IFNγ+ ABCs

%IFNγ+

**** p<0.0001

Uninfected   Infected

Figure 3 Analysis of ABC infection status with fluorescent γHV68 strain

Female C57BL/6(J) mice (6 to 8-week-old at infection) infected i.p. with γHV68 or

γHV68.H2bYFP. 8 days p.i. spleen collected and processed for flow cytometry. (**A**)

Representative flow cytometry plots displaying yellow fluorescent protein (YFP) expression (x-

axis) versus side-scatter (SSC, y-axis) from mice infected with γHV68 or γHV68.H2bYFP with

mean ± SEM. Samples gated on live, single lymphocytes, and then T cells (CD3$^+$), total B cells

(CD19$^+$), previously activated B cells (CD19$^+$IgD$^-$), or ABCs (CD19$^+$CD11c$^+$T-bet$^+$). Two samples

per group concatenated for the purpose of visualization. Representative of two experiments.

(**B**) Proportion of cell subsets positive for YFP expression from γHV68-infected mice (filled

circles) and γHV68.H2bYFP-infected mice (open triangles). Each data point represents an

individual mouse. (**C**) Proportion of YFP$^+$ cells that are non-B cells (CD19$^-$, white), non-ABC B

cells (CD19$^+$CD11c$^-$T-bet$^-$, grey), or ABCs (CD19$^+$CD11c$^+$T-bet$^+$, black). Data n=6 mice per group,

data compiled from two experiments, representative of three experiments. (**D**) Representative

flow plot of ABCs (CD19$^+$CD11c$^+$T-bet$^+$) in the spleen gated for YFP (γHV68.H2bYFP) and IFNγ

with mean ± SEM. Dot plot displays the proportion of either uninfected (YFP$^-$) or infected (YFP$^+$)

ABCs in the spleen that are positive for IFNγ. (**B, D**) Data presented as mean ± SEM, analyzed by

Mann-Whitney test. Each data point represents an individual mouse. P-values indicated as

asterixis as follows: ****p<0.0001.

Figure 4

730
731

Figure 4 ABCs are not required for controlling lytic infection and achieving latency

Tbx21$^{fl/fl}$Cd19$^{+/+}$ (Ctrl, filled grey circles) and *Tbx21$^{fl/fl}$Cd19$^{cre/+}$* (KO, open blue squares) mice

were infected i.p. with γHV68 or ACRTA-γHV68. 6 or 35 days p.i., spleens collected and

processed for RNA extraction, DNA extraction, and flow cytometry. (**A**) Experimental scheme

for data shown in panels B, C, D, and F. (B) Proportion of ABCs (CD11c$^+$T-bet$^+$) of previously

activated B cells (CD19$^+$IgD$^-$) in the spleen in Ctrl and KO mice 6 and 35 days post-γHV68

infection. (**C-D**) Splenic γHV68 viral load (copies *Orf50* per million cells) in Ctrl (filled circles) and

KO (open squares) mice as determined by qPCR day 6 and 35 p.i., with limit of detection

indicated by dotted line. (**E**) Splenic γHV68 viral load (copies *Orf50* per million cells) in Ctrl and

KO mice infected with γHV68 or ACRTA-γHV68 35 days p.i.. (**F**) Relative expression in the spleen

of *Orf73*, *Orf50*, and *Orf68* in Ctrl and KO mice at day 35 p.i.. Each data point represents an

individual mouse. Data presented as mean ± SEM, analyzed by Mann-Whitney test (**B-D, F**) or

Kruskal-Wallis H test (**E**).

Figure 5

Figure 5 Mice without ABCs display a dysregulated anti-γHV68 antibody response, though no

alterations to the cell populations infected with γHV68 or γHV68-responding T cells,

compared to mice with ABCs

(**A-C**) *Tbx21$^{fl/fl}$Cd19$^{+/+}$* (Ctrl, filled grey circles) and *Tbx21$^{fl/fl}$Cd19$^{cre/+}$* (KO, open blue squares)

mice were infected i.p. with γHV68 for 6 or 35 days. N=5 per group, representative of 2

experiments. (**D**) Ctrl and KO mice were infected i.p. with γHV68.H2bYFP for 8 days. N=6-8 per

group, data compiled from 2 experiments. (**E-H**) Ctrl and KO mice were infected i.p. with γHV68

for 6 or 35 days, at which point spleens were collected for flow cytometry. Representative of 2

experiments. Each data point represents an individual mouse. Data presented as mean ± SEM.

Analyzed by two-way ANOVA (**A-C**) or Mann-Whitney test (**D-H**). P-values indicated as asterixis

as follows: ****$p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$.

Figure 6

763
764

Figure 6 ABC knockout mice are more susceptible to reactivation

(**A**) *Tbx21*$^{fl/fl}$*Cd19*$^{+/+}$ (Ctrl, filled grey circles) and *Tbx21*$^{fl/fl}$*Cd19*$^{cre/+}$ (KO, open blue squares) mice

were infected i.p. with γHV68. 35 days p.i., spleens were collected and processed for

reactivation assay. N=6 mice per group, data compiled from two independent experiments. (**B**)

Experimental scheme for data shown in panels C-I. Ctrl and KO mice were infected i.p. with

γHV68 for 35 days and then challenged with either LCMV (Armstrong) or CVB4 for 10 to 4 days,

respectively. (**C-I**) Spleens collected, processed, and DNA and RNA was extracted from total

splenocytes following RBC lysis. (**C, D**) Relative expression of LCMV (**C**) and CVB4 (**D**) between

Ctrl and KO mice measured by RT-qPCR. (**E, F**) Copies of γHV68 detected by qPCR in the spleen

per million cells following challenge with LCMV or CVB4. (**G-I**) Relative expression in the spleen

of *Orf73*, *Orf50*, and *Orf68* in Ctrl and KO mice. Each data point represents an individual mouse.

Data presented as mean ± SEM, analyzed by Mann-Whitney test (**A, C-D, G-I**) or one-way

ANOVA (**E, F**). P-values indicated as asterixis as follows: ** $p<0.01$, * $p<0.05$.