Nerve-mediated amputation-induced stem cell activation primes distant appendages for future
regeneration events in axolotl

Duygu Payzin-Dogru[1]*, Sarah E. Wilson[1]*, Steven J. Blair[1], Burcu Erdogan[1], Shifa Hossain[1], Louis
Cammarata[2], Juan Velazquez Matos[1], Alan Wong[1], Julia Losner[1], Sangwon Min[1], Hani Singer[1], Jessica L.
Whited[1,3,4]

[1]Department of Stem Cell and Regenerative Biology, Harvard University, 7 Divinity Avenue, Cambridge,
MA, USA, 02138

[2]Department of Statistics, Harvard University

[3]The Broad Institute

[4]The Harvard Stem Cell Institute

Correspondence: Jessica L. Whited, jessica_whited@harvard.edu

* These authors contributed equally to this work

Summary

Animals exhibit extreme diversity in regenerative ability. This likely reflects different, lineage-specific
selective pressures in their evolutionary histories, but how specific molecular features of regenerative
programs help solve species-specific challenges has not been examined in detail. Here we discover that,
in the highly-regenerative axolotl salamander, a conserved, body-wide stem cell activation response
triggered in response to limb removal primes undisturbed limbs for regeneration upon subsequent
amputation. This response should be particularly useful to salamanders, which frequently lose limbs in
response to cannibalism. We further demonstrate the body-wide response requires both peripheral
nervous system input at these distant sites and mTOR signaling. We defined gene expression changes
within the nerves and nearby tissues, harboring responsive stem cells, leading to identification of
candidate genetic pathways influencing distant stem cell activation following amputation. Functional
experimentation confirmed a requirement for adrenergic signaling in amputation-induced activation of
distant stem cells. These findings reveal a direct link between systemic cellular activation responses to
local tissue damage and overall regenerative ability. Similar systemic activation responses to tissue
removal have been observed in animals with widely differing regenerative abilities (e.g., planaria to
mice), suggesting that it is the responses downstream of these signals, likely sculpted by differing
evolutionary pressures, that ultimately distinguish regenerators from non-regenerators.

Keywords: regeneration, stem cells, axolotl, limb, systemic responses, peripheral nervous system

Main

Loss of a body part provokes cellular and molecular changes at the injury site and systemically
[1]. Most research in vertebrate models has focused on local responses, within the remaining "stump"
tissues, as cells within these tissues contribute directly to the construction of the new body part. Yet,
understanding the mechanism and consequence of systemic responses is critical as they may contribute
to regeneration and because comparing them across species may reveal insights related to the evolution
of regeneration [2]. The axolotl, a highly-regenerative salamander for which molecular genetic tools
now exist [3], provides an extraordinary model for studying regeneration events due to its ability to
regenerate complex tissues and organs such as limbs, hearts, and parts of the nervous system. Following
amputation in axolotl, a structure essential for regeneration called the blastema forms at the
amputation site [4]. The blastema is largely composed of a dense population of highly-proliferative
progenitor cells that ultimately undergo differentiation, repatterning, and growth to replace the lost
structure. Our current understanding of limb regeneration has focused primarily on local events at the
injury site. However, the discovery of a body-wide response to amputation in axolotls [5], in which many
distant stem cells are activated to re-enter the cell cycle, raises the possibility that systemic activation
may be a component of the regeneration process on which local growth responses are elaborated.
Indeed, the activation of stem cells distant from the site of injury is evolutionarily conserved amongst
many other species, both super-regenerators (such as planarians) and those with more limited natural
regenerative prowess (such as mice) [1, 6-8]. The extensive conservation of this distant-activation
phenomenon suggests it is an evolutionarily ancient response and raises the question of whether it is
used differently in species with differing regenerative potentials, and, if so, how this relates to specific
evolutionary pressures these species face.

Here, we investigate the consequence and mechanism of systemic stem cell activation following
amputation in axolotls. We identify potential selective advantages of systemic activation and discover a
distant stem cell-priming program driving rapid activation of progenitors and faster proliferation of
blastema cells in animals recently challenged with contralateral amputations. Our results highlight
mTOR signaling and innervation as critical mediators of systemic activation, with a vital role in the
regenerative process. We also use transcriptomics to identify additional candidate signaling pathways
controlling systemic activation and implicate adrenergic signaling as a key regulator of systemic
activation. Our work reveals the impact evolutionarily conserved systemic cues play in the regenerative
process with implications for our broader understanding of the evolution of regenerative competency.

Systemically-activated cells are primed for future regeneration events

We recently reported systemic cell cycle re-entry, defined to be "activation," following limb
amputation in axolotl, and identified muscle satellite stem cells to constitute a sizeable fraction of these
systemically-active cells [5]. We now demonstrate that among the other cells activated in this system
are epidermal stem cells, as defined by their basal position in the tissue and expression of Keratin 10
(K10) (Extended Data Fig. 1). We first reported that systemically-activated cells undergo at least one
round of cell division, as evidenced by their expression of phosphorylated Histone H3 [5]. We performed
a similar single-pulse EdU experiment, but with a long chase (six weeks). This experiment revealed that
systemically-activated cells form small clones, typically 2-4 cells in size (Extended Data Fig. 2), indicating
that the response is limited to one or two cell divisions.

To determine if systemic cell activation confers a selective advantage by enhancing the
regenerative ability of axolotls, we compared regeneration rates of limbs of systemically-activated
animals to those of controls (Fig. 1a). Systemically-activated animals, which received contralateral
amputations 7 days prior, exhibited faster blastema growth (blastema area) compared to control
animals only receiving a single set of amputations (Fig. 1b, c). The limbs of systemically-activated
animals differentiate faster than those of controls, entering the palette stage faster, with outlines of the
future digits appearing sooner than in controls, and internal cartilage nodules clearly visible (Fig. 1b, d-e)
[9]. Systemic activation of cells in limbs contralateral to an amputation primes them for future
regeneration events, enabling the activated limbs to initially coalesce a blastema population of
progenitors and then also differentiate them more quickly, ultimately leading to faster regeneration.
87

mTOR signaling is required for systemic activation

The upregulation of mTOR signaling during regeneration and wound healing is conserved across
many species such as planaria, zebrafish, mice, and axolotl [5, 10-13]. In planaria, mTORC1 is necessary
for proliferation of neoblast stem cells to promote regeneration [11], and a specific subset of neoblasts
characterized as "slow-cycling" during homeostasis relies on mTOR signaling to mount a regenerative
response [14]. Furthermore, experimental inhibition of mTOR signaling can counteract
hyperproliferation of neoblasts in response to amputation caused by RNAi against *smg1*, a PI3-Kinase-
related kinase [10]. Zebrafish caudal fin regeneration also requires mTOR signaling for progenitor cell
proliferation, although activation of this pathway in tissues or cells distant to amputation has not been
examined in detail [12]. In mice, which repair discrete tissues such as muscle but cannot regenerate full
appendages, local tissue injury has been demonstrated to lead to systemic changes as distant stem cells
are provoked to enter an "alert" phase of the cell cycle characterized by increased nuclear size and
genome replication [8]. The "alert" state sensitizes distant stem cells to respond more quickly to similar
injuries in the future [8]. The conserved activation of mTOR across regenerative models led us to explore
mTOR signaling as a requirement for both local and systemic stem cell activation during limb
regeneration in axolotl. We sought to answer this question by inhibiting mTORC1 signaling via
rapamycin. Rapamycin-treated animals produced smaller blastemas compared to controls (Extended
Data Fig. 3). Treatment with rapamycin wholly blocked cell-cycle activation of stem cells systemically in
contralateral limbs (Fig. 2a-b). These findings indicate that mTOR signaling is required for stem cell
activation at both the local injury site, as well as at distant locations.

Overlap of cells engaging in mTOR signaling and those actively proliferating in systemically
activated limbs is scarce [5], implying that non-cell-autonomous effects may control mTOR mediated
activation of stem cells at locations distant to the site of injury. In an attempt to identify the conduit
transmitting the essential non-cell-autonomous mTOR signal throughout the axolotl body, we examined
mTOR activation elsewhere in systemically-activated animals. We analyzed the brains of systemically-
activated axolotls, and found robust overlap between expression of phosphorylated s6 Kinase (pS6), a
marker for mTOR activation, and NeuN, a marker post-mitotic neurons [15], within the ipsilateral
anterior telencephalon, suggesting significant mTOR activation in this region of the brain (Extended Data
Fig. 4), and prompting us to explore the nervous system more generally. We examined mTOR activation
in potential neuronal processes upstream and downstream of the brain, with relation to limb, such as
local peripheral nervous system ganglia. We found that the dorsal root ganglia, which contain the cell
bodies of peripheral sensory nerves, are positive for pS6 signal in homeostasis, and that signal intensity

increases specifically in the DRG serving the amputated limb (Extended Data Fig. 5). These data imply
peripheral nerves may be important for the mediating systemic responses.

Nerves drive systemic activation

Nerve input in the stumps of amputated limbs is an essential requirement for limb regeneration
in axolotl and newt [16, 17]. It is also a critical local input in other regenerative contexts in many species,
such as heart regeneration in zebrafish and neonatal mice [18]. Recently, the nervous system has been
shown to mobilize several stem cell niches in mammalian systems [19, 20]. The upregulation of mTOR
signaling in both the central and peripheral nervous systems of systemically activated animals led us to
hypothesize that nerves may be the potential agent communicating systemic cues and signals across the
body of the axolotl post-amputation (Extended Data Fig. 4 and 5). To test this hypothesis, we transected
nerve bundles innervating limbs contralateral to amputations and assayed for successful systemic
activation in denervated and sham-operated states (Fig. 3a). Nerve transections performed in intact
animals did not induce systemic cell cycle re-entry (Fig. 3b-c). Denervation of limbs contralateral to
amputations completely impeded the activation of cells in those appendages demonstrating limb
innervation is a requirement in appendages for systemic activation (Fig. 3b-c),

Using cell and tissue morphology and location, we classified responding tissues in sham and
denervated contralateral limbs to examine the relationship between innervation and cell-type-specific
activation. Tissues were classified into three categories: epidermis, skeletal elements, and other internal
tissues. Skeletal and internal soft tissues in sham amputated animals displayed a significant increase in
EdU+ nuclei compared to intact controls (Fig. 3d-f). Denervated limbs in both intact and amputated
states did not exhibit an increase in EdU+ nuclei compared to unamputated controls (Fig. 3d-f). Our
results suggest that nerve-mediated systemic activation of distant progenitors is not tissue type specific,
indicating peripheral nervous system input is likely required for systemic activation of all limb tissue
types in axolotl.

Transcriptomic analyses reveal pathways mediating systemic activation

We next sought to identify candidate factors regulating nerve-mediated systemic cell activation.
We performed bulk-tissue RNA sequencing on peripheral nerves, excluding cell bodies, and surrounding
tissues in the contralateral limbs of denervated and sham-operated regenerating animals (Fig. 4a). We
uncovered 273 differentially-expressed transcripts between intact and transected nerves, and 82
differentially-expressed transcripts between activated and unactivated surrounding tissue samples
(Supplementary File 1). Our RNA-seq analysis uncovered many genes with known roles in regeneration
and cell proliferation, both in axolotl and other regeneration models. Among the transcripts upregulated
in transected nerve ends, we identified hepatocyte growth factor (Hgf), a systemic factor shown to
stimulate quiescent distant tissue stem cells [8, 21] and transcripts encoding kazal-type serine protease
inhibitor 2 (Kazald2) and von Willebrand factor D and EGF domains (VWDE), both of which are blastema
cell markers required for axolotl limb regeneration [22, 23]. In surrounding tissue samples of
systemically activated animals, we discovered transcripts encoding PCNA (proliferating cell nuclear
antigen) to be more highly expressed compared to denervated limbs in which systemic activation was
blocked. Conversely, we found downregulation of transcripts encoding serine/threonine-protein kinase
Lats1. Inhibition of Lats1, a negative regulator of YAP1 in the Hippo signaling pathway, promotes
proliferation in murine cardiomyocytes and also cells in the inner ear [24].

We then directed our attention to candidate pathways controlling distant cell activation which
have not been extensively studied in the context of limb regeneration. We prioritized investigating
transcripts of genes known to encode proteins in signaling pathways, with established chemical
inhibitors, for further functional study. Among these candidates was α-2A adrenergic receptor (Adra2a),
with transcripts 7.4-fold upregulated in transected nerve ends compared to controls. Adra2a activates
hepatic stellate cells in mouse hepatocellular carcinoma models [25] and its silencing has been shown to
increase cell proliferation in cervical cancer cell lines through inhibition of the PI3K/AKT/mTOR pathway
[26]. We explored the potential role of this gene in systemic cell cycle re-entry by inhibiting the function
of its encoded protein with yohimbine, a well-studied Adra2a antagonist [27-29]. In animals treated
with yohimbine, limbs contralateral to amputation had a significantly lower fraction of EdU+ cells
compared to vehicle-treated controls (Fig. 4b-c). Thus, we conclude that adrenergic signaling is required
for distant cell activation in response to amputation, and we propose this signal originates in the nerve.

Other candidate pathways we have tested include insulin and Wnt signaling. Our RNA-seq
dataset revealed that the expression of canonical Wnt signaling pathway transcription factor lymphoid
enhancer binding factor 1 (Lef1) is 5.9-fold increased in severed nerve ends. Intriguingly, Lef1 has
recently been shown to prime adult skin to enhance regeneration outcomes in mice [30] and is
expressed in the wound epidermis of regenerating zebrafish fins [31]. While Lef1 itself has not been
directly studied in the context of axolotl limb regeneration, Wnt signaling has been shown to regulate
limb regeneration in axolotl and *Xenopus* and fin regeneration in zebrafish [32]. Surprisingly, the
inhibition of Wnt signaling by the tankyrase inhibitor IWR-1 did not block systemic cell cycle re-entry
(Extended Data Fig. 6a-b), suggesting that the Wnt requirement may act locally, at the amputation site,
in axolotls. We also found transcripts encoding insulin-like growth factor 1 receptor (Igf1r) to be 19.2-
fold upregulated in tissues surrounding the nerve in denervated samples compared to sham-operated
controls. Igf1r is required for axonal regeneration of retinal ganglion cells from rats, and has been
implicated in progenitor activation mechanisms in several regeneration studies [33-35]. Inhibition of
Igf1r by an ATP-competitive inhibitor, NVP-ADW742 [35], blocked distant cell activation in limbs
contralateral to amputation, indicating that Igf-1 signaling is also required for this process (Extended
Data Fig. 6a-b). Taken together, our data suggest that nerve-mediated Adra2a signaling and Igf-1
signaling in the responding cells mediate systemic cell activation in response to amputation injury in
axolotl.

Discussion

Limb blastemas form at the site of amputation in salamanders, which has led the field to focus
on events occurring at the injury site. Early localized irradiation studies demonstrated that only cells
within the remaining limb stump are required to divide for regeneration [36]. While this lends support
to a model wherein only cells at the amputation site give rise to the blastema, understanding how cells
interpret injury signals in response to amputation is crucial to understanding regeneration. The
traditional view postulates that progenitor cells in the injured limb are activated upon amputation,
migrate to the tip of the stump, and proliferate to form the blastema. However, our earlier work
demonstrating that injury signals activate distant tissue progenitors [5] supports a model where cell
activation and blastema cell commitment are two separate events, and predicts that some activated
cells do not become blastema cells. This possibility led us to ask whether there could be an advantage to
having this system-wide activation that is later refined to the injury site. Here, we show that global
activation primes distant progenitors to respond faster in the event of an additional injury (Fig. 4d). This

"jump start" could serve as a selective advantage in salamanders, many species of which lay eggs in high
concentrations in small vernal pools [37], leading to dense aggregates of siblings and other conspecifics.
Salamanders developing in environments with limited space and resources are cannibalistic [38-41],
often resulting in amputation injuries to more than one limb in juveniles. Becoming poised for a future
injury upon losing a limb would allow the animal to quickly respond to subsequent loss of another limb,
and to produce new functional limbs, which is essential for their survival.

Our work provides mechanistic insights into how injury signals are transmitted to distant
locations. The nervous system has recently been implicated in mammalian systems in stem cell
mobilization [19, 20] and our work supports the idea that nerves can directly communicate with resident
progenitor cells to initiate cell cycle re-entry. Studies of other regeneration models have also implicated
circulation as a means to transduce injury signals [8, 13, 21]. While our work does not exclude the
possibility that factors circulating in the blood may also transmit injury signals, here we report a role of
nerves. Future work will help uncover whether similar signals are also carried through the blood in
axolotls. Our findings should prompt a re-evaluation of the role of peripheral nerves in the regenerative
process *per se*, including within the amputated limb itself, because it indicates a role for nerves in the
primary decision cells make to re-enter the cell cycle rather than simply serving to maintain proliferation
of cells that have independently made this choice.

Regeneration of complex body parts is often explained as an all-or-nothing phenomenon for a
given species. Yet, evidence that partial responses can occur, such as adult frogs regenerating
cartilaginous spikes missing many internal tissues and lacking appropriate patterning [42, 43], and lizards
regenerating tails lacking dorsal-ventral polarity and neuronal cell bodies [44-47], challenges this view.
Our data regarding systemic cellular activation in axolotl argues this feature may be a component of the
regenerative process and that it has a potentially selectable consequence of enhancing future
regenerative responses of distant tissues. Combined with previous reports of systemic activation
following injury in mice, we argue that a paradigm worth considering is that both species typically
classified as non-regenerators and as regenerators engage in a similar initial post-amputation response
that activates prospective progenitor cells throughout the body. This systemic response may therefore
be ancestral, present in mammals as a vestigial component of a broader regeneration program. In this
scenario, perhaps the key critical difference between the two types of animals is their ability to harness
this cellular activation at the site of injury, coalescing activated cells into a blastema, and sustaining
proliferation there long enough to spur regeneration. These issues have direct implications for
development of regenerative medicine approaches because a pool of proliferative progenitors is an
essential pre-requisite of coaxing a limb to regenerate without resorting to cellular transplantation. We
predict future therapies may harness endogenous stem cells that have been naturally and successfully
primed for appendage regeneration but require molecular instruction to transition to a blastema-like
state and, once instructed, these cells may organize a response more akin to salamander limb
regeneration.

Methods

Animal Experimentation

All animal experimentation was approved by and conducted in accordance with Harvard University's
Institutional Animal Care and Use Committee. Leucistic axolotls were used for all animal
experimentation except RNA sequencing and maintained as previously described [22]. For RNA
sequencing, *βIIItub:GAP43-EGFP* transgenic animals were used [48]. Animals in all experiments were
both age and size matched. Animals were anesthetized in 0.1% (w/v) tricaine prior to all procedures
involving amputation, denervation, or injection. After all surgical procedures, axolotls were allowed to
recover overnight in 0.5% (w/v) sulfamerazine. All limb amputations were performed at the mid-
stylopod and the bone was then trimmed back from the amputation plane. For forelimb denervations
the brachial plexus nerves were transected according to [49] and for hindlimb denervations the sciatic
nerves were transected according to [50]. Sham operations as controls for denervations were performed
by anesthetizing animals and cutting the skin open with sterile scissors, locating the nerve bundles with
closed sterile forceps, but not cutting the nerves.

Drug treatments and EdU incorporation

Rapamycin was dissolved into a 25 mg/ml stock solution in ethanol. Rapamycin treatments were
delivered via both water treatments at a concentration of 1 µM for 16 days with water changed daily,
and through daily intraperitoneal injections at a concentration of 5 mg/kg/day for 16 days, both starting
two days before amputation. All intraperitoneal injections were administered at a volume of 20µL/g in a
vehicle solution comprised of APBS with 5% Tween80, 5% PEG-400. All stock solutions were frozen at
−80 °C in 1-mL aliquots. Yohimbine (Y3125, Sigma-Aldrich) was dissolved into a 10 mg/mL stock solution
with sterilized water. Yohimbine was administered daily at a dose of 1mg/kg for 14 days via
intraperitoneal injection. A stock solution of IWR-1 (I0161, Sigma-Aldrich) was created by dissolving
5mg/mL in DMSO. IWR-1 was delivered daily at a dose of 10mg/kg/day for 9 days via intraperitoneal
injection. A stock solution of NVP-ADW742 (5247, Tocris) was created by dissolving 2mg/mL in DMSO.
NVP-ADW742 was injected daily at a dose of 10mg/kg/day for 14 days.

5-ethynyl-2'-deoxyuridine (EdU) was purchased from ThermoFisher and stock solutions were created by
dissolving the powder in dimethyl sulfoxide as instructed by the manufacturer. Following
anesthetization, axolotls were administered 20 µL/g of 400 µM EdU in 0.7X PBS via intraperitoneal
injection 18 hours before tissue harvest.

Skeletal and tissue preparations

For tissue sections, post-collection, tissue samples were fixed in 4% paraformaldehyde in DEPC-treated
PBS overnight at 4°C. Tissues were then cryopreserved in a sucrose gradient following a wash in DEPC-
PBS and incubated in 30% sucrose/DEPC-PBS at 4°C overnight. Tissues were then embedded and frozen
on dry ice in Tissue-Tek O.C.T. Compound (Sakura) and stored at −80°C. All blocks were sectioned at
16µm thickness using a Leica CM 1950 cryotome and sections were stored at −80°C until use.

For skeletal preparations, post-collection, limbs incubated overnight in 95% ethanol at room
temperature and then incubated in 100% acetone under the same conditions. Skeletal elements were
visualized with alcian blue and alizarin red S stains followed by a clearing of non-skeletal tissue with
potassium hydroxide.

Immunofluorescence

Sections were rehydrated with PBS and then permeabilized with 0.5% Triton-X-100 PBS. Post-
permeabilization, slides were rinsed twice with PBS and EdU visualization was performed via sulfo-
cyanine 3 azide (Lumiprobe). Post-EdU visualization slides were washed twice with PBS. Staining and
labeling with any additional antibodies was conducted post-EdU labeling. Primary antibodies used were
rabbit anti-phospho-S6 ribosomal protein (Ser235/236, 1:200; Cell Signaling 4858), mouse anti-NeuN
(1:200; Abcam ab104224), and rabbit anti-keratin 10 (BioLegend 905401). Secondary antibodies used
were Cy3-goat anti-mouse (1:250, Jackson ImmunoResearch 115-165-146), Alexa Fluor 488-Goat Anti-
Rabbit (1:250, Jackson ImmunoResearch, 111-545-003) and Cy3-goat anti-rabbit (1:250, Jackson
ImmunoResearch 111-165-003).

Sections stained with all antibodies were blocked with 10% serum from the host species of the
secondary antibody in 0.1% Triton-X-100 in PBS. Primary and secondary antibodies were diluted in in 2%
BSA, 0.1% Triton-X-100 in PBS. Sections labeled with anti-pS6 and NeuN underwent an antigen retrieval
step after permeabilization, in which slides were boiled in 0.1 M sodium citrate (pH 6.0) and then rinsed
once in PBS, before EdU was visualized, followed by blocking and primary antibody incubation. All slides
were incubated with DAPI (Sigma D9542) before cover slipping.

Western blot

299 Samples were harvested and blinded, protein extracted and western blots performed as in [22]. Primary
300 antibodies used were anti-pS6 (1:2000, Cell Signaling Technology 4858), anti-S6 (1:1000, Cell Signaling
301 Technology 2317S), and α-tubulin (1:2000, Sigma-Aldrich T9026); secondary antibodies used were goat
302 anti-rabbit HRP (1:5000, Jackson ImmunoResearch 111-035-003) and goat anti-mouse HRP (1:5000, Bio-
303 Rad 1721019).

Imaging and quantification

Imaging was performed on a Zeiss Axio Scan.Z1 automated slide scanner, Leica M165 FC equipped with
Leica DFC310 FX camera, and a Nikon Eclipse Ni microscope with a DS-Ri2 camera. Quantification of all
images was conducted blindly. For imaging of intact contralateral limbs, representative sections
displaying all tissue types (bone and cartilage, soft internal tissues, and epidermis) were selected. For
the tissue type analysis of EdU+ nuclei, tissues were classified as epidermis, "skeletal elements" (bone,
cartilage), and "soft tissues" (muscle, joint, tendon, ligaments, dermis, vasculature, and nerves) based
on nuclear morphology and location as previously described (Whited et al, 2013). All cell counts were
conducted utilizing CellProfiler [51]. Measurements of blastema area were performed using Fiji [52].

Statistical analysis

Statistical analysis of blastema differentiation was conducted using a Fisher's exact test where $p < 0.05$
was considered to be significant. For statistical analysis of the number of digit outlines, a Chi-square test
comparing no digits, 2 digits and 3+ digits distribution between activated and control groups was utilized
in which $p ≤ 0.05$ significant. Statistical analysis of the blastema area and length was performed using an
unpaired, two-tailed Welch's t-test, with Holm-Šídák correction for multiple hypothesis testing, and an
adjusted p-value < 0.05 was considered significant.

For the quantification of %EdU+ nuclei in the limbs of animals in the denervation, long-chase EdU, and
inhibitor experiments the average of both the forelimb and hindlimb were taken and valued as a single
data point. Multiple unpaired, two-tailed Welch's t-tests were used to analyze effect of mTOR inhibition

and denervation, with Bonferroni-Dunn correction for multiple hypothesis testing. For the remaining
experiments, a Welch's ANOVA was used to compare more than 2 groups, or an unpaired two-tailed
Welch's t-test, used to compare exactly two groups, were conducted assuming unequal variance. The
ANOVA was conducted under the parameters of post-hoc pairwise t-tests with Dunnett T3 correction for
multiple hypothesis testing. For all statistical analyses an adjusted-p-value < 0.05 was considered
significant.

RNA isolation, sequencing, and computational analysis

βIIItub:GAP43-EGFP transgenic animals were used for RNA sequencing experiments to allow for easily
identifiable collections of tissues [48]. RNA was extracted from tissue samples with Trizol (15596018,
Ambion) and chloroform (288306, Sigma Aldrich). RNA was then isolated from tissue samples using a
Zymo Clean Up and Concentrator kit (R1013). RNA samples were processed at the Bauer Core Facility at
Harvard University using the Illumina NextSeq 500. Paired End reads were demultiplexed with
bcl2fastq2 v2.2.

The reads were trimmed for quality and adapters with Trimmomatic v0.39 [53] with the following
parameters: LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36. An index of the previously
published axolotl transcriptome [22] was created using Kallisto v0.45.1 [54]. Sequence based bias
correction was performed using Kallisto with 200 bootstraps across 20 threads. The transcript-level
abundances from Kallisto were summarized and imported into R as a matrix of counts using tximport
v1.0.3 [55].

Differentially expressed genes were determined by estimating fold change and dispersion using DESeq2
[56]. Genes were considered significant if they had a logarithmic fold change greater than 1.5 and
adjusted p value of 5%. These significant genes were visualized using pheatmap v1.0.12 [57]. Pathway
analysis was performed in R using pathfinder [58].

Data availability

The sequencing data associated with this study have been deposited in the Gene Expression Omnibus
(GEO) with the accession code GSE192477.

References

1.   Ricci, L. and M. Srivastava, *Wound-induced cell proliferation during animal regeneration.* Wiley
Interdiscip Rev Dev Biol, 2018. **7**(5): p. e321.
2.   Losner, J., K. Courtemanche, and J.L. Whited, *A cross-species analysis of systemic mediators of
repair and complex tissue regeneration.* NPJ Regen Med, 2021. **6**(1): p. 21.
3.   Haas, B.J. and J.L. Whited, *Advances in Decoding Axolotl Limb Regeneration.* Trends Genet, 2017.
33(8): p. 553-565.
4.   McCusker, C., S.V. Bryant, and D.M. Gardiner, *The axolotl limb blastema: cellular and molecular
mechanisms driving blastema formation and limb regeneration in tetrapods.* Regeneration (Oxf),
2015. **2**(2): p. 54-71.
5.   Johnson, K., et al., *Systemic cell cycle activation is induced following complex tissue injury in
axolotl.* Dev Biol, 2018. **433**(2): p. 461-472.

6. Wenemoser, D. and P.W. Reddien, *Planarian regeneration involves distinct stem cell responses*
to wounds and tissue absence. Dev Biol, 2010. **344**(2): p. 979-91.
7. Srivastava, M., et al., *Whole-body acoel regeneration is controlled by Wnt and Bmp-Admp*
signaling. Curr Biol, 2014. **24**(10): p. 1107-13.
8. Rodgers, J.T., et al., *mTORC1 controls the adaptive transition of quiescent stem cells from G0 to*
G(Alert). Nature, 2014. **510**(7505): p. 393-6.
9. Maier, C.E. and M. Singer, *The effect of light on forelimb regeneration in the newt.* J Exp Zool,
1977. **202**(2): p. 241-4.
10. González-Estévez, C., et al., *SMG-1 and mTORC1 act antagonistically to regulate response to*
injury and growth in planarians. PLoS Genet, 2012. **8**(3): p. e1002619.
11. Tu, K.C., B.J. Pearson, and A. Sanchez Alvarado, *TORC1 is required to balance cell proliferation*
and cell death in planarians. Dev Biol, 2012. **365**(2): p. 458-69.
12. Hirose, K., et al., *Mechanistic target of rapamycin complex 1 signaling regulates cell*
proliferation, cell survival, and differentiation in regenerating zebrafish fins. BMC Dev Biol, 2014.
14: p. 42.
13. Haller, S., et al., *mTORC1 Activation during Repeated Regeneration Impairs Somatic Stem Cell*
Maintenance. Cell Stem Cell, 2017. **21**(6): p. 806-818 e5.
14. Molinaro, A.M., N. Lindsay-Mosher, and B.J. Pearson, *Identification of TOR-responsive slow-*
cycling neoblasts in planarians. EMBO Rep, 2021. **22**(3): p. e50292.
15. Gusel'nikova, V.V. and D.E. Korzhevskiy, *NeuN As a Neuronal Nuclear Antigen and Neuron*
Differentiation Marker. Acta Naturae, 2015. **7**(2): p. 42-7.
16. Todd, J.T., *On the process of reproduction of the members of the aquatic salamander. .* The
Quarterly Journal of Science, Literature, and the Arts. , 1823. **16**: p. 84-96.
17. Farkas, J.E. and J.R. Monaghan, *A brief history of the study of nerve dependent regeneration.*
Neurogenesis (Austin), 2017. **4**(1): p. e1302216.
18. Mahmoud, A.I., et al., *Nerves Regulate Cardiomyocyte Proliferation and Heart Regeneration.* Dev
Cell, 2015. **34**(4): p. 387-99.
19. Gao, X., et al., *Nociceptive nerves regulate haematopoietic stem cell mobilization.* Nature, 2021.
589(7843): p. 591-596.
20. Zhang, B., et al., *Hyperactivation of sympathetic nerves drives depletion of melanocyte stem*
cells. Nature, 2020. **577**(7792): p. 676-681.
21. Rodgers, J.T., et al., *HGFA Is an Injury-Regulated Systemic Factor that Induces the Transition of*
Stem Cells into GAlert. Cell Rep, 2017. **19**(3): p. 479-486.
22. Bryant, D.M., et al., *A Tissue-Mapped Axolotl De Novo Transcriptome Enables Identification of*
Limb Regeneration Factors. Cell Rep, 2017. **18**(3): p. 762-776.
23. Leigh, N.D., et al., *von Willebrand factor D and EGF domains is an evolutionarily conserved and*
required feature of blastemas capable of multitissue appendage regeneration. Evol Dev, 2020.
24. Kastan, N., et al., *Small-molecule inhibition of Lats kinases may promote Yap-dependent*
proliferation in postmitotic mammalian tissues. Nat Commun, 2021. **12**(1): p. 3100.
25. Chen, P., et al., *Dexmedetomidine promotes the progression of hepatocellular carcinoma*
through hepatic stellate cell activation. Exp Mol Med, 2020. **52**(7): p. 1062-1074.
26. Wang, W., X. Guo, and H. Dan, *alpha2A-Adrenergic Receptor Inhibits the Progression of Cervical*
Cancer Through Blocking PI3K/AKT/mTOR Pathway. Onco Targets Ther, 2020. **13**: p. 10535-
10546.
27. Blanchard, R.J., et al., *Yohimbine potentiates active defensive responses to threatening stimuli in*
Swiss-Webster mice. Pharmacol Biochem Behav, 1993. **44**(3): p. 673-81.
28. Ito, K., et al., *Endogenous alpha2A-Adrenoceptor-Operated Sympathoadrenergic Tones*
Attenuate Insulin Secretion via cAMP/TRPM2 Signaling. Diabetes, 2017. **66**(3): p. 699-709.

29. Mantsch, J.R., et al., *Involvement of noradrenergic neurotransmission in the stress- but not*
cocaine-induced reinstatement of extinguished cocaine-induced conditioned place preference in
mice: role for beta-2 adrenergic receptors. Neuropsychopharmacology, 2010. **35**(11): p. 2165-78.
30. Phan, Q.M., et al., *Lef1 expression in fibroblasts maintains developmental potential in adult skin*
to regenerate wounds. Elife, 2020. **9**.
31. Poss, K.D., J. Shen, and M.T. Keating, *Induction of lef1 during zebrafish fin regeneration.* Dev
Dyn, 2000. **219**(2): p. 282-6.
32. Kawakami, Y., et al., *Wnt/beta-catenin signaling regulates vertebrate limb regeneration.* Genes
Dev, 2006. **20**(23): p. 3232-7.
33. Tonkin, J., et al., *Monocyte/Macrophage-derived IGF-1 Orchestrates Murine Skeletal Muscle*
Regeneration and Modulates Autocrine Polarization. Mol Ther, 2015. **23**(7): p. 1189-1200.
34. Schertzer, J.D., et al., *Modulation of insulin-like growth factor (IGF)-I and IGF-binding protein*
interactions enhances skeletal muscle regeneration and ameliorates the dystrophic pathology in
mdx mice. Am J Pathol, 2007. **171**(4): p. 1180-8.
35. Mitsiades, C.S., et al., *Inhibition of the insulin-like growth factor receptor-1 tyrosine kinase*
activity as a therapeutic strategy for multiple myeloma, other hematologic malignancies, and
solid tumors. Cancer Cell, 2004. **5**(3): p. 221-30.
36. Butler, E.G., and O'Brien, J. P., *Effects of localized x-radiation on regeneration of the urodele*
limb. Anatomial Record, 1942. **84**(4): p. 407-413.
37. Rothenberger, M.B., Vera, M. K., Germanoski, D., Ramirez, E. , *Comparing amphibian habitat*
quality and functional success among natural, restored, and created vernal pools. Restoration
Ecology, 2019. **27**(4).
38. Pfennig, D.W. and J.P. Collins, *Kinship affects morphogenesis in cannibalistic salamanders.*
Nature, 1993. **362**(6423): p. 836-8.
39. Collins, J.P., Cheek, J. E. , *Effect of food and density on development of typical and cannibalistic*
salamander larvae in Ambystoma tigrrinum nebulosum. Ameican Zoologist, 1983. **23**: p. 77-84.
40. Thompson, S., et al., *Probability of Regenerating a Normal Limb After Bite Injury in the Mexican*
Axolotl (Ambystoma mexicanum). Regeneration (Oxf), 2014. **1**(3): p. 27-32.
41. Bothe, V., K. Mahlow, and N.B. Frobisch, *A histological study of normal and pathological limb*
regeneration in the Mexican axolotl Ambystoma mexicanum. J Exp Zool B Mol Dev Evol, 2021.
336(2): p. 116-128.
42. Dent, J.N., *Limb regeneration in larvae and metamorphosing individuals of the South African*
clawed toad. J Morphol, 1962. **110**: p. 61-77.
43. Yakushiji, N., H. Yokoyama, and K. Tamura, *Repatterning in amphibian limb regeneration: A*
model for study of genetic and epigenetic control of organ regeneration. Seminars in Cell &
Developmental Biology, 2009. **20**(5): p. 565-574.
44. Egar, M., S.B. Simpson, and M. Singer, *The growth and differentiation of the regenerating spinal*
cord of the lizard, Anolis carolinensis. J Morphol, 1970. **131**(2): p. 131-51.
45. Simpson, S.B., Jr., *Morphology of the regenerated spinal cord in the lizard, Anolis carolinensis.* J
Comp Neurol, 1968. **134**(2): p. 193-210.
46. Simpson, S.B., Jr., *Studies on regeneration of the lizard's tail.* Am Zool, 1970. **10**(2): p. 157-65.
47. Lozito, T.P. and R.S. Tuan, *Lizard tail skeletal regeneration combines aspects of fracture healing*
and blastema-based regeneration. Development, 2016. **143**(16): p. 2946-57.
48. Khattak, S., et al., *Germline transgenic methods for tracking cells and testing gene function*
during regeneration in the axolotl. Stem Cell Reports, 2013. **1**(1): p. 90-103.
49. Schotté, O.E. and E.G. Butler, *Morphological effects of denervation and amputation of limbs in*
urodele larvae. Journal of Experimental Zoology, 1941. **87**(2): p. 279-322.

50.   Kropf, N., et al., *Sciatic nerve injury model in the axolotl: functional, electrophysiological, and
radiographic outcomes.* J Neurosurg, 2010. **112**(4): p. 880-9.
51.   McQuin, C., et al., *CellProfiler 3.0: Next-generation image processing for biology.* PLOS Biology,
2018. **16**(7): p. e2005970.
52.   Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nature Methods,
2012. **9**(7): p. 676-682.
53.   Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina sequence
data.* Bioinformatics, 2014. **30**(15): p. 2114-20.
54.   Bray, N.L., et al., *Near-optimal probabilistic RNA-seq quantification.* Nat Biotechnol, 2016. **34**(5):
p. 525-7.
55.   Soneson, C., M.I. Love, and M.D. Robinson, *Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences.* F1000Res, 2015. **4**: p. 1521.
56.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.
57.   Kolde, R., *pheatmap: pretty heatmaps. R package version 1.0. 12. 2019*. 2019.
58.   Ulgen, E., O. Ozisik, and O.U. Sezerman, *pathfindR: An R Package for Comprehensive
Identification of Enriched Pathways in Omics Data Through Active Subnetworks.* Front Genet,
2019. **10**: p. 858.

Acknowledgements

This work was supported by The Richard and Susan Smith Family Odyssey Award (J.L.W.), the NIH New
Innovator Award DP2HD087953 (J.L.W.) and Harvard University Faculty of Arts and Sciences (J.L.W.). We
thank William Ye, Tyler Mourey, Kelly Dooling and Rui Qun Miao for animal care. We thank Matthew
Warman, Marc Kirschner, Ya-Chieh Hsu, Davie Van Vactor, Rich Lee, Paul Garrity, and members of the
Whited, Chiu, Greer and Lehoczky labs for discussions and comments on the manuscript. We thank Yulia
Swartz for sharing reagents and assisting in data interpretation, Bobby Groves and Angelo Osofsky for
technical assistance, and the Bauer Core Facility at Harvard University for sequencing.

Author Contributions

J.L.W., D.P.-D and S.E.W. conceived of the project. D.P.-D, S.E.W., B.E., S.H., J.L., J.V.M, and J.L.W.
designed the experiments; D.P.-D. S.E.W., and J.L.W. wrote the paper; S.J.B. performed computational
analyses; L.C. performed statistical analyses; D. P.-D., S.E.W., B.E., S.H., J.V.M., A.W., J.L., S.M and H.S.
performed sample preparation, experiments, and experimental data analyses. All authors contributed to
manuscript editing.
491
492 **Competing Interests**
493
494 J.L.W. is a co-founder of Matice Biosciences. Other authors declare no conflicts of interest.
495
496 **Figures**

a

Control

+7 days → not activated → amputate both right limbs →

Activated

amputate both left limbs → +7 days → activated → amputate both right limbs →

Compare regeneration rates of right limbs

b

	7 dpa	14 dpa	21 dpa	25 dpa	25 dpa

Control

Activated

c

P = 0.0264

Blastema Area (mm^2)

Control   Activated

d

P = 0.004

Number of limbs

Control   Activated

■ Differentiation
▨ No Differentiation

e

P = 0.0014

Number of digit outlines per limb

Control   Activated

□ 4
▨ 3
▨ 2
■ 0

Figure 1

Figure 1. Systemically-activated cells are primed for future regeneration events.
a, Experimental design. We compared regeneration rates of systemically-activated and naïve, control
limbs. Pink dots, activated, EdU positive nuclei. **b**, Representative images of regenerating systemically-
activated or control limbs. **c-e**, Quantification of blastema area at 14 days post-amputation **(c)**,
differentiating limbs **(d)**, and number of digit outlines **(e)** at 25 days post-amputation (n=15 animals for
each condition in blastema area measurement, analyzed by two-tailed Welch's t-test, n = 30 limbs for
each condition in differentiation and digit outline quantification, analyzed by Fisher's exact test for
differentiation and Chi-square test for digit outlines). All data shown as mean ± SD. Scale bars, 2 mm.
dpa, days post-amputation. EdU, 5-ethynyl-2'-deoxyuridine.
511

Figure 2

Figure 2. mTOR signaling is required for systemic activation. a, pS6 and EdU staining of blastema, intact (homeostatic) and contralateral limb tissue sections from animals treated with saline, vehicle, or rapamycin at 14 dpa. **b**, Quantification of EdU positive nuclei at 14 dpa. (n = 12 limbs for intact saline, intact vehicle and intact rapamycin groups, n = 10 limbs for amputated saline and amputated vehicle, and n = 8 limbs for amputated rapamycin groups, analyzed by unpaired two-tailed Welch's t-tests with Bonferroni-Dunn correction). All data shown as mean ± SD. Scale bar, 500 µm. rapa, rapamycin. Int, intact. contra, contralateral. dpa, days post-amputation. pS6**,** phospho-S6 ribosomal protein; EdU, 5-ethynyl-2'-deoxyuridine.

Figure 3

Figure 3. Nerves are required for systemic activation. a, Experimental design. Nerve bundles innervating limbs contralateral to unilateral amputations were transected or sham-operated, and animals were assayed for systemic activation. Animals receiving unilateral denervations or sham-operations with no amputations were used as controls. **b**, EdU and DAPI staining of intact and contralateral limb tissue sections of denervated and sham-operated animals in the intact state or at 14 days post amputation. **c-f**, Quantification of percent EdU positive nuclei in denervated and sham-operated limbs **(c)**, epidermis **(d)**, skeletal elements **(e)**, and soft internal tissues **(f)** at 14 dpa. (n = 11 animals for sham, n = 10 animals for denervated, n = 13 animals for sham amputated and denervated amputated, analyzed by unpaired two-tailed Welch's t-tests with Bonferroni-Dunn correction). All data shown as mean ± SD. Scale bar, 500 µm. den, denervated. int, intact. contra, contralateral. EdU, 5-ethynyl-2'-deoxyuridine.

Figure 4

Figure 4. Transcriptome analyses reveal pathways regulating systemic activation. Nerves servicing limbs contralateral to unilateral amputations were transected or sham-operated. Nerve ends and the immediate tissue surrounding them were collected 14 days post amputation. **a**, Differential gene expression analysis identifies transcripts that are enriched in nerve ends (left) and responding tissues (right). **b**, EdU and DAPI staining of limbs contralateral to unilateral amputations treated with yohimbine or vehicle control, harvested at 7 days post amputation. **c**, Quantification of percent EdU positive nuclei in contralateral limbs in **(b)** (n = 15 animals for each group, analyzed by unpaired two-tailed Welch's t-test). All data shown as mean ± SD. Scale bar, 500 μm. EdU, 5-ethynyl-2'-deoxyuridine. **d**, Model summarizing the main findings of the study. Limbs activated by a prior contralateral amputation regenerate faster than naïve, control limbs. This effect is mediated by nerves, mTOR signaling, and adrenergic signaling.

Extended Data Figure 1

545   **Extended Data Figure 1. A subset of activated cells are epidermal stem cells.** Keratin 10, EdU and DAPI
546   staining of limbs contralateral to amputation harvested 14 days post amputation, 18 hours after EdU
547   administration**.** Data shown as mean ± SD. Scale bars, 200 μm. K10, keratin 10. EdU, 5-ethynyl-2'-
548   deoxyuridine.
549
550

Extended Data Figure 2

551   **Extended Data Figure 2. Activated cells in distant limbs form clones. a,** EdU and DAPI staining of intact
552   limbs (left) and limbs contralateral to amputation (right) harvested 6 weeks after EdU administration. **b,**
553   Quantification of percent EdU positive nuclei of intact and contralateral limbs. (n = 15 animals for intact
554   and n = 12 animals for contralateral group, analyzed by unpaired two-tailed Welch's t-test). Data shown
555   as mean ± SD. Scale bar, 500 µm. EdU, 5-ethynyl-2'-deoxyuridine.
556

a

Vehicle | Rapamycin

b

Control
Rapamycin

Length of Blastema (mm)

days post amputation

P = 0.0365

c

Vehicle

Rapamycin

Extended Data Figure 3

Extended Data Figure 3. Rapamycin-treated animals produce smaller blastemas.

a, Regenerating limbs at 7 dpa treated with rapamycin or vehicle control. Arrowheads mark the

amputation plane. **b**, Blastema growth curves of rapamycin or vehicle treated animals. (n = 10 animals

per group, analyzed by unpaired two-tailed Welch's t-test with Holm-Šídák correction). **c**,

Representative skeletal preparations of regenerated limbs with Alcian blue and Alizarin red at 10 weeks

post amputation. Scale bars, 0.5 mm.

Intact **Regenerating**

Extended Data Figure 4

565 **Extended Data Figure 4. mTOR signaling is upregulated in the brains of animals with regenerating**
566 **limbs.** pS6 and NeuN staining of anterior telencephalon sections of animals with intact (left) or
567 regenerating (right) limbs harvested at 7 days post amputation. Scale bar, 200 μm. pS6**,** phospho-S6
568 ribosomal protein. NeuN, neuronal nuclear protein.

a

same animal

| Intact | Regenerating | Contralateral |

pS6 DAPI

NeuN DAPI

pS6 NeuN DAPI

b

Regenerating   Contralateral

1 2 3   1 2 3

pS6

total S6

α-tubulin

Extended Data Figure 5

Extended Data Figure 5. mTOR signaling is differentially regulated in the dorsal root ganglia of sensory
nerves innervating the regenerating limb. a, pS6 and NeuN staining of dorsal root ganglia servicing
intact, regenerating, and contralateral limbs at 7 days post amputation. **b**, Western blot of pS6 and total
S6 protein levels in dorsal root ganglia servicing regenerating and contralateral limbs harvested at 3 days
post amputation. Alpha-tubulin serves as loading control. Lanes with same numbers correspond to
dorsal root ganglia harvested from the same animal. Scale bar, 200 μm. pS6**,** phospho-S6 ribosomal
protein. NeuN, neuronal nuclear protein.

a

Vehicle | IWR-1 | NVP-ADW742

EdU DAPI

b

P = 0.0263

P = 0.9501

%EdU+ nuclei

Vehicle | IWR-1 | NVP-ADW742

Extended Data Figure 6

Extended Data Figure 6. Test of additional candidate pathways regulating systemic activation. a, EdU and DAPI staining of limbs contralateral to unilateral amputations treated with tankyrase inhibitor IWR-1, Igf1r inhibitor NVP-ADW742 or vehicle control, harvested at 7 days post amputation. **b,** Quantification of percent EdU positive nuclei of contralateral limbs in **(a)**. (n = 11 animals for vehicle control, n = 14 animals for IWR-1 and NVP-ADW742, analyzed by Welch's ANOVA with Dunnett's T3 correction, W = 4.883). All data shown as mean ± SD. Scale bar, 500 μm. EdU, 5-ethynyl-2'-deoxyuridine.