1   **Adrenergic signaling stimulates body-wide stem cell activation for limb regeneration**
2
3

4   Duygu Payzin-Dogru[1#], Sarah E. Wilson[1#], Steven J. Blair[1], Aaron M. Savage[1], Emil Kriukov[2,3], Victor
5   Cat[1], Louis Cammarata[4], Burcu Erdogan[1], Shifa Hossain[1], Noah Lopez[1], Julia Losner[1], Juan Velazquez
6   Matos[1], Sangwon Min[1], Kelly Dooling[1], Bobby Groves[1], Alan Y. Wong[1], Hani Singer[1], Petr Baranov[2,3],
7   Isaac M. Chiu[5], Brian J. Haas[6], Jessica L. Whited[1,6,7*]
8
[1] Department of Stem Cell and Regenerative Biology, Harvard University, Harvard University, 7 Divinity
Ave., Cambridge, MA, USA, 02138
[2] Schepens Eye Research Institute of Massachusetts Eye and Ear, Boston, MA, USA, 02114
[3] Department of Ophthalmology, Harvard Medical School, Boston, MA, USA, 02114
[4] Department of Statistics, Harvard University, Cambridge, MA, USA, 02138
[5] Department of Immunology, Blavatnik Institute, Harvard Medical School, Boston, MA, USA, 02115
[6] The Broad Institute of Harvard and MIT, Cambridge, MA, USA, 02138
[7] Harvard Stem Cell Institute, Harvard University, Cambridge, MA, USA, 02138
17
18   [*] Correspondence: Jessica L. Whited, jessica_whited@harvard.edu
19
20   [#] These authors contributed equally to this work

21
22
23   **SUMMARY**
24

Many species throughout the animal kingdom naturally regenerate complex body parts following
amputation. Most research in appendage regeneration has focused on identifying mechanisms that
influence cell behaviors in the remaining stump tissue, immediately adjacent to the injury site. Roles for
activation steps that occur outside of the injury site remain largely unexplored, yet they may be critical for
successful regeneration. Here, we discovered a role for adrenaline in stimulating a body-wide stem cell
activation response to amputation that drives limb regeneration. Notably, this systemic response is
mediated by engagement of the peripheral nervous system (PNS) and adrenergic signaling at both the
injury site and in distant, uninjured tissues. This work challenges the predominant conceptual framework
considering the injury site alone in the regenerative response, arguing instead for body-wide stem cell
activation as a priming step upon which molecular cues at the injury site then build tissue.

35
36
37   Keywords: regeneration, systemic responses, adrenaline, mTOR, stem cells, progenitor cells, limb,
38   amputation
39

INTRODUCTION

Replacing large body parts following severe injury is a long-term goal of regenerative medicine, and one which is likely to stand on first elucidating how other animals execute this enormous task. Salamander limbs are anatomically quite similar to human limbs, yet they can be spontaneously regenerated throughout life following amputation.[1,2] Recent studies have revealed some clues about the transcriptional regulation that accompanies salamander limb regeneration, along with evidence that particular genes and molecules are important to the process.[3-7] Yet, these kinds of studies have thus far not provided critical clues about how organismal-level signaling throughout the body might be critical to regeneration. Salamanders grow an essential and outwardly-visible tissue primordium called a "blastema," which contains the progenitor cells required to grow a whole limb, while humans and other mammals do not.

Prior studies have shown that invertebrates undergo broad proliferation body-wide prior to the response becoming restricted to the site of amputation, at which point blastema formation becomes apparent.[8-11] Broad activation followed by local blastema formation as a strategy for regeneration could also underlie limb regeneration in vertebrates, but thus far this hypothesis has not been tested. If some species do use this strategy to effectively regenerate limbs, the discrete activation and blastema specification steps could be examined in detail in mammals to determine which components might occur in mammals despite their overall inability to naturally regenerate limbs. Here, we explored these concepts and effectively decoupled body-wide activation responses from subsequent blastema formation during limb regeneration. We find that injury-induced activation of adrenergic signaling critically mediates both local and systemic progenitor cell proliferation to prime body-wide tissue-resident cells for limb regeneration.

RESULTS

Amputation primes distant tissues for faster future regeneration

We recently discovered that following axolotl limb amputation, cells located throughout the body, including within contralateral unamputated limbs, proliferate, in a "systemic activation" response.[12] We also showed that systemic activation is an amputation-injury-induced response *per se*, rather than a consequence of the regeneration process, by amputating limbs and surgically blocking their regeneration.[12] Here, we first sought to understand what the consequences of systemic activation are to animals that have recently undergone amputation. We hypothesized that distant tissues harboring these systemically-activated cells might exhibit altered regeneration potentials or kinetics. We therefore challenged axolotl limbs in a systemically-active state to a local amputation 7 days later (**Figure 1A**). We found that limbs engaged in systemic activation regenerate faster than those limbs on naïve animals that had never undergone amputation (**Figure 1B-C**). We observed this effect to be significant at the early, initiation stages of regeneration (**Figure 1C**), when the blastema is being built. The effect continued through later stages of regeneration, when the blastema commences differentiation (**Figure 1D**) and digits become discernable (**Figure 1E**). We also tested whether molecular markers of blastema formation are accelerated following amputation of primed limbs versus naïve limbs. We performed HCR-FISH to detect *Prrx1* mRNA expression[13] at 5 days post-amputation (dpa) (**Figure 1F**), and we observed a significant increase in the signal intensity of this blastema-cell marker in the primed limbs versus naïve limbs (**Figure 1F-G**). This data supports the conclusion that the consequence of amputation-induced systemic activation is a priming of distant tissues for faster regeneration when subjected to future local injury.

We also investigated how many divisions cells undergo following their systemic activation and how long the priming effect lasts. Using a short-pulse (EdU) and long-chase strategy, we found most

89 labeled clones in limbs to be composed of 2-8 cells (**Figure 1H**), indicating amputation-induced
90 proliferation may last only a few cell cycles. Concordantly, we found there to be no difference in the
91 fraction of EdU+ cells by 4 weeks post-amputation (**Figure 1I**). We also found that systemically-activated
92 limbs challenged with local amputation 4 weeks after initial activation regenerated at the same rate as
93 naïve limbs, indicating the effect is relatively short-lived (**Figure 1J-K**).
94
95
96 **Systemically-activated cell types are shared with proliferating cell types in homeostasis**
97
Defining which cells are activated is key to developing hypotheses about how systemic activation
relates to regeneration. We therefore asked which types of cells become systemically activated in
response to amputation. Using immunohistochemistry, we also found epidermal stem cells to participate
in the response (**Figure 1M**). To account for the full repertoire of activated cells that we observed, we
turned to a more agnostic method of addressing this question. We used FACS and the nuclear stain
Hoecsht to sort cells with 4C nuclear content from systemically-activated limbs and from homeostatic
limbs (**Figure 2A-B**). We identified cell types in both samples via single-cell RNAseq and downstream
clustering analyses (**Figure 2C**, **Table S1**). The distribution of marker gene expression across clusters is
shown in **Figure S1A**. In the systemically-activated samples, we found multiple blood cell types, including
myeloid progenitors, T cells, two clusters of hematopoietic progenitors, and two clusters of red blood cells
(**Figure 2C**). We also identified skeletal muscle progenitors, two types of endothelial cells, basal
epidermal cells, and four other types of epidermal cells (**Figure 2C**). Previously, we had identified satellite
cells, stem cell populations that support muscle regeneration in many vertebrates,[14] as being among
those cells that become activated,[12] but due to the limitations of our dissociation protocol, we did not
recover them here. We parsed eight subtypes of activated fibroblasts (**Figure 2C**). Intriguingly, the same
clustering algorithm applied to the 4C cells purified from homeostatic animals produced the exact same
number and repertoire of cell clusters (**Figure. 2C**). This result indicates that axolotls systemically activate
the same cell types that cycle during homeostasis, boosting the number of these cells actively in S-phase
after amputation.
We performed differential gene expression analysis between the two conditions for each cell
cluster (**Table S2**). We focused on fibroblasts because they are known to play a central role in
contributing to blastema formation during limb regeneration,[15,16] and connective tissues have been shown
to exhibit more plasticity across the limb regeneration process than other tissues.[17,18] We identified 19
genes to be significantly (p<0.05) different in expression in systemically-activated type IV fibroblasts
versus 4C type IV fibroblasts from homeostatic animals (**Figure 2E**), while an additional 44 genes were
differentially expressed with 0.05<p<0.1. We detected no significant differences in gene expression in
other fibroblast cell types between systemically-activated and correlating 4C cells from homeostatic
animals (other than a single unannotated gene whose expression differs among the two conditions in
fibroblast II). Other cell types for which we uncovered significant differential gene expression included one
type of epidermal cells and one type of endothelial cells; red blood cells; myeloid progenitor and
hematopoietic progenitor cells (**Figure 2D**). For the type IV fibroblasts, 8/19 genes with p<0.05 and 20/44
genes with 0.05<p<0.1 are predicted to encode ribosomal subunits (**Figure 2E**, **Table S2**). 27/28 of these
ribosomal subunits were more highly expressed in systemically-activated fibroblast IV cells than their 4C
homeostatic counterparts. The most significant differentially expressed gene (p<0.0059) to display this
relationship is predicted to encode Snail (**Figure 2E**), a factor most known for its promotion of epithelial-
to-mesenchymal transition (EMT) in a variety of contexts,[19-21] was expressed 1.6-fold higher in
systemically-activated fibroblast IV cells. We also found the p53 inhibitor Sestrin 1,[22] which also regulates
mTORC1 and mTORC2 in other species by acting as a nutrient sensor,[23] to be similarly significantly
upregulated (p<0.0059), with a 2-fold enrichment in the systemically-activated fibroblast IV cells
compared to their 4C homeostatic counterparts.

To gain insights into possible cell state trajectories, we used the 4C scRNA-seq dataset to perform cell fate reconstruction analysis using RNA Velocity (**Figure 2F**).[24] In both homeostatic and systemically-activated conditions, we found evidence for fibroblast II and IV populations potentially transitioning to fibroblast I cell state, which might be enhanced in the systemically-activated samples (**Figure 2F, Figure S1C**). Genes that are most likely driving these changes are highlighted in **Figure S1D**; notably, nearly all of these genes remain unannotated in the axolotl genome, with no known high-confidence orthologs. Similarly, applying gene dynamics analysis to 4C cells dataset identified a handful of transcripts with unique regulation in the kinetics of RNA transcription and splicing (**Figure S1E**), although we again found almost all of these genes to be unannotated.

We additionally analyzed the single cells not undergoing systemic activation, those with 2C nuclear content (**Figure 2G**). We did not find any difference in the repertoire of cell types from these two conditions, 2C cells from homeostatic versus 2C cells from activated limb samples (**Figure 2G, Figure S1D**). Major transcriptional differences were observed mainly within the epidermal cell populations and small secretory cell populations (which reside in skin), indicating that a global transcriptional reprogramming of skin may occur downstream of amputation (**Figure 2H, Table S3**). Collectively, the data indicate that amputation induces a subset of cell types throughout the body that ordinarily can proliferate during homeostasis to become hyper-proliferative, thus becoming systemically activated.

mTOR signaling is required for systemic activation and priming

Our previous work demonstrated that mTOR signaling is upregulated in distant tissues, such as contralateral limbs, during systemic activation.[12] We find this interesting because this pathway can be activated following injury in mice[25,26] despite the fact that they do not regenerate limbs. Additionally, we used CellChat[27] and our scRNA-seq dataset to identify possible signaling pathways that may drive differences between 4C cells in the systemically-activated versus homeostatic conditions, and we found substantial differences in Hepatocyte growth factor (HGF) signaling, which acts to stimulate mTOR signaling in mice and in human cells (**Figure 3A**).[25,26,28] We therefore tested whether axolotls use mTOR signaling for systemic activation and priming. We treated axolotls with the mTOR inhibitor rapamycin,[29] and we found that rapamycin wholly blocked systemic activation of cells in contralateral limbs, as indicated by significantly fewer EdU+ nuclei (**Figure 3B-C**). This finding led us to ask if mTOR signaling is also required for priming of distant tissues for faster regeneration. We tested this hypothesis (**Figure 3D**), and we found that mTOR signaling is required for priming axolotl limbs for faster regeneration following amputation elsewhere (**Figure 3E**). This requirement is evident both at the early stages of blastema initiation (**Figure 3F**) and at later stages when limbs start to differentiate (**Figure 3G**). Separately, we also asked whether mTOR is required for limb regeneration, in the context of a single amputation. We found that treating axolotls with rapamycin for a time period of two days prior to through 7 days after limb amputation led to a diminished rate of early blastema growth (**Figure S2A-B**), but, ultimately, these limbs regenerated and were outwardly normal at 10 weeks post-amputation (**Figure S2C**). While mTOR activation in mice has been shown to be mediated via molecular messengers in the circulation,[25,26] we were interested in understanding whether mTOR signaling may be activated within the nervous system to regulate systemic activation in axolotls. We examined mTOR activation in dorsal root ganglia (DRG), which include somatosensory neurons that innervate and receive sensory input from the limbs as well as non-neuronal cells such as fibroblasts and satellite glia (**Figure 3H**).[30] While there was no significant difference in the proliferation of cells in the DRGs following amputation (**Figure 3I**), we found that DRGs from both amputated limbs and contralateral limbs showed higher levels of mTOR activation as indicated by pS6+ cells than those from homeostatic animals (**Figure 3J**).

Peripheral nervous system is a conduit for systemic proliferative response to amputation

Determining the mode by which amputation-induced signals spread throughout the body is
important for unraveling the molecular mechanisms of the systemic activation response. We hypothesized
that the peripheral nervous system (PNS) might be a conduit for this type of information, as it transduces
sensory information to the brain and spinal cord, thereby facilitating communication between distant
locations in the body. This hypothesis was strengthened by the upregulation in mTOR signaling in DRGs
we observed after amputation. We tested whether systemic activation requires peripheral innervation at
either the responding site (**Figure 4A-C**) and/or the amputation site (**Figure 4D-F**) using surgical
denervation. We found that systemic activation did not occur when these distant sites were denervated at
the time that contralateral limbs were amputated (**Figure 4A-C**). As a control, we found systemic
activation still occurred following amputation when these distant sites (contralateral limbs) underwent
sham operations. We parsed the contralateral tissues into three broad bins—epidermis, skeletal
elements, and internal (non-skeletal, soft) tissues—and quantified the extent of systemic activation within
each (**Figure 4D-F**). This analysis revealed each of these three bins of tissue types requires innervation
for cellular activation. We also found that when limbs were denervated then immediately amputated, the
amputation no longer provoked systemic activation in distant sites (contralateral limbs) (**Figure 4G-I**).
Sham surgeries in which limbs were not denervated prior to their amputation still provoked the systemic
activation response. These results demonstrate that peripheral innervation of both the amputated limb
and the distantly-responding limb are required for systemic activation of progenitor cells. Therefore, both
neuronal input that senses injury and neuronal output that mediates efferent signaling to tissues could
regulate systemic activation. To explore molecular pathways that may underlie PNS dependence in
responding tissues, we performed two types of bulk RNA-seq analyses. We compared gene expression in
limb nerve bundles from animals actively engaged in systemic activation to those who failed to
systemically activate due to denervation (**Figure 4J**). We identified 273 genes with substantially different
expression between these two conditions (**Figure 4J**, **Table S5**). We also performed differential gene
expression analysis between samples derived from the soft tissue of systemically-activated limbs and the
soft tissue of limbs that were denervated and therefore not activated (**Figure 4K**). This analysis identified
82 differentially-expressed genes (**Figure 4K**, **Table S5**).
214
215 **a-Adrenergic signaling drives systemic activation and priming**
216
We next used the candidate genes revealed by the RNA-seq of nerve and surrounding tissue to
functionally test pathways to determine if they are required for systemic activation. We prioritized well-
studied pathways with robust pharmacological inhibitors. *Adra2a*, which encodes a-2A adrenergic
receptor, was 7.4-fold upregulated in transected nerve samples, which are deficient in stimulating
systemic activation, compared to intact nerves (**Figure 4J**, **Table S5**), implicating a possible role for
adrenergic signaling in the systemic activation response to amputation. Separately, we investigated
adrenergic receptor signaling in our scRNA-seq dataset, and we found many adrenergic receptors to be
widely expressed in both homeostasis and activated conditions (**Figure S3**). The primary neurotransmitter
mediators of adrenergic signaling are the stress-related hormones epinephrine and norepinephrine.[31] We
therefore quantified levels of these molecules in axolotl plasma using ELISA. We found that while
circulating levels of norepinephrine are similar between unamputated axolotls and axolotls at four hours
post-amputation (**Figure 5A**), the amputated animals exhibited approximately 2-fold more epinephrine in
plasma (**Figure 5B**). To address function, we treated axolotls with yohimbine, a well-studied Adra2a
antagonist,[32,33] to test for an adrenergic signaling requirement in systemic activation. We found that in
animals treated with yohimbine, limbs contralateral to amputation did not undergo systemic activation,
demonstrating that adrenergic signaling is required for systemic stem cell activation in response to
amputation (**Figure 5C-D**). We then asked whether adrenergic signaling is also required for priming
distant tissues for future regeneration (**Figure 5E**). We found that when axolotls were treated with
yohimbine through the time period 2 days before to 7 days after amputation, priming was diminished

(**Figure 5F**), with both initial blastema growth (**Figure 5G**) and limb differentiation (**Figure 5H**) occurring
more slowly. We tested whether stimulating adrenergic signaling via the agonist clonidine might enhance
priming (**Figure 5I**), and we found evidence that this pathway can indeed be stimulated to accelerate the
regeneration of distant tissues that have been systemically activated (**Figure 5J-K**). Our finding that a2-
adrenergic-receptor signaling mediates systemic activation prompted us to ask whether b-adrenergic-
receptor signaling also contributes to systemic activation. We therefore used an identical treatment
regimen to administer the non-selective b-adrenergic receptor antagonist propranolol [34] to axolotls
(**Figure 5L-N**). We found that while propranolol treatment led to a significantly decreased fraction of cells
actively in S-phase within blastemas of regenerating limbs (**Figure 5L-M**), it had no effect on systemic
activation (**Figure 5L, 5N**). These results indicate that a-adrenergic signaling is required for the systemic
activation response to amputation in axolotls, but b-adrenergic signaling is only required for the localized
cellular proliferation response at the amputation site.
In parallel, we also tested whether systemic activation requires insulin and Wnt signaling as both
Igfr1 and Lef1 were differentially regulated (in surrounding tissue and nerves, respectively) in denervated
limbs versus sham-operated limbs following amputation **(Table S5)**. CellChat analysis also supported
differential regulation of Insulin/IGF signaling between homeostatic and activated states in the 4C
populations (**Figure S4A-B**). Inhibition of Igf1r by NVP-ADW742,[35] an ATP-competitive inhibitor, blocked
systemic activation in limbs contralateral to amputation, indicating Igf-1 signaling is also necessary for this
process (**Figure S4 C-D**). Although Wnt signaling has been implicated in a variety of regenerative events
across many species,[36-40] we found it to be dispensable for systemic activation in axolotl as the Wnt
inhibitor IWR-1[41] did not alter the number of cells that become systemically activated in response to
amputation (**Figure S4**).
258
259 **Chromatin accessibility changes associated with systemic activation**
260
We hypothesized that cells might be altering their chromatin accessibility in response to
amputation injury as a means to shift into an activated state, primed for regeneration. To address this
possibility, we used our FACS strategy to isolate 4C cells from the homeostatic limb, systemically-
activated limb, and regenerating limb samples including the blastema and wound epidermis. We pooled
cells from each of these three conditions and performed ATAC-seq, and we identified 140 regulatory
regions across 119 genes (**Figure 6**). Principal component analysis of these three populations revealed
that the three conditions cluster separately with respect to PCA 1 and 2 (**Figure 6A**).
Most peaks with differential accessibility mapped to unannotated genes (**Figure 6B**). However,
we uncovered several differences in chromatin accessibility near or within annotated genes across the
three conditions. For example, we only recovered peaks from 4C cells isolated from systemically-
activated limbs, but not 4C cells from homeostatic or regenerating samples for *Foxo1* (**Figure 6C**). We
also found *Dyrk2* loci to be more accessible in the activated 4C samples compared to 4C homeostatic
and blastema samples (**Figure 6C**). DYRK2 is a direct downregulator of two key regulators of G1/S
transition, c-JUN and c-MYC, suggesting that it might be involved in maintaining the stem cells of the
activated limbs in a quiescent state.[42] These data suggest chromatin may be opened in the vicinity of
genes key for regulating the systemically-activated state, while it may be closed when cells proliferate
during homeostasis and while in the blastema cell state.
278
279
280
281 **DISCUSSION**
282
283   Our results highlight a previously-unappreciated role for body-wide activation of cell types that
284 cycle during organismal homeostasis and their repurposing for pro-regenerative responses during

vertebrate limb regeneration. Our data support a model in which the mechanism of systemic activation requires peripheral innervation at both the injury and distant responding sites. Our work identifies a more primary role for peripheral innervation in the initial stimulation of progenitor cells to re-enter the cell cycle. This role for PNS is complementary to the previously-appreciated roles for PNS as a source of mitogens in growing nascent blastemas to become larger[43-47] and as a positive regulator of limb size during regeneration.[48] We investigated the mechanisms whereby systemic activation occurs, and we discovered adrenergic signaling and mTOR pathways as critical mediators of both systemic activation and priming for future regeneration.

Our work provides molecular support for vertebrate limb regeneration being predicated on the activity of cells with shared use during homeostasis. Future studies will be required to understand the downstream pathways that these cells use to become poised to promote limb regeneration following an amputation. How cells sense their location in space and respond accordingly may depend on their proximity to other extrinsic cues, such as the wound epidermis that forms specifically at the injury site. We postulate that while the initial proliferative response to amputation is global in axolotls **(Figure 6E),** the sustained proliferation of activated cells and their migration out of tissue niches may be driven by wound-epidermis factors. This prediction aligns with the finding that suturing salamander limbs closed after amputation blocks formation of this molecularly-specialized, transient wound epidermis and, thus, outward blastema formation.[49] However, when sutures were removed after 5 weeks, the wound epidermis formed and limbs grew visible blastemas and fully regenerated—at a faster rate than a fresh amputation.[49] In essence, this experiment temporally decoupled the amputation injury from blastema formation, and suggesting that cells may have been poised by the injury and held in that state until receiving signals from wound epidermis. Our study provides molecular support for this model.

Several outstanding questions remain for understanding the implications of our findings. Additional experimentation will be required to address the relative contributions of sympathetic neurons locally versus the hypothalamic-pituitary-adrenal (HPA) axis as the source of the catecholamines important for regeneration. While we showed that the systemic activation and priming effects are transient, and that they disappear within several weeks, the mechanism whereby they are extinguished remains to be determined, and the determining the role of epigenetics in this process requires more experimentation. Our ATAC-seq analysis identified the promoter region of the *Dyrk2* loci to be more accessible in the 4C cells of the activated limb compared to homeostatic limbs, suggesting this kinase could be important to keep the activated stem cells in a quiescent state in the absence of injury-site specific signals to commit to a blastema cell fate. We also identified loci of interest whose homologs have relevance to stem cell populations in other species. For example, Tetraspanins have been implicated as markers of planarian neoblast subsets with enhanced differentiation potentials,[50] and FOXO1 is required to maintain pluripotency in human embryonic stem cells.[51] Importantly, the fork head transcription factors act downstream of insulin signaling to promote longevity and cell survival,[52,53] in line with our findings that suggest insulin/IGF1R signaling is required for systemic activation (**Figure 4S**). Addressing hypotheses about how the cells we have identified in axolotl might relate to planarian neoblasts and mammalian stem cells will require additional experimentation. Additionally, while systemic activation may have short-term benefits to the organism by acting to prime other tissues for regeneration, the consequences of this system over the individual lifespan need to be evaluated. Repeated injury and chronic stimulation of this system may come at a cost. While we can imagine scenarios whereby coupling systemic activation to limb regeneration might be evolutionarily advantageous, for example in high-density settings among cannibalistic species,[54] addressing these possibilities will require controlled experimentation.

Our work is likely to have important implications for defining where in the process of limb regeneration mammals have evolved and/or retained additional molecular controls that restrict natural regenerative ability. As mice have been shown to engage in a similar response following more limited local injuries,[25] the possibility that the first step of limb regeneration—in which cycling progenitor cells shift from a homeostatic function to a regenerative function—is operational and salient in mammals. Whether

an important role for the peripheral nervous system in orchestrating the body-wide activation response in mammals also exists, perhaps intertwined with the circulation, remains to be determined. Additional studies will also be required to reveal if either, or both, axolotls and mice reserve some subset of these homeostatic progenitor cell types exclusively or preferentially for regeneration. Elucidating how some of these activated progenitors enter the blastema cell state will be critical. Our work predicts that activating the relevant cell types and directing them toward a regenerative response could be stimulated by harnessing adrenergic and mTOR signaling in a mammalian setting, providing a major advancement toward the goal of therapeutic limb regeneration in the future. Our findings may also provide important reference points for medical applications outside of the amputee population, for example, in the context of "polytrauma," in which a patient suffers severe wounds in several different body parts.[55]

ACKNOWLEDGMENTS

This work was supported by The Richard and Susan Smith Family Odyssey Award (J.L.W.), the NIH New Innovator Award DP2HD087953 (J.L.W.), NICHD R01HD095494 (J.L.W.), NSF-CAREER award (J.L.W.), Harvard University Faculty of Arts and Sciences (J.L.W.), Harvard/MIT Joint Basic Program in Neuroscience (J.L.W. and I.M.C.), the Human Frontiers Science Program Long-term Postdoctoral Fellowship (A.M.S.), the Harvard Program for Research in Science and Engineering (S.H. and J.L.), the Harvard College Research Program (S.H. and A.Y.W.), and the Harvard Stem Cell Institute (A.W.). We thank Dr. Elaine Fuchs' lab for sharing reagents. We thank Samantha Payne, Solsa Cariba, Sydney Chambule, Jason Buenrostro, Fabiana Duarte, Juliana Babu, Jeramiah Smith, Nataliya Timoshevskaya, Tomás Gomes, and Adnan Abouelela for technical assistance. We thank William Ye, Tyler Mourey, and Quinley Miao for animal care. We thank Ya-Chieh Hsu, Matthew Warman, Marc Kirschner, Davie Van Vactor, Paola Arlotta, Rich Lee, Paul Garrity, and members of the Whited, Chiu, Greer and Lehoczky labs for discussions and comments on the manuscript. We thank Yulia Swartz for sharing reagents and assisting in data interpretation, and the Bauer Core Facility at Harvard University for cell sorting and sequencing.

AUTHOR CONTRIBUTIONS

D.P.-D, S.E.W., B.E., S.H., J.L., J.V.M, I.M.C., and J.L.W. designed the experiments; D.P.-D. S.E.W., and J.L.W. wrote the paper; S.J.B. , E.K., and B.J.H. designed and performed computational analyses; L.C. performed statistical analyses; D. P.-D., S.E.W., A.M.S., B.E., N.L., S.M., B.G., K.D., S.H., J.V.M., A.W., J.L., and H.S. performed sample preparation, experiments, and experimental data analyses. All authors contributed to manuscript editing.

DECLARATION OF INTERESTS

MAIN FIGURE TITLES AND LEGENDS

Figure 1. Systemically-activated cells are primed for future regeneration events.
(A) Experimental design. We compared regeneration rates of systemically-activated and naïve, control limbs. Green dots, activated, EdU positive nuclei. **(B)** Representative images of regenerating systemically-activated or control limbs. **(C-E)** Quantification of blastema area at 14 dpa **(C)**, differentiating

limbs **(D),** and number of digit outlines **(E)** at 25 dpa (n=15 animals for each condition in blastema area measurement, n = 30 limbs for each condition in differentiation and digit outline quantification). **(F)** *Prrx1* HCR-FISH was performed on tissue sections at 5 dpa following local amputation of naïve limbs (no previous amputation) and primed limbs (previously amputated on the contralateral side 14 days earlier). **(G)** Quantification of *Prrx1* mean fluorescence intensity across entire tissue sections (n = 8 animals for each group). **(H)** EdU and DAPI staining of intact limbs (left) and limbs contralateral to amputation (right) harvested 6 weeks after EdU administration. **(I)** Quantification of percent EdU positive nuclei of intact and contralateral limbs in **(H),** (n = 15 animals for intact and n = 12 animals for contralateral group). **(J)** Representative images of systemically-activated and naïve limbs amputated 28 days post-contralateral amputation. **(K-L)** Quantification of blastema area at 7 dpa **(K)** and number of digit outlines at 27 dpa **(L)** (n = 19 animals for control, n = 20 animals for activated for blastema area measurement, n = 38 limbs for control and n = 40 limbs for systemically-activated animals in digit outline quantification). **(M)** Keratin 5 (left), Keratin 10 (right) staining of limbs contralateral to amputation harvested 12 and 14 days post-amputation, respectively, 18 hours after EdU administration. All data shown as mean ± SD. Statistical significance was determined using unpaired two-tailed Welch's t-test in **(C)**, **(G)**, **(I)** and **(K)**; Fisher's exact test in **(D)**; and Fisher's exact test with Freeman-Halton extension in **(E)** and **(L)**. Scale bars, 2 mm in **(B)** and **(D)**; 500 µm in **(H)**; 200 µm in **(M).** dpa, days post-amputation. K5, keratin 5. K10, keratin 10. EdU, 5-ethynyl-2′- deoxyuridine.

Figure 2. Systemically-activated cells represent the same broad cell types as cells proliferating during homeostasis.   (A) Experimental design to identify systemically activated cells and compare to proliferating cells during homeostasis. Dividing (4C) cells of systemically-activated and naïve, control limbs were isolated by sorting dissociated cells according to their DNA content. **(B)** Representative FACS distribution following Hoecsht stain of homeostatic and systemically-activated limb cell samples. **(C)** UMAP plot of 4C cells collected from homeostatic control limbs (n = 4) and systemically-activated limbs (n = 4). **(D)** Heat map depicting statistically-significant differential gene expression between 4C cells in homeostatic versus activated samples. Only cell types with differential gene expression are shown. **(E)** Dot plot showing relative frequency of expression (dot size) and expression level (color) for differentially expressed genes across the eight fibroblast cell populations in homeostatic (H) versus activated (A) samples. **(F)** Cell fate reconstruction analysis of 4C cells in homeostatic and systemically activated limbs. **(G)** UMAP plot of 2C cells collected from homeostatic control limbs (n = 3) and systemically-activated limbs (n = 4). **(H)** Heat map depicting statistically-significant differential gene expression between 2C cells in homeostatic versus activated samples. Only cell types with differential gene expression are shown. See also **Figure S1.**

Figure 3. mTOR signaling is required for systemic activation and priming. (A) Chord diagrams of CellChat analysis of HGF signaling between cell types identified by scRNA-seq in Figure 2. **(B)** pS6 and EdU staining of blastema, intact (homeostatic) and contralateral limb tissue sections from animals treated with saline, vehicle, or rapamycin at 14 dpa. **(C)** Quantification EdU positive nuclei at 14 dpa (n = 12 limbs for intact saline, intact vehicle and intact rapamycin groups, n = 10 limbs for amputated saline and amputated vehicle, and n = 8 limbs for amputated rapamycin group). **(D)** Experimental design to test requirement for mTOR in priming.   Animals were treated with rapamycin or vehicle solution by daily injections starting from 3 days before to 7 days post unilateral amputations. The treatments were then ceased, and the uninjured contralateral limbs were unilaterally amputated. The regeneration rates of the second set of amputations were compared between treatment groups. **(E)** Representative images of regenerating contralateral limbs of rapamycin and vehicle solution-treated animals. **(F)** Quantification of blastema area at 7 days post-amputation. **(G)** Quantification of the number of digit outlines at 23 days post-amputation (n=20 animals for vehicle and rapamycin; n = 40 limbs for vehicle and rapamycin for each condition in digit outline quantification). The same group of vehicle control animals were used in Figure 5 (I-J). **(H)** pS6 and EdU staining of dorsal root ganglia servicing homeostatic, amputated, and contralateral limbs at 7 days post amputation (n = 20 for homeostatic and amputated, and n = 18 for contralateral). **(I-J)** Quantification of EdU positive **(I)** and pS6 positive **(J)** nuclei for DRGs in **(H)**. All data shown as mean ± SD. Statistical significance was determined using unpaired two-tailed Welch's t-test

with Bonferroni-Dunn correction in **(C)**; two-tailed Welch's ANOVA with Dunnett's T3 correction for multiple hypothesis testing in **(F)**, **(I)** and **(J)**; and Fisher's exact test with Freeman-Halton extension in **(G)**. Scale bars, 500 μm in **(B)**; 2 mm in **(E)**; and 100 μm in **(H)**. rapa, rapamycin. Int, intact. contra, contralateral. dpa, days post-amputation. pS6, phospho-S6 ribosomal protein; EdU, 5-ethynyl-2′-deoxyuridine. See also **Figure S2, Tables S1-S4.**

Figure 4. PNS at both amputation site and responding site are required for systemic activation and priming. (A) Experimental design. Nerve bundles innervating limbs contralateral to unilateral amputations were transected or sham-operated, and animals were assayed for systemic activation. Animals receiving unilateral denervations or sham-operations with no amputations were used as controls. **(B)** EdU and DAPI staining of intact and contralateral limb tissue sections of denervated and sham-operated animals in the intact state or at 14 days post amputation. **(C)** Quantification of percent EdU positive nuclei in denervated and sham-operated at 14 dpa. **(D-F)** Breakdown of tissue types represented in **(C)**. **(D)** Epidermis. **(E)** Skeletal elements. **(F)** Internal, non-skeletal tissues. (n = 11 animals for sham, n = 10 animals for denervated, n = 13 animals for sham amputated and denervated amputated, **(G)** Experimental design. Nerve bundles were transected or sham-operated in limbs that were immediately amputated thereafter. **(H)** EdU and DAPI staining of intact and contralateral limb tissue sections of denervated and sham-operated animals in the intact state or at 14 days post amputation. **(I)** Quantification of percent EdU positive nuclei in denervated and sham-operated at 14 dpa. (n = 18 animals for sham, n = 16 for sham amputated, n = 11 for denervated, and n = 12 for denervated amputated. **(J-K)** Differential gene expression analysis identifies transcripts that are enriched in nerve ends **(J)** and responding tissues **(K)**. All data shown as mean ± SD. Statistical significance was determined using unpaired two-tailed Welch's t-tests with Bonferroni-Dunn correction. Scale bars, 500 μm. den, denervated. int, intact. contra, contralateral. EdU, 5-ethynyl-2′-deoxyuridine. See also **Table S5**.

Figure 5. α-Adrenergic signaling is required for systemic activation and priming. (A) Plasma levels of norepinephrine at homeostasis and 4 hours post-amputation, as determined by ELISA. **(B)** Plasma levels of epinephrine at homeostasis and 4 hours post-amputation, as determined by ELISA. **(C)** EdU and DAPI staining of limbs contralateral to unilateral amputations treated with yohimbine or vehicle control, harvested at 7 days post amputation. **(D)** Quantification of percent EdU positive nuclei in contralateral limbs in **(C)** (n = 15 animals for each group, analyzed by unpaired two-tailed Welch's t-test). All data shown as mean ± SD. Scale bar, 500 μm. EdU, 5-ethynyl-2′-deoxyuridine. **(E)** Experimental design. Animals were treated with Adra2a inhibitor yohimbine or vehicle solution by daily injections starting from 2 days before to 7 days post unilateral amputations. The treatments were then ceased, and the uninjured contralateral limbs were unilaterally amputated. The regeneration rates of the second set of amputations were compared between treatment groups. **(F)** Representative images of regenerating contralateral limbs of yohimbine and vehicle solution treated animals. **(G)** Quantification of blastema area at 14 days post-amputation (n=15 animals for each condition). **(H)** Quantification of the number of digit outlines at 25 days post-amputation (n = 30 limbs for each condition). **(I)** Experimental design. Animals were treated with Adra2a agonist clonidine or vehicle solution by daily injections starting from 3 days before to 7 days post unilateral amputations. The treatments were then ceased, and the uninjured contralateral limbs were unilaterally amputated. The regeneration rates of the second set of amputations were compared between treatment groups. **(J)** Representative images. **(K)** Quantification of blastema area at 7 days post-amputation. (n = 20 animals for vehicle and n = 18 animals for clonidine). The same group of vehicle control animals were used in Figure 3 (E-G). **(L-N)** EdU and DAPI staining of limbs contralateral to unilateral amputations treated with propranolol or vehicle control, harvested at 7 days post-amputation. **(L)** Representative images. **(M)** Quantification of percent EdU positive nuclei within the blastema area of the initially-amputated limbs (n = 15 animals for vehicle and n = 10 for propranolol). **(N)** Quantification of percent EdU positive nuclei within contralateral limbs (n = 15 animals for vehicle and n = 10 for propranolol). Statistical significance was determined using paired two-tailed Welch's t-tests in **(A)** and **(B)**; unpaired two-tailed Welch's t-tests in **(D)**, **(G)**, **(M)** and **(N)**; Welch's ANOVA with Dunnett's T3 correction in **(K)**; Fisher's exact test with Freeman-Halton extension in **(H)**. Scale bars, 500 μm in **(C)** and **(L)**; 2 mm in **(F)** and **(J)**. hpa, hours post-amputation. dpa, days post-amputation. YOH, yohimbine. See also **Figure S3**.

Figure 6. (A) Principal component analysis of chromatin accessibility of 4C cells in homeostatic, activated, and regenerating limbs. **(B)** Heatmap showing relative abundances accessible regions in 4C cells of homeostatic, activated, and regenerating limbs. **(C-D)** Genome browser tracks showing accessible regions of the Foxo1 **(C)** and Dyrk2 **(D)** loci. **(E)** Two-step model for blastema cell specification. In step 1, some cells (magenta nuclei) from a field of possible progenitor cells re-enter the cell cycle (activation) and begin to proliferate in response to amputation. We postulate this step is systemic, occurring throughout the body. In step 2, some subset of activated progenitor cells later converts to the blastema cell state (blue cells). We postulate this step is local to the site of amputation. The subsequent migration of blastema cells to the distal-most tip of the stump creates a visible blastema (yellow).

STAR Methods

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Jessica Whited (jessica_whited@harvard.edu).

Materials availability
This study did not generate new unique reagents.

Data and code availability
- All bulk RNA-sequencing data sets generated during this study are available at Gene Expression Omnibus (GEO), accession number GSE192477.
- All single-cell RNA-sequencing datasets generated in this study are available at Gene Expression Omnibus (GEO), accession numbers GSE232083 (4C) and GSE232082 (2C).
- All bulk ATAC-sequencing datasets generated in this study are available at Gene Expression Omnibus (GEO), accession number GSE232080.
- All original code has been deposited at Zenodo and is publicly available as of the date of publication. The DOI is listed in the key resources table.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Animals
All animal experimentation was approved by and conducted in accordance with Harvard University's Institutional Animal Care and Use Committee. Leucistic axolotls were used for all animal experimentation except RNA sequencing and maintained as previously described.[5] For RNA sequencing, *βIIItub:GAP43-EGFP* transgenic animals were used.[56] Animals in all experiments were both age and size matched. Animals were anesthetized in 0.1% (w/v) tricaine prior to all procedures involving amputation, denervation, or injection. After all surgical procedures, axolotls were allowed to recover overnight in 0.5% (w/v) sulfamerazine. All limb amputations were performed at the mid-stylopod and the bone was then trimmed back from the amputation plane. For forelimb denervations the brachial plexus nerves were transected as described previously,[57] and for hindlimb denervations the sciatic nerves were transected as described[58]. Sham operations as controls for denervations were performed by anesthetizing animals and

541 cutting the skin open with sterile scissors, locating the nerve bundles with closed sterile forceps, but not
542 cutting the nerves.
543
544 **METHOD DETAILS**
545
Denervations and amputations
Animals in all experiments were both age and size matched. Animals were anesthetized in 0.1% (w/v)
tricaine prior to all procedures involving amputation, denervation, or injection. After all surgical
procedures, axolotls were allowed to recover overnight in 0.5% (w/v) sulfamerazine. All limb amputations
were performed at the mid-stylopod and the bone was then trimmed back from the amputation plane. For
forelimb denervations the brachial plexus nerves were transected according to Schotté et al. [57] and for
hindlimb denervations the sciatic nerves were transected according to Kropf et al. [58] Sham operations as
controls for denervations were performed by anesthetizing animals and cutting the skin open with sterile
scissors, locating the nerve bundles with closed sterile forceps, but not cutting the nerves.
555
Drug treatments and EdU incorporation
Rapamycin was dissolved into a 25 mg/ml stock solution in ethanol. Rapamycin treatments were
delivered via both water treatments at a concentration of 1 µM for 16 days with water changed daily, and
through daily intraperitoneal injections at a concentration of 5 mg/kg/day for 16 days, both starting two
days before amputation. All intraperitoneal injections were administered at a volume of 20µL/g in a
vehicle solution comprised of APBS with 5% Tween80, 5% PEG-400. All stock solutions were frozen at
−80 °C in 1-mL aliquots. Yohimbine (Y3125, Sigma-Aldrich) was dissolved into a 10 mg/mL stock solution
with sterilized water. Yohimbine was administered daily at a dose of 1mg/kg for 14 days via
intraperitoneal injection. A stock solution of IWR-1 (I0161, Sigma-Aldrich) was created by dissolving
5mg/mL in DMSO. IWR-1 was delivered daily at a dose of 10mg/kg/day for 9 days via intraperitoneal
injection. A stock solution of NVP-ADW742 (5247, Tocris) was created by dissolving 2mg/mL in DMSO.
NVP-ADW742 was injected daily at a dose of 10mg/kg/day for 14 days. Clonidine (C7897, Sigma-Aldrich)
was dissolved in water at 7.5 mM stock concentration and was delivered via water treatments at 30 µM. A
stock solution of propranolol (P0884, Sigma-Aldrich) was created by dissolving 100 mM in water.
Propranolol was delivered via water treatments at 5 µM. 5-ethynyl-2′-deoxyuridine (EdU) was purchased
from ThermoFisher and stock solutions were created by dissolving the powder in dimethyl sulfoxide as
instructed by the manufacturer. Following anesthetization, axolotls were administered 20 µL/g of 400 µM
EdU in 0.7X PBS via intraperitoneal injection 18 hours before tissue harvest.
574
Skeletal and tissue preparations
For tissue sections, post-collection, tissue samples were fixed in 4% paraformaldehyde in DEPC-treated
PBS overnight at 4°C. Tissues were then cryopreserved in a sucrose gradient following a wash in DEPC-
PBS and incubated in 30% sucrose/DEPC-PBS at 4°C overnight. Tissues were then embedded and
frozen on dry ice in Tissue-Tek O.C.T. Compound (Sakura) and stored at −80°C. All blocks were
sectioned at 16µm thickness using a Leica CM 1950 cryotome and sections were stored at −80°C until
use.
582
583 For skeletal preparations, post-collection, limbs were incubated overnight in 95% ethanol at room
584 temperature and then incubated in 100% acetone under the same conditions. Skeletal elements were
585 visualized with alcian blue and alizarin red S stains followed by a clearing of non-skeletal tissue with
586 potassium hydroxide.
587
588
589

Immunofluorescence and HCR-FISH

Sections were rehydrated with PBS and then permeabilized with 0.5% Triton-X-100 PBS. Post-permeabilization, slides were rinsed twice with PBS and EdU visualization was performed via sulfo-cyanine 3 azide (Lumiprobe). Post-EdU visualization slides were washed twice with PBS. Staining and labeling with any additional antibodies was conducted post-EdU labeling. Primary antibodies used were rabbit anti-phospho-S6 ribosomal protein (Ser235/236, 1:200; Cell Signaling 4858), and rabbit anti-keratin 10 (BioLegend 905401). Secondary antibodies used were Cy3-goat anti-mouse (1:250, Jackson ImmunoResearch 115-165-146), Alexa Fluor 488-Goat Anti-Rabbit (1:250, Jackson ImmunoResearch, 111-545-003) and Cy3-goat anti-rabbit (1:250, Jackson ImmunoResearch 111-165-003).

Sections stained with all antibodies were blocked with 10% serum from the host species of the secondary antibody in 0.1% Triton-X-100 in PBS. Primary and secondary antibodies were diluted in in 2% BSA, 0.1% Triton-X-100 in PBS. Sections labeled with anti-pS6 and NeuN underwent an antigen retrieval step after permeabilization, in which slides were boiled in 0.1 M sodium citrate (pH 6.0) and then rinsed once in PBS, before EdU was visualized, followed by blocking and primary antibody incubation. All slides were incubated with DAPI (Sigma D9542) before cover slipping. HCR-FISH was performed as described previously.[59]

Plasma isolation and ELISA

Blood was collected from axolotls as described previously.[60] Efferent blood vessels were exposed by placing the animals on its back. Winged infusion sets were used to collect the blood. Whole blood was placed into EDTA coated tubes and spun down. The supernatant fraction (plasma) was used for downstream analysis. Animals were allowed to recover overnight in fresh axolotl water. Epinephrine and norepinephrine ELISAs were performed using Abnova ELISA kit KA1877.

Imaging and quantification

Imaging was performed on a Zeiss Axio Scan.Z1 automated slide scanner, Leica M165 FC equipped with Leica DFC310 FX camera, and a Nikon Eclipse Ni microscope with a DS-Ri2 camera. Quantification of all images was conducted blindly. For imaging of intact contralateral limbs, representative sections displaying all tissue types (bone and cartilage, soft internal tissues, and epidermis) were selected. For the tissue type analysis of EdU+ nuclei, tissues were classified as epidermis, "skeletal elements" (bone, cartilage), and "soft tissues" (muscle, joint, tendon, ligaments, dermis, vasculature, and nerves) based on nuclear morphology and location as previously described (Whited et al, 2013). All cell counts were conducted utilizing CellProfiler.[61] Measurements of blastema area were performed using Fiji.[62] For *Prrx1* HCR signal quantifications, the wound epidermis was first masked from each image. *Prrx1* fluorescence intensity was normalized to the area of the blastema as measured by the DAPI signal.

Cell dissociation, FACS isolation, and single-cell RNA-sequencing

To isolate single cells, 12-16 homeostatic or systemically-activated limbs at 14 days post-amputation were pooled for each biological replicate and dissociated in 4 ml of dissociation buffer (100 mg/ml liberase diluted in 0.7X PBS with 0.5 U/ml DNase I) for 30 min at room temperature with gentle mixing. Suspensions were filtered through 70 mm strainers to remove cell clumps and debris, and strainers were washed with 1 ml of 0.7X PBS. Cells were spun down at 300 g at room temperature and resuspended in 1 ml of 0.7X PBS-0.04% BSA. Cells were then stained with 1 mg/ml Hoecsht, calcein AM, and propidium iodide for isolation of alive, dividing (4C) and non-dividing (2C) cells by fluorescence activated cell sorting. The purified cells were then sequenced at the Bauer Core Facility at Harvard University using an Illumina NovaSeq 6000 [https://www.illumina.com]. Paired end reads were demultiplexed with bcl2fastq2 (V2.2.1) (RRRID:SCR_015058).

ATAC-seq Library Prep and Sequencing

Ten thousand sorted 4C cells from homeostatic, systemically-activated, and regenerating limbs were resuspended in 5 µL PBS. 42.5 µL of transposition buffer (38.8mM tris-acetate, 77.6mM potassium acetate, 11.8mM magnesium acetate, 0.12% NP-40, 18.8% DMF, 0.47% protease inhibitor cocktail) was added to the cells, mixed, and incubated 10 min at room temperature. 2.5 µL of pre-loaded Tn5 (Diagenode cat# C01070012) was added to the reaction mixture. The transposition reaction was carried out at 37°C for 30 min with shaking at 300 rpm. Following transposition, the samples were purified through Direct-zol RNA MiniPrep kit (Zymo Research cat# R2050). Purified library fragments were amplified with custom primers 1 and 2 using the following conditions: 72°C for 5min; 98°C for 30s; and thermocycling at 98°C for 10s, 72°C for 30s and 72°C for 1min. The libraries were amplified for five cycles. Soon after, 5µL aliquot of the PCR reaction was added with 10µL of the PCR cocktail with SYBR Green (0.6Xfinal concentration). 20 cycles of PCR was performed to determine the additional number of cycles needed for the remaining 45 µL reaction. The libraries were purified using Direct-zol RNA MiniPrep kit. Purified libraries were sequenced at The Bauer Core Facility at Harvard University using an Illumina NovaSeq 6000 [https://www.illumina.com]. Paired End reads were demultiplexed with bcl2fastq2 (V2.2.1) (RRID: SCR_015058).

Bulk RNA isolation, sequencing, and computational analyses

βIIItub:GAP43-EGFP transgenic animals were used for RNA sequencing experiments to allow for easily identifiable collections of tissues.[56] RNA was extracted from tissue samples with Trizol (15596018, Ambion) and chloroform (288306, Sigma Aldrich). RNA was then isolated from tissue samples using a Zymo Clean Up and Concentrator kit (R1013). RNA samples were processed at the Bauer Core Facility at Harvard University using the Illumina NextSeq 500. Paired End reads were demultiplexed with bcl2fastq2 v2.2.

QUANTIFICATION AND STATISTICAL ANALYSIS

Minigenome

A custom reference "minigenome" was created by first performing in silico normalization on the sequenced reads by sample with Trinity (V2.12.0)[63] with the following parameters: --CPU 10 --max_memory 50G --no_run_inchworm --min_kmer_cov 2 --trimmomatic --seqType fq, before assembling them with parameters: --CPU 10 --max_memory 50G --min_kmer_cov 3 --seqType fa --no_normalize_reads. This transcriptome was then aligned to the entire axolotl genome (V6.0.0) [https://www.axolotl-omics.org/assemblies][64] using minimap2 (V2.24)[65] and the parameters -ax splice --secondary=no -O6,24 -B4 -L -t 20 -cs -ub -G 100000 to create genome intervals of the genes present in the transcriptome. The intervals were converted from the original SAMtools (V1.15.1)[66] format to BED using BEDOPS (V2.4.25).[67] Finally, the minigenome and annotation was created from the intervals using ShrinkGenome [https://github.com/brianjohnhaas/ShrinkGenome]. The complete axolotl mitochondrial genome [https://www.ncbi.nlm.nih.gov/nuccore/NC_005797][68] was then appended to this minigenome. We prevented competitive alignments to mitochondrial contributors downstream by masking their genomic loci using the maskFastaFromBed function from BEDTools (V2.30.0).[69]

Single-cell RNA-sequencing read quantification

We used Kallisto (V0.48.0)[70] to create an index, as well as pseudoalign the RNA-seq data before creating BUS [71] files by biological replicante. The BUS files were sorted and converted into text using bustools (V0.41.0)[72] using default parameters before converting the BUS files into sparse matrices of gene by cell data in R using BUSpaRse (V1.9.0).[73] The gene by cell sparse matrices were converted into Seurat[74] objects in R, filtering out cells with less than 200 transcripts as well as any feature that was in less than 3

cells, before merging the sample replicates into either homeostatic samples or systemically activated
samples, which were merged into a single data structure, while maintaining their original metadata. For
QC, cells were scored for their complexity with the following formula cell complexity = $\log_{10}$(# of genes in
cell) / $\log_{10}$(# of UMI in cell). If a cell had less than 1,000 UMI while also both expressing less than 350
features and a cell complexity score less than 0.75 it was not considered a candidate for analysis and
removed from the dataset.
695
Cell clustering and visualization
Clustering the cells was performed with Seurat (V4.1.0). Default parameters were used in the
FindVariableFeatures function of Seurat to determine the number of principal components. We used a
resolution of 1.0 to cluster the cells with 17 dimensions. We generated markers for the clusters and found
that one of the clusters consisted primarily of mitochondrial markers. This cluster was eliminated and the
cell clustering pipeline was repeated. We performed homotypic doublet proportion estimation to predict
and remove doublets using DoubletFinder (V2.0.3)[74] as described at [https://github.com/chris-mcginnis-
ucsf/DoubletFinder]. The cell clusters were visualized with the reduction method UMAP. Clusters were
identified by unique markers in each cluster.
705
Differential gene expression analysis for single-cell RNA sequencing
707 We searched for differentially expressed genes between the homeostatic and systemically activated cells
708 using DESeq2 (V1.36.0).[75] The contrast in the results was configured by condition, where homeostatic
709 limb was the baseline and systemically activated limb was the contrasting condition. The results were
710 determined to be significant if the adjusted p value was less than 0.1. Statistically significant results were
711 visualized with ggplot2 (V3.3.6).[76]
712
CellChat analysis
To obtain the information related to incoming and outgoing clusters ligand-receptors interactions, we used
the CellChat package[77] for 4C datasets using the previously generated Seurat objects. A custom axolotl
ligand-receptor reference was generated using the minigenome. The database included information on
ligands and receptors, and for orthologous gene repeats that have different positions in the genome and
different annotations, labeled as "Gene_ik", where k is the number of the repeats. The database was
reconstructed following the ligand-receptor pairs used for the mouse CellChat database with the gene
repeats included as separated ligand-receptor pairs (L1-R1, L1_i1_R1, etc.).
After generating the database, we followed the standard CellChat pipeline
[https://github.com/sqjin/CellChat/]. To avoid the projection reducing the dropout effects of signaling
genes, for possible zero expression of subunits of ligands/receptors, we used raw.use = FALSE setting in
computeCommunProb function. We also used population.size = TRUE parameter to consider the effect of
cell proportion in each cell group in the probability calculation. The chord diagrams were generated using
netVisual_aggregate() function.
727
RNA Velocity
Upon the clusters to states annotation, we utilized Cell ranger, samtools, velocyto, and scvelo packages
to perform the RNA Velocity analysis[78]. For the previously generated Cellranger output folder, we used
the samtools sort function in the samtools package (-t CB -O BAM) to create the
cellsorted_possorted_genome_bam.bam file from the possorted_genome_bam.bam file. The generated
file was used for making the .loom file upon utilizing the velocyto run10x function of the velocyto package,
using the minigenome. We tgeb converted the .h5Seurat annotated objects to .h5ad objects to transfer
them from R to the Python environment. We used the scvelo package[79] for the RNA Velocity analysis. We
merged .loom files with the respective data files previously generated, processed, and annotated. The
following files were filtered and normalized. The first and second order moments (means and uncentered

variances) were computed among nearest neighbors in PCA space with the scvelo.pp.moments()
function. Then we estimated velocity (scvelo.tl.velocity()), built the velocity_graph
(scvelo.tl.velocity_graph()) and projected the velocities using scvelo.pl.velocity_embedding_stream() and
scvelo.pl.velocity_embedding_grid() on UMAP (basis = 'umap') embedding. With that, we quantified
pseudotime, latent time, and built the latent-time-based heatmaps with gene RNA abundance analysis.
743
Bulk RNA-seq dataset processing
745 The reads were trimmed for quality and adapters with Trimmomatic v0.39[80] with the following parameters:
746 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36. An index of the previously published
747 axolotl transcriptome[5] was created using Kallisto v0.45.1[70]. Sequence based bias correction was
748 performed using Kallisto with 200 bootstraps across 20 threads. The transcript-level abundances from
749 Kallisto were summarized and imported into R as a matrix of counts using tximport v1.0.3[81].
750
751 Differentially expressed genes were determined by estimating fold change and dispersion using
752 DESeq2[75]. Genes were considered significant if they had a logarithmic fold change greater than 1.5 and
753 adjusted p value of 5%. These significant genes were visualized using pheatmap v1.0.12.[82] Pathway
754 analysis was performed in R using pathfinder.[83]
755
Chunked Genome
A custom "chunked genome" was needed for compatibility with tools used to process the alignments. We
broke the large chromosomes of the axolotl genome (V6.0.0) [https://www.axolotl-omics.org/assemblies][68]
into chunks smaller than 500M using ChunkGenome [https://github.com/brianjohnhaas/ChunkGenome].
ChunkGenome was also used to create a new feature coordinates from the original GTF. The complete
axolotl mitochondrial genome [https://www.ncbi.nlm.nih.gov/nuccore/NC_005797][68] was appended to this
chunked genome. We prevented competitive alignments to mitochondrial contributors downstream by
masking their genomic loci using the maskFastaFromBed function from BEDTools (V2.30.0)
[https://github.com/arq5x/bedtools2].[69]
765
Read processing for ATAC-seq
The sequencing data was aligned to the reference using Bowtie2 (V2.3.2)[84] using the parameters '--no-
mixed -X 2000 --dovetail --no-discordant'. Samtools (V1.10)[85] was used to keep the alignments sorted
and indexed throughout the processing. We removed all alignments to mitochondria by simply using the
grep -v chrM before filtering out the low quality alignments using Samtools view -q 30. The Picard Toolkit
[https://broadinstitute.github.io/picard/] was used to remove PCR and optical duplicates using Picard's
MarkDuplicates with the parameters 'ASSUME_SORTED=true REMOVE_DUPLICATES=true
SORTING_COLLECTION_SIZE_RATIO=0.2'. Coverage tracks of the alignments were generated with
deepTools[86] bamCoverage with parameters '--outFileFormat=bigwig --normalizeUsing CPM -p 16 --
binSize 20'.
776
ATAC-seq Peak Calling
Sites of genomic enrichment were found using Genrich (V0.5) (https://github.com/jsh58/Genrich#atacseq)
in ATAC-seq mode using default arguments '-j -q 0.05', the results of which were converted to bigwig
format using bedGraphToBigWig[87]. We evaluated the consistency of the Genrich generated peaks using
Irreproducible Discovery Rate (IDR) (V2.0.4.2)[88]. We filtered these conservative peaks even further as
described in the guidelines and practices of the ENCODE consortia[88]. We counted these peaks with high
rank consistency by first extending the coordinates of the peaks with bedtools (V2.30.0)[69] function slop
with argument '-b 500' before merging all the pooled replicates with bedtools merge, then finally using the
intersect function with parameters '-wo'.

Annotating and visualizing the ATAC data

We used ChIPseeker (V1.32.1)[89] to annotate the peaks. The counts were put together into a genes by peak count matrix and differential expression was observed using DESeq2[75]. Pheatmap (V1.9.12)[82] was used to visualize the normalized counts. We clustered the samples based on their similarity using pcaExplorer (V2.22.0)[90]. Genome browser tracks of all the peaks were plotted against the total coverage of all of the reads using pyGenomeTracks (V3.8).[91]

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Marker genes used to identify clusters in single-cell RNA sequencing of 4C and 2C cells (Related to Figure 2)
(A) Dot plots showing the relative frequency of expression (dot size) and expression level (color) for marker genes across all clusters of 4C cells. **(B)** Latent time-ordered heatmaps demonstrating variable genes across 4C fibroblast clusters. **(C)** Gene dynamics analysis shows the top 15 genes across the 4C cells with the most unique RNA abundance portraits. X-axis shows the spliced and the y-axis shows the unspliced transcripts for the gene, the steady state marked with the line. Each dot represents a single cell, with colors describing the clusters indicated on the right. **(D)** Dot plots showing the relative frequency of expression (dot size) and expression level (color) for marker genes across all clusters of 4C cells.

Figure S2. Inhibition of mTOR signaling with rapamycin delays regeneration (Related to Figure 3)
(A) Regenerating limbs at 7 dpa treated with rapamycin or vehicle control. Arrowheads mark the amputation plane. **(B)** Blastema growth curves of rapamycin or vehicle treated animals (n = 10 animals per group). Statistical significance was determined using unpaired two-tailed Welch's t-test with Holm-Šídák correction). **(C)** Representative skeletal preparations of regenerated limbs with Alcian blue and Alizarin red at 10 weeks post-amputation. Scale bars, 0.5 mm.

Figure S3. Adrenergic receptors expression in proliferating fibroblasts (Related to Figure 5) (Related to Figure 4)
Dot plot showing the relative frequency of expression (dot size) and expression level (color) for adrenergic receptors across all clusters of 4C cells.

Figure S4. Effects of inhibiting Wnt and Igf1r signaling on systemic activation (Related to Figure 4)
(A-B) Chord diagrams of CellChat analysis of Igf1r signaling **(A)** and insulin signaling **(B)** between 4C cell types identified by scRNA-seq in Figure 2. **(C)** EdU and DAPI staining of limbs contralateral to unilateral amputations treated with tankyrase inhibitor IWR-1, Igf1r inhibitor NVP-ADW742 or vehicle control, harvested at 7 days post amputation. **(D)** Quantification of percent EdU positive nuclei of contralateral limbs in (A) (n = 11 animals for vehicle control, n = 14 animals for IWR-1 and NVP-ADW742). All data shown as mean ± SD. Statistical significance was determined using Welch's ANOVA with Dunnett's T3 correction, W = 4.883. Scale bar, 500 µm. NVP, NVP-ADW742. EdU, 5-ethynyl-2′-deoxyuridine.

SUPPLEMENTAL EXCEL TABLE LEGENDS

Table S1. List of transcripts expressed by each cluster of 4C cells in homeostatic and systemically-activated limbs (Related to Figure 2).

838   **Table S2.** Differentially expressed transcripts expressed by each cluster of 4C cells in homeostatic and
839   systemically-activated limbs (Related to Figure 2).
840
841   **Table S3.** List of transcripts expressed by each cluster of 2C cells in homeostatic and systemically-
842   activated limbs (Related to Figure 2).
843
844   **Table S4.** Differentially expressed transcripts expressed by each cluster of 2C cells in homeostatic and
845   systemically-activated limbs (Related to Figure 2).
846
847   **Table S5.** List of differentially expressed transcripts between sham-operated and denervated forelimbs
848   contralateral to unilateral amputations (Related to Figure 4).
849
850   **Table S6.** List of abundant peaks identified across 4C cells of homeostatic, activated, and regenerating
851   limbs and differentially abundant peaks across pairwise comparisons. (Related to Figure 6).
852
853
854

References

1.   Spallanzani, L. (1769 [Italian: 1768].). Reproductions of the legs in the aquatic salamander. In An
Essay on Animal Reproductions [Prodromo di un opera da imprimersi sopra la riproduzioni
anamali], (Becket and de Hondt), pp. 68-72.
2.   Simon, A., and Tanaka, E.M. (2013). Limb regeneration. Wiley Interdiscip Rev Dev Biol *2*, 291-
300. 10.1002/wdev.73.
3.   Voss, S.R., Palumbo, A., Nagarajan, R., Gardiner, D.M., Muneoka, K., Stromberg, A.J., and
Athippozhy, A.T. (2015). Gene expression during the first 28 days of axolotl limb regeneration I:
Experimental design and global analysis of gene expression. Regeneration (Oxford, England) *2*,
120-136. 10.1002/reg2.37.
4.   Stewart, R., Rascon, C.A., Tian, S., Nie, J., Barry, C., Chu, L.F., Ardalani, H., Wagner, R.J.,
Probasco, M.D., Bolin, J.M., et al. (2013). Comparative RNA-seq analysis in the unsequenced
axolotl: the oncogene burst highlights early gene expression in the blastema. PLoS computational
biology *9*, e1002936. 10.1371/journal.pcbi.1002936.
5.   Bryant, D.M., Johnson, K., DiTommaso, T., Tickle, T., Couger, M.B., Payzin-Dogru, D., Lee, T.J.,
Leigh, N.D., Kuo, T.H., Davis, F.G., et al. (2017). A Tissue-Mapped Axolotl De Novo
Transcriptome Enables Identification of Limb Regeneration Factors. Cell reports *18*, 762-776.
10.1016/j.celrep.2016.12.063.
6.   Gerber, T., Murawala, P., Knapp, D., Masselink, W., Schuez, M., Hermann, S., Gac-Santel, M.,
Nowoshilow, S., Kageyama, J., Khattak, S., et al. (2018). Single-cell analysis uncovers
convergence of cell identities during axolotl limb regeneration. Science *362*.
10.1126/science.aaq0681.
7.   Leigh, N.D., Dunlap, G.S., Johnson, K., Mariano, R., Oshiro, R., Wong, A.Y., Bryant, D.M., Miller,
B.M., Ratner, A., Chen, A., et al. (2018). Transcriptomic landscape of the blastema niche in
regenerating adult axolotl limbs at single-cell resolution. Nat Commun *9*, 5153. 10.1038/s41467-
018-07604-0.
8.   Ricci, L., and Srivastava, M. (2018). Wound-induced cell proliferation during animal regeneration.
Wiley Interdiscip Rev Dev Biol *7*, e321. 10.1002/wdev.321.
9.   Wenemoser, D., and Reddien, P.W. (2010). Planarian regeneration involves distinct stem cell
responses to wounds and tissue absence. Developmental biology *344*, 979-991.
10.1016/j.ydbio.2010.06.017.
10.   Newmark, P.A., and Sanchez Alvarado, A. (2000). Bromodeoxyuridine specifically labels the
regenerative stem cells of planarians. Dev Biol *220*, 142-153. 10.1006/dbio.2000.9645.
11.   Salo, E., and Baguna, J. (1984). Regeneration and pattern formation in planarians. I. The pattern
of mitosis in anterior and posterior regeneration in Dugesia (G) tigrina, and a new proposal for
blastema formation. J Embryol Exp Morphol *83*, 63-80.
12.   Johnson, K., Bateman, J., DiTommaso, T., Wong, A.Y., and Whited, J.L. (2018). Systemic cell
cycle activation is induced following complex tissue injury in axolotl. Developmental biology *433*,
461-472. 10.1016/j.ydbio.2017.07.010.
13.   Satoh, A., Gardiner, D.M., Bryant, S.V., and Endo, T. (2007). Nerve-induced ectopic limb
blastemas in the Axolotl are equivalent to amputation-induced blastemas. Dev Biol *312*, 231-244.
10.1016/j.ydbio.2007.09.021.
14.   Dumont, N.A., Bentzinger, C.F., Sincennes, M.C., and Rudnicki, M.A. (2015). Satellite Cells and
Skeletal Muscle Regeneration. Compr Physiol *5*, 1027-1059. 10.1002/cphy.c140068.
15.   Muneoka, K., Holler-Dinsmore, G.V., and Bryant, S.V. (1985). A quantitative analysis of
regeneration from chimaeric limb stumps in the axolotl. J Embryol Exp Morphol *90*, 1-12.
16.   Muneoka, K., Fox, W.F., and Bryant, S.V. (1986). Cellular contribution from dermis and cartilage
to the regenerating limb blastema in axolotls. Dev Biol *116*, 256-260.
17.   Kragl, M., Knapp, D., Nacu, E., Khattak, S., Maden, M., Epperlein, H.H., and Tanaka, E.M.
(2009). Cells keep a memory of their tissue origin during axolotl limb regeneration. Nature *460*,
60-65. nature08152 [pii]
10.1038/nature08152.
18.   McCusker, C.D., Diaz-Castillo, C., Sosnik, J., A, Q.P., and Gardiner, D.M. (2016). Cartilage and
bone cells do not participate in skeletal regeneration in Ambystoma mexicanum limbs.
Developmental biology *416*, 26-33. 10.1016/j.ydbio.2016.05.032.

19. Carver, E.A., Jiang, R., Lan, Y., Oram, K.F., and Gridley, T. (2001). The mouse snail gene encodes a key regulator of the epithelial-mesenchymal transition. Mol Cell Biol *21*, 8184-8188. 10.1128/MCB.21.23.8184-8188.2001.

20. Barbera, M.J., Puig, I., Dominguez, D., Julien-Grille, S., Guaita-Esteruelas, S., Peiro, S., Baulida, J., Franci, C., Dedhar, S., Larue, L., and Garcia de Herreros, A. (2004). Regulation of Snail transcription during epithelial to mesenchymal transition of tumor cells. Oncogene *23*, 7345-7354. 10.1038/sj.onc.1207990.

21. Lamouille, S., Xu, J., and Derynck, R. (2014). Molecular mechanisms of epithelial-mesenchymal transition. Nat Rev Mol Cell Biol *15*, 178-196. 10.1038/nrm3758.

22. Buckbinder, L., Talbott, R., Seizinger, B.R., and Kley, N. (1994). Gene regulation by temperature-sensitive p53 mutants: identification of p53 response genes. Proc Natl Acad Sci U S A *91*, 10640-10644. 10.1073/pnas.91.22.10640.

23. Haidurov, A., and Budanov, A.V. (2020). Sestrin family - the stem controlling healthy ageing. Mech Ageing Dev *192*, 111379. 10.1016/j.mad.2020.111379.

24. La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., Lidschreiber, K., Kastriti, M.E., Lonnerberg, P., Furlan, A., et al. (2018). RNA velocity of single cells. Nature *560*, 494-498. 10.1038/s41586-018-0414-6.

25. Rodgers, J.T., King, K.Y., Brett, J.O., Cromie, M.J., Charville, G.W., Maguire, K.K., Brunson, C., Mastey, N., Liu, L., Tsai, C.R., et al. (2014). mTORC1 controls the adaptive transition of quiescent stem cells from G0 to G(Alert). Nature *510*, 393-396. 10.1038/nature13255.

26. Rodgers, J.T., Schroeder, M.D., Ma, C., and Rando, T.A. (2017). HGFA Is an Injury-Regulated Systemic Factor that Induces the Transition of Stem Cells into GAlert. Cell reports *19*, 479-486. 10.1016/j.celrep.2017.03.066.

27. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.H., Myung, P., Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using CellChat. Nat Commun *12*, 1088. 10.1038/s41467-021-21246-9.

28. Takeuchi, K., Shibamoto, S., Nagamine, K., Shigemori, I., Omura, S., Kitamura, N., and Ito, F. (2001). Signaling pathways leading to transcription and translation cooperatively regulate the transient increase in expression of c-Fos protein. J Biol Chem *276*, 26077-26083. 10.1074/jbc.M102704200.

29. Brown, E.J., Albers, M.W., Shin, T.B., Ichikawa, K., Keith, C.T., Lane, W.S., and Schreiber, S.L. (1994). A mammalian protein targeted by G1-arresting rapamycin-receptor complex. Nature *369*, 756-758. 10.1038/369756a0.

30. Renthal, W., Tochitsky, I., Yang, L., Cheng, Y.C., Li, E., Kawaguchi, R., Geschwind, D.H., and Woolf, C.J. (2020). Transcriptional Reprogramming of Distinct Peripheral Sensory Neuron Subtypes after Axonal Injury. Neuron *108*, 128-144 e129. 10.1016/j.neuron.2020.07.026.

31. Rao, Y. (2019). The First Hormone: Adrenaline. Trends Endocrinol Metab *30*, 331-334. 10.1016/j.tem.2019.03.005.

32. Jabir, N.R., Firoz, C.K., Zughaibi, T.A., Alsaadi, M.A., Abuzenadah, A.M., Al-Asmari, A.I., Alsaieedi, A., Ahmed, B.A., Ramu, A.K., and Tabrez, S. (2022). A literature perspective on the pharmacological applications of yohimbine. Ann Med *54*, 2861-2875. 10.1080/07853890.2022.2131330.

33. Tam, S.W., Worcel, M., and Wyllie, M. (2001). Yohimbine: a clinical review. Pharmacol Ther *91*, 215-243. 10.1016/s0163-7258(01)00156-5.

34. Black, J.W., Crowther, A.F., Shanks, R.G., Smith, L.H., and Dornhorst, A.C. (1964). A New Adrenergic Betareceptor Antagonist. Lancet *1*, 1080-1081. 10.1016/s0140-6736(64)91275-9.

35. Mitsiades, C.S., Mitsiades, N.S., McMullan, C.J., Poulaki, V., Shringarpure, R., Akiyama, M., Hideshima, T., Chauhan, D., Joseph, M., Libermann, T.A., et al. (2004). Inhibition of the insulin-like growth factor receptor-1 tyrosine kinase activity as a therapeutic strategy for multiple myeloma, other hematologic malignancies, and solid tumors. Cancer Cell *5*, 221-230. 10.1016/s1535-6108(04)00050-9.

36. Kawakami, Y., Rodriguez Esteban, C., Raya, M., Kawakami, H., Marti, M., Dubova, I., and Izpisua Belmonte, J.C. (2006). Wnt/beta-catenin signaling regulates vertebrate limb regeneration. Genes & development *20*, 3232-3237. 10.1101/gad.1475106.

37. Aztekin, C., Hiscock, T.W., Marioni, J.C., Gurdon, J.B., Simons, B.D., and Jullien, J. (2019).
Identification of a regeneration-organizing cell in the Xenopus tail. Science *364*, 653-658.
10.1126/science.aav9996.
38. Srivastava, M., Mazza-Curll, K.L., van Wolfswinkel, J.C., and Reddien, P.W. (2014). Whole-body
acoel regeneration is controlled by Wnt and Bmp-Admp signaling. Current biology : CB *24*, 1107-
1113. 10.1016/j.cub.2014.03.042.
39. Lander, R., and Petersen, C.P. (2016). Wnt, Ptk7, and FGFRL expression gradients control trunk
positional identity in planarian regeneration. Elife *5*. 10.7554/eLife.12850.
40. Petersen, C.P., and Reddien, P.W. (2009). A wound-induced Wnt expression program controls
planarian regeneration polarity. Proc Natl Acad Sci U S A *106*, 17061-17066.
10.1073/pnas.0906823106.
41. Chen, B., Dodge, M.E., Tang, W., Lu, J., Ma, Z., Fan, C.W., Wei, S., Hao, W., Kilgore, J.,
Williams, N.S., et al. (2009). Small molecule-mediated disruption of Wnt-dependent signaling in
tissue regeneration and cancer. Nat Chem Biol *5*, 100-107. 10.1038/nchembio.137.
42. Taira, N., Mimoto, R., Kurata, M., Yamaguchi, T., Kitagawa, M., Miki, Y., and Yoshida, K. (2012).
DYRK2 priming phosphorylation of c-Jun and c-Myc modulates cell cycle progression in human
cancer cells. The Journal of Clinical Investigation *122*, 859-872. 10.1172/jci60818.
43. Todd, J.T. (1823). On the process of reproduction of the members of the aquatic salamander. .
The Quarterly Journal of Science, Literature, and the Arts. *16*, 84-96.
44. Singer, M. (1946). The nervous system and regeneration of the forelimb of adult Triturus; the
influence of number of nerve fibers, including a quantitative study of limb innervation. J Exp Zool
101, 299-337.
45. Singer, M. (1952). The influence of the nerve in regeneration of the amphibian extremity. Q Rev
Biol *27*, 169-200.
46. Brockes, J.P. (1987). The nerve dependence of amphibian limb regeneration. The Journal of
experimental biology *132*, 79-91.
47. Farkas, J.E., and Monaghan, J.R. (2017). A brief history of the study of nerve dependent
regeneration. Neurogenesis (Austin) *4*, e1302216. 10.1080/23262133.2017.1302216.
48. Wells, K.M., Kelley, K., Baumel, M., Vieira, W.A., and McCusker, C.D. (2021). Neural control of
growth and size in the axolotl limb regenerate. Elife *10*. 10.7554/eLife.68584.
49. Tassava, R.A., and Loyd, R.M. (1977). Injury requirement for initiation of regeneration of newt
limbs which have whole skin grafts. Nature *268*, 49-50. 10.1038/268049a0.
50. Zeng, A., Li, H., Guo, L., Gao, X., McKinney, S., Wang, Y., Yu, Z., Park, J., Semerad, C., Ross,
E., et al. (2018). Prospectively Isolated Tetraspanin(+) Neoblasts Are Adult Pluripotent Stem
Cells Underlying Planaria Regeneration. Cell *173*, 1593-1608 e1520. 10.1016/j.cell.2018.05.006.
51. Zhang, X., Yalcin, S., Lee, D.F., Yeh, T.Y., Lee, S.M., Su, J., Mungamuri, S.K., Rimmele, P.,
Kennedy, M., Sellers, R., et al. (2011). FOXO1 is an essential regulator of pluripotency in human
embryonic stem cells. Nat Cell Biol *13*, 1092-1099. 10.1038/ncb2293.
52. Ogg, S., Paradis, S., Gottlieb, S., Patterson, G.I., Lee, L., Tissenbaum, H.A., and Ruvkun, G.
(1997). The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity
signals in C. elegans. Nature *389*, 994-999. 10.1038/40194.
53. Brunet, A., Bonni, A., Zigmond, M.J., Lin, M.Z., Juo, P., Hu, L.S., Anderson, M.J., Arden, K.C.,
Blenis, J., and Greenberg, M.E. (1999). Akt promotes cell survival by phosphorylating and
inhibiting a Forkhead transcription factor. Cell *96*, 857-868. 10.1016/s0092-8674(00)80595-4.
54. Collins, J.P., Cheek, J.E. (1983). Effect of Food and Density on Development of Typical and
Cannibalistic Salamander Larvae in Ambystoma tigrinum nebulosum. Integr Comp Biol *23*, 77-84.
55. Keel, M., and Trentz, O. (2005). Pathophysiology of polytrauma. Injury *36*, 691-709.
10.1016/j.injury.2004.12.037.
56. Khattak, S., Schuez, M., Richter, T., Knapp, D., Haigo, S.L., Sandoval-Guzman, T., Hradlikova,
K., Duemmler, A., Kerney, R., and Tanaka, E.M. (2013). Germline transgenic methods for
tracking cells and testing gene function during regeneration in the axolotl. Stem cell reports *1*, 90-
103. 10.1016/j.stemcr.2013.03.002.
57. Schotté, O.E., and Butler, E.G. (1941). Morphological effects of denervation and amputation of
limbs in urodele larvae. Journal of Experimental Zoology *87*, 279-322.
https://doi.org/10.1002/jez.1400870207.

58. Kropf, N., Krishnan, K., Chao, M., Schweitzer, M., Rosenberg, Z., and Russell, S.M. (2010).
Sciatic nerve injury model in the axolotl: functional, electrophysiological, and radiographic
outcomes. J Neurosurg *112*, 880-889. 10.3171/2008.10.Jns08222.
59. Lovely, A.M., Duerr, T.J., Stein, D.F., Mun, E.T., and Monaghan, J.R. (2022). Hybridization Chain
Reaction Fluorescence In Situ Hybridization (HCR-FISH) in Ambystoma mexicanum Tissue. In
Salamanders: Methods and Protocols, (Springer), pp. 109-122.
60. Debuque, R.J., and Godwin, J.W. (2015). Methods for axolotl blood collection, intravenous
injection, and efficient leukocyte isolation from peripheral blood and the regenerating limb.
Methods Mol Biol *1290*, 205-226. 10.1007/978-1-4939-2495-0_17.
61. McQuin, C., Goodman, A., Chernyshev, V., Kamentsky, L., Cimini, B.A., Karhohs, K.W., Doan,
M., Ding, L., Rafelski, S.M., Thirstrup, D., et al. (2018). CellProfiler 3.0: Next-generation image
processing for biology. PLOS Biology *16*, e2005970. 10.1371/journal.pbio.2005970.
62. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-
image analysis. Nature Methods *9*, 676-682. 10.1038/nmeth.2019.
63. Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan,
L., Raychowdhury, R., Zeng, Q., et al. (2011). Full-length transcriptome assembly from RNA-Seq
data without a reference genome. Nat Biotechnol *29*, 644-652. 10.1038/nbt.1883.
64. Schloissnig, S., Kawaguchi, A., Nowoshilow, S., Falcon, F., Otsuki, L., Tardivo, P.,
Timoshevskaya, N., Keinath, M.C., Smith, J.J., Voss, S.R., and Tanaka, E.M. (2021). The giant
axolotl genome uncovers the evolution, scaling, and transcriptional control of complex gene loci.
Proceedings of the National Academy of Sciences of the United States of America *118*.
10.1073/pnas.2017176118.
65. Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics *34*, 3094-
3100. 10.1093/bioinformatics/bty191.
66. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and
Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-
2079. 10.1093/bioinformatics/btp352.
67. Neph, S., Kuehn, M.S., Reynolds, A.P., Haugen, E., Thurman, R.E., Johnson, A.K., Rynes, E.,
Maurano, M.T., Vierstra, J., Thomas, S., et al. (2012). BEDOPS: high-performance genomic
feature operations. Bioinformatics *28*, 1919-1920. 10.1093/bioinformatics/bts277.
68. Arnason, U., Gullberg, A., Janke, A., Joss, J., and Elmerot, C. (2004). Mitogenomic analyses of
deep gnathostome divergences: a fish is a fish. Gene *333*, 61-70. 10.1016/j.gene.2004.02.014.
69. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics *26*, 841-842. 10.1093/bioinformatics/btq033.
70. Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol *34*, 525-527. 10.1038/nbt.3519.
71. Gao, F., da Veiga Beltrame, E., Gehring, J.A., Edljarn Hjoerleifsson, K.E., Lu, L., Melsted, P.,
Ntranos, V., Pachter, L., and Svensson, V. (2019). The BUS Format for Single-Cell RNA-Seq
Processing and Analysis. J Biomol Tech *30*, S62-S62.
72. Melsted, P., Booeshaghi, A.S., Liu, L., Gao, F., Lu, L., Min, K.H.J., da Veiga Beltrame, E.,
Hjörleifsson, K.E., Gehring, J., and Pachter, L. (2021). Modular, efficient and constant-memory
single-cell RNA-seq preprocessing. Nat Biotechnol *39*, 813-818. 10.1038/s41587-021-00870-2.
73. Moses L, P.L. (2022). BUSpaRse: kallisto | bustools R utilities.
74. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J., Wilk,
A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell
184, 3573-3587.e3529. 10.1016/j.cell.2021.04.048.
75. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome biology *15*, 550. 10.1186/s13059-014-0550-
8.
76. Wickham, H. (2009). Elegant graphics for data analysis (ggplot2). Applied Spatial Data Analysis
R.
77. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H., Myung, P., Plikus,
M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using CellChat.
Nature communications *12*, 1088. 10.1038/s41467-021-21246-9.

78.   La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., Lidschreiber, K.,
1076   Kastriti, M.E., Lönnerberg, P., Furlan, A., et al. (2018). RNA velocity of single cells. Nature *560*,
1077   494-498. 10.1038/s41586-018-0414-6.

1078   79.   Bergen, V., Lange, M., Peidli, S., Wolf, F.A., and Theis, F.J. (2020). Generalizing RNA velocity to
1079   transient cell states through dynamical modeling. Nature Biotechnology *38*, 1408-1414.
1080   10.1038/s41587-020-0591-3.

1081   80.   Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
1082   sequence data. Bioinformatics *30*, 2114-2120. 10.1093/bioinformatics/btu170.

1083   81.   Soneson, C., Love, M.I., and Robinson, M.D. (2015). Differential analyses for RNA-seq:
1084   transcript-level estimates improve gene-level inferences. F1000Res *4*, 1521.
1085   10.12688/f1000research.7563.2.

1086   82.   Kolde, R. (2019). pheatmap: Pretty Heatmaps. R package version 1.0. 12. R Packag. version 1.0
1087   *8*.

1088   83.   Ulgen, E., Ozisik, O., and Sezerman, O.U. (2019). pathfindR: An R Package for Comprehensive
1089   Identification of Enriched Pathways in Omics Data Through Active Subnetworks. Front Genet *10*,
1090   858. 10.3389/fgene.2019.00858.

1091   84.   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
1092   Methods *9*, 357-359. 10.1038/nmeth.1923.

1093   85.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
1094   Durbin, R., and Subgroup, G.P.D.P. (2009). The Sequence Alignment/Map format and SAMtools.
1095   Bioinformatics *25*, 2078-2079. 10.1093/bioinformatics/btp352.

1096   86.   Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dündar,
1097   F., and Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data
1098   analysis. Nucleic acids research *44*, W160-165. 10.1093/nar/gkw257.

1099   87.   Kent, W.J., Zweig, A.S., Barber, G., Hinrichs, A.S., and Karolchik, D. (2010). BigWig and BigBed:
1100   enabling browsing of large distributed datasets. Bioinformatics *26*, 2204-2207.
1101   10.1093/bioinformatics/btq351.

1102   88.   Li, Q., Brown, J.B., Huang, H., and Bickel, P.J. (2011). Measuring reproducibility of high-
1103   throughput experiments. The Annals of Applied Statistics *5*, 1752-1779, 1728.

1104   89.   Yu, G., Wang, L.-G., and He, Q.-Y. (2015). ChIPseeker: an R/Bioconductor package for ChIP
1105   peak annotation, comparison and visualization. Bioinformatics *31*, 2382-2383.
1106   10.1093/bioinformatics/btv145.

1107   90.   Marini, F., and Binder, H. (2019). pcaExplorer: an R/Bioconductor package for interacting with
1108   RNA-seq principal components. BMC Bioinformatics *20*, 331. 10.1186/s12859-019-2879-1.

1109   91.   Lopez-Delisle, L., Rabbani, L., Wolff, J., Bhardwaj, V., Backofen, R., Grüning, B., Ramírez, F.,
1110   and Manke, T. (2020). pyGenomeTracks: reproducible plots for multivariate genomic
1111   datasets Bioinformatics *37*, 422-423. 10.1093/bioinformatics/btaa692.

1112   92.   Carpenter, A.E., Jones, T.R., Lamprecht, M.R., Clarke, C., Kang, I.H., Friman, O., Guertin, D.A.,
1113   Chang, J.H., Lindquist, R.A., Moffat, J., et al. (2006). CellProfiler: image analysis software for
1114   identifying and quantifying cell phenotypes. Genome biology *7*, R100. 10.1186/gb-2006-7-10-
1115   r100.

1116   93.   Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq
1117   quantification. Nature Biotechnology *34*, 525-527. 10.1038/nbt.3519.

1118

Figure 1

Figure 4

Figure 5

Figure 6

Figure S1, related to Figure 2

A

Vehicle   Rapamycin

B

B

Vehicle

Rapamycin

Figure S2, related to Figure 3

Figure S3, related to Figure 5

Figure S4, related to Figure 4