The evolution and biological correlates of hand preferences in anthropoid primates

Kai R. Caspar[1,2*], Fabian Pallasdies[3], Larissa Mader[1], Heitor Sartorelli[4], Sabine Begall[1]

[1]Department of General Zoology, University of Duisburg-Essen, Universitaetsstraße 5, 45141 Essen, Germany

[2]Department of Game Management and Wildlife Biology, Faculty of Forestry and Wood Sciences, Czech
University of Life Sciences, Praha, Czech Republic.

[3]Institute for Theoretical Biology, Department of Biology, Humboldt-Universität zu Berlin, Berlin, Germany

[4]Independent researcher

Abstract

The evolution of human right-handedness has been intensively debated for decades. Manual
lateralization patterns in non-human primates have the potential to elucidate evolutionary
determinants of human handedness. However, restricted species samples and inconsistent
methodologies have so far limited comparative phylogenetic studies. By combining original data with
published literature reports, we assembled data on hand preferences for standardized object
manipulation in 1,786 individuals from 38 species of anthropoid primates, including monkeys, apes,
and humans. Based on that, we employ quantitative phylogenetic methods to test prevalent
hypotheses on the roles of ecology, brain size and tool use in primate handedness evolution. We
confirm that human right-handedness represents an unparalleled extreme among anthropoids and
found taxa displaying significant population-level handedness to be rare. Species-level direction of
manual lateralization was largely uniform among non-human primates and did not notably correlate
with any of the selected biological predictors, nor with phylogeny. In contrast, we recovered highly
variable patterns of hand preference strength, which show signatures of both ecology and phylogeny.
In particular, terrestrial primates tend to display weaker hand preferences than arboreal species. These
results challenge popular ideas on primate handedness evolution, especially the postural origins
hypothesis. Furthermore, they point to a potential adaptive benefit of disparate lateralization strength
in primates, a measure of hand preference that has often been overlooked in the past. Finally, our data
show that human lateralization patterns do not align with trends found among other anthropoids,
suggesting that unique selective pressures gave rise to the unusual hand preferences displayed by our
species.

Key words: handedness, laterality, hemispheric dominance, tube task, comparative cognition, postural
origins hypothesis

Introduction

Pronounced right-handedness is a universal trait among extant human populations (Coren & Porac,
1977; Raymond & Pontier, 2004; Faurie et al., 2005) and might be an ancient attribute of the genus
Homo (Toth, 1985; Lozano et al., 2017). Whereas a significant expression of manual lateralization at
population-level is not exclusive to humans, the universal proportion of approximately 85-95 % right-
handers in our species appears to be an unmatched extreme among primates (Meguerditchian et al.,
2013). Furthermore, individual humans tend to be strongly handed with ambiguous hand
preferences being extremely rare (Cochet & Vauclair, 2012), which is also unusual when compared to
many other primate lineages (Westergaard & Suomi, 1996; Hopkins et al., 2011). Therefore, both

strength and direction of population-level manual lateralization in humans are widely considered to
be remarkable.

Handedness is a behavioral consequence of functional asymmetries in the brain (Amunts et al., 1996;
Häberling & Corballis, 2016; Sha et al., 2021). Accordingly, right-handedness results from unilateral
specializations in the left hemisphere and vice versa. In anthropoid primates, which encompass
monkeys, apes, and humans, asymmetries of the precentral gyrus in the primary motor cortex show
a particular association with individual hand preferences (Yousry et al., 1997; Phillips & Sherwood,
2005; Dadda et al., 2006; Sha et al., 2021). Nevertheless, the proximate reasons for the expression of
individual handedness in humans and other anthropoids, including its genetic basis and the influence
of brain areas located outside of the motor cortex, are by no means fully understood (Rogers, 2009;
Hopkins et al. 2013; Ocklenburg et al., 2014; Schmitz et al., 2017a; Richards et al., 2021). For this
paper, however, we exclusively focus on the evolutionary underpinnings of population-level manual
lateralization.

The origins of pronounced population-level right-handedness in the human lineage have traditionally
been linked to the emergence of complex communication mediated by manual gestures and
language, which are also predominantly processed in the left hemisphere (Corballis, 1991; Annett,
2002; Meguerditchian et al., 2013; Ocklenburg et al., 2014; Prieur et al., 2019). However, current
evidence suggests that manual and language lateralization are not nearly as tightly correlated as
traditionally believed and functional ties between these phenomena remain unidentified (Fitch &
Braccini, 2013; Ocklenburg et al., 2014; Schmitz et al., 2017b). It has also been established that
various nonhuman primates show significant asymmetries in hand-use at the individual and
population-level in a variety of tasks including manual communicative gestures and bimanual actions
(MacNeilage, 2007; Hopkins et al., 2013; Meguerditchian et al., 2013; Regaiolli et al., 2016).
Nevertheless, reports of significant population-level biases are largely confined to a comparatively
small number of species and the distribution of individuals across hand preference categories is
always far more balanced than in humans. For instance, olive baboons (*Papio anubis*), Western
gorillas (*Gorilla gorilla*), and chimpanzees (*Pan troglodytes*) have all been reported to show a
significant population-level right hand bias for bimanual manipulation but the portion of right-
handers among these species is only around 50 % (Vauclair et al., 2005; Hopkins et al., 2011 – note
that this skew is significant due to a notable portion of ambipreferent individuals among these
species). Still, it is essential to discern what underlies these comparatively weak population-level
hand preference patterns that emerged across the primate order to unravel the origins of
pronounced right-handedness in our species (MacNeilage, 2007).

The most influential conjecture to explain how primate hand preference patterns evolved is the
postural origins hypothesis (POH) (MacNeilage et al., 1987; MacNeilage, 2007). Considering galagos
as models, the POH assumes that hypothetical insectivorous primate ancestors exhibited a right-
hand bias to support their body on vertical substrates, while the left-hand specialized for fast
grasping movements, the so-called ballistic reaching (Ward, 1998). Based on this, the POH predicts
that with the emergence of anthropoid primates, which exhibit arboreal quadrupedalism and more
refined digit control, the right hand became adopted to manipulate objects during foraging
(MacNeilage et al., 1987). Hence, it proposes that all anthropoids share a right-hand bias for
manipulation, which would find its most extreme expression in humans (MacNeilage, 2007;
MacNeilage et al., 1987). In the anatomically less derived strepsirrhines, the left hand is instead
expected to be dominant (MacNeilage, 2007). However, the POH has been drastically modified by
more recent studies (Hopkins et al., 2011; Meguerditchian et al., 2013; Morino et al., 2017). The
novel interpretation proposes (in conflict with the original POH) that arboreal monkeys and apes
should display a left-hand bias for manipulation. Their right hand would provide the necessary

postural support, retaining the hypothesized ancestral primate pattern laid out by MacNeilage et al.
(1987). Terrestrial lineages, however, would no longer be bound to reserve the right hand for posture
stabilization and are expected to evolve right hand preferences for fine motor skills, eventually giving
rise to the human condition. Hence, the novel POH expects that left-handedness is prevalent in
arboreal groups, while a preponderance of right-handedness should be restricted to terrestrial
primate species (Meguerditchian et al., 2013). In contrast, the original POH expects to find a right-
handedness trend in all anthropoids, regardless of their ecology. Both versions of the POH exclusively
rely on correlational evidence from behavioral studies on hand preferences in different primate
groups.

In addition to these considerations, it has been prominently proposed that foraging-related
extractive tool use facilitated the evolution of hand preferences in humans to allow for more efficient
object handling (Kimura, 1979; Frost, 1980), but this idea has also been extended to nonhuman
primates that habitually use tools (Cashmore et al., 2008; Prieur et al., 2019). Moreover,
neuroanatomical studies demonstrated that the expression of overall neural lateralization and
hemispheric independence positively correlates with absolute brain size in primates (Rilling & Insel,
1999; Karolis et al., 2019; Ardesch et al., 2021). Such a scaling relation was already hypothesized by
Ringo et al. (1994) and implies that the strength of individual handedness could also be tied to
absolute brain size (Hopkins, 2013b); the concept does, however, not concern the direction of hand
preferences. Hence, it can be hypothesized that larger-brained primates should evolve greater
manual lateralization strength to reduce the amount of interhemispheric communication needed to
accomplish manipulative tasks. Complementary to these predictions, there is evidence that absolute
brain size is positively correlated with foot preference strength for grasping among parrot species
(Kaplan & Rogers, 2021; but note that this study did not control for phylogeny; see also Brown and
Magat, 2011).

Thus testable hypotheses on primate hand preference evolution have long been established, but
none of them has so far been tested within quantitative evolutionary frameworks (Hopkins, 2013a).
Studies in comparative cognition increasingly rely on phylogenetically informed modelling to
estimate how and when specific behaviors evolved (MacLean et al., 2012; ManyPrimates et al., 2019;
Krasheninnikova et al., 2020). Such approaches can provide estimates of ancestral states and allow
researchers to quantify the pace as well as the influence of phylogeny and ecological variables on
cognitive evolution (MacLean et al., 2012). However, to yield meaningful results, dense taxonomic
sampling of species and a consistent testing scheme are required (Freckleton et al., 2002;
Krasheninnikova et al., 2020). For research on primate hand preferences, this means that subjects
from different species need to engage in the same experimental task to assess lateralization.
Standardization is particularly important since both the strength and the direction of individual hand
preferences can be task-dependent and because the expression of manual lateralization correlates
positively with motor complexity (Vauclair et al., 2005; Blois-Heulin et al., 2007; Lilak & Phillips, 2008;
Meguerditchian et al., 2015; Caspar et al., 2018). Bimanual actions that involve both hands and which
spontaneously occur during food manipulation or tool use are particularly suitable to detect hand
preferences (Meguerditchian et al., 2013), while non-gestural unimanual actions such as grasping
often do not elicit pronounced group or individual-level lateralized responses (Papademetriou et al.,
2005; Rogers, 2009; Hopkins et al., 2013a). For comparative approaches in primates, tool handling is
only of limited use as a model behavior, because habitual tool use is only found in few lineages
(Musgrave & Sanz, 2018). Therefore, bimanual actions related to foraging present themselves as
suitable candidates for comparative studies on hand preferences in these animals.

A simple and widely applicable experiment to determine anthropoid primate hand preferences is the
so-called tube task (Hopkins, 1995): A subject is handed a PVC tube filled with desired food. To

extract it, one hand must hold the tube while the other has to engage in the more complicated action of retrieving the food mesh, thereby revealing biases in hand use dominance. Results from the tube task have been demonstrated to robustly correlate with hemispheric asymmetries in various primate groups (New World monkeys: Phillips & Sherwood, 2005; Old World monkeys: Margiotoudi et al., 2019; apes: Hopkins & Cantalupo, 2004; Dadda et al., 2006) and to be intraindividually consistent across re-tests, even if these were separated by several years (Hopkins et al., 2001). Furthermore, its simplicity allows uniform testing of a wide range of species in captive as well as natural settings (Zhao et al., 2012).

Here we compiled a comprehensive multispecies tube task dataset to test pertaining hypotheses on the evolution of primate manual lateralization by means of phylogenetically informed modelling. This way we provide a broad comparative perspective on the origins of human right-handedness in the context of object manipulation.

Materials and Methods

Subjects

We analyzed the expression of hand preferences for object manipulation in the tube task, as well as potential factors influencing their evolution, for a dataset of anthropoid primates (infraorder Simiiformes: New World monkeys (Platyrrhini), Old World monkeys (Cercopithecoidea), and apes (Hominoidea)) from 38 species. Data from 501 individuals belonging to 26 primate species were collected in the tube task paradigm (see below) between September 2017 and May 2020 in 39 institutions in Europe, Brazil, and Indonesia (Table 1). Of these species, 14 had never been tested in the tube task before. Additional datasets were drawn from the literature, resulting in a total sample of 1,786 individuals from 38 species and 20 genera, covering all anthropoid primate families except Aotidae. Data for humans were derived from Cochet and Vauclair (2012). In this study, participants had to use one hand to repeatedly retrieve pieces of paper out of a plastic cylinder while the other one had to tilt and stabilize the receptacle. We considered this bimanual testing paradigm as functionally equivalent to the tube task. Our complete study sample with respective data sources is included in Supplementary Table 1.

We classified the tested subjects into two age categories, adults (n = 1,307, sexually mature individuals) and subadults; the latter being comprised by infants (n = 9, individuals that had not yet been weaned) and juveniles (n = 456, weaned individuals that have not reached sexual maturity). If previous tube task studies assigned age categories to their subjects, we adopted this classification for the individuals concerned. In other cases and for our original data, age classification followed life history data from Harvey and Clutton-Brock (1985). The taxonomy and nomenclature we apply follows Mittermeier, Rylands, and Wilson (2013) with the following exceptions: The recently diverging sister species pairs *Cercopithecus diana* and *C. roloway* (Diana and Roloway monkeys) as well as *Nomascus leucogenys* and *N. siki* (white-cheeked gibbons) are treated here as one respective taxonomic unit and data were pooled to obtain larger sample sizes (species identity of subjects is noted in Suppl. Table 1). Because the hand preference literature on orangutans (*Pongo* spp.) did not consider the species status of the individuals concerned, we analyzed respective data on the genus level. In other cases, we carefully checked the current taxonomic status of subjects drawn from the literature and tried to avoid the inclusion of interspecific hybrids. This was particularly relevant for data on lab-housed tufted capuchins (*Sapajus* spp.). If the species or hybrid status of animals was ambiguous, we did not consider them for our analyses (e.g., capuchins in Westergaard and Suomi, 1996).

Although available for testing at most of the institutions we visited, lemurs could not be included into
the study. All tested genera (*Eulemur*, *Hapalemur*, *Propithecus*, *Varecia*), failed to manually remove
food mash from the tube, despite eagerly licking it up from the ends. Only a single subject, a female
Eulemur rubriventer, succeeded (Suppl. Table 2). White-faced sakis (*Pithecia pithecia*) and Javan
langurs (*Trachypithecus auratus*) were also reluctant to engage in the task, so that the final sample
for these species is smaller than expected from their abundance at the institutions visited. Apart
from the species considered for analysis, nine individuals belonging to miscellaneous taxa were
sampled in the study (data included in Suppl. Table 1).

Experimental procedure and data scoring

All species were uniformly tested in the established bimanual tube task paradigm (Hopkins, 1995).
Due to the pronounced differences in body size between the observed species, PVC tubes of varying
length and diameter were employed (Fig. 1). Lion tamarins (*Leontopithecus*) were presented with
small-sized tubes that were 5 cm long and had an inner diameter of 1 cm. Capuchins and sakis
(*Sapajus*, *Pithecia*) received 10 cm x 2 cm medium-sized tubes and all remaining species large tubes
measuring 10 cm x 2.5 cm. The tubes were filled with various food incentives, which differed
dependent on the nutrition regimes enacted by the respective institutions. Among preferred food
items for cercopithecines, gibbons, spider monkeys, and capuchins were oatmeal mixed with banana
mash, soaked pellets, and boiled carrots (but note that the latter did not appeal to *Cercopithecus* and
Ateles). Geladas (*Theropithecus gelada*) exclusively received boiled carrots. Langurs (*Semnopithecus*,
Trachypithecus) and sakis were preferably tested with boiled rice, and tubes for the latter were also
stowed with nuts as an additional incentive. Lion tamarins received tubes filled with pure banana
mash or commercial tamarin pie. Primates were preferably tested within their social groups. A
separation of individuals was only undertaken in exceptional cases when it was necessary to
counteract social tension created by the presentation of the tubes. Dependent on the constructional
restraints of the enclosures, tubes were either placed into a separated part of the enclosure before
the primates could enter or were handed over directly through the wire mesh. In the latter case, the
hand that the experimenter used to offer the tube was noted (Suppl. Table 2). To check whether the
hand used by the experimenter to offer the tube had an effect on the directional hand preferences of
the tested primates, we ran a linear mixed effect model employing a binomial link function. No effect
on the recovered hand preferences in the respective sessions was found (t = -1.31, SE = 0.02, *p* =
0.191).

The tube tasks were recorded with digital cameras and scored from the video footage. For each
subject, we obtained a minimum of 30 bimanual insertions (one hand is holding the tube, the other
one is retrieving food; mean number of insertions: 50.66 ± 20.98, range: 30-155) in at least six bouts
(uninterrupted manipulation sequences, as defined by Morino et al. (2017); mean number of bouts:
20.60 ± 11.13, range: 6-82). Literature data for individual subjects had to match or exceed these
thresholds to be included into the analysis. Unimanual or foot-assisted insertions were not scored
and were, whenever possible, also carefully excluded from the literature data. We noted the digit
used to extract the food as well as the body posture of the manipulating individual (sitting, crouched
bipedal stance, erect bipedal stance, suspended (clinging to wire mesh or other substrates without
the hands stabilizing posture, always tail-assisted in spider monkeys)). The vast majority of responses
were observed in a sitting position (*n* = 22,993; 90.6 %). Due to this imbalance, because posture-
related information was mostly unavailable for literature data, and since its influence on manual
laterality already received great research attention in the past (Sanford et al., 1984; Westergaard et
al., 1998; Blois-Heulin et al., 2007; Laurence et al., 2011), we refrained from including posture effects
into our analyses. Nevertheless, for potential future use by other researchers, we decided to include

this measure, alongside information on digit use ("N.A." if ambiguous in the respective footage)
during manipulation, in Supplementary Table 2.

Table 1: Composition of the study sample, listing taxonomic identity, sex, age, and origin of subjects. See cited studies for locations of individuals drawn from

the literature.

Family	Species	# Subjects tested	# Subjects drawn from literature*	Total sample	# adult females	# adult males	# subadult females	# subadult males	# unsexed subadults	Locations for subjects in this study
Atelidae	*Ateles fusciceps*	37	9[a]	46	30	11	3	2	0	Berlin (Zoo), Doué-la-Fontaine, Landau, Mulhouse, Munich, Osnabrück, Wuppertal
Atelidae	*Ateles geoffroyi*	9	14[b]	23	12	9	0	2	0	Basel, Karlsruhe
Atelidae	*Ateles hybridus*	18		18	10	7	0	1	0	Doué-la-Fontaine, Erfurt, Frankfurt, Neuwied, Stuttgart
Callitrichidae	*Leontopithecus chrysomelas*	30		30	11	16	2	1	0	Apeldoorn, Karlsruhe, Magdeburg, Mulhouse, São Paulo, Stuttgart, Wuppertal
Callitrichidae	*Leontopithecus chrysopygus*	15		15	6	9	0	0	0	São Paulo
Callitrichidae	*Leontopithecus rosalia*	28		28	7	16	0	5	0	Apeldoorn, Basel, Doué-la-Fontaine, Duisburg, Frankfurt, Heidelberg, Landau, Magdeburg, Münster, São Paulo
Cebidae	*Saimiri sciureus*		36[c]	36	14	16	5	1	0	
Cebidae	*Sapajus apella*		25[d, e]	25	10	11	0	4	0	
Cebidae	*Sapajus flavius*	3	18[e]	21	7	9	2	3	0	São Paulo
Cebidae	*Sapajus xanthosternos*	16	18[e]	34	11	19	1	2	1	Apeldoorn, Magdeburg, Münster, Overloon
Cercopithecidae	*Cercocebus torquatus*	18	13[f]	31	15	13	1	2	0	Apeldoorn, Berlin (Tierpark), Karlsruhe, Münster
Cercopithecidae	*Cercopithecus diana/roloway*	20		20	9	7	3	1	0	Amsterdam, Berlin (Tierpark), Doué-la-Fontaine, Duisburg, Heidelberg, Liberec, Mulhouse, Osnabrück
Cercopithecidae	*Cercopithecus neglectus*	12	13[f, g]	25	8	12	1	4	0	Bekesbourne, Duisburg, Hannover, Overloon
Cercopithecidae	*Macaca fascicularis*	12	8[h]	20	13	7	0	0	0	Basel, Hamm, Mönchengladbach
Cercopithecidae	*Macaca nemestrina*	29		29	12	15	0	1	1	Arnhem, Bali, Berlin (Tierpark), Gelsenkirchen, Osnabrück
Cercopithecidae	*Macaca silenus*	35		35	16	17	1	1	0	Apeldoorn, Bekesbourne, Berlin (Zoo), Cologne, Dresden, Duisburg, Hodenhagen, Rheine
Cercopithecidae	*Macaca sylvanus*	15	9[i, j]	24	11	12	0	1	0	Aachen, Rheine
Cercopithecidae	*Macaca tonkeana***		14[k]	14	NA	NA	NA	NA	NA	
Cercopithecidae	*Mandrillus sphinx*	32		32	14	7	4	7	0	Amsterdam, Berlin (Zoo), Dresden, Hamm, Hodenhagen
Cercopithecidae	*Papio anubis*		84[l]	84	48	22	5	9	0	
Cercopithecidae	*Papio hamadryas*	24		24	14	10	0	0	0	Cologne, Frankfurt, Krefeld
Cercopithecidae	*Pygathrix cinerea*		18[m]	18	7	11	0	0	0	
Cercopithecidae	*Rhinopithecus roxellana*		24[n]	24	8	5	8	3	0	

Family	Species									Location
Cercopithecidae	*Semnopithecus entellus*	30		30	17	7	4	2	0	Apeldoorn, Berlin (Zoo), Gelsenkirchen, Hannover, Heidelberg
Cercopithecidae	*Theropithecus gelada*	38		38	20	11	4	3	0	Bekesbourne, Berlin (Tierpark), Magdeburg, Rheine, Stuttgart
Cercopithecidae	*Trachypithecus auratus*	8		8	3	0	3	2	0	Bali, Stuttgart
Cercopithecidae	*Trachypithecus hatinhensis*		18[m]	18	8	10	0	0	0	
Hominidae	*Gorilla gorilla*		76[o]	76	22	18	19	17	0	
Hominidae	*Homo sapiens*		127[p]	127	71	56	0	0	0	
Hominidae	*Pan paniscus*		118[o]	118	29	23	35	31	0	
Hominidae	*Pan troglodytes*		536[o]	536	186	138	110	102	0	
Hominidae	*Pongo* sp.		47[o]	47	17	12	9	9	0	
Hylobatidae	*Hylobates lar*	16	20[q, r, s]	36	14	18	2	2	0	Berlin (Tierpark), Cologne, Doué-la-Fontaine, Landau, Stuttgart, Ulm, Wuppertal
Hylobatidae	*Hylobates moloch*	22		22	8	5	4	5	0	Bekesbourne, Lympne, Munich
Hylobatidae	*Nomascus gabriellae*	6	4[r]	10	5	3	0	2	0	Arnhem, Doué-la-Fontaine
Hylobatidae	*Nomascus leucogenys /siki*	7	19[q,r,t]	26	15	7	1	3	0	Apeldoorn, Frankfurt, Osnabrück
Hylobatidae	*Symphalangus syndactylus*	14	17[q,r]	31	12	11	4	4	0	Arnhem, Bekesbourne, Berlin (Zoo), Dortmund, Doué-la-Fontaine, Munich, Arnhem, Hodenhagen, Osnabrück
Pitheciidae	*Pithecia pithecia*	7		7	4	3	0	0	0	Basel, Dresden, Krefeld
Total***		**501**	**1285**	**1786**	**724**	**583**	**231**	**232**	**2**	

* Fulfilling our criteria
** Ages unknown, sex derived from given names
*** Not including *M. tonkeana* in sex and age specific categories

[a] Nelson and Boeving (2015), [b]Motes Rodrigo et al. (2018), [c]Meguerditchian et al. (2012), [d]Phillips et al. (2007); [e]de Andrade and de Sousa (2018), [f]Maille et al. (2013),

[g]Schweitzer et al. (2007), [h]Chatagny et al. (2013), [i]Schmitt et al. (2008), [j]Regaiolli et al. (2018), [k]Canteloup et al. (2013), [l]Vauclair et al. (2005), [m]Cubí and Llorente (2021), [n]Zhao

et al. (2012), [o]Hopkins et al. (2011), [p]Cochet and Vauclair (2012), [q]Morino et al. (2017), [r]Caspar et al. (2018), [s]Spoelstra (2021), [t]Fan et al. (2017).

Figure 1: Various anthropoid primates engaging in the tube task. A: Golden lion tamarin
(*Leontopithecus rosalia*) manipulating a small tube at Zoo Frankfurt, Germany. B: White-handed
gibbon (*Hylobates lar*) handling a large tube at Bioparc de Doué-la-Fontaine, France. Note that the
thumb is used to probe into the tube, an insertion pattern characteristic of gibbons. C: Yellow-
breasted capuchin (*Sapajus xanthosternos*) engaging in the task with a medium-sized tube while
adopting an erect bipedal stance at ZooParc Overloon, the Netherlands. Photographs by Kai R.
Caspar.

Statistics

Data were analyzed in R (R Core Team, 2020). Preferably, analyses were performed on insertion data
(also called frequencies) instead of manipulations bouts to approximate laterality. Unfortunately, not
all available tube task studies provided insertion data (Maille et al., 2013; Spoelstra, 2021), so that in
the final dataset, estimates of manual lateralization based on insertions and bouts had to be mixed
for three species (*Cercocebus torquatus*, *Cercopithecus neglectus*, and *Hylobates lar*). However, since
previous work demonstrated that hand preferences derived from bouts and insertions are highly
correlated and non-conflicting, we do not consider this a confounding factor for our analysis (Hopkins
et al., 2001; Hopkins, 2013a).

For quantifying lateralized responses on the individual level, we calculated handedness indices (HI)
for all subjects as well as the corresponding binomial z scores to allow grouping into hand preference
categories. HI is a descriptive index that can range from -1 (all manipulations left-handed) to 1 (all
right-handed) and is calculated via the formula HI = (R − L)/ (R+L). The z score, on the other hand,
indicates whether there is a statistically significant bias in hand use. Following established criteria
(Hopkins, 2013a), we rated subjects with z score values higher than 1.96 as right-handed, those with
values lower than −1.96 as left-handed, and the remaining ones as ambipreferent. We use the term
"ambipreferent" here instead of "ambidextrous" or "ambiguously handed" to indicate a lack of
preferences, because the latter two expressions have clear non-synonymous definitions when

applied to human, but are not consistently used in the non-human primate literature (Hopkins et al.,
2013). At the species level, we used the mean HI of subjects as a measure of lateralization direction
and the mean of absolute HI values (MeanAbsHI) as a measure of strength.

We applied one-sample t-tests to each species and genus sample encompassing data for at least 15
individuals to check whether HI distributions were significantly skewed at the population-level.
Additionally, the chi-square goodness-of-fit test was employed to test if the numbers of left- and
right-handers as well as ambipreferent individuals differed from a baseline distribution. Earlier
studies performed the goodness-of-fit test against the null hypothesis of a chance distribution of the
three hand preference categories (Vauclair et al., 2005). Due to our large multispecies sample, we
could adopt a different approach: For each of the three major clades studied (Cercopithecoidea,
Hominoidea, and Platyrrhini), we calculated the mean frequencies of individuals being either
ambipreferent or handed. We then assumed an equal probability of handed individuals to be either
right or left-handed and this way calculated a clade-specific baseline. Distributions for each species
were then compared to this clade-specific average. For the Hominoidea, we excluded humans for the
calculation of the baseline, because their evidently extreme lateralization bias would have otherwise
skewed the results. Bonferroni correction was employed to address multiple testing.

We employed the phytools package version 0.7-70 (Revell, 2012) to visualize evolutionary patterns,
quantify phylogenetic signals (by employing Pagel's λ – Freckleton et al., 2002; the null hypothesis of
no phylogenetic signal was tested by means of the likelihood ratio test) and to calculate maximum
likelihood ancestral state estimates (Suppl. Table 3), each separately for direction and strength of
lateralization. Time-calibrated primate phylogenies were derived from the 10kTrees website (Arnold
et al., 2010). Three species in our study were not included in the respective database, *Ateles*
hybridus, *Sapajus flavius*, and *Trachypithecus hatinhensis*. We therefore replaced *A. hybridus* and *T.*
hatinhensis in the tree with respective sister taxa, namely *A. belzebuth* and *T. francoisi* (Morales-
Jimenez et al., 2015; Roos et al., 2019), for which data were provided. This way the topology and
branch lengths of the tree could be kept equal to a model which would have included the actual
species we studied. The same could not be done for *S. flavius*, so that it was amended manually in
the respective trees by relying on divergence dating from Lima et al. (2018).

We computed phylogenetic generalized least squares (PGLS) regression models to test the effects of
different biological variables on hand preferences while addressing phylogeny (correlation structure:
Pagel's λ; model fit: maximum likelihood). The R packages *ape* (Paradis, 2019), *nlme* (Pinheiro, 2020),
and *MuMIn* (Barton, 2020) were used for model creation and evaluation. We used multi-model
inference to test how well hypothesis-derived predictors could explain HI measures on the species
level. Predictor-based models were ranked against a NULL model to estimate their explanatory
power and identify the best-performing one (*dredge* function in *MuMIn*). Second order Akaike's
information criteria (AICc) and respective Akaike weights were used to evaluate model components.
The normality of model residuals was checked by applying the Shapiro-Wilk test. We relied on the
conditional model average to assess effects of individual predictors. We selected the following
variables as model predictors to address established hypotheses on the evolution on primate hand
preferences: Ecology (terrestrial vs. arboreal), occurrence of habitual foraging-related tool use in the
wild (binarily coded), and endocranial volume (numeric, log-transformed) of females (see Suppl.
Table 4 for predictor data and respective references). A potential multicollinearity of predictors was
checked by computing their variable inflation factors. We ran the models on the species means for HI
and AbsHI, respectively, resulting in two separate analyses for direction and strength of population-
level laterality. All species (*n* = 38) were included into the strength (AbsHI) analysis. For direction, we
only considered species for which we had at least 15 sampled individuals, resulting in a more
restricted sample (*n* = 34). Since humans are both extreme outliers in regard to absolute brain size

and hand preference direction, we decided to compute a second direction model to identify
potential biases that might derive from their inclusion. Thus, this second model on hand preference
direction encompassed a species sample of $n = 33$.

Finally, we employed Bayesian phylogenetic multilevel models by aid of the *brms* package (Bürkner,
2017) to check whether individual HI and AbsHI were influenced by age and sex (excluding unsexed
individuals; $n = 2$). Analyses were run for the complete sample as well as within the three
superordinate clades, thus resulting in a total of eight models, four for each of the two response
variables. We assumed a notable effect on lateralization patterns when the model's 95% credible
intervals of intercept and respective regression coefficients did not overlap with zero. Default priors
were used. Modelling results, including the number of chains and iterations performed are
summarized and visualized in Supplementary File 1. Due to their highly derived handedness patterns,
we again excluded humans from these analyses to avoid skewing the results. To further avoid bias,
we also removed chimpanzees from the models, since they are vastly overrepresented in our sample
(31% of total individuals and 47% of all subadults). Finally, as no age data on Tonkean macaques
(*Macaca tonkeana*) were available, this species was not featured in these analyses as well. This left
us with $n = 1,107$ individuals in the full dataset, $n = 366$ in the hominoid, $n = 459$ in the
cercopithecoid and $n = 282$ in the platyrrhine subsample.

Results

Lateralization strength (MeanAbsHI) but not direction (MeanHI) displays a strong phylogenetic signal
among anthropoids (Figure 2). Accordingly, MeanAbsHI ($\lambda = 0.87$, $p_{\text{likelihood-ratio test}} = 0.04$) varied
substantially between lineages but was largely stable within groups of closely related taxa. MeanHI
on the other hand, showed random fluctuations ($\lambda < 0.001$; $p_{\text{likelihood-ratio test}} = 1$). Species-level
distributions of hand preferences are visualized in Figure 3 and summarized in Table 2.

Ancestral hand preference strength was modelled to have been similarly moderate in the stem
lineages of hominoids (AbsHI = 0.605) and cercopithecoids (AbsHI = 0.626), while it was inferred to
be higher in the ancestral platyrrhines (AbsHI = 0.738; Fig. 2, Suppl. Tab. 3). We found hand
preference strength to be particularly weak among some species in the Papionina clade (baboons
and their relatives) and to be least expressed in the gelada (*Theropithecus gelada* – MeanAbsHI =
0.257). The strongest preferences were found in humans (MeanAbsHI = 0.943), langurs of the genus
Trachypithecus (MeanAbsHI = 0.868), and spider monkeys (*Ateles* spp. – MeanAbsHI = 0.831). In
consequence, these taxa included only very few, if any, ambipreferent individuals. The South Asian
colobines *Rhinopithecus* and *Trachypithecus* were the only genera in which no ambipreferent
individuals were found. Direction of manual lateralization was far more uniform across the
anthropoid radiation than strength (Fig. 2; Suppl. Tab. 3). No species approached the extreme
direction bias of humans, and only 2 of 37 non-human species exhibited a significant population-level
bias as indicated by the one-sample t-test after correcting for multiple testing (4 of 37 when no
correction was applied), namely gorillas and chimpanzees. Thus, such biases were restricted to the
African ape lineage. After correcting for multiple testing, intraspecific frequencies of ambipreferent
individuals, right-, and left-handers, as indicated by the chi-square goodness-of-fit test, differed
significantly from the superordinate taxon estimate in just two cases. These were humans and
chimpanzees, which are both right-handed at the population-level (Tab. 2). When omitting
correction, respective biases were notably more frequent and found in 11 non-human species across
the three major clades studied. Species- and genus-level results closely corresponded to each other.
However, different from the species-level results, the corrected chi-square goodness-of-fit test for
intrageneric patterns also found lion tamarins (*Leontopithecus*) and snub-nosed monkeys

(*Rhinopithecus*) to exhibit significantly derived handedness patterns. In the former genus,
ambipreferent individuals occur more frequently than in other platyrrhines, while the sample for the
latter completely lacked ambipreferent animals and included comparatively fewer right-handed ones
than are found among the remaining cercopithecoids.

Our selected predictors for PGLS models showed a mixed performance in explaining expression
patterns of hand preference strength and direction in anthropoids (Tab. 3, Fig. 4). For the initial
model on lateralization direction, we found a significant effect of ecology, with terrestrial species
displaying a right-hand bias compared to arboreal ones (p = 0.04; Tab. 3A). However, when humans
were removed from the model, this effect was merely recovered as a non-significant trend (p = 0.07;
Tab. 3B).. Other predictors had no significant effect on lateralization direction, regardless of whether
humans were included in the model or not (p > 0.19, Tab. 3A, B). When humans were included (Tab.
4A), models encompassing the component ecology as well as some considering brain size
outperformed the lateralization direction null model. When humans were not considered in the
analysis, the null model was solely outperformed by one that exclusively included the component
ecology, and only slightly so (Δ AICc = 1.07; Tab. 4B), indicating a notable bias derived from the
extreme right handedness found in our species. Thus, whereas habitual tool use and absolute brain
size clearly do not influence the direction of lateralization among anthropoids in general, the
analyses provide evidence for a weak but detectable effect of ecology. Such ecological signatures
were found to be of greater relevance for patterns of lateralization strength. Here, a significant
negative effect of a terrestrial lifestyle was found (p = 0.038; see Tab. 3C). In line with that, models
including the component ecology consistently outperformed the null model, which was not the case
for those including only tool use and/or brain size (Tab. 4C). Thus, terrestrial anthropoids show
weaker hand preferences than arboreal ones while there is no correlation with brain size or habitual
tool use for this trait.

At the individual level, Bayesian models showed that neither age, nor sex had an influence on
lateralization direction when the total sample was concerned (the respective credible intervals
overlapped with zero; Suppl. File 1). However, we recovered a notable effect of age on lateralization
direction the hominoid subsample exclusively, with subadults tending more towards left-handedness
than adults (credible interval = -0.28 – -0.01; Suppl. File 1). We observed a different pattern for
lateralization strength. Here, an effect of age but again not sex was recovered for both the total
sample (credible interval = -0.11 – -0.04) and the hominoid subsample (credible interval = -0.16 – -
0.02), with weaker lateralization in subadults compared to adults. When examined separately, this
effect was not detectable in neither platyrrhines nor cecopithecoids this effect was not detectable
(credible intervals overlapping with zero; Suppl. File 1).

Figure 2: A color-coded phylogeny of hand preferences in anthropoid primates. The strength (A) and direction (B) of laterality, expressed by the mean absolute handedness index (MeanAbsHI) and the mean handedness index (MeanHI; 0 is marked by the black bar on the color scale), respectively, calculated for each species and inferred for each tree node by maximum likelihood estimates. Silhouettes by Kai R. Caspar, except *Ateles* (by Yan Wong, public domain) and *Homo* (public domain)

Figure 3: Violin plots of hand preference distribution in 22 genera of anthropoid primates and the genus-specific expression of three potential biological correlates (ecology, habitual foraging-related tool use, and absolute brain size). Attributions only apply to the species that represent the respective genus within our sample. Color coding: Ecology – green: arboreal, yellow: terrestrial; Habitual tool use – grey: present; white: absent. Brain size is visualized here as the log-transformed genus average of female endocranial volume.

401   **Table 2**: Hand preferences of anthropoid species as recovered by the tube task. * Results remain significant after Bonferroni correction.

Species	*n*	# left (%)	# right (%)	# ambipreferent (%)	MeanHI	MeanAbsHI	Species direction bias (HI), *p* value	Species L/R/A distribution, *p* value	*n*Genus	Genus direction bias (HI), *p* value	Genus L/R/A distribution, *p* value
Ateles fusciceps	46	20 (43.5)	22 (47.8)	4 (8.7)	0.063	0.798	0.618	0.389	87	0.759	0.060
Ateles geoffroyi	23	10 (43.5)	11 (47.8)	2 (8.7)	0.061	0.829	0.748	0.624			
Ateles hybridus	18	13 (72.2)	5 (27.8)	0	-0.377	0.917	0.086	**0.022**			
Cercocebus torquatus	31	13 (41.9)	11 (35.5)	7 (22.6)	-0.029	0.665	0.832	0.915	31	0.832	0.915
Cercopithecus diana	20	7 (35)	9 (45)	4 (20)	0.178	0.755	0.339	0.865	45	0.572	0.125
Cercopithecus neglectus	25	17 (68)	6 (24)	2 (8)	-0.258	0.621	0.061	**0.011**			
Gorilla gorilla	76	17 (22.4)	41 (53.9)	18 (23.7)	0.248	0.541	**< 0.001***	**0.007**	76	**< 0.001***	**0.007**
Homo sapiens	127	12 (9.5)	111 (87.4)	4 (3.1)	0.761	0.943	**< 0.001***	**< 0.001***	127	**< 0.001***	**< 0.001***
Hylobates lar	36	17 (47.2)	16 (44.5)	3 (8.3)	-0.011	0.614	0.924	0.182	58	0.612	0.125
Hylobates moloch	22	11 (50)	8 (36.4)	3 (13.6)	-0.115	0.799	0.540	0.552			
Leontopithecus chrysomelas	30	11 (36.7)	11 (36.7)	8 (26.6)	0.151	0.514	0.171	0.273	73	0.241	**0.001***
Leontopithecus chrysopygus	15	4 (26.7)	5 (33.3)	6 (40)	0.039	0.350	0.744	**0.037**			
Leontopithecus rosalia	28	11 (39.3)	8 (28.6)	9 (32.1)	0.022	0.502	0.850	0.052			
Macaca fascicularis	20	9 (45)	9 (45)	2 (10)	-0.036	0.684	0.835	0.445	102	0.692	0.785
Macaca nemestrina	29	10 (34.5)	11 (37.9)	8 (27.6)	0.035	0.527	0.768	0.730			
Macaca silenus	35	16 (45.7)	11 (31.4)	8 (22.9)	-0.051	0.467	0.596	0.625			
Macaca sylvanus	24	10 (41.7)	10 (41.7)	4 (16.6)	-0.025	0.670	0.873	0.835			
Macaca tonkeana	14	5 (35.7)	3 (21.4)	6 (42.9)	-0.057	0.543	0.753	0.133			
Mandrillus sphinx	32	7 (21.9)	11 (34.4)	14 (43.7)	0.034	0.389	0.701	**0.008**	32	0.701	**0.008**
Nomascus gabriellae	10	5 (50)	2 (20)	3 (30)	-0.173	0.618	0.465	0.436	36	0.593	0.805
Nomascus leucogenys	26	9 (34.6)	11 (42.3)	6 (23.1)	-0.031	0.555	0.818	0.869			
Pan paniscus	118	50 (42.4)	51 (43.2)	17 (14.4)	0.044	0.529	0.431	0.237	654	**< 0.001***	**< 0.001***
Pan troglodytes	536	155 (28.9)	266 (49.6)	115 (21.5)	0.133	0.507	**< 0.001***	**< 0.001***			
Papio anubis	84	27 (32.1)	41 (48.8)	16 (19.1)	0.108	0.527	0.102	0.189	108	NA	0.218
Papio hamadryas	24	7 (29.2)	9 (37.5)	8 (33.4)	0.066	0.408	0.533	0.348			

Pithecia pithecia	7	4 (57.1)	1 (14.3)	2 (28.6)	-0.385	0.934	0.312	0.306	7	NA	0.306
Pongo sp.	47	27 (57.5)	9 (19.1)	11 (23.4)	-0.225	0.487	**0.006**	**0.012**	47	**0.006**	**0.012**
Pygathrix cinerea	18	6 (33.3)	9 (50)	3 (16.7)	0.165	0.499	0.268	0.634	18	0.268	0.634
Rhinopithecus roxellana	24	17 (70.8)	7 (29.2)	0	-0.319	0.729	**0.040**	**0.002**	24	**0.040**	**0.002***
Saimiri sciureus	36	21 (58.4)	14 (38.9)	1 (2.7)	-0.119	0.757	0.382	**0.044**	36	0.382	**0.044**
Sapajus apella	25	11 (44)	10 (40)	4 (16)	0.016	0.651	0.917	0.976	80	0.922	0.932
Sapajus flavius	21	9 (42.9)	8 (38.1)	4 (19)	-0.130	0.769	0.495	0.900			
Sapajus xanthosternos	34	15 (44.1)	14 (41.2)	5 (14.7)	0.089	0.677	0.492	0.965			
Semnopithecus entellus	30	18 (60%)	7 (23.4)	5 (16.6)	-0.184	0.560	0.110	0.061	30	0.110	0.061
Symphalangus syndactylus	31	12 (38.7)	8 (25.8)	11 (35.5)	-0.048	0.482	0.663	0.093	31	0.663	0.093
Theropithecus gelada	38	11 (28.95)	11 (28.95)	16 (42.1)	0.053	0.257	0.326	**0.010**	38	0.326	**0.010**
Trachypithecus auratus	8	6 (75)	2 (25)	0	-0.256	0.984	0.499	0.092	26	0.153	0.006
Trachypithecus hatinhensis	18	11 (61.1)	7 (38.9)	0	-0.248	0.817	0.219	**0.047**			

Figure 4: Visualization of PGLS coefficient estimates (including 95% confidence intervals) for the
influence of brain size, tool use, and ecology on lateralization direction (A, B) as well as strength (C) in
anthropoid primates. Two models for lateralization direction were computed, one including (A) the
other one excluding humans (B).

Table 3: Conditional average of PGLS model coefficients for lateralization strength and direction in
anthropoid primate species. VIF = variable inflation factor.

A: Conditional PGLS model average for lateralization direction, including humans				
Predictor	**Estimate**	**Std. Error**	**VIF**	***p* value**
Ecology (Terrestrial lifestyle)	0.152	0.072	1.798	0.041
Tool use (present)	0.111	0.081	1.164	0.191
Log$_{10}$ brain size	0.050	0.043	1.978	0.263
B: Conditional PGLS model average for lateralization direction, excluding humans				
Predictor	**Estimate**	**Std. Error**	**VIF**	***p* value**
Ecology (Terrestrial lifestyle)	0.11	0.057	1.419	0.072
Log$_{10}$ brain size	-0.020	0.037	1.405	0.603
Tool use (present)	0.011	0.069	1.098	0.882
C: Conditional PGLS model average for lateralization strength				
Predictor	**Estimate**	**Std. Error**	**VIF**	***p* value**
Ecology (Terrestrial lifestyle)	-0.143	0.067	1.499	**0.038**
Tool use (present)	0.054	0.070	1.164	0.459
Log$_{10}$ brain size	0.034	0.047	1.612	0.476

415 **Table 4:** Results of PGLS model averaging for lateralization direction (considering the inclusion and
416 exclusion of humans) and strength. Null models are shown in italics. Df. = Degrees of freedom. AICc =
417 Second order Akaike information criterion.

A: PGLS model for lateralization direction, including humans				
Components	**Df.**	**AICc**	**Δ AICc**	**Weight**
Ecology, tool use	4	-13.50	0	0.26
Ecology	3	-13.47	0.03	0.26
Brain size	3	-12.16	1.33	0.14

Ecology, brain size	4	-11.93	1.57	0.12
Ecology, tool use, brain size	5	-10.96	2.54	0.07
(NULL)	*2*	*-10.39*	*3.11*	*0.06*
Tool use, brain size	4	-10.28	3.21	0.05
Tool use	3	-9.70	3.80	0.04

B: PGLS model for lateralization direction, excluding humans

Components	Df.	AICc	Δ	Weight
Ecology	*3*	*-30.37*	*0*	*0.36*
(NULL)	*2*	*-29.30*	*1.07*	*0.21*
Ecology, brain size	4	-28.59	1.78	0.15
Ecology, tool use	4	-27.83	2.54	0.10
Brain size	3	-26.90	3.48	0.06
Tool use	3	-26.88	3.49	0.06
Ecology, tool use, brain size	5	-26.04	4.34	0.04
Tool use, brain size	4	-24.31	6.07	0.02

C: PGLS model for lateralization strength

Components	Df.	AICc	Δ AICc	Weight
Ecology	3	-9.88	0	0.37
Ecology, brain size	4	-8.79	1.09	0.21
Ecology, tool use	4	-8.16	1.71	0.16
(NULL)	*2*	*-7.28*	*2.60*	*0.10*
Ecology, brain size, tool use	5	-6.31	3.57	0.06
Tool use	3	-5.80	4.07	0.05
Brain size	3	-5.13	4.75	0.03
Tool use, brain size	4	-3.93	5.94	0.02

Discussion

General

Our study provides the first quantitative phylogenetic perspectives on hand preferences in monkeys, apes, and humans. While population-level lateralization strength is highly varied among anthropoid primates and often distinctly expressed in specific lineages, direction fluctuates irrespective of phylogeny and appears comparatively uniform. Indeed, significant population-level biases are notably rare. After expanding the sample size for some species in which such biases haven been previously reported based on the tube task (siamang – Morino et al., 2017; de Brazza's monkey – Schweitzer et al., 2007), we failed to replicate significant deviations from a chance distribution (even if not correcting for multiple testing). The only taxon in which significant hand use biases for bimanual manipulation occur frequently, is constituted by the great apes and humans. Nevertheless, since sample sizes for species in this group are consistently and conspicuously large, statistical analyses performed on them (in particular the frequently applied one-sample t-test) will have higher

power compared to tests done on taxa represented by a smaller number of individuals. It is therefore
premature to assume that hominids display qualitatively different population-level lateralization
patterns than other primates. Looking at the population-level frequencies of right-handed, left-
handed, and ambipreferent individuals, cercopithecoids such as the golden snub-nosed monkey (70.8
% left-handers, 0 % ambipreferents, MeanHI: -0.319) and de Brazza's monkey (68 % left-handers, 8 %
ambipreferents, MeanHI: -0.258) approach a human-like skew more than any great ape species does,
albeit in the contrary direction to lateralization in our species (approximated herein as encompassing
87.3% right-handers, 3.1 % ambipreferents, MeanHI: 0.761). Whether the hand preference patterns
recovered for these monkeys are indeed representative needs to be clarified in future studies, which
should expand the available samples. The finding that subadult non-human hominoids, but not other
anthropoids, are notably stronger left-handed than adults, likely results from the fact that juveniles
are more weakly lateralized overall (see below) and from the composition of our Bayesian model
sample. Therein, Western gorillas and bonobos, both of which tend to be right-handed as adults
(Hopkins et al., 2011), are well-represented across age groups. Again, we suggest that an expanded
and more balanced dataset could level out the lateralization differences between (great) apes and
other anthropoids.

The fact that population-level hand preference fluctuates without phylogenetic and only marginal
ecological signatures among anthropoids, suggests that there are no strong directional selective
pressures acting on this trait, different from what pertaining hypotheses predict (see below). On the
other hand, population-level lateralization strength is more variable but consistent among closely
related taxa, thus exhibiting a strong phylogenetic signal. Our finding that hand preference strength
is generally weaker in juveniles compared to adults replicates results from several studies relying on
smaller sample sizes (e.g., Westergaard & Suomi, 1996; Zhao et al., 2012). PGLS modelling
demonstrated a significant negative effect of a terrestrial lifestyle on hand preference strength,
indicating an influence of ecology. It appears intuitive that terrestrial taxa tend to be less lateralized
than arboreal ones, since the latter often need to flexibly stabilize their body in the canopy, for
instance while foraging. Accordingly, one hand will be preferably used to provide such support, but
whether the left or the right one adopts this function seems to be arbitrary. The fact that these
lateralization tendencies were found in zoo-housed primates that often adopt locomotor regimes
very different from their wild conspecifics (e.g., captive spider monkeys and gibbons spend
considerable amounts of time moving and feeding on the ground), suggest a significant innate
component to these patterns. However, within ecologically uniform groups, the variability of hand
preference strength can still be notable, at times even among closely related taxa (compare e.g.,
Javan gibbon and siamang), pointing at yet unidentified biological influences being at play. Given the
great variability of lateralization strength among anthropoids and its ties to phylogeny as well as
ecology, this aspect of manual lateralization and its evolution should receive more research attention
in the future. In the past, most work and evolutionary considerations regarding primate handedness
have instead focused on lateralization direction, surely for anthropocentric reasons. As we attempt
to show here, however, the explanatory power of these in parts very long-lived conjectures appears
to be limited.

Testing prevalent hypotheses

Our data does not unambiguously support any of the tested hypotheses on hand preference
evolution in primates. The traditional postural origins hypothesis (POH) assumes right-hand
tendencies for manipulation across anthropoid taxa. However, we found that anthropoid population-
level lateralization is in most cases not notably shifted into either direction, with a slight majority of
species displaying a weak left-hand bias (20 of 38 species). It is important to note that the recovered
correlation between arboreality and hand preference strength does not support any version of the

POH, as they focus exclusively on lateralization direction. The novel POH assumes that terrestrial
non-human primates tend to be right-handed, while arboreal ones tend to be left-handed, a claim
only marginally supported by our data. While the mean HIs of terrestrial and arboreal non-human
species are indeed dissimilar, they do not deviate strongly from zero and there is a notable overlap in
variation between the two ecological groups (MeanHI$_{arboreal}$: -0.08, SD: 0.16; MeanHI$_{terrestrial}$: 0.04, SD:
0.10). Thus, the predictive power of the hypothesis is comparatively weak. Admittedly, improved
sampling, both within and across species might consolidate this pattern in the future, so that we do
not want to dismiss the POH at this point. However, even if its predictions should become validated
eventually, the hypothesis does still not provide an evolutionary mechanism for why population-level
hand preference patterns should be coupled to positional behavior. This represents a considerable
conceptual shortcoming that needs to be addressed in the future.

Furthermore, the evolutionary scenario proposed by both versions of the POH is outdated and
should not be perpetuated without explicitly stating its shortcomings. According to the POH, small-
bodied bushbabies (genus *Galago*) are suitable models for early primates, since they would be "the
most direct descendants of the earliest forms" (MacNeilage, 2007). Because contemporary studies
suggested a left-hand bias for prey grasping in bushbabies, such a pattern was also assumed for
primate ancestors in the paper that introduced the original POH (MacNeilage et al., 1987). However,
these assumptions have always been speculative and become problematic in light of more recent
data. First, there is no convincing evidence for preferably left-handed grasping in the genus *Galago,*
or other galagids, anymore (Papademetriou et al., 2005). Second, bushbabies represent a remarkably
derived radiation of strepsirrhines (Tab Rasmussen & Nekaris, 1998) and are thus no suitable
ecological models for the common ancestor of modern primates. Current evidence suggests that
both the earliest primates and the ancestors of the anthropoid clade studied herein, were
omnivorous arboreal quadrupeds with moderate leaping ability (Silcox et al., 2009; Gebo, 2011;
Sussman, Tab Rasmussen, & Raven, 2013) and possibly diurnal habits (Tan et al., 2005; Ankel-Simons
& Rasmussen, 2008). Thus, they were extremely different from extant galagos. Our results suggest
that the common ancestor of anthropoids did not display a notable population-level hand preference
for manipulation, nor that such biases are common among extant monkeys and apes. While lemurs
(different from galagos) indeed appear to show a consistent left-hand preference for unimanual
reaching, this pattern is not recovered in bimanual tasks (Papademetriou et al., 2005; Regaiolli et al.,
2016; Batist & Mayhew, 2020), again contradicting the POH (MacNeilage, 2007). In conclusion, the
unsupported evolutionary scenario proposed by the POH and its rather fragile empirical basis lead us
to challenge its status as a keystone idea in the current discourse on the evolution of primate manual
lateralization.

We also found no effects of foraging-related tool use on neither direction, nor strength of
lateralization although our sample represented all primate lineages that include habitual tool users
(Musgrave & Sanz, 2018). Surprisingly, we also did not recover notable influences of absolute brain
size on hand preferences. An effect on lateralization strength was expected both on theoretical
considerations (Ringo et al., 1994) and empirical evidence from studies investigating intra- and
interspecific covariation of brain size and overall cortical lateralization (Kong et al., 2018; Ardesch et
al., 2021). Why do anthropoid hand preferences not conform to these predictions? We cannot
provide a satisfying answer to this question. It is possible that the effects of increased overall brain
lateralization on hand preference expression turn out to be unexpectedly weak and are masked by
yet unidentified neurologic factors.

All in all, none of the hypotheses on primate handedness that we addressed were corroborated by
our data. Nevertheless, when discussing conflicts between prevalent ideas and our results, we also
need to address general limitations of our approach. For instance, we equate general hand

preferences with tube task results. Although the tube task represents one of the best available
behavioral assays for brain lateralization (Dadda et al., 2006; Margiotoudi et al., 2019), it is obvious
that other hand use situations, such as communicative gesturing, need to be considered to arrive at a
holistic understanding of primate hand use evolution (Hopkins et al., 2013; Becker et al., 2022). Such
an approach could also test how variable hand use consistency across contexts is among primates
and whether ecological variables have an influence here as well. At the moment, it appears as if
there is comparatively little consistency in manual preferences across different hand use situations in
non-human primates (Laska, 1996; Lilak & Phillips, 2008; Marchant & McGrew, 2013; Caspar et al.,
2018), but most studies so far compare tasks of varying complexity in a few model species and cases
of consistent hand use ("true handedness") have indeed been reported (Diamond & McGrew, 1994;
Hopkins et al., 2013; but note neuroanatomical evidence for a dissociation of lateralization in non-
communicative actions and gestures in baboons – Becker et al., 2022).

Another limitation is posed by our sample composition. Both the number of species and subjects per
species need to be increased to verify the patterns communicated here. In particular, additional
sampling of the speciose New World monkey families Pitheciidae and Callitrichidae, tarsiers, and
strepsirrhines would be desirable, to make inferences more robust. This is especially the case for
predictions about hand preference patterns in early crown-group primates. Our experience suggests
that at least pitheciids and lemurs only reluctantly engage in the tube task so that it might be
advisable to apply different bimanual testing schemes in these groups. In lemurs, puzzle boxes have
been employed as such: The animals open the lid of a box with one hand while the other one is
retrieving food stored within (Regaiolli et al., 2016; Batist & Mayhew, 2020). Future studies need to
check the functional equivalence of this approach with the tube task (which is not a trivial question,
compare e.g., Lilak and Phillips, 2008) to establish a set of behavioral assays that could be employed
to study hand preferences in the whole primate order. These methods might then also be applied to
other dexterous and ecologically variable mammalian groups, such as musteloid carnivorans
(Kitchener, 2017), to test hypotheses on the evolution of manual laterality across a wider
phylogenetic margin. Finally, we need to acknowledge the limitations of our phylogenetic modelling
approach. In particular, the binary coding of *ecology* (arboreal vs terrestrial) obviously simplifies the
remarkable spectrum of positional behaviors found among anthropoids. Future approaches might
explore alternative strategies to statistically code this multifaceted variable.

The evolutionary issue of human handedness

In line with previous research, we found human right-handedness to be unparalleled among
primates. We want to stress, however, that humans only deviate markedly from all other taxa in
direction and not in strength of lateralization for bimanual manipulation. When it comes to the
latter, the human condition is approached in groups such as leaf monkeys and spider monkeys.
Perhaps surprisingly, handedness strength in great apes is modest in comparison (Table 2). Still,
humans are highly deviant among predominately terrestrial primates in displaying such strong
individual hand preferences. Whether this difference relates to bipedal locomotion, which has often
been championed as a correlate of human handedness (Westergaard et al., 1998; Cashmore et al.,
2008; Prieur et al., 2019;), is open for debate. Since no other extant primate shows similar
adaptations to terrestrial bipedalism, the validity of this assumption is hard to test in the framework
of comparative approaches (but see Giljov et al., 2015). Quadrupedal primates tend to exhibit
stronger hand preferences when adopting the relatively instable bipedal posture (Westergaard et al.,
1998). Whether this finding has any evolutionary implications remains unclear and it should be
emphasized that although humans are bipeds, a high percentage of complex manual actions,
including numerous examples of bimanual manipulation and tool use, are not (and never have been)

habitually performed in a bipedal posture. In any case, our results suggest that bipedalism is at least
not a prerequisite to evolve strong hand preferences in anthropoid primates.

When turning to lateralization direction, the statement of Corballis (1987) remains valid: some non-
primate vertebrates approach humans more closely in population-level handedness than their simian
relatives do. Apart from humans, extreme forms of vertebrate limb use biases are known from
parrots (Kaplan & Rogers, 2021) and ground-living kangaroos (Giljov et al., 2015). Why these very
different groups converge in this respect remains enigmatic. So why do humans stand out among the
primate order when it comes to handedness direction? The limited insights gained by comparative
behavioral studies, including this one, do indeed suggest that the extreme right-handedness of
humans is a trait that evolved due to unique neurophysiological demands not experienced by other
primates. Frost (1980) already pointed out that humans' outstanding proficiency in tool use and
manufacture should be considered a significant influence on handedness evolution. Thus, not
foraging-related tool use per se, but the unique way in which it became immersed into complex
human behaviors might have influenced overall brain lateralization in our lineage. In line with that,
areas of the prefrontal cortex involved in motor cognition are structurally derived in humans and
differ significantly from their homologs in apes and monkeys (Hecht et al., 2015a; Barrett et al.,
2020). However, specializations of both the right and the left hemisphere are determining human-
specific tool use proficiency and motor planning, apparently with particular involvement of the right
inferior frontal gyrus (Ramayya et al., 2010; Hecht et al., 2015a; Hecht et al., 2015b). Postulating that
hominin tool use and right-handedness evolved in tandem is therefore not straight-forward.

Besides that, there is of course the notion of coevolution between language and handedness, which
might explain human-specific patterns of population-level manual lateralization. For this hypothesis
to be convincing, the development and function of neural substrates controlling vocal behavior and
those regulating manual motor control would need to be uniquely intertwined in humans. Indeed,
the connectivity of the arcuate fasciculus, a tract critically involved in language processing and
production, is highly derived in humans, suggesting important qualitative differences to other species
(Rilling et al., 2008; but see Barrett et al., 2020 for other elements relevant for language production
which are conserved across catarrhine primates). Nevertheless, how such traits could functionally
relate to population-level handedness remains totally unclear. In fact, despite the popularity of the
idea, a link between handedness and language processing that goes beyond superficial left-
hemisphere collateralization in right handers (not even in the majority of left-handers) is far from
evident (Fitch & Braccini, 2013). To defend an evolutionary connection between these phenomena,
pleiotropic or otherwise functionally linked genes influencing the development of both language
areas and those related to handedness would need to be identified. So far, this has not been
accomplished and current evidence suggests that language and handedness are largely independent
on various structural levels (Ocklenburg et al., 2014; Schmitz et al., 2017b). Hence, despite the
continuing efforts to unravel the evolution of human right-handedness, including the ones made by
us herein, it remains an essentially unsolved issue of human cognitive evolution.

Conclusions

We recover highly variable patterns of hand preference strength in anthropoid primates, which
correlate with ecology and phylogeny. In contrast to this, no phylogenetic or ecological signal was
found for lateralization direction, and few species exhibit significant population-level hand
preferences. We tested three pertaining conjectures on primate handedness evolution, the POH, tool
use and brain size hypotheses, but none were corroborated by our data. Novel ideas on the evolution
of primate hand preference should put a stronger focus on manual lateralization strength rather than
direction to address the phylogenetic patterns described herein. However, additional datasets on
primates and potentially non-primate mammals are needed to robustly inform respective

hypotheses. By relying on standardized testing paradigms, such as the tube task, researchers can effectively build on our as well as others results and expand multispecies datasets for further comparative phylogenetic studies. Although we are convinced that such approaches could significantly improve our understanding of general trends in the evolution of primate hand preferences, the unusual autapomorphic handedness pattern of humans will very likely require explanations that cannot be derived from such comparative behavioral studies. The evolutionary underpinnings of handedness expression in our species remain enigmatic.

Acknowledgements

We would like to thank the curators and animal keepers at the following institutions for the permission to study their animals and for their support during data collection:

Allwetterzoo Münster, Apenheul (Apeldoorn), Artis Amsterdam, Bali Wildlife Rescue Center, Bioparc de Doué-la-Fontaine, Burger's Zoo (Arnhem), Erlebnis-Zoo Hannover, Howletts Wild Animal Park (Bekesbourne), Naturzoo Rheine, Parc Zoologique et Botanique de Mulhouse, Port Lympne Wild Animal Park (Lympne), Serengeti-Park Hodenhagen, Tierpark Aachen, Tierpark Berlin, Tierpark Hamm, Tierpark Hellabrunn (Munich), Tierpark Mönchengladbach, Tierpark Ulm, Wilhelma Stuttgart, Zoo Basel, Zoo Berlin, Zoo Dortmund, Zoo Dresden, Zoo Duisburg, Zoo Frankfurt, Zoo Heidelberg, Zoo Köln (Cologne), Zoo Krefeld, Zoo Landau in der Pfalz, Zoo Liberec, Zoo Magdeburg, Zoo Neuwied, Zoo Osnabrück, Zoo Wuppertal, Zoológico de São Paulo, Zoologischer Stadtgarten Karlsruhe, ZOOM Erlebniswelt Gelsenkirchen, ZooParc Overloon, and Zoopark Erfurt.

We are also indebted to Hélène Cochet, Julia Fischer, William D. Hopkins, Adrien Meguerditchian, and Luca Morino for sharing raw data on bimanual task preferences in humans, barbary macaques, great apes, olive baboons, and gibbons, respectively.

Daniel Issel is greatly acknowledged for collecting data on pig-tailed macaques and Javan langurs in Bali and Petra Bolechová for doing so for Diana monkeys at Zoo Liberec. Finally, we would like to thank Miriam Lindenmeier and Larissa Günther for crucial assistance during data collection at various locations and Alex DeCasien as well as Adrien Meguerditchian for their constructive and insightful reviews on earlier versions of this manuscript.

This research was funded by the German Society for Mammalian Biology (Deutsche Gesellschaft für Säugetierkunde). KRC was supported by a Ph.D. fellowship of the German Academic Scholarship Foundation (Studienstiftung des deutschen Volkes e.V.).

References

Amunts, K., Schlaug, G., Schleicher, A., Steinmetz, H., Dabringhaus, A., Roland, P. E., & Zilles, K. (1996). Asymmetry in the human motor cortex and handedness. *NeuroImage, 4*(3), 216-222. doi:https://doi.org/10.1006/nimg.1996.0073

Ankel-Simons, F., & Rasmussen, D. T. (2008). Diurnality, nocturnality, and the evolution of primate visual systems. *American Journal of Physical Anthropology, 137*(S47), 100-117. doi:https://doi.org/10.1002/ajpa.20957

Annett, M. (2002). *Handedness and Brain Asymmetry: The Right Shift Theory*. Hove: Psychology Press.

Ardesch, D. J., Scholtens, L. H., de Lange, S. C., Roumazeilles, L., Khrapitchev, A. A., Preuss, T. M., . . . van den Heuvel, M. P. (2021). Scaling principles of white matter connectivity in the human and nonhuman primate brain. *Cerebral Cortex*, bhab384. doi:10.1093/cercor/bhab384

Arnold, C., Matthews, L. J., & Nunn, C. L. (2010). The 10kTrees website: A new online resource for primate phylogeny. *Evolutionary Anthropology: Issues, News, and Reviews, 19*(3), 114-118. doi:https://doi.org/10.1002/evan.20251

Barrett, R. L. C., Dawson, M., Dyrby, T. B., Krug, K., Ptito, M., Arceuil, H., . . . Catani, M. (2020). Differences in frontal network anatomy across primate species. *The Journal of Neuroscience, 40*(10), 2094. doi:10.1523/JNEUROSCI.1650-18.2019

Barton, K. (2020). MuMIn: Multi-model inference. R package version 1.43.17. Retrieved from https://CRAN.R-project.org/package=MuMIn

Batist, C. H., & Mayhew, J. A. (2020). Lateralization in seven lemur species when presented with a novel cognitive task. *American Journal of Physical Anthropology, 172*(2), 270-279. doi:https://doi.org/10.1002/ajpa.24037

Becker, Y., Claidière, N., Margiotoudi, K., Marie, D., Roth, M., Nazarian, B., Anton, J. L., Coulon, O., & Meguerditchian, A. (2022). Broca's cerebral asymmetry reflects gestural communication's lateralisation in monkeys (*Papio anubis*). *eLife, 11*, e70521. https://doi.org/10.7554/eLife.70521

Blois-Heulin, C., Bernard, V., & Bec, P. (2007). Postural effect on manual laterality in different tasks in captive grey-cheeked mangabey (*Lophocebus albigena*). *Journal of Comparative Psychology, 121*(2), 205-213. doi:10.1037/0735-7036.121.2.205

Brown, C., & Magat, M. (2011). The evolution of lateralized foot use in parrots: a phylogenetic approach. *Behavioral Ecology, 22*(6), 1201-1208. doi:10.1093/beheco/arr114

Bürkner, P. (2017). brms: an R package for Bayesian multilevel models using stan. *Journal of Statistical Software*, 80(1), 1–28. doi: 10.18637/jss.v080.i01.

Canteloup, C., Vauclair, J., & Meunier, H. (2013). Hand preferences on unimanual and bimanual tasks in Tonkean macaques (*Macaca tonkeana*). *American Journal of Physical Anthropology, 152*(3), 315-321. doi:https://doi.org/10.1002/ajpa.22342

Cashmore, L., Uomini, N., & Chapelain, A. (2008). The evolution of handedness in humans and great apes: a review and current issues. *Journal of Anthropological Sciences 86*, 7-35.

Caspar, K. R., Mader, L., Pallasdies, F., Lindenmeier, M., & Begall, S. (2018). Captive gibbons (Hylobatidae) use different referential cues in an object-choice task: insights into lesser ape cognition and manual laterality. *PeerJ, 6*, e5348. doi:10.7717/peerj.5348

Chatagny, P., Badoud, S., Kaeser, M., Gindrat, A.-D., Savidan, J., Fregosi, M., . . . Rouiller, E. M. (2013). Distinction between hand dominance and hand preference in primates: a behavioral investigation of manual dexterity in nonhuman primates (macaques) and human subjects. *Brain and Behavior, 3*(5), 575-595. doi:https://doi.org/10.1002/brb3.160

Cochet, H., & Vauclair, J. (2012). Hand preferences in human adults: non-communicative actions versus communicative gestures. *Cortex, 48*(8), 1017-1026. doi:https://doi.org/10.1016/j.cortex.2011.03.016

Corballis, M. C. (1987). Straw monkeys. *Behavioral and Brain Sciences, 10*(2), 269-270. doi:10.1017/S0140525X00047762

Corballis, M. C. (1991). *The Lopsided Ape: Evolution of the Generative Mind*. Oxford: Oxford University Press.

Coren, S., & Porac, C. (1977). Fifty centuries of right-handedness: the historical record. *Science, 198*(4317), 631. doi:10.1126/science.335510

Cubí, M., & Llorente, M. (2021). Hand preference for a bimanual coordinated task in captive hatinh langurs (*Trachypithecus hatinhensis*) and grey-shanked douc langurs (*Pygathrix cinerea*). *Behavioural Processes, 187*, 104393. doi:https://doi.org/10.1016/j.beproc.2021.104393

Dadda, M., Cantalupo, C., & Hopkins, W. D. (2006). Further evidence of an association between handedness and neuroanatomical asymmetries in the primary motor cortex of chimpanzees (*Pan troglodytes*). *Neuropsychologia, 44*(12), 2582-2586. doi:https://doi.org/10.1016/j.neuropsychologia.2006.03.037

de Andrade, A. C., & de Sousa, A. B. (2018). Hand preferences and differences in extractive foraging in seven capuchin monkey species. *American Journal of Primatology, 80*(8), e22901. doi:https://doi.org/10.1002/ajp.22901

715   Diamond, A. C., & McGrew, W. C. (1994). True handedness in the cotton-top tamarin (*Saguinus*
716   *oedipus*)? *Primates, 35*(1), 69-77. doi:10.1007/BF02381487
717   Fan, P., Liu, C., Chen, H., Liu, X., Zhao, D., Zhang, J., & Liu, D. (2017). Preliminary study on hand
718   preference in captive northern white-cheeked gibbons (*Nomascus leucogenys*). *Primates,*
719   *58*(1), 75-82. doi:10.1007/s10329-016-0573-8
720   Faurie, C., Schiefenhövel, W., le Bomin, S., Billiard, S., & Raymond, M. (2005). Variation in the
721   frequency of left-handedness in traditional societies. *Current Anthropology, 46*(1), 142-147.
722   doi:10.1086/427101
723   Fitch, W. T., & Braccini, S. N. (2013). Primate laterality and the biology and evolution of human
724   handedness: a review and synthesis. *Annals of the New York Academy of Sciences, 1288*(1),
725   70-85. doi:https://doi.org/10.1111/nyas.12071
726   Freckleton, R. P., Harvey, P. H., & Pagel, M. (2002). Phylogenetic analysis and comparative data: a
727   test and review of evidence. *The American Naturalist, 160*(6), 712-726. doi:10.1086/343873
728   Frost, G. (1980). Tool behavior and the origins of laterality. *Journal of Human Evolution, 9*, 447-459.
729   Gebo, D. L. (2011). Vertical clinging and leaping revisited: vertical support use as the ancestral
730   condition of strepsirrhine primates. *American Journal of Physical Anthropology, 146*(3), 323-
731   335. doi:https://doi.org/10.1002/ajpa.21595
732   Giljov, A., Karenina, K., Ingram, J., & Malashichev, Y. (2015). Parallel emergence of true handedness
733   in the evolution of marsupials and placentals. *Current Biology, 25*(14), 1878-1884.
734   doi:https://doi.org/10.1016/j.cub.2015.05.043
735   Häberling, I. S., & Corballis, M. C. (2016). Cerebellar asymmetry, cortical asymmetry and handedness:
736   two independent networks. *Laterality, 21*(4-6), 397-414.
737   doi:10.1080/1357650X.2015.1110161
738   Harvey, P. H., & Clutton-Brock, T. H. (1985). Life history variation in primates. *Evolution, 39*(3), 559-
739   581. doi:https://doi.org/10.1111/j.1558-5646.1985.tb00395.x
740   Hecht, E. E., Gutman, D. A., Bradley, B. A., Preuss, T. M., & Stout, D. (2015a). Virtual dissection and
741   comparative connectivity of the superior longitudinal fasciculus in chimpanzees and humans.
742   *NeuroImage, 108*, 124-137. doi:https://doi.org/10.1016/j.neuroimage.2014.12.039
743   Hecht, E. E., Gutman, D. A., Khreisheh, N., Taylor, S. V., Kilner, J., Faisal, A. A., . . . Stout, D. (2015b).
744   Acquisition of Paleolithic toolmaking abilities involves structural remodeling to inferior
745   frontoparietal regions. *Brain Structure and Function, 220*(4), 2315-2331. doi:10.1007/s00429-
746   014-0789-6
747   Hopkins, W. D. (1995). Hand preferences for a coordinated bimanual task in 110 chimpanzees (*Pan*
748   *troglodytes*): cross-sectional analysis. *Journal of Comparative Psychology, 109*(3), 291-297.
749   doi:10.1037/0735-7036.109.3.291
750   Hopkins, W. D. (2013a). Comparing human and nonhuman primate handedness: challenges and a
751   modest proposal for consensus. *Developmental Psychobiology, 55*(6), 621-636.
752   doi:https://doi.org/10.1002/dev.21139
753   Hopkins, W. D. (2013b). Neuroanatomical asymmetries and handedness in chimpanzees (*Pan*
754   *troglodytes*): a case for continuity in the evolution of hemispheric specialization. *Annals of*
755   *the New York Academy of Sciences, 1288*(1), 17-35. doi:10.1111/nyas.12109
756   Hopkins, W. D., Adams, M. J., & Weiss, A. (2013). Genetic and environmental contributions to the
757   expression of handedness in chimpanzees (*Pan troglodytes*). *Genes, Brain and Behavior,*
758   *12*(4), 446-452. doi:https://doi.org/10.1111/gbb.12044
759   Hopkins, W. D., & Cantalupo, C. (2004). Handedness in chimpanzees (*Pan troglodytes*) is associated
760   with asymmetries of the primary motor cortex but not with homologous language areas.
761   *Behavioral Neuroscience, 118*(6), 1176-1183. doi:10.1037/0735-7044.118.6.1176
762   Hopkins, W. D., Fernandez-Carriba, S., Wesley, M. J., Hostetter, A., Pilcher, D., & Poss, S. (2001). The
763   use of bouts and frequencies in the evaluation of hand preferences for a coordinated
764   bimanual task in chimpanzees (*Pan troglodytes*): an empirical study comparing two different
765   indices of laterality. *Journal of Comparative Psychology, 115*(3), 294-299. doi:10.1037/0735-
766   7036.115.3.294

767   Hopkins, W. D., Gardner, M., Mingle, M., Reamer, L., & Schapiro, S. J. (2013). Within- and between-
768   task consistency in hand use as a means of characterizing hand preferences in captive
769   chimpanzees (*Pan troglodytes*). *Journal of Comparative Psychology, 127*(4), 380-391.
770   doi:10.1037/a0031071
771   Hopkins, W. D., Phillips, K. A., Bania, A., Calcutt, S. E., Gardner, M., Russell, J., . . . Schapiro, S. J.
772   (2011). Hand preferences for coordinated bimanual actions in 777 great apes: implications
773   for the evolution of handedness in hominins. *Journal of Human Evolution, 60*(5), 605-611.
774   doi:10.1016/j.jhevol.2010.12.008
775   Kaplan, G., & Rogers, L. J. (2021). Brain size associated with foot preferences in Australian parrots.
776   *Symmetry, 13*(5). doi:10.3390/sym13050867
777   Karolis, V. R., Corbetta, M., & Thiebaut de Schotten, M. (2019). The architecture of functional
778   lateralisation and its relationship to callosal connectivity in the human brain. *Nature*
779   *Communications, 10*(1), 1417. doi:10.1038/s41467-019-09344-1
780   Kimura, D. (1979). Neuromotor mechanisms in the evolution of human communication. In H. D. S. M.
781   Paleigh (Ed.), *Neurobiology of social communication in primates: An evolutionary perspective*
782   (pp. 1197-1220). Cambridge: Academic Press.
783   Kitchener, A. C., Meloro, C., Williams, T.M. (2017). Form and function of the musteloids. In D. W.
784   Macdonald, Newman, C., Harrington, L. (Ed.), *The Biology and Conservation of Wild*
785   *Musteloids* (pp. 98-135). Oxford: Oxford University Press.
786   Kong, X.-Z., Mathias, S. R., Guadalupe, T., Group, E. L. W., Glahn, D. C., Franke, B., . . . Francks, C.
787   (2018). Mapping cortical brain asymmetry in 17,141 healthy individuals worldwide via the
788   ENIGMA Consortium. *Proceedings of the National Academy of Sciences, 115*(22), E5154.
789   doi:10.1073/pnas.1718418115
790   Krasheninnikova, A., Chow, P. K. Y., & von Bayern, A. M. P. (2020). Comparative cognition: practical
791   shortcomings and some potential ways forward. *Canadian Journal of Experimental*
792   *Psychology/Revue canadienne de psychologie expérimentale, 74*(3), 160-169.
793   doi:10.1037/cep0000204
794   Laska, M. (1996). A study of correlates of hand preferences in squirrel monkeys (*Saimiri sciureus*).
795   *Primates, 37*(4), 457-465. doi:10.1007/BF02381379
796   Laurence, A., Wallez, C., & Blois-Heulin, C. (2011). Task complexity, posture, age, sex: Which is the
797   main factor influencing manual laterality in captive *Cercocebus torquatus torquatus*?
798   *Laterality, 16*(5), 586-606. doi:10.1080/1357650X.2010.501338
799   Lilak, A. L., & Phillips, K. A. (2008). Consistency of hand preference across low-level and high-level
800   tasks in Capuchin monkeys (Cebus apella). *American Journal of Primatology, 70*(3), 254-260.
801   doi:https://doi.org/10.1002/ajp.20485
802   Lima, M. G. M., Silva-Júnior, J. d. S. e., Černý, D., Buckner, J. C., Aleixo, A., Chang, J., . . . Lynch Alfaro,
803   J. W. (2018). A phylogenomic perspective on the robust capuchin monkey (*Sapajus*)
804   radiation: First evidence for extensive population admixture across South America. *Molecular*
805   *Phylogenetics and Evolution, 124*, 137-150. doi:https://doi.org/10.1016/j.ympev.2018.02.023
806   Lozano, M., Estalrrich, A., Bondioli, L., Fiore, I., Bermúdez de Castro, J.-M., Arsuaga, J. L., . . . Frayer, D.
807   W. (2017). Right-handed fossil humans. *Evolutionary Anthropology: Issues, News, and*
808   *Reviews, 26*(6), 313-324. doi:https://doi.org/10.1002/evan.21554
809   MacLean, E. L., Matthews, L. J., Hare, B. A., Nunn, C. L., Anderson, R. C., Aureli, F., . . . Wobber, V.
810   (2012). How does cognition evolve? Phylogenetic comparative psychology. *Animal cognition,*
811   *15*(2), 223-238. doi:10.1007/s10071-011-0448-8
812   MacNeilage, P. F. (2007). Present status of the postural origins theory. In *The evolution of*
813   *hemispheric specialization in primates*, W. D. Hopkins (Ed.) American Society of
814   Primatologists, pp. 59–91. Oxford: Elsevier.
815   MacNeilage, P. F., Studdert-Kennedy, M. G., & Lindblom, B. (1987). Primate handedness
816   reconsidered. *Behavioral and Brain Sciences, 10*(2), 247-263.
817   doi:10.1017/S0140525X00047695
818   Maille, A., Belbeoc'h, C., Rossard, A., Bec, P., & Blois-Heulin, C. (2013). Which are the features of the
819   TUBE task that make it so efficient in detecting manual asymmetries? An investigation in two

Cercopithecine species (*Cercopithecus neglectus* and *Cercocebus torquatus*). *Journal of*
Comparative Psychology, 127(4), 436-444. doi:10.1037/a0032227

ManyPrimates, Altschul, D. M., Beran, M. J., Bohn, M., Caspar, K. R., Fichtel, C., . . . Grebe, N. M.
(2019). Collaborative open science as a way to reproducibility and new insights in primate
cognition research. *Japanese Psychological Review, 62*(3), 205-220.

Marchant, L. F., & McGrew, W. C. (2013). Handedness is more than laterality: lessons from
chimpanzees. *Annals of the New York Academy of Sciences, 1288*(1), 1-8.
doi:https://doi.org/10.1111/nyas.12062

Margiotoudi, K., Marie, D., Claidière, N., Coulon, O., Roth, M., Nazarian, B., . . . Meguerditchian, A.
(2019). Handedness in monkeys reflects hemispheric specialization within the central sulcus.
An in vivo MRI study in right- and left-handed olive baboons. *Cortex, 118*, 203-211.
doi:https://doi.org/10.1016/j.cortex.2019.01.001

Meguerditchian, A., Donnot, J., Molesti, S., Francioly, R., & Vauclair, J. (2012). Sex difference in
squirrel monkeys' handedness for unimanual and bimanual coordinated tasks. *Animal*
Behaviour, 83(3), 635-643. doi:https://doi.org/10.1016/j.anbehav.2011.12.005

Meguerditchian, A., Phillips, K. A., Chapelain, A., Mahovetz, L. M., Milne, S., Stoinski, T., . . . Hopkins,
W. D. (2015). Handedness for unimanual grasping in 564 great apes: the effect on grip
morphology and a comparison with hand use for a bimanual coordinated task. *Frontiers in*
Psychology, 6, 1794-1794. doi:10.3389/fpsyg.2015.01794

Meguerditchian, A., Vauclair, J., & Hopkins, W. D. (2013). On the origins of human handedness and
language: A comparative review of hand preferences for bimanual coordinated actions and
gestural communication in nonhuman primates. *Developmental Psychobiology, 55*(6), 637-
650. doi:https://doi.org/10.1002/dev.21150

Mittermeier, R. A., Rylands, A. B., & Wilson, D. E. (2013). *Handbook of the Mammals of the World: 3.*
Primates. Barcelona: Lynx Ediciones.

Morales-Jimenez, A. L., Disotell, T., & Di Fiore, A. (2015). Revisiting the phylogenetic relationships,
biogeography, and taxonomy of spider monkeys (genus *Ateles*) in light of new molecular
data. *Molecular Phylogenetics and Evolution, 82*, 467-483.
doi:https://doi.org/10.1016/j.ympev.2014.09.019

Morino, L., Uchikoshi, M., Bercovitch, F., Hopkins, W. D., & Matsuzawa, T. (2017). Tube task hand
preference in captive hylobatids. *Primates, 58*(3), 403-412. doi:10.1007/s10329-017-0605-z

Motes Rodrigo, A., Ramirez Torres, C. E., Hernandez Salazar, L. T., & Laska, M. (2018). Hand
preferences in two unimanual and two bimanual coordinated tasks in the black-handed
spider monkey (*Ateles geoffroyi*). *Journal of Comparative Psychology, 132*(2), 220-229.
doi:10.1037/com0000110

Musgrave, S., & Sanz, C. (2018). Tool use in nonhuman primates. In *The International Encyclopedia of*
Anthropology, H. Callan (Ed.). doi:https://doi.org/10.1002/9781118924396.wbiea2063

Nelson, E. L., & Boeving, E. R. (2015). Precise digit use increases the expression of handedness in
Colombian spider monkeys (*Ateles fusciceps rufiventris*). *American Journal of Primatology,*
77(12), 1253-1262. doi:https://doi.org/10.1002/ajp.22478

Ocklenburg, S., Beste, C., Arning, L., Peterburs, J., & Güntürkün, O. (2014). The ontogenesis of
language lateralization and its relation to handedness. *Neuroscience & Biobehavioral*
Reviews, 43, 191-198. doi:https://doi.org/10.1016/j.neubiorev.2014.04.008

Papademetriou, E., Sheu, C.-F., & Michel, G. F. (2005). A meta-analysis of primate hand preferences,
particularly for reaching. *Journal of Comparative Psychology, 119*(1), 33-48.
doi:10.1037/0735-7036.119.1.33

Paradis, E. S., K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary analyses
in R. *Bioinformatics, 35*, 526-528.

Phillips, K. A., & Sherwood, C. C. (2005). Primary motor cortex asymmetry is correlated with
handedness in capuchin monkeys (*Cebus apella*). *Behavioral Neuroscience, 119*(6), 1701-
1704. doi:10.1037/0735-7044.119.6.1701

Phillips, K. A., Sherwood, C. C., & Lilak, A. L. (2007). Corpus callosum morphology in capuchin monkeys is influenced by sex and handedness. *PLOS ONE, 2*(8), e792. doi:10.1371/journal.pone.0000792

Pinheiro J., B. D., DebRoy S., Sarkar D., R Core Team. (2020). nlme: linear and nonlinear mixed effects models. R package version 3.1-148. Retrieved from https://CRAN.R-project.org/package=nlme

Prieur, J., Lemasson, A., Barbu, S., & Blois-Heulin, C. (2019). History, development and current advances concerning the evolutionary roots of human right-handedness and language: Brain lateralisation and manual laterality in non-human primates. *Ethology, 125*(1), 1-28. doi:https://doi.org/10.1111/eth.12827

Ramayya, A. G., Glasser, M. F., & Rilling, J. K. (2010). A DTI investigation of neural substrates supporting tool use. *Cerebral Cortex, 20*(3), 507-516. doi:10.1093/cercor/bhp141

Raymond, M., & Pontier, D. (2004). Is there geographical variation in human handedness? *Laterality, 9*(1), 35-51. doi:10.1080/13576500244000274

R Core Team (2020). R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Retrieved from https://www.R-project.org/

Regaiolli, B., Spiezio, C., & Hopkins, W. D. (2016). Hand preference on unimanual and bimanual tasks in strepsirrhines: The case of the ring-tailed lemur (*Lemur catta*). *American Journal of Primatology, 78*(8), 851-860. doi:https://doi.org/10.1002/ajp.22549

Regaiolli, B., Spiezio, C., & Hopkins, W. D. (2018). Hand preference on unimanual and bimanual tasks in Barbary macaques (*Macaca sylvanus*). *American Journal of Primatology, 80*(3), e22745. doi:https://doi.org/10.1002/ajp.22745

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution, 3*(2), 217-223. doi:10.1111/j.2041-210X.2011.00169.x

Richards, G., Beking, T., Kreukels, B. P. C., Geuze, R. H., Beaton, A. A., & Groothuis, T. (2021). An examination of the influence of prenatal sex hormones on handedness: Literature review and amniotic fluid data. *Hormones and Behavior, 129*, 104929. doi:https://doi.org/10.1016/j.yhbeh.2021.104929

Rilling, J. K., Glasser, M. F., Preuss, T. M., Ma, X., Zhao, T., Hu, X., & Behrens, T. E. J. (2008). The evolution of the arcuate fasciculus revealed with comparative DTI. *Nature Neuroscience, 11*(4), 426-428. doi:10.1038/nn2072

Rilling, J. K., & Insel, T. R. (1999). Differential expansion of neural projection systems in primate brain evolution. *NeuroReport, 10*(7), 1453-1459, doi:10.1097/00001756-199905140-00012

Ringo, J. L., Doty, R. W., Demeter, S., & Simard, P. Y. (1994). Time is of the essence: a conjecture that hemispheric specialization arises from interhemispheric conduction delay. *Cerebral Cortex, 4*(4), 331-343. doi:10.1093/cercor/4.4.331

Rogers, L. J. (2009). Hand and paw preferences in relation to the lateralized brain. *Philosophical Transactions of the Royal Society B: Biological Sciences, 364*(1519), 943-954. doi:10.1098/rstb.2008.0225

Roos, C., Liedigk, R., Thinh, V. N., Nadler, T., & Zinner, D. (2019). The hybrid origin of the Indochinese gray langur *Trachypithecus crepusculus*. *International Journal of Primatology, 40*(1), 9-27. doi:10.1007/s10764-017-0008-4

Sanford, C., Guin, K., & Ward, J. P. (1984). Posture and laterality in the bushbaby (*Galago senegalensis*). *Brain, Behavior and Evolution, 25*(4), 217-224. doi:10.1159/000118867

Schmitt, V., Melchisedech, S., Hammerschmidt, K., & Fischer, J. (2008). Hand preferences in Barbary macaques (*Macaca sylvanus*). *Laterality, 13*(2), 143-157. doi:10.1080/13576500701757532

Schmitz, J., Metz, G. A. S., Güntürkün, O., & Ocklenburg, S. (2017a). Beyond the genome—Towards an epigenetic understanding of handedness ontogenesis. *Progress in Neurobiology, 159*, 69-89. doi:https://doi.org/10.1016/j.pneurobio.2017.10.005

Schmitz, J., Lor, S., Klose, R., Güntürkün, O., & Ocklenburg, S. (2017b). The functional genetics of handedness and language lateralization: insights from gene ontology, pathway and disease association analyses. *Frontiers in Psychology, 8*, 1144. Retrieved from https://www.frontiersin.org/article/10.3389/fpsyg.2017.01144

Schweitzer, C., Bec, P., & Blois-Heulin, C. (2007). Does the complexity of the task influence manual
laterality in de Brazza's monkeys (*Cercopithecus neglectus*)? *Ethology, 113*(10), 983-994.
doi:https://doi.org/10.1111/j.1439-0310.2007.01405.x

Sha, Z., Pepe, A., Schijven, D., Carrión-Castillo, A., Roe, J. M., Westerhausen, R., . . . Francks, C. (2021).
Handedness and its genetic influences are associated with structural asymmetries of the
cerebral cortex in 31,864 individuals. *Proceedings of the National Academy of Sciences,*
118(47), e2113095118. doi:10.1073/pnas.2113095118

Silcox, M. T., Bloch, J. I., Boyer, D. M., Godinot, M., Ryan, T. M., Spoor, F., & Walker, A. (2009).
Semicircular canal system in early primates. *Journal of Human Evolution, 56*(3), 315-327.
doi:https://doi.org/10.1016/j.jhevol.2008.10.007

Spoelstra, K. (2021). *Lateralized behavior in white-handed gibbons (Hylobates lar).* Master Thesis.
Linköping University, Linköping.

Sussman, R. W., Tab Rasmussen, D., & Raven, P. H. (2013). Rethinking primate origins again.
American Journal of Primatology, 75(2), 95-106. doi:https://doi.org/10.1002/ajp.22096

Tab Rasmussen, D., & Nekaris, K. (1998). Evolutionary history of lorisiform primates. *Folia*
Primatologica, 69(Suppl. 1), 250-285. doi:10.1159/000052716

Tan, Y., Yoder, A. D., Yamashita, N., & Li, W.-H. (2005). Evidence from opsin genes rejects nocturnality
in ancestral primates. *Proceedings of the National Academy of Sciences of the United States*
of America, 102(41), 14712. doi:10.1073/pnas.0507042102

Toth, N. (1985). Archaeological evidence for preferential right-handedness in the lower and middle
pleistocene, and its possible implications. *Journal of Human Evolution, 14*(6), 607-614.
doi:https://doi.org/10.1016/S0047-2484(85)80087-7

Vauclair, J., Meguerditchian, A., & Hopkins, W. D. (2005). Hand preferences for unimanual and
coordinated bimanual tasks in baboons (*Papio anubis*). *Cognitive Brain Research, 25*(1), 210-
216. doi:https://doi.org/10.1016/j.cogbrainres.2005.05.012

Ward, J. P. (1998). Left hand advantage for prey capture in the galago (*Galago moholi*). *International*
Journal of Comparative Psychology, 11(4), 173-184.

Westergaard, G. C., Kuhn, H. E., & Suomi, S. J. (1998). Bipedal posture and hand preference in
humans and other primates. *Journal of Comparative Psychology, 112*(1), 55-64.
doi:10.1037/0735-7036.112.1.55

Westergaard, G. C., & Suomi, S. J. (1996). Hand preference for a bimanual task in tufted capuchins
(*Cebus apella*) and rhesus macaques (*Macaca mulatta*). *Journal of Comparative Psychology,*
110(4), 406-411. doi:10.1037/0735-7036.110.4.406

Yousry, T. A., Schmid, U. D., Alkadhi, H., Schmidt, D., Peraud, A., Buettner, A., & Winkler, P. (1997).
Localization of the motor hand area to a knob on the precentral gyrus. A new landmark.
Brain, 120(1), 141-157. doi:10.1093/brain/120.1.141

Zhao, D., Hopkins, W. D., & Li, B. (2012). Handedness in nature: first evidence on manual laterality on
bimanual coordinated tube task in wild primates. *American Journal of Physical Anthropology,*
148(1), 36-44. doi:https://doi.org/10.1002/ajpa.22038

A

Platyrrhini
Pithecia pithecia
Saimiri sciureus
Saguinus fuscicollis
Saguinus oedipus
Saguinus labiatus
Leontopithecus chrysomelas
Leontopithecus chrysopygus
Ateles geoffroyi
Ateles fusciceps
Ateles cubicollis

Hominoidea
Homo sapiens
Pan troglodytes
Pan paniscus
Gorilla gorilla
Nomascus gabriellae
Nomascus gabriellae
Symphalangus syndactylus
Hylobates lar
Hylobates moloch
Hoolock hoolock
Nasalis larvatus
Semnopithecus entellus
Rhinopithecus roxellana
Pygathrix nemaeus
Trachypithecus cristatus
Papio anubis
Theropithecus gelada
Mandrillus sphinx
Cercocebus atys
Macaca fuscicularis
Macaca nemestrina
Macaca mulatta
Macaca fascicularis
Macaca silenus
Papio papio
Theropithecus gelada
Mandrillus sphinx
Cercopithecus neglectus
Cercopithecus diana

Cercopithecoidea

MeanAbsHI
0.257 ——— 0.984

Time (Mya)
50 40 30 20 10 0

B

Platyrrhini

Hominoidea

Cercopithecoidea

MeanHI
-0.377 ——— 0.761

Coefficient estimates (incl. 95% CI)