The evolution and biological correlates of hand preferences in anthropoid primates

Kai R. Caspar[1,2*], Fabian Pallasdies[3], Larissa Mader[1], Heitor Sartorelli[4], Sabine Begall[1]

[1]Department of General Zoology, University of Duisburg-Essen, Universitätsstraße 5, 45141 Essen, Germany

[2]Department of Game Management and Wildlife Biology, Faculty of Forestry and Wood Sciences, Czech

University of Life Sciences, Praha, Czech Republic

[3]Institute for Theoretical Biology, Department of Biology, Humboldt-Universität zu Berlin, Berlin, Germany

[4]Independent researcher

Abstract

The evolution of human right-handedness has been intensively debated for decades. Manual
lateralization patterns in non-human primates have the potential to elucidate evolutionary
determinants of human handedness. However, restricted species samples and inconsistent
methodologies have so far limited comparative phylogenetic studies. By combining original data with
published literature reports, we assembled data on hand preferences for standardized object
manipulation in 1,786 individuals from 38 species of anthropoid primates, including monkeys, apes,
and humans. Based on that, we employ quantitative phylogenetic methods to test prevalent
hypotheses on the roles of ecology, brain size and tool use in primate handedness evolution. We
confirm that human right-handedness represents an unparalleled extreme among anthropoids and
found taxa displaying population-level handedness to be rare. Species-level direction of manual
lateralization was largely uniform among non-human primates and did not strongly correlate with any
of the selected biological predictors, nor with phylogeny. In contrast, we recovered highly variable
patterns of hand preference strength, which show signatures of both ecology and phylogeny. In
particular, terrestrial primates tend to display weaker hand preferences than arboreal species. These
results challenge popular ideas on primate handedness evolution, especially the postural origins
hypothesis. Furthermore, they point to a potential adaptive benefit of disparate lateralization strength
in primates, a measure of hand preference that has often been overlooked in the past. Finally, our data
show that human lateralization patterns do not align with trends found among other anthropoids,
suggesting that unique selective pressures gave rise to the unusual hand preferences of our species.

Key words: handedness, laterality, hemispheric dominance, tube task, comparative cognition, postural
origins hypothesis

Introduction

Pronounced right-handedness is a universal trait among extant human populations (Coren & Porac,
1977; Raymond & Pontier, 2004; Faurie et al., 2005) and might be an ancient attribute of the genus
Homo (Toth, 1985; Lozano et al., 2017). Whereas a significant expression of manual lateralization at
population-level is not exclusive to humans, the universal proportion of approximately 85-95 % right-
handers in our species appears to be an unmatched extreme among primates (Meguerditchian et al.,
2013). Furthermore, individual humans tend to be strongly handed with ambiguous hand
preferences being extremely rare (Cochet & Vauclair, 2012), which is also unusual when compared to
many other primate lineages (Westergaard & Suomi, 1996; Hopkins et al., 2011). Therefore, both

strength and direction of population-level manual lateralization in humans are widely considered to
be remarkable.

Handedness is a behavioral consequence of functional asymmetries in the brain (Amunts et al., 1996;
Häberling & Corballis, 2016; Sha et al., 2021). Accordingly, right-handedness results from unilateral
specializations in the left hemisphere and vice versa. In anthropoid primates, which encompass
monkeys, apes, and humans, asymmetries of the precentral gyrus in the primary motor cortex show
a particular association with individual hand preferences (Yousry et al., 1997; Phillips & Sherwood,
2005; Dadda et al., 2006; Sha et al., 2021). Nevertheless, the proximate reasons for the expression of
individual handedness in humans and other anthropoids, including its genetic basis and the influence
of brain areas located outside of the motor cortex, are by no means fully understood (Rogers, 2009;
Hopkins et al. 2013; Ocklenburg et al., 2014; Schmitz et al., 2017a; Richards et al., 2021). For this
paper, however, we exclusively focus on the evolutionary underpinnings of population-level manual
lateralization.

The origins of pronounced population-level right-handedness in the human lineage have traditionally
been linked to the emergence of complex communication mediated by manual gestures and
language, which are also predominantly processed in the left hemisphere (Corballis, 1991; Annett,
2002; Meguerditchian et al., 2013; Ocklenburg et al., 2014; Prieur et al., 2019). However, current
evidence suggests that manual and language lateralization are not nearly as tightly correlated as
traditionally believed and functional ties between these phenomena remain unidentified (Fitch &
Braccini, 2013; Ocklenburg et al., 2014; Schmitz et al., 2017b). It has also been established that
various nonhuman primates show significant asymmetries in hand-use at the individual and
population-level in a variety of tasks including manual communicative gestures and bimanual actions
(MacNeilage, 2007; Hopkins et al., 2013; Meguerditchian et al., 2013; Regaiolli et al., 2016).
Nevertheless, reports of significant population-level biases are largely confined to a comparatively
small number of species and the distribution of individuals across hand preference categories is
always far more balanced than in humans. For instance, olive baboons (*Papio anubis*), Western
gorillas (*Gorilla gorilla*), and chimpanzees (*Pan troglodytes*) have all been reported to show a
significant population-level right hand bias for bimanual manipulation but the portion of right-
handers among these species is only around 50 % (Vauclair et al., 2005; Hopkins et al., 2011 – note
that this skew is significant due to a notable portion of ambipreferent individuals among these
species). Still, it is essential to discern what underlies these comparatively weak population-level
hand preference patterns that emerged across the primate order to unravel the origins of
pronounced right-handedness in our species (MacNeilage, 2007).

The most influential conjecture to explain how primate hand preference patterns evolved is the
postural origins hypothesis (POH) (MacNeilage et al., 1987; MacNeilage, 2007). Considering galagos
as models, the POH assumes that hypothetical insectivorous primate ancestors exhibited a right-
hand bias to support their body on vertical substrates, while the left-hand specialized for fast
grasping movements, the so-called ballistic reaching (Ward, 1998). Based on this, the POH predicts
that with the emergence of anthropoid primates, which exhibit arboreal quadrupedalism and more
refined digit control, the right hand became adopted to manipulate objects during foraging
(MacNeilage et al., 1987). Hence, it proposes that all anthropoids share a right-hand bias for
manipulation, which would find its most extreme expression in humans (MacNeilage, 2007;
MacNeilage et al., 1987). In the anatomically less derived strepsirrhines, the left hand is instead
expected to be dominant (MacNeilage, 2007). However, the POH has been drastically modified by
more recent studies (Hopkins et al., 2011; Meguerditchian et al., 2013; Morino et al., 2017). The
novel interpretation proposes (in conflict with the original POH) that arboreal monkeys and apes
should display a left-hand bias for manipulation. Their right hand would provide the necessary

postural support, retaining the hypothesized ancestral primate pattern laid out by MacNeilage et al.
(1987). Terrestrial lineages, however, would no longer be bound to reserve the right hand for posture
stabilization and are expected to evolve right hand preferences for fine motor skills, eventually giving
rise to the human condition. Hence, the novel POH expects that left-handedness is prevalent in
arboreal groups, while a preponderance of right-handedness should be restricted to terrestrial
primate species (Meguerditchian et al., 2013). In contrast, the original POH expects to find a right-
handedness trend in all anthropoids, regardless of their ecology. Both versions of the POH exclusively
rely on correlational evidence from behavioral studies on hand preferences in different primate
groups.

In addition to these considerations, it has been prominently proposed that foraging-related
extractive tool use facilitated the evolution of hand preferences in humans to allow for more efficient
object handling (Kimura, 1979; Frost, 1980), but this idea has also been extended to nonhuman
primates that habitually use tools (Cashmore et al., 2008; Prieur et al., 2019). Moreover,
neuroanatomical studies demonstrated that the expression of overall neural lateralization and
hemispheric independence positively correlates with absolute brain size in primates (Rilling & Insel,
1999; Karolis et al., 2019; Ardesch et al., 2021). Such a scaling relation was already hypothesized by
Ringo et al. (1994) and implies that the strength of individual handedness could also be tied to
absolute brain size (Hopkins, 2013b); the concept does, however, not concern the direction of hand
preferences. Hence, it can be hypothesized that larger-brained primates should evolve greater
manual lateralization strength to reduce the amount of interhemispheric communication needed to
accomplish manipulative tasks. Complementary to these predictions, there is evidence that absolute
brain size is positively correlated with foot preference strength for grasping among parrot species
(Kaplan & Rogers, 2021; but note that this study did not control for phylogeny; see also Brown and
Magat, 2011).

Thus testable hypotheses on primate hand preference evolution have long been established, but
none of them has so far been tested within quantitative evolutionary frameworks (Hopkins, 2013a).
Studies in comparative cognition increasingly rely on phylogenetically informed modelling to
estimate how and when specific behaviors evolved (MacLean et al., 2012; ManyPrimates et al., 2019;
Krasheninnikova et al., 2020). Such approaches can provide estimates of ancestral states and allow
researchers to quantify the pace as well as the influence of phylogeny and ecological variables on
cognitive evolution (MacLean et al., 2012). However, to yield meaningful results, dense taxonomic
sampling of species and a consistent testing scheme are required (Freckleton et al., 2002;
Krasheninnikova et al., 2020). For research on primate hand preferences, this means that subjects
from different species need to engage in the same experimental task to assess lateralization.
Standardization is particularly important since both the strength and the direction of individual hand
preferences can be task-dependent and because the expression of manual lateralization correlates
positively with motor complexity (Vauclair et al., 2005; Blois-Heulin et al., 2007; Lilak & Phillips, 2008;
Meguerditchian et al., 2015; Caspar et al., 2018). Bimanual actions that involve both hands and which
spontaneously occur during food manipulation or tool use are particularly suitable to detect hand
preferences (Meguerditchian et al., 2013), while non-gestural unimanual actions such as grasping
often do not elicit pronounced group or individual-level lateralized responses (Papademetriou et al.,
2005; Rogers, 2009; Hopkins et al., 2013a). For comparative approaches in primates, tool handling is
only of limited use as a model behavior, because habitual tool use is only found in few lineages
(Musgrave & Sanz, 2018). Therefore, bimanual actions related to foraging present themselves as
suitable candidates for comparative studies on hand preferences in these animals.

A simple and widely applicable experiment to determine anthropoid primate hand preferences is the
so-called tube task (Hopkins, 1995): A subject is handed a PVC tube filled with desired food. To

extract it, one hand must hold the tube while the other has to engage in the more complicated action
of retrieving the food mesh, thereby revealing biases in hand use dominance. Results from the tube
task have been demonstrated to robustly correlate with hemispheric asymmetries in various primate
groups (New World monkeys: Phillips & Sherwood, 2005; Old World monkeys: Margiotoudi et al.,
2019; apes: Hopkins & Cantalupo, 2004; Dadda et al., 2006) and to be intraindividually consistent
across re-tests, even if these were separated by several years (Hopkins et al., 2001). Furthermore, its
simplicity allows uniform testing of a wide range of species in captive as well as natural settings (Zhao
et al., 2012).

Here we compiled a comprehensive multispecies tube task dataset to test pertaining hypotheses on
the evolution of primate manual lateralization by means of phylogenetically informed modelling. This
way we provide a broad comparative perspective on the origins of human right-handedness in the
context of object manipulation.

Results

Lateralization strength (MeanAbsHI) but not direction (MeanHI) displays a strong phylogenetic signal
among anthropoids (Figure 1). Accordingly, MeanAbsHI (λ = 0.89, $p_{\text{likelihood-ratio test}}$ = 0.03) varied
substantially between lineages but was often stable within groups of closely related taxa. MeanHI on
the other hand, showed random fluctuations around zero (λ < 0.001; $p_{\text{likelihood-ratio test}}$ = 1), and notable
population-level shifts towards either right- or left-hand preferences were uncommon (see below).
Species-level distributions of hand preferences are summarized in Table 1 and are visualized at
genus-level in Figure 2.

Ancestral hand preference strength was modelled to have been similarly moderate in the stem
lineages of hominoids (AbsHI = 0.606) and cercopithecoids (AbsHI = 0.627), while it was inferred to
be higher in the ancestral platyrrhines (AbsHI = 0.740; Figure 1, Suppl. File 1). We found hand
preference strength to be particularly weak among some species in the Papionina clade (baboons
and their relatives) and to be least expressed in the gelada (*Theropithecus gelada* – MeanAbsHI =
0.257). The strongest preferences were found in humans (MeanAbsHI = 0.943), langurs of the genus
Trachypithecus (MeanAbsHI = 0.868), and spider monkeys (*Ateles* spp. – MeanAbsHI = 0.831). Very
pronounced individual preferences were also found in white-faced sakis (*Pithecia pithecia* –
MeanAbsHI = 0.934) but our sample size for this species was notably small (*n* = 7). In consequence, all
aforementioned taxa included only very few, if any, ambipreferent individuals. Besides the saki genus
Pithecia, the South Asian colobines *Rhinopithecus* and *Trachypithecus* were the only genera in which
no ambipreferent individuals were found. Direction of manual lateralization was far more uniform
across the anthropoid radiation than strength (Figure 1; Suppl. File 1). No species approached the
extreme direction bias of humans (compare Figure 2), and only 2 of 37 non-human species exhibited
a significant population-level bias as indicated by the one-sample t-test after correcting for multiple
testing (4 of 37 when no correction was applied), namely gorillas and chimpanzees. Thus, such biases
were restricted to the African ape lineage. After correcting for multiple testing, intraspecific
frequencies of ambipreferent individuals, right-, and left-handers, as indicated by the chi-square
goodness-of-fit test, differed significantly from the superordinate taxon estimate in five species.
These were humans and chimpanzees, which are both right-handed at the population-level, and the
predominately left-handed golden snub-nosed monkeys (*Rhinopithecus roxellana*), as well as geladas
and the black lion tamarin (*Leontopithecus chrysopygus*), both of which encompass a large
proportion of ambipreferent individuals (Tab. 1). When omitting correction, respective biases were
notably more frequent and found in 12 non-human species across the three major clades studied.
Species- and genus-level results closely corresponded to each other (Tab. 1).

Our selected predictors for PGLS models showed a mixed and overall weak performance in explaining
expression patterns of hand preference strength and direction in anthropoids (Table 2, Figure 3). For
the initial model on lateralization direction, we found a significant effect of ecology, with terrestrial
species displaying a right-hand bias compared to arboreal ones (p = 0.04; Table 2A). However, when
humans were removed from the model, this effect was merely recovered as a non-significant trend
(p = 0.07; Table 2B). Other predictors had no significant effect on lateralization direction, regardless
of whether humans were considered in the model or not (p > 0.2, Tables 2A and 2B). When humans
were included (Table 3A), models encompassing the component ecology as well as some considering
brain size outperformed the lateralization direction null model. When humans were omitted from
the analysis, the null model was solely outperformed by one that exclusively included the ecological
component, and only slightly so (Δ AICc = 1.23; Table 3B), indicating a notable bias derived from the
extreme right-handedness found in our species. Thus, whereas habitual tool use and absolute brain
size clearly do not influence the direction of lateralization among anthropoids in general, the
analyses provide evidence for a weak but detectable effect of ecology in non-human taxa. Such
ecological signatures were found to be of somewhat greater relevance for patterns of lateralization
strength. Here, a significant negative effect of a terrestrial lifestyle was found (p = 0.04; see Table
2C). In line with that, models including the component ecology consistently outperformed the null
model, which was not the case for those including only tool use and/or brain size (Table 3C). Still,
even the accuracy of the model relying on ecology alone exceeded that of the null model only
moderately (Δ AICc = 2.53; Table 3C). Thus, terrestrial anthropoids tend to show weaker hand
preferences than arboreal ones while there is no correlation with brain size or habitual tool use for
this trait.

At the individual level, Bayesian models showed that neither age, nor sex had an influence on
lateralization direction when the total sample was concerned (the respective credible intervals
overlapped with zero; Suppl. File 2). However, we recovered a notable effect of age on lateralization
direction in the hominoid subsample exclusively, with subadults tending more towards left-
handedness than adults (credible interval = -0.28 – -0.01; Suppl. File 2). We observed a different
pattern for lateralization strength. Here, an effect of age but again not sex was recovered for both
the total sample (credible interval = -0.11 – -0.05) and the hominoid subsample (credible interval = -
0.16 – -0.02), with weaker lateralization in subadults compared to adults. Such an influence of age
was not detectable in neither platyrrhines nor cercopithecoids when these taxa were separately
considered (credible intervals overlapping with zero; Suppl. File 2).

Figure 1: A color-coded phylogeny of hand preferences in anthropoid primates. The strength (A) and direction (B) of laterality, expressed by the mean absolute handedness index (MeanAbsHI) and the mean handedness index (MeanHI; 0 is marked by the black bar on the color scale), respectively, calculated for each species and inferred for each tree node by maximum likelihood estimates. Silhouettes by Kai R. Caspar, except *Ateles* (by Yan Wong, public domain) and *Homo* (public domain)

Figure 2: Violin plots of hand preference distribution in 22 genera of anthropoid primates and the genus-specific expression of three potential biological correlates (ecology, habitual foraging-related tool use, and absolute brain size). Attributions only apply to the species that represent the respective genus within our sample. Color coding: Ecology – green: arboreal, yellow: terrestrial; Habitual tool use – grey: present; white: absent. Brain size is visualized here as the log-transformed genus average of female endocranial volume.

226 **Table 1**: Hand preferences of anthropoid species as recovered by the tube task. * Results remain significant after Bonferroni correction.

Species	N	# left (%)	# right (%)	# ambipreferent (%)	MeanHI	MeanAbsHI	Species direction bias (HI), p value	Species L/R/A distribution, p value	nGenus	Genus direction bias (HI), p value	Genus L/R/A distribution, p value
Ateles fusciceps	46	20 (43.5)	22 (47.8)	4 (8.7)	0.063	0.798	0.618	0.288	87	0.759	**0.031**
Ateles geoffroyi	23	10 (43.5)	11 (47.8)	2 (8.7)	0.061	0.829	0.748	0.536			
Ateles hybridus	18	13 (72.2)	5 (27.8)	0	-0.377	0.917	0.086	**0.018**			
Cercocebus torquatus	31	13 (41.9)	11 (35.5)	7 (22.6)	-0.029	0.665	0.832	0.836	31	0.832	0.836
Cercopithecus diana	20	7 (35)	10 (50)	3 (15)	0.178	0.755	0.339	0.836	45	0.572	0.222
Cercopithecus neglectus	25	14 (56)	7 (28)	4 (16)	-0.258	0.621	0.061	0.140			
Gorilla gorilla	76	17 (22.4)	41 (53.9)	18 (23.7)	0.248	0.541	**< 0.001***	**0.007**	76	**< 0.001***	**0.007**
Homo sapiens	127	12 (9.5)	111 (87.4)	4 (3.1)	0.761	0.943	**< 0.001***	**< 0.001***	127	**< 0.001***	**< 0.001***
Hylobates lar	36	17 (47.2)	16 (44.5)	3 (8.3)	-0.011	0.614	0.924	0.182	58	0.612	0.125
Hylobates moloch	22	11 (50)	8 (36.4)	3 (13.6)	-0.115	0.799	0.540	0.552			
Leontopithecus chrysomelas	30	7 (23.3)	12 (40)	11 (36.7)	0.151	0.514	0.171	**0.012**	73	0.241	**< 0.001***
Leontopithecus chrysopygus	15	3 (20)	4 (26.7)	8 (53.3)	0.039	0.350	0.744	**0.001***			
Leontopithecus rosalia	28	10 (35.7)	8 (28.6)	10 (35.7)	0.022	0.502	0.850	**0.033**			
Macaca fascicularis	20	8 (45)	10 (45)	2 (10)	-0.036	0.684	0.835	0.233	102	0.692	0.863
Macaca nemestrina	29	9 (31)	11 (37.9)	9 (31)	0.035	0.527	0.768	0.750			
Macaca silenus	35	14 (40)	9 (25.7)	12 (34.3)	-0.051	0.467	0.596	0.328			
Macaca sylvanus	24	12 (50)	10 (41.7)	2 (8.3)	-0.025	0.670	0.873	0.129			
Macaca tonkeana	14	5 (35.7)	3 (21.4)	6 (42.9)	-0.057	0.543	0.753	0.291			
Mandrillus sphinx	32	6 (18.8)	10 (31.2)	16 (50)	0.034	0.389	0.701	**0.006**	32	0.701	**0.006**
Nomascus gabriellae	10	5 (50)	2 (20)	3 (30)	-0.173	0.618	0.465	0.436	36	0.539	0.805
Nomascus leucogenys	26	9 (34.6)	11 (42.3)	6 (23.1)	-0.031	0.555	0.818	0.869			
Pan paniscus	118	50 (42.4)	51 (43.2)	17 (14.4)	0.044	0.529	0.431	0.237	654	**< 0.001***	**< 0.001***
Pan troglodytes	536	155 (28.9)	266 (49.6)	115 (21.5)	0.133	0.507	**< 0.001***	**< 0.001***			
Papio anubis	84	27 (32.1)	41 (48.8)	16 (19.1)	0.108	0.527	0.102	0.073	108	0.079	0.239
Papio hamadryas	24	6 (25)	7 (29.2)	11 (45.8)	0.066	0.408	0.533	0.082			

	N								N		
Pithecia pithecia	7	5 (71.4)	2 (28.6)	0	-0.385	0.934	0.312	0.221	7	NA	0.221
Pongo sp.	47	27 (57.5)	9 (19.1)	11 (23.4)	-0.225	0.487	**0.006**	**0.012**	47	**0.006**	**0.012**
Pygathrix cinerea	18	6 (33.3)	10 (55.6)	2 (11.1)	0.165	0.499	0.268	0.196	18	0.268	0.196
Rhinopithecus roxellana	24	17 (70.8)	7 (29.2)	0	-0.319	0.729	**0.040**	**< 0.001***	24	**0.040**	**< 0.001***
Saimiri sciureus	36	21 (58.4)	14 (38.9)	1 (2.7)	-0.119	0.757	0.382	**0.031**	36	0.382	**0.031**
Sapajus apella	25	11 (44)	10 (40)	4 (16)	-0.028	0.687	0.854	0.961	80	0.922	0.905
Sapajus flavius	21	10 (47.6)	7 (33.3)	4 (19)	-0.130	0.769	0.495	0.755			
Sapajus xanthosternos	34	14 (41.2)	15 (44.1)	5 (14.7)	0.089	0.677	0.492	0.906			
Semnopithecus entellus	30	15 (50)	7 (23.4)	8 (26.6)	-0.184	0.560	0.110	0.315	30	0.110	0.315
Symphalangus syndactylus	31	11 (35.5)	9 (29)	11 (35.5)	-0.048	0.482	0.663	0.118	31	0.663	0.118
Theropithecus gelada	38	6 (15.8)	6 (15.8)	26 (68.4)	0.053	0.257	0.326	**< 0.001***	38	0.326	**< 0.001***
Trachypithecus auratus	8	5 (62.5)	3 (37.5)	0	-0.256	0.984	0.499	0.176	26	0.153	0.004
Trachypithecus hatinhensis	18	11 (61.1)	7 (38.9)	0	-0.248	0.817	0.219	**0.023**			

Figure 3: Visualization of PGLS coefficient estimates (including 95% confidence intervals) for the
influence of brain size, tool use, and ecology on lateralization direction (A, B) as well as strength (C) in
anthropoid primates. Two models for lateralization direction were computed, one including (A), the
other one excluding humans (B).

Table 2: Conditional average of PGLS model coefficients for lateralization strength and direction in
anthropoid primate species. VIF = variable inflation factor.

A: Conditional PGLS model average for lateralization direction, including humans				
Predictor	**Estimate**	**Std. Error**	**VIF**	***p* value**
Ecology (Terrestrial lifestyle)	0.153	0.072	1.499	0.040
Tool use (present)	0.104	0.082	1.164	0.220
Log$_{10}$ brain size	0.050	0.043	1.612	0.254
B: Conditional PGLS model average for lateralization direction, excluding humans				
Predictor	**Estimate**	**Std. Error**	**VIF**	***p* value**
Ecology (Terrestrial lifestyle)	0.108	0.056	1.419	0.067
Log$_{10}$ brain size	-0.020	0.037	1.405	0.601
Tool use (present)	0.003	0.068	1.098	0.962
C: Conditional PGLS model average for lateralization strength				
Predictor	**Estimate**	**Std. Error**	**VIF**	***p* value**
Ecology (Terrestrial lifestyle)	-0.143	0.067	1.813	**0.040**
Tool use (present)	0.060	0.070	1.235	0.402
Log$_{10}$ brain size	0.035	0.047	1.997	0.468

240 **Table 3:** Results of PGLS model averaging for lateralization direction (considering the inclusion and
241 exclusion of humans) and strength. Null models are shown in italics. Df. = Degrees of freedom. AICc =
242 Second order Akaike information criterion.

A: PGLS model for lateralization direction, including humans				
Components	**Df.**	**AICc**	**Δ AICc**	**Weight**
Ecology	3	-13.60	0	0.28
Ecology, tool use	4	-13.37	0.23	0.25
Brain size	3	-12.22	1.38	0.14

Ecology, brain size	4	-12.07	1.54	0.13
Ecology, tool use, brain size	5	-10.87	2.73	0.07
(NULL)	*2*	*-10.37*	*3.23*	*0.05*
Tool use, brain size	4	-10.20	3.41	0.05
Tool use	3	-9.48	4.12	0.04

B: PGLS model for lateralization direction, excluding humans

Components	Df.	AICc	Δ	Weight
Ecology	3	-30.60	0	0.37
(NULL)	*2*	*-29.37*	*1.23*	*0.20*
Ecology, brain size	4	-28.82	1.78	0.15
Ecology, tool use	4	-28.01	2.59	0.10
Tool use	3	-26.98	3.62	0.06
Brain size	3	-26.97	3.63	0.06
Ecology, tool use, brain size	5	-26.16	4.44	0.04
Tool use, brain size	4	-24.42	6.18	0.02

C: PGLS model for lateralization strength

Components	Df.	AICc	Δ AICc	Weight
Ecology	3	-8.84	0	0.35
Ecology, brain size	4	-7.85	0.99	0.21
Ecology, tool use	4	-7.33	1.51	0.16
(NULL)	*2*	*-6.31*	*2.53*	*0.10*
Ecology, brain size, tool use	5	-5.46	3.38	0.06
Tool use	3	-5.05	3.80	0.05
Brain size	3	-4.14	4.71	0.03
Tool use, brain size	4	-3.19	5.66	0.02

Discussion

General

Our study provides the first quantitative phylogenetic perspectives on hand preferences in monkeys, apes, and humans. While population-level lateralization strength is highly varied among anthropoid primates and often distinctly expressed in specific lineages, direction fluctuates irrespective of phylogeny and appears comparatively uniform. Indeed, significant population-level biases in the latter are notably rare, both at the species and genus level. After expanding the sample size for some species in which such biases have been previously reported based on the tube task (siamang – Morino et al., 2017; de Brazza's monkey – Schweitzer et al., 2007), we failed to replicate significant deviations from a chance distribution (even if not correcting for multiple testing). The only taxon in which significant hand use biases for bimanual manipulation occur frequently, is constituted by the great apes and humans. Nevertheless, since sample sizes for species in this group are consistently and conspicuously large, statistical analyses performed on them (in particular the commonly applied

one-sample t-test) will have higher power compared to tests done on taxa represented by a smaller
number of individuals. It is therefore premature to assume that hominids display qualitatively
different population-level lateralization patterns than other primates based on this statistical
parameter. In fact, we would like to stress that due to this issue, simply determining the presence or
absence of significant population-level handedness is of little merit for comparative studies, since
species with vastly different hand preference distributions might fall into either category dependent
on the available sample sizes (Figure 2). Looking at the population-level frequencies of right-handed,
left-handed, and ambipreferent individuals, non-hominoid species such as the golden snub-nosed
monkey (70.8 % left-handers, 0 % ambipreferents, MeanHI: -0.319) and the brown spider monkey
(72.2 % left-handers, 0 % ambipreferents, MeanHI: -0.377) approach a human-like skew more than
any great ape species does, albeit in the contrary direction to lateralization in our species
(approximated herein as encompassing 87.3% right-handers, 3.1 % ambipreferents, MeanHI: 0.761).
Whether the hand preference patterns recovered for these so far only poorly-sampled monkeys are
indeed representative needs to be clarified by future studies encompassing greater numbers of
individuals.

Our finding that hand preference strength is generally weaker in juveniles compared to adults
replicates results from several studies relying on smaller sample sizes (e.g., Westergaard & Suomi,
1996; Zhao et al., 2012). The result that subadult non-human hominoids, but not other anthropoids,
are notably stronger left-handed than adults, likely derives from the aforementioned fact that
juveniles are more weakly lateralized overall in conjunction with the composition of our sample for
Bayesian models. Therein, Western gorillas and bonobos (*Pan paniscus*), both of which tend to be
right-handed as adults (Hopkins et al., 2011), are well-represented across age groups. Again, we
suggest that an expanded dataset encompassing both an increased number of individuals and
species could potentially level out the lateralization differences between (great) apes and other
anthropoids observed in our study.

The fact that population-level hand preference fluctuates without phylogenetic and with rather weak
ecological signatures among anthropoids, suggests that there are no strong directional selective
pressures acting on this trait, different from what pertaining hypotheses predict (see below). On the
other hand, population-level lateralization strength is more variable but consistent among closely
related taxa, thus exhibiting a strong phylogenetic signal. PGLS modelling demonstrated a significant
negative effect of a terrestrial lifestyle on hand preference strength, indicating a relevant but
previously undescribed influence of ecology. It appears intuitive that terrestrial taxa tend to be less
lateralized than arboreal ones, since the latter often need to flexibly stabilize their body in the
canopy, for instance while foraging. Accordingly, one hand will be preferably used to provide such
support, but whether the left or the right one adopts this function seems to be arbitrary. The fact
that these lateralization tendencies were found in zoo-housed primates that often adopt locomotor
regimes very different from their wild conspecifics (e.g., captive spider monkeys and gibbons spend
considerable amounts of time moving and feeding on the ground), suggest a significant innate
component to these patterns. However, within ecologically uniform groups, the variability of hand
preference strength can still be notable, at times even among closely related taxa (compare e.g.,
Javan gibbon and siamang), pointing at important but yet unidentified biological influences being at
play as well. Accordingly, the explanatory power of PGLS models for lateralization strength, which
considered the component ecology, only moderately, although consistently, exceeded that of a null
model (compare Tab. 3C). Nevertheless, given the great variability of lateralization strength among
anthropoids and its ties to phylogeny as well as ecology, this aspect of manual lateralization should
receive more research attention in the future. In the past, most work and evolutionary
considerations regarding primate handedness have instead focused on lateralization direction, surely

for anthropocentric reasons. As we attempt to show here, however, the explanatory power of these
in parts very long-lived conjectures appears to be limited.

Testing prevalent hypotheses

Our data does not unambiguously support any of the tested hypotheses on hand preference
evolution in primates. The traditional postural origins hypothesis (POH) assumes right-hand
tendencies for manipulation across anthropoid taxa. However, we found that anthropoid population-
level lateralization is in most cases not notably shifted into either direction, with a slight majority of
species displaying a weak left-hand bias (21 of 38 species). It is important to note that the recovered
correlation between arboreality and hand preference strength does not corroborate any version of
the POH, as they focus exclusively on lateralization direction. The novel POH assumes that terrestrial
non-human primates tend to be right-handed, while arboreal ones tend to be left-handed, a
prediction that gains weak support from our data. While the mean HIs of terrestrial and arboreal
non-human species appear dissimilar, they do not deviate strongly from zero and there is a notable
overlap in variation between the two ecological groups (MeanHI$_{arboreal}$: -0.08, SD: 0.16;
MeanHI$_{terrestrial}$: 0.04, SD: 0.10). Thus, the predictive power of the hypothesis is comparatively low,
which is reflected by our modelling results (Tables 2 and 3). Admittedly, improved sampling, both
within and across species might consolidate this pattern in the future, so that we certainly do not
want to dismiss relevant effects of ecology on lateralization direction at this point.

However, even if its predictions should become confirmed eventually, this would by no means
necessarily validate the theoretical framework of the novel POH. First, the hypothesis does not
provide an evolutionary mechanism for why population-level hand preference patterns should be
coupled with positional behavior. This represents a considerable conceptual shortcoming that needs
to be addressed in the future. Second, it is important to note that the evolutionary scenario
proposed by both versions of the POH is outdated and should not be perpetuated without explicitly
stating its flaws. According to the POH, small-bodied bushbabies (genus *Galago*) are suitable models
for early primates, since they would be "the most direct descendants of the earliest forms"
(MacNeilage, 2007). Because contemporary studies suggested a left-hand bias for prey grasping in
bushbabies, such a pattern was also assumed for primate ancestors in the paper that introduced the
original POH (MacNeilage et al., 1987). However, these assumptions have always been speculative
and become problematic in light of more recent data. Importantly, there is no convincing evidence
for preferably left-handed grasping in the genus *Galago,* or other galagids, anymore (Papademetriou
et al., 2005). Furthermore, bushbabies represent a remarkably derived radiation of strepsirrhines
(Tab Rasmussen & Nekaris, 1998) and are thus no suitable ecological models for the common
ancestor of modern primates. Current evidence suggests that both the earliest primates and the
ancestors of the anthropoid clade studied herein, were omnivorous arboreal quadrupeds with
moderate leaping ability (Silcox et al., 2009; Gebo, 2011; Sussman, Tab Rasmussen, & Raven, 2013)
and possibly diurnal habits (Tan et al., 2005; Ankel-Simons & Rasmussen, 2008). Thus, they were
extremely different from extant galagos. Our results indicate that the common ancestor of
anthropoids did not display notable population-level hand preferences for manipulation, nor that
such biases are common among extant monkeys and apes. Their occurence among strepsirrhines still
needs to be comparatively studied. While lemurs (different from galagos) indeed appear to show a
consistent left-hand preference for unimanual reaching, this pattern is not recovered in bimanual
tasks (Papademetriou et al., 2005; Regaiolli et al., 2016; Batist & Mayhew, 2020), again contradicting
the POH (MacNeilage, 2007). In conclusion, the evolutionary scenario proposed by both versions of
the POH is unsupported by current data. Furthermore, our findings contradict the predictions of the
original POH and only provide a very fragile empirical basis for the novel version of the hypothesis.

This leads us to challenge the status of the POH as a keystone idea in the discourse on the evolution
of primate manual lateralization.

We also found no effects of foraging-related tool use on neither direction, nor strength of
lateralization although our sample represented all primate lineages that include habitual tool users
(Musgrave & Sanz, 2018). Surprisingly, we also did not recover notable influences of absolute brain
size on hand preferences. An effect on lateralization strength was expected both on theoretical
considerations (Ringo et al., 1994) and empirical evidence from studies investigating intra- and
interspecific covariation of brain size and overall cortical lateralization (Kong et al., 2018; Ardesch et
al., 2021). Why do anthropoid hand preferences not conform to these predictions? We cannot
provide a satisfying answer to this question. It is possible that the effects of increased overall brain
lateralization on hand preference expression turn out to be unexpectedly weak and are masked by
yet unidentified neurologic factors.

All in all, none of the hypotheses on primate handedness that we addressed were clearly
corroborated by our data. Nevertheless, when discussing conflicts between prevalent ideas and our
results, we also need to address general limitations of our study framework. For instance, we equate
general hand preferences with tube task results. Although the tube task represents one of the best
available behavioral assays for brain lateralization (Dadda et al., 2006; Margiotoudi et al., 2019), it is
obvious that other hand use situations, such as communicative gesturing, need to be considered to
arrive at a holistic understanding of primate hand use evolution (Hopkins et al., 2013; Becker et al.,
2022). Such an approach could also test how variable hand use consistency across contexts is among
primates and whether ecological variables have an influence here as well. At the moment, it appears
as if there is comparatively little consistency in manual preferences across different hand use
situations in non-human primates (Laska, 1996; Lilak & Phillips, 2008; Marchant & McGrew, 2013;
Caspar et al., 2018), but most studies so far compare tasks of varying complexity in a few model
species and cases of consistent hand use ("true handedness") have indeed been reported (Diamond
& McGrew, 1994; Hopkins et al., 2013; but note neuroanatomical evidence for a dissociation of
lateralization in non-communicative actions and gestures in baboons – Becker et al., 2022).

Another limitation is posed by our sample composition. Both the number of species and subjects per
species need to be increased to further validate the patterns communicated here. In particular,
extended sampling of the speciose New World monkey families Pitheciidae and Callitrichidae would
be desirable, to make the inferences presented herein more robust. To allow for proper predictions
on hand preferences in anthropoid ancestors and early crown-group primates, additional tests with
tarsiers and strepsirrhines would be crucial. Our experience suggests that at least pitheciids and
lemurs only reluctantly engage in the tube task so that it might be advisable to apply different
bimanual testing schemes in these groups. In lemurs, puzzle boxes have been employed as such: The
animals open the lid of a box with one hand while the other one is retrieving food stored within
(Regaiolli et al., 2016; Batist & Mayhew, 2020). Future studies need to check the functional
equivalence of this approach with the tube task (which is not a trivial question, compare e.g., Lilak
and Phillips, 2008) to establish a set of behavioral assays that could be employed to study hand
preferences in the whole primate order. These methods might then also be applied to other
dexterous and ecologically variable mammalian groups, such as musteloid carnivorans (Kitchener,
2017), to test hypotheses on the evolution of manual laterality across a wider phylogenetic margin.
Finally, we need to acknowledge the limitations of our phylogenetic modelling approach. In
particular, the binary coding of *ecology* (arboreal vs terrestrial) obviously simplifies the remarkable
spectrum of positional behaviors found among anthropoids. Future studies might explore alternative
strategies to statistically code this multifaceted variable.

The evolutionary issue of human handedness

In line with previous research, we found human right-handedness to be unparalleled among
primates. We want to stress, however, that humans only deviate markedly from all other taxa in
direction and not in strength of lateralization for bimanual manipulation. When it comes to the
latter, the human condition is at least approached by groups such as leaf monkeys and spider
monkeys. Perhaps surprisingly, handedness strength in great apes is modest in comparison (Table 1).
Still, humans are highly deviant among predominately terrestrial primates in displaying such strong
individual hand preferences. Whether this difference relates to bipedal locomotion, which has often
been championed as a correlate of human handedness (Westergaard et al., 1998; Cashmore et al.,
2008; Prieur et al., 2019;), is open for debate. Since no other extant primate shows similar
adaptations to terrestrial bipedalism, the validity of this assumption is hard to test in the framework
of comparative approaches (but see Giljov et al., 2015). Quadrupedal primates tend to exhibit
stronger hand preferences when adopting the relatively instable bipedal posture (Westergaard et al.,
1998). Whether this finding has any evolutionary implications remains unclear and it should be
emphasized that although humans are bipeds, a high percentage of complex manual actions,
including numerous examples of bimanual manipulation and tool use, are not (and never have been)
habitually performed in a bipedal posture. In any case, our results suggest that bipedalism is at least
not a prerequisite to evolve strong hand preferences in anthropoid primates.

When turning to lateralization direction, the statement of Corballis (1987) remains valid: some non-
primate vertebrates approach humans more closely in population-level handedness than their simian
relatives do. Apart from humans, extreme forms of vertebrate limb use biases are known from
parrots (Kaplan & Rogers, 2021) and ground-living kangaroos (Giljov et al., 2015). Why these very
different groups converge in this respect remains enigmatic. So why do humans stand out among the
primate order when it comes to handedness direction? The limited insights gained by comparative
behavioral studies, including this one, do indeed suggest that the extreme right-handedness of
humans is a trait that evolved due to unique neurophysiological demands not experienced by other
primates. Frost (1980) already pointed out that humans' outstanding proficiency in tool use and
manufacture should be considered a significant influence on handedness evolution. Thus, not
foraging-related tool use per se, but the unique way in which it became immersed into complex
human behaviors might have influenced overall brain lateralization in our lineage. In line with that,
areas of the prefrontal cortex involved in motor cognition are structurally derived in humans and
differ significantly from their homologs in apes and monkeys (Hecht et al., 2015a; Barrett et al.,
2020). However, specializations of both the right and the left hemisphere are determining human-
specific tool use proficiency and motor planning, apparently with particular involvement of the right
inferior frontal gyrus (Ramayya et al., 2010; Hecht et al., 2015a; Hecht et al., 2015b). Postulating that
hominin tool use and right-handedness evolved in tandem is therefore not straight-forward.

Besides that, there is of course the notion of coevolution between language and handedness, which
might explain human-specific patterns of population-level manual lateralization. For this hypothesis
to be convincing, the development and function of neural substrates controlling vocal behavior and
those regulating manual motor control would need to be uniquely intertwined in humans. Indeed,
the connectivity of the arcuate fasciculus, a tract critically involved in language processing and
production, is highly derived in humans, suggesting important qualitative differences to other species
(Rilling et al., 2008; but see Barrett et al., 2020 for other elements relevant for language production
which are conserved across catarrhine primates). Nevertheless, how such traits could functionally
relate to population-level handedness remains totally unclear. In fact, despite the popularity of the
idea, a link between handedness and language processing that goes beyond superficial left-
hemisphere collateralization in right handers (not even in the majority of left-handers) is far from
evident (Fitch & Braccini, 2013). To defend an evolutionary connection between these phenomena,
pleiotropic or otherwise functionally linked genes influencing the development of both language

areas and those related to handedness would need to be identified. So far, this has not been
accomplished and current evidence suggests that language and handedness are largely independent
on various structural levels (Ocklenburg et al., 2014; Schmitz et al., 2017b). Hence, despite the
continuing efforts to unravel the evolution of human right-handedness, including the ones made by
us herein, it remains an essentially unsolved issue of human cognitive evolution.

Conclusions

We recover highly variable patterns of hand preference strength in anthropoid primates, which
correlate with ecology and phylogeny. In contrast to this, no phylogenetic signal and weaker
ecological effects were found for lateralization direction, and few species exhibit significant
population-level hand preferences. We tested three pertaining conjectures on primate handedness
evolution, the POH, tool use and brain size hypotheses, but none were unambiguously corroborated
by our data. Novel ideas on the evolution of primate hand preference should put a stronger focus on
manual lateralization strength rather than direction to address the phylogenetic patterns described
herein. However, additional datasets on primates and potentially non-primate mammals are needed
to robustly inform respective hypotheses. By relying on standardized testing paradigms, such as the
tube task, researchers can effectively build on our as well as others results and expand multispecies
datasets for further comparative phylogenetic studies. Although we are convinced that such
approaches could significantly improve our understanding of general trends in the evolution of
primate hand preferences, the unusual autapomorphic handedness pattern of humans will very likely
require explanations that cannot be derived from such comparative behavioral studies. The
evolutionary underpinnings of handedness expression in our species remain enigmatic.

Materials and Methods

Subjects

We analyzed the expression of hand preferences for object manipulation in the tube task, as well as
potential factors influencing their evolution, for a dataset of anthropoid primates (infraorder
Anthropoidea: New World monkeys (Platyrrhini), Old World monkeys (Cercopithecoidea), and apes
(Hominoidea)) from 38 species. Data from 501 individuals belonging to 26 primate species were
collected in the tube task paradigm (see below) between September 2017 and May 2020 in 39
institutions in Europe, Brazil, and Indonesia (Table 1). Of these species, 14 had never been tested in
the tube task before. Additional datasets were drawn from the literature, resulting in a total sample
of 1,786 individuals from 38 species and 20 genera, covering all anthropoid primate families except
Aotidae. Data for humans were derived from Cochet and Vauclair (2012). In this study, participants
had to use one hand to repeatedly retrieve pieces of paper out of a plastic cylinder while the other
one had to tilt and stabilize the receptacle. We considered this bimanual testing paradigm as
functionally equivalent to the tube task. Our complete study sample with annotated respective
literature sources and raw data on manual lateralization (itemized at the individual level and also at
the insertion-level for data generated in this study) can be viewed at Dryad
(https://doi.org/10.5061/dryad.8sf7m0crv).

We classified the tested subjects into two age categories, adults (n = 1,307, sexually mature
individuals) and subadults; the latter being comprised by infants (n = 9, individuals that had not yet
been weaned) and juveniles (n = 456, weaned individuals that have not reached sexual maturity). If
previous tube task studies assigned age categories to their subjects, we adopted this classification for
the individuals concerned. In other cases and for our original data, age classification followed life
history data from Harvey and Clutton-Brock (1985). The taxonomy and nomenclature we apply

follows Mittermeier, Rylands, and Wilson (2013) with the following exceptions: The recently
diverging sister species pairs *Cercopithecus diana* and *C. roloway* (Diana and Roloway monkeys) as
well as *Nomascus leucogenys* and *N. siki* (white-cheeked gibbons) are treated here as one respective
taxonomic unit and data were pooled to obtain larger sample sizes (exact species identity of subjects
is annotated in the raw data). Because the hand preference literature on orangutans (*Pongo* spp.) did
not consider the species status of the individuals concerned, we analyzed respective data on the
genus level. In other cases, we carefully checked the current taxonomic status of subjects drawn
from the literature and tried to avoid the inclusion of interspecific hybrids. This was particularly
relevant for data on lab-housed tufted capuchins (*Sapajus* spp.). If the species or hybrid status of
animals was ambiguous, we did not consider them for our analyses (e.g., capuchins in Westergaard
and Suomi, 1996).

Although available for testing at most of the institutions we visited, lemurs could not be included into
the study. All tested genera (*Eulemur*, *Hapalemur*, *Propithecus*, *Varecia*), failed to manually remove
food mash from the tube, despite eagerly licking it up from the ends. Only a single subject, a female
Eulemur rubriventer, succeeded. White-faced sakis (*Pithecia pithecia*) and Javan langurs
(*Trachypithecus auratus*) were also reluctant to engage in the task, so that the final sample for these
species is smaller than expected from their abundance at the institutions visited. Apart from the
species considered for analysis, nine individuals belonging to miscellaneous taxa were sampled in the
study.

All experimental procedures strictly adhered to the guidance of the responsible zoo staff, were
approved by the institutional boards in charge and complied to the applicable animal welfare and
testing regulations of the countries they were performed in. No further ethical permissions had to be
obtained.

Experimental procedure and data scoring

All species were uniformly tested in the established bimanual tube task paradigm (Hopkins, 1995).
Due to the pronounced differences in body size between the observed species, PVC tubes of varying
length and diameter were employed (Figure 4). Lion tamarins (*Leontopithecus*) were presented with
small-sized tubes that were 5 cm long and had an inner diameter of 1 cm. Capuchins and sakis
(*Sapajus*, *Pithecia*) received 10 cm x 2 cm medium-sized tubes and all remaining species large tubes
measuring 10 cm x 2.5 cm. The tubes were filled with various food incentives, which differed
dependent on the nutrition regimes enacted by the respective institutions. Among preferred food
items for cercopithecines, gibbons, spider monkeys, and capuchins were oatmeal mixed with banana
mash, soaked pellets, and boiled carrots (but note that the latter did not appeal to *Cercopithecus* and
Ateles). Geladas (*Theropithecus gelada*) exclusively received boiled carrots. Langurs (*Semnopithecus*,
Trachypithecus) and sakis were preferably tested with boiled rice, and tubes for the latter were also
stowed with nuts as an additional incentive. Lion tamarins received tubes filled with pure banana
mash or commercial tamarin pie. Primates were preferably tested within their social groups. A
separation of individuals was only undertaken in exceptional cases when it was necessary to
counteract social tension created by the presentation of the tubes. Dependent on the constructional
restraints of the enclosures, tubes were either placed into a separated part of the enclosure before
the primates could enter or were handed over directly through the wire mesh. In the latter case, the
hand that the experimenter used to offer the tube was noted. To check whether the hand used by
the experimenter to offer the tube had an effect on the directional hand preferences of the tested
primates, we ran a linear mixed effect model employing a binomial link function. No effect on the
recovered hand preferences in the respective sessions was found (t = -1.31, SE = 0.02, *p* = 0.191).

The tube tasks were recorded with digital cameras and scored from the video footage. For each
subject, we obtained a minimum of 30 bimanual insertions (one hand is holding the tube, the other
one is retrieving food; mean number of insertions: 50.66 ± 20.98, range: 30-155) in at least six bouts
(uninterrupted manipulation sequences, as defined by Morino et al. (2017); mean number of bouts:
20.60 ± 11.13, range: 6-82). Literature data for individual subjects had to match or exceed these
thresholds to be included into the analysis. Unimanual or foot-assisted insertions were not scored
and were, whenever possible, also carefully excluded from the literature data. We noted the digit
used to extract the food as well as the body posture of the manipulating individual (sitting, crouched
bipedal stance, erect bipedal stance, suspended (clinging to wire mesh or other substrates without
the hands stabilizing posture, always tail-assisted in spider monkeys)). The vast majority of responses
were observed in a sitting position (n = 22,993; 90.6 %). Due to this imbalance, because posture-
related information was mostly unavailable for literature data, and since its influence on manual
laterality already received great research attention in the past (Sanford et al., 1984; Westergaard et
al., 1998; Blois-Heulin et al., 2007; Laurence et al., 2011), we refrained from including posture effects
into our analyses. Nevertheless, for potential future use by other researchers, we decided to include
this measure, alongside information on digit use ("N.A." if ambiguous in the respective footage)
during manipulation, alongside the respective raw data.

Table 4: Composition of the study sample, listing taxonomic identity, sex, age, and origin of subjects. See cited studies for locations of individuals drawn from
the literature.

Family	Species	# Subjects tested	# Subjects drawn from literature*	Total sample	# adult females	# adult males	# subadult females	# subadult males	# unsexed subadults	Locations for subjects in this study
Atelidae	*Ateles fusciceps*	37	9[a]	46	30	11	3	2	0	Berlin (Zoo), Doué-la-Fontaine, Landau, Mulhouse, Munich, Osnabrück, Wuppertal
Atelidae	*Ateles geoffroyi*	9	14[b]	23	12	9	0	2	0	Basel, Karlsruhe
Atelidae	*Ateles hybridus*	18		18	10	7	0	1	0	Doué-la-Fontaine, Erfurt, Frankfurt, Neuwied, Stuttgart
Callitrichidae	*Leontopithecus chrysomelas*	30		30	11	16	2	1	0	Apeldoorn, Karlsruhe, Magdeburg, Mulhouse, São Paulo, Stuttgart, Wuppertal
Callitrichidae	*Leontopithecus chrysopygus*	15		15	6	9	0	0	0	São Paulo
Callitrichidae	*Leontopithecus rosalia*	28		28	7	16	0	5	0	Apeldoorn, Basel, Doué-la-Fontaine, Duisburg, Frankfurt, Heidelberg, Landau, Magdeburg, Münster, São Paulo
Cebidae	*Saimiri sciureus*		36[c]	36	14	16	5	1	0	
Cebidae	*Sapajus apella*		25[d, e]	25	10	11	0	4	0	
Cebidae	*Sapajus flavius*	3	18[e]	21	7	9	2	3	0	São Paulo
Cebidae	*Sapajus xanthosternos*	16	18[e]	34	11	19	1	2	1	Apeldoorn, Magdeburg, Münster, Overloon
Cercopithecidae	*Cercocebus torquatus*	18	13[f]	31	15	13	1	2	0	Apeldoorn, Berlin (Tierpark), Karlsruhe, Münster
Cercopithecidae	*Cercopithecus diana/roloway*	20		20	9	7	3	1	0	Amsterdam, Berlin (Tierpark), Doué-la-Fontaine, Duisburg, Heidelberg, Liberec, Mulhouse, Osnabrück
Cercopithecidae	*Cercopithecus neglectus*	12	13[f, g]	25	8	12	1	4	0	Bekesbourne, Duisburg, Hannover, Overloon
Cercopithecidae	*Macaca fascicularis*	12	8[h]	20	13	7	0	0	0	Basel, Hamm, Mönchengladbach
Cercopithecidae	*Macaca nemestrina*	29		29	12	15	0	1	1	Arnhem, Bali, Berlin (Tierpark), Gelsenkirchen, Osnabrück
Cercopithecidae	*Macaca silenus*	35		35	16	17	1	1	0	Apeldoorn, Bekesbourne, Berlin (Zoo), Cologne, Dresden, Duisburg, Hodenhagen, Rheine
Cercopithecidae	*Macaca sylvanus*	15	9[i, j]	24	11	12	0	1	0	Aachen, Rheine
Cercopithecidae	*Macaca tonkeana***		14[k]	14	NA	NA	NA	NA	NA	
Cercopithecidae	*Mandrillus sphinx*	32		32	14	7	4	7	0	Amsterdam, Berlin (Zoo), Dresden, Hamm, Hodenhagen
Cercopithecidae	*Papio anubis*		84[l]	84	48	22	5	9	0	
Cercopithecidae	*Papio hamadryas*	24		24	14	10	0	0	0	Cologne, Frankfurt, Krefeld
Cercopithecidae	*Pygathrix cinerea*		18[m]	18	7	11	0	0	0	
Cercopithecidae	*Rhinopithecus roxellana*		24[n]	24	8	5	8	3	0	

Family	Species									Location
Cercopithecidae	*Semnopithecus entellus*	30		30	17	7	4	2	0	Apeldoorn, Berlin (Zoo), Gelsenkirchen, Hannover, Heidelberg
Cercopithecidae	*Theropithecus gelada*	38		38	20	11	4	3	0	Bekesbourne, Berlin (Tierpark), Magdeburg, Rheine, Stuttgart
Cercopithecidae	*Trachypithecus auratus*	8		8	3	0	3	2	0	Bali, Stuttgart
Cercopithecidae	*Trachypithecus hatinhensis*		18[m]	18	8	10	0	0	0	
Hominidae	*Gorilla gorilla*		76[o]	76	22	18	19	17	0	
Hominidae	*Homo sapiens*		127[p]	127	71	56	0	0	0	
Hominidae	*Pan paniscus*		118[o]	118	29	23	35	31	0	
Hominidae	*Pan troglodytes*		536[o]	536	186	138	110	102	0	
Hominidae	*Pongo* sp.		47[o]	47	17	12	9	9	0	
Hylobatidae	*Hylobates lar*	16	20[q, r, s]	36	14	18	2	2	0	Berlin (Tierpark), Cologne, Doué-la-Fontaine, Landau, Stuttgart, Ulm, Wuppertal
Hylobatidae	*Hylobates moloch*	22		22	8	5	4	5	0	Bekesbourne, Lympne, Munich
Hylobatidae	*Nomascus gabriellae*	6	4[r]	10	5	3	0	2	0	Arnhem, Doué-la-Fontaine
Hylobatidae	*Nomascus leucogenys /siki*	7	19[q,r,t]	26	15	7	1	3	0	Apeldoorn, Frankfurt, Osnabrück
Hylobatidae	*Symphalangus syndactylus*	14	17[q,r]	31	12	11	4	4	0	Arnhem, Bekesbourne, Berlin (Zoo), Dortmund, Doué-la-Fontaine, Munich, Arnhem, Hodenhagen, Osnabrück
Pitheciidae	*Pithecia pithecia*	7		7	4	3	0	0	0	Basel, Dresden, Krefeld
Total***		**501**	**1285**	**1786**	**724**	**583**	**231**	**232**	**2**	

* Fulfilling our criteria

** Ages unknown, sex derived from given names

*** Not including *M. tonkeana* in sex and age specific categories

[a] Nelson and Boeving (2015), [b]Motes Rodrigo et al. (2018), [c]Meguerditchian et al. (2012), [d]Phillips et al. (2007); [e]de Andrade and de Sousa (2018), [f]Maille et al. (2013),

[g]Schweitzer et al. (2007), [h]Chatagny et al. (2013), [i]Schmitt et al. (2008), [j]Regaiolli et al. (2018), [k]Canteloup et al. (2013), [l]Vauclair et al. (2005), [m]Cubí and Llorente (2021), [n]Zhao

et al. (2012), [o]Hopkins et al. (2011), [p]Cochet and Vauclair (2012), [q]Morino et al. (2017), [r]Caspar et al. (2018), [s]Spoelstra (2021), [t]Fan et al. (2017).

Figure 4: Various anthropoid primates engaging in the tube task. A: Golden lion tamarin
(*Leontopithecus rosalia*) manipulating a small tube at Zoo Frankfurt, Germany. B: White-handed
gibbon (*Hylobates lar*) handling a large tube at Bioparc de Doué-la-Fontaine, France. Note that the
thumb is used to probe into the tube, an insertion pattern characteristic of gibbons. C: Yellow-
breasted capuchin (*Sapajus xanthosternos*) engaging in the task with a medium-sized tube while
adopting an erect bipedal stance at ZooParc Overloon, the Netherlands. Photographs by Kai R.
Caspar.

Statistics

Data were analyzed in R (R Core Team, 2020). Preferably, analyses were performed on insertion data
(also called frequencies) instead of manipulations bouts to approximate laterality. Unfortunately, not
all available tube task studies provided insertion data (Maille et al., 2013; Spoelstra, 2021), so that in
the final dataset, estimates of manual lateralization based on insertions and bouts had to be mixed
for certain species (*Cercocebus torquatus*, *Cercopithecus neglectus*, *Hylobates lar*, *Nomascus*
leucogenys, and *Sapajus* spp.). However, since previous work demonstrated that hand preferences
derived from bouts and insertions are highly correlated and non-conflicting, we do not consider this a
confounding factor for our analysis (Hopkins et al., 2001; Hopkins, 2013a).

For quantifying lateralized responses on the individual level, we calculated handedness indices (HI)
for all subjects as well as the corresponding binomial z scores to allow grouping into hand preference
categories. HI is a descriptive index that can range from -1 (all manipulations left-handed) to 1 (all
right-handed) and is calculated via the formula HI = (R − L)/ (R+L). The z score, on the other hand,
indicates whether there is a statistically significant bias in hand use. Following established criteria
(Hopkins, 2013a), we rated subjects with z score values higher than 1.96 as right-handed, those with
values lower than −1.96 as left-handed, and the remaining ones as ambipreferent. We use the term
"ambipreferent" here instead of "ambidextrous" or "ambiguously handed" to indicate a lack of
preferences, because the latter two expressions have clear non-synonymous definitions when

applied to human, but are not consistently used in the non-human primate literature (Hopkins et al.,
2013). At the species level, we used the mean HI of subjects as a measure of lateralization direction
and the mean of absolute HI values (MeanAbsHI) as a measure of strength.

We applied one-sample t-tests to each species and genus sample encompassing data for at least 15
individuals to check whether HI distributions were significantly skewed at the population-level.
Additionally, the chi-square goodness-of-fit test was employed to test if the numbers of left- and
right-handers as well as ambipreferent individuals differed from a baseline distribution. Earlier
studies performed the goodness-of-fit test against the null hypothesis of a chance distribution of the
three hand preference categories (Vauclair et al., 2005). Due to our large multispecies sample, we
could adopt a different approach: For each of the three major clades studied (Cercopithecoidea,
Hominoidea, and Platyrrhini), we calculated the mean frequencies of individuals being either
ambipreferent or handed. We then assumed an equal probability of handed individuals to be either
right or left-handed and this way calculated a clade-specific baseline. Distributions for each species
were then compared to this clade-specific average. For the Hominoidea, we excluded humans for the
calculation of the baseline, because their evidently extreme lateralization bias would have otherwise
skewed the results. Bonferroni correction was employed to address multiple testing.

We employed the phytools package version 0.7-70 (Revell, 2012) to visualize evolutionary patterns,
quantify phylogenetic signals (by employing Pagel's λ – Freckleton et al., 2002; the null hypothesis of
no phylogenetic signal was tested by means of the likelihood ratio test) and to calculate maximum
likelihood ancestral state estimates (Suppl. File. 1), each separately for direction and strength of
lateralization. Time-calibrated primate phylogenies were derived from the 10kTrees website (Arnold
et al., 2010). Three species in our study were not included in the respective database, *Ateles*
hybridus, *Sapajus flavius*, and *Trachypithecus hatinhensis*. We therefore replaced *A. hybridus* and *T.*
hatinhensis in the tree with respective sister taxa, namely *A. belzebuth* and *T. francoisi* (Morales-
Jimenez et al., 2015; Roos et al., 2019), for which data were provided. This way the topology and
branch lengths of the tree could be kept equal to a model which would have included the actual
species we studied. The same could not be done for *S. flavius*, so that it was amended manually in
the respective trees by relying on divergence dating from Lima et al. (2018).

We computed phylogenetic generalized least squares (PGLS) regression models to test the effects of
different biological variables on hand preferences while addressing phylogeny (correlation structure:
Pagel's λ; model fit: maximum likelihood). The R packages *ape* (Paradis, 2019), *nlme* (Pinheiro, 2020),
and *MuMIn* (Barton, 2020) were used for model creation and evaluation. We used multi-model
inference to test how well hypothesis-derived predictors could explain HI measures on the species
level. Predictor-based models were ranked against a NULL model to estimate their explanatory
power and identify the best-performing one (*dredge* function in *MuMIn*). Second order Akaike's
information criteria (AICc) and respective Akaike weights were used to evaluate model components.
The normality of model residuals was checked by applying the Shapiro-Wilk test. We relied on the
conditional model average to assess effects of individual predictors. We selected the following
variables as model predictors to address established hypotheses on the evolution on primate hand
preferences: Ecology (terrestrial vs. arboreal), occurrence of habitual foraging-related tool use in the
wild (binarily coded), and endocranial volume (numeric, log-transformed) of females (see Suppl. File
3 for predictor data and respective references). A potential multicollinearity of predictors was
checked by computing their variable inflation factors. We ran the models on the species means for HI
and AbsHI, respectively, resulting in two separate analyses for direction and strength of population-
level laterality. All species (*n* = 38) were included into the strength (AbsHI) analysis. For direction, we
only considered species for which we had at least 15 sampled individuals, resulting in a more
restricted sample (*n* = 34). Since humans are both extreme outliers in regard to absolute brain size

and hand preference direction, we decided to compute a second direction model to identify
potential biases that might derive from their inclusion. Thus, this second model on hand preference
direction encompassed a species sample of n = 33.

Finally, we employed Bayesian phylogenetic multilevel models by aid of the *brms* package (Bürkner,
2017) to check whether individual HI and AbsHI were influenced by age and sex (excluding unsexed
individuals; n = 2). Analyses were run for the complete sample as well as within the three
superordinate clades, thus resulting in a total of eight models, four for each of the two response
variables. We assumed a notable effect on lateralization patterns when the model's 95% credible
intervals of intercept and respective regression coefficients did not overlap with zero. Default priors
were used. Modelling results, including the number of chains and iterations performed are
summarized and visualized in Supplementary File 2. Due to their highly derived handedness patterns,
we again excluded humans from these analyses to avoid skewing the results. To further avoid bias,
we also removed chimpanzees from the models, since they are vastly overrepresented in our sample
(31% of total individuals and 47% of all subadults). Finally, as no age data on Tonkean macaques
(*Macaca tonkeana*) were available, this species was not featured in these analyses as well. This left
us with n = 1,107 individuals in the full dataset, n = 366 in the hominoid, n = 459 in the
cercopithecoid and n = 282 in the platyrrhine subsample.

Supplementary Material

Suppl. File 1: Estimates of ancestral manual lateralization patterns in anthropoid taxa.

Suppl. File 2: Results and visualization of Bayesian models to infer effect of sex and age cohort on
individual-level hand preference patterns

Suppl. File 3: Species-level predictors used for PGLS modelling.

Data Availability

Raw data on manual lateralization used and/or generated in this study have been made available
through the public repository Dryad as detailed in the transparent reporting form. The link for
accessing these datasets is: https://doi.org/10.5061/dryad.8sf7m0crv

The R code for the statistical analyses presented here can be accessed via:
https://github.com/fpallasdies/CasparEtAl2022PrimateHandPreference.git

Acknowledgements

We would like to thank the curators and animal keepers at the following institutions for the
permission to study their animals and for their support during data collection:

Allwetterzoo Münster, Apenheul (Apeldoorn), Artis Amsterdam, Bali Wildlife Rescue Center, Bioparc
de Doué-la-Fontaine, Burger's Zoo (Arnhem), Erlebnis-Zoo Hannover, Howletts Wild Animal Park
(Bekesbourne), Naturzoo Rheine, Parc Zoologique et Botanique de Mulhouse, Port Lympne Wild
Animal Park (Lympne), Serengeti-Park Hodenhagen, Tierpark Aachen, Tierpark Berlin, Tierpark
Hamm, Tierpark Hellabrunn (Munich), Tierpark Mönchengladbach, Tierpark Ulm, Wilhelma Stuttgart,
Zoo Basel, Zoo Berlin, Zoo Dortmund, Zoo Dresden, Zoo Duisburg, Zoo Frankfurt, Zoo Heidelberg, Zoo
Köln (Cologne), Zoo Krefeld, Zoo Landau in der Pfalz, Zoo Liberec, Zoo Magdeburg, Zoo Neuwied, Zoo

Osnabrück, Zoo Wuppertal, Zoológico de São Paulo, Zoologischer Stadtgarten Karlsruhe, ZOOM Erlebniswelt Gelsenkirchen, ZooParc Overloon, and Zoopark Erfurt.

We are also indebted to Hélène Cochet, Julia Fischer, William D. Hopkins, Adrien Meguerditchian, and Luca Morino for sharing raw data on bimanual task preferences in humans, barbary macaques, great apes, olive baboons, and gibbons, respectively.

Daniel Issel is greatly acknowledged for collecting data on pig-tailed macaques and Javan langurs in Bali and Petra Bolechová for doing so for Diana monkeys at Zoo Liberec. Finally, we would like to thank Miriam Lindenmeier and Larissa Günther for crucial assistance during data collection at various locations and Alex DeCasien as well as Adrien Meguerditchian for their constructive and insightful reviews on earlier versions of this manuscript.

This research was funded by the German Society for Mammalian Biology (Deutsche Gesellschaft für Säugetierkunde). KRC was supported by a Ph.D. fellowship of the German Academic Scholarship Foundation (Studienstiftung des deutschen Volkes e.V.).

References

Amunts, K., Schlaug, G., Schleicher, A., Steinmetz, H., Dabringhaus, A., Roland, P. E., & Zilles, K. (1996). Asymmetry in the human motor cortex and handedness. *NeuroImage, 4*(3), 216-222. doi:https://doi.org/10.1006/nimg.1996.0073

Ankel-Simons, F., & Rasmussen, D. T. (2008). Diurnality, nocturnality, and the evolution of primate visual systems. *American Journal of Physical Anthropology, 137*(S47), 100-117. doi:https://doi.org/10.1002/ajpa.20957

Annett, M. (2002). *Handedness and Brain Asymmetry: The Right Shift Theory*. Hove: Psychology Press.

Ardesch, D. J., Scholtens, L. H., de Lange, S. C., Roumazeilles, L., Khrapitchev, A. A., Preuss, T. M., . . . van den Heuvel, M. P. (2021). Scaling principles of white matter connectivity in the human and nonhuman primate brain. *Cerebral Cortex*, bhab384. doi:10.1093/cercor/bhab384

Arnold, C., Matthews, L. J., & Nunn, C. L. (2010). The 10kTrees website: A new online resource for primate phylogeny. *Evolutionary Anthropology: Issues, News, and Reviews, 19*(3), 114-118. doi:https://doi.org/10.1002/evan.20251

Barrett, R. L. C., Dawson, M., Dyrby, T. B., Krug, K., Ptito, M., Arceuil, H., . . . Catani, M. (2020). Differences in frontal network anatomy across primate species. *The Journal of Neuroscience, 40*(10), 2094. doi:10.1523/JNEUROSCI.1650-18.2019

Barton, K. (2020). MuMIn: Multi-model inference. R package version 1.43.17. Retrieved from https://CRAN.R-project.org/package=MuMIn

Batist, C. H., & Mayhew, J. A. (2020). Lateralization in seven lemur species when presented with a novel cognitive task. *American Journal of Physical Anthropology, 172*(2), 270-279. doi:https://doi.org/10.1002/ajpa.24037

Becker, Y., Claidière, N., Margiotoudi, K., Marie, D., Roth, M., Nazarian, B., Anton, J. L., Coulon, O., & Meguerditchian, A. (2022). Broca's cerebral asymmetry reflects gestural communication's lateralisation in monkeys (*Papio anubis*). *eLife*, *11*, e70521. https://doi.org/10.7554/eLife.70521

Blois-Heulin, C., Bernard, V., & Bec, P. (2007). Postural effect on manual laterality in different tasks in captive grey-cheeked mangabey (*Lophocebus albigena*). *Journal of Comparative Psychology, 121*(2), 205-213. doi:10.1037/0735-7036.121.2.205

Brown, C., & Magat, M. (2011). The evolution of lateralized foot use in parrots: a phylogenetic approach. *Behavioral Ecology, 22*(6), 1201-1208. doi:10.1093/beheco/arr114

Bürkner, P. (2017). brms: an R package for Bayesian multilevel models using stan. *Journal of Statistical Software*, 80(1), 1–28. doi: 10.18637/jss.v080.i01.

Canteloup, C., Vauclair, J., & Meunier, H. (2013). Hand preferences on unimanual and bimanual tasks in Tonkean macaques (*Macaca tonkeana*). *American Journal of Physical Anthropology, 152*(3), 315-321. doi:https://doi.org/10.1002/ajpa.22342

Cashmore, L., Uomini, N., & Chapelain, A. (2008). The evolution of handedness in humans and great apes: a review and current issues. *Journal of Anthropological Sciences 86*, 7-35.

Caspar, K. R., Mader, L., Pallasdies, F., Lindenmeier, M., & Begall, S. (2018). Captive gibbons (Hylobatidae) use different referential cues in an object-choice task: insights into lesser ape cognition and manual laterality. *PeerJ, 6*, e5348. doi:10.7717/peerj.5348

Chatagny, P., Badoud, S., Kaeser, M., Gindrat, A.-D., Savidan, J., Fregosi, M., . . . Rouiller, E. M. (2013). Distinction between hand dominance and hand preference in primates: a behavioral investigation of manual dexterity in nonhuman primates (macaques) and human subjects. *Brain and Behavior, 3*(5), 575-595. doi:https://doi.org/10.1002/brb3.160

Cochet, H., & Vauclair, J. (2012). Hand preferences in human adults: non-communicative actions versus communicative gestures. *Cortex, 48*(8), 1017-1026. doi:https://doi.org/10.1016/j.cortex.2011.03.016

Corballis, M. C. (1987). Straw monkeys. *Behavioral and Brain Sciences, 10*(2), 269-270. doi:10.1017/S0140525X00047762

Corballis, M. C. (1991). *The Lopsided Ape: Evolution of the Generative Mind*. Oxford: Oxford University Press.

Coren, S., & Porac, C. (1977). Fifty centuries of right-handedness: the historical record. *Science, 198*(4317), 631. doi:10.1126/science.335510

Cubí, M., & Llorente, M. (2021). Hand preference for a bimanual coordinated task in captive hatinh langurs (*Trachypithecus hatinhensis*) and grey-shanked douc langurs (*Pygathrix cinerea*). *Behavioural Processes, 187*, 104393. doi:https://doi.org/10.1016/j.beproc.2021.104393

Dadda, M., Cantalupo, C., & Hopkins, W. D. (2006). Further evidence of an association between handedness and neuroanatomical asymmetries in the primary motor cortex of chimpanzees (*Pan troglodytes*). *Neuropsychologia, 44*(12), 2582-2586. doi:https://doi.org/10.1016/j.neuropsychologia.2006.03.037

de Andrade, A. C., & de Sousa, A. B. (2018). Hand preferences and differences in extractive foraging in seven capuchin monkey species. *American Journal of Primatology, 80*(8), e22901. doi:https://doi.org/10.1002/ajp.22901

Diamond, A. C., & McGrew, W. C. (1994). True handedness in the cotton-top tamarin (*Saguinus oedipus*)? *Primates, 35*(1), 69-77. doi:10.1007/BF02381487

Fan, P., Liu, C., Chen, H., Liu, X., Zhao, D., Zhang, J., & Liu, D. (2017). Preliminary study on hand preference in captive northern white-cheeked gibbons (*Nomascus leucogenys*). *Primates, 58*(1), 75-82. doi:10.1007/s10329-016-0573-8

Faurie, C., Schiefenhövel, W., le Bomin, S., Billiard, S., & Raymond, M. (2005). Variation in the frequency of left-handedness in traditional societies. *Current Anthropology, 46*(1), 142-147. doi:10.1086/427101

Fitch, W. T., & Braccini, S. N. (2013). Primate laterality and the biology and evolution of human handedness: a review and synthesis. *Annals of the New York Academy of Sciences, 1288*(1), 70-85. doi:https://doi.org/10.1111/nyas.12071

Freckleton, R. P., Harvey, P. H., & Pagel, M. (2002). Phylogenetic analysis and comparative data: a test and review of evidence. *The American Naturalist, 160*(6), 712-726. doi:10.1086/343873

Frost, G. (1980). Tool behavior and the origins of laterality. *Journal of Human Evolution, 9*, 447-459.

Gebo, D. L. (2011). Vertical clinging and leaping revisited: vertical support use as the ancestral condition of strepsirrhine primates. *American Journal of Physical Anthropology, 146*(3), 323-335. doi:https://doi.org/10.1002/ajpa.21595

Giljov, A., Karenina, K., Ingram, J., & Malashichev, Y. (2015). Parallel emergence of true handedness in the evolution of marsupials and placentals. *Current Biology, 25*(14), 1878-1884. doi:https://doi.org/10.1016/j.cub.2015.05.043

Häberling, I. S., & Corballis, M. C. (2016). Cerebellar asymmetry, cortical asymmetry and handedness:
two independent networks. *Laterality, 21*(4-6), 397-414.
doi:10.1080/1357650X.2015.1110161
Harvey, P. H., & Clutton-Brock, T. H. (1985). Life history variation in primates. *Evolution, 39*(3), 559-
581. doi:https://doi.org/10.1111/j.1558-5646.1985.tb00395.x
Hecht, E. E., Gutman, D. A., Bradley, B. A., Preuss, T. M., & Stout, D. (2015a). Virtual dissection and
comparative connectivity of the superior longitudinal fasciculus in chimpanzees and humans.
NeuroImage, 108, 124-137. doi:https://doi.org/10.1016/j.neuroimage.2014.12.039
Hecht, E. E., Gutman, D. A., Khreisheh, N., Taylor, S. V., Kilner, J., Faisal, A. A., . . . Stout, D. (2015b).
Acquisition of Paleolithic toolmaking abilities involves structural remodeling to inferior
frontoparietal regions. *Brain Structure and Function, 220*(4), 2315-2331. doi:10.1007/s00429-
014-0789-6
Hopkins, W. D. (1995). Hand preferences for a coordinated bimanual task in 110 chimpanzees (*Pan
troglodytes*): cross-sectional analysis. *Journal of Comparative Psychology, 109*(3), 291-297.
doi:10.1037/0735-7036.109.3.291
Hopkins, W. D. (2013a). Comparing human and nonhuman primate handedness: challenges and a
modest proposal for consensus. *Developmental Psychobiology, 55*(6), 621-636.
doi:https://doi.org/10.1002/dev.21139
Hopkins, W. D. (2013b). Neuroanatomical asymmetries and handedness in chimpanzees (*Pan
troglodytes*): a case for continuity in the evolution of hemispheric specialization. *Annals of
the New York Academy of Sciences, 1288*(1), 17-35. doi:10.1111/nyas.12109
Hopkins, W. D., Adams, M. J., & Weiss, A. (2013). Genetic and environmental contributions to the
expression of handedness in chimpanzees (*Pan troglodytes*). *Genes, Brain and Behavior,
12*(4), 446-452. doi:https://doi.org/10.1111/gbb.12044
Hopkins, W. D., & Cantalupo, C. (2004). Handedness in chimpanzees (*Pan troglodytes*) is associated
with asymmetries of the primary motor cortex but not with homologous language areas.
Behavioral Neuroscience, 118(6), 1176-1183. doi:10.1037/0735-7044.118.6.1176
Hopkins, W. D., Fernandez-Carriba, S., Wesley, M. J., Hostetter, A., Pilcher, D., & Poss, S. (2001). The
use of bouts and frequencies in the evaluation of hand preferences for a coordinated
bimanual task in chimpanzees (*Pan troglodytes*): an empirical study comparing two different
indices of laterality. *Journal of Comparative Psychology, 115*(3), 294-299. doi:10.1037/0735-
7036.115.3.294
Hopkins, W. D., Gardner, M., Mingle, M., Reamer, L., & Schapiro, S. J. (2013). Within- and between-
task consistency in hand use as a means of characterizing hand preferences in captive
chimpanzees (*Pan troglodytes*). *Journal of Comparative Psychology, 127*(4), 380-391.
doi:10.1037/a0031071
Hopkins, W. D., Phillips, K. A., Bania, A., Calcutt, S. E., Gardner, M., Russell, J., . . . Schapiro, S. J.
(2011). Hand preferences for coordinated bimanual actions in 777 great apes: implications
for the evolution of handedness in hominins. *Journal of Human Evolution, 60*(5), 605-611.
doi:10.1016/j.jhevol.2010.12.008
Kaplan, G., & Rogers, L. J. (2021). Brain size associated with foot preferences in Australian parrots.
Symmetry, 13(5). doi:10.3390/sym13050867
Karolis, V. R., Corbetta, M., & Thiebaut de Schotten, M. (2019). The architecture of functional
lateralisation and its relationship to callosal connectivity in the human brain. *Nature
Communications, 10*(1), 1417. doi:10.1038/s41467-019-09344-1
Kimura, D. (1979). Neuromotor mechanisms in the evolution of human communication. In H. D. S. M.
Paleigh (Ed.), *Neurobiology of social communication in primates: An evolutionary perspective*
(pp. 1197-1220). Cambridge: Academic Press.
Kitchener, A. C., Meloro, C., Williams, T.M. (2017). Form and function of the musteloids. In D. W.
Macdonald, Newman, C., Harrington, L. (Ed.), *The Biology and Conservation of Wild
Musteloids* (pp. 98-135). Oxford: Oxford University Press.
Kong, X.-Z., Mathias, S. R., Guadalupe, T., Group, E. L. W., Glahn, D. C., Franke, B., . . . Francks, C.
(2018). Mapping cortical brain asymmetry in 17,141 healthy individuals worldwide via the

ENIGMA Consortium. *Proceedings of the National Academy of Sciences, 115*(22), E5154.
doi:10.1073/pnas.1718418115

Krasheninnikova, A., Chow, P. K. Y., & von Bayern, A. M. P. (2020). Comparative cognition: practical
shortcomings and some potential ways forward. *Canadian Journal of Experimental*
Psychology/Revue canadienne de psychologie expérimentale, 74(3), 160-169.
doi:10.1037/cep0000204

Laska, M. (1996). A study of correlates of hand preferences in squirrel monkeys (*Saimiri sciureus*).
Primates, 37(4), 457-465. doi:10.1007/BF02381379

Laurence, A., Wallez, C., & Blois-Heulin, C. (2011). Task complexity, posture, age, sex: Which is the
main factor influencing manual laterality in captive *Cercocebus torquatus torquatus*?
Laterality, 16(5), 586-606. doi:10.1080/1357650X.2010.501338

Lilak, A. L., & Phillips, K. A. (2008). Consistency of hand preference across low-level and high-level
tasks in Capuchin monkeys (Cebus apella). *American Journal of Primatology, 70*(3), 254-260.
doi:https://doi.org/10.1002/ajp.20485

Lima, M. G. M., Silva-Júnior, J. d. S. e., Černý, D., Buckner, J. C., Aleixo, A., Chang, J., . . . Lynch Alfaro,
J. W. (2018). A phylogenomic perspective on the robust capuchin monkey (*Sapajus*)
radiation: First evidence for extensive population admixture across South America. *Molecular*
Phylogenetics and Evolution, 124, 137-150. doi:https://doi.org/10.1016/j.ympev.2018.02.023

Lozano, M., Estalrrich, A., Bondioli, L., Fiore, I., Bermúdez de Castro, J.-M., Arsuaga, J. L., . . . Frayer, D.
W. (2017). Right-handed fossil humans. *Evolutionary Anthropology: Issues, News, and*
Reviews, 26(6), 313-324. doi:https://doi.org/10.1002/evan.21554

MacLean, E. L., Matthews, L. J., Hare, B. A., Nunn, C. L., Anderson, R. C., Aureli, F., . . . Wobber, V.
(2012). How does cognition evolve? Phylogenetic comparative psychology. *Animal cognition,*
15(2), 223-238. doi:10.1007/s10071-011-0448-8

MacNeilage, P. F. (2007). Present status of the postural origins theory. In *The evolution of*
hemispheric specialization in primates, W. D. Hopkins (Ed.) American Society of
Primatologists, pp. 59–91. Oxford: Elsevier.

MacNeilage, P. F., Studdert-Kennedy, M. G., & Lindblom, B. (1987). Primate handedness
reconsidered. *Behavioral and Brain Sciences, 10*(2), 247-263.
doi:10.1017/S0140525X00047695

Maille, A., Belbeoc'h, C., Rossard, A., Bec, P., & Blois-Heulin, C. (2013). Which are the features of the
TUBE task that make it so efficient in detecting manual asymmetries? An investigation in two
Cercopithecine species (*Cercopithecus neglectus* and *Cercocebus torquatus*). *Journal of*
Comparative Psychology, 127(4), 436-444. doi:10.1037/a0032227

ManyPrimates, Altschul, D. M., Beran, M. J., Bohn, M., Caspar, K. R., Fichtel, C., . . . Grebe, N. M.
(2019). Collaborative open science as a way to reproducibility and new insights in primate
cognition research. *Japanese Psychological Review, 62*(3), 205-220.

Marchant, L. F., & McGrew, W. C. (2013). Handedness is more than laterality: lessons from
chimpanzees. *Annals of the New York Academy of Sciences, 1288*(1), 1-8.
doi:https://doi.org/10.1111/nyas.12062

Margiotoudi, K., Marie, D., Claidière, N., Coulon, O., Roth, M., Nazarian, B., . . . Meguerditchian, A.
(2019). Handedness in monkeys reflects hemispheric specialization within the central sulcus.
An in vivo MRI study in right- and left-handed olive baboons. *Cortex, 118*, 203-211.
doi:https://doi.org/10.1016/j.cortex.2019.01.001

Meguerditchian, A., Donnot, J., Molesti, S., Francioly, R., & Vauclair, J. (2012). Sex difference in
squirrel monkeys' handedness for unimanual and bimanual coordinated tasks. *Animal*
Behaviour, 83(3), 635-643. doi:https://doi.org/10.1016/j.anbehav.2011.12.005

Meguerditchian, A., Phillips, K. A., Chapelain, A., Mahovetz, L. M., Milne, S., Stoinski, T., . . . Hopkins,
W. D. (2015). Handedness for unimanual grasping in 564 great apes: the effect on grip
morphology and a comparison with hand use for a bimanual coordinated task. *Frontiers in*
Psychology, 6, 1794-1794. doi:10.3389/fpsyg.2015.01794

Meguerditchian, A., Vauclair, J., & Hopkins, W. D. (2013). On the origins of human handedness and
language: A comparative review of hand preferences for bimanual coordinated actions and

gestural communication in nonhuman primates. *Developmental Psychobiology, 55*(6), 637-
650. doi:https://doi.org/10.1002/dev.21150
Mittermeier, R. A., Rylands, A. B., & Wilson, D. E. (2013). *Handbook of the Mammals of the World: 3.*
Primates. Barcelona: Lynx Ediciones.
Morales-Jimenez, A. L., Disotell, T., & Di Fiore, A. (2015). Revisiting the phylogenetic relationships,
biogeography, and taxonomy of spider monkeys (genus *Ateles*) in light of new molecular
data. *Molecular Phylogenetics and Evolution, 82*, 467-483.
doi:https://doi.org/10.1016/j.ympev.2014.09.019
Morino, L., Uchikoshi, M., Bercovitch, F., Hopkins, W. D., & Matsuzawa, T. (2017). Tube task hand
preference in captive hylobatids. *Primates, 58*(3), 403-412. doi:10.1007/s10329-017-0605-z
Motes Rodrigo, A., Ramirez Torres, C. E., Hernandez Salazar, L. T., & Laska, M. (2018). Hand
preferences in two unimanual and two bimanual coordinated tasks in the black-handed
spider monkey (*Ateles geoffroyi*). *Journal of Comparative Psychology, 132*(2), 220-229.
doi:10.1037/com0000110
Musgrave, S., & Sanz, C. (2018). Tool use in nonhuman primates. In *The International Encyclopedia of*
Anthropology, H. Callan (Ed.). doi:https://doi.org/10.1002/9781118924396.wbiea2063
Nelson, E. L., & Boeving, E. R. (2015). Precise digit use increases the expression of handedness in
Colombian spider monkeys (*Ateles fusciceps rufiventris*). *American Journal of Primatology,*
77(12), 1253-1262. doi:https://doi.org/10.1002/ajp.22478
Ocklenburg, S., Beste, C., Arning, L., Peterburs, J., & Güntürkün, O. (2014). The ontogenesis of
language lateralization and its relation to handedness. *Neuroscience & Biobehavioral*
Reviews, 43, 191-198. doi:https://doi.org/10.1016/j.neubiorev.2014.04.008
Papademetriou, E., Sheu, C.-F., & Michel, G. F. (2005). A meta-analysis of primate hand preferences,
particularly for reaching. *Journal of Comparative Psychology, 119*(1), 33-48.
doi:10.1037/0735-7036.119.1.33
Paradis, E. S., K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary analyses
in R. *Bioinformatics, 35*, 526-528.
Phillips, K. A., & Sherwood, C. C. (2005). Primary motor cortex asymmetry is correlated with
handedness in capuchin monkeys (*Cebus apella*). *Behavioral Neuroscience, 119*(6), 1701-
1704. doi:10.1037/0735-7044.119.6.1701
Phillips, K. A., Sherwood, C. C., & Lilak, A. L. (2007). Corpus callosum morphology in capuchin
monkeys is influenced by sex and handedness. *PLOS ONE, 2*(8), e792.
doi:10.1371/journal.pone.0000792
Pinheiro J., B. D., DebRoy S., Sarkar D., R Core Team. (2020). nlme: linear and nonlinear mixed effects
models. R package version 3.1-148. Retrieved from https://CRAN.R-
project.org/package=nlme
Prieur, J., Lemasson, A., Barbu, S., & Blois-Heulin, C. (2019). History, development and current
advances concerning the evolutionary roots of human right-handedness and language: Brain
lateralisation and manual laterality in non-human primates. *Ethology, 125*(1), 1-28.
doi:https://doi.org/10.1111/eth.12827
Ramayya, A. G., Glasser, M. F., & Rilling, J. K. (2010). A DTI investigation of neural substrates
supporting tool use. *Cerebral Cortex, 20*(3), 507-516. doi:10.1093/cercor/bhp141
Raymond, M., & Pontier, D. (2004). Is there geographical variation in human handedness? *Laterality,*
9(1), 35-51. doi:10.1080/13576500244000274
R Core Team (2020). R: A language and environment for statistical computing. Vienna: R Foundation
for Statistical Computing. Retrieved from https://www.R-project.org/
Regaiolli, B., Spiezio, C., & Hopkins, W. D. (2016). Hand preference on unimanual and bimanual tasks
in strepsirrhines: The case of the ring-tailed lemur (*Lemur catta*). *American Journal of*
Primatology, 78(8), 851-860. doi:https://doi.org/10.1002/ajp.22549
Regaiolli, B., Spiezio, C., & Hopkins, W. D. (2018). Hand preference on unimanual and bimanual tasks
in Barbary macaques (*Macaca sylvanus*). *American Journal of Primatology, 80*(3), e22745.
doi:https://doi.org/10.1002/ajp.22745

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other things).
Methods in Ecology and Evolution, 3(2), 217-223. doi:10.1111/j.2041-210X.2011.00169.x
Richards, G., Beking, T., Kreukels, B. P. C., Geuze, R. H., Beaton, A. A., & Groothuis, T. (2021). An
examination of the influence of prenatal sex hormones on handedness: Literature review and
amniotic fluid data. *Hormones and Behavior, 129*, 104929.
doi:https://doi.org/10.1016/j.yhbeh.2021.104929
Rilling, J. K., Glasser, M. F., Preuss, T. M., Ma, X., Zhao, T., Hu, X., & Behrens, T. E. J. (2008). The
evolution of the arcuate fasciculus revealed with comparative DTI. *Nature Neuroscience,
11*(4), 426-428. doi:10.1038/nn2072
Rilling, J. K., & Insel, T. R. (1999). Differential expansion of neural projection systems in primate brain
evolution. *NeuroReport, 10*(7), 1453-1459, doi:10.1097/00001756-199905140-00012
Ringo, J. L., Doty, R. W., Demeter, S., & Simard, P. Y. (1994). Time is of the essence: a conjecture that
hemispheric specialization arises from interhemispheric conduction delay. *Cerebral Cortex,
4*(4), 331-343. doi:10.1093/cercor/4.4.331
Rogers, L. J. (2009). Hand and paw preferences in relation to the lateralized brain. *Philosophical
Transactions of the Royal Society B: Biological Sciences, 364*(1519), 943-954.
doi:10.1098/rstb.2008.0225
Roos, C., Liedigk, R., Thinh, V. N., Nadler, T., & Zinner, D. (2019). The hybrid origin of the Indochinese
gray langur *Trachypithecus crepusculus*. *International Journal of Primatology, 40*(1), 9-27.
doi:10.1007/s10764-017-0008-4
Sanford, C., Guin, K., & Ward, J. P. (1984). Posture and laterality in the bushbaby (*Galago
senegalensis*). *Brain, Behavior and Evolution, 25*(4), 217-224. doi:10.1159/000118867
Schmitt, V., Melchisedech, S., Hammerschmidt, K., & Fischer, J. (2008). Hand preferences in Barbary
macaques (*Macaca sylvanus*). *Laterality, 13*(2), 143-157. doi:10.1080/13576500701757532
Schmitz, J., Metz, G. A. S., Güntürkün, O., & Ocklenburg, S. (2017a). Beyond the genome—Towards
an epigenetic understanding of handedness ontogenesis. *Progress in Neurobiology, 159*, 69-
89. doi:https://doi.org/10.1016/j.pneurobio.2017.10.005
Schmitz, J., Lor, S., Klose, R., Güntürkün, O., & Ocklenburg, S. (2017b). The functional genetics of
handedness and language lateralization: insights from gene ontology, pathway and disease
association analyses. *Frontiers in Psychology, 8*, 1144. Retrieved from
https://www.frontiersin.org/article/10.3389/fpsyg.2017.01144
Schweitzer, C., Bec, P., & Blois-Heulin, C. (2007). Does the complexity of the task influence manual
laterality in de Brazza's monkeys (*Cercopithecus neglectus*)? *Ethology, 113*(10), 983-994.
doi:https://doi.org/10.1111/j.1439-0310.2007.01405.x
Sha, Z., Pepe, A., Schijven, D., Carrión-Castillo, A., Roe, J. M., Westerhausen, R., . . . Francks, C. (2021).
Handedness and its genetic influences are associated with structural asymmetries of the
cerebral cortex in 31,864 individuals. *Proceedings of the National Academy of Sciences,
118*(47), e2113095118. doi:10.1073/pnas.2113095118
Silcox, M. T., Bloch, J. I., Boyer, D. M., Godinot, M., Ryan, T. M., Spoor, F., & Walker, A. (2009).
Semicircular canal system in early primates. *Journal of Human Evolution, 56*(3), 315-327.
doi:https://doi.org/10.1016/j.jhevol.2008.10.007
Spoelstra, K. (2021). *Lateralized behavior in white-handed gibbons (Hylobates lar).* Master Thesis.
Linköping University, Linköping.
Sussman, R. W., Tab Rasmussen, D., & Raven, P. H. (2013). Rethinking primate origins again.
American Journal of Primatology, 75(2), 95-106. doi:https://doi.org/10.1002/ajp.22096
Tab Rasmussen, D., & Nekaris, K. (1998). Evolutionary history of lorisiform primates. *Folia
Primatologica, 69*(Suppl. 1), 250-285. doi:10.1159/000052716
Tan, Y., Yoder, A. D., Yamashita, N., & Li, W.-H. (2005). Evidence from opsin genes rejects nocturnality
in ancestral primates. *Proceedings of the National Academy of Sciences of the United States
of America, 102*(41), 14712. doi:10.1073/pnas.0507042102
Toth, N. (1985). Archaeological evidence for preferential right-handedness in the lower and middle
pleistocene, and its possible implications. *Journal of Human Evolution, 14*(6), 607-614.
doi:https://doi.org/10.1016/S0047-2484(85)80087-7

Vauclair, J., Meguerditchian, A., & Hopkins, W. D. (2005). Hand preferences for unimanual and
982   coordinated bimanual tasks in baboons (*Papio anubis*). *Cognitive Brain Research, 25*(1), 210-
983   216. doi:https://doi.org/10.1016/j.cogbrainres.2005.05.012

984   Ward, J. P. (1998). Left hand advantage for prey capture in the galago (*Galago moholi*). *International
985   Journal of Comparative Psychology, 11*(4), 173-184.

986   Westergaard, G. C., Kuhn, H. E., & Suomi, S. J. (1998). Bipedal posture and hand preference in
987   humans and other primates. *Journal of Comparative Psychology, 112*(1), 55-64.
988   doi:10.1037/0735-7036.112.1.55

989   Westergaard, G. C., & Suomi, S. J. (1996). Hand preference for a bimanual task in tufted capuchins
990   (*Cebus apella*) and rhesus macaques (*Macaca mulatta*). *Journal of Comparative Psychology,
991   110*(4), 406-411. doi:10.1037/0735-7036.110.4.406

992   Yousry, T. A., Schmid, U. D., Alkadhi, H., Schmidt, D., Peraud, A., Buettner, A., & Winkler, P. (1997).
993   Localization of the motor hand area to a knob on the precentral gyrus. A new landmark.
994   *Brain, 120*(1), 141-157. doi:10.1093/brain/120.1.141

995   Zhao, D., Hopkins, W. D., & Li, B. (2012). Handedness in nature: first evidence on manual laterality on
996   bimanual coordinated tube task in wild primates. *American Journal of Physical Anthropology,
997   148*(1), 36-44. doi:https://doi.org/10.1002/ajpa.22038

Coefficient estimates (incl. 95% CI)