Cell-free chromatin particles released from dying cells inflict mitochondrial damage and ROS production in living cells

Gorantla V Raghuram[Ť], Bhabesh Kumar Tripathy[Ť], Kartikeya Avadhani[Ť], Snehal Shabrish, Naveen Kumar Khare, Relestina Lopes, Kavita Pal, Indraneel Mittra[*]

Translational Research Laboratory
Advanced Centre for Treatment, Research and Education in Cancer
Tata Memorial Centre
Kharghar, Navi Mumbai – 410210, India and
Homi Bhabha National Institute, Anushakti Nagar, Mumbai –400094

[Ť] These three investigators contributed equally to the work.

[*] Author for correspondence: Indraneel Mittra; imittra@actrec.gov.in

Abstract

mtDNA damage and the resultant oxidative stress are associated with neurodegenerative diseases, ageing and cancer. However, what triggers mtDNA damage remains unclear. We have reported that cell-free chromatin particles (cfChPs) that are released from the billions of cells that die in the body every day can readily enter into healthy cells and damage their DNA. We show here that cfChPs isolated from sera of healthy individuals, or those that are released from dying cells, inflict direct physical damage mtDNA leading to marked activation of ROS. The latter could be abrogated by concurrent treatment with three different cfChPs deactivating agents. Given that 1×10^{9}-1×10^{12} cells die in the body every day, our findings suggest that cfChPs from dying cells are major physiological triggers for mtDNA damage and ROS production. Deactivation of cfChPs may provide a novel therapeutic approach to retard ageing and associated degenerative conditions that have been linked to oxidative stress.

Introduction

Mitochondrial electron transport chain during oxidative phosphorylation is thought to be the predominant source of intracellular ROS[1]. In addition, external factors such as UV radiation, chemicals, heat and pH stress can also lead to production of mitochondrial ROS[2–5]. Under normal physiological conditions, ROS levels are regulated by a balance between generation of ROS and their elimination by several antioxidant enzymes and non-enzymatic defence systems[6]. Oxidative stress can arise when there is excess ROS production or its inefficient elimination. One of the major triggers of excess ROS production is damage to mitochondria, especially to mtDNA[7,8]. Excess ROS production can cause irreversible damage to mtDNA, proteins and membrane lipids resulting in mitochondrial dysfunction and ultimately to cell death[9]. Chronic mitochondrial dysfunction is associated with ageing and several degenerative conditions such as diabetes, cardiovascular diseases, neurodegenerative disorders and cancer[10–14].

Several hundred billion to a trillion cells die in the body every day[15,16] and the cell-free chromatin particles (cfChPs) that are released from them enter into the extracellular compartments, including into the circulation[17]. We have earlier reported that cfChPs can readily enter into the cells and damage their DNA[18,19]. In this study we provide evidence that cfChPs are a major source of ROS production by their ability to readily enter into healthy cells and inflict damage to mitochondria, especially to mtDNA. We also provide evidence that mtDNA damage and ROS production can be minimized by treatment with cfChPs inactivating agents suggesting therapeutic possibilities for multiple conditions mentioned above that are associated with oxidative stress.

Results

cfChPs are readily internalized by NIH3T3 cells

cfChPs isolated from sera of healthy individuals were fluorescently dually labelled in their DNA with Platinum Bright 550 and in their histones with ATTO-488 and added to cultured NIH3T3 cells (10 ng equivalent of DNA). Abundant fluorescent signals were detected in the cytoplasm of the treated cells at 4 h (**Fig. 1**). All further experiments described below were conducted at 4 h.

cfChPs inflict mtDNA damage: studies on whole cells

In order to investigate whether the internalized cfChPs induced mtDNA damage, NIH3T3 cells were treated with cfChPs (10ng) and the treated cells were immune stained with antibodies against γH2AX and pATM. The cells were simultaneously stained with MitoTracker Red to identify mitochondria. Figure 2 shows that numerous fluorescence γH2AX and pATM signals could be detected in the cytoplasm of treated cells which were highly significantly higher than those detected in untreated control cells ($p < 0.0001$) **(Fig. 2)**.

cfChPs inflict mtDNA damage: studies using isolated mitochondria

Mitochondria were isolated from NIH3T3 cells that had been treated with cfChPs (10ng) for 4h. Isolated mitochondria were clearly identifiable as DAPI positive particles which co-localized with MitoTracker Red **(Fig. 3a)**. The isolated mitochondria were next analysed in immune FISH experiments using fluorescent antibodies against histone H4 and a human whole genomic DNA probe. Negative control experiments, not shown here, were done to confirm that the DNA FISH probe did not cross hybridized with mouse DNA. Strictly co-localising fluorescent signals of histone H4, human DNA with DAPI indicated that cfChPs were in close contact with

mouse mitochondria represented by DAPI positive signals **(Fig. 3b)**. Dual staining of the isolated mitochondria with antibodies against γH2AX and MitoTracker Red and pATM and MitoTracker Red detected strictly co-localising signals indicating that cfChPs that were associated with mitochondria had inflicted mtDNA damage **(Figs 3c and d)**. When compared with isolated mitochondria from untreated NIH3T3 cells, the number of γH2AX and pATM signals in mitochondria isolated from treated cells were found to be significantly elevated (p<0.001 and p<0.0001 respectively).

cfChPs damage other components of mitochondria

We next examined whether, in addition to mtDNA damage, cfChPs inducing damage to other components of mitochondria.

Ultra structural changes: When examined under transmission electron microscopy (TEM), the untreated control cells were found to harbour mitochondria that were normal and elongated in shape. Mitochondria of the cfChPs treated cells, in contrast, were mainly rounded in appearance. The latter has been reported to signify mitochondrial damage[20] (**Fig. 4a**). Proportion of elongated and round mitochondria in control and treated cells are depicted as histograms. Quantitative estimation of the ratio of round versus elongated mitochondria showed a three-fold increase in favour of round mitochondria indicating increased mitochondrial damage in the treated cells (**Fig. 4a**).

Increase in mitochondrial mass and mitochondrial shape: It has been reported that a cell up-regulates mitochondrial biogenesis in response to mitochondrial damage in the form of increased mitochondrial mass[21]. MitoTracker Green has been widely used to assess mitochondrial mass as it stains the mitochondria irrespective of its membrane potential[22]. A marked increase in MitoTracker Green fluorescence was

observed in cells treated with cfChPs indicating that mitochondrial mass had increased as a result of damage inflicted by cfChPs treatment (**Fig. 4b**). MitoTracker red is known to detect changes in mitochondrial shape which correlates with mitochondrial damage[20,23]. Confocal microscopy of cfChPs treated cells stained with MitoTracker Red generated diffuse fluorescence signals which have been reported to represent mitochondrial fragmentation[20]. Untreated control cells on the other hand exhibited a distinct network of mitochondria (**Fig. 4b).** Flow cytometric analysis of cells dually labelled with MitoTracker Green and Red was performed to determine the relative propotion of functional mitochondria (MitoTracker Green**high** and MitoTracker Red**high**) and dysfunctional mitochondria (MitoTracker Green**high** and MitoTracker Red**low**) in response to cfChPs treatment. Quantitative analysis revealed an increased ratio of MitoTracker Green**high** and MitoTracker Red**low** population in cfChPs treated cells compared to untreated cells. These findings provided additional evidence of mitochondrial dysfunction (**Fig. 4b**)

Upregulation of mitochondrial outer membrane protein TOM20: Translocase of the outer membrane 20 (TOM20) acts as an import receptor that belongs to the family of TOM proteins and allows movement of proteins across the mitochondrial outer membrane into the inner membrane space[24]. Upregulation of TOM20 is a sign of dysfunctional mitochondria with increased biogenesis and excess import of proteins[25]. We observed a significant increase in expression of TOM20 in cfChPs treated cells when compared to controls under confocal microscopy (**Fig. 4c**). Quantitative analysis expressed as MFI per cell showed a three-fold increase in TOM20 expression in cfChPs treated cells when compared to control untreated cells (p<0.01) (**Fig. 4c**).

<u>Altered mitochondrial membrane potential</u>: Low mitochondrial membrane potential

has been reported to be associated with mitochondrial damage[26]. We investigated

mitochondrial depolarization as an indicator of mitochondrial damage using JC-1 (5,

5', 6, 6'-tetrachloro-1, 1', 3, 3'-tetraethyl benzimidazolylcarbocyanine iodide) - a

mitochondrial membrane fluorescent dye. JC-1 accumulates as aggregates in

healthy mitochondria which are known to have a high membrane potential and emit

red fluorescence. On the other hand, in dysfunctional mitochondria, JC-1 appears as

monomers and emits green fluorescence since they have a relatively low membrane

potential[27]. Representative images given in **Fig. 4d** show that while control cells have

predominantly red signals, cfChPs treated cells show abundance of green signals in

addition to red signals. Quantitative analysis of the data revealed a highly significant

increase in ratio of green to red fluorescence signals in cells treated with cfChPs

(p<0.001) (**Fig. 4d**). These results were validated using flow cytometry which

revealed a two-fold increase in ratio of green to red fluorescence in cfChPs treated

cells compared to untreated cells (p<0.05). Taken together, these results indicate

that cfChPs had induced damage to the mitochondria of the treated cells, leading to

decreased mitochondrial membrane potential.

cfChPs induce ROS production from mitochondria

Mitochondrial damage, especially mtDNA damage, is a major cause of excess ROS

production[7,8]. One of the primary forms of ROS generated in the mitochondria are

superoxide radicals[9], which can be readily detected by MitoSOX Red, a

triphenylphosphonium (TPP+)-linked DHE compound that is selectively oxidized by

superoxide radicals leading to emission of red fluorescence[28]. To investigate whether

mitochondrial damage, including mtDNA damage that we have described above had

led to increased ROS production, NIH3T3 cells were treated with cfChPs (10ng) for

4h followed by staining of the cells with MitoSOX Red. A marked increase in MitoSOX fluorescence was detected upon microscopic examination (p<0.0001) (**Fig. 5a)**. These results were validated using flow cytometry which detected a two-fold increase in MFI of MitoSox Red in cfChPs treated cells compared to untreated cells (p< 0.05). These data clearly indicated that cfChPs can induce increased ROS production resulting from damage to mitochondria, especially damage to mtDNA.

cfChPs activate the antioxidant enzyme superoxide dismutase

The next question that we addressed was whether excess ROS production following treatment of NIH3T3 cells with cfChPs had led to an increase in the antioxidant enzyme SOD-1. The latter was investigated both by IF and flow cytometry. Fluorescence microscopy of cells treated with cfChPs (10ng) for 4h detected a highly significant increase in SOD-1 production (p<0.0001) (**Fig. 5b)**. Flow cytometry analysis which also showed increased SOD-1 expression in treated cells (p<0.05).

Experiments using conditioned media from dying NIH3T3 cells

We next conducted experiments based on our earlier report that cfChPs that are spontaneously released from dying cells can be readily internalised by healthy cells[19]. These experiments involved treating NIH3T3 cells with conditioned medium containing cfChPs released from hypoxia induced dying NIH3T3 cells. The method for generating cfChPs containing conditioned medium has been described in the Material and Methods section. Supplementary figure 1 shows that cfChPs that were released from dually fluorescently labelled dying NIH3T3 cells were readily internalised by recipient healthy NIH3T3 cells. Conditioned medium containing cfChPs released from hypoxia induced dying NIH3T3 cells were collected and applied to live NIH3T3 cells. As a first step, we performed a dose response experiment wherein we added increasing volumes of cfChPs containing conditioned

medium to NIH3T3 cells followed by assay for MitoSOX Red expression as a marker of ROS production. Since MitoSox Red generates the by-product mitoethidium which intercalates in DNA and generates nuclear fluorescence, nuclei were gated and nuclear fluorescence was excluded from MFI analysis. Optimum ROS production was seen with a volume of 50 µl of culture media (**Supplementary Fig. 2**). All further experiments were performed using 50µl of conditioned medium. In order to confirm that cfChPs in the conditioned medium were indeed responsible for ROS production, conditioned media were pre-treated with the three different cfChPs-deactivating agent viz. anti-histone antibody complexed nanoparticles (CNPs), DNase I and a novel pro-oxidant combination of resveratrol and copper (R-Cu). As noted earlier, 50 µl of conditioned medium induced a marked activation of ROS in NIH3T3 cells which was abrogated by pre-treatment of the conditioned medium with all three cfChPs inactivating agents (p<0.0001) **(Fig. 6a)**. This experiment was repeated using flow cytometry which also detected down regulation of ROS production by cfChPs inhibitors **(Fig. 6b).** It should be pointed out that the degree of reduction detected by flow cytometry was somewhat less than that of obtained by analysis by fluorescent microscopy, since the nuclear fluorescence could not be gated and excluded from analysis in flow-cytometry experiments.

Discussion

Several hundred billion to a trillion cells die in the body every day[15,16], and cfChPs that are released from the dying cells enter into the extra-cellular compartments, including into the circulation[17]. We have earlier reported that circulating cfChPs, or those that are released locally from dying cells, can readily enter into healthy cells

wherein they induce dsDNA breaks, inflammation and apoptotic responses[18,19]. The

current study was based on the hypothesis that, in addition to DNA damage, the

internalised cfChPs would also induce damage to mitochondria. We have shown in

this study that this is indeed the case. Treatment of NIH3T3 mouse fibroblast cells

with cfChPs isolated from healthy human individuals led to activation of H2AX and

ATM signals which co-localized with those of mitochondria identified by staining with

MitoTracker Red. This was further confirmed in experiments using isolated

mitochondria from cfChPs treated NIH3T3 cells in which we show that mtDNA

damage is caused by physical association of cfChPs to mtDNA. We speculate that

the physical association of cfChPs with mitochondria is the consequence of a

chemical interaction between the positively charged histones of cfChPs and the

negatively charged mtDNA. The binding of cfChPs to mitochondria leads to multiple

other signs of mitochondrial damage including ultra-structural changes, changes in

mitochondrial mass, shape and mitochondrial function. These events lead to

activation of ROS production and consequent up-regulation of the anti-oxidant

enzyme SOD-1. We propose that excess ROS production sets in motion a vicious

cycle of more mtDNA damage and production of ROS leading to oxidative stress

which is associated with several pathological conditions such as neurodegenerative

diseases, ageing and cancer[7,29,30]. The mechanism of cfChPs induced mitochondrial

damage and ROS production as discussed above is graphically illustrated in

Supplementary Fig. 3.

We demonstrate that cfChPs induced ROS production can be greatly reduced by

three different cfChPs inactivating agents. Of the three agents, a combination of the

widely used nutraceuticals resveratrol (R) and copper (Cu) hold therapeutic promise.

We have shown that combining small quantities of R and Cu leads to production of

oxygen radicals which are capable of deactivating cfChPs with multiple therapeutic

effects[31–38].

Oxidative stress resulting from increased ROS production is known to be the

underlying cause of many serious disorders[10–14]. However, what causes excess ROS

production has remained elusive till now. We have shown in this study that

mitochondrial damage is inflicted by extraneous agents in the form of cfChPs. Given

that 1×10^9-1×10^{12} cells die in the body every day, our findings suggest that cfChPs

from dying cells are major physiological triggers for mtDNA damage and ROS

production. Deactivation of cfChPs may provide a novel therapeutic approach to

retard ageing and associated degenerative conditions that have been linked to

oxidative stress. In this context, we have recently reported that prolonged

administration of resveratrol and copper can retard multiple biological hallmarks of

ageing and neurodegeneration in C57Bl/6 mice[38].

Materials and Methods

Ethics approval

Institutional Ethics Committee (IEC) approval was obtained for collecting blood samples from healthy volunteers and signed informed consent was obtained on consent forms approved by IEC (Approval no.900520).

Reagents and antibodies

Commercial sources and catalogue numbers of reagents and antibodies used in this study are given in Supplementary Table 1.

Isolation of cfChPs from sera of healthy donors

Isolation of cfChPs from sera of healthy individuals was performed according to a protocol described by us earlier[18]. The isolated cfChPs when examined under electron microscopy revealed a beads-on-a-string appearance typical of chromatin[18]. The amount of DNA in cfChPs was estimated by PicoGreen quantification assay, and cfChPs concentration was expressed in terms of their DNA content.

Fluorescent dual labelling of cfChPs

cfChPs were fluorescently dually labelled in their DNA by Platinum Bright 550 (red) and in their histone H4 with ATTO-488 (green) according to a protocol described by us earlier[18].

Cell culture

NIH3T3 embryonic mouse fibroblast cells were obtained from the American Type Culture Collection (ATCC, USA) and grown in Dulbecco's Modified Eagle's Medium (DMEM) (Gibco, Catalog No.12800-017) containing 10% bovine calf serum (Cytiva HyClone, Catalog No. SH30073) and maintained at 37□°C in an atmosphere of 5%

CO_2 and air. For microscopy experiments, cells were seeded at a density of 1×10^5 on

coverslips in 1.5 ml DMEM; for flow cytometry, 2×10^5 cells were seeded in 1.5 ml

DMEM and incubated overnight before conducting the experiments.

Fluorescence and confocal microscopy

For fluorescence microscopy cells were imaged using Spectral Bio-Imaging System

(Applied Spectral Imaging). Images were captured using a 40x air objective. All

images were captured with the same exposure time set to ensure uniform

fluorescence intensity for purposes of comparison. For confocal microscopy, images

were acquired using a 63x oil objective on Zeiss LSM 780 laser scanning

microscope (Carl Zeiss, Germany) and Leica SP8 confocal imaging system (Leica

Microsystems, Wetzlar, Germany). Mean fluorescence intensity (MFI) of images was

measured using ImageJ (W. Rasband, Bethesda, USA).

Flow cytometry

For flow cytometry, cells were acquired on Attune NxT flow cytometer (Thermo

Fisher Scientific, USA) and analysis was performed using *FlowJo™ v10.6 Software*.

NIH3T3 cells were gated on forward and side scatter parameters. At least 20,000

NIH3T3 cells were acquired and the threshold between negative and positive

expression was defined by the fluorescence minus one (FMO) method. Median

fluorescence intensity (MFI) of respective dye was then compared by histogram

analysis of untreated and treated cells.

Isolation of mitochondria

NIH3T3 cells were treated with 10 ng cfChPs for 4 h and mitochondria were isolated

from the treated cells using Mitochondria Isolation Kit (Supplementary Table 1) as

per the instruction manual. Mitochondria were similarly isolated from control untreated cells.

Fluorescent staining of isolated mitochondria

a) MitoTracker Red CMXRos Staining

Isolated mitochondria were cytospun onto a slide and fixed in 2% paraformaldehyde (PFA) / 1.5% glutaraldehyde (GA) fixative mixture for 15 min on ice followed by neutralisation in 0.3 M Glycine buffer for 30 min at room temperature. The fixed smears were stained with 100nM MitoTracker Red CMXRos for 15 min at room temperature and stained with DAPI (1µg/ml) in Hanks Balanced Salt Solution (HBSS). Images were acquired on the Spectral Bio-Imaging System (Applied Spectral Imaging).

b) Immuno-FISH for Histone and Human DNA

Isolated mitochondria were fixed as described above and processed for immune-staining with anti-histone H4 IgG and appropriate secondary antibodies (Supplementary Table 1). The immunostained smears were fixed with 2% PFA - 1.5% GA mixture for 5 min on ice and processed for FISH using custom synthesized Human DNA probe (Supplementary Table 1). Briefly, immune-stained slides were hybridised overnight with Human DNA probe at 37°C and washed once with 0.4X sodium saline Citrate (SSC) at 72°C (± 2°C) for 2 mins and 4X SSC in Tween-20 twice at room temperature for 5 mins each. The slides were mounted with DAPI (1µg/ml) in HBSS and images were acquired on the Spectral Bio-Imaging System (Applied Spectral Imaging).

Detection of mtDNA damage by immunofluorescence

Mitochondrial DNA damage in the form of activation of H2AX and ATM was

evaluated on both whole cells and isolated mitochondria.

<u>Whole Cells:</u> Cells were stained with the mitochondrial stain MitoTracker Red

CMXRos and processed for immune-staining for γ-H2AX and p-ATM using

appropriate antibodies (Supplementary table 1). The cells were counterstained with

Vecta-shield DAPI and mounted on slides. Images were acquired using Spectral Bio-

Imaging System (Applied Spectral Imaging). Mean fluorescence intensity (MFI) in the

cytoplasm was measured after gating out the nuclei fluorescence.

<u>Isolated mitochondria:</u>   DNA damage in isolated mitochondria were assessed by

standard IF procedure as described above. Images were acquired under Spectral

Bio-Imaging System (Applied Spectral Imaging).

Assessment of mitochondrial integrity by electron microscopy

Exponentially growing NIH3T3 cells (4×10^6) were treated with cfChPs (10 ng) for 4 h.

Control and treated cells were fixed with 3% glutaraldehyde in 0.1 M sodium

cacodylate buffer (pH 7.2) for 2 h at 4°C followed by 1% osmium tetroxide in 0.1 M

sodium cacodylate buffer for 1 h at 4°C. Samples were then dehydrated using

alcohol at 4°C and embedded in araldite resin to make ultrathin sections that were

mounted onto EM grids for imaging. Grids were examined under a JEM1400-Plus

transmission electron microscope (JEOL, Japan).

Assessment of mitochondrial mass using MitoTracker Green FM

<u>Fluorescence microscopy:</u> Following treatment with cfChPs (10 ng) for 4 h, NIH3T3

cells were stained with 100 nM MitoTracker Green FM diluted in 2ml of Hank's

balanced salt solution (HBSS, pH 7.4) for 15 minutes at 37°C. Cells were washed

with HBSS, counterstained with Hoechst, mounted on glass slides and imaged using

Spectral Bio-Imaging System.

<u>Flow cytometry</u>: Cells were stained with 20nM MitoTracker Green FM diluted in 2ml

of Hank's balanced salt solution (HBSS, pH 7.4) for 15 minutes at 37°C. Cells were

washed twice with pre-warm HBSS, collected in FACs tubes using cell scraper.

Samples were acquired on Attune NxT flow cytometer (ThermoFisher Scientific,

USA)

360 Assessment of mitochondrial shape using MitoTracker Red CMXRos

<u>Fluorescence microscopy:</u> Following treatment with cfChPs (10 ng) for 4 h, NIH3T3

cells were stained with MitoTracker Red CMXRos (100nM) diluted in 2ml of Hank's

balanced salt solution (HBSS, pH 7.4) for 15 minutes at 37°C. Cells were then

washed thrice with PBS, fixed using 4% para-formaldehyde (PFA), counterstained

with Vecta-shield DAPI and mounted onto glass slides. Images were acquired on the

LSM 780 confocal microscope.

<u>Flow cytometry</u>: Cells were stained with MitoTracker Red CMXRos (100nM) diluted

in 2ml of Hank's balanced salt solution (HBSS, pH 7.4) for 15 minutes at 37°C. Cells

were washed twice with pre-warm HBSS, collected in FACs tubes using cell scraper.

Samples were acquired on Attune NxT flow cytometer (ThermoFisher Scientific,

USA).

372 Assessment of mitochondrial membrane protein TOM20.

NIH3T3 cells were treated with cfChPs (10 ng) for 4 h and immunostaining for

TOM20 was performed by fixing the cells with 4% PFA for 15 minutes at 37°C

followed by permeabilization with 0.2% Triton X-100 for 30 min and blocking with 3%

BSA for 1 h. Cells were immune-stained using anti-rabbit TOM20 primary antibody

and appropriate secondary anti-rabbit FITC antibody followed by mounting with

Vecta-shield DAPI. Images were acquired on the Leica confocal SP8 imaging

system.

Assessing mitochondrial membrane potential using JC-1 dye

Fluorescence microscopy: After treatment with cfChPs (10 ng) for 4 h, NIH3T3 cells were stained with JC-1 dye in HBSS and incubated at 37°C for 15 min. The cells were washed twice with pre-warmed HBSS. Cells were counterstained with Hoechst, mounted on glass slides and cell-associated fluorescence was detected using Spectral Bio-Imaging System (Applied Spectral Imaging).

Flow cytometry: Cells were stained with JC-1 dye in HBSS and incubated at 37°C for 15 min. Cells were washed twice with pre-warm HBSS, collected in FACs tubes using cell scraper. Samples were acquired on Attune NxT flow cytometer (ThermoFisher Scientific, USA).

Assessment of mitochondrial ROS production using MitoSOX Red

Fluorescence microscopy: NIH3T3 cells were treated with cfChPs (10 ng) for 4 h and stained with 0.5 µM MitoSOX Red. Cells were in 2 ml of HBSS for 15 minutes at 37°C, washed with warm buffer. Cells were counter stained with Hoechst, mounted on glass slides and cell-associated fluorescence was measured using Spectral Bio-Imaging System (Applied Spectral Imaging).

Flow cytometry: Cells were stained with 0.5 µM MitoSOX Red diluted in 2 ml of HBSS for 15 minutes at 37°C, washed with pre-warm HBSS and then were collected in FACs tubes using cell scraper. Samples were acquired on Attune NxT flow cytometer (ThermoFisher Scientific, USA).

Assessment of superoxide dismutase (SOD-1) expression

Fluorescence microscopy: cfChPs treated NIH3T3 cells were fixed with 4% paraformaldehyde (PFA) for 15 minutes at 37°C and were then immune-stained using primary antibody against SOD-1 and appropriate secondary antibody according to the protocol described above. Cells were mounted with Vecta-shield

DAPI, and images were acquired using Spectral Bio-Imaging System (Applied Spectral Imaging). <u>Flow cytometry</u>: Cells were fixed with 4% paraformaldehyde (PFA) for 15 minutes at 37°C and were then stained using primary antibody against SOD-1 and appropriate secondary antibody. Cells were acquired on Attune NxT flow cytometer (ThermoFisher Scientific, USA).

Procedure for collecting conditioned media from hypoxia induced dying NIH3T3 cells

A dual chamber system was used to generate culture medium containing cfChPs released from dying cells. NIH3T3 cells were seeded at a density of 2 x 10^5 on ThinCert® Cell Culture Inserts (pore size 400 nm) containing 1.5 ml of DMEM and were placed in a six-well culture plate. Cells were incubated at 37 °C for 24 h. The six-well plate with Thincert® Inserts was then transferred to a hypoxia chamber with 1% O_2 to induce hypoxic cell death. After 48h, the plate was removed from the hypoxia chamber and the 700 µl of DMEM media was added to the lower chamber of the six-well palte below the ThinCert® Inserts and kept for 48 h at 37°C in normoxic conditions to allow cfChPs <400 nm in size to seep into the medium in the lower chamber. The medium in the lower chamber was collected and divided into aliquots and stored at -80°C until further use.

Assessment of mitochondrial ROS production using conditioned medium

Aliquots were thawed for use in the dose / volume response experiments which involved addition of increasing volumes of the culture medium to NIH3T3 cells, and the level of ROS was measured by MitoSOX Red staining. A dose / volume of 50 µl was found to be the most effective in generating ROS (Supplementary Fig. 1). Once the optimal dose / volume was ascertained, aliquots were thawed for experiments involving the use of three different cfChPs deactivators viz. pullulan-histone antibody

430   nanoconjugates (CNPs), DNase I and a novel pro-oxidant combination of small

431   quantities of resveratrol and copper (R-Cu) (please see below).

Pullulan-histone antibody nanoconjugates (CNPs)

433   Pullulan-histone antibody nanoconjugates (complexed nanoparticles, CNPs) were

434   synthesized according to the protocol described by us earlier[39].

435   **DNase I.** DNase I from bovine pancreas was procured from Sigma-Aldrich

436   (Catalogue No. DN25-1G).

Resveratrol-copper (R-Cu)

438   Resveratrol (R) is a plant polyphenol with antioxidant properties[40]. However, R acts

439   as a pro-oxidant in the presence of copper (Cu) by reducing Cu (II) to Cu (I) thereby

440   generating a free radical[41,42]. R-Cu can deactivate cfChPs in *in vitro*[19,34]. The involved

441   concentrations of R and Cu used in this study were 1mM and 0.0001mM

442   respectively.

Experiments using conditioned medium with or without cfChPs deactivators

444   Conditioned media were pre-treated with the above three cfChPs deactivators for 1h

445   at 37°C prior to addition to NIH3T3 cells. ROS production in NIH3T3 cells were

446   measured at 4h using MitoSox Red (protocol mentioned above).

Statistical analysis

448   All data are presented as Mean $\pm$ Standard Error of Mean (SEM). Statistical

449   analysis was performed using GraphPad Prism 8 (GraphPad Software, Inc., USA,

450   Version 8.0). Data were compared using Student's t-test (two tailed, unpaired) and

451   one-way ANNOVA. $p < 0.05$ was taken as the level of significance.

Data Availability:

453   All data are contained within the manuscript. Any additional data will be provided on

454   reasonable request.

Legends to figures

Fig. 1: Rapid uptake of cfChPs by NIH3T3 cells at 4h. NIH3T3 cells were treated with cfChPs (10ng) isolated from sera of healthy individuals which had been dually labelled in their DNA (Platinum Bright 550) and in their histones (ATTO-488). Images were acquired on Applied Spectral bio-imaging system. Dually labelled cfChPs are clearly seen in the cytoplasm. The particles appear relatively large which is most likely due to linking-up of multiple cfChPs to form concatameres as described by us earlier[18].

Fig. 2. **cfChPs treatment of NIH3T3 cells induces mitochondrial DNA damage as observed at 4h. a.** Representative images of untreated control and cfChPs (10ng) treated cells dually stained with MitoTracker red dye and antibody against γH2AX (green). Expression of γH2AX is visibly increased in the mitochondria of cfChPs treated cells (arrows). Quantitative analysis of MFI was performed after gating and excluding the nuclear fluorescence. Analysis illustrates significantly higher MFI for γH2AX in treated cells as compared to untreated controls. It should be noted that cfChPs have induced the expected activation of H2AX in the nuclei of the treated cells as reported by us earlier[18]. **b.** Representative images of untreated control and cfChPs (10ng) treated cells dually stained with MitoTracker red dye and antibody against pATM (green). Expression of pATM is visibly increased in the mitochondria of cfChPs treated cells (arrows). Quantitative analysis of MFI was performed after gating and excluding the nuclear fluorescence. Analysis illustrates significantly higher MFI for pATM in treated cells as compared to untreated controls. It should be noted that cfChPs have induced the expected activation of ATM in the nuclei of the treated cells as reported by us earlier[18]. Experiments were repeated twice and all images were acquired under the same settings. Results represent mean ± SEM values. Scale Bar - 5 µm. Data were analysed using Student's t-test. *** p<0.001, **** p<0.0001.

Fig. 3. Mitochondria isolated from cfChPs treated cells show intimate association with cfChPs. a. Confirmation that isolated mitochondria represented by

Mito-Tracker Red co-localize with DAPI signals; **b.** Immuno-FISH performed on
isolated mitochondria using antibody against histone H4 and a human DNA FISH
probe show co-localization of cfChPs with mitochondria represented by DAPI. **c.**
Mitochondria isolated from cfChPs treated cells show phosphorylated H2AX signals
which co-localize with mitochondria represented by Mito-Tracker Red. Few γH2AX
signals are seen in untreated control cells. Histograms represent quantitative
analysis of co-localizing signals in control and treated cells. **d.** Mitochondria isolated
from cfChPs treated cells show phosphorylated ATM signals which co-localize with
mitochondria represented by Mito-Tracker Red. Few pATM signals are seen in
untreated control cells. Histograms represent quantitative analysis of co-localizing
signals in control and treated cells. Results are represented as mean ± SEM values
and data were analysed using Student's☐t-☐test. ******* $p<0.001$; ******** $p<0.0001$.

Fig. 4: Various mitochondrial abnormalities induced by cfChPs in treated cells
as observed at 4 h. a. Representative EM images of untreated and cfChPs (10ng)
treated cells reveals ultra-structural changes in mitochondria. Control cells contain
elongated mitochondria (red arrows), whereas cfChPs treated cells harbour mostly
round shaped mitochondria (red arrows) (Scale bar - 1 μm). ELM= Elongated
Mitochondria, RM= Round Mitochondria. Histograms represent quantitative analysis
of ultrastructural changes and the ratio of round to elongated mitochondria in treated
and control cells. **b.** Evaluation of mitochondrial dysfunction by IF and flow cytometry
using mitochondrial dyes. Representative microphotographs of mitochondrial mass
(MitoTracker Green) and of mitochondrial dysfunction (MitoTracker red) in cfChPs
(10ng) treated and control cells. Quantitative analysis (MFI) of the ratio of
mitochondrial mass and mitochondrial dysfunction by flow cytometry. The latter
shows a significant increase in ratio of green to red fluorescence in cfChPs treated
cells as compared to untreated cells. **c.** Representative confocal microphotographs
of mitochondrial outer membrane marker TOM20 in cfChPs treated (10ng) and
untreated cells. Expression of TOM20 is visibly higher in cfChPs treated cells
compared to untreated cells. (Scale Bar - 20 μm, Pseudo colour - Orange).
Quantitative analysis (MFI) of TOM20 expression in control versus cfChPs treated
cells exhibits a significant increase in MFI in treated cells indicating loss of
mitochondrial membrane integrity as compared to control cells. **d.** Representative

confocal microphotographs of mitochondrial membrane potential marker JC-1 in
cfChPs treated (10ng) and untreated cells. Quantitative analysis (MFI) shows
increased green to red fluorescence ratio signifying depolarized mitochondrial
membrane. Quantitative MFI analysis by flow cytometry reveals higher ratio of green
to red fluorescence in cfChPs treated cells compared to control cells. All the
experiments were repeated twice. Results are represented as mean ± SEM values
and data were analysed using Student's□t-□test. * =p<0.05, ** =p<0.01, ***
=p<0.005.

Fig. 5: Treatment of NIH3T3 cells with cfChPs induces oxidative stress as
detected at 4h. a. Representative fluorescence microscopy images of control and
cfChPs treated (10ng) cells stained with MitoSOX Red shows significant activation of
ROS production in treated cells as compared to untreated cells (Scale Bar = 5 μm).
Results of quantitative analysis of MFI (after gating the nuclei to exclude nuclear
fluorescence) are shown as histograms. Flow cytometric analysis of MitoSox Red
confirms significant increase in treated cells compared to control cells.   **b.**
Representative fluorescence microscopy images of control and cfChPs (10ng)
treated cells stained with superoxide dismutase-1 (SOD-1) antibody shows
significant increase in activation of SOD-1 in treated cells as compared to untreated
cells (Scale Bar = 5 μm). Results of quantitative analysis (MFI) are shown as
histograms. Flow cytometric analysis of SOD-1 confirms significant increase in
treated cells compared to control cells. All the experiments were repeated twice.
Results are represented as mean ± SEM values and data were analysed using
Student's□t-□test. * =p<0.05, *** =p<0.005.

Fig. 6. Activation of ROS production in NIH3T3 cells treated with conditioned
medium containing cfChPs from hypoxia induced dying NIH3T3 cells in
presence and absence of inhibitors of cfChPs. a. Representative fluorescence
microscopy images showing activation of ROS production at 4h detected by MitoSox
Red in NIH3T3 cells following treatment with conditioned medium (50μl) from dying
NIH3T3 cells (upper panel). ROS activation is inhibited by concurrent treatment with
CNPs, DNase I and R-Cu (lower panel). Results of quantitative analysis (MFI) are

given as histograms. Since MitoSox Red generates the by-product mitoethidium which intercalates in DNA and generates nuclear fluorescence, nuclei were gated and nuclear fluorescence was excluded from MFI analysis. **b.** Representative flow cytometry plots showing activation of ROS production represented by MitoSox Red in NIH3T3 cells following treatment with conditioned medium (50μl) from dying NIH3T3 cells. ROS activation is inhibited by concurrent treatment with CNPs, DNase I and R-Cu. Results of quantitative analysis (MFI) are given as histograms. It should be noted that the degree of reduction detected by flow cytometry is less than that of obtained by microscopy analysis since the nuclear fluorescence could not be gated and excluded from analysis in flow-cytometry experiments. Experiments were repeated twice. Results are represented as mean ± SEM values and data were analysed using one way ANNOVA (GraphPad Prism 8).

Legends to supplementary figures

S1 Fig. Representative fluorescence microscopy images showing bystander uptake of dually labelled cfChPs by NIH3T3 cells at 4h. Donor NIH3T3 cells were dually labelled in their DNA with BrdU (red) and histones by CellLight® Histone 2B-GFP (green). Cells were kept in hypoxic chamber for 48h and the conditioned medium (50µl) containing dually labelled cfChPs were added to healthy recipient NIH3T3 cells.

S2 Fig. Dose / volume response analysis of activation of ROS production as detected by MitoSOX at 4h. NIH3T3 cells treated with increasing volume of conditioned medium containing cfChPs released from hypoxia induced dying NIH3T3 cells. Maximum activation of MitoSOX is seen with 50µl of culture medium.

S3 Fig. Graphical illustration showing cfChPs that circulate in blood (left hand image) and those that are released locally from dying cells (right hand image) can readily enter into healthy cells to damage their mitochondria leading to ROS production.

References

1. Zhao, R. Z., Jiang, S., Zhang, L. & Yu, Z. bin. Mitochondrial electron transport chain, ROS generation and uncoupling (Review). *International Journal of Molecular Medicine* vol. 44 Preprint at https://doi.org/10.3892/ijmm.2019.4188 (2019).

2. de Jager, T. L., Cockrell, A. E. & du Plessis, S. S. Ultraviolet light induced generation of reactive oxygen species. Ultraviolet Light in Human Health. *Diseases and Environment* 15–23 (2017).

3. Hortelano, S. *et al.* Nitric oxide induces apoptosis via triggering mitochondrial permeability transition. *FEBS Lett* **410**, (1997).

4. Slimen, I. B. *et al.* Reactive oxygen species, heat stress and oxidative-induced mitochondrial damage. A review. *International Journal of Hyperthermia* vol. 30 Preprint at https://doi.org/10.3109/02656736.2014.971446 (2014).

5. Selivanov, V. A. *et al.* The role of external and matrix pH in mitochondrial reactive oxygen species generation. *Journal of Biological Chemistry* **283**, (2008).

6. Birben, E., Sahiner, U. M., Sackesen, C., Erzurum, S. & Kalayci, O. Oxidative stress and antioxidant defense. *World Allergy Organization Journal* vol. 5 Preprint at https://doi.org/10.1097/WOX.0b013e3182439613 (2012).

7. Guo, C. Y., Sun, L., Chen, X. P. & Zhang, D. S. Oxidative stress, mitochondrial damage and neurodegenerative diseases. *Neural Regen Res* **8**, (2013).

8. Nissanka, N. & Moraes, C. T. Mitochondrial DNA damage and reactive oxygen species in neurodegenerative disease. *FEBS Letters* vol. 592 Preprint at https://doi.org/10.1002/1873-3468.12956 (2018).

9. Halliwell, Barry & John MC Gutteridge. *Free radicals in biology and medicine.* (Oxford university press, 2015).

10. Cui, Hang, Yahui Kong & Hong Zhang. Oxidative stress, mitochondrial dysfunction, and aging. *J Signal Transduct* 2012 (2012).

11. Itoh, K., Nakamura, K., Iijima, M. & Sesaki, H. Mitochondrial dynamics in neurodegeneration. *Trends Cell Biol* **23**, 64–71 (2013).

12. Parish, R. & Petersen, K. F. Mitochondrial dysfunction and type 2 diabetes. *Curr Diab Rep* **5**, 177–183 (2005).

13. Ballinger, S. W. Mitochondrial dysfunction in cardiovascular disease. *Free Radic Biol Med* **38**, 1278–1295 (2005).

620   14.   Boland, M. L., Chourasia, A. H. & Macleod, K. F. Mitochondrial dysfunction in
621   cancer. *Front Oncol* 292 (2013).

622   15.   Fliedner, T. M., Graessle, D., Paulsen, C. & Reimers, K. Structure and function
623   of bone marrow hemopoiesis: mechanisms of response to ionizing radiation
624   exposure. *Cancer Biother Radiopharm* **17**, 405–426 (2002).

625   16.   Sender, R. & Milo, R. The distribution of cellular turnover in the human body.
626   *Nat Med* **27**, 45–48 (2021).

627   17.   Holdenrieder, S. & Stieber, P. Clinical use of circulating nucleosomes. *Crit Rev
628   Clin Lab Sci* **46**, 1–24 (2009).

629   18.   Mittra, I. *et al.* Circulating nucleic acids damage DNA of healthy cells by
630   integrating into their genomes. *J Biosci* **40**, 91–111 (2015).

631   19.   Mittra, I. *et al.* Cell-free chromatin from dying cancer cells integrate into
632   genomes of bystander healthy cells to induce DNA damage and inflammation.
633   *Cell Death Discov* **3**, 17015 (2017).

634   20.   Zorov, D. B. *et al.* Lessons from the discovery of mitochondrial fragmentation
635   (fission): A review and update. *Cells* vol. 8 Preprint at
636   https://doi.org/10.3390/cells8020175 (2019).

637   21.   Apostolova, N. *et al.* Enhanced oxidative stress and increased mitochondrial
638   mass during efavirenz-induced apoptosis in human hepatic cells. *Br J
639   Pharmacol* **160**, 2069–2084 (2010).

640   22.   Marques-Santos, L. F., Oliveira, J. G. P., Maia, R. C. & Rumjanek, V. M.
641   Mitotracker green is a P-glycoprotein substrate. *Biosci Rep* **23**, 199–212
642   (2003).

643   23.   Samudio, I. *et al.* 2-Cyano-3, 12-dioxooleana-1, 9-dien-28-imidazolide (CDDO-
644   Im) directly targets mitochondrial glutathione to induce apoptosis in pancreatic
645   cancer. *Journal of Biological Chemistry* **280**, 36273–36282 (2005).

646   24.   Dudek, J., Rehling, P. & van der Laan, M. Mitochondrial protein import:
647   common principles and physiological networks. *Biochimica et Biophysica Acta
648   (BBA)-Molecular Cell Research* **1833**, 274–285 (2013).

649   25.   Franco-Iborra, S. *et al.* Defective mitochondrial protein import contributes to
650   complex I-induced mitochondrial dysfunction and neurodegeneration in
651   Parkinson's disease. *Cell Death Dis* **9**, 1122 (2018).

652   26.   Popkov, V. A. *et al.* Mitodiversity. *Biochemistry (Moscow)* **80**, 532–541 (2015).

653   27.   Sivandzade, F., Bhalerao, A. & Cucullo, L. Analysis of the Mitochondrial
654   Membrane Potential Using the Cationic JC-1 Dye as a Sensitive Fluorescent
655   Probe. *Bio Protoc* **9**, (2019).

656   28.   Johnson-Cadwell, L. I., Jekabsons, M. B., Wang, A., Polster, B. M. & Nicholls,
657   D. G. Mild Uncoupling does not decrease mitochondrial superoxide levels in
658   cultured cerebellar granule neurons but decreases spare respiratory capacity
659   and increases toxicity to glutamate and oxidative stress. *J Neurochem* **101**,
660   1619–1631 (2007).

661   29.   Harman, D. The aging process. *Proceedings of the National Academy of*
662   *Sciences* **78**, 7124–7128 (1981).

663   30.   Copeland, W. C., Wachsman, J. T., Johnson, F. M. & Penta, J. S.
664   Mitochondrial DNA alterations in cancer. *Cancer Invest* **20**, 557–569 (2002).

665   31.   Mittra, I. *et al.* Prevention of chemotherapy toxicity by agents that neutralize or
666   degrade cell-free chromatin. *Annals of Oncology* **28**, (2017).

667   32.   Agarwal, A. *et al.* A novel pro-oxidant combination of resveratrol and copper
668   reduces transplant related toxicities in patients receiving high dose melphalan
669   for multiple myeloma (RESCU 001). *PLoS One* **17**, (2022).

670   33.   Ostwal, V. *et al.* A pro-oxidant combination of resveratrol and copper reduces
671   chemotherapy-related non-haematological toxicities in advanced gastric
672   cancer: results of a prospective open label phase II single-arm study (RESCU
673   III study). *Medical Oncology* **40**, 17 (2022).

674   34.   Kirolikar, S. *et al.* Prevention of radiation-induced bystander effects by agents
675   that inactivate cell-free chromatin released from irradiated dying cells. *Cell*
676   *Death Dis* **9**, 1142 (2018).

677   35.   Mittra, I. *et al.* Cell-free chromatin particles released from dying host cells are
678   global instigators of endotoxin sepsis in mice. *PLoS One* **15**, e0229017 (2020).

679   36.   Mittra, I. *et al.* Resveratrol and Copper for treatment of severe COVID-19: an
680   observational study (RESCU 002). *medRxiv* 2020–2027 (2020).

681   37.   Pilankar, A. *et al.* A pro-oxidant combination of resveratrol and copper down-
682   regulates hallmarks of cancer and immune checkpoints in patients with
683   advanced oral cancer: Results of an exploratory study (RESCU 004). *Front*
684   *Oncol* 5061 (2022).

685   38.   Pal, K. *et al.* A pro-oxidant combination of resveratrol and copper down-
686   regulates multiple biological hallmarks of ageing and neurodegeneration in
687   mice. *Sci Rep* **12**, 17209 (2022).

39. Rekha, M. R. *et al.* Pullulan-histone antibody nanoconjugates for the removal
of chromatin fragments from systemic circulation. *Biomaterials* **34**, (2013).

40. Zhang, L. X. *et al.* Resveratrol (RV): A pharmacological review and call for
further research. *Biomedicine and Pharmacotherapy* **143**, (2021).

41. Fukuhara, K. & Miyata, N. Resveratrol as a new type of DNA-cleaving agent.
Bioorganic and Medicinal Chemistry Letters **8**, (1998).

42. Fukuhara, K. *et al.* Structural basis for DNA-cleaving activity of resveratrol in
the presence of Cu(II). *Bioorganic and Medicinal Chemistry* **14**, (2006).

Funding:

This study was supported by the Department of Atomic Energy, Government of India, through its grant CTCTMC to Tata Memorial Centre awarded to IM.

Acknowledgement:

The authors thank Mr. Ashish Pawar for his help in preparing manuscript.

Author contributions:

G.V.R., B.K.T., K.A., S.S., N.K.K., R.L., K.P. conducted the experiment; G.V.R., B.K.T., K.A., S.S. conducted data analysis; G.V.R. supervised the experiments; G.V.R., B.K.T., K.A., S.S., I.M. wrote the paper; I.M. conceived the idea, procured funding, supervised the experiments, wrote the paper and approved the final manuscript.

Competing Interest Statement:

The authors declare no competing interests

Fig. 1

	Pt Bright Red (DNA)	Atto-488 Green (Histone)	Merged

Scale bar = 20 μm

Fig. 2

a γ-H2AX

b pATM

Fig. 3

Fig. 4

a

EM

c

TOM-20

b

MitoTracker Green & Red

d

JC-1

Fig. 5

a

MitoSOX Red

Fig. 6

a

Control

Conditioned Medium

Conditioned Medium + CNPs

Conditioned Medium + DNase I

Conditioned Medium + R-Cu

Scale bar = 10 μm

ANOVA
p=0.0001

MFI per cell ($\times10^4$)

Control   CM   CM+CNPs   CM+DNase   CM+R-Cu

b

Only cells

Control

Conditioned Medium

Conditioned Medium + CNPs

Conditioned Medium + DNase I

Conditioned Medium + R-Cu

10^3 10^4 10^6

ANOVA
p=0.0251

MFI

Control   CM   CM+CNPs   CM+DNase   CM+R-Cu