DRP1-mediated mitochondrial fission is essential to maintain cristae morphology and bioenergetics

Gabriella L. Robertson[1], Stellan Riffle[1], Mira Patel[1], Andrea Marshall[2], Heather Beasley[2], Edgar Garza Lopez[2], Jianqiang Shao[3], Zer Vue[2], Antentor Hinton Jr.[2], Jason Mears[4], and Vivian Gama[1,5,6,7]

[1]Vanderbilt University, Cell and Developmental Biology, Nashville, TN.
[2]Vanderbilt University, Molecular Physiology and Biophysics, Nashville, TN.
[3]Central Microscopy Research Facility, University of Iowa, Iowa City, IA.
[4]Case Western Reserve University, Department of Pharmacology, Cleveland, OH.
[5]Vanderbilt University, Vanderbilt Center for Stem Cell Biology, Nashville, TN
[6]Vanderbilt University, Vanderbilt Brain Institute, Nashville, TN

[7]Corresponding author

Correspondence should be addressed to:

Vivian Gama
PMB407935
465 21st Avenue South, 4150A
Vanderbilt University
Nashville, TN 37240-7935
Tel: (615) 875-9490
vivian.gama@vanderbilt.edu

Running title: DRP1 mutations associated with EMPF1

Keywords: fibroblasts, DRP1, mitochondria, peroxisomes, cristae, glycolysis, oxidative phosphorylation

Abstract

Mitochondria and peroxisomes are both dynamic signaling organelles that constantly undergo

fission. While mitochondrial fission is known to coordinate cellular metabolism, proliferation, and

apoptosis, the physiological relevance of peroxisome dynamics and the implications for cell fate

are not fully understood. DRP1 (dynamin-related protein 1) is an essential GTPase that executes

both mitochondrial and peroxisomal fission. Patients with *de novo* heterozygous missense

mutations in the gene that encodes DRP1, *DNM1L*, present with encephalopathy due to defective

mitochondrial and peroxisomal fission (EMPF1). EMPF1 is a devastating neurodevelopmental

disease with no effective treatment. To interrogate the mechanisms by which DRP1 mutations

cause cellular dysfunction, we utilized human-derived fibroblasts from patients with mutations in

DRP1 who present with EMPF1. As expected, patient cells display elongated mitochondrial

morphology and lack of fission. Patient cells display a lower coupling efficiency of the electron

transport chain, increased proton leak, and upregulation of glycolysis. In addition to these

metabolic abnormalities, mitochondrial hyperfusion results in aberrant cristae structure and

hyperpolarized mitochondrial membrane potential, both of which are tightly linked to the changes

in metabolism. Peroxisome structure is also severely elongated in patient cells and results in a

potential functional compensation of fatty acid oxidation. Understanding the mechanism by which

DRP1 mutations cause these metabolic changes will give insight into the role of mitochondrial

dynamics in cristae maintenance and the metabolic capacity of the cell, as well as the disease

mechanism underlying EMPF1.

Introduction

Mitochondrial dynamics are controlled and executed by a vast group of proteins and other

organelles. The underlying morphology of the mitochondria is an important determinant for the

ability of this organelle to regulate a variety of cellular processes, such as apoptosis, cell fate, lipid

biosynthesis, and epigenetic modifications (Bayraktar et al., 2019; Iwata et al., 2020; Khacho et

al., 2016; King et al., 2021; Prigione and Adjaye, 2011; Rasmussen et al., 2020; Vantaggiato et

al., 2019). Mitochondrial fusion is primarily catalyzed by the large GTPases Mitofusin 1 and

Mitofusin 2 at the outer mitochondrial membrane (Chen et al., 2003). The GTPase OPA1 is

responsible for executing fusion of the inner mitochondrial membrane (Alexander et al., 2000;

Delettre et al., 2000). Mitochondrial fission, mediated through the large GTPase DRP1 (Dynamin-

related protein 1), is orchestrated by a complex sequence of events (Fonseca et al., 2019;

Smirnova et al., 2001). The first steps of mitochondrial fission include contact with the

endoplasmic reticulum (ER) and replication of the mitochondrial genome (Friedman et al., 2011;

Lewis et al., 2016). This is followed by contact and pre-constriction by actin filaments, which are

associated with the ER membrane (Chakrabarti et al., 2018; Hatch et al., 2016, 2014; Ji et al.,

2017, 2015; Li et al., 2015). Fission is also regulated through adaptor proteins at the outer

mitochondrial membrane, including MiD49, MiD51, MFF, and FIS1, however, their distinct roles

in fission are not completely understood (Losó n et al., 2013; Osellame et al., 2016; Palmer et al.,

2013). After the association with adaptor proteins, DRP1 then forms a 12-dimer ring around the

mitochondria and hydrolyzes GTP to GDP causing a confirmational change in the ring that

executes fission (Chang and Blackstone, 2007; Fröhlich et al., 2013; Nagashima et al., 2020).

DRP1 is essential for mitochondrial fission and without functional DRP1, the mitochondrial

network becomes hyperfused. In addition to the effects on mitochondrial morphology, DRP1 also

mediates the fission of peroxisomes (Koch et al., 2003). Peroxisomes are highly specialized

organelles vital for the beta oxidation of very-long-chain fatty acids, detoxification of reactive

oxygen species (ROS), production of cholesterol, bile acids, and plasmalogens, which contribute

to the phospholipid content in the brain white matter (Chandel, 2021; Ding et al., 2021; Ferreira

et al., 2021; Lodhi and Semenkovich, 2014; Ravi et al., 2021; Schrader et al., 2020; Sugiura et

al., 2017). The adaptor proteins MFF and FIS1 are also involved in DRP1 recruitment to sites of

peroxisome fission (Gandre-Babbe and Van Der Bliek, 2008; Itoyama et al., 2013; Koch and

Brocard, 2012). The contacts between mitochondria and peroxisomes are essential for linking

peroxisomal β-oxidation and mitochondrial ATP generation by the ETC, maintenance of redox

homeostasis, and lipid transfer (Chandel, 2021; Fan et al., 2016; Fransen et al., 2017). Thus,

peroxisomal and mitochondrial metabolic activities are functionally interconnected. Consistent

with this, peroxisomal disorders caused by mutations in genes involved in the biogenesis and

function of peroxisomes, are often associated with mitochondrial dysfunction (Tanaka et al.,

2019). This communication may be particularly important during brain development and metabolic

stress. However, the molecular mechanisms that facilitate this communication are still poorly

understood.

In mouse models, global DRP1 knockout is embryonic lethal (Ishihara et al., 2009; Wakabayashi

et al., 2009). With the advent of exome sequencing, patients with *de novo* mutations in the gene

that encodes DRP1, *DNM1L (Dynamin 1 Like)*, have been identified (Chao et al., 2016; Fahrner

et al., 2016; Keller et al., 2021; Liu et al., 2021; Longo et al., 2020; Nasca et al., 2016; Nolan et

al., 2019; Ryan et al., 2018; Sheffer et al., 2016; Vandeleur et al., 2019; Waterham et al., 2007).

The disease associated with mutations in *DNM1L* is known as encephalopathy due to defective

mitochondrial and peroxisomal fission-1 (EMPF1). Most reported mutations are heterozygous

missense   mutations.   Patients   present   with   heterogeneous   symptoms   including

neurodevelopmental delay, seizures, muscle abnormalities, and ataxia. Patient fibroblasts show

very elongated mitochondrial networks and elevated lactate levels, suggesting an impairment in

oxidative phosphorylation (Liu et al., 2021; Whitley et al., 2018). Currently, there is no effective

treatment or cure for patients effected by DRP1 mutations. The life expectancy for most patients

with DRP1 mutations is childhood to early adolescence (Ryan et al., 2018). Although other

proteins and organelles are involved in fission, DRP1 has been shown to be indispensable for

mitochondrial fission (Csordás et al., 2006; Fonseca et al., 2019; Kamerkar et al., 2018; Li et al.,

2015; Nagashima et al., 2020; Smirnova et al., 2001), yet the cellular consequences of DRP1

dysfunction are not well understood. In this study, we aimed to characterize the effects of two

mutations in DRP1 identified in patients with EMPF1, the G32A mutation, which lies in the catalytic

GTPase domain of DRP1, and the R403C mutation, which lies in the stalk domain (Fröhlich et al.,

2013), on mitochondrial network morphology, cristae ultrastructure, and related metabolic

adaptations, to gain fundamental insight into the cellular adaptations to dysfunctional organelle

fission and the pathophysiology of EMPF1.

Here, we resolve and quantify mitochondrial and peroxisomal morphological features associated

with EMPF1-associated mutations and reveal their impact on cellular metabolism, to put into

perspective the clinical findings that were previously described (Fahrner et al., 2016; Liu et al.,

2021; Ryan et al., 2018). We found that the inability of DRP1 to fragment the mitochondria in

patient-derived fibroblasts leads not only to the hyperfusion phenotype previously described, but

also to delayed cell death in response to apoptotic signals, aberrant mitochondrial cristae

structure, and hyperpolarized mitochondrial membrane potential. Metabolically, the changes in

mitochondrial and peroxisomal morphology result in inefficiency of the electron transport chain

coupling oxygen consumption with ATP production and upregulation of glycolysis. Altogether,

these data indicate that mitochondrial and peroxisomal fission influence dynamics of the inner

mitochondrial membrane and the metabolic state of the cell.

Results

DRP1 variants found in patients have dominant negative effects that alter the expression

of critical proteins at the mitochondria

Light microscopy studies have shown the profound alterations of mitochondrial fission in cells

from patients harboring mutations in DRP1 (Ryan et al., 2018; Whitley et al., 2018). These studies

highlighted the clinical relevance of mitochondrial fission in neural development. We quantified

the morphology of the mitochondria in patient-derived primary fibroblasts at super resolution level

using structured illumination microscopy (SIM). Both patient fibroblast lines showed increased

mitochondrial volume, surface area and major axis length, consistent with a hyperfused

mitochondrial morphology, as reported for DRP1 null cells (Fonseca et al., 2019; Kamerkar et al.,

2018; Smirnova et al., 2001) (Fig. 1A-B). Live-cell imaging of mitochondria stained with

MitoTracker confirmed that mitochondrial fission events are rare in patient cells compared to

control fibroblasts (Supplementary Fig. 1A). High resolution three-dimensional images of control

and patient fibroblasts were obtained using serial block facing-scanning electron microscopy

(SBF-SEM) (Garza-Lopez et al., 2021). Manual segmentation and stack reconstruction of the

mitochondrial networks validated the increased length and volume of the mitochondrial networks

as well as increased mitochondrial complexity index (MCI) in patient fibroblasts (Fig. 1C-D). MCI

is a quantitative measurement of network complexity, which takes into account both volume and

surface area of the mitochondria (MCI = $SA^3 / 16\pi^2V^2$) (Faitg et al., 2021; Vincent et al., 2019).

Both mutations in DRP1 result in significant alterations of the mitochondrial network with G32A

fibroblasts showing slightly more severe elongation than R403C (Fig. 1D and Supplementary Fig.

1B).

Both mutations in DRP1 are found to be heterozygous in patients, suggesting that mutant DRP1

has a dominant effect that can interfere with the ability of wild-type DRP1 to fragment the

mitochondria. Previous studies have demonstrated that the GFP can interfere with the activity of

DRP1 (Montecinos-Franjola et al., 2020). Thus, we generated mCherry tagged- constructs

containing the patient derived mutations and expressed them in human fibroblasts homozygous

for endogenous wild-type DRP1. Overexpression of mCherry-DRP1-G32A and mCherry-DRP1-

R403C by transient transfection in wild-type fibroblasts demonstrated that, as previously

described (Fahrner et al., 2016; Ryan et al., 2018; Whitley et al., 2018), the mitochondrial

hyperfusion seen in patient cells is not due to the genetic background and that these mutations

act in a dominant negative manner (Fig. 1E-F). While both mutations of DRP1 significantly

impaired DRP1 function in human fibroblasts, the G32A show a slightly more severe phenotype

than R403C.

Hyperfusion of the mitochondrial membrane due to sequestration and inactivation of DRP1 is

thought to lead to unopposed fusion events that increase the severity of mitochondrial elongation

(Palmer et al., 2013). We examined the effects of DRP1 mutations on the protein expression

levels of mitochondrial dynamics proteins. Both Mitofusin 1 and Mitofusin 2 were expressed at

lower levels in patient cells than in control cells. However, there was no significant difference in

the expression of OPA1 (Fig. 2A). These data show that there is a reduction in the expression of

mitochondrial outer membrane fusion proteins in patient cells, suggesting that active

mitochondrial fission promotes protein stability at the outer mitochondrial membrane. These data

also indicate that aberrant fission caused by DRP1 mutations may not alter fusion dynamics at

the inner mitochondrial membrane.

Phosphorylation of DRP1 at Serine 616 has been proposed to be a critical determinant of DRP1

recruitment and activity (Koch and Brocard, 2012; Taguchi et al., 2007; Valera-Alberni et al.,

2021). Whether mutations in DRP1 found in patients with EMPF1 affect the levels of

phosphorylation at Serine 616 has not been explored. We examined DRP1 levels in patient

derived fibroblasts. Although there was no significant difference in total DRP1 expression, G32A

patient cells expressed significantly less pS616-DRP1 (Fig. 2B). Since G32A mutation impairs

DRP1 phosphorylation to a greater extent than the R403C mutation, the GTPase domain of DRP1

may be structurally or functionally involved in interactions with kinases responsible for DRP1

phosphorylation. We wondered whether the expression of one copy of mutant DRP1 in patient

fibroblasts affected the recruitment of endogenous pS616-DRP1 to the mitochondria. To test this,

we quantified the recruitment of phosphorylated DRP1 to the outer mitochondrial membrane using

super-resolution microscopy. On average, there was no significant difference in the recruitment

of pS616-DRP1 to the mitochondria in patient cells and control cells (Fig. 2C-D). While in all cell

lines, pS616-DRP1 was found in close contact to the mitochondrial membrane, in the patient

fibroblasts this interaction appeared less transient. These results point to generalized stalling of

the fission events in patient fibroblasts.

The onset of apoptosis is accompanied by mitochondrial remodeling, stable association of DRP1

to the mitochondrial membrane, and increased fragmentation of the mitochondrial membrane

(Martinou and Youle, 2006; Wasiak et al., 2007). In DRP1 deficient cells, this remodeling of

mitochondria is significantly reduced and the onset of cell death is delayed (Lee et al., 2004). To

determine the effects of mitochondrial stress on mitochondrial fragmentation in patient fibroblasts,

we quantified changes in mitochondrial fission in response to CCCP treatment. As previously

shown using RNAi experiments in HeLa cells, patient fibroblasts showed reduced mitochondrial

fragmentation induced by CCCP compared to control cells (Fig. 2E-F). Although some

mitochondrial fragmentation may occur, mitochondrial volume and major axis length are

significantly larger in patient cells treated with CCCP than control cells. Upon apoptosis, BAX

translocates to the mitochondria and cytochrome c is released to the cytosol (Smaili et al., 2001).

While in DRP1 depleted cells BAX is recruited to the mitochondria, cytochrome c release is

delayed. Consistent with this, patient fibroblasts show higher cell survival when treated with CCCP

and Etoposide; as well as, lower caspase-3 activity than control fibroblasts when treated with STS

(Supplementary Fig. 2A-B). Interestingly, basal levels of both BAX and BAK are significantly

reduced in patient fibroblasts with mutations in DRP1 (Supplementary Fig. 2C). Levels of the anti-

apoptotic protein MCL-1 were significantly reduced in G32A fibroblasts but not in R403C

fibroblasts, while BCL-XL expression levels remained relatively constant in all cell lines. Thus,

patient fibroblasts may display a compensatory adaptation to reduce the basal levels of BAX/BAK

activation. Whether this reduction in the ability to undergo apoptosis in response to mitochondrial

stress is a widespread phenomenon in all tissues of EMPF1 patients and/or whether apoptosis

inhibition results in deleterious accumulation of mitochondrial and nuclear DNA encoded

mutations remains unexplored. No significant alteration in the protein expression of DRP1

adaptors was detected (Supplementary Fig. 2D)

Mitochondrial hyperfusion in DRP1 mutant fibroblasts leads to higher glycolytic rate and

increased mitochondrial membrane potential

Mitochondrial morphology is tightly linked to metabolic function (Dorn, 2015; Wai and Langer,

2016); thus, we assessed the effects of DRP1 patient mutations on cellular metabolism by

measuring rates of glycolysis and oxidative phosphorylation (OXPHOS) in cells using a Seahorse

XF Analyzer. The Seahorse analyzer measures extracellular oxygen consumption as a readout

of OXPHOS and extracellular acidification as a readout of glycolysis. Basal and maximal oxygen

consumption was not significantly different in mutant cells compared to control, suggesting no

major defects in ATP production (Fig. 3A). However, proton leak was significantly higher in the

G32A patient cells and coupling efficiency was significantly lower in both patient lines (Fig. 3B).

Therefore, although basal ATP production is not perturbed in the mutant cells, the electron

transport chain is not effectively coupling substrate oxidation and ATP synthesis. Extracellular

acidification rate was significantly increased in both patient lines, consistent with an upregulation

of glycolysis (Fig. 3A) and the hyperlactacidemia observed in more than 50% of patients with

mutations in DRP1 (Liu et al., 2021). A glycolytic rate assay confirmed that both basal and

compensatory glycolysis are higher in both patient lines (Fig. 3C). Thus, although the cells are

capable of producing ATP, DRP1 mutations cause a defect in OXPHOS efficiency and metabolic

dependency on glycolysis (Fig. 3D).

The upregulation of glycolysis with no accompanied increase in oxidative phosphorylation led us

to interrogate a potential impairment in pyruvate transport. Two proteins are essential for

mitochondrial pyruvate transport in humans, Mitochondrial Pyruvate Carrier 1 and 2 (MPC1 and

MPC2) (Bricker et al., 2012; Herzig et al., 2012). MPC1 and MPC2 associate to form an ~150-

kilodalton complex in the inner mitochondrial membrane. We examined the expression of MPC1

and MPC2, using Western blot, which revealed a significant decrease in MPC2 expression in the

R403C cells (Fig. 3E). Real time quantitative PCR (RT-qPCR) gene expression analysis showed

a significant downregulation of *MPC2* expression in R403C cells and *MPC1* in both patient lines

(Fig. 3F). Taken together these data suggest that the upregulation in glycolysis seen in DRP1

mutant patient fibroblasts is a compensatory mechanism for impaired pyruvate import into the

mitochondria and/or decreased coupling efficiency of the ETC.

The major contributor of the proton motive force of the ETC is the mitochondrial membrane

potential, which drives oxidative phosphorylation. Thus, changes in OXPHOS coupling efficiency

can point to perturbation of the mitochondrial membrane potential and the maintenance of key ion

gradients. We investigated this possibility using the fluorescent sensor Tetramethylrhodamine,

ethyl ester (TMRE). TMRE is a cell-permeant, cationic, red-orange fluorescent dye that is

sequestered by polarized mitochondria. We found that mitochondrial membrane potential as

reported by TMRE was significantly increased in patient fibroblasts (Fig. 4), indicative of

hyperpolarization of the mitochondria.

DRP1 mutations alter mitochondrial cristae morphology

Cristae topology readjustments participate in energy balance control (Eramo et al., 2020; Kondadi

et al., 2020). The hyperpolarization of the mitochondrial membrane detected in patient fibroblasts

lead us to assess the effects of DRP1 mutations on cristae morphology. We used a previously

optimized method to quantify and analyze cristae morphology changes from transmission electron

microscopy (TEM) (Eisner et al., 2017; Lam et al., 2021). In this method, cristae abundance is

calculated by manually quantifying the number of cristae per mitochondrion, and a score between

0 (worst) and 4 (best) is assigned based on the morphology: (0—no well-defined cristae, 1—no

cristae in more than ~75% of the mitochondrial area, 2—no cristae in approximately ~25% of

mitochondrial area, 3—many cristae (over 75% of area) but irregular, 4—many regular cristae)

(Supplementary. Fig 3A). TEM of patient fibroblasts confirmed the mitochondrial elongation

observed in SIM (Supplementary Fig. 3B-D) and revealed differences in the cristae structure (Fig.

5A). The quantification of cristae morphology demonstrated a reduced cristae score and

decreased cristae number relative to mitochondrial length in DRP1-deficient cells compared with

the control cells (Fig. 5B). These data link aberrant outer mitochondrial membrane fission caused

by DRP1 mutations to generalized cristae structural changes. The structure and respiratory

function of mitochondrial cristae are maintained by the m̲itochondrial contact sites and c̲ristae

o̲rganizing s̲ystem (MICOS) complex – a large complex of scaffolding and membrane shaping

proteins localized at the inner mitochondrial membrane. Given the reduced cristae score, we

examined expression of some key components of the MICOS complex. *MIC19, MIC25,* and

MIC60 are all reduced in either one or both patient lines at the mRNA level; however, protein

expression is not significantly changed (Supplementary. Fig. 4). The data suggest that the

structural abnormalities seen in patient fibroblasts are not due to changes in the expression of

proteins of the MICOS complex. Additional biochemical assays are needed to assess the

functionality of these proteins in the context of EMPF1.

Patient fibroblasts display hyperfused peroxisomes

In addition to catalyzing mitochondrial fission, DRP1 also participates in peroxisomal fission

(Kamerkar et al., 2018). While fission of peroxisomes has been demonstrated to be involved in

de novo peroxisomal biogenesis, the functional consequences of peroxisomal fission are not

completely understood. Previous reports using EMPF1 patient fibroblasts have not reported

elongation of peroxisomes, but DRP1 null cells have been reported to have hyperfused

peroxisomes (Kamerkar et al., 2018). To interrogate the effects of DRP1 mutations on peroxisome

morphology, we performed super-resolution microscopy. SIM revealed that both mutations in

DRP1 result in significant elongation of peroxisomes (Fig. 6A). Quantification of the peroxisome

structure shows an increase in volume, surface area, and length of peroxisomes (Fig. 6B). These

data confirm that DRP1 mutations found in patients with EMPF1 result in impaired peroxisome

fission.

Since peroxisomes and mitochondria are hubs for the regulation of fatty acid metabolism, we first

tested whether the dramatic changes in morphology would alter the expression of key genes

involved in fatty acid oxidation in patient fibroblasts. The expression of these genes have

previously been reported to increase in a DRP1 phosphomimetic mouse model (Valera-Alberni et

al., 2021). RT-qPCR analysis showed that the expression of carnitine palmitoyltransferase I,

CPT1, and acyl-coA dehydrogenase medium chain, *ACADM*, are significantly lower in patient

cells compared to control (Fig. 6C). CPT1 is associated with the outer mitochondrial membrane

and is responsible for catalyzing the transfer of the acyl group of acyl-CoA to L-carnitine. Fatty

acids are then able to move from the cytosol into the inner membrane space of the mitochondria

and the matrix, where fatty acid oxidation occurs. *ACADM* encodes for the medium-chain-acyl-

CoA dehydrogenase (MCAD) protein, which catalyzes the first step of mitochondrial fatty acid

beta-oxidation. Thus, the reduction in *CPT1* and *ACADM* expression may hinder the ability of the

mitochondria to carryout fatty acid oxidation.

To examine the ability of patient fibroblasts to perform fatty acid oxidation, we used the Seahorse

analyzer to quantify oxygen consumption rates of patient fibroblasts treated with either vehicle or

etomoxir, a CPT1 inhibitor. The cells were cultured with minimal glucose and glutamine and high

L-carnitine and palmitate to promote fatty acid oxidation over glycolysis or glutaminolysis. Control

fibroblasts treated with etomoxir, an inhibitor of CPT1, show a reduction in maximal respiration

following FCCP injection. In contrast, the G32A patient-derived cells show no change in OCR with

etomoxir treatment and the R403C patient-derived cells show an increase in maximal respiration

following FCCP injection (Fig. 6D and Supplementary Fig. 5). These data indicate that in contrast

to the reliance of control cells to long chain fatty acid oxidation under conditions of maximal

respiration, both patient lines have lost this dependency and adapted to the inhibition of long-

chain fatty acid import into mitochondria under these conditions. These data show that fibroblasts

with DRP1 mutations appear to have compensatory adaptations to alterations in long-chain fatty

acid oxidation.

Discussion

DRP1 is an essential gene for embryonic development (Ishihara et al., 2009; Wakabayashi et al., 2009). RNAi based approaches in human cell culture models have been used to explore the impact of DRP1 loss on mitochondrial fission and downstream processes such as apoptosis and mitophagy (Kleele et al., 2021; Otera et al., 2016; Smirnova et al., 2001). Although loss of DRP1 results in mitochondrial hyperfusion, it's not fully understood how mutations in DRP1 can impact cellular metabolic function, cristae structure, and/or peroxisomal function. The analysis of organelle morphology in fibroblast lines derived from patients with EMPF1 that harbor one wild-type copy of DRP1 and one missense copy of DRP1, demonstrate a dominant negative effect of mutant DRP1 that results in stalling of the fission machinery that has a wide impact on cellular function.

Mitochondrial fission is essential for cellular processes such as apoptosis, mitophagy, and mitochondrial biogenesis. However, how mitochondrial fission helps maintain cristae structure and/or cellular metabolism have not been thoroughly explored. Our studies revealed that while in EMPF1 patient lines, DRP1 is recruited to the mitochondria, the fission event is incomplete, and thus, mitochondrial fission is paused. This stalling of mitochondrial fission results in dramatic changes to cristae morphology, notably a decrease in cristae score and number, indicative of immature and poorly formed cristae. Changes in redox signaling can lead to cysteine oxidation of MIC60 (Li et al., 2021; Wasilewski and Chacinska, 2021), an inner mitochondrial membrane protein responsible for cristae structure integrity. Thus, the lack of proper mitochondrial fission in patient cells may disrupt redox signaling, which may in turn be responsible for the aberrant cristae morphology detected in patient cells.

Detailed metabolic analysis of patient fibroblasts showed that these cells have decreased coupling efficiency and upregulation of glycolysis, suggesting that the functional impact of stalled fission is electron transport chain inefficiency. This inefficiency may be due to the changes in

cristae structure, since cristae structure is essential for F_0F_1–ATPase to function effectively (Afzal

et al., 2021; Cogliati et al., 2013; Davies et al., 2011; Wilkens et al., 2013). Previous studies

revealed that there are distinct fission signatures that are key determinants of the decision of

mitochondria to proliferate or degrade (Kleele et al., 2021). Peripheral fission events were shown

to be preceded by a decrease in membrane potential and proton motive force. We found that the

elongated mitochondria from patient fibroblasts have a hyperpolarized membrane potential, which

could result from inefficient function of the F_0F_1–ATPase. Under certain conditions, the F_0F_1–

ATPase can pump protons out of the mitochondrial matrix into the intermembrane space in a

reverse-mode action, causing ΔΨm hyperpolarization. Alternatively, hyperpolarization may be

due to complex I inhibition or VDAC (voltage-dependent anion channel) opening, which can cause

hyperpolarization independently of F_0F_1–ATPase reverse-mode action (Forkink et al., 2014). A

comprehensive analysis of the functionality of the ETC complexes would be needed to elucidate

the connection between fission stalling events and mitochondrial hyperpolarization. Upregulation

of glycolysis in these patient fibroblasts could result in subsequent glucose depletion locally within

the cells, which could in turn cause ΔΨm hyperpolarization. Carbon-tracing could help reveal

whether all of the products of glycolysis are efficiently shuttled into the TCA cycle or if

intermediates are lost in the process. Investigating the function of uncoupling proteins such as

UCP2 and UCP3 may elucidate whether the decreased efficiency is a byproduct or cause of

mitochondrial dysfunction, such as increased ΔΨm, in these cells. Our finding that cells with

DRP1 mutations have profound alterations in cristae morphology but apparent maintenance of

overall ATP production, raises the interesting possibility that DRP1 mutant cells have evolved

specialized mechanisms to cope with electron leakage and damaging reactive oxygen species. A

landmark study from Spinelli and collaborators demonstrated the ability of complex II to work in

reverse in some tissues with the intrinsic capacity to use fumarate as a terminal electron acceptor

(Spinelli et al., 2021). It is plausible that complex II reversal has evolved as a mechanism to

maintain mitochondrial function in cells with alterations in fission caused by DRP1 mutations.

The existence of two mechanistically and functionally distinct types of fission at the mitochondria

and the peroxisomes have not been clearly elucidated. In our study, we demonstrate that patient

fibroblasts with mutations in DRP1 also have elongated peroxisomal morphology. Previous

studies in mammalian cell culture revealed that peroxisome membrane elongation precedes

membrane constriction and final membrane fission (Delille et al., 2010; Schrader et al., 2016).

Whether peroxisomes have a specialized fusion machinery is still under debate. However, we

speculate that under particular metabolic conditions peroxisomes, particularly those in contact

with mitochondria, may be able to fuse to respond to cellular metabolic needs. Elongation of the

peroxisomes in various human disorders has been associated with alterations in fatty acid

oxidation (Waterham et al., 2016). However; in these cells, peroxisomal function cannot be fully

dissociated from changes in mitochondrial morphology. Our model (Figure 7) suggests that the

mechanistic and functional connection between peroxisome remodeling and metabolic

homeostasis could be dysregulated in mitochondrial diseases such EMPF1. We propose that in

this context, human diseases could provide insight into the causal mechanisms underlying basic

cellular processes, such as organelle fission. The detailed elucidation of DRP1-centric

mechanisms will reveal the connection between organelle dynamics and cell fate. Several

questions remain: Does stalling of mitochondrial fission affect the association/dissociation of other

proteins at point of fission? Is peripheral fission also inhibited in patient fibroblasts? If that is the

case, what are the consequences for mitophagy and mitochondrial DNA integrity? What are the

effects of peroxisomal elongation on fatty acid metabolism? How is cell division and mitochondrial

segregation affected by DRP1 mutations? How are cell fate transitions affected by DRP1

mutations? Answering these fundamental questions would provide insight into the mechanisms

of mitochondrial fission under the lens of this rare disease caused by dysregulated mitochondrial

and peroxisomal homeostasis, and inform the field on more rational and specific therapeutic

targets for these devastating neurodevelopmental disorders.

STAR Methods

Cell Culture

Human fibroblasts were thawed and plated directly in maintenance medium consisting of MEM (Gibco #11095098) with 10% FBS (Sigma-Aldrich #F2442) and non-essential amino acids (Gibco # 11-140-050). Cells were maintained at 37°C and 5% CO_2 with media changes every other day. Cells were passaged as needed using TrypLE (Thermo Fisher #12-064-013). For imaging experiments, cells were re-plated onto #1.5 glass-bottom 35 mm dishes (Cellvis #D35C4-20-1.5-N).

Cell Treatments

All treatments were added directly to fibroblasts in maintenance medium. The electron transport chain decoupler CCCP (Sigma-Aldrich #C2759-250MG) was added to cells at a concentration of 1 µM. Cells were treated for 2 hours prior to fixation and imaging. All stock solutions were prepared in DMSO.

Plasmid Transfection

Plasmid encoding mCherry-DRP1 (Addgene #49152) was transfected using FuGENE (Promega #E2311) as described in the manufacturer protocol. Patient mutations were introduced to plasmids using the In Fusion cloning kit (Takara Bio #638909) as described in the manufacturer protocol. Cells were maintained until optimal transfection efficiency was reached before cells were imaged.

Immunofluorescence

For immunofluorescence, cells were fixed with 4% paraformaldehyde for 20 min and permeabilized in 1% Triton-X-100 for 10 min at room temperature. After blocking in 10% BSA, cells were treated with primary and secondary antibodies using standard methods. Cells were mounted in Fluoromount-G (Electron Microscopy Sciences #17984-25) prior to imaging. Primary

antibodies used include mouse anti-Mitochondria (Abcam #ab92824), rabbit anti-Tom20 (Cell

Signaling Technology #42406S), rabbit anti-DRP1 (Cell Signaling Technology #8570S), rabbit

anti-HSP60 (Cell Signaling Technologies # 12165S), mouse anti-PMP70 (Thermo Fisher # MA5-

31368), and rabbit anti-pDRP1 S616 (Cell Signaling Technology #3455S). Secondary antibodies

used include donkey anti-mouse Alexa Fluor-488 (Thermo Fisher Scientific #A21202), donkey

anti-rabbit Alexa Fluor-488 (Thermo Fisher Scientific #A21206), and donkey anti-mouse Alexa

Fluor-546 (Thermo Fisher Scientific #A10036), and donkey anti-rabbit Alexa Fluor-546 (Thermo

Fisher Scientific #A10040).

Western Blot

Gel samples were prepared by mixing cell lysates with LDS sample buffer (Life Technologies,

#NP0007) and 2-Mercaptoethanol (BioRad #1610710) and boiled at 95°C for 5 minutes. Samples

were resolved on 4-20% Mini-PROTEAN TGX precast gels (BioRad #4561096) and transferred

onto PVDF membrane (BioRad #1620177). Antibodies used for Western blotting are as follows:

rabbit anti-DRP-1 (Cell Signaling Technologies #8570S), rabbit anti-pDRP-1 S616 (Cell Signaling

Technologies #3455S), rabbit anti-OPA1 (Cell Signaling Technologies #67589S), mouse anti-

MFN1 (Abcam #ab57602, mouse anti-MFN2 (Abcam #ab56889), rabbit anti-MFF (Proteintech

#17090-1-AP), rabbit anti-MiD49 (Proteintech #16413-1-AP), rabbit anti-MiD51 (Proteintech

#20164-1-AP), and rabbit anti-FIS1 (Proteintech #10956-1-AP), and mouse anti-HSP70

(Invitrogen #MA3-006).

Seahorse XF Analysis

Fibroblasts were plated onto Seahorse XF96 V3 PS cell culture microplates 2 days before the

assay at 50,000 cells per well. One hour prior to the assay, media was switched to XF DMEM

media containing 1 mM pyruvate, 2 mM glutamine, and 10 mM glucose. For the Seahorse XF

Mito Stress Test, oxygen consumption rate (OCR) was measured sequentially after addition of

1.0 µM oligomycin, 0.5 µM FCCP, and 0.5 µM rotenone plus Antimycin A. Protein leak and

coupling efficiency were calculated using OCR. For the Seahorse XF Glycolytic Rate Assay, basal

glycolytic rate was measured by extracellular acidification, and compensatory glycolysis was

measured after addition of 0.5 μM rotenone plus antimycin A. Treatment with 50 mM 2-DG was

used as a negative control. For the Seahorse XF Fuel Flex Test, glutamine dependency was

measured by sequential addition of 3.0 μM BPTES followed by 4.0 μM Etomoxir plus 2.0 μM

UK5099. Fatty acid dependency was measured by sequential addition of 4.0 μM Etomoxir

followed by 3.0 μM BPTES plus 2.0 μM UK5099. Glucose dependency was measured by

sequential addition of 2.0 μM UK5099 followed by 3.0 μM BPTES plus 4.0 μM Etomoxir. For the

Seahorse XF Long Chain Fatty Acid Oxidation Stress Test, 24 hours prior to assay cells were

switched to base media containing 0.5 mM glucose, 1.0 mM glutamine, 0.5 mM XF L-carnitine,

and 1.0 % fetal bovine serum. One hour prior to the assay, media was switched to XF DMEM

media containing 2 mM glucose, 0.5 mM L-Carnitine, and Palmitate-BSA. Oxygen consumption

rate (OCR) was measured sequentially after addition of 4.0 μM etomoxir, 1.5 μM oligomycin, 2.0

μM FCCP, and 0.5 μM rotenone plus Antimycin A.

RNA Extraction and cDNA Synthesis

Cells were washed with PBS and scraped into 1000μl Trizol reagent. 200μl of chloroform was

added and the samples were incubated at room temperature for 5 minutes. The samples were

spun down at 12,000 g and the aqueous phase was collected. 500μl of isopropanol was added

to precipitate RNA and the sample was incubated for 25 minutes at room temperature, followed

by centrifugation at 12,000 g for 10 min at 4°C. The RNA pellet was washed with 75% ethanol,

semi-dried, and resuspended in 30μl of DEPC water. After quantification and adjusting the volume

of all the samples to 1μg/μl, the samples were treated with DNAse (New England Biolabs, #

M0303). 10μl of this volume was used to generate cDNA using the manufacturer's protocol

(Thermo Fisher, #4368814).

Quantitative RT PCR (RT-qPCR)

1ug of cDNA sample was used to run RT-qPCR for the primers mentioned in the table.

QuantStudio 3 Real-Time PCR machine, SYBR green master mix (Thermo Fisher, #4364346),

and manufacturer instructions were used to set up the assay.

Image Acquisition

Super-resolution images were acquired using a Nikon SIM microscope equipped with a 1.49 NA

100x Oil objective an Andor DU-897 EMCCD camera. Live-imaging experiments experiments

were acquired on a Nikon Eclipse Ti-E spinning disk confocal microscope equipped with a 1.40

NA 60X Oil objective and an Andor DU-897 EMCCD camera. Image processing and quantification

was performed using Fiji.

Statistical Analysis

All experiments were performed with a minimum of 3 biological replicates. Statistical significance

was determined by one-way or repeated measures ANOVA as appropriate for each experiment.

GraphPad Prism v8.1.2 was used for all statistical analysis and data visualization. Error bars in

all bar graphs represent standard error of the mean unless otherwise noted in the figure. No

outliers were removed from the analyses. Quantification of mitochondrial morphology was

performed in NIS-Elements (Nikon); briefly, we segmented mitochondria in 3D and performed

skeletonization of the resulting 3D mask. Skeleton major axis length, volume, and surface area

measurements were exported into Excel.

Sample Preparation for Transmission Electron Microscope (TEM) Analysis

Human fibroblasts were immediately washed in ice-cold saline. Samples were processed as

described by (Lam et. al., 2021, Pereira et al., 2017). Cells were fixed in 2.5% glutaraldehyde in

sodium cacodylate buffer for 1 hour at 37 °C, embedded in 2% agarose, postfixed in buffered 1%

osmium tetroxide, stained in 2% uranyl acetate, and dehydrated with an ethanol graded series.

After embedding in EMbed-812 resin, 80 nm sections were cut on an ultramicrotome and stained

with 2% uranyl acetate and lead citrate. Images were acquired on a JEOL JEM-1230

Transmission Electron Microscope, operating at 120 kV. Random images of human fibroblasts

were obtained.

TEM Analysis

Measurements of mitochondrial area, circularity, and length were performed using the multi-

measure region of interest (ROI) tool in ImageJ (Lam et. al., 2021, Parra et al., 2014). To measure

cristae morphology, we used ROIs in ImageJ to determine the cristae area, circulatory index,

number, volume, and cristae score. The NIH ImageJ software was used to manually trace and

analyze all mitochondria or cristae using the freehand told in the ImageJ application (Lam et. al.,

2021, Parra et al., 2014). To assign a mitochondrion a cristae score, cristae abundance and form

were evaluated. A score between 0 and 4 (0 – no sharply defined crista, 1 – greater than 50% of

mitochondrial area without cristae, 2 – greater than 25% of mitochondrial area without cristae, 3

– many cristae but irregular, 4 – many regular cristae) was assigned (Eisner et al., 2017).

Serial Block Facing-Scanning Electron Microscopy (SBF-SEM) Processing

Human fibroblasts were fixed in 2% glutaraldehyde in 0.1 M cacodylate buffer and processed

using a heavy metal protocol adapted from a previously published protocol (Courson et al., 2021;

Mustafi et al., 2014). Human fibroblasts were immersed in 3% potassium ferrocyanide and 2%

osmium tetroxide (1 hour at 4 °C), followed by filtered 0.1% thiocarbohydrazide (20 min), 2%

osmium tetroxide (30 min), and left overnight in 1% uranyl acetate at 4 °C (several de-ionized

H2O washes were performed between each step). The next day samples were immersed in a

0.6% lead aspartate solution (30 min at 60 °C and dehydrated in graded acetone (as described

for TEM). Human fibroblasts were impregnated with epoxy Taab 812 hard resin, embedded in

fresh resin, and polymerized at 60 °C for 36–48 hours. After polymerization, the block was

sectioned for TEM to identify the area of interest, then trimmed to 0.5 mm × 0.5 mm and glued to

an aluminum pin. The pin was placed into an FEI/Thermo Scientific Volumescope 2 scanning

electron microscope (SEM). The scope is a state-of-the-art SBF imaging system. For 3D EM reconstruction, thin (0.09 µm) serial sections, 300 to 400 per block, were obtained from the blocks that were processed for conventional TEM. Serial sections were collected onto formvar coated slot grids (Pella, Redding CA), stained, and imaged, as described above. The segmentation of SBF-SEM reconstructions was performed by manually tracing structural features through sequential slices of the micrograph block. Images were collected from 30–50 serial sections, which were then stacked, aligned, and visualized using Amira to make videos and quantify volumetric structures.

Statistical Analysis

The results are presented as the mean ± standard error of the mean. Data were analyzed using an unpaired T-test. If more than two groups were compared, one-way analysis of variance (ANOVA) was performed, and significance was assessed using Fisher's protected least significance difference test. GraphPad PRISM and Statplus software package was used for t-tests and ANOVA analyses (SAS Institute: Cary, NC, USA). For all statistical analyses, a significant difference was accepted when P < 0.05.

Acknowledgements

We would like to thank the patients and families for permission to publish this work. We thank Dr. Eric Payne and Dr. Suzanne Hoppins for providing the patient-derived fibroblast lines used in this paper. We would also like to thank Dr. Evan Krystofiak for providing EM expertise, and Dr. Matthew Tyska, Dr. Uri Manor, and Dr. Kari Seedle, for providing advice and expertise with high-resolution microscopy. This work was supported by 1R35 GM128915-01NIGMS (VG), 1RF1MH123971-01 (VG); the Precision Medicine and Mental Health Initiative sponsored by the Vanderbilt Brain Institute (VG), 1R21CA227483-01A1NCI (VG), F99NS125829 (GLR), HHMI Gilliam (GLR), and CBMS T32 (GLR). All SIM and spinning disk confocal microscopy imaging and image analysis were performed in part through the use of the Vanderbilt Cell Imaging Shared

535   Resource, which is supported by NIH grants 1S10OD012324-01 and 1S10OD021630-01. The

537   **Author Contributions**

Figure Legends

Figure 1. DRP1 patient-derived fibroblasts have elongated mitochondrial morphology. (A)

Representative images of mitochondrial morphology in patient-derived fibroblasts using

structured illumination microscopy (n=3, 7 cells per replicate). (B) Quantification of mitochondrial

volume, surface area, and major axis length. Analyzed using one-way ANOVA followed by

Dunnett's multiple comparison test. (C) Representative sbf-SEM 3D reconstructed images. Each

mitochondrion is colored a unique color. (D) Quantification of sbf-SEM for 3D mitochondrial

length, volume, and complexity index. Analyzed using one-way ANOVA followed by Dunnett's

multiple comparison test. (E) Representative images of mitochondrial morphology in control

fibroblasts transfected with either DRP1^{WT}-mCherry, DRP1^{G32A}-mCherry, or DRP1^{R403C}-mCherry

using structured illumination microscopy (n=3, 7 cells per replicate). (F) Quantification of

mitochondrial volume, surface area, and major axis length. Analyzed using one-way ANOVA

followed by Dunnett's multiple comparison test.

Figure 2. Mutant G32A and R403C DRP1 are recruited to mitochondria. i. (A) Western blot of

total protein lysate isolated from patient fibroblasts and probed for mitochondrial fusion proteins.

Representative image of three independent biological replicates. (B) Western blot of total protein

lysate isolated from patient fibroblasts and probed for mitochondrial fission proteins. Band density

is normalized to loading control (HSP70) and relative to control cells. Representative image of

three independent biological replicates. (C) Representative images of mitochondrial and DRP1 in

patient-derived fibroblasts using structured illumination microscopy (n=3, 7 cells per replicate).

Top image is a zoom of white box from bottom image. (D) Quantification of DRP1 puncta that

overlap with mitochondria and volume of DRP1 puncta. Analyzed using one-way ANOVA followed

by Dunnett's multiple comparison test. (E) Representative images of mitochondrial morphology in

patient-derived fibroblasts treated with either DMSO or CCCP for 1 hours using structured

illumination microscopy (n=3, 7 cells per replicate). (F) Quantification of mitochondrial volume,

surface area, and major axis length. Analyzed using one-way ANOVA followed by Dunnett's

multiple comparison test.

Figure 3. Mitochondrial defects lead to inefficient coupling of the electron transport chain

and higher glycolytic rate. (A) Oxygen consumption rate and extracellular acidification rate in

patient fibroblasts as measured using Seahorse Mitochondrial stress test (n=3, 20 wells per

replicate). Oligomycin applied at 20 minutes, FCCP applied at 40 minutes, and

rotenone/Antimycin A applied at 60 minutes. Analyzed using two-way ANOVA followed by Šídák's

multiple comparisons test. (B) Proton leak and coupling efficiency calculated using oxygen

consumption rate. (C) Basal glycolysis of patient fibroblasts measure using Seahorse Glycolytic

Rate Assay. Compensatory glycolysis measured following application of rotenone/Antimycin A.

(D) Glucose, glutamine, and fatty acid dependency of the electron transport chain measured using

Seahorse Fuel Flex Assay. Glucose dependency measured after application of UK5099,

glutamine dependency measured after application of BPTES, and fatty acid dependency measure

after application of Etomoxir. (E) qRT-PCR analysis of gene expression in fibroblasts relative to

control and normalized to two housekeeping genes (*GPI* and *GAPDH)*. Quantifications are shown

for three independent biological replicates. (F) Western blot of total protein lysate isolated from

patient fibroblasts and probed for mitochondrial pyruvate carriers. Band density is normalized to

loading control (HSP70) and relative to control cells. Representative image of three independent

biological replicates.

Figure 4. DRP1 patient fibroblasts have more hyperpolarized mitochondrial membrane

potential. (A) Representative live-cell imaging of patient-derived fibroblasts treated with 500nM

TMRE for 10 minutes. Cells were imaged every 15 seconds for 15 minutes using spinning disk

confocal (n=3, 7 cells per replicate). Scale bar: 5 μm.

Figure 5. Cristae morphology is altered in patient cells. (A) Transmission electron microscopy

of patient fibroblasts. 3 representative images per line shown. (B) Quantification of cristae from

TEM images for cristae score and cristae number relative to mitochondrial length. Analyzed using

one-way ANOVA followed by Dunnett's multiple comparison test.

Figure 6. Patient fibroblasts have elongated peroxisomes. (A) Representative images of

peroxisomal morphology in patient-derived fibroblasts using structured illumination microscopy

(n=3, 7 cells per replicate). (B) Quantification of peroxisomal volume, surface area, and major

axis length. Analyzed using one-way ANOVA followed by Dunnett's multiple comparison test. (C)

qRT-PCR analysis of gene expression in fibroblasts relative to control and normalized to two

housekeeping genes (*GPI* and *GAPDH)*. Quantifications are shown for three independent

biological replicates. (D) Oxygen consumption rate in patient fibroblasts as measured using

Seahorse Analyzer XFe96 (n=3, 20 wells per replicate). Etomoxir applied after 20 minutes,

oligomycin applied at 40 minutes, FCCP applied at 60 minutes, and rotenone/Antimycin A applied

at 80 minutes. Analyzed using two-way ANOVA followed by Šídák's multiple comparisons test.

Figure 7: Current model of the cellular responses to EMPF1 associated DRP1 mutations.

EMPF1- associated mutations do not appear to alter the capacity of DRP1 to be recruited to points

of fission, but rather cause stalling of the fission events. Pausing peroxisomal and mitochondrial

fission results in alterations of cristae structure, changes in mitochondrial membrane potential,

decreased mitochondrial coupling efficiency, delayed cell death response, and overall metabolic

adaptations.

References

Afzal N, Lederer WJ, Jafri MS, Mannella CA. 2021. Effect of crista morphology on mitochondrial ATP output: A computational study. *Curr Res Physiol* **4**:163–176. doi:10.1016/J.CRPHYS.2021.03.005

Alexander C, Votruba M, Pesch UEA, Thiselton DL, Mayer S, Moore A, Rodriguez M, Kellner U, Leo-Kottler B, Auburger G, Bhattacharya SS, Wissinger B. 2000. OPA1, encoding a dynamin-related GTPase, is mutated in autosomal dominant optic atrophy linked to chromosome 3q28. *Nat Genet 2000 262* **26**:211–215. doi:10.1038/79944

Bayraktar EC, Baudrier L, Özerdem C, Lewis CA, Chan SH, Kunchok T, Abu-Remaileh M, Cangelosi AL, Sabatini DM, Birsoy K, Chen WW. 2019. MITO-Tag Mice enable rapid isolation and multimodal profiling of mitochondria from specific cell types in vivo. *Proc Natl Acad Sci U S A* **116**:303–312. doi:10.1073/pnas.1816656115

Bricker DK, Taylor EB, Schell JC, Orsak T, Boutron A, Chen YC, Cox JE, Cardon CM, Van Vranken JG, Dephoure N, Redin C, Boudina S, Gygi SP, Brivet M, Thummel CS, Rutter J. 2012. A mitochondrial pyruvate carrier required for pyruvate uptake in yeast, Drosophila, and humans. *Science* **337**:96–100. doi:10.1126/SCIENCE.1218099

Chakrabarti R, Ji WK, Stan R V., Sanz J de J, Ryan TA, Higgs HN. 2018. INF2-mediated actin polymerization at the ER stimulates mitochondrial calcium uptake, inner membrane constriction, and division. *J Cell Biol* **217**:251–268. doi:10.1083/JCB.201709111

Chandel NS. 2021. Lipid Metabolism. *Cold Spring Harb Perspect Biol* **13**. doi:10.1101/CSHPERSPECT.A040576

Chang C, Blackstone C. 2007. Drp1 phosphorylation and mitochondrial regulation. *EMBO Rep* **8**:1088–1089. doi:10.1038/sj.embor.7401118

Chao YH, Robak LA, Xia F, Koenig MK, Adesina A, Bacino CA, Scaglia F, Bellen HJ, Wangler

MF. 2016. Missense variants in the middle domain of DNM1L in cases of infantile

encephalopathy alter peroxisomes and mitochondria when assayed in Drosophila. *Hum Mol*

Genet **25**:1846–1856. doi:10.1093/HMG/DDW059

Chen H, Detmer SA, Ewald AJ, Griffin EE, Fraser SE, Chan DC. 2003. Mitofusins Mfn1 and Mfn2

coordinately regulate mitochondrial fusion and are essential for embryonic development. *J*

Cell Biol **160**:189–200. doi:10.1083/JCB.200211046

Cogliati S, Frezza C, Soriano ME, Varanita T, Quintana-Cabrera R, Corrado M, Cipolat S, Costa

V, Casarin A, Gomes LC, Perales-Clemente E, Salviati L, Fernandez-Silva P, Enriquez JA,

Scorrano L. 2013. Mitochondrial Cristae Shape Determines Respiratory Chain

Supercomplexes Assembly and Respiratory Efficiency. *Cell* **155**:160–171.

doi:10.1016/J.CELL.2013.08.032

Csordás G, Renken C, Várnai P, Walter L, Weaver D, Buttle KF, Balla T, Mannella CA, Hajnóczky

G. 2006. Structural and functional features and significance of the physical linkage between

ER and mitochondria. *J Cell Biol* **174**:915–921. doi:10.1083/jcb.200604016

Davies KM, Strauss M, Daum B, Kief JH, Osiewacz HD, Rycovska A, Zickermann V, Kühlbrandt

W. 2011. Macromolecular organization of ATP synthase and complex I in whole

mitochondria. *Proc Natl Acad Sci U S A* **108**:14121–14126.

doi:10.1073/PNAS.1103621108/-/DCSUPPLEMENTAL

Delettre C, Lenaers G, Griffoin JM, Gigarel N, Lorenzo C, Belenguer P, Pelloquin L, Grosgeorge

J, Turc-Carel C, Perret E, Astarie-Dequeker C, Lasquellec L, Arnaud B, Ducommun B,

Kaplan J, Hamel CP. 2000. Nuclear gene OPA1, encoding a mitochondrial dynamin-related

protein, is mutated in dominant optic atrophy. *Nat Genet 2000 262* **26**:207–210.

doi:10.1038/79936

Delille HK, Agricola B, Guimaraes SC, Borta H, Lüers GH, Fransen M, Schrader M. 2010.

Pex11pbeta-mediated growth and division of mammalian peroxisomes follows a maturation

pathway. *J Cell Sci* **123**:2750–2762. doi:10.1242/JCS.062109

Ding L, Sun W, Balaz M, He A, Klug M, Wieland S, Caiazzo R, Raverdy V, Pattou F, Lefebvre P,

Lodhi IJ, Staels B, Heim M, Wolfrum C. 2021. Peroxisomal β-oxidation acts as a sensor for

intracellular fatty acids and regulates lipolysis. *Nat Metab* **3**:1648–1661.

doi:10.1038/S42255-021-00489-2

Dorn GW. 2015. Mitochondrial dynamism and heart disease: changing shape and shaping

change. *EMBO Mol Med* **7**:865–877. doi:10.15252/emmm.201404575

Eisner V, Cupo RR, Gao E, Csordás G, Slovinsky WS, Paillard M, Cheng L, Ibetti J, Chen SRW,

Chuprun JK, Hoek JB, Koch WJ, Hajnóczky G. 2017. Mitochondrial fusion dynamics is robust

in the heart and depends on calcium oscillations and contractile activity. *Proc Natl Acad Sci*

U S A **114**:E859–E868. doi:10.1073/PNAS.1617288114/-/DCSUPPLEMENTAL

Eramo MJ, Lisnyak V, Formosa LE, Ryan MT. 2020. The "mitochondrial contact site and cristae

organising system" (MICOS) in health and human disease. *J Biochem* **167**:243–255.

doi:10.1093/JB/MVZ111

Fahrner JA, Liu R, Perry MS, Klein J, Chan DC. 2016. A novel de novo dominant negative

mutation in DNM1L impairs mitochondrial fission and presents as childhood epileptic

encephalopathy. *Am J Med Genet A* **170**:2002–2011. doi:10.1002/AJMG.A.37721

Faitg J, Lacefield C, Davey T, White K, Laws R, Kosmidis S, Reeve AK, Kandel ER, Vincent AE,

Picard M. 2021. 3D neuronal mitochondrial morphology in axons, dendrites, and somata of

the aging mouse hippocampus. *Cell Rep* **36**.

doi:10.1016/J.CELREP.2021.109509/ATTACHMENT/28BF2465-59D0-4B68-BCDD-

D3F180754B5F/MMC1.PDF

Fan J, Li X, Issop L, Culty M, Papadopoulos V. 2016. ACBD2/ECI2-Mediated Peroxisome-

Mitochondria Interactions in Leydig Cell Steroid Biosynthesis. *Mol Endocrinol* **30**:763–782.

doi:10.1210/ME.2016-1008

Ferreira AR, Marques M, Ramos B, Kagan JC, Ribeiro D. 2021. Emerging roles of peroxisomes

in viral infections. *Trends Cell Biol*. doi:10.1016/J.TCB.2021.09.010

Fonseca TB, Sánchez-Guerrero Á, Milosevic I, Raimundo N. 2019. Mitochondrial fission requires

DRP1 but not dynamins. *Nature*. doi:10.1038/s41586-019-1296-y

Forkink M, Manjeri GR, Liemburg-Apers DC, Nibbeling E, Blanchard M, Wojtala A, Smeitink JAM,

Wieckowski MR, Willems PHGM, Koopman WJH. 2014. Mitochondrial hyperpolarization

during chronic complex I inhibition is sustained by low activity of complex II, III, IV and V.

Biochim Biophys Acta - Bioenerg **1837**:1247–1256. doi:10.1016/J.BBABIO.2014.04.008

Fransen M, Lismont C, Walton P. 2017. The Peroxisome-Mitochondria Connection: How and

Why? *Int J Mol Sci* **18**. doi:10.3390/IJMS18061126

Friedman JR, Lackner LL, West M, DiBenedetto JR, Nunnari J, Voeltz GK. 2011. ER tubules mark

sites of mitochondrial division. *Science (80-)* **334**:358–362.

doi:10.1126/SCIENCE.1207385/SUPPL_FILE/FRIEDMAN.SOM.REV1.PDF

Fröhlich C, Grabiger S, Schwefel D, Faelber K, Rosenbaum E, Mears J, Rocks O, Daumke O.

2013. Structural insights into oligomerization and mitochondrial remodelling of dynamin 1-

like protein. *EMBO J* **32**:1280–1292. doi:10.1038/EMBOJ.2013.74

Gandre-Babbe S, Van Der Bliek AM. 2008. The novel tail-anchored membrane protein Mff

controls mitochondrial and peroxisomal fission in mammalian cells. *Mol Biol Cell* **19**:2402–

2412.   doi:10.1091/MBC.E07-12-

1287/ASSET/IMAGES/LARGE/ZMK0060885640008.JPEG

Garza-Lopez E, Vue Z, Katti P, Neikirk K, Biete M, Lam J, Beasley HK, Marshall AG, Rodman
TA, Christensen TA, Salisbury JL, Vang L, Mungai M, AshShareef S, Murray SA, Shao J,
Streeter J, Glancy B, Pereira RO, Dale Abel E, Hinton A, Jr, Lopez G. 2021. Protocols for
Generating Surfaces and Measuring 3D Organelle Morphology Using Amira. *Cells 2022, Vol*
11, Page 65 **11**:65. doi:10.3390/CELLS11010065

Hatch AL, Gurel PS, Higgs HN. 2014. Novel roles for actin in mitochondrial fission. *J Cell Sci*
127:4549–4560. doi:10.1242/JCS.153791

Hatch AL, Ji WK, Merrill RA, Strack S, Higgs HN. 2016. Actin filaments as dynamic reservoirs for
Drp1 recruitment. *Mol Biol Cell* **27**:3109–3121. doi:10.1091/MBC.E16-03-0193

Herzig S, Raemy E, Montessuit S, Veuthey JL, Zamboni N, Westermann B, Kunji ERS, Martinou
JC. 2012. Identification and functional expression of the mitochondrial pyruvate carrier.
Science **337**:93–96. doi:10.1126/SCIENCE.1218530

Ishihara N, Nomura M, Jofuku A, Kato H, Suzuki SO, Masuda K, Otera H, Nakanishi Y, Nonaka
I, Goto YI, Taguchi N, Morinaga H, Maeda M, Takayanagi R, Yokota S, Mihara K. 2009.
Mitochondrial fission factor Drp1 is essential for embryonic development and synapse
formation in mice. *Nat Cell Biol* **11**:958–966. doi:10.1038/ncb1907

Itoyama A, Michiyuki S, Honsho M, Yamamoto T, Moser A, Yoshida Y, Fujiki Y. 2013. Mff
functions with Pex11pβ and DLP1 in peroxisomal fission. *Biol Open* **2**:998.
doi:10.1242/BIO.20135298

Iwata R, Casimir P, Vanderhaeghen P. 2020. Mitochondrial dynamics in postmitotic cells regulate
neurogenesis. *Science (80-)* **369**:858–862. doi:10.1126/science.aba9760

Ji WK, Chakrabarti R, Fan X, Schoenfeld L, Strack S, Higgs HN. 2017. Receptor-mediated Drp1
oligomerization on endoplasmic reticulum. *J Cell Biol* **216**:4123–4139.

doi:10.1083/JCB.201610057

Ji WK, Hatch AL, Merrill RA, Strack S, Higgs HN. 2015. Actin filaments target the oligomeric maturation of the dynamin GTPase Drp1 to mitochondrial fission sites. *Elife* **4**. doi:10.7554/eLife.11553

Kamerkar SC, Kraus F, Sharpe AJ, Pucadyil TJ, Ryan MT. 2018. Dynamin-related protein 1 has membrane constricting and severing abilities sufficient for mitochondrial and peroxisomal fission. *Nat Commun* **9**:1–15. doi:10.1038/s41467-018-07543-w

Keller N, Paketci C, Edem P, Thiele H, Yis U, Wirth B, Karakaya M. 2021. De novo DNM1L variant presenting with severe muscular atrophy, dystonia and sensory neuropathy. *Eur J Med Genet* **64**. doi:10.1016/J.EJMG.2020.104134

Khacho M, Clark A, Svoboda DS, Azzi J, MacLaurin JG, Meghaizel C, Sesaki H, Lagace DC, Germain M, Harper ME, Park DS, Slack RS. 2016. Mitochondrial Dynamics Impacts Stem Cell Identity and Fate Decisions by Regulating a Nuclear Transcriptional Program. *Cell Stem Cell* **19**:232–247. doi:10.1016/j.stem.2016.04.015

King WT, Axelrod CL, Zunica ERM, Noland RC, Davuluri G, Fujioka H, Tandler B, Pergola K, Hermann GE, Rogers RC, López-Domènech S, Dantas WS, Stadler K, Hoppel CL, Kirwan JP. 2021. Dynamin-related protein 1 regulates substrate oxidation in skeletal muscle by stabilizing cellular and mitochondrial calcium dynamics. *J Biol Chem* **297**:101196. doi:10.1016/J.JBC.2021.101196

Kleele T, Rey T, Winter J, Zaganelli S, Mahecic D, Perreten Lambert H, Ruberto FP, Nemir M, Wai T, Pedrazzini T, Manley S. 2021. Distinct fission signatures predict mitochondrial degradation or biogenesis. *Nature* **593**:435–439. doi:10.1038/S41586-021-03510-6

Koch A, Thiemann M, Grabenbauer M, Yoon Y, McNiven MA, Schrader M. 2003. Dynamin-like

752 protein 1 is involved in peroxisomal fission. *J Biol Chem* **278**:8597–8605.

753 doi:10.1074/JBC.M211761200

754 Koch J, Brocard C. 2012. PEX11 proteins attract Mff and human Fis1 to coordinate peroxisomal

755 fission. *J Cell Sci* **125**:3813–3826. doi:10.1242/JCS.102178/263234/AM/PEX11-

756 PROTEINS-ATTRACT-MFF-AND-HFIS1-TO-COORDINATE

757 Kondadi AK, Anand R, Hänsch S, Urbach J, Zobel T, Wolf DM, Segawa M, Liesa M, Shirihai OS,

758 Weidtkamp-Peters S, Reichert AS. 2020. Cristae undergo continuous cycles of membrane

759 remodelling in a MICOS-dependent manner. *EMBO Rep* **21**:e49776.

760 doi:10.15252/EMBR.201949776

761 Lam J, Katti P, Biete M, Mungai M, Ashshareef S, Neikirk K, Lopez EG, Vue Z, Christensen TA,

762 Beasley HK, Rodman TA, Murray SA, Salisbury JL, Glancy B, Shao J, Pereira RO, Abel ED,

763 Hinton, A., Jr. 2021. A Universal Approach to Analyzing Transmission Electron Microscopy

764 with ImageJ. *Cells 2021, Vol 10, Page 2177* **10**:2177. doi:10.3390/CELLS10092177

765 Lee YJ, Jeong SY, Karbowski M, Smith CL, Youle RJ. 2004. Roles of the mammalian

766 mitochondrial fission and fusion mediators Fis1, Drp1, and Opa1 in apoptosis. *Mol Biol Cell*

767 **15**:5001–5011. doi:10.1091/MBC.E04-04-0294

768 Lewis SC, Uchiyama LF, Nunnari J. 2016. ER-mitochondria contacts couple mtDNA synthesis

769 with Mitochondrial division in human cells. *Science (80-)* **353**.

770 doi:10.1126/SCIENCE.AAF5549

771 Li L, Conradson DM, Bharat V, Kim MJ, Hsieh CH, Minhas PS, Papakyrikos AM, Durairaj AS,

772 Ludlam A, Andreasson KI, Partridge L, Cianfrocco MA, Wang X. 2021. A mitochondrial

773 membrane-bridging machinery mediates signal transduction of intramitochondrial oxidation.

774 *Nat Metab* **3**:1242–1258. doi:10.1038/S42255-021-00443-2

Li S, Xu S, Roelofs BA, Boyman L, Jonathan Lederer W, Sesaki H, Karbowski M. 2015. Transient
assembly of F-actin on the outer mitochondrial membrane contributes to mitochondrial
fission. *J Cell Biol* **208**:109–123. doi:10.1083/jcb.201404050

Liu Xingmiao, Zhang Z, Li D, Lei M, Li Q, Liu Xiaojun, Zhang P. 2021. DNM1L-Related
Mitochondrial Fission Defects Presenting as Encephalopathy: A Case Report and Literature
Review. *Front Pediatr* **9**. doi:10.3389/FPED.2021.626657

Lodhi IJ, Semenkovich CF. 2014. Peroxisomes: a nexus for lipid metabolism and cellular
signaling. *Cell Metab* **19**:380–392. doi:10.1016/J.CMET.2014.01.002

Longo F, Benedetti S, Zambon AA, Sora MGN, Di Resta C, De Ritis D, Quattrini A, Maltecca F,
Ferrari M, Previtali SC. 2020. Impaired turnover of hyperfused mitochondria in severe axonal
neuropathy due to a novel DRP1 mutation. *Hum Mol Genet* **29**:177–188.
doi:10.1093/HMG/DDZ211

Losón OC, Song Z, Chen H, Chan DC. 2013. Fis1, Mff, MiD49, and MiD51 mediate Drp1
recruitment in mitochondrial fission. *Mol Biol Cell* **24**:659–667. doi:10.1091/MBC.E12-10-
0721

Martinou JC, Youle RJ. 2006. Which came first, the cytochrome c release or the mitochondrial
fission? *Cell Death Differ* **13**:1291–1295. doi:10.1038/SJ.CDD.4401985

Montecinos-Franjola F, Bauer BL, Mears JA, Ramachandran R. 2020. GFP fluorescence tagging
alters dynamin-related protein 1 oligomerization dynamics and creates disassembly-
refractory puncta to mediate mitochondrial fission. *Sci Reports 2020 101* **10**:1–16.
doi:10.1038/s41598-020-71655-x

Nagashima S, Tábara LC, Tilokani L, Paupe V, Anand H, Pogson JH, Zunino R, McBride HM,
Prudent J. 2020. Golgi-derived PI(4)P-containing vesicles drive late steps of mitochondrial

division. *Science (80-)* **367**:1366–1371. doi:10.1126/science.aax6089

Nasca A, Legati A, Baruffini E, Nolli C, Moroni I, Ardissone A, Goffrini P, Ghezzi D. 2016. Biallelic

Mutations in DNM1L are Associated with a Slowly Progressive Infantile Encephalopathy.

Hum Mutat **37**:898–903. doi:10.1002/HUMU.23033

Nolan DA, Chen B, Michon AM, Salatka E, Arndt D. 2019. A Rasmussen encephalitis,

autoimmune encephalitis, and mitochondrial disease mimicker: expanding the *DNM1L*-

associated intractable epilepsy and encephalopathy phenotype. *Epileptic Disord* **21**:112–

116. doi:10.1684/EPD.2019.1036

Osellame LD, Singh AP, Stroud DA, Palmer CS, Stojanovski D, Ramachandran R, Ryan MT.

2016. Cooperative and independent roles of the Drp1 adaptors Mff, MiD49 and MiD51 in

mitochondrial fission. *J Cell Sci* **129**:2170–2181. doi:10.1242/JCS.185165/-/DC1

Otera H, Miyata N, Kuge O, Mihara K. 2016. Drp1-dependent mitochondrial fission via MiD49/51

is essential for apoptotic cristae remodeling. *J Cell Biol* **212**:531–544.

doi:10.1083/jcb.201508099

Palmer CS, Elgass KD, Parton RG, Osellame LD, Stojanovski D, Ryan MT. 2013. Adaptor

proteins MiD49 and MiD51 can act independently of Mff and Fis1 in Drp1 recruitment and

are specific for mitochondrial fission. *J Biol Chem* **288**:27584–27593.

doi:10.1074/JBC.M113.479873

Prigione A, Adjaye J. 2011. Modulation of mitochondrial biogenesis and bioenergetic metabolism

upon in vitro and in vivo differentiation of human ES and iPS cells. *Int J Dev Biol*.

doi:10.1387/ijdb.103198ap

Rasmussen ML, Taneja N, Neininger AC, Wang L, Robertson GL, Riffle SN, Shi L, Knollmann

BC, Burnette DT, Gama V. 2020. MCL-1 Inhibition by Selective BH3 Mimetics Disrupts

Mitochondrial Dynamics Causing Loss of Viability and Functionality of Human
Cardiomyocytes. *iScience* **23**. doi:10.1016/j.isci.2020.101015

Ravi A, Palamiuc L, Loughran RM, Triscott J, Arora GK, Kumar A, Tieu V, Pauli C, Reist M, Lew
RJ, Houlihan SL, Fellmann C, Metallo C, Rubin MA, Emerling BM. 2021. PI5P4Ks drive
metabolic homeostasis through peroxisome-mitochondria interplay. *Dev Cell* **56**:1661-
1676.e10. doi:10.1016/J.DEVCEL.2021.04.019

Ryan CS, Fine AL, Cohen AL, Schiltz BM, Renaud DL, Wirrell EC, Patterson MC, Boczek NJ, Liu
R, Babovic-Vuksanovic D, Chan DC, Payne ET. 2018. De Novo DNM1L Variant in a
Teenager With Progressive Paroxysmal Dystonia and Lethal Super-refractory Myoclonic
Status Epilepticus. *J Child Neurol* **33**:651–658. doi:10.1177/0883073818778203

Schrader M, Costello JL, Godinho LF, Azadi AS, Islinger M. 2016. Proliferation and fission of
peroxisomes - An update. *Biochim Biophys Acta* **1863**:971–983.
doi:10.1016/J.BBAMCR.2015.09.024

Schrader M, Kamoshita M, Islinger M. 2020. Organelle interplay-peroxisome interactions in health
and disease. *J Inherit Metab Dis* **43**:71–89. doi:10.1002/JIMD.12083

Sheffer R, Douiev L, Edvardson S, Shaag A, Tamimi K, Soiferman D, Meiner V, Saada A. 2016.
Postnatal microcephaly and pain insensitivity due to a de novo heterozygous DNM1L
mutation causing impaired mitochondrial fission and function. *Am J Med Genet Part A*
170:1603–1607. doi:10.1002/AJMG.A.37624

Smaili SS, Hsu YT, Sanders KM, Russell JT, Youle RJ. 2001. Bax translocation to mitochondria
subsequent to a rapid loss of mitochondrial membrane potential. *Cell Death Differ* **8**:909–
920. doi:10.1038/SJ.CDD.4400889

Smirnova E, Griparic L, Shurland DL, van der Bliek AM. 2001. Dynamin-related protein Drp1 is

required for mitochondrial division in mammalian cells. *Mol Biol Cell* **12**:2245–56.

doi:10.1091/mbc.12.8.2245

Spinelli JB, Rosen PC, Sprenger H-G, Puszynska AM, Mann JL, Roessler JM, Cangelosi AL,

Henne A, Condon KJ, Zhang T, Kunchok T, Lewis CA, Chandel NS, Sabatini DM. 2021.

Fumarate is a terminal electron acceptor in the mammalian electron transport chain. *Science*

374:1227–1237. doi:10.1126/SCIENCE.ABI7495

Sugiura A, Mattie S, Prudent J, Mcbride HM. 2017. Newly born peroxisomes are a hybrid of

mitochondrial and ER-derived pre-peroxisomes. *Nature* **542**:251–254.

doi:10.1038/NATURE21375

Taguchi N, Ishihara N, Jofuku A, Oka T, Mihara K. 2007. Mitotic phosphorylation of dynamin-

related GTPase Drp1 participates in mitochondrial fission. *J Biol Chem* **282**:11521–11529.

doi:10.1074/jbc.M607279200

Tanaka H, Okazaki T, Aoyama S, Yokota M, Koike M, Okada Y, Fujiki Y, Gotoh Y. 2019.

Peroxisomes control mitochondrial dynamics and the mitochondrion-dependent apoptosis

pathway. *J Cell Sci* **132**. doi:10.1242/JCS.224766

Valera-Alberni M, Joffraud M, Miro-Blanch J, Capellades J, Junza A, Dayon L, Núñez Galindo A,

Sanchez-Garcia JL, Valsesia A, Cercillieux A, Söllner F, Ladurner AG, Yanes O, Cantó C.

2021. Crosstalk between Drp1 phosphorylation sites during mitochondrial remodeling and

their impact on metabolic adaptation. *Cell Rep* **36**.

doi:10.1016/J.CELREP.2021.109565/ATTACHMENT/3EF18C2E-C1FE-434A-A87F-

E1E1BD407021/MMC1.PDF

Vandeleur D, Chen C V., Huang EJ, Connolly AJ, Sanchez H, Moon-Grady AJ. 2019. Novel and

lethal case of cardiac involvement in DNM1L mitochondrial encephalopathy. *Am J Med*

Genet Part A **179**:2486–2489. doi:10.1002/AJMG.A.61371

Vantaggiato C, Castelli M, Giovarelli M, Orso G, Bassi MT, Clementi E, De Palma C. 2019. The

Fine Tuning of Drp1-Dependent Mitochondrial Remodeling and Autophagy Controls

Neuronal Differentiation. *Front Cell Neurosci* **13**:120. doi:10.3389/fncel.2019.00120

Vincent AE, White K, Davey T, Philips J, Ogden RT, Lawess C, Warren C, Hall MG, Ng YS,

Falkous G, Holden T, Deehan D, Taylor RW, Turnbull DM, Picard M. 2019. Quantitative 3D

Mapping of the Human Skeletal Muscle Mitochondrial Network. *Cell Rep* **26**:996-1009.e4.

doi:10.1016/J.CELREP.2019.01.010/ATTACHMENT/FB7711C2-65A0-4155-8EB8-

6FD4440E627D/MMC1.PDF

Wai T, Langer T. 2016. Mitochondrial Dynamics and Metabolic Regulation. *Trends Endocrinol*

Metab. doi:10.1016/j.tem.2015.12.001

Wakabayashi J, Zhang Z, Wakabayashi N, Tamura Y, Fukaya M, Kensler TW, Iijima M, Sesaki

H. 2009. The dynamin-related GTPase Drp1 is required for embryonic and brain

development in mice. *J Cell Biol* **186**:805–816. doi:10.1083/jcb.200903065

Wasiak S, Zunino R, McBride HM. 2007. Bax/Bak promote sumoylation of DRP1 and its stable

association with mitochondria during apoptotic cell death. *J Cell Biol* **177**:439–450.

doi:10.1083/JCB.200610042

Wasilewski M, Chacinska A. 2021. MIC60 relays oxidation to block mitophagy. *Nat Metab 2021*

39 **3**:1146–1147. doi:10.1038/s42255-021-00460-1

Waterham HR, Ferdinandusse S, Wanders RJA. 2016. Human disorders of peroxisome

metabolism and biogenesis. *Biochim Biophys Acta - Mol Cell Res* **1863**:922–933.

doi:10.1016/J.BBAMCR.2015.11.015

Waterham HR, Koster J, van Roermund CWT, Mooyer PAW, Wanders RJA, Leonard J V. 2007.

A lethal defect of mitochondrial and peroxisomal fission. *N Engl J Med* **356**:1736–1741.

doi:10.1056/NEJMOA064436

Whitley BN, Lam C, Cui H, Haude K, Bai R, Escobar L, Hamilton A, Brady L, Tarnopolsky MA,

Dengle L, Picker J, Lincoln S, Lackner LL, Glass IA, Hoppins S. 2018. Aberrant Drp1-

mediated mitochondrial division presents in humans with variable outcomes. *Hum Mol Genet*

27:3710–3719. doi:10.1093/hmg/ddy287

Wilkens V, Kohl W, Busch K. 2013. Restricted diffusion of OXPHOS complexes in dynamic

mitochondria delays their exchange between cristae and engenders a transitory mosaic

distribution. *J Cell Sci* **126**:103–116. doi:10.1242/JCS.108852

Control

G32A

R403C

0 seconds   45 seconds   90 seconds   135 seconds   180 seconds

Intensity Scale

Hyperfused peroxisome

Hyperfused mitochondria

EMPF1-associated DRP1 mutations

Stalled
fission events

Hyperpolarized mitochondrial
membrane potential

Altered
cristae structure

Decreased
coupling efficiency

Delayed
apoptotic response