Peter D. Fowler

Comparative Medicine & Integrative Biology

Michigan State University

784 Wilson Rd., Room G100

East Lansing, Michigan 48824

fowlerpe@msu.edu

Northward Expansion of *Amblyomma americanum* (Acari: Ixodidae) into Southern Michigan

Peter D. Fowler[1,6] , S. Nguyentran[2], L. Quatroche[2], M.L. Porter[1], V. Kobbekaduwa[1], S. Tippin[3], Guy

Miller[3], E. Dinh[4], E. Foster[4,7], and J.I. Tsao[2,5]

[1]Comparative Medicine & Integrative Biology, Michigan State University, East Lansing, MI 48824

USA, [2]Department of Fisheries & Wildlife, Michigan State University, East Lansing, MI 48824 USA,

[3]Berrien County Health Department, Benton Harbor, MI 49023 USA, [4]Michigan Department of Health

and Human Services, Lansing, MI 48933 USA, [5]Department of Large Animal Clinical Sciences,

Michigan State University, East Lansing, MI 48824, USA.

[6]Corresponding author: fowlerpe@msu.edu

[7]Current address: Division of Vector-Borne Diseases, National Center for Emerging and Zoonotic

Infectious Diseases, Centers for Disease Control and Prevention, Fort Collins, CO 80521, USA.

24   **Abstract**

Amblyomma americanum (Linnaeus) (Acari: Ixodidae) (lone star tick) is an aggressive, generalist

parasite that vectors numerous important human and animal pathogens. In recent decades its geographic

range has expanded northwards from endemic regions in the southeastern and southcentral US. In 2019

five questing *A. americanum* comprising two life stages were detected at one site in Berrien County, in

southwestern Michigan, satisfying one CDC criterium for an established population for the first time in

the state. To better characterize the northern extent of emerging *A. americanum*, we conducted active

surveillance (i.e., drag sampling) in summer 2020 throughout Michigan's southern counties and detected

one adult *A. americanum* from each of six widespread sites, including where they had been detected in

2019. A larger established population was identified at another site in Berrien County, which yielded

691 *A. americanum* comprising three life stages. Questing tick phenologies at this site were similar to

that reported for other regions. Statewide surveillance in 2021 revealed no *A. americanum* outside of

Berrien County, but establishment criteria were met again at the two sites where established populations

were first detected respectively in 2019 and 2020. These observations may represent the initial

successful invasion of *A. americanum* into Michigan. Data from passive (1999-2020) and active

surveillance (2004-2021) efforts, including a domestic animal sentinel program (2015-2018), are

reported to provide context for this nascent invasion. Continued active surveillance is needed to help

inform the public, medical professionals, and public health officials of the health risks associated with

this vector.

Introduction

Amblyomma americanum (Linnaeus) (Acari: Ixodidae), also known as the lone star tick or turkey tick, is

a generalist parasite known for its aggressive behavior (Stafford et al. 2018, Molaei et al. 2019). Aside

from being a nuisance, it can bear negative health consequences. Heavy infestations of *A. americanum*

on wildlife can cause morbidity and mortality (Bolte et al. 1970, Stafford et al. 2018). *Amblyomma*

americanum is also known to vector a number of pathogens resposnsible for disease in both humans and

domestic animals (Jaworski et al. 2017, Madison-Antenucci et al. 2020, Kennedy and Marshall 2021)

and bites from this tick have also been linked with the recently discovered alpha-gal syndrome (AS),

otherwise known as the tick-associated red meat allergy (Cabezas-Cruz et al. 2019, Crispell et al. 2019,

de la Fuente et al. 2020, Sharma and Karim 2021), which, if serious, can result in anaphylaxis (Pattanaik

et al. 2018).

Since the mid-1900s, the geographic range of *A. americanum* has been expanding northwards from

endemic areas in the south central and southeastern US (Paddock and Yabsley 2007, Springer et al.

2014, Sonenshine 2018, Molaei et al. 2019). Management practices beginning in the early 20[th] century

to recover and increase forests and wildlife resulted in the growth and spread of its main host, the white-

tailed deer (*Odocoileus virginianus*), and possibly the wild turkey (*Meleagris gallopavo*) (Paddock and

Yabsley 2007). This increase in woodland habitats created more habitat not only for host species but

also for *A. americanum* itself by providing leaf litter suitable to protect the tick from desiccation

(Paddock and Yabsley 2007, Tsao et al. 2021). Additionally, the trend of warmer and shorter winters

beginning in late 20[th] century associated with anthropogenic climate change may be providing additional

opportunities for this tick to become established in new areas farther north (Sonenshine 2018). Models

suggest that the northward invasion of this tick is likely to continue in the coming decades (Springer et

al. 2015, Ludwig et al. 2016, Raghavan et al. 2019, Sagurova et al. 2019). Detecting, further

characterizing, and refining predictions about the northward expansion can help prepare the public and

medical professionals for the emergence of diseases associated with this vector.

Recently, breeding populations of *A. americanum* have been documented in southern New England

(Stafford et al. 2018, Telford et al. 2019) and in states surrounding the Great Lakes, established

populations of have been spreading northwards in Illinois (Gilliam et al. 2020, Lyons et al. 2021),

Indiana (Wojan et al. 2021), and Ohio (Fitak et al. 2014). Although there have been increases in *A.*

americanum detected by the public in Wisconsin (Christenson et al. 2017) and southern Ontario (Nelder

et al. 2019), established populations have not yet been declared in those areas (at the time of manuscript

submission). As of July 2019, the Minnesota Department of Public Health has not reported any

established populations of *A. americanum* although they have been reported by residents since 1998

(https://www.health.state.mn.us/diseases/tickborne/longstarstickreportsmap.pdf, accessed 10/6/21), and

established populations have been identified in nearby counties to the west in South Dakota (Black et al.

2021) and to the south in Iowa (Springer et al. 2014).

Although *A. americanum* in Michigan have been documented by passive surveillance since the mid-

1980s (Walker et al. 1998, MDHHS 2021a) there is a lack of evidence supporting the existence of any

established breeding populations (Walker et al. 1998). Here we present such data for one county in

Michigan. To better understand the context of this invasion, we also summarize detections of *A.*

americanum reported in historical passive and active surveillance efforts dating back to 2004 designed

to detect *Ixodes scapularis* (Say) because there is a high degree of overlap between *A. americanum* and

I. scapularis in habitat use and temporal activity (Lindsay et al. 1998, Kollars et al. 2000, Guerra et al.

2002, Allan 2009, Centers for Disease Control and Prevention 2020). We further include data from

active surveillance also focused on detecting *I. scapularis* carried out by the Berrien County Health

Department (BCHD) from 2019-2021. Finally, we present data from a broad statewide surveillance

study carried out in 2010 in order to define the range of the then emerging *I. scapularis*.

Methods

Passive surveillance

Records of resident-submitted ticks to the Michigan Department of Agriculture began in 1999. In 2015

the Michigan Department of Health and Human Services (MDHHS) took over the program, which

continues to the present. Data were retrieved from the public database on the MDHHS Michigan

Environmental Public Health Tracking website (MDHHS 2021a). Ticks are reported by the submitter's

most likely county of exposure or county of residence when the county of exposure is uncertain.

Submitted ticks were identified to species using dichotomous morphological keys (Brinton et al. 1965,

Sonenshine 1979, Keirans and Litwak 1989, Durden and Keirans 1996, Faulkner and Reinhard 2014,

Dubie et al. 2017, Egizi et al. 2019). Although the program was in operation in 2014, data were not

available for this analysis.

Active surveillance by drag sampling

Site selection. From 2004-2016, much of the sampling by Michigan State University (MSU) was

conducted to track the invasion of *I. scapularis* mainly in the western Lower Peninsula (Hamer et al.

2010), but this was expanded to southeastern Michigan beginning in 2015. In 2010 in order to better

define the range of *I. scapularis* surveillance was broadened to cover 21 counties across Michigan's

lower penninsula. Beginning in 2017, surveillance became more widespread across the state in order to

sample counties that previously had been sampled less frequently or not at all. In 2019 BCHD began

conducting surveillance for questing *I. scapularis*. Although the specific monitoring objectives varied

from year to year, site selection and sampling protocols were similar between MSU and BCHD. Public

land (e.g., city, county and state parks, state forests, state wildlife game areas, and national forests) with

suitable habitat for *I. scapularis* were selected. Much like *I. scapularis, A. americanum* is frequently

found in deciduous forests containing leaf litter that helps protect ticks from desiccation (Hair and

Howell 1970, Childs and Paddock 2003, Allan 2009, Kensinger and Allan 2011). In 2020, the number of

sites per county sampled by MSU was increased relative to that in prior years in order to increase the

likelihood of detecting *A. americanum* in southern Michigan: fifty-five primary sites were selected after

consulting local public health departments, local parks departments and other regional authorities to

identify sites with suitable habitat and public access. Sites with insufficient area to sample (less than

1000 meters) or unsuitable habitat (little to no deciduous forest, or lack of a managed trail) upon first

visits were not repeated.

Surveillance methods and sampling frequency. Active surveillance by drag sampling (Falco and Fish

1992) aimed at detecting the spread of *I. scapularis* has been ongoing in Michigan since 2002 (Foster

2004). Based on CDC guidelines for the active surveillance of questing prostriate (Eisen et al. 2019) and

metastriate ticks (Centers for Disease Control and Prevention 2020), a minimum of 1,000 meters were

sampled per visit per site by dragging a 1-m^2 white corduroy cloth along trail edges in contact with

understory and leaf litter where ticks were likely questing. In larger parks, two transects with a minimum

of 800 meters each were sampled in different areas of the park. Sampling was performed on rain-free

days throughout the day unless relative humidity dropped below 40%. Collected ticks were placed in 70-

90% ethanol and brought back to the lab for species identification using dichotomous morphological

keys (Brinton et al. 1965, Sonenshine 1979, Keirans and Litwak 1989, Durden and Keirans 1996,

Faulkner and Reinhard 2014, Dubie et al. 2017, Egizi et al. 2019).

In 2004-2019 and 2021, because the main objective of our surveillance was to document the distribution

of *I. scapularis* and Lyme disease risk, the bulk of the sampling effort generally occurred from May –

July to target the nymphal stage of this tick. This time frame also encompasses part of adult and much of

the nymphal host-seeking period of *A. americanum*, according to reported phenologies in other states

(Sonenshine 1979, Schulze et al. 1986, Kollars et al. 2000, Telford et al. 2019). Sites were sampled at

least once during these years. In 2020, COVID-19 restrictions delayed sampling efforts by MSU, but

sites with appropriate habitat and area were sampled a minimum of three times from June 8th - July

23rd, 2020. Sites where *A. americanum* was detected during the first three visits were sampled an

additional time (i.e., four times total) to attempt to detect multiple life stages or multiple specimens.

Figures were generated in ArcGIS Pro v. 2.8.3 and compiled in Adobe Illustrator v. 24.1.2.

Active surveillance by sampling companion canines

From 2015-2018 veterinary clinics and/or shelters throughout Michigan were recruited to survey

companion canines for ticks during routine health checks. Veterinary workers were asked to inspect

thoroughly a minimum of 60 patients (3 per day) over a period of at least 30 days when adult *I.*

scapularis ticks would be active in Michigan, targeting April – June. *Amblyomma americanum*

frequently parasitizes companion canines (Sonenshine 1979, Saleh et al. 2019) and this timeframe

overlaps the activity period of the adult *A. americanum* as well as part of that of the nymph. Only

canines who had not traveled more than 30 miles from their residence during the 10 days prior to their

health visit were included in the study. Clinics were instructed to report data on all canines inspected,

including those who did not present with ticks. All ticks were stored in 70% ethanol and submitted to

Michigan State University for identification as described above. The activities herein are considered

standard veterinary care and involve samples collected using non-invasive techniques; an approved

exemption (4/20/15) from filing an animal use form is on file with the MSU Institutional Animal Care

and Use Committee.

Characterization of the phenology of A. americanum

Because both adults and nymphs were detected at one site, Grand Mere State Park (GMSP), early in the

surveillance period in 2020, weekly sampling beyond the surveillance window was performed there to

detect larvae as well as to describe the host-seeking phenology of *A. americanum*. Sampling in the fall

ceased after two consecutive visits failed to yield any *A. americanum*. Host-seeking phenologies for

each tick life stage was characterized by determining the density of ticks per site visit in ticks per m^2 and

displaying the percentage of the total tick density for the season for each respective visit.

Results

Passive Surveillance

Figure 1 shows the numbers of ticks submitted by county to the Michigan Department of Agriculture

from 1999-2014 and MDHHS from 2015-2020 excluding 2014 for which data were not available.

Figure 2 shows the total number of ticks submitted from 2009 to 2020; *A. americanum* submissions

accounted for 1.8% (0.6 – 3.0% range) of all ticks submitted each year. Submissions were variable in

space and time with no apparent trends especially given the spatial distribution of Michigan residents.

Active Surveillance

Drag sampling (2004-2021)

Between 2004 and 2016, 628,140 m^2 were sampled from 35 counties located in the Lower Peninsula

(Allegan, Barry, Bay, Benzie, Berrien, Calhoun, Cass, Charlesvoix, Clinton, Ingham, Ionia, Iosco, Iron,

Isabella, Jackson, Kalamazoo, Kent, Lapeer, Leelanau, Livingston, Manistee, Mason, Montcalm,

Muskegon, Oakland, Oceana, Osceola, Ottawa, Roscommon, Saginaw, Sanilac, St. Clair, Tuscola, Van

Buren, Wayne) (Hamer et al. 2010 (2004-2008); Tsao, unpublished data (2009-2016)).   Four *A.*

americanum were recovered, all from southwestern Michigan: one adult female in Barry County (2004),

one adult female from Allegan County (2005), one adult female in Van Buren County (2005), and one

nymph in Barry County (2007) (Table 1).

Figure 3 shows the distribution of sampling effort across Michigan from 2017-2021 including the efforts

by BCHD (2019-2021, Berrien County). Ingham County has a site marked by a yellow star which was

sampled at regular intervals throughout the season in order to monitor regional activity of the target

species, normally *I. scapularis;* only the first three visits to this site are included in the effort

calculations as these efforts do not represent surveillance across the county and would skew effort

reported in the figure (Figure 3). From a total of 212,989 m^2 sampled in 2017, only one adult male *A.*

americanum was recovered (Barry County). From 467,688 m^2 sampled in 2018, no *A. americanum* was

detected. From 511,330 m^2 sampled in 2019, five *A. americanum* of two life stages (1 adult male and 4

nymphs) were recovered at a single site in Berrien County, fulfilling one of the CDC criteria for an

established population (Centers for Disease Control and Prevention 2020). BCHD sampled 12,000 m^2,

that same year (2019) but no other *A. americanum* were detected. In 2020, a total of 265,745 m^2 were

sampled at 55 sites across 24 counties in southern Michigan by MSU between June 8[th] and July 23[rd]. A

total of 2,918 ticks were collected (Table 2) comprised of *I. scapularis* (90.3%), *Dermacentor variabilis*

(Say) (7.8%), *A. americanum* (1.2%), *I. dentatus* (Marx) (0.01%), and *Haemaphysalis leporispalustris*

(Packard) (0.1%). Of the 55 sites sampled, *A. americanum* was found at 7 sites in five counties across

southern Michigan (from west to east: Berrien, Branch, Cass, Ingham, and Huron). Only one site,

located in Berrien County, yielded more than one *A. americanum* and multiple life stages. An additional

25,000 m^2 were sampled from May 11th to July 30th in 2020 by the Berrien County Health Department,

which revealed a single *A. americanum* from two sites. In 2021, statewide surveillance by MSU for *I.*

scapularis resumed; 373,378 m^2 were sampled and an additional 38,000 m^2 were sampled by BCHD.

Amblyomma americanum was only detected in Berrien County, where multiple ticks were detected at

two sites, Grand Mere State Park (GMSP) and Warren Dunes State Park (WDSP).

Berrien County Active Surveillance

Table 3 summarizes the total sampling effort in Berrien County and numbers of *A. americanum* detected

from 2019 – 2021 at all the sites that were sampled. Four parks were sampled all three years; two other

parks were sampled only once. From the four parks sampled each year, *A. americanum* was detected at

three sites (Table 3). At one site, Warren Dunes State Park (WDSP), the status of *A. americanum* was

established in 2019 and was confirmed in 2021. Surprisingly only one *A. americanum* was found at this

site in 2020 despite similar effort. At a second site, Grand Mere State Park (GMSP), *A. americanum* was

not detected in 2019, but then multiple stages, including larvae, were detected in both 2020 and 2021,

suggesting the presence of an established and breeding population. At the third site, Love Creek Nature

Center, *A. americanum* were detected in only one year, in which two adults were detected thus not

reaching the threshold criteria for an established population. Total sampling effort per site for all three

years ranged from 26,003 m^2 to 57,165 m^2, and the total distance sampled at WDSP and GMSP was

~1.5-2 times more than the other two sites, where no *A. americanum* were detected or reported.

 Sampling companion canines (2015-2018)

Study periods varied over the four years, but all included the periods when adult and nymphal *A.*

americanum would be expected to be active (Table 4). At least 300 animals representing 16 – 69

counties per year were examined for a total of 4,097 animals examined. No *A. americanum* were

detected in three of the four study years. In 2016, when 2,510 individuals were examined, a single adult

female *A. americanum* was detected from a dog residing in Muskegon County.

Characterization of the phenology of A. americanum *at GMSP*

A total of 448 *A. americanum* were sampled at GMSP between June 11[th] and October 12[th] 2020,

including 409 larvae, 32 nymphs, 3 adult males, and 2 adult females. Seasonal activity for *A.*

americanum and the now endemic *I. scapularis* (for comparison) are presented in Figure 4. *Amblyomma*

americanum adults were detected only in June. Nymphs were active from June to September but were

most active in June. Larvae were first detected in August but peaked in September and were not detected

in October. *Ixodes scapularis* adults were detected in June, late September, and October, with greatest

numbers in October. Nymphal *I. scapularis* were detected from June to September and peaked in June.

Larval *I. scapularis* were detected throughout the study period, with a first peak in July followed by a

second peak in September.

Discussion

Since the mid-1980s, detections of *A. americanum* in Michigan have been recorded in passive surveys

(Walker et al. 1998, MDHHS 2021a); however, until recently, active surveillance efforts only rarely

detected *A. americanum*. Applying the CDC's operational definition for established tick populations

(Springer et al. 2014), we report the first records of established populations of *A. americanum* in

Michigan, and according to CDC convention, the lone star tick status of Berrien County should be

considered established since 2019. To provide evidence that this invasion is recent and geographically

limited we collated the results of prior surveillance efforts in Michigan comprising passive surveillance

for ticks in general and active surveillance for blacklegged ticks specifically. Weekly surveillance at one

site where a breeding population of *A. americanum* was detected allowed us to characterize the seasonal

activity of this newly establishing tick in Michigan, which appears to be similar to that in other areas

where lone star ticks are established.

Previous active surveillance in Michigan (2004-2019)

Between 2004 and 2018, 841,129 m^2 were sampled throughout the state and only four adult and one

nymphal *A. americanum* were detected despite sampling in suitable habitats during phenologically

appropriate times of the year (Hamer et al. 2010; Tsao, unpublished data). While these efforts may have

missed some cryptic populations, especially in less sampled regions (e.g, the Upper Peninsula), the

findings suggest that there were no widespread established populations of *A. americanum* within

Michigan during this timeframe. Furthermore, examination of > 4,000 companion canines over four

years (2015-2018) across Michigan resulted in the detection of one *A. americanum* tick attached to one

dog, who resided in a county in the Lower Peninsula along Lake Michigan.

In 2019, drag sampling revealed five *A. americanum* of two life stages (one adult male and four

nymphs) all from Warren Dunes State Park (WDSP). We took immediate notice of this finding because

detecting five ticks at one site in one summer starkly contrasts with detecting a total of five ticks from

three counties over fifteen years. We also took notice because prior detections mainly were comprised

of single adult ticks, whereas at WDSP in 2019, 4 of 5 ticks were nymphs. For nymphs to be detected in

an area, they may have derived from locally hatched larvae that fed on local hosts, thereby representing

a locally reproducing population. On the other hand, they may have derived from larvae that had

hatched elsewhere and then were introduced into the area by dispersing hosts. For example, animals,

such as deer, coyote, or birds, dispersing from northwest Indiana or other areas in southwest Berrien

County with undetected lone star tick populations may have introduced larvae. In either of these cases,

introduction of larvae would be from within the same local region, suggesting the successful invasion

lone star ticks in Michigan. Considering that lone star tick larvae generally host-seek in late summer

(Telford et al. 2019), it is unlikely that migratory songbirds would have introduced larvae from the

endemic regions in southeastern or south central U.S. or even populations in southern Indiana or Illinois,

as fall migrating songbirds are traveling in the opposite direction. Although it is possible for nymphs to

be dispersed northwards by spring migrating songbirds, as spring migration overlaps with the early part

of the nymphal host-seeking period, multiple successful events as well as surviving the local abiotic

conditions would need to follow, for these engorged nymphs to result in local larval ticks (e.g., molting,

finding an adult host, finding a mate, ovipositing, and larval hatch). A high volume of migrating birds

landing and/or stopping over in an area may elevate the chances of such introductions. However,

introductions of dispersing larvae, nymphs or adults from nearby established populations might be more

likely.

Targeted surveillance for A. americanum *in Michigan in 2020 and statewide surveillance in 2021*

Given the finding of multiple life stages of *A. americanum* at WDSP in southwestern Michigan, and

given the presence of established *A. americanum* populations in northwestern Indiana counties that

border southwestern Michigan, (L. Green, Indiana Department of Public Health (IDOH) pers. comm.)

surveillance conducted by MSU in 2020 focused on southern counties. *Amblyomma americanum* were

detected at a greater number of sites and counties in 2020 than in any previous year, and this may have

been due in part to greater sampling effort per county compared with that in previous years. First,

although sampling effort per site was similar in 2020 compared to 2019 (approximately 3-4 visits/site), it

was generally greater compared to that in prior years. Second, in 2020 we attempted to increase the

concentration of sites from ~1-2 sites per county to 3-5 sites per county. Even with this increased

coverage of southern Michigan, only one county (Berrien) yielded more than one site with a *A.*

americanum tick detection, and only one site, Grand Mere State Park (GMSP) in Berrien County,

revealed more than one life stage. It is not surprising that sampling a greater density of sites may result

in a greater number of detections of *A. americanum*. Although the sample size is small (n = 5),

interestingly, the detections outside of Berrien County were all adults, which may represent adventitious

nymphs dispersed by from endemic areas in the south central and southeastern U.S. by spring migrating

songbirds (as discussed in Stafford et al. 2018). At one of the sites outside Berrien County where an

adult *A. americanum* was detected in 2020, ~ bi-weekly sampling May – October in 2019 and 2021 did

not result in any detections of *A. americanum*.

A. americanum tick emergence in Berrien County

In the early 2000s Berrien County was sampled frequently because it is also where established

populations of *I. scapularis* were first detected in the Lower Peninsula (Foster et al. 2004). Because of

the limited resources and the need to monitor other areas where the establishment status of *I. scapularis*

was less known, Berrien County was not sampled as intensively (i.e., number of sites and frequency of

visits per year) by MSU after the early 2000s. Beginning in 2019, sampling effort in Berrien County

increased in large part due to the initiation of surveillance by BCHD as part of a MDHHS-funded

program to increase surveillance efforts for invasive *I. scapularis* and pathogens in Michigan. MSU

efforts also increased in 2020 following the discovery of multiple life stages of *A. americanum* at WDSP

in 2019. Subsequently, multiple life stages of *A. americanum* were detected in two years for two sites,

WDSP and GMSP.

Although *A. americanum* were collected in each year (2019-2021) at WDSP, the tick collections only

met one of the CDC criteria for established populations (detecting at least two life stages in one calendar

year) in 2019 and 2021. The numbers of ticks collected each year were low, never achieving the other

criterium of detecting at least 6 ticks. In contrast, data from GMSP suggest that *A. americanum* either

were not yet established at GMSP in 2019 or were present at an abundance below our threshold of

detection, but then appear to be steadily increasing in 2020 and 2021. Taken together, established

populations, according to the CDC criteria and applied at the site level, have been detected in Berrien

County for three consecutive years.

No established populations of *A. americanum* were detected outside WDSP and GMSP, suggesting that

either further spread has not yet occurred, or that the level of sampling effort was inadequate to detect

low density emerging populations. Future research may investigate differences in abiotic and biotic

factors among sites to understand variation in initial establishment success of *A. americanum*. For

instance, both WDSP and GMSP are located along Lake Michigan, whereas other sites are found further

inland. Proximity to Indiana's established populations, from which Berrien County's *A. americanum*

likely originated, may have increased invasion success. Interestingly WDSP, where *A. americanum*

appears to be establishing at a slower rate compared to GMSP, is closer to the Indiana populations. In

fact, WDSP is ~48 km from the Indiana Dunes National Park and Indiana Dunes State Park, where *A.*

americanum are abundant (L. Green, IDOH, pers. Comm., see below).

Definition of established populations and importance of abiotic factors for establishment and invasion

The criteria for considering a tick species as being established in a given county is important to consider

as we seek to understand their spread. Established tick populations as defined by the CDC involves

either finding more than one life stage, or finding at least six ticks, in that county in a calendar year

(Dennis et al. 1998, Springer et al. 2014, Nelder et al. 2019). Passive surveillance data also have been

used to fulfill this criteria (Dennis et al. 1998, Springer et al. 2014). While community-submitted ticks

can be an important source of data for informing risk, they can be unreliable for determining whether a

tick population is established in a given area, if travel history is not taken into account and tick exposure

may have occurred elsewhere in a tick endemic area. This may be a particularly important point for

aggressive ticks such as *A. americanum* which can have multiple ticks feeding on a single host. A single

dog traveling to an endemic area can easily bring back enough ticks to consider a county established.

Amblyomma americanum submitted to the MDHHS for the last decade fulfill CDC criteria for some

Michigan counties in some years, but the numbers vary widely from year to year, and no county is

consistently represented (Figures 1-2). Additionally, active surveillance from those same years do not

provide evidence for any established populations in those areas. More stringent criteria have been

proposed by Nelder et al. 2019, specifically requiring that active surveillance is needed to confirm an

established population. They further suggest that a population be confirmed for three consecutive years

where at least six specimens from a given locale be collected, including at least one drag-sampled tick

and at least one nymph. While this may seem like excessive criteria, it may be warranted, especially

when establishment data are being used to train ecological niche models used to predict the northward

spread of this vector in the coming years (e.g, Springer et al. 2015, Raghavan et al. 2019).

After lone star ticks are introduced into an area, both biotic and abiotic factors will affect establishment and invasion success. Given the generalist nature of *A. americanum*, and because one of their main hosts – the white-tailed deer – is very common and abundant in Michigan, invading *A. americanum* are not likely to be host-limited and instead may be more limited by mate-finding success, which will be overcome by increasing propagule pressure and off-host survivorship success. Abiotic factors, therefore, likely play a particularly critical role in population establishment along the northern edge of its distribution. These include foliage composition, temperature, humidity, day length and seasonality (Ludwig et al. 2016). Cold winters are known to limit the survivorship of overwintering ticks (Semtner and Hair 1976, Kaizer et al. 2015, Linske et al. 2019, Bacon et al. 2021), so it might not be purely coincidence that both in New England and the northern central U.S., populations of *A. americanum* are emerging initially along coastal habitats, which have more moderate winters relative to areas further inland. It is unknown how *A. americanum* may have been introduced into northwest Indiana, whether via terrestrial hosts or migrating birds, but there are records of low numbers of *A. americanum* collected via a combination of passive and active surveillance from the three northwest counties bordering Lake Michigan since the late 1990s and evidence of established populations since 2005 (L. Green, Indiana Department of Health, pers. Comm.). The Indiana Department of Health (IDOH) has been drag sampling *A. americanum* consistently for the last decade at the Indiana Dunes National Park and Indiana Dunes State Park, where there are robust populations (L. Green, IDOH pers. Comm.). Besides the moderating effect of Lake Michigan on winter conditions, the oak upland coastal dune forests along Lake Michigan provide sheltered areas with highly suitable microhabitat (i.e., thick leaf litter) that further insulate ticks from cold and desiccation (especially in the "valleys" on the leeward sides of dunes). *Ixodes scapularis* in Michigan's Lower Peninsula likely invaded from northwestern Indiana and emerged faster northwards along Lake Michigan than it did farther inland (Hamer et al. 2010). Given *A.*

americanum's lower tolerance for cold winters, the invasion of this tick might follow a similar pattern.

Additionally, if introduced, *A. americanum* may establish and emerge on the eastern side of the state

along the Lake Erie and Lake Huron coastlines where the proximity to the lakes help create more

moderate conditions facilitating winter survival.

Models have recently predicted that climate change resulting in longer summers and milder winters may

increase the survival and therefore establishment probability of *A. americanum* farther north (Raghavan

et al. 2019, Sagurova et al. 2019); correspondingly, these models predict that much of southern

Michigan is already suitable for *A. americanum* tick establishment (Raghavan et al. 2019, Sagurova et

al. 2019). It should be noted, however, that some ecological niche models (Springer et al. 2015,

Raghavan et al. 2019) rely on passive surveillance data (Springer et al. 2014) that resulted in

categorizing some Michigan counties as having established *A. americanum* populations, which as

discussed previously, have not been confirmed through active surveillance (Walker et al. 1998).

Sagurova et al. (2019), however, uses a mechanistic model based on regional climate and tick

development rates, which does not rely on these data, and still predicts that *A. americanum* are likely to

become established in Michigan in the immediate future.

Seasonal activity of A. americanum *emerging in Michigan and implications for public health*

Frequent sampling at GMSP in 2020 allowed us insight into the phenologies of the three questing life

stages of *A. americanum* in southwestern Michigan (Figure 3). Unfortunately, due to COVID-19 related

restrictions, surveillance by MSU was unable to begin until 6/12/20 (although BCHD started

surveillance 5/17/20, they did not detected any *A. americanum* at this site). The phenological trends,

however, were consistent with that described in endemic areas further south and east (Sonenshine 1979,

Schulze et al. 1986, Kollars et al. 2000, Carroll 2011, Gilliam et al. 2018). The seasonal patterns

observed here are based on very low tick numbers. Assuming established *A. americanum* populations

will continue to increase in abundance, continued monitoring encompassing early spring through fall

will result in a greater confidence in observed patterns. Increased abundance also may result in broader

activity periods than currently observed. For instance, perhaps more nymphs will be observed later in

the summer, and some larvae may be detected in early summer as had been noted recently on Long

Island and Massachusetts (Telford et al. 2019).

Although there are differences in phenologies between *A. americanum* and *I. scapularis,* the activity

periods for all three life stages of *A. americanum* coincide with one of the major activity periods (i.e.,

larvae and adults) or overlap almost entirely (i.e., nymphs) with that of the respective life stages of *I.*

scapularis. Coincident phenologies have practical implications for public health. First, efforts to survey

for *I. scapularis*, which generally target the nymphal host-seeking period because of their preeminent

epidemiological role, should also detect nymphal *A. americanum*, and to a lesser extent adults, thereby

increasing the likelihood of detecting and characterizing emerging *A. americanum* in Michigan. Second

and similarly, public health messaging that typically is targeted in spring to prevent and mitigate Lyme

disease (e.g., promoting tick prevention and recognition of signs of a tick-borne disease) will also help

reduce risk and negative impacts of *A. americanum*-associated diseases. Third, it is important for the

public and for healthcare workers to recognize the different tick species and how they are associated

with different diseases to prevent misdiagnosis and treatment of tick-borne diseases (e.g., over-diagnosis

and treatment of Lyme disease and under-diagnosis and treatment of an *A. americanum*-associated

disease) (Egizi et al. 2017). For example, post-tick antibiotic prophylaxis of Lyme disease (Lantos et al.

2021) can be provided for patients who detect a nymphal or adult *I. scapularis* that has fed for at least 36

hours and which had been acquired from a high Lyme disease incidence area, such as Berrien County

(MDHHS 2021b). The coincident activity of *I. scapularis* and *A. americanum* nymphs, however, creates

an additional challenge to determine the species of the small feeding nymphs, which thus may affect the

decision to recommend post-tick antibiotic prophylaxis.

Additional considerations for public health in emerging areas

Amblyomma americanum is of special concern in emerging areas as many of its associated diseases may

not be familiar to medical professionals in the region. Pathogens of concern for human health that *A.*

americanum are known to vector have been summarized in a recent review by Madison-Antenucci et al.

2020 and include *Ehrlichia chaffeensis* the causative agent of human monocytic ehrlichiosis (HME), *E.*

ewingii, and the rare but highly fatal Heartland virus and Bourbon virus (Madison-Antenucci et al.

2020). For example, the numbers of *A. americanum* submitted to a passive surveillance program in

central New Jersey have been increasing over the last decade and have overtaken that of *I. scapularis* in

central New Jersey (Egizi et al. 2017, 2019). Yet relative to the estimated human encounter rates of *E.*

chaffeensis-infected *A. americanum* to *Borrelia burgdorferi*-infected *I. scapularis* , there are

significantly fewer expected reported cases of HME relative to Lyme disease, which may suggest an

under-reporting or mis-diagnosis of HME cases (Egizi et al. 2017). Similarly, an increase in mild cases

of spotted fever rickettsias (potentially caused by *Rickettsia amblyommatis*, which have been detected in

a high proportion of *A. Americanum*) should be expected, but should not be mistaken for the much more

severe Rocky Mountain spotted fever (Behravesh et al. 2016). One of the most concerning conditions

associated with *A. americanum* is the recently discovered alpha-gal syndrome (AS). In *A. americanum*

endemic areas, AS is the leading diagnosed cause of anaphylaxis (Pattanaik et al. 2018). One recent case

study (Houchens et al. 2021) published in the *New England Journal of Medicine*, titled "Hunting for a

Diagnosis," describes a Michigan resident (with no travel history outside of the state) who arrived at an

emergency room with severe anaphylaxis. After four days of hospitalization and multiple tests, the

patient was diagnosed with AS. The details in this report demonstrate the potential severity of AS and

the difficulty in diagnosing this unusual condition, especially in areas where *A. americanum* is newly

emerging and awareness of AS may not be prevalent in the medical community (Houchens et al. 2021).

Conclusion

Models have predicted that *A. americanum* would expand its range northward into Michigan (Molaei et

al. 2019, Sagurova et al. 2019), but until 2019, active surveillance had not detected any established

population. Here we report both an established *A. americanum* population at the northern extent of its

range as well as the seasonal activity. Negative surveillance data in preceding years suggest that these

findings most likely represent a nascent invasion rather than a long established but undetected

population. Phenological patterns observed also provide insights into the public health risks posed by

this tick over the course of the year and can help inform optimal time frames for future surveillance

efforts. Due to limited resources and time, collected ticks have not yet been submitted for pathogen

testing. Given similar host and habitat requirements, if the previous invasion of *I. scapularis* into

Michigan is any indication (Hamer et al. 2010, Lantos et al. 2017, MDHHS 2020), we can expect to see

A. americanum expanding its range in Michigan in the coming years. Educating public health officials

and the medical community for both humans and companion animals about the diseases associated with

this tick and how to differentiate them from other endemic tick-borne diseases is recommended.

Acknowledgements

We thank S. Hamer, G.J. Hickling, and E.D. Walker for their contributions to beginning the long-term
surveillance studies of invasive ticks in Michigan from which subsequent efforts have grown. Similarly,

we thank E.D. Walker for providing additional insight into the data from Walker et al. 1998 as well as
having coordinated and published that valuable work, which provides a baseline from which to compare
subsequent distributions of ticks over time. For assistance in the field and lab, we thank S. Altus, I.
Arsnoe, M. Bammer, A. Dunivant, K. Fake*, S. Froehlich, M. Gleason*, G. Grzesiak, S. Haupt*, D.
Houvener, S. Kim, J. Kryda*, A. Larson, A. Luchenbill, M. Orbain, G. Pang, J.Pastori, M. Regalado, B.
Reith, M. Rice, M. (Rosen) Clayson, J. Schroeder, M. Soja, N. Spala, J. Stych, A. Talbott, M. Volk, H.
Waters, K. Wickens, B. Wilson*, A. Yackley, G. Yarandi, S. Zohr, and many other assistants who
contributed to earlier, published work. We thank our agency partners who variously started and continue
to maintain the Michigan Department of Health and Human Services (MDHHS) community tick
submission program; developed, funded, and facilitated active invasive vector surveillance efforts
conducted by local health departments; and provided suggestions for MSU active surveillance efforts
based on public health risks. They include R. Eisen, M. Poplar, R. Reik, J. Sidge, K. Signs, and M.G.
Stobierski. We furthermore thank L. Green from the Indiana Department of Health (IDOH) for
providing historical knowledge and guidance on sampling for *A. americanum* in northern Indiana.
Permission to sample properties was graciously provided by the Michigan Department of Natural
Resource, National Park Service, many county parks offices, nature centers, land conservancies,
universities, and other public and private entities. We are grateful for many sources of funding
supporting the work presented here. First and foremost, the majority of the active surveillance efforts for
invasive ticks in the Tsao lab has been generously supported by the Michigan Lyme Disease Association
(MLDA), facilitated by the L. Lobes. Drag surveillance by the Tsao lab in 2019 was largely funded by
Contract E20193274-00 from MDHHS. Funding for the companion animal surveillance study was
provided by the College of Veterinary Medicine Endowed Companion Animal Fund RT082792, the
MLDA, and greatly facilitated by L. Penman and Boehringer Ingelheim Animal Health. Drag
surveillance by the Berrien County Health Department was funded in part by the MDHHS, and the
MDHHS community tick submission program is supported with funds from the Epidemiology and
Laboratory Capacity (ELC) for Prevention and Control of Emerging Infectious Diseases Cooperative
Agreement from the Centers for Disease Control and Prevention, award number CDC-RFA-CK19-1904.
We thank the Michigan State University (MSU) Comparative Medicine and Integrative Biology
program for fellowship funding for PF, MP, and VK. MP, VK, and JT also have been funded by the
CDC Midwest Center of Excellence in Vector-Borne Diseases U50723K866 under a subcontract to
Michigan State University from the University of Wisconsin (L. Bartholomay, S. Paskewitz, principal
investigators). This publication was supported by Cooperative Agreement #U01 CK000505, funded by
the Centers for Disease Control and Prevention including funding for SN, LQ, and several other research
fellows listed above with asterisks (*) who contributed to surveillance efforts. Its contents are solely the
responsibility of the authors and do not necessarily represent the official views of the Centers of Disease
Control and Prevention or the Department of Health and Human Services.
519

References

Allan, B. F. 2009. Influence of prescribed burns on the abundance of amblyomma americanum (Acari:
Ixodidae) in the missouri ozarks. J. Med. Entomol. 46: 1030–1036.

Bacon, E. A., H. Kopsco, P. Gronemeyer, N. Mateus-Pinilla, and R. L. Smith. **2021**. Effects of
Climate on the Variation in Abundance of Three Tick Species in Illinois. J. Med. Entomol. 1–10.
Behravesh, C. B., C. D. Paddock, F. S. Dahlgren, R. J. Eisen, and Y. P. Springer. **2016**. Expanding
Range of Amblyomma americanum and Simultaneous Changes in the Epidemiology of Spotted
Fever Group Rickettsiosis in the United States. Am. J. Trop. Med. Hyg. 94: 35–42.
Black, H., R. Potts, J. Fiechtner, J. E. Pietri, and H. B. Britten. **2021**. Establishment of Amblyomma
americanum populations and new records of Borrelia burgdorferi-infected Ixodes scapularis in
South Dakota. J. Vector Ecol. 46: 143–147.
Bolte, J., J. Hair, and J. Fletcher. **1970**. White-Tailed Deer Mortality Following Tissue Destruction
Induced by Lone Star Ticks Author (s): John R . Bolte , Jakie A . Hair and Joe Fletcher Published
by□: Wiley on behalf of the Wildlife Society Stable URL□: http://www.jstor.org/stable/3798861
Acce. J. Parasitol. 34: 546–552.
Brinton, E. ., D. . Beck, and D. . Allred. **1965**. Identification of the adults, nymphs and larvae of ticks
of the genus dermacentor koch (Ixodidae) in the western united states. Brigham Young Univ. 5: 1–
42.
Cabezas-Cruz, A., A. Hodžić, P. Román-Carrasco, L. Mateos-Hernández, G. G. Duscher, D. K.
Sinha, W. Hemmer, I. Swoboda, A. Estrada-Peña, and J. De La Fuente. **2019**. Environmental
and molecular drivers of the α-Gal syndrome. Front. Immunol. 10: 1–12.
Carroll, J. F. 2011. An Increasing Presence: The Lone Star Tick, Amblyomma americanum (Linnaeus)
(Acari: Ixodidae) in Maryland. 5: 66–76.
Centers for Disease Control and Prevention. **2020**. Guide to the Surveillance of Metastriate Ticks (
Acari□: Ixodidae) and their Pathogens in the United States. Div. Vector-Borne Dis. CDC. Atlanta
Ft. Collins.
Childs, J. E., and C. D. Paddock. **2003**. THE ASCENDANCY OF AMBLYOMMA AMERICANUM
AS A VECTOR OF PATHOGENS AFFECTING HUMANS IN THE UNITED STATES *. Annu.
Rev. Entomol. 48: 307–337.
Christenson, M., X. Lee, S. Larson, D. H. Johnson, J. Jensen, M. Meller, S. Paskewitz, and L. Foil.
2017. Occurrence of Amblyomma Americanum (Acari: Ixodidae) and human infection with
ehrlichia chaffeensis in Wisconsin, 2008-2015. J. Med. Entomol. 54: 752–756.
Crispell, G., S. P. Commins, S. A. Archer-Hartman, S. Choudhary, G. Dharmarajan, P. Azadi,
and S. Karim. **2019**. Discovery of alpha-gal-containing antigens in North American tick species
believed to induce red meat allergy. Front. Immunol. 10.
Dennis, D. T., T. S. Nekomoto, J. C. Victor, W. S. Paul, and J. Piesman. **1998**. Reported Distribution
of Ixodes scapularis and Ixodes pacificus (Acari: Ixodidae) in the United States. J. Med. Entomol.
35: 629–638.
Dubie, T. R., R. Grantham, L. Coburn, and B. H. Noden. **2017**. Pictorial Key for Identification of
Immature Stages of Common Ixodid Ticks Found in Pastures in Oklahoma. Southwest. Entomol.
42: 1–14.
Durden, L. A., and J. E. Keirans. **1996**. Nymphs of the genus Ixodes (Acari: Ixodidae) of the United
States: taxonomy, identification key, distribution, hosts, and medical/veterinary importance.,
Nymphs Genus Ixodes (Acari Ixodidae) United States Taxon. Identif. Key, Distrib. Hosts,
Medical/Veterinary Importance. Entomological Society of America, Lanham, Maryland.
Egizi, A., N. H. Fefferman, and R. A. Jordan. **2017**. Relative risk for ehrlichiosis and lyme disease in
an area where vectors for both are sympatric, New Jersey, Usa. Emerg. Infect. Dis. 23: 939–945.
Egizi, A. M., R. G. Robbins, L. Beati, S. Nava, C. R. Evans, J. L. Occi, and D. M. Fonseca. **2019**. A
pictorial key to differentiate the recently detected exotic Haemaphysalis longicornis Neumann,

1901 (Acari, Ixodidae) from native congeners in North America. Zookeys. 818: 117–128.

Eisen, R. J., L. Eisen, C. B. Graham, A. Hojgaard, P. S. Mead, G. Kersch, S. Karpathy, C. D.
Paddock, H. Savage, B. L. Herwaldt, and R. Bradbury. **2019**. Surveillance for Ixodes scapularis
and pathogens found in this tick species in the United States. Centers Dis. Control Prev. 1–34.

Falco, R. C., and D. Fish. **1992**. A comparison of methods for sampling the deer tick, Ixodes dammini,
in a Lyme disease endemic area. Exp. Appl. Acarol. 14: 165–173.

Faulkner, C. T., and K. J. Reinhard. **2014**. A retrospective examination of paleoparasitology and its
establishment in the journal of parasitology. J. Parasitol. 100: 253–259.

Fitak, R. R., D. J. Kelly, M. K. Daniels, J. Jiang, A. L. Richards, and P. A. Fuerst. **2014**. The
prevalence of rickettsial and ehrlichial organisms in Amblyomma americanum ticks collected from
Ohio and surrounding areas between 2000 and 2010. Ticks Tick. Borne. Dis. 5: 797–800.

Foster, E. 2004. Ixodes scapularis (Acari: Ixodidae) and Borrelia burgdorferi in southwest Michigan:
population ecology and verification of a geographic risk model.

Gilliam, B., P. Gronemeyer, S. Chakraborty, F. Winata, L. A. Lyons, C. Miller-Hunt, H. C. Tuten,
S. Debosik, D. Freeman, M. O'Hara-Ruiz, and N. Mateus-Pinilla. **2020**. Impact of Unexplored
Data Sources on the Historical Distribution of Three Vector Tick Species in Illinois. J. Med.
Entomol. 57: 872–883.

Gilliam, M. E., W. T. Rechkemmer, K. W. McCravy, and S. E. Jenkins. **2018**. The influence of
prescribed fire, habitat, and weather on Amblyomma americanum (Ixodida: Ixodidae) in West-
Central Illinois, USA. Insects. 9.

Guerra, M., E. Walker, C. Jones, S. Paskewitz, M. Roberto Cortinas, L. B. Ashley Stancil, M.
Bobo, and U. Kitron. **2002**. Predicting the risk of Lyme disease: Habitat suitability for Ixodes
scapularis in the north central United States. Emerg. Infect. Dis. 8: 289–297.

Hair, J. A., and D. E. Howell. **1970**. Lone star ticks: their biology and control in Ozark recreation
areas. Bulletin B□;679.

Hamer, S. A., J. I. Tsao, E. D. Walker, and G. J. Hickling. **2010**. Invasion of the lyme disease vector
ixodes scapularis: Implications for Borrelia burgdorferi endemicity. Ecohealth. 7: 47–63.

Houchens, N., S. Hartley, S. P. Commins, D. Claar, and S. Saint. **2021**. Hunting for a Diagnosis. N.
Engl. J. Med. 384: 462–467.

ISDH. **2021**. ISDH: Amblyomma americanum. 2021. (https://www.in.gov/isdh/28009.htm).

Jaworski, D. C., C. Cheng, A. D. S. Nair, and R. R. Ganta. **2017**. Amblyomma americanum ticks
infected with in vitro cultured wild-type and mutants of Ehrlichia chaffeensis are competent to
produce infection in naïve deer and dogs. Ticks Tick. Borne. Dis. 8: 60–64.

Kaizer, A. M., S. A. Foré, H. J. Kim, and E. C. York. **2015**. Modeling the biotic and abiotic factors
that describe the number of active off-host Amblyomma americanum larvae. J. Vector Ecol. 40: 1–
10.

Keirans, J. E., and T. R. Litwak. **1989**. Pictorial key to the adults of hard ticks, family Ixodidae
(Ixodida: Ixodoidea), east of the Mississippi River. J. Med. Entomol. 26: 435–448.

Kennedy, A. C., and E. Marshall. **2021**. Lone star ticks (Amblyomma americanum): An emerging
threat in delaware. Delaware J. Public Heal. 7.

Kensinger, B. J., and B. F. Allan. **2011**. Efficacy of dry ice-baited traps for sampling Amblyomma
americanum (Acari: Ixodidae) varies with life stage but not habitat. J. Med. Entomol. 48: 708–711.

Kollars, T. M., J. H. Oliver, L. A. Durden, and P. G. Kollars. **2000**. Host Associations and Seasonal
Activity of Amblyomma americanum (Acari: Ixodidae) in Missouri. J. Parasitol. 86: 1156.

de la Fuente, J., A. Cabezas-Cruz, and I. Pacheco. **2020**. Alpha-Gal Syndrome: challenges to
understanding sensitization and clinical reactions to alpha-gal. Expert Rev. Mol. Diagn. 00: 1–7.

Lantos, P. M., J. Rumbaugh, L. K. Bockenstedt, Y. T. Falck-Ytter, M. E. Aguero-Rosenfeld, P. G.
Auwaerter, K. Baldwin, R. R. Bannuru, K. K. Belani, W. R. Bowie, J. A. Branda, D. B.
Clifford, F. J. DiMario, J. J. Halperin, P. J. Krause, V. Lavergne, M. H. Liang, H. C.
Meissner, L. E. Nigrovic, J. (Jay) J. Nocton, M. C. Osani, A. A. Pruitt, J. Rips, L. E.
Rosenfeld, M. L. Savoy, S. K. Sood, A. C. Steere, F. Strle, R. Sundel, J. Tsao, E. E. Vaysbrot,
G. P. Wormser, and L. S. Zemel. **2021**. Clinical Practice Guidelines by the Infectious Diseases
Society of America (IDSA), American Academy of Neurology (AAN), and American College of
Rheumatology (ACR): 2020 Guidelines for the Prevention, Diagnosis and Treatment of Lyme
Disease. Clin. Infect. Dis. 72: e1–e48.
Lantos, P. M., J. Tsao, L. E. Nigrovic, P. G. Auwaerter, V. G. Fowler, F. Ruffin, E. Foster, and G.
Hickling. **2017**. Geographic expansion of Lyme disease in Michigan, 2000-2014. Open Forum
Infect. Dis. 4: 3–7.
Lindsay, L. R., I. K. Barker, G. A. Surgeoner, S. A. McEwen, T. J. Gillespie, and E. M. Addison.
1998. Survival and Development of the Different Life Stages of Ixodes scapularis (Acari: Ixodidae)
Held within Four Habitats on Long Point, Ontario, Canada. J. Med. Entomol. 35: 189–199.
Linske, M., S. Williams, K. Stafford, C. Lubelczyk, E. Henderson, M. Welch, and P. Teel. **2019**.
Determining Effects of Winter Weather Conditions on Adult Amblyomma americanum (Acari:
Ixodidae) Survival in Connecticut and Maine, USA. Insects. 11: 13.
Ludwig, A., H. S. Ginsberg, G. J. Hickling, and N. H. Ogden. **2016**. A dynamic population model to
investigate effects of climate and climate-independent factors on the lifecycle of Amblyomma
americanum (Acari: Ixodidae). J. Med. Entomol. 53: 99–115.
Lyons, L. A., M. E. Brand, P. Gronemeyer, N. Mateus-Pinilla, M. O. H. Ruiz, C. M. Stone, H. C.
Tuten, and R. L. Smith. **2021**. Comparing Contributions of Passive and Active Tick Collection
Methods to Determine Establishment of Ticks of Public Health Concern within Illinois. J. Med.
Entomol. 58: 1849–1864.
Madison-Antenucci, S., L. D. Kramer, L. L. Gebhardt, and E. Kauffman. **2020**. Emerging Tick-
Borne Diseases. Clin. Microbiol. Rev. 33.
MDHHS. **2020**. Michigan trends in tickborne Disease 2016-2020, Emerg. Zoonotic Infect. Dis.
MDHHS. **2021a**. MiTracking Michigan Environmental Public Health Tracking.
(https://mitracking.state.mi.us/?bookmark=112).
MDHHS. **2021b**. MDHHS Michigan Disease Mapper: Zoonotic and Vector Borne.
(http://www.mcgi.state.mi.us/midiseasemapper/mcgi.html).
Molaei, G., E. A. H. Little, S. C. Williams, and K. C. Stafford. **2019**. Bracing for the Worst — Range
Expansion of the Lone Star Tick in the Northeastern United States. N. Engl. J. Med. 381: 2189–
2192.
Nelder, M. P., C. B. Russell, K. M. Clow, S. Johnson, J. S. Weese, K. Cronin, F. Ralevski, C. M.
Jardine, and S. N. Patel. **2019**. Occurrence and distribution of Ambylomma americanum as
determined by passive surveillance in Ontario, Canada (1999–2016). Ticks Tick. Borne. Dis. 10:
146–155.
Paddock, C. D., and M. J. Yabsley. **2007**. Ecological Havoc, the Rise of White-Tailed Deer, and the
Emergence of Amblyomma americanum-Associated Zoonoses in the United States, pp. 289–324.
In .
Pattanaik, D., P. Lieberman, J. Lieberman, T. Pongdee, and A. T. Keene. **2018**. The changing face
of anaphylaxis in adults and adolescents. Ann. Allergy, Asthma Immunol. 121: 594–597.
Raghavan, R. K., A. Townsend Peterson, M. E. Cobos, R. Ganta, and D. Foley. **2019**. Current and
Future Distribution of the Lone Star Tick, Amblyomma americanum (L.) (Acari: Ixodidae) in

661   North America. PLoS One. 14: 1–13.

662 **Sagurova, I., A. Ludwig, N. H. Ogden, Y. Pelcat, G. Dueymes, and P. Gachon**. **2019**. Predicted
663   northward expansion of the geographic range of the tick vector amblyomma americanum in North
664   America under future climate conditions. Environ. Health Perspect. 127: 1–15.

665 **Saleh, M. N., K. D. Sundstrom, K. T. Duncan, M. M. Ientile, J. Jordy, P. Ghosh, and S. E. Little**.
666   **2019**. Show us your ticks: A survey of ticks infesting dogs and cats across the USA. Parasites and
667   Vectors. 12: 1–11.

668 **Schulze, T. L., G. S. Bowen, M. F. Lakat, W. E. Parkin, and J. K. Shisler**. **1986**. Seasonal abundance
669   and hosts of Ixodes dammini (Acari: Ixodidae) and other Ixodid ticks from an endemic lyme
670   disease focus in New Jersey, USA. J. Med. Entomol. 23: 105–109.

671 **Semtner, P. J., and J. A. Hair**. **1976**. The Ecology and Behavior of the Lone Star Tick (Acarina:
672   Ixodidae): X. Adult Overwintering and survival in woodlots1. J. Med. Entomol. 13: 216–217.

673 **Sharma, S. R., and S. Karim**. **2021**. Tick Saliva and the Alpha-Gal Syndrome: Finding a Needle in a
674   Haystack. Front. Cell. Infect. Microbiol. 11: 1–13.

675 **Sonenshine, D. E. 1979**. Insects of Virginia.13. Ticks of Virginia (Acari, Metastigmata), Virginia
676   Polytech. Inst. Res. Div. Bull.

677 **Sonenshine, D. E. 2018**. Range expansion of tick disease vectors in north america: Implications for
678   spread of tick-borne disease. Int. J. Environ. Res. Public Health.

679 **Springer, Y. P., L. Eisen, L. Beati, A. M. James, and R. J. Eisen**. **2014**. Spatial distribution of
680   counties in the continental United States with records of occurrence of amblyomma americanum
681   (Ixodida: Ixodidae). J. Med. Entomol. 51: 342–351.

682 **Springer, Y. P., C. S. Jarnevich, D. T. Barnett, A. J. Monaghan, and R. J. Eisen**. **2015**. Modeling
683   the present and future geographic distribution of the lone star tick, amblyomma americanum
684   (ixodida: Ixodidae), in the continental United States. Am. J. Trop. Med. Hyg. 93: 875–890.

685 **Stafford, K. C., G. Molaei, E. A. H. Little, C. D. Paddock, S. E. Karpathy, and A. M. Labonte**.
686   **2018**. Distribution and establishment of the lone star tick in Connecticut and implications for range
687   expansion and public health. J. Med. Entomol. 55: 1561–1568.

688 **Telford, S. R., J. Buchthal, and P. Elias**. **2019**. Early questing by lone star tick larvae, New York and
689   Massachusetts, USA, 2018. Emerg. Infect. Dis.

690 **Tsao, J. I., S. A. Hamer, S. Han, J. L. Sidge, and G. J. Hickling**. **2021**. The Contribution of Wildlife
691   Hosts to the Rise of Ticks and Tick-Borne Diseases in North America. J. Med. Entomol. 58: 1565–
692   1587.

693 **Walker, E. D., M. G. Stobierski, M. L. Poplar, T. W. Smith, A. J. Murphy, P. C. Smith, S. M.
694   Schmitt, T. M. Cooley, and C. M. Kramer**. **1998**. Geographic Distribution of Ticks (Acari:
695   Ixodidae) in Michigan, with Emphasis on Ixodes scapularis and Borrelia burgdorferi. J. Med.
696   Entomol. 35: 872–882.

697 **Wojan, C., T. Thrasher, E. Lacey, and K. Clay**. **2021**. Distribution, Dynamics, and Diversity of
698   Questing Ticks in the Lower Midwest. J. Med. Entomol. 1–10.

Table 1. Questing *Amblyomma americanum* detected by active surveillance (drag sampling) in Michigan from 2004 – 2021 by life stage and county. Sampling in 2004-2016 was focused on the western side of the state (with a few exceptions – 2010 was broadened to better identify the range of *I. scapularis*). Beginning in 2017 active surveillance was carried out statewide except in 2020 when surveillance efforts focused on southern counties. See text for details. Only years where *A. americanum* were detected are included.

| Year | County | No. of *Amblyomma americanum* detected | | | |
		Adult Male	Adult Female	Nymph	Larvae
2004	Barry	-	1	-	-
2005	Allegan	-	1	-	-
	Van Buren	-	1	-	-
2007	Barry	-	-	1	-
2017	Van Buren	1	-	-	-
2019	Berrien	1	-	4	-
	Muskegon	-	1	-	-
2020	Berrien	1	3	32	409
	Branch	1	-	-	-
	Huron	1	-	-	-
	Ingham	1	-	-	-
2021	Berrien	5	1	45	50
Total		11	8	82	459

Table 2. Total numbers of each tick species and life stage collected by drag sampling by county from June 8th – July 23rd, 2020 in southern Michigan. AM = Adult Male; AF = Adult Female; N = Nymph; L = Larva; *H. lep.* = *Haemaphysalis leporispalustris; I. dentatus = Ixodes dentatus; I. cookei = Ixodes cookei; D. variabilis = Dermacentor variabilis; Ixodes* spp. = Unidentifiable *Ixodes* species due to damage to specimen.

County	Effort (m²)	*Ixodes scapularis*				*Dermacentor variabilis*			*Amblyomma americanum*			Other
		AM	AF	N	L	AM	AF	N	AM	AF	N	
Allegan	5,381	1	7	8	-	7	8	-	-	-	-	
Barry	5,625	-	-	16	23	-	-	-	-	-	-	
Berrien	42,039	26	26	278	309	22	32	4	3	3	26	*I. dentatus* 1L, 3N
Branch	12,529	-	-	2	-	1	1	-	1	-	-	
Calhoun	10,855	5	3	52	1	3	2	-	-	-	-	*I. dentatus* 2N
Cass	25,161	-	3	128	835	1	8	-	-	1	-	*I. dentatus* 1N
Clinton	1,870	1	-	4	1	-	-	-	-	-	-	
Hillsdale	8,060	-	-	8	31	4	4	-	-	-	-	
Huron	10,277	1	4	3	-	-	-	-	1	-	-	*H. lep.* 1N
Ingham	4,700	8	7	250	82	-	-	-	1	-	-	
Jackson	16,513	3	-	36	9	1	3	-	-	-	-	
Kalamazoo	12,850	2	3	184	31	1	3	-	-	-	-	*I. dentatus* 3N
Lapeer	10,999	-	-	3	2	-	1	-	-	-	-	*I. dentatus* 1N
Lenawee	20,487	-	-	3	-	12	20	-	-	-	-	*I. cookei* 1AF
Monroe	11,336	-	-	-	-	10	7	-	-	-	-	*I. dentatus* 1N
Oakland	5,678	1	1	3	-	1	-	-	-	-	-	
Sanilac	1,560											
St. Clair	4,986	-	1	28	3	-	-	-	-	-	-	
St. Joseph	9,300	-	-	63	-	3	7	-	-	-	-	*I. dentatus* 1N
Tuscola	4,296	-	-	5	-	-	-	-	-	-	-	*Ixodes* spp. 1L
Van Buren	5,732	5	3	52	6	21	30	-	-	-	-	*H. lep.* 1N
Washtenaw	10,230	5	3	25	5	1	1	-	-	-	-	*I. dentatus* 1N
Wayne	26,806	1	-	21	6	3	5	-	-	-	-	*D. variabilis* 1L
Total	265,745	59	61	1172	1344	91	132	4	6	4	26	19

Table 3. Active surveillance effort for Berrien County from 2019-2021and status level for *Amblyomma americanum* detections (per site per year) as defined by the CDC as more than 6 ticks or more than one life stage found in a given calendar year (Springer et al. 2014). Total effort for each site is reported for each year and includes sampling performed by both Michigan State University and the Berrien County Health Department. Two sites were sampled only for one year. Total numbers of *A. americanum* are reported for each site for each year. (Note: the lone star status of Berrien County as a whole would be considered established as of 2019, given the findings at Warren Dunes State Park.)

Site Year	# of visits	Area dragged (m²)	Total *Amblyomma americanum* collected			Establishment status
			larvae	nymphs	adults	
Grand Mere State Park						
2019	6	6,600	0	0	0	
2020	24	36,365	409	32	5	Established
2021	12	14,200	50	42	5	Established
total		57,165				
Warren Dunes State Park						
2019	9	14,820	0	4	1	Established
2020	9	12,671	0	0	1	Reported
2021	11	12,858	0	4	1	Established
total		40,349				
Love Creek Nature Center						
2019	2	2,000	0	0	0	Not detected
2020	11	16,232	0	0	2	Reported
2021	10	12,000	0	0	0	Not detected
total		30,232				
Madeline Bertrand County Park						
2019	2	2,600	0	0	0	Not detected
2020	10	13,803	0	0	0	Not detected
2021	9	9,600	0	0	0	Not detected
total		26,003				
Galien River County Park						
2020	4	6667	0	0	0	Not detected
total		6,667				
Lincoln Township Beach						
2019	1	1000	0	0	0	Not detected
total		1,000				

734 **Table 4.** On-host *Amblyomma americanum* detected by active surveillance of companion canines with
735 no travel history in Michigan from 2015 – 2018 by participating veterinary clinics or shelters.
736

	Year			
	2015	2016	2017	2018
Dates animals were examined	5/1 – 6/5	2/10 – 6/3	4/3 – 7/26	5/13 - 7/27
No. clinics/shelters submitting data	30	48	26	8
No. of animals examined	391	2519	887	300
No. of counties represented	31	69	42	16
No. of animals infested with *A. americanum*	0	1[a]	0	0

737 [a]One adult female *A. americanum* was detected from an animal residing in Muskegon County.
738
739

Figure captions

Figure 1. Numbers of *Dermacentor variabilis, Ixodes scapularis, Amblyomma americanum,* and other species submitted annually to the Michigan Department of Agriculture (1999 – 2013) and Michigan Department of Health and Human Services (2015 – 2020). On average, 3.2% ±1.7% SD of all ticks submitted annually were *A. americanum*. Data source: MiTracking- Michigan Environmental Public Health Tracking; no data are available for 2014.

 Figure 2. *Amblyomma americanum* submitted annually to the Michigan Department Agriculture (2009-2013) and the Michigan Department of Health and Human Services (MDHHS) from 2015-2020. Data source: MiTracking- Michigan Environmental Public Health Tracking. Numbers in each county represent the number of community- submitted *A. americanum* by county of residence. No data were available for 2014. * Human population heatmap for each county is given for reference based on 2020 census date (collated by Esri 2021) with lighter yellow being lower population and darker orange being higher population.

Figure 3. Drag sampling effort by county (m^2) from 2017-2021. Black dots mark field sampling sites. Yellow dots with red circles mark sites where *A. americanum* were found. A) Sampling effort for 2017, 2018, 2019, 2021; B) Sampling effort for 2020 from June 8 – July 23 (See Table 1). The yellow square marks the established population of *A. americanum* at Grand Mere State Park (detected in this study). Yellow star marks a site sampled regularly as a reference for *I. scapularis* in order to ensure they are active regionally when sampling other sites. Only the first three visits for this site (i.e., during the nymphal host-seeking period) were included in the effort calculations for each year so as not to skew overall county effort.

Figure 4. Phenology of each questing life stage of A) *Amblyomma americanum* and B) *Ixodes scapularis* at Grand Mere State Park (Berrien County) from 6/11/20 – 10/10/20 (ended after 2 consecutive visits in the fall where no *A. americanum* were detected). See the yellow square on the map in Figure 2 for the location of the field site.

772
773
774
775 **Figure 1.** Numbers of *Dermacentor variabilis, Ixodes scapularis, Amblyomma americanum,* and other
776 species submitted annually to the Michigan Department of Agriculture (1999 – 2013) and Michigan
777 Department of Health and Human Services (2015 – 2020). On average, 3.2% ±1.7% SD of all ticks
778 submitted annually were *A. americanum.* Data source: MiTracking- Michigan Environmental Public
779 Health Tracking; no data are available for 2014.
780

781
782

Figure 2. Amblyomma americanum submitted annually to the Michigan Department Agriculture (2009-
2013) and the Michigan Department of Health and Human Services (MDHHS) from 2015-2020. Data
source: MiTracking- Michigan Environmental Public Health Tracking. Numbers in each county
represent the number of community- submitted *A. americanum* by county of residence. No data were
available for 2014. * Human population heatmap for each county is given for reference based on 2020
census date (collated by Esri 2021) with lighter yellow being lower population and darker orange being
higher population.

Figure 3. Drag sampling effort by county (m^2) from 2017-2021. Black dots mark field sampling sites.
Yellow dots with red circles mark sites where *A. americanum* were found. A) Sampling effort for 2017,
2018, 2019, 2021; B) Sampling effort for 2020 from June 8 – July 23 (See Table 1). The yellow square
marks the established population of *A. americanum* at Grand Mere State Park (detected in this study).
Yellow star marks a site sampled regularly as a reference for *I. scapularis* in order to ensure they are
active regionally when sampling other sites. Only the first three visits for this site (i.e., during the
nymphal host-seeking period) were included in the effort calculations for each year so as not to skew
overall county effort.

801
802

Figure 4. Phenology of each questing life stage of A) *Amblyomma americanum* and B) *Ixodes scapularis* at Grand Mere State Park (Berrien County) from 6/11/20 – 10/10/20 (ended after 2 consecutive visits in the fall where no *A. americanum* were detected). See the yellow square on the map in Figure 3 for the location of the field site.

807