1

Epithelial MHC class II directs microbiota-specific intestinal immune homeostasis

Emily M. Eshleman[1,3], Tzu-Yu Shao[2,3,4], Vivienne Woo[1,3], Taylor Rice[1,3], Jordan Whitt[1,3], Laura

Engleman[1,3], Sing Sing Way[2,3], and Theresa Alenghat[1,3]

[1]Division of Immunobiology, [2]Division of Infectious Disease, [3]Center for Inflammation and

Tolerance, [4]Immunology Graduate Program, Cincinnati Children's Hospital Medical Center and

the University of Cincinnati College of Medicine, Cincinnati, OH 45229 USA

CORRESPONDENCE: Theresa Alenghat, 3333 Burnet Ave, MLC 7038, Tel: 513-803-7498, Fax:

513-636-5355, theresa.alenghat@cchmc.org

Abstract

Dysregulated immune responses to resident microbes promote pathologic inflammation, however,

the mechanisms instructing commensal-specific T cells remain poorly understood. Here, we find

that non-hematopoietic intestinal epithelial cells (IECs) represent the primary cells expressing

major histocompatibility complex (MHC) II at the intestinal host-microbiota interface.

Interestingly, epithelial MHCII and commensal-specific CD4$^+$ T cells were concurrently induced

by post-natal microbiota colonization, provoking the hypothesis that epithelial MHCII regulates

local commensal-specific CD4$^+$ T cells. While MHCII on classical antigen presenting cells directs

expansion of antigen-specific CD4$^+$ T cells, loss of IEC-intrinsic MHCII surprisingly led to

elevated commensal-specific CD4$^+$ T cells in the intestine. Further, epithelial MHCII expression

actively limited accumulation of antigen-specific CD4$^+$ T cells in adult mice. Expansion of

commensal-specific Th17 cells was restricted by epithelial MHCII, and remarkably mice lacking

epithelial MHCII were highly susceptible to microbiota-triggered inflammation. Collectively,

these data indicate that impaired epithelial MHCII-T cell regulation within mucosal tissues alters

microbiota-specific immunity and predisposes to chronic inflammation.

Introduction

The mammalian gastrointestinal tract is home to trillions of microorganisms, collectively termed the microbiota, that form beneficial symbiotic relationships with host cells and play a significant role in health and disease. Extensive experiments utilizing broad-spectrum antibiotic or germ-free (GF) mouse models, which lack the microbiota, have demonstrated the necessity of a diverse and intact microbiota on mediating intestinal homeostasis[1,2]. These studies have highlighted that the microbiota directly interact with host cells to influence normal host physiology and tissue development. Digestion, intestinal barrier function, and immune cell development all rely on signals from the microbiota. Further, changes in the containment or diversity of the microbiota have been associated with the development of several chronic diseases including inflammatory bowel disease (IBD), asthma, cancer, and diabetes[1,3].

Intestinal immune responses are tightly regulated to allow protective immunity against invading pathogens, while limiting responses to innocuous commensals. It is well appreciated that the microbiota plays an important role in shaping CD4$^+$ T cell differentiation[4,5]. Commensal bacteria have been shown to drive regulatory T cell differentiation[6], and microbiota-reactive effector and memory T cells are present in both mice and humans[7–11]. Studies have demonstrated that commensal-specific T cells can contribute to intestinal homeostasis by promoting barrier protective cytokines and providing cross-reactivity to pathogens[10,12,13]. However, aberrant immune responses to the microbiota also promote inflammatory conditions, such as IBD[1,14–16]. In mouse models of colitis, intestinal dysbiosis drives inflammation in part by stimulating microbiota-reactive CD4$^+$ T cells[17–19]. Further, recent work found that microbiota-specific CD4$^+$ T cells in IBD patients were functionally altered and produced more pro-inflammatory cytokines compared

to healthy controls[10]. However, the mechanisms directing this critical balance between

commensal-specific effector and regulatory T cells in the intestine remain poorly understood.

Auto-reactive CD4[+] T cell responses are limited by presentation of self-antigens in the thymus by

cells expressing major histocompatibility complex-II (MHCII)[20–22]. In contrast, microbiota-

responsive CD4[+] T cells are unlikely to encounter their specific antigen within the thymus, and are

thus not subject to negative selection before entering peripheral tissues[23,24]. Previous work

described that negative selection of activated commensal-specific T cells can be mediated by

MHCII-expressing group 3 innate lymphoid cells (ILC3)[25,26]. However, ILC3s are relatively rare

cells at the luminal surface of the intestine. In contrast, intestinal epithelial cells (IECs) are

uniquely positioned at the direct interface between the microbiota and host immune cells and

express necessary machinery for antigen uptake, processing, and presentation[27–31]. While MHCII

expression by IECs was described over two decades ago, studies have only recently started to

investigate its function in inflammation and tissue homeostasis[32–35]. Results from models of

intestinal inflammation have demonstrated that epithelial MHCII regulation can be either

protective or detrimental depending on the inflammatory context[33–36]. However, it remains

unknown whether IEC expression of MHCII functionally regulates microbiota-triggered immune

responses.

Surprisingly, we found that IECs were the dominant cells expressing MHCII at the intestinal

luminal interface. Antibiotic-based approaches and germ-free mouse models demonstrated that

IEC-intrinsic MHCII expression was dramatically upregulated by the microbiota following birth.

Interestingly, this induction of MHCII was concurrent with the expansion of microbiota-specific

CD4$^+$ T cells in post-natal mice. This relationship provoked the hypothesis that IEC-intrinsic

MHCII may regulate immune responses to the microbiota. In contrast to a predicted activating role

of MHCII, we unexpectedly found that IEC-specific loss of MHCII *in vivo* resulted in increased

accumulation of microbiota-specific intestinal CD4$^+$ T cells. Further, IEC-intrinsic MHCII actively

limited commensal-specific CD4$^+$ T cells. Mice lacking IEC-intrinsic MHCII expression also

exhibited increased accumulation of RORγt$^+$ and decreased FoxP3$^+$ commensal-specific CD4$^+$ T

cells and were more susceptible to spontaneous, microbiota-triggered intestinal inflammation.

Taken together, these data reveal that microbiota colonization promotes intestinal tolerance by

inducing epithelial MHCII expression and thereby limiting local expansion of microbiota-specific

T cells.

Results

Commensal bacteria induce intestinal epithelial MHCII expression.

Antigen presentation via MHCII is critical for instructing CD4$^+$ T cell responses, however we have

limited understanding for how commensal-specific immune responses are established and

maintained in mucosal tissues colonized by microbes. Hematopoietic-derived antigen presenting

cells can regulate commensal-specific responses[37,38]. Surprisingly, though, we found that intestinal

epithelial cells (IECs), and not CD45$^+$ hematopoietic cells, expressed the majority of MHCII at the

intestinal-microbiota interface (**Fig. 1a, b**). IECs integrate microbial cues and instruct downstream

immune responses, thus, we next assessed whether the microbiota plays a role in regulating IEC-

intrinsic MHCII expression. Interestingly, IECs isolated from conventionally-housed (CNV) mice

displayed higher expression of *H2-Ab1* and *H2-Aa*, which encode MHCII, compared to IECs

isolated from GF mice (**Fig. 1c**). Consistent with this, surface MHCII protein expression was also

highly enriched in IECs from CNV mice compared to GF mice (**Fig. 1d, e**), whereas MHCII

expression on hematopoietic cells at the luminal interface was not altered by microbiota

(**Supplementary Fig. 1a**).

Microbiota colonization begins at birth with the complexity and density of microbiota increasing

most significantly during early life and into adulthood[39–41]. To next determine whether IEC-

intrinsic MHCII expression is modulated with microbiota colonization, IECs were assessed for

H2-Ab1 expression during development. Young wildtype pups with normal microbiota

colonization exhibited epithelial *H2-Ab1* expression on IECs that increased into adulthood,

whereas pups reared under GF conditions demonstrated low epithelial *H2-Ab1* expression that was

minimally altered with age (**Fig. 1f**), indicating that the microbiota strongly controls the induction

of epithelial MHCII expression with age. These results were recapitulated following antibiotic-

induced microbiota depletion (**Fig. 1g**), as pups born to mothers treated with antibiotics failed to

induce epithelial *H2-Ab1* expression (**Fig. 1h, i**). Taken together, these data demonstrate that post-

natal microbiota colonization triggers elevated epithelial MHCII expression in the intestine.

Epithelial MHCII expression can be regulated by local immune cells[33,42–44]. Therefore, to evaluate

whether MHCII is modulated in IECs in the absence of these cues, intestinal epithelial organoids

were derived from the colon of wildtype mice. These three-dimensional organoid cultures possess

physiologically relevant intestinal features including a polarized epithelial monolayer but are

devoid of immune cells (**Supplementary Fig. 1b**). Interestingly, incubation of intestinal organoids

with luminal contents collected from the colon of CNV mice robustly upregulated MHCII

expression (**Supplementary Fig. 1c**). However, contents collected from GF mice failed to induce

similar expression of MHCII (**Supplementary Fig. 1c**), indicating that microbiota are required for

triggering MHCII on IECs and that this can occur without external signals from other cells. Heat-

killing bacteria in intestinal contents of CNV mice completely ablated this induction

(**Supplementary Fig. 1d**), suggesting that MHCII expression is actively regulated by live bacteria

or heat-liable bacterial products. Past studies have indicated that IFNγ signaling can induce IEC

expression of MHCII[33,42–44]. However, microbiota exposure of IFNγ-deficient organoids still

induced MHCII expression (**Supplementary Fig. 1e**), further supporting that the microbiota

regulates epithelial MHCII in part via IEC-intrinsic mechanisms.

Commensal-specific CD4[+] T cells expand following post-natal microbial colonization.

Recent studies have highlighted that microbiota exposure early in life is required for effective

immune education and that disruption to the microbiota during this early window of development

may increase susceptibility to inflammatory disorders[40,41,45–47]. Therefore, to investigate whether

microbiota instruct development of commensal-specific T cell responses, intestinal CD4[+] T cells

were analyzed for reactivity to the class II-commensal flagella antigen-specific cBir1 tetramer.

CNV mice exhibited significantly higher numbers of cBir1[+] CD4[+] T cells in the large intestine

compared to GF mice that lack commensal microbes (**Fig. 2a, b**). Consistent with analyses by

others[48,49], intestinal CD4[+] T cells were reduced in GF mice relative to CNV mice (**Fig. 2c**) and

displayed a more naïve phenotype with decreased CD44 and increased CD62L expression

(**Supplementary Fig. 2a, b**). However, day 15 GF and CNV pups did not display differences in

CD4[+] T cell numbers in the large intestine (**Fig. 2c**). Thus, microbiota-derived signals are

necessary for accumulation and maturation of intestinal CD4[+] T cell responses. Interestingly, while

commensal-specific cBir1[+] CD4[+] T cells were expanded in adult CNV mice relative to adult GF

mice, commensal-specific CD4$^+$ T cells were similar between GF and CNV pups (**Fig. 2d, e**).

Collectively, these data suggest that the microbiota direct development of commensal-specific T

cell responses in the intestine with age.

Mice lacking IEC-intrinsic MHCII exhibit elevated microbiota-specific CD4+ T cells.

The temporal association between microbiota exposure, IEC-intrinsic MHCII expression, and

expansion of commensal bacterial-specific CD4$^+$ T cells provoked the hypothesis that IEC-

intrinsic MHCII expression may mediate the relationship between microbiota colonization and

regulation of microbiota-specific T cells. To begin to test this hypothesis, an IEC-specific MHCII

knockout mouse model was generated by crossing C57Bl/6 floxed H2-Ab1 mice (MHCII^{FF}) to

mice expressing cre recombinase downstream of the villin promoter[50,51]. Significant reduction in

MHCII expression in IECs was confirmed by mRNA (**Fig. 3a**) and surface protein (**Fig. 3b**)

analyses. Importantly, loss of MHCII expression was specific to IECs, as CD45$^+$ cells in the

intestine retained MHCII surface expression (**Fig. 3c**). Loss of IEC-intrinsic MHCII had minimal

effect on total CD4$^+$ T cell numbers in the intestine (**Fig. 3d**). However, MHCII$^{\Delta IEC}$ mice

surprisingly displayed elevated numbers of commensal-specific cBir1$^+$ CD4$^+$ T cells, relative to

floxed littermate control mice (**Fig. 3e**), suggesting that expression of MHCII by IECs regulates

microbiota-specific T cell homeostasis in the intestine.

Contrary to the canonical role of MHCII in promoting CD4$^+$ T cell expansion, mice with loss of

IEC-intrinsic MHCII surprisingly exhibited elevated commensal-specific CD4$^+$ T cells. Therefore,

to investigate the mechanism by which epithelial MHCII regulates tissue-intrinsic T cells,

proliferation of microbiota-specific T cells was compared in MHCII$^{\Delta IEC}$ mice (**Supplementary**

Fig. 3a). Loss of IEC-intrinsic MHCII expression had little effect on Ki67$^+$ levels in cBir1$^+$CD4$^+$

T cells (**Supplementary Fig. 3a**), suggesting that epithelial MHCII does not restrict proliferation

of commensal-specific T cells. Therefore, to next test whether antigen-experienced CD4$^+$ T cells

exhibited altered apoptosis, AnnexinV$^+$ levels were compared. Interestingly, loss of epithelial

MHCII expression resulted in decreased AnnexinV$^+$ CD4$^+$ T cells in the intestine compared to

littermate controls (**Supplementary Fig. 3b**). These data suggest that IEC-intrinsic MHCII

expression promotes deletion of local commensal-specific CD4$^+$ T cells and, therefore, these cells

accumulate in the intestine of MHCII$^{\Delta IEC}$ mice relative to control mice.

Epithelial MHCII actively regulates commensal-specific T cells in the intestine.

To evaluate whether IEC-intrinsic MHCII actively regulates CD4$^+$ T cell responses, we employed

a fungal commensal model that uses a recombinant *C. albicans* expressing 2W1S$_{55-68}$ variant of I-

Ea epitope (**Fig. 4a**)[52–54]. MHCII^{FF} and MHCII$^{\Delta IEC}$ mice colonized for 2 weeks with *C. albicans*-

2W1S$^+$ displayed similar levels of *C. albicans* intestinal colonization (**Fig. 4b**) with no evidence

of dissemination or infection. 2W1S$^+$ CD4$^+$ T cells were detected in the large intestine of mice

colonized with *C. albicans*-2W1S$^+$ (**Fig. 4c**). Interestingly, MHCII$^{\Delta IEC}$ mice displayed increased

numbers of *C. albicans*-2W1S-specific CD4$^+$ T cells relative to MHCII^{FF} mice (**Fig. 4c, d**),

indicating that IEC-intrinsic MHCII actively controls commensal-specific CD4$^+$ T cell levels in

the intestine of adult mice. Both MHCII^{FF} and MHCII$^{\Delta IEC}$ mice exhibited *Candida*-specific CD4$^+$

T cells that were characterized predominantly as FoxP3$^+$ Tregs, and fewer RORγt$^+$ Th17 cells (**Fig.**

4e). However, differentiation of *Candida*-specific CD4$^+$ T cells harvested from MHCII$^{\Delta IEC}$ mice

exhibited an elevated Th17 and diminished Treg profile relative to antigen-specific CD4$^+$ T cells

from MHCII^{FF} mice (**Fig. 4e, f**). Together, these results demonstrate that IEC-intrinsic MHCII

expression actively regulates the number and phenotype of commensal-specific CD4$^+$ T cells in

the intestine.

MHCII expression on IECs limits microbiota-induced intestinal inflammation.

Dysregulated commensal-specific CD4$^+$ T cells have been found in patients with intestinal

inflammation[10]. Remarkably, when MHCII$^{\Delta IEC}$ mice and littermate MHCII^{FF} controls were

permitted to age longer, MHCII$^{\Delta IEC}$ mice exhibited an increased incidence of rectal prolapse (**Fig.**

5a), a clinical symptom of intestinal inflammation. Consistent with this, levels of the inflammatory

biomarker, lipocalin-2, was elevated in the intestine of MHCII$^{\Delta IEC}$ mice relative to control floxed

mice (**Fig. 5b**), while barrier permeability was similar between these genotypes (**Supplementary**

Fig. 4a, b). Interestingly, MHCII$^{\Delta IEC}$ mice displaying an increased propensity to prolapse also

demonstrated the highest amount of commensal-specific cBir1$^+$CD4$^+$ T cells within these cohorts

(**Fig. 5c**).

In addition, similar to findings with the shorter *C. albicans*-2W1S$^+$ commensal model, the ratio of

FoxP3$^+$ Tregs to RORγt$^+$ Th17 cells was significantly altered in MHCII$^{\Delta IEC}$ mice (**Fig. 5d, e**). The

majority of cBir1$^+$CD4$^+$ T cells were FoxP3$^+$ Tregs (**Fig. 5d**), however, loss of IEC-intrinsic

MHCII resulted in a reduction of Tregs and an increase in Th17 commensal-reactive T cells

relative to the ratio observed in control MHCII^{FF} mice (**Fig. 5d, e**). IL-10 and IL-17 expression in

the large intestine were also correspondingly shifted in MHCII$^{\Delta IEC}$ mice compared to MHCII^{FF}

mice (**Fig. 5f, g**). This Treg to Th17 skewing of commensal-specific CD4$^+$ T cells and cytokine

expression was more evident in MHCII$^{\Delta IEC}$ mice that developed intestinal inflammation and

prolapse (**Fig. 5c-g**).

To test the requirement for the microbiota in this spontaneous intestinal inflammation, MHCII^{FF}

and MHCII$^{\Delta IEC}$ mice were compared to littermates of both genotypes in the context of broad-

spectrum antibiotics that significantly deplete commensal bacteria (**Supplementary Fig. 4c**).

Consistent with earlier results, MHCII$^{\Delta IEC}$ mice exhibited an increased prevalence of rectal

prolapse (**Fig. 5h**). However, depletion of the microbiota prevented spontaneous intestinal

inflammation in MHCII$^{\Delta IEC}$ mice (**Fig. 5h**). Furthermore, accumulation of commensal-specific

Th17 cells did not occur following microbiota depletion (**Fig. 5i**) and loss of microbiota inhibited

the IL-17 elevation observed in microbiota-replete MHCII$^{\Delta IEC}$ mice (**Fig. 5j**). Collectively, these

data support that regulation by epithelial MHCII promotes T cell tolerance to resident microbes

and limits development of microbiota-sensitive Th17-mediated intestinal inflammation.

Discussion

The microbiota is essential for the development and education of the host immune system.

However, inappropriate immune reactions to the microbiota have been associated with several

chronic inflammatory conditions. In this study, we found that upregulation of epithelial MHCII

expression in the large intestine coincided with initial microbiota colonization and expansion of

commensal-specific T cells, indicating that IECs may play an important role in early-life immune

education during development. Further, deletion of IEC-intrinsic MHCII expression led to

accumulation of commensal-specific Th17 cells in the large intestine that reflected impaired

apoptosis of commensal-specific T cells. Loss of IEC-intrinsic MHCII also elevated IL-17 in the

tissue and increased susceptibility to microbiota-sensitive intestinal inflammation.

(**Supplementary Fig. 5**).

Interestingly, patients with active IBD have functionally distinct microbiota-reactive CD4[+] T cells,

compared to those found in healthy controls, and secrete higher levels of IL-17[10,55]. IL-17 has been

linked to development and exacerbation of several autoimmune and inflammatory conditions,

including IBD[56]. Furthermore, mouse models with disrupted IL-17 signaling are protected from

intestinal inflammation[57,58], supporting a pathogenic role for IL-17 in intestinal inflammation.

However, clinical trials depleting IL-17 in IBD patients have been ineffective, and in some cases

worsened disease[59]. This pleotropic role for IL-17 in regulating intestinal homeostasis suggests

that targeting specific IL-17 producers, instead of broadly neutralizing IL-17 itself, may be a more

effective treatment. In fact, work has shown that therapeutics which target commensal-specific

Th17 cells, while retaining IL-17 production from other cell types, reduced intestinal

inflammation[60]. Our data suggests that promoting epithelial MHCII expression may further restrict

pro-inflammatory commensal-specific T cells and promote healthy intestinal homeostasis.

MHCII[ΔIEC] mice also exhibited a concurrent decrease in commensal-specific FoxP3[+] Treg cells

and intestinal IL-10 expression. Interestingly, colonization with a new microbe resulted in active

skewing of commensal-specific T cells toward Th17 cells in mice that lack epithelial MHCII,

whereas wildtype littermate controls developed more commensal-specific Treg cells. Therefore,

epithelial MHCII is necessary to control the delicate balance between suppressive and pro-

inflammatory antigen-specific T cells in the intestine. The requirement for IEC-intrinsic MHCII

in promoting commensal-reactive Tregs and limiting Th17 cells in the colon is consistent with

recent findings that epithelial MHCII expression in the small intestine affects IL-10 production by

local intraepithelial lymphocytes[34]. Taken together, these findings support that MHCII expression

specifically by epithelial cells is a dominant mechanism at the luminal surface that limits

commensal-specific CD4[+] T cells and dampens pathogenic responses to the microbiota. Given the

role for MHCII in ILC3s[25,26], complimentary and essential roles for antigen presentation pathways

in non-classical cells seem to be essential for establishing and maintaining healthy intestine-

intrinsic tissue homeostasis.

Crosstalk between the microbiota and mammalian immune cells involves communication via

pattern recognition receptor engagement and microbiota-derived metabolite signaling. While

MHCII expression in the intestine has been linked to the microbiota[61], we found that live microbes,

or heat-labile microbial components, are necessary for MHCII induction on these cells. Immune

cells can induce epithelial MHCII, however we determined that IECs can also intrinsically sense

the microbiota to upregulate MHCII expression by employing intestinal organoid cultures that lack

immune cells or IFNγ. Therefore, microbiota-dependent epithelial MHCII regulation also occurs,

in part, through direct communication that is independent of immune cells. Disruption to canonical

pattern recognition pathways leads to loss of intestinal barrier integrity and increased

inflammation[62–64] and the toll-like receptor signaling adaptors MyD88/TRAF have been shown to

induce epithelial MHCII expression, particularly in the small intestine[33]. However, further

investigation will be needed to decipher the host and microbial-derived pathways that are

responsible for epithelial MHCII regulation.

Recent studies have begun to investigate epithelial MHCII from multiple approaches. Loss of

epithelial MHCII protected mice that were subjected to models of T cell driven intestinal

inflammation and graft versus host disease[33,35]. Our findings highlight a broader role in which

epithelial cells are the primary MHC class II-expressing cells in the intestine controlling

microbiota-specific T cells and intestinal pathology induced by commensal bacteria

(**Supplementary Fig. 5**). Therefore, this work uncovers a fundamental new tenet for immune

regulation whereby IEC-intrinsic MHCII is largely induced by the microbiota to dampen self-

directed immune responses in early life. Antigen-presenting immune cells are traditionally viewed

to be most important in regulation of antigen specificity of CD4 T cells. However, these findings

reveal a central and essential epithelial-T cell mechanism that controls mucosal immunity

specifically to commensal microbes, and suggest that this regulatory pathway can be targeted to

treat microbiota-triggered inflammatory conditions.

Methods

Mice. Conventionally-housed C57Bl/6J mice were purchased from Jackson Laboratories and maintained in our specific-pathogen free colony at Cincinnati Children's Hospital Medical Center (CCHMC). Germ-free (GF) mice were maintained in flexible isolators in the CCHMC Gnotobiotic Mouse Facility, fed autoclaved feed and water, and monitored for absence of microbes. H2-Ab1(MHCII)^{FF} were purchased from Jackson and maintained at CCHMC. MHCII^{FF} mice were crossed to C57Bl/6J-Villin-Cre to generate MHCII$^{\Delta IEC}$ mice. Sex- and age-matched littermate controls were used for all studies. Mice were housed up to 4-per cage in a ventilated cage system in a 12-h light/dark cycle, with free access to water and food. For antibiotic treatment, pregnant mice were provided water supplemented with 1-mg/mL colistin (Sigma), 1-mg/mL ampicillin (Sigma) and 5-mg/mL streptomycin (Sigma) at mid-gestation[65]. Mothers and pups were maintained on antibiotic-water until harvest as noted. For MHCII^{FF} and MHCII$^{\Delta IEC}$ mice, pups were placed on antibiotic-water at weaning and maintained on antibiotics for 16-weeks. Antibiotic-water was refreshed every 7-10 days. *Candida albicans* colonization was conducted as previously described[52,53]. Briefly, mice were pre-treated with ampicillin-water (1-mg/mL) two days prior to colonization and maintained on ampicillin-water for the duration of the experiment. Mice were gavaged with recombinant *C. albicans* expressing 2W1S$_{55-68}$ peptide[54]. Mice were monitored for colonization by plating fecal CFUs and were harvested 14-days post-colonization. All mouse studies were conducted with approval by Animal Care and Use Committees at CCHMC. These protocols follow standards enacted by the United States Public Health Services and Department of Agriculture. All experiments followed standards set forth by Animal Research: Reporting of In Vivo Experiments (ARRIVE).

Cell Isolation. The large intestine was harvested, opened, and washed in PBS. For IECs, tissue

was placed in pre-warmed strip buffer (PBS, 5% FBS, 1mM EDTA, 1mM DTT) and incubated at

37°C at 45-degree angle with shaking at 180-rpm for 15-min. For lamina propria isolation, tissue

was washed with PBS to remove EDTA and DTT then incubated in pre-warmed digestion buffer

(RPMI with 1-mg/mL Collagenase/Dispase (Sigma)) at 37°C at 45-degree angle with shaking at

180-rpm for 30-min. After incubation, the tissue was vortexed and passed through a 70µM cell

strainer.

Flow cytometry. Cells were stained for flow cytometry using the following antibodies diluted in

FACS Buffer (2% FBS, 0.01% Sodium Azide, PBS): BV711-anti-CD326 (EpCAM) (Clone:G8.8,

BioLegend), BUV395-anti-CD45.2 (Clone:104, BD Biosciences), APC-anti-MHCII

(Clone:M5/114.15.2, eBioscience), APC-eFluor-780-anti-CD4 (Clone:RM4-5, eBioscience), Pe-

Cy7-anti-CD44 (Clone:IM7, eBioscience), AlexaFluor-647-anti-RORγt (Clone:Q31-378, BD

Biosciences), PacificBlue-anti-FoxP3 (Clone:FJK-16s, eBiosciences), PerCP-eFluor710-anti-

CD8a (Clone:53-6.7, eBiosciences), PerCP-eFluor710-anti-B220 (Clone:RA3-6B2,

eBiosciences), PerCP-eFluor710-anti-Ly6G (Clone:1A8-Ly6g, eBiosciences), PerCP-Cy5.5-anti-

CD11b (Clone:M1/70, eBiosciences), PerCP-Cy5.5-anti-CD11c (Clone:N418, eBiosciences),

PacificBlue-anti-Ki67 (Clone:SolA15, eBiosciences). Dead cells were excluded with the Fixable

Aqua Dead Cell Stain Kit (Invitrogen). The BD Fix/Perm kit was used for intracellular staining.

Cells were stained with 488-congugated-AnnexinV diluted in AnnexinV staining buffer (BD

Pharamgen). Class-II restricted tetramers (cBir1 and 2W1S) were PE-conjugated and were

generated and provided by the NIH tetramer core. Samples were acquired on the BD LSRFortessa

and analyzed with FlowJo Software (Treestar).

Organoid cultures. Murine organoids were generated from colonic crypts isolated from GF and

CNV, or B6.IFNγ[-/-66] mice[67]. Briefly, the colon was opened, cleaned, and cut into 1-cm pieces.

Tissue was incubated in chelation buffer (2mM EDTA in PBS) for 30-min at 4°C with rotation

then transferred into shaking buffer (PBS, 43.3mM sucrose, 54.9mM sorbitol) and shaken by hand.

Crypts were plated in Matrigel (Corning) with organoid culture media (60% Advanced

DMEM/F12 supplemented with 10mM HEPES, 2mM L-glutamate, 40% L-WRN conditioned

media, 1xN2, 1xB27, 50-ng/mL murine EGF, and 10μM Y-27632) overlaid. Media was changed

every 2-3 days. For microbial stimulation, stool was collected from either GF or CNV mice and

homogenized in PBS using a TissueLyser (Qiagen) at a concentration of 100-mg of stool

weight/mL of PBS. Organoid cultures were treated with 100-μL of intestinal contents for 24-hrs.

Heat-killing was performed at 95°C for 1-hr. After incubation, cultures were washed with PBS,

Matrigel was disrupted, and treated with TrypLE (Gibco) for 3-5 mins to dissociate into single-

cell suspension. Cells were stained for flow cytometry. For immunostaining, organoids were plated

in 8-well u-slide (ibidi) and fixed in 4% paraformaldehyde for 20-min. Organoids were

permeabilized for 20-min in PBS containing 0.5% TritonX-100 followed by a 30-min incubation

in Blocking Buffer (PBS, 0.2% TritonX-100, 0.05% Tween, 1% BSA, and 1% normal goat serum).

Organoids were stained overnight at 4°C in blocking buffer containing AlexaFluor488-anti-

EpCAM (Clone:G8.8, BioLegend). Organoids were washed and imaged using a confocal

microscope (Nikon).

RNA and qPCR analysis. RNA was isolated using the RNeasy Kit (Qiagen). For RNA from whole

tissue, samples were homogenized in TRIzol. Choroform was added for phase separation and RNA

was precipitated by mixing with isopropanol. cDNA was synthesized using the Verso reverse transcriptase kit (Thermo Fisher) following the manufacturer's protocol. Real-time PCR was performed using SYBR green (Applied Biosystems) and analyzed using the following primers sequences. HPRT forward 5'- GATTAGCGATGAACCAGGT-3', HPRT reverse 5'- CCTCCCATCTCCTTCATGACA-3', H2-Ab1 forward 5'- TGTGAGTCCTGGTGACTGCCATTA-3', H2-Ab1 reverse 5'- TCGCCCATGAACTGGTACACGAAA-3', H2-Aa forward 5'- CAACCGTGACTATTCCTTCC-3', H2-Aa reverse 5'- CCACAGTCTCTGTCAGCTC-3', IL-10 forward 5'- GAGAGCTGCAGGGCCCTTTGC-3', IL-10 reverse 5'- CTCCCTGGTTTCTCTTCCCAAGACC-3', IL-17 forward 5'- ACCGCAATGAAGACCCTGAT-3', IL-17 reverse 5'- TCCCTCCGCATTGACACA-3'.

Lipocalin-2 ELISA and FITC analyses. Fecal pellets were homogenized in PBS at a concentration of 100-mg/mL then centrifuged at high-speed for 10-mins. Supernatants were collected and lipocalin-2 levels were determined using mouse Lipocalin-2/NGAL (R&D Systems) following manufacturer's instructions. For FITC permeability studies, mice were fasted for at least 4-hrs before being gavaged with FITC-dextran (0.6 mg/g of mouse body weight) diluted in PBS. Serum was collected 4-hrs post gavage and fluorescent intensity was measured (excitation 485nm/emission 530nm).

Statistical analysis. Results are expressed as mean ± SEM. Statistical significance was determined with the Student's t-test or one-way ANOVA with all data meeting the assumptions of the statistical test used. Results were considered significant at $*p < 0.05$; $**p < 0.01$; $***p < 0.001$,

****p < 0.0001. Statistical significance was calculated using Prism version 7.0 (GraphPad

Software).

Acknowledgements. We thank the Way, Qualls, and Deshmukh labs for useful discussions and

members of the Alenghat lab for critical reading of the manuscript. We thank CCHMC Veterinary

Services, Research Flow Cytometry Core, Confocal Imaging Core, and Pathology Research Core

for services and technical assistance. This research is supported by the National Institutes of Health

(DK114123, DK116868 to T.A., and F32AI147591 to E.M.E.), and a Kenneth Rainin Foundation

award to T.A. T.A. holds an Investigator in the Pathogenesis of Infectious Disease Award from

the Burroughs Wellcome Fund. This project is supported in part by PHS grant P30 DK078392 and

the CCHMC Trustee Award and Procter Scholar's Program.

Author Contributions. Conceptualization, E.M.E., T.A.; Methodology, E.M.E, T.-Y.S., V.W.,

L.E., J.W., S.S.W., T.A.; Investigation, E.M.E., J.W., T.-Y.S., V.W., T.R.; Formal Analysis,

E.M.E, T.-Y.S., T.R.; Writing, Review, and Editing, E.M.E., T.A.

Fig. 1. Commensal bacteria induce intestinal epithelial MHCII expression. (**a-b**) MHCII[+] expressing cells at colonic luminal surface. (**c**) mRNA expression of MHCII genes H2-Ab1 and H2-Aa in IECs isolated from large intestine of GF and CNV mice. (**d**) Surface expression and (**e**) MFI of MHCII on EpCAM[+] cells. (**f**) mRNA expression of H2-Ab1 in IECs isolated from large intestine of GF and CNV mice at multiple timepoints. (**g**) Relative 16S rRNA gene levels in intestinal contents. (**h**) IEC H2-Ab1 mRNA expression and (**i**) MHCII surface expression (gated on CD45[-], EpCAM[+]) in control and antibiotic (ABX)-treated mice. Data are representative of at least three independent experiments, 3-4 mice per group, per timepoint. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Fig. 2. Commensal-specific CD4[+] T cells expand following post-natal microbial colonization. (**a-b**) Number of cBir1[+] tetramer-specific CD4[+] T cells in large intestine of adult GF and CNV mice. (**c**) Frequency of CD4[+] T cells and (**d-e**) number of cBir1[+] tetramer-specific CD4[+] T cells isolated from large intestine of neonatal and adult GF and CNV mice. cBir1[+] tetramer cells are gated on live, CD45[+], lineage (CD11b, B220, Ly6G, CD11c, CD8a)[-], CD4[+]. Data represent at least two independent experiments, 3-4 mice per group. *p<0.05, **p<0.01, ***p<0.001, ns=not significant.

Fig. 3. Mice lacking IEC-intrinsic MHCII exhibit elevated microbiota-specific CD4[+] T cells. (**a**) H2-Ab1 mRNA expression in IECs isolated from the large intestine of MHCII[FF] and

MHCII$^{\Delta\text{IEC}}$ mice. (**b**) Surface MHCII expression in EpCAM$^+$, CD45$^-$ cells from MHCII$^{\text{FF}}$ and

MHCII$^{\Delta\text{IEC}}$ mice. (**c**) Surface MHCII expression in EpCAM$^-$, CD45$^+$ cells from MHCII$^{\text{FF}}$ and

MHCII$^{\Delta\text{IEC}}$ mice. (**d**) Frequency of total CD4$^+$ and (**e**) number of cBir1$^+$ tetramer-specific CD4$^+$ T

cells isolated from large intestine of MHCII$^{\text{FF}}$ and MHCII$^{\Delta\text{IEC}}$ mice. Gated on live, CD45$^+$, lineage

(CD11b, B220, Ly6G, CD11c, CD8a)$^-$, CD4$^+$. Data represent at least three independent

experiments, 3-4 mice per group. *p<0.05, **p<0.01, ***p<0.001.

Fig. 4. Epithelial MHCII actively regulates commensal-specific T cells in the intestine. (**a**)

Schematic of *C. albicans* commensal colonization. (**b**) Colony forming units (CFUs) of *C. albicans*

in stool from MHCII$^{\text{FF}}$ and MHCII$^{\Delta\text{IEC}}$ mice. (**c, d**) Number of *C. albicans*-2W1S$^+$-specific CD4$^+$

cells in large intestine of MHCII$^{\text{FF}}$ and MHCII$^{\Delta\text{IEC}}$ mice. (**e**) Frequency of FoxP3$^+$ and (**f**) RORγt$^+$

C. albicans-2W1S$^+$-specific CD4$^+$ cells in large intestine of MHCII$^{\text{FF}}$ and MHCII$^{\Delta\text{IEC}}$ mice. Gated

on live, CD45$^+$, lineage (CD11b, B220, Ly6G, CD11c, CD8a)$^-$, CD4$^+$. Data represent at least two

independent experiment, 3-4 mice per group. *p<0.05.

Fig. 5. MHCII expression on IECs limits microbiota-induced intestinal inflammation. (**a**)

Frequency of rectal prolapse in MHCII$^{\text{FF}}$ and MHCII$^{\Delta\text{IEC}}$ mice. (**b**) Lipocalin-2 levels in stool of

MHCII$^{\text{FF}}$, MHCII$^{\Delta\text{IEC}}$, and prolapsed MHCII$^{\Delta\text{IEC}}$ mice. (**c**) cBir1$^+$-specific CD4$^+$ T cells in large

intestine of MHCII$^{\text{FF}}$, MHCII$^{\Delta\text{IEC}}$, and prolapsed MHCII$^{\Delta\text{IEC}}$ mice. (**d**) Frequency of FoxP3$^+$ and

(**e**) RORγt$^+$ cBir1$^+$-specific CD4$^+$ T cells. (**f**) IL-10 and (**g**) IL-17 mRNA in large intestine of

MHCII$^{\text{FF}}$, MHCII$^{\Delta\text{IEC}}$, and prolapsed MHCII$^{\Delta\text{IEC}}$ mice. (**h**) Frequency of rectal prolapse in control

and antibiotic (ABX)-treated MHCII$^{\text{FF}}$ and MHCII$^{\Delta\text{IEC}}$ mice. (**i**) Frequency of RORγt$^+$ cBir1$^+$-

specific CD4$^+$ T cells in large intestine. (**j**) IL-17 mRNA expression in large intestine of control

and ABX-treated MHCII^{FF} and MHCII$^{\Delta IEC}$ mice. Data are representative of at least two

independent experiments, 3-6 mice per group. *$p<0.05$, **$p<0.01$, ***$p<0.001$, ns=not significant.

References

1. Hill, D. A. & Artis, D. Intestinal bacteria and the regulation of immune cell homeostasis. *Annu. Rev. Immunol.* **28**, 623–667 (2010).

2. Belkaid, Y. & Harrison, O. J. Homeostatic Immunity and the Microbiota. *Immunity* **46**, 562–576 (2017).

3. Kayama, H., Okumura, R. & Takeda, K. Interaction between the Microbiota, Epithelia, and Immune Cells in the Intestine. *Annu. Rev. Immunol.* **38**, 23–48 (2020).

4. Honda, K. & Littman, D. R. The microbiota in adaptive immune homeostasis and disease. *Nature* **535**, 75–84 (2016).

5. Brown, E. M., Kenny, D. J. & Xavier, R. J. Gut Microbiota Regulation of T Cells During Inflammation and Autoimmunity. *Annu. Rev. Immunol.* **37**, 599–624 (2019).

6. Honda, K. & Littman, D. R. The Microbiome in Infectious Disease and Inflammation. *Annu. Rev. Immunol.* **30**, 759–795 (2012).

7. Duchmann, R. *et al.* T cell specificity and cross reactivity towards enterobacteria, Bacteroides, Bifidobacterium, and antigens from resident intestinal flora in humans. *Gut* **44**, 812–818 (1999).

8. Duchmann, R. *et al.* Tolerance exists towards resident intestinal flora but is broken in active inflammatory bowel disease (IBD). *Clin. Exp. Immunol.* **102**, 448–455 (1995).

9. Ergin, A. *et al.* Impaired peripheral Th1 CD4+ T cell response to escherichia coli proteins in patients with crohn's disease and ankylosing spondylitis. *J. Clin. Immunol.* **31**, 998–1009 (2011).

10. Hegazy, A. N. *et al.* Circulating and Tissue-resident CD4+ T Cells With Reactivity to Intestinal Microbiota Are Abundant in Healthy Individuals and Function is Altered During Inflammation. *Gastroenterology* **153**, 1320–1337 (2017).

11. Hand, T. W. *et al.* Acute Gastrointestinal Infection Induces Long-Lived Microbiota-Specific T Cell Responses. *Science (80-.).* **337**, 1553–1556 (2012).

12. Su, L. F., Kidd, B. A., Han, A., Kotzin, J. J. & Davis, M. M. Virus-Specific CD4+ Memory-Phenotype T Cells Are Abundant in Unexposed Adults. *Immunity* **38**, 373–383 (2013).

13. Campion, S. L. *et al.* Proteome-wide analysis of HIV-specific naive and memory CD4+ T cells in unexposed blood donors. *J. Exp. Med.* **211**, 1273–1280 (2014).

14. Kaser, A., Zeissig, S. & Blumberg, R. S. Inflammatory Bowel Disease. *Annu. Rev. Immunol.* **28**, 573–621 (2010).

15. Tindemans, I., Joosse, M. E. & Samsom, J. N. Dissecting the Heterogeneity in T-Cell Mediated Inflammation in IBD. *Cells* **9**, 110 (2020).

16. Clemente, J. C., Ursell, L. K., Parfrey, L. W. & Knight, R. The impact of the gut microbiota on human health: An integrative view. *Cell* **148**, 1258–1270 (2012).

17. Belkaid, Y., Bouladoux, N. & Hand, T. W. Effector and memory T cell responses to commensal bacteria. *Trends Immunol.* **34**, 299–306 (2013).

18. Maloy, K. J. & Powrie, F. Intestinal homeostasis and its breakdown in inflammatory bowel disease. *Nature* **474**, 298–306 (2011).

19. Cong, Y. *et al.* CD4+ T cells reactive to enteric bacterial antigens in spontaneously colitic C3H/HeJBir mice: Increased T helper cell type 1 response and ability to transfer disease. *J. Exp. Med.* **187**, 855–864 (1998).

20. Blackman, M., Kappler, J. & Marrack, P. The role of the T cell receptor in positive and negative selection of developing T cells. *Science (80-.).* **248**, 1335–1341 (1990).

21. Mathis, D. & Benoist, C. Aire. *Annu. Rev. Immunol.* **27**, 287–312 (2009).

22. Klein, L., Kyewski, B., Allen, P. M. & Hogquist, K. A. Positive and negative selection of
the T cell repertoire: What thymocytes see (and don't see). *Nat. Rev. Immunol.* **14**, 377–
391 (2014).

23. Cong, Y., Feng, T., Fujihashi, K., Schoeb, T. R. & Elson, C. O. A dominant, coordinated
T regulatory cell-IgA response to the intestinal microbiota. *Proc. Natl. Acad. Sci. U. S. A.*
106, 19256–19261 (2009).

24. Ai, T. L., Solomon, B. D. & Hsieh, C. S. T-cell selection and intestinal homeostasis.
Immunol. Rev. **259**, 60–74 (2014).

25. Hepworth, M. R. *et al.* Innate lymphoid cells regulate CD4+ T-cell responses to intestinal
commensal bacteria. *Nature* **498**, 113–117 (2013).

26. Hepworth, M. R. *et al.* Group 3 innate lymphoid cells mediate intestinal selection of
commensal bacteria-specific CD4+ T cells. *Science (80-.).* **348**, 1031–1035 (2015).

27. Büning, J. *et al.* Antigen targeting to MHC class II-enriched late endosomes in colonic
epithelial cells: trafficking of luminal antigens studied in vivo in Crohn's colitis patients.
Faseb J. **20**, 359–361 (2006).

28. Bland, P. W. & Warren, L. G. Antigen presentation by epithelial cells of the rat small
intestine I.Kinetics, antigen specificity and blocking by anti-Ia antisera. *Immunology* **58**,
1–7 (1986).

29. Mayer, L. & Shlien, R. Evidence of function of Ia molecules on gut epithelial cells in
man. *J. Exp. Med.* **166**, 1471–1483 (1987).

30. Hershberg, R. M. *et al.* Intestinal epithelial cells use two distinct pathways for HLA class
II antigen processing. *J. Clin. Invest.* **100**, 204–15 (1997).

31. Hershberg, R. M. *et al.* Highly polarized HLA class II antigen processing and presentation
by human intestinal epithelial cells. *J. Clin. Invest.* **102**, 792–803 (1998).

32. Biton, M. *et al.* T Helper Cell Cytokines Modulate Intestinal Stem Cell Renewal and
Differentiation. *Cell* **175**, 1307-1320.e22 (2018).

33. Koyama, M. *et al.* MHC Class II Antigen Presentation by the Intestinal Epithelium
Initiates Graft-versus-Host Disease and Is Influenced by the Microbiota. *Immunity* **51**,
885-898.e7 (2019).

34. Tuganbaev, T. *et al.* Diet Diurnally Regulates Small Intestinal Microbiome-Epithelial-
Immune Homeostasis and Enteritis. *Cell* **182**, 1441-1459.e21 (2020).

35. Jamwal, D. R. *et al.* Intestinal Epithelial Expression of MHCII Determines Severity of
Chemical, T-Cell–Induced, and Infectious Colitis in Mice. *Gastroenterology* **159**, 1342-
1356.e6 (2020).

36. Thelemann, C. *et al.* Interferon-γ induces expression of MHC class II on intestinal
epithelial cells and protects mice from colitis. *PLoS One* **9**, 1–10 (2014).

37. Loschko, J. *et al.* Absence of MHC class II on cDCs results in microbial-dependent
intestinal inflammation. *J. Exp. Med.* **213**, 517–534 (2016).

38. Maggio-Price, L. *et al.* Lineage targeted MHC-II transgenic mice demonstrate the role of
dendritic cells in bacterial-driven colitis. *Inflamm. Bowel Dis.* **19**, 174–184 (2013).

39. Aagaard, K., Stewart, C. J. & Chu, D. Una destinatio, viae diversae. *EMBO Rep.* **17**,
1679–1684 (2016).

40. Russell, S. L. *et al.* Early life antibiotic-driven changes in microbiota enhance
susceptibility to allergic asthma. *EMBO Rep.* **13**, 440–447 (2012).

41. Gensollen, T., Iyer, S. S., Kasper, D. L. & Blumberg, R. S. How colonization by

microbiota in early life shapes the immune system. *Science (80-.).* **352**, 539–544 (2016).

42. Blumberg, R. S. *et al.* Antigen presentation by intestinal epithelial cells. *Immunol. Lett.*
69, 7–11 (1999).

43. Matsumoto, S. *et al.* Gamma delta TCR-bearing intraepithelial lymphocytes regulate class
II major histocompatibility complex molecule expression on the mouse small intestinal
epithelium. *Epithelial Cell Biol.* **4**, 163–170 (1995).

44. Matsumoto, S. *et al.* Physiological roles of γδ T-cell receptor intraepithelial lymphocytes
in cytoproliferation and differentiation of mouse intestinal epithelial cells. *Immunology* **97**,
18–25 (1999).

45. Al Nabhani, Z. *et al.* A Weaning Reaction to Microbiota Is Required for Resistance to
Immunopathologies in the Adult. *Immunity* **50**, 1276-1288.e5 (2019).

46. Wang, T. *et al.* Early life antibiotic exposure and host health: Role of the microbiota–
immune interaction. *Semin. Perinatol.* **44**, 151323 (2020).

47. Gray, J. *et al.* Intestinal commensal bacteria mediate lung mucosal immunity and promote
resistance of newborn mice to infection. *Sci. Transl. Med.* **9**, 1–14 (2017).

48. Chung, H. *et al.* Gut immune maturation depends on colonization with a host-specific
microbiota. *Cell* **149**, 1578–1593 (2012).

49. Visekruna, A. *et al.* Intestinal development and homeostasis require activation and
apoptosis of diet-reactive T cells. *J. Clin. Invest.* **129**, 1972–1983 (2019).

50. Hashimoto, K., Joshi, S. K. & Koni, P. A. A conditional null allele of the major
histocompatibility IA-beta chain gene. *Genesis* **32**, 152–153 (2002).

51. El Marjou, F. *et al.* Tissue-specific and inducible Cre-mediated recombination in the gut
epithelium. *Genesis* **39**, 186–193 (2004).

52. Jiang, T. T. *et al.* Commensal Fungi Recapitulate the Protective Benefits of Intestinal
Bacteria. *Cell Host Microbe* **22**, 809-816.e4 (2017).

53. Shao, T. Y. *et al.* Commensal Candida albicans Positively Calibrates Systemic Th17
Immunological Responses. *Cell Host Microbe* **25**, 404-417.e6 (2019).

54. Igyártó, B. Z. *et al.* Skin-Resident Murine Dendritic Cell Subsets Promote Distinct and
Opposing Antigen-Specific T Helper Cell Responses. *Immunity* **35**, 260–272 (2011).

55. Annunziato, F. *et al.* Phenotypic and functional features of human Th17 cells. *J. Exp.*
Med. **204**, 1849–1861 (2007).

56. Ouyang, W., Kolls, J. K. & Zheng, Y. The Biological Functions of T Helper 17 Cell
Effector Cytokines in Inflammation. *Immunity* **28**, 454–467 (2008).

57. Schmidt, E. G. W. *et al.* TH17 cell induction and effects of IL-17A and IL-17F blockade
in experimental colitis. *Inflamm. Bowel Dis.* **19**, 1567–1576 (2013).

58. Tang, C. *et al.* Suppression of IL-17F, but not of IL-17A, provides protection against
colitis by inducing T reg cells through modification of the intestinal microbiota. *Nat.*
Immunol. **19**, 755–765 (2018).

59. Hueber, W. *et al.* Secukinumab, a human anti-IL-17A monoclonal antibody, for moderate
to severe Crohn's disease: unexpected results of a randomised, double-blind placebo-
controlled trial. *Gut* **61**, 1693–1700 (2012).

60. Withers, D. R. *et al.* Transient inhibition of ROR-γt therapeutically limits intestinal
inflammation by reducing TH17 cells and preserving group 3 innate lymphoid cells. *Nat.*
Med. **22**, 319–323 (2016).

61. El Aidy, S. *et al.* Temporal and spatial interplay of microbiota and intestinal mucosa drive
establishment of immune homeostasis in conventionalized mice. *Mucosal Immunol.* **5**,

567–579 (2012).

62.   Rakoff-Nahoum, S., Pglino, J., Eslami-Varzaneh, F., Edberg, S. & Medzhitov, R.
Recognition of comensal microflora by toll-like receptors in required for intestinal
homeostasis. *Cell* **118**, 229–241 (2004).

63.   Rhee, S. H. *et al.* Pathophysiological role of Toll-like receptor 5 engagement by bacterial
flagenllin in colonic inflammation. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 13610–13615
(2005).

64.   Rose, W. A. I., Sakamoto, K. & Leifer, C. A. TLR9 is important for protection against
intestinal damage and for intestinal repair. *Sci. Rep.* **2**, 1–9 (2012).

65.   Vétizou, M. *et al.* Anticancer immunotherapy by CTLA-4 blockade relies on the gut
microbiota. *Science (80-.).* **350**, 1079–1084 (2015).

66.   Dalton, D. K. *et al.* Multiple defects of immune cell function in mice with disrupted
interferon-γ genes. *Science (80-.).* **259**, 1739–1742 (1993).

67.   Waddell, A., Vallance, J. E., Hummel, A., Alenghat, T. & Rosen, M. J. IL-33 Induces
Murine Intestinal Goblet Cell Differentiation Indirectly via Innate Lymphoid Cell IL-13
Secretion. *J. Immunol.* **202**, ji1800292 (2018).

Fig. 1. Microbiota colonization induces epithelial MHCII expression.

Fig. 2. Commensal-specific CD4+ T cell responses expand with microbiota colonization.

Fig. 3. Microbiota-specific CD4+ T cells are elevated in the colon of MHCII$^{\Delta IEC}$ mice.

Fig. 4. Epithelial MHCII actively regulates microbiota-specific T cells in the colon.

Fig. 5. Epithelial MHCII expression limits microbiota-dependent intestinal inflammation.