Genetically-linked simultaneous overexpression of multiple

herbicide-metabolizing genes for broad-spectrum resistance in an agricultural

weed *Echinochloa phyllopogon*

Hiroe Suda[1], Tomomi Kubo[1], Yusuke Yoshimoto[1], Keisuke Tanaka[2], Satoru Tanaka[3],

Akira Uchino[4], Satoshi Azuma[5], Makoto Hattori[5], Takuya Yamaguchi[6], Masahiro

Miyashita[1], Tohru Tominaga[1], Satoshi Iwakami[1]

[1] Graduate School of Agriculture, Kyoto University, Kitashirakawa-Oiwake-cho, Sakyo-ku,

Kyoto 606-8502, Japan

[2] NODAI Genome Research Center, Tokyo University of Agriculture, 1-1-1 Sakuragaoka,

Setagaya-ku, Tokyo 156-8502, Japan

[3] Faculty of Regional Environment Science, Tokyo University of Agriculture, 1-1-1

Sakuragaoka, Setagaya-ku, Tokyo 156-8502, Japan

[4] Central Region Agricultural Research Center, National Agriculture and Food Research

Organization, Tsu 514-2392, Japan

[5] Niigata Agricultural Research Institute Crop Research Center, 857 Nagakura-machi,

Nagaoka, Niigata 940-0826, Japan

[6] Biotechnology Research Center and Department of Biotechnology, Toyama Prefectural

University, 5180 Kurokawa, Imizu, Toyama 939-0398, Japan

Corresponding Author: Satoshi Iwakami

Tel: +81 75 753 6064

Email: iwakami.satoshi.2v@kyoto-u.ac.jp

Summary

• Previous research unveiled that the overexpression of catalytically promiscuous

CYP81A cytochrome P450s underlies the multiple-herbicide resistance (MHR) in a

Californian population of *Echinochloa phyllopogon*. However, it does not fully

accommodate the resistance to diverse herbicides in MHR *E. phyllopogon* although

the genetic inheritance of MHR was suggested as under a single gene control.

• We investigated the high-level resistance to diclofop-methyl in MHR *E. phyllopogon*.

Detailed diclofop-methyl metabolism was analyzed, followed by gene expression

study and functional characterization of P450 genes. The generality of the MHR

mechanism was investigated using another MHR line.

• The MHR line rapidly produced two distinct hydroxylated-diclofop-acid, only one of

which was the major metabolite produced by CYP81A12/21. Gene expression study

identified the genetically linked overexpression of a novel gene *CYP709C69* with

CYP81A12/21 in the MHR line. The gene conferred diclofop-methyl resistance in

plants and produced another hydroxylated-diclofop-acid in yeast. The activity was

observed in some CYP709C in plants. Unlike the broad substrate-specificity in

CYP81As, CYP709C69 showed narrow substrate-specificity. The overexpression of

the CYP81A and CYP709C69 was also observed in another MHR line.

• The present findings establish a novel concept that genetically-linked simultaneous

overexpression of herbicide-metabolizing genes enhances and broadens the profile

of metabolic resistance in weeds.

Key words: cytochrome P450, herbicide resistance, evolution, cross-resistance,

NIH-shift

Introduction

Heavy reliance on herbicide for weed control in agricultural fields have led to the evolution of herbicide resistance in weeds (Powles & Yu, 2010). The devastating impact on agriculture can be brought about by enhanced herbicide-metabolism since weeds with the mechanism often exhibit a large scope of resistance to chemically unrelated herbicides (Yu & Powles, 2014). Cytochrome P450 monooxygenases (P450s) have been assigned as a major component of metabolism-based resistance in weeds. Studies over the past 20 years have clearly shown that some P450s especially from crops carry metabolizing activity to some herbicides, leading to the protection plants from herbicide injuries (Dimaano & Iwakami, 2021). The involvement of P450s in weed resistance is beginning to be understood at the molecular level, but it has not been fully outlined. In particular, how multiple-herbicide resistance (MHR) is realized needs to be studied in more detail.

Echinochloa phyllopogon (syn. *Echinochloa oryzicola*) is an allotetraploid (2x=4x=36) and predominantly self-fertilizing noxious weed in rice cultivation (Yamasue, 2001). The first case of resistance in *E. phyllopogon* was reported from California in the late 90s' where the resistant populations exhibited resistance not only to the herbicides used in the fields at that time but also to many other herbicides commercialized later (Nandula *et al.*, 2019). Intensive analyses of this MHR population have led to the discovery that the overexpression of two CYP81A P450s is responsible for the cross-resistance to many herbicides such as acetolactate synthase (ALS) inhibitors (Iwakami *et al.*, 2014a; Dimaano *et al.*, 2020), acetyl-CoA carboxylase (ACCase) inhibitors (Iwakami *et al.*, 2019), a 1-deoxy-D-xylulose 5-phosphate synthase (DXS) inhibitor (Guo *et al.*, 2019), and a synthetic auxin (Chayapakdee, 2019). The two P450s,

CYP81A12 and CYP81A21, exhibit broad substrate specificity, i.e. catalytic promiscuity,

against diverse herbicide chemistries, accommodating the resistance to multiple

herbicides in the *E. phyllopogon* population (Dimaano *et al.*, 2020). Through genetic

segregation study, the overexpression of the two P450s was suggested to be controlled

by an unidentified single element (Iwakami *et al.*, 2014a). Recently, similar cases have

been reported in other weeds, *Lolium rigidum* and *Echinochloa crus-galli*, where

multiple-herbicide resistance is also mediated by the overexpression of

herbicide-metabolizing CYP81A P450s (Han *et al.*, 2021; Pan *et al.*, 2022). These

studies provided molecular evidence that the overexpression of promiscuous enzyme at

least partly accommodates MHR that is often accompanied in weeds with enhanced

herbicide metabolism.

While the identification of the responsible genes advanced the understanding of

metabolic resistance in weeds, MHR in the *E. phyllopogon* population is not fully

understood, among which is the high-level resistance to ACCase inhibitor

diclofop-methyl. While MHR line exhibit high-level resistance to diclofop-methyl, the

diclofop-methyl metabolizing activity of the overexpressed P450 CYP81A12/21 was not

high enough (Iwakami *et al.*, 2019). Considering that diclofop-methyl resistance is

inherited following the Mendelian segregation ratio as in other herbicides (Iwakami *et al.*,

2019), we reasoned that an additional element specific to diclofop-methyl metabolism

underlies the high-level diclofop-methyl resistance. Here, we report that the

overexpression of a novel diclofop-methyl metabolizing P450, CYP709C69, is involved

in high-level resistance to diclofop-methyl. In contrast to the catalytic promiscuity

observed in CYP81As, CYP709C69 exhibited little metabolic activity to other herbicides

except that it may be involved in the activation of a DXS inhibitor clomazone. This study

highlights that genetically-linked simultaneous overexpression of herbicide-metabolizing

P450s with different enzymatic properties is involved in MHR in weeds. The concept is

distinct from the accumulations of multiple genes for broad or high-level resistance and

serves as a model for future studies of metabolic resistance in weeds.

Materials and Methods

Origin of *E. phyllopogon*

Sensitive (S) and MHR lines (i.e. 401 and 511, respectively) originated from the

Sacramento Valley in California (Fischer *et al.*, 2000; Tsuji *et al.*, 2003) were used. The

F6 progeny of the two lines was derived from a single seed descent method (Iwakami *et*

al., 2014a). The herbicide responses of F6 lines were evaluated previously (Iwakami *et*

al., 2014a; Guo *et al.*, 2019; Iwakami *et al.*, 2019; Chayapakdee *et al.*, 2020). Seeds of

suspected S (Eoz1804) and resistant (Eoz1814) populations were collected at the paddy

fields in Niigata prefecture in 2017. A plant of each population was cultivated and

subjected to self-pollination, which was repeated twice before use. The full-length

sequences of ALS genes (two copies) and carboxytransferase domain of ACCase genes

(four copies) were amplified in a copy-specific way and directly sequenced as previously

described (Iwakami *et al.*, 2012).

Analysis of diclofop metabolites in plants

E. phyllopogon were cultured hydroponically to around 2.5 leaf-stage at 25°C with a 12-h

photoperiod (approximately 300 μmol $m^{-2}s^{-1}$) as described previously (Iwakami *et al.*,

2019). The shoots of hydroponically cultured *E. phyllopogon* (2.5 leaf stage) were dipped

in 30 μM diclofop-methyl solution supplemented with Tween 20 (0.01%) for 30 min. The

shoots of 10 plants (approximately 300 mg) were collected at 0, 1, 3, and 6 h after

application. On collection, they were rinsed in distilled water containing 20% methanol

and 0.2% Triton X-100, and snap-frozen in liquid nitrogen. The plant tissue was ground

into powder in liquid nitrogen with a mortar and pestle and extracted with 6 ml of 80%

cold methanol, followed by centrifugation at 9,000 g for 10 min at 4°C. The pellet was

further extracted twice with 2 ml of 80% cold methanol. The extract was combined,

evaporated to dryness, and dissolved in 1 ml of 0.1 M sodium acetate buffer (pH 5.0).

The solution was treated with 0.3 mg of almond β-glucosidase (BGH-101, Toyobo,

Osaka, Japan) and incubated for 24 h at 37°C. After the addition of 1 ml of acetonitrile,

the solution was vortexed, left for 15 min at 4°C, and then centrifuged at 1,700 g for 15

min. The supernatant was collected and evaporated to remove acetonitrile. The sample

was loaded onto a Sep-Pak C18 cartridge (Waters, Tokyo, Japan) and washed with 10

ml of water containing 0.1% formic acid, followed by elution with 4 ml of acetonitrile. The

fraction eluted with acetonitrile was evaporated to dryness and redissolved in acetonitrile

(0.5 ml/300 mg of fresh weight of plant tissue), which was subjected to mass

spectrometric analysis.

Mass spectrometric analysis

Hydroxylated metabolites of diclofop-acid were analyzed using a liquid

chromatography-tandem mass spectrometer (Shimadzu LCMS-8030, Kyoto, Japan)

equipped with an electrospray ionization source. The MS parameters used were as

follows: interface voltage of 4.5 kV, desolvation line temperature of 250°C, heat block

temperature of 400°C, nebulizing gas (N_2) of 2.0 l min^{-1}, and drying gas at 15 l min^{-1}.

Multiple reaction monitoring (MRM) was used to quantitate the hydroxylated metabolites

of diclofop-acid under the following conditions: negative ion mode with MRM transition of

m/z 341 to m/z 269 (CE= 15 V, Q1 Pre Bias= 15 V, Q3 Pre Bias= 19 V). Separation of

analytes was carried out using a reversed-phase column, TSK gel ODS-100V (2 mm ID

x 150 mm, 3 µm, Tosoh, Tokyo, Japan). The column was eluted with a linear gradient

from 30 to 90% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A

(0.1% formic acid in water) for 20 min at a flow rate of 0.2 ml min^{-1} at 40°C. The injection

volume was 5 µL.

RNA extraction and cDNA synthesis

Total RNA was extracted using an RNeasy Plant Mini Kit (Qiagen, Tokyo, Japan)

followed by the application of TURBO DNA-*free* kit (Thermo Fisher Scientific, Tokyo,

Japan) to eliminate genomic DNA. Total RNA (1 µg) was reverse-transcribed using

ReverTra Ace (Toyobo, Osaka, Japan) according to the manufacturer's instructions.

RNA-seq analysis

RNAs extracted from the shoot of 2.5-leaf stage plants of S and MHR lines were used for

RNA-seq analysis with four biological replications. RNA-Seq libraries were prepared with

TruSeq RNA Library Preparation Kit v2 (Illumina, USA), followed by pair-end sequencing

on HiSeq 2500 sequencer (Illumina). Reads were filtered using TRIMMOMATIC (ver

0.36) (Bolger *et al.*, 2014) with the following options: TRAILING:25, SLIDINGWINDOW:4,

MINLEN:80. The reads were mapped against coding sequences of the draft genome of

E. phyllopogon (Ye *et al.*, 2020) using bowtie2 (ver 2.3.5.1) with the following options: '-X

900–very-sensitive–no-discordant–no-mixed–dpad 0–gbar 99999999'. The mapped

reads were counted using RSEM (Li & Dewey, 2011). Differential expression analysis

was performed using edgeR (ver 3.28.0) with glmQLFTest (Robinson *et al.*, 2010) in R

(R Core Team, 2016). Contigs were annotated by BLASTX (Blast+ v.2.9.0) (Camacho *et*

al., 2009) against rice protein sequences (*Oryza sativa* MSU release 7) and rice P450

protein sequences in Cytochrome P450 Homepage (https://drnelson.uthsc.edu/).

Real-time PCR

Real-time PCR was performed as described previously (Tanigaki *et al.*, 2021) using

primers listed in Table S1. Some of the primer sets were developed elsewhere

(Czechowski *et al.*, 2005; Iwakami *et al.*, 2014b; Tanigaki *et al.*, 2021). Eukaryotic

translation initiation factor 4B (*EIF4B*) was used as an internal control gene and data was

analyzed by the ΔΔCt method (Schmittgen & Livak, 2008).

Isolation of P450 genes

The full-length sequence of each P450 gene was amplified from the cDNA of *E.*

phyllopogon (line 511) using primers listed in Table S1 with KOD FX Neo (TOYOBO) or

PrimeSTAR GXL DNA Polymerase (Takara). The amplicons were subcloned using

pGEM-T Easy Vector Systems Kit (Promega, Tokyo, Japan) or Zero Blunt PCR Cloning

Kit (Thermo Fisher Scientific, Tokyo, Japan) according to the manufacturer's instructions.

Plant transformation

The coding region of each P450 gene was cloned into pCAMBIA1390 vector with the

In-Fusion DH Cloning Kit (TaKaRa, Kusatsu, Japan) or SLiCE reaction (Motohashi,

2015) using primers listed in Table S1. The binary construct was introduced into

Agrobacterium tumefaciens strain EHA105 using the freeze and thaw method (Hofgen &

Willmitzer, 1988). Rice calli (*Oryza sativa* cv. Nipponbare) were transformed as

previously described (Toki, 1997; Iwakami *et al.*, 2019). *Arabidopsis thaliana* (ecotype

Columbia-0) was transformed by floral dip method (Clough & Bent, 1998) with the

selection of hygromycin B (20 mg l^{-1}). T3 homozygous lines were selected based on the

segregation ratio.

Herbicide sensitivity assay

Independently transformed calli were placed on N6D solid media supplemented with a

herbicide as described previously (Iwakami *et al.*, 2019). The calli were cultured at 30°C

for 3 weeks. For the *A. thaliana* sensitivity assay, the surface-sterilized seeds were

placed on Murashige and Skoog solid media (Murashige & Skoog, 1962). After three

days of stratification at 4°C, the media were placed at a climate chamber with 22°C, 12-h

photoperiod, and 70 μE m^{-2} s^{-1} light intensity for 12 days. In the case of the

diclofop-methyl assay, the plates were kept in the chamber for 14 days. Chlorophyll was

extracted and measured as described previously (Endo *et al.*, 2021). The herbicides and

the doses were summarized in Table S2.

The diclofop-methyl response of *E. phyllopogon* was determined as described

previously (Iwakami *et al.*, 2019). Briefly, hydroponically cultured plants were dipped in

diclofop-methyl solution for 30 min as described above. The plants were further

cultivated for nine days. Dose-response curves were drawn with the drc package (ver

3.0) (Ritz *et al.*, 2015).

Heterologous expression of P450s in yeast and whole-cell herbicide metabolism

assay

Expression in yeast (*Saccharomyces cerevisiae*) was conducted using WAT11 strain and

pYeDP60 vector system (Pompon *et al.*, 1996). The coding regions of other P450s were

cloned into the pYeDP60 vector with the yeast Kozak sequence inserted as described

before (Iwakami *et al.*, 2019). WAT11 was transformed using the lithium acetate method

(Ito *et al.*, 1983). A whole-cell assay for diclofop-methyl metabolism was performed

according to Iwakami *et al.* (2019) with the modification of diclofop-methyl concentration

to 300 µM. After incubation for 24 h, yeast cells were centrifuged at 13,500 *g* for 10 min.

The supernatant was cleaned up using a Sep-Pak C18 cartridge as described above and

subjected to liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis.

Structure determination of diclofop metabolites

The yeast expressing *CYP81A12* or *CYP709C69* in whole-cell assay buffer (350 ml) was

incubated with diclofop-methyl (300 µM) for 24 h. The culture was centrifuged at 1,500 *g*

for 5 min to collect the supernatant. The yeast cells were resuspended in 1/10 volume of

the buffer without diclofop-methyl. After incubation for an additional 12 h, the supernatant

was collected by centrifugation, which was repeated twice. The whole procedure was

repeated 30 times. After combining all the supernatants, the pH of the solution was

adjusted to 1 by hydrochloric acid. The solution was extracted four times with diethyl

ether (0.7 volume). After evaporating diethyl ether *in vacuo*, the resultant residue was

dissolved in acetonitrile. High-performance liquid chromatography (HPLC) (LC-10ADvp

and SPD-M10Avp, Shimadzu) separation was carried out on a reversed-phase column

(TSK gel ODS-100, 4.6 mm ID x 250 mm, 5 µm, Tosoh, Tokyo, Japan) with the following

conditions: linear gradient from 30 to 72% mobile phase B (0.1% formic acid in

acetonitrile) in mobile phase A (0.1% formic acid in water) for 35 min at a flow rate of 0.8

ml/min at 40°C. The elution was monitored by UV absorbance at 215 and 280 nm. The

fractions containing each hydroxylated metabolite of diclofop-acid were evaporated to

dryness and dissolved in deuterated chloroform. ^{1}H-nuclear magnetic resonance (NMR)

spectra of each metabolite were recorded on a Bruker Ascend 400 MHz spectrometer

(Billerica, MA, USA) with tetramethylsilane as an internal standard.

Phylogenetic analysis of *CYP81A* and *CYP709C*

Nucleotide sequences of CYP709C subfamilies of rice were obtained from *Oryza sativa*

MSU release 7 (Ouyang *et al.*, 2007), and RAP-DB (Sakai *et al.*, 2013). Deduced protein

sequences were aligned with MAFFT (ver 7.475) (Katoh & Standley, 2013). The

alignments were used to estimate evolutionary topology in MEGA X (Kumar *et al.*, 2018)

using Neighbor-Joining method. JTT matrix-based method was used to compute the

evolutionary distances. Bootstrap values (1,000 replicates) are shown next to the

branches.

Synteny analysis

The genome of *E. haploclada* (Ye *et al.*, 2020) was used for the synteny analysis. The

local synteny around the *CYP81A*s and *CYP709C*s was visualized with clinker (ver

0.0.21) (Gilchrist & Chooi, 2021).

Results

Two distinct hydroxylated metabolites of diclofop-acid are highly accumulated in

MHR *E. phyllopogon*

To gain clues for the high-level resistance to diclofop-methyl in the MHR line, we

investigated the metabolism of diclofop-methyl in *E. phyllopogon*. Diclofop-methyl is a

pro-herbicide that is activated by endogenous esterase as is the case for most

aryloxyphenoxy-propanoate herbicides (Wenger *et al.*, 2012). Following the

de-esterification, the resultant active form diclofop-acid is inactivated by P450-mediated

hydroxylation followed by further glucosylation (Fig. 1a) (Shimabukuro *et al.*, 1979, 1987;

McFadden *et al.*, 1989). In the metabolism scheme, diclofop-acid hydroxylation is the

rate-limiting step for diclofop-methyl inactivation in wheat, which is also suggested in

other plants (Owen, 2000). Our previous study showed that both the MHR and S lines

converted the majority of diclofop-methyl into diclofop-acid by 3 h after treatment for 30

min (Iwakami *et al.*, 2019). In this study, we further analyzed the dynamics of putative OH

metabolites of diclofop-acid in the extract of plants prepared as before (Fig. 1b). Two

peaks with retention times of 11.3 and 12.4 min (M1 and M2, respectively), which

correspond to putative OH metabolites, were detected by LC-MS/MS analysis using a

MRM mode (Fig. 1c). The signal intensity of M2 was higher than that of M1 under this

condition. In line with our study, a variety of hydroxylated diclofop-acid metabolites were

reported in wheat (Tanaka *et al.*, 1990; Zimmerlin & Durst, 1992), suggesting the

presence of at least partially overlapping metabolic pathway for diclofop-methyl between

wheat and *E. phyllopogon*.

We next compared the amount of each hydroxylated metabolite between the MHR

and S lines. To accurately evaluate the hydroxylation activity, possible glucosylated

metabolites were converted into hydroxylated diclofop-acid by adding β-glucosidase to

the extract of the diclofop-methyl treated plants (Fig. 1a), which was then subjected to

comparative quantification between the lines. As a result, the amounts of both

hydroxylated metabolites were higher in MHR plants at 0 and 1 HAT although no

significant difference was observed at 3 and 6 HAT (Fig. 1d). Diclfop-methyl follows

complexed metabolic pathways after glucosylation in plants (Shimabukuro *et al.*, 1987;

Han *et al.*, 2013), which should have masked the difference of these metabolites

between the lines later than 3 HAT.

To investigate whether the metabolites are produced by CYP81A12 and

CYP81A21, the P450s responsible for the resistance to the majority of herbicides in the

MHR line (Dimaano *et al.*, 2020), we analyzed the whole cell diclofop-methyl metabolism

of yeast carrying *CYP81A12/21*. In the yeast system, another peak (M3), which was not

clearly separated from M1, was observed in addition to M1 and M2. Notably, the

chromatographic profile of M1 and M2 differed between *E. phyllopogon* and yeast: the

intensity of M2 was higher in *E. phyllopogon*, while that of M1 was prominently higher in

yeast (Fig. 1e). The result suggests that an additional enzyme that catalyzes

hydroxylation to form M2 is involved in diclofop-methyl resistance in the MHR line.

Candidate genes for high-level diclofop-methyl resistance were identified by gene

expression study

To identify the additional diclofop-hydroxylating enzyme, we performed RNA-seq

analysis using the recently released draft genome of *E. phyllopogon* with 66,521 gene

models (Ye *et al.*, 2020). Among the gene models, we identified 549 putative P450 genes.

A transcriptome comparison detected some P450 genes highly expressed in the MHR

line. A closer look at the sequences of these P450 genes found that at least two of them

(Contig817_pilon.380 and Contig 222_pilon.21) were misannotated due to the existence

of similar genes next to each other (Fig. S1). Thus, we replaced the respective coding

sequences with two correct sequences and mapped the reads to the revised

transcriptome.

Finally, we detected 463 genes that were highly expressed in the MHR line (Fig.

2a, b, Fig. S2). Among the genes, 18 P450 genes were detected including previously

reported multiple-herbicide metabolizing *CYP81A12* and *CYP81A21* taking the top four

and five positions, respectively, among all the P450 genes in the MHR line (Fig. 2a, c).

Four of them respectively encode a truncated protein: Contig168_pilon.233

(*CYP81A25P*), Contig1257_pilon.115 (*CYP81A19P*), Contig222_pilon21.b (*CYP704A*),

and Contig774_pilon.23 (*CYP71E*); thus, they were excluded from further analyses.

The candidate overexpressed genes were further screened by real-time PCR for

their co-segregation with MHR using the F6 recombinant inbred lines from a cross

between the MHR and S lines (Fig. 2d). The analysis revealed that *CYP72A122*,

CYP72A252v1, *CYP72A252v2*, and *CYP709C69* co-segregated with diclofop-methyl

resistance (Fig. 2e), suggesting that they are the prime candidates for metabolite

M2-producing P450(s). Thus, we subjected the four genes to subsequent functional

characterization. We also tested the function of *CYP92A198* and *CYP78C* as we failed to

evaluate them by real-time PCR due to failures in designing appropriate primers.

331 Identification of novel diclofop-methyl metabolizing P450

For the evaluation of the diclofop-metabolizing activity of each gene, we used the rice

transformation system as employed previously (Iwakami *et al.*, 2019). We expressed the

six candidate genes in rice calli with the pCAMBIA1390 vector, where the transgene was

placed under the control of Cauliflower mosaic virus 35S promoter. The calli were tested

for growth on the media supplemented with diclofop-methyl. Among the six genes, only

CYP709C69 conferred marked resistance at the lethal dose of diclofop-methyl (0.4 µM)

to rice calli (Fig. S3a). No difference was observed in the coding sequence of

CYP709C69 between the S and MHR lines. When 12 independent transformed lines

were subjected to diclofop-methyl, all the lines with *CYP709C69* stopped growing at 6

µM (Fig. 3a). The diclofop-methyl resistance level was equivalent to the cases of

CYP81A12 and *CYP81A21*. Decrease in diclofop-methyl sensitivity was also observed in

a *A. thaliana* line transformed with *CYP709C69* as well as those with *CYP81A12* and

CYP81A21 (Fig. S4).

We then transformed *CYP709C69* in yeast and investigated whole cell

diclofop-methyl metabolism as performed previously (Iwakami *et al.*, 2019). LC-MS/MS

analysis revealed that M2 was produced as the main peak (Fig. 3b). To determine the

hydroxylation position in the OH-metabolites of diclofop-acid, we purified M1 and M2

from yeast culture applied with diclofop-methyl using HPLC, which were then subjected

to NMR analysis. Since the NMR structural analysis of three OH-metabolites of

diclofop-methyl has been reported previously (Tanaka *et al.*, 1990), we compared the

chemical shift patterns in the aromatic regions of M1 and M2 with the reported data. The

results showed that M1 has a similar pattern with the metabolite containing a

2,5-dichloro-4-hydroxyphenoxy   moiety   and   M2   with   that   containing   a

2,4-dichloro-5-hydroxyphenoxy moiety (Fig. S5). The position of one of the chlorine

atoms in M1 is different from the original position, which is thought to be formed by

intramolecular migration during enzymatic hydroxylation of the aromatic ring (so-called

NIH-shift) as previously reported (Fig. 3c) (Tanaka *et al.*, 1990). This implies that

CYP81A12 and CYP81A21 mediate the NIH-shift reaction in diclofop metabolism. In this

study, we could not determine the structure of M3 due to the limited amount available

from CYP81A expressing yeast (Fig. 1e, 3b). Considering that wheat produces the

hydroxylated diclofop with a 2,3-dichloro-4-hydroxyphenoxy moiety (Tanaka *et al.*, 1990)

in addition to the metabolites shown above, M3 may carry this moiety (Fig. 3c) although

further experiments are required.

Diclofop-acid metabolizing activity is partially conserved among CYP709C P450s

CYP709C was previously characterized in wheat, where the isolated CYP709C1

harbors sub-terminal hydroxylation activity of C18 fatty acids (Kandel *et al.*, 2005) (Fig.

4a). Importantly, the *in vitro* microsome assay purified from yeast showed that

CYP709C1 is inactive to diclofop-acid, raising a possibility that the observed

diclofop-acid metabolizing activity in CYP709C69 may be specific to this P450. To

investigate the functional conservation, we further tested other CYP709Cs in *E.*

phyllopogon although no association with resistance was observed in the gene

expression analyses (Fig. 2e). Since *E. phyllopogon* is an alletetraploid, a majority of

genes have a close relative, as observed for CYP709C68 and Contig960_pilon.355,

CYP709C70 and CYP709C71, CYP709C67 and Contig1380_pilon.11. However, the

counterpart for CYP709C69 was missing in the genome (Fig. 4a). We first tested

CYP709C68, one of the highly similar CYP709Cs to CYP709C69. The yeast expressing

CYP709C68 produced a small amount of M2 metabolite (Fig. 4b). The rice calli

expressing *CYP709C68* exhibited no prominent difference in diclofop-methyl sensitivity

(Fig. S3b), which may indicate its low activity against diclofop-methyl. Further study is

required to determine its relative activity to CYP709C69 since the expression levels were

not determined both in the yeast and rice calli. Four more genes, i.e., *CYP709C67,*

CYP709C68, CYP709C70, CYP709C71, and Contig960_pilon.355 were also tested in

rice calli system. However, the growth of the calli on diclofop-methyl media did not

markedly differ from *GFP* expressing calli (Fig. S3b). The genome holds six more

putative *CYP709C* genes (Contig1380_pilon.11, Contig12_pilon.1, Contig291_pilon.387,

Contig1182_pilon.1, Contig291_pilon.396, and Contig120_pilon.313). However, the

annotated protein sequences other than Contig1380_pilon.11 and Contig291_pilon.387

were too short (<331) for a canonical P450 sequence.

We further evaluated the function of the *CYP709C* gene in rice. Seven genes were

annotated as *CYP709C* in rice (Fig. 4a). Of these, CYP709C5 and CYP709C9, which

are relatively similar to CYP709C69, were tested in the yeast expression system. Both

enzymes produced significant amount of M2 metabolite (Fig. 4b) although the activity

cannot be directly compared with CYP709C69. Taken together, these results suggest

that the ability to metabolize diclofop is at least partially conserved in CYP709C

members.

CYP709C69 has a narrow substrate specificity

To investigate the role of CYP709C69 in the metabolism of other herbicides, we tested

the herbicide sensitivity of *CYP709C69*-expressing *A. thaliana*. We also used

CYP709C69-expressing rice calli for the evaluation of its metabolizing activity to ACCase

inhibitors. Among a total of 46 herbicides from at least 15 modes of action, a DXS

inhibitor clomazone was the only herbicide where a marked alteration in the sensitivity of

the transgenic plants was observed (Fig. S6). The *A. thaliana* lines expressing

CYP709C69 became more vulnerable to clomazone in inverse proportion to the

expression level (Fig. 5a, b), in contrast to the *A. thaliana* transformed with *CYP81A*s

(Guo *et al.*, 2019). Clomazone is a pro-herbicide where P450 was proposed as the

enzyme to convert 5-OH clomazone, and to further convert to the active form 5-keto

clomazone (Fig. 5c) (Nandula *et al.*, 2019). Thus, CYP709C69 may be involved in either

of the sequential bio-activation reaction.

Distantly located herbicide-metabolizing genes are simultaneously overexpressed

That the overexpression of the three P450 genes co-segregated in the F6 progenies

suggest that a single causal element underlies the regulation. To further examine the

possibility, we analyzed an additional 16 F6 lines, where the linkage of overexpression of

CYP81A12/21 with MHR was reported (Chayapakdee *et al.*, 2020). In these additional

lines, the overexpression of *CYP709C69* also linked with MHR (Fig. S7), supporting the

simultaneous regulation of the three genes in the MHR line.

To gain insights into the mechanism of overexpression of the three

herbicide-metabolizing P450 genes (*CYP81A12*, *CYP81A21*, and *CYP709C69*), we

estimated the genomic loci of the three genes. Since the draft genome of *E. phyllopogon*

is contig-based, the physical relationships of the three P450 genes in the chromosome

are unclear. Thus, we allocated the loci of the three P450 genes in the

chromosome-scale genome of diploid *Echinochloa* sp. (*E. haploclada*) (Ye *et al.*, 2020).

The *CYP81A* and *CYP709C* loci respectively correspond to chromosomes 1 and 3 in the

E. haploclada genome (Fig. 6a). The synteny around the *CYP81A* regions (Contig168

and Contig1257) and *CYP709C* regions (Contig817 and Contig960) is highly conserved

in the *E. haploclada* genome (Fig. 6b), implying that the contigs are fragments of

homeologous chromosomes. Importantly, the synteny structure shows that *CYP81A12*

and *CYP81A21* represent homeologous relationship as hinted previously from their

phylogenetic relationships and herbicide-metabolizing functions (Iwakami *et al.*, 2014a;

Dimaano *et al.*, 2020). A homeologous gene for *CYP709C69* was not identified on

Contig960, which is consistent with the phylogenetic analysis (Fig. 4a). Considering that

the MHR including diclofop-methyl resistance is inherited as a Mendelian trait (Iwakami

et al., 2014a, 2019) and that the overexpression of the three genes is genetically linked

(Fig. 2, Fig. S7), a single genetic element is suggested to be behind the MHR of the

Californian population of *E. phyllopogon*. The causal element may directly or indirectly

regulate the expression of the P450s (Fig. 6c).

Overexpression of the three P450 genes was observed in an MHR population

found in Japan

Finally, we analyzed the generality of the overexpression of the three genes in another

MHR population. Recently, a resistant population (Eoz1814) was identified from a paddy

field in Niigata prefecture, Japan, the first case of herbicide resistant *E. phyllopogon* in

Japan (Fig. 7a). Greenhouse experiments with commercial formulations of herbicides

revealed that the resistant line purified from the population shows the decreased

sensitivity to an ACCase inhibitor (cyhalofop-butyl), ALS inhibitors (pyrimisulfan,

propyrisulfuron, and pinoxaden) and a synthetic auxin (quinclorac) (Fig. S8) as in the

case of the MHR population found in California (Ruiz-Santaella *et al.*, 2006;

Chayapakdee *et al.*, 2020; Dimaano *et al.*, 2020). We further investigated the

diclofop-methyl sensitivity of the Japanese MHR line together with an S line from Japan

and MHR and S lines from California. In this assay, the Japanese MHR showed similar

resistance level with the Californian MHR line (Fig. 7b). Analyses of the genes encoding

herbicide target-sites, ACCase (*ACCase1* to *ACCase4*) and ALS (*ALS1* and *ALS2*), in

Eoz1814 found no mutations involved in TSR mechanisms. These observations suggest

that the geographically distinct two MHR populations share the resistance mechanism.

We then analyzed the transcription of *CYP81A12*, *CYP81A21*, and *CYP706C69*

by real-time PCR. The MHR line from Japan also exhibited significantly higher mRNA

levels although they are less pronounced compared to the MHR line from California (Fig.

7c). The results suggest that the activation of the three P450 genes underlies the MHR in

the Japanese population of *E. phyllopogon*. Meanwhile, similar diclofop-methyl

resistance levels in the two MHR lines despite the difference in mRNA levels may imply

the presence of additional mechanism(s) in the Japanese line.

Discussion

Previous studies have uncovered that *E. phyllopogon* carries enzymes with broad

substrate specificity, namely CYP81A12 and CYP81A21, that cause resistance to many

herbicides ranging from multiple modes of action when overexpressed (Iwakami *et al.*,

2019). On the other hand, the MHR *E. phyllopogon* from California exhibits resistance

even to some herbicides to which the CYP81As have little or no activity, making the

overexpression of CYP81As not the prime explanation to the expression of MHR trait

(Dimaano *et al.*, 2020, 2022). In this study, by focusing on the discrepancy between the

resistance level of diclofop-methyl in the Californian MHR line and the metabolic activity

of CYP81A12/21 to diclofop-methyl, we revealed that the overexpression of a novel

P450 gene, *CYP709C69*, was genetically linked with MHR in all the 32 crossed progeny,

where the overexpression of *CYP81A12* and *CYP81A21* has also been observed.

CYP709C69 carries diclofop-acid metabolizing activity similar to CYP81A12/21 although

the positions primarily hydroxylated differ between CYP709C69 and CYP81A12/21. Both

hydroxylated metabolites accumulated more in the MHR line than the S line, which is

most likely caused by the overexpressed CYP709C69 and CYP81A12/21. Thus, the

cooperative action of the three P450s on the active form of the herbicide underlies the

high-level resistance to diclofop-methyl resistance in the MHR line.

Unlike the broad substrate specificity of CYP81A12/21, CYP709C69 only

metabolized clomazone among the 46 herbicides that were further tested: *A. thaliana*

expressing this gene exhibited higher sensitivity to clomazone. Regardless of the high

substrate specificity, clomazone metabolism in particular needs to be further investigated.

Firstly, the study will accumulate molecular insight about plant genes for herbicide

activation, where only a single gene to activate ACCase herbicides was reported

(Cummins & Edwards, 2004). Secondly, the role of herbicide-activating enzymes is

important in the context of herbicide-resistance evolution. Reduced herbicide-activation

is associated with triallate resistance in *Avena fatua* found in Montana (Kern *et al.*, 1996),

which can be caused by the disruption of a gene with similar function of CYP709C69. It

is interesting to investigate whether clomazone resistance can be evolved by knocking

out *CYP709C69*. Lastly, the increased clomazone sensitivity in *A. thaliana* implies that

resistant   weeds   with   enhanced   herbicide   metabolism   may   exhibit   negative

cross-resistance to some herbicides depending on the properties of the causal enzymes.

The insight further highlights the importance of the findings. However, it is of note that the

MHR line from California is resistant to clomazone (Yasuor *et al.*, 2008) despite the

overexpression of CYP709C69. This is likely explained by the clomazone metabolizing

activity of CYP81A12/21. Our previous studies showed that *A. thaliana* expressing

CYP81A12 or *CYP81A21* became resistant to clomazone (Guo *et al.*, 2019) and that

they oxidize clomazone to an unidentified metabolite (Dimaano *et al.*, 2020). The higher

expression of *CYP81A12/21* than *CYP709C69* in the MHR line (Fig. 2c) suggests that

inactivating forces (CYP81A12/21) are the greater than activating forces (CYP709C69).

The present study represents a major step forward toward a complete

understanding of the plant metabolism of the "model herbicide" diclofop-methyl, which is

one of the well-understood herbicides in biochemistry in planta. We disclosed the genes

producing the two of the three hydroxylated diclofop-acid metabolites found in wheat in

the early 90's (Tanaka *et al.*, 1990; Zimmerlin & Durst, 1992). CYP81As and CYP709Cs

in both *E. phyllopogon* and rice produced M1 and M2, respectively, suggesting the

conserved function in other species. The genes in wheat/barley, major crops for

diclofop-methyl usage, and in *L. rigidum*, a major weed well-known for the

metabolism-based resistance to diclofop-methyl, are worthy of investigation. In our

experiments, we observed that yeast expressing CYP81A (possibly CYP709C as well)

produced a small peak of M3, which likely corresponds to a metabolite with the

2,3-dichloro-4-hydroxyphenoxy moiety detected in wheat (Fig. 3c). The reason why M3

was not detected in *E. phyllopogon* is unclear, but this could be due to differences in the

experimental systems between yeast and *E. phyllopogon*: In the yeast whole cell assay,

the cells were treated with a 300 µM diclofop-methyl solution for 24 h, while *E.*

phyllopogon was treated with a 30 µM diclofop-methyl solution for 30 min. Thus, the

yeast system subjected much higher concentration of the substrate to the enzymes,

possibly forcing a metabolic reaction to produce M3. It is also possible that M3 is more

susceptible to further metabolization in *E. phyllopogon* than M1 and M2, reducing the

actual amount of M3 and M3-glucoside than those of M1 and M2 in *E. phyllopogon*.

Analysis of other species in addition to the characterization of the M3 metabolite will

provide insight into these possibilities.

Trough the detailed diclofop-methyl metabolism study, we disclosed that

CYP81As mediate chlorine migration reaction in the process of diclofop hydroxylation.

The hydroxylation of the aromatic ring can cause the migration of the original functional

group such as halogen, alkyl group, and carboxyl group, to the adjacent carbon, which is

called NIH-shift (Zhao *et al.*, 2018). P450s are among the major enzymes involved in

NIH-shift although the specific P450 members that cause NIH-shift are rarely reported.

2,4-D and diclofop-methyl are the two major herbicides involving NIH-shift in their

metabolism *in planta* (Dimaano & Iwakami, 2021). The biological influence of the

NIH-shift reactions in these herbicides is not well-understood. The reactions may

influence the accessibility of the enzymes governing the following reactions of the

metabolites, which may further influence plant herbicide sensitivity.

This study provides another aspect in the mechanism of MHR in agricultural

weeds: MHR can be caused by the result of genetically-linked simultaneous activation of

genes encoding herbicide-detoxifying enzymes with different enzymatic properties. The

idea differs from a mere stepwise accumulation of multiple mechanisms to broaden the

spectrum of the resistance as suggested in other species (e.g. Preston *et al.*, 1996;

Giacomini *et al.*, 2020). Similar concept is suggested in the mode of action of herbicide

safener, chemical agents that reduce the phytotoxicity of herbicides to crop plants, where

a concerted upregulation of multiple herbicide-metabolism related genes has been

proposed as the mechanism (Riechers *et al.*, 2010) although no clear genetic

investigations for the concerted upregulation has been provided. Thus, it is possible that

the safener-mediated concerted regulations are caused by the stimulations to the

multiple independent gene regulation networks, which is clearly distinct from the concept

of genetically-linked simultaneous activation of herbicide metabolizing genes provided

here. Intriguingly, we observed the co-segregations of overexpression and MHR in at

least three more P450 genes in the F6 lines, i.e., *CYP72A122*, *CYP72A252v1*, and

CYP72A252v2 (Fig. 2e). If any of the genes carry herbicide-metabolizing function, the

scope of resistance of the MHR line would be expanded.

The three P450 overexpression may correspond to a constitutive activation of

some plant defense against biotic or abiotic stress, and to the expression of one or more

natural compound pathway(s). CYP81As in maize are involved in the biosynthesis of

zealexins, antimicrobial substances (Ding *et al.*, 2020), where a cluster of genes

including *CYP81A* and other P450 genes are activated in response to pathogen

infestation, leading to zealexin biosynthesis. Given a similar metabolic regulation system

is conserved in *E. phyllopogon*, the overexpression of the three P450 genes may be a

result of constitutive overexpression of plant defense system. Elucidation of the

regulatory system and the endogenous function of each P450 may provide insight into

the fitness costs associated with resistance expression and hints for the control of MHR

plants that are hardly controlled by herbicide chemistries.

Another interesting finding is that overexpression of the three P450s was also

found in the Japanese line. It is unlikely that herbicide selection pressure acted on

CYP709C69, since diclofop and clomazone, which were the only herbicides that

CYP709C69 metabolized in our study, have never been registered in rice cropping in

Japan. Thus, it is more likely that other selection pressures, such as ALS inhibitors, have

selected for a mutation that activate the regulatory system for expression of genes such

as *CYP81A12* and *CYP81A21* that metabolize these herbicides. The fact that the

Japanese line has the same diclofop-methyl resistance trait as the MHR line from

California, even though they were selected not by diclofop-methyl, supports the idea that

the gene regulatory system proposed in the Californian line is a common endogenous

system shared by other *E. phyllopogon*. In herbicide resistance, molecular convergence

is frequently observed in the case of target-site resistance (Baucom, 2019). On the other

hand, the convergence in metabolic resistance remains poorly understood. Elucidation of

the causal mechanisms that confer resistance in these MHR *E. phyllopogon* is expected

to provide new insights in understanding the mechanisms of stress tolerance evolution in

plants.

Acknowledgements

We thank Dr. David Nelson for naming the P450s, Dr. Niña Gracel Dimaano for English

critique, and Yuuri Nakamura for technical assistance. RNA-seq was supported by the

NODAI Genome Research Center, Tokyo University of Agriculture under Grant

Cooperative Research of the Genome Research for Bio-Resource. Computations were

partially performed on the NIG supercomputer at ROIS National Institute of Genetics.

This work was supported by a JSPS KAKENHI grant no. 15H06072, 17K15234,

19H02955 (to S.I.), and in part by the Program for the Development of Next-generation

Leading Scientists with Global Insight (L-INSIGHT), sponsored by the Ministry of

Education, Culture, Sports, Science and Technology (MEXT), Japan.

Author contributions

Conceptualization, S.I.; Research design, S.I., H.S., M.M., Y.Y., T.Y., K.T., S.T., T.T.;

RNA-sequencing, K.T.; Bioinformatics, S.I., H.S., T.K.; LC-MS/MS and NMR, H.S., Y.Y.,

M.M.; Japanese line characterization, K.T., A.U., S.A., M.H.; Gene characterization, H.S.,

S.I., T.Y.; Writing, S.I., H.S., M.M.; Visualization, S.I., H.S., T.K.; Supervision, S.I., T.T.;

Funding acquisition, S.I.

Data availability

The mRNA-Seq data have been deposited in the DDBJ Sequence Read Archive (DRA)

database (accession number DRA013092).

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Baucom RS. **2019**. Evolutionary and ecological insights from herbicide-resistant weeds:

what have we learned about plant adaptation, and what is left to uncover? *The New*

phytologist **223**: 68–82.

Bolger AM, **Lohse M**, **Usadel B**. **2014**. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**: 2114–2120.

Camacho C, **Coulouris G**, **Avagyan V**, **Ma N**, **Papadopoulos J**, **Bealer K**, **Madden TL**.

2009. BLAST+: architecture and applications. *BMC Bioinformatics* **10**: 421.

Chayapakdee P. **2019**. *Molecular mechanism of quinclorac resistance in*

multiple-Herbicide resistant Echinochloa phyllopogon. PhD thesis, University of Tsukuba,

Tsukuba, Japan.

Chayapakdee P, **Sunohara Y**, **Endo M**, **Yamaguchi T**, **Fan L**, **Uchino A**, **Matsumoto**

H, **Iwakami S**. **2020**. Quinclorac resistance in *Echinochloa phyllopogon* is associated

with reduced ethylene synthesis rather than enhanced cyanide detoxification by

beta-cyanoalanine synthase. *Pest Management Science* **76**: 1195–1204.

Clough SJ, **Bent AF**. **1998**. Floral dip: a simplified method for *Agrobacterium*-mediated

transformation of *Arabidopsis thaliana*. *The Plant Journal* **16**: 735–743.

Cummins I, **Edwards R**. **2004**. Purification and cloning of an esterase from the weed

black-grass (*Alopecurus myosuroides*), which bioactivates aryloxyphenoxypropionate

herbicides. *The Plant Journal* **39**: 894–904.

Czechowski T, **Stitt M**, **Altmann T**, **Udvardi MK**, **Scheible WR**. **2005**. Genome-wide

identification and testing of superior reference genes for transcript normalization in

Arabidopsis. *Plant Physiology* **139**: 5–17.

Dimaano NG, **Iwakami S**. **2021**. Cytochrome P450-mediated herbicide metabolism in

plants: current understanding and prospects. *Pest Management Science* **77**: 22–32.

Dimaano NG, **Tominaga T**, **Iwakami S**. **2022**. Thiobencarb resistance mechanism is

distinct from CYP81A-based cross-resistance in late watergrass (*Echinochloa*

phyllopogon). *Weed Science* **70**: 160–166.

Dimaano NG, **Yamaguchi T**, **Fukunishi K**, **Tominaga T**, **Iwakami S**. **2020**. Functional

characterization of cytochrome P450 CYP81A subfamily to disclose the pattern of

cross-resistance in *Echinochloa phyllopogon*. *Plant Molecular Biology* **102**: 403–416.

Ding Y, **Weckwerth PR**, **Poretsky E**, **Murphy KM**, **Sims J**, **Saldivar E**, **Christensen**

SA, **Char SN**, **Yang B**, **Tong AD**, *et al.* **2020**. Genetic elucidation of interconnected

antibiotic pathways mediating maize innate immunity. *Nature Plants* **6**: 1375–1388.

Endo M, **Iwakami S**, **Toki S**. **2021**. Precision genome editing in plants via gene targeting

and subsequent break-induced single-strand annealing. *Plant Biotechnology Journal* **19**:

563–574.

Fischer AJ, **Ateh CM**, **Bayer DE**, **Hill JE**. **2000**. Herbicide-resistant *Echinochloa*

oryzoides and *E. phyllopogon* in California *Oryza sativa* fields. *Weed Science* **48**:

225–230.

Giacomini DA, **Patterson EL**, **Kupper A**, **Beffa R**, **Gaines TA**, **Tranel PJ**. **2020**.

Coexpression clusters and allele-specific expression in metabolism-based herbicide

resistance. *Genome Biology and Evolution* **12**: 2267–2278.

Gilchrist CLM, **Chooi Y-H**. **2021**. Clinker & clustermap.js: Automatic generation of gene

cluster comparison figures. *Bioinformatics* **37**: 2473–2475.

Guo F, **Iwakami S**, **Yamaguchi T**, **Uchino A**, **Sunohara Y**, **Matsumoto H**. **2019**. Role

of CYP81A cytochrome P450s in clomazone metabolism in *Echinochloa phyllopogon*.

Plant Science **283**: 321–328.

Han H, **Yu Q**, **Beffa R**, **Gonzalez S**, **Maiwald F**, **Wang J**, **Powles SB**. **2021**.

Cytochrome P450 CYP81A10v7 in *Lolium rigidum* confers metabolic resistance to

herbicides across at least five modes of action. *The Plant journal* **105**: 79–92.

Han H, **Yu Q**, **Cawthray GR**, **Powles SB**. **2013**. Enhanced herbicide metabolism

induced by 2,4-D in herbicide susceptible *Lolium rigidum* provides protection against

diclofop-methyl. *Pest Management Science* **69**: 996–1000.

Hofgen R, **Willmitzer L**. **1988**. Storage of competent cells for *Agrobacterium*

transformation. Nucleic acids research **16**: 9877.

Ito H, **Fukuda Y**, **Murata K**, **Kimura A**. **1983**. Transformation of intact yeast cells treated

with alkali cations. *Journal of Bacteriology* **153**: 163–168.

Iwakami S, **Endo M**, **Saika H**, **Okuno J**, **Nakamura N**, **Yokoyama M**, **Watanabe H**,

Toki S, **Uchino A**, **Inamura T**. **2014a**. Cytochrome P450 CYP81A12 and CYP81A21 are

associated with resistance to two acetolactate synthase inhibitors in *Echinochloa*

phyllopogon. Plant Physiology **165**: 618–629.

Iwakami S, **Kamidate Y**, **Yamaguchi T**, **Ishizaka M**, **Endo M**, **Suda H**, **Nagai K**,

Sunohara Y, **Toki S**, **Uchino A**, *et al.* **2019**. CYP81A P450s are involved in concomitant

cross-resistance to acetolactate synthase and acetyl-CoA carboxylase herbicides in

Echinochloa phyllopogon. New Phytologist **221**: 2112–2122.

Iwakami S, **Uchino A**, **Kataoka Y**, **Shibaike H**, **Watanabe H**, **Inamura T**. **2014b**.

Cytochrome P450 genes induced by bispyribac-sodium treatment in a multiple-herbicide

resistant biotype of *Echinochloa phyllopogon. Pest Management Science* **70**: 549–558.

Iwakami S, **Uchino A**, **Watanabe H**, **Yamasue Y**, **Inamura T**. **2012**. Isolation and

expression of genes for acetolactate synthase and acetyl-CoA carboxylase in

Echinochloa phyllopogon, a polyploid weed species. *Pest Management Science* **68**:

1098–1106.

Kandel S, **Morant M**, **Benveniste I**, **Blee E**, **Werck-Reichhart D**, **Pinot F**. **2005**.

Cloning, functional expression, and characterization of CYP709C1, the first sub-terminal

hydroxylase of long chain fatty acid in plants. Induction by chemicals and methyl

jasmonate. *The Journal of Biological Chemistry* **280**: 35881–35889.

Katoh K, **Standley DM**. **2013**. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Molecular Biology and Evolution* **30**:

772–780.

Kern AJ, **Peterson DM**, **Miller EK**, **Colliver CC**, **Dyer WE**. **1996**. Triallate resistance in

Avena fatua L. Is due to reduced herbicide activation. *Pesticide Biochemistry and*

Physiology **56**: 163–173.

Kumar S, **Stecher G**, **Li M**, **Knyaz C**, **Tamura K**. **2018**. MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. *Molecular biology and evolution* **35**:

1547–1549.

Li B, **Dewey CN**. **2011**. RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome. *BMC Bioinformatics* **12**: 323.

McFadden JJ, **Frear DS**, **Mansager ER**. **1989**. Aryl hydroxylation of diclofop by a

cytochrome P450 dependent monooxygenase from wheat. *Pesticide Biochemistry and*

Physiology **34**: 92–100.

Motohashi K. **2015**. A simple and efficient seamless DNA cloning method using SLiCE

from *Escherichia coli* laboratory strains and its application to SLiP site-directed

mutagenesis. *BMC Biotechnology* **15**: 47.

Murashige T, **Skoog F**. **1962**. A revised medium for rapid growth and bio assays with

tobacco tissue cultures. *Physiologia Plantarum* **15**: 473–497.

Nandula VK, **Riechers DE**, **Ferhatoglu Y**, **Barrett M**, **Duke SO**, **Dayan FE**,

Goldberg-Cavalleri A, **Tétard-Jones C**, **Wortley DJ**, **Onkokesung N**, *et al.* **2019**.

Herbicide metabolism: crop selectivity, bioactivation, weed resistance, and regulation.

Weed Science **67**: 149–175.

Ouyang S, **Zhu W**, **Hamilton J**, **Lin H**, **Campbell M**, **Childs K**, **Thibaud-Nissen F**,

Malek RL, **Lee Y**, **Zheng L**. **2007**. The TIGR rice genome annotation resource:

improvements and new features. *Nucleic acids research* **35**: D883–D887.

Owen WJ. **2000**. Herbicide metabolism as a basis for selectivity. In: Roberts T, ed.

Metabolsim of Agrachemicals in Plants. Chichester: John Wiley & Sons Ltd., 211–258.

Pan L, **Guo Q**, **Wang J**, **Shi L**, **Yang X**, **Zhou Y**, **Yu Q**, **Bai L**. **2022**. CYP81A68 confers

metabolic resistance to ALS and ACCase-inhibiting herbicides and its epigenetic

regulation in *Echinochloa crus-galli*. *Journal of Hazardous Materials* **428**: 128225.

Pompon D, **Louerat B**, **Bronine A**, **Urban P**. **1996**. Yeast expression of animal and

plant P450s in optimized redox environments. *Methods in Enzymology* **272**: 51–64.

Powles SB, **Yu Q**. **2010**. Evolution in action: plants resistant to herbicides. *Annual*

Review of Plant Biology **61**: 317–347.

Preston C, **Tardif FJ**, **Christopher JT**, **Powles SB**. **1996**. Multiple resistance to

dissimilar herbicide chemistries in a biotype of *Lolium rigidum* due to enhanced activity of

several herbicide degrading enzymes. *Pesticide Biochemistry and Physiology* **54**:

123–134.

R Core Team,. **2016**. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Riechers DE, **Kreuz K**, **Zhang Q**. **2010**. Detoxification without intoxication: herbicide
safeners activate plant defense gene expression. *Plant physiology* **153**: 3–13.

Ritz C, **Baty F**, **Streibig JC**, **Gerhard D**. **2015**. Dose-response analysis using R. *PloS*
one **10**: e0146021.

Robinson MD, **McCarthy DJ**, **Smyth GK**. **2010**. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**:
139–140.

Ruiz-Santaella JP, **De Prado R**, **Wagner J**, **Fischer AJ**, **Gerhards R**. **2006**. Resistance
mechanisms to cyhalofop-butyl in a biotype of *Echinochloa phyllopogon* (Stapf) Koss.
from California. *Journal of plant diseases and protection: scientific journal of the German*
Phytomedical Society **XX**: 95–100.

Sakai H, **Lee SS**, **Tanaka T**, **Numa H**, **Kim J**, **Kawahara Y**, **Wakimoto H**, **Yang C-C**,
Iwamoto M, **Abe T**, *et al.* **2013**. Rice Annotation Project Database (RAP-DB): an
integrative and interactive database for rice genomics. *Plant & Cell Physiology* **54**: e6.

Schmittgen TD, **Livak KJ**. **2008**. Analyzing real-time PCR data by the comparative C(T)
method. *Nature Protocols* **3**: 1101–1108.

Shimabukuro RH, **Walsh WC**, **Hoerauf RA**. **1979**. Metabolism and selectivity of
diclofop-methyl in wild oat and wheat. *Journal of Agricultural and Food Chemistry* **27**:
615–623.

Shimabukuro RH, **Walsh WC**, **Jacobson A**. **1987**. Aryl-*O*-glucoside of diclofop: a
detoxication product in wheat shoots and wild oat cell suspension culture. *Journal of*
Agricultural and Food Chemistry **35**: 393–397.

Tanaka FS, **Hoffer BL**, **Shimabukuro RH**, **Wien RG**, **Walsh WC**. **1990**. Identification of
the isomeric hydroxylated metabolites of methyl 2-[4-(2, 4-dichlorophenoxy) phenoxy]
propanoate (diclofop-methyl) in wheat. *Journal of Agricultural and Food Chemistry* **38**:
559–565.

Tanigaki S, **Uchino A**, **Okawa S**, **Miura C**, **Hamamura K**, **Matsuo M**, **Yoshino N**, **Ueno**
N, **Toyama Y**, **Fukumi N**, *et al.* **2021**. Gene expression shapes the patterns of parallel
evolution of herbicide resistance in the agricultural weed *Monochoria vaginalis*. *New*
Phytologist **232**: 928–940.

Toki S. **1997**. Rapid and efficient *Agrobacterium*-mediated transformation in rice. *Plant*
Molecular Biology Reporter **15**: 16–21.

Tsuji R, **Fischer AJ**, **Yoshino M**, **Roel A**, **Hill JE**, **Yamasue Y**. **2003**. Herbicide-resistant
late watergrass (*Echinochloa phyllopogon*): similarity in morphological and amplified
fragment length polymorphism traits. *Weed Science* **51**: 740–747.

Wenger J, **Niderman T**, **Mathew C**. **2012**. Acetyl-CoA carboxylase inhibitors. In: Krämer
W, Schirmer U, Jeschke P, Matthias W, eds. Modern Crop Protection Compounds,
Second, Revised and Enlarged Edition. Weinheim, Germany: Wiley-VCH, 447–477.

Yamasue Y. **2001**. Strategy of *Echinochloa oryzicola* Vasing. for survival in flooded rice.
Weed Biology and Management **1**: 28–36.

Yasuor H, **Milan M**, **Eckert JW**, **Fischer AJ**. **2012**. Quinclorac resistance: a concerted

hormonal and enzymatic effort in *Echinochloa phyllopogon*. *Pest Management Science*

68: 108–115.

Yasuor H, **Tenbrook PL**, **Tjeerdema RS**, **Fischer AJ**. **2008**. Responses to clomazone

and 5-ketoclomazone by *Echinochloa phyllopogon* resistant to multiple herbicides in

Californian rice fields. *Pest Management Science* **64**: 1031–1039.

Ye CY, **Wu D**, **Mao L**, **Jia L**, **Qiu J**, **Lao S**, **Chen M**, **Jiang B**, **Tang W**, **Peng Q**, *et al.*

2020. The genomes of the allohexaploid *Echinochloa crus-galli* and its progenitors

provide insights into polyploidization-driven adaptation. *Molecular Plant* **13**: 1298–1310.

Yu Q, **Powles S**. **2014**. Metabolism-based herbicide resistance and cross-resistance in

crop weeds: a threat to herbicide sustainability and global crop production. *Plant*

Physiology **166**: 1106–1118.

Zhao H, **Xu Y**, **Lin S**, **Spain JC**, **Zhou N-Y**. **2018**. The molecular basis for the

intramolecular migration (NIH shift) of the carboxyl group during para-hydroxybenzoate

catabolism. *Molecular Microbiology* **110**: 411–424.

Zimmerlin A, **Durst F**. **1992**. Aryl hydroxylation of the herbicide diclofop by a wheat

cytochrome p-450 monooxygenase: substrate specificity and physiological activity. *Plant*

Physiology **100**: 874–881.

Figure legends

Figure 1. Characterization of diclofop-acid metabolism in multiple-herbicide resistant (MHR) *Echinochloa phyllopogon*

(a) Diclofop-methyl metabolism in plants. The enzymes estimated to be involved in each metabolism steps are shown. (b) Schematic representation of sample preparation for diclofop-methyl metabolism study. The shoots were dipped in diclofop-methyl solution (30 µM) for 30 min, which were collected and stored at 0, 1, 3, or 6 h after treatment (HAT). (c) Detection of OH-diclofop-acid in the MHR line of *E. phyllopogon* on LC-MS/MS. The extract of 3 HAT was analyzed. (d) Comparison of M1 and M2 metabolite amounts between sensitive (S) and MHR lines. Diclofop-acid-*O*-Gluc was converted to OH-diclofop-acid with β-glucosidase as shown in (a). The error bars, SEM. * and ** show significant differences of $P < 0.05$ and $P < 0.01$, respectively (Student's *t*-test). (e) Whole cell diclofop metabolism assay using yeast harboring empty vector (pYeDP60), *CYP81A12,* or *CYP81A21*.

Figure 2. Candidate genes for high-level diclofop-methyl resistance in *Echinochloa phyllopogon* **were identified by gene expression study**

(a) The annotations of P450 genes that were upregulated in the MHR line. Genes are ordered according to MHR expression rank. Progeny association study was not conducted on Contig490_pilon.258 and Contig1194_pilon.509 since the appropriate primers were not designed. Highlight in gray indicates that the gene passed the tests. [a], calculated in edgeR. [b], Iwakami *et al*. (2014a). [c], Iwakami *et al*. (2019). (b) The number of differentially expressed genes. (c) Expression rank of P450 genes in the transcriptome of MHR line. P450s genes are shown in yellow and red. Red plots represent differentially

expressed P450 genes. Larger Red plots are the genes shown in grey in Fig. 2a. (d) F6

lines used in the association study. (e) Progeny association of the mRNA levels in the

shoots (four plants in bulk for each line) of the candidate P450 genes. mRNA levels were

quantified by real-time PCR in 16 F6 lines that were previously characterized (Iwakami *et*

al., 2019; Chayapakdee *et al.*, 2020). The values for *CYP71C35* in the S and some F6

lines line were set as 40 cycles since the signal was not detected. F6-S, sensitive F6

lines; F6-R, resistant F6 lines.

Figure 3. Identification of novel diclofop-methyl metabolizing P450

(a) Diclofop-methyl sensitivity of rice calli transformed with P450 genes of *Echinochloa*

phyllopogon. Twelve independent calli were grown for three weeks. (b) Whole cell

diclofop-methyl metabolism assay using yeast harboring empty vector (pYeDP60),

CYP81A12, or *CYP709C69.* (c) The role of P450s in diclofop-acid hydroxylation. The

structures of M1 and M2 were determined by nuclear magnetic resonance analysis. The

arrows indicate migration of chlorine. The structure of M3 metabolite was estimated to

have a 2,3-dichloro-4-hydroxyphenoxy moiety as reported in wheat (Tanaka *et al.*, 1990).

Figure 4. Diclofop-methyl metabolizing activity of CYP709Cs

(a) Phylogenetic tree of CYP709Cs in *Echinochloa phyllopogon* and rice, and

CYP709C1 from wheat. CYP709C1 did not metabolize diclofop-acid in in vitro assay

(Kandel *et al.*, 2005). The gene ID or accession number is shown next to CYP name

when CYP names are available. The annotation of Contig817_pilon.380.a and

Contig817_pilon.380.b is shown in Fig. S1. (b) Whole cell diclofop-methyl metabolism

assay using yeast harboring empty vector (pYeDP60), *CYP709C5, CYP709C9,*

CYP709C68, and *CYP709C69.*

Figure 5. *CYP709C69* increased clomazone sensitivity in *Arabidopsis thaliana*

(a) Clomazone (0.3 µM) response of *A. thaliana* lines transformed with *CYP709C69.* The

mRNA level of each line was evaluated in Fig. S4. (b) Association of transcript level of

CYP709C69 (n=4) and chlorophyll content in the 0.3 µM clomazone treated *A. thaliana*

lines. Linear regression line is shown in pin with shaded regions representing the 95%

confidence interval. (c) The pro-herbicide clomazone is converted to the active form

5-keto clomazone through 5-OH clomazone in plants. The involvement of P450s were

suggested (Nandula *et al.*, 2019). CYP709C69 may be involved in either of the reaction.

Figure 6. The genetic model of multiple-herbicide resistance in *Echinochloa*

phyllopogon

(a) The loci of *CYP81A12*, *CYP81A21*, and *CYP709C69* orthologs in diploid *E.*

haploclada. (b) The local synteny of *CYP81A* and *CYP709C* regions in the draft genome

of *E. phyllopogon* (2n=4X=36) and *E. haploclada* (2n=2X=18) (Ye *et al.*, 2020). (c) A

working model for multiple-herbicide resistance in *E. phyllopogon.* A causal mutation

activates the expression of at least three herbicide-metabolizing P450s through unknown

gene-regulating systems in *E. phyllopogon.* Arrow with solid line, inactivation. T-bar,

activation.

Figure 7. Overexpression of the three P450 genes was observed in a

multiple-herbicide resistant (MHR) population found in Japan

(a) Sampling sites of the five populations. Eoz1804 and Eoz1814 are sensitive (S) and

MHR, respectively. The two Californian populations (401 and 511) were used as controls.

(b) Diclofop-methyl response of the four lines. The plant appearance nine days after

diclofop-methyl treatment (100 µM) is shown. (c) Transcription of the three P450 genes

in the shoot of 2.5-leaf stage plants. Statistically significant differences evaluated using

Tukey's HSD test are indicated by different letters (n=4).

Supplementary data

Table S1. Primers used in this study

a, Primer concentration in real-time PCR. b, Developed in Tanigaki *et al.*, (2021). c, Developed in Czechowski *et al.*, (2005). d, Developed in Iwakami *et al.*, (2014b).

Table S2. The summary of herbicides used for the functional characterizations of *CYP709C69*

The herbicide classifications follow the HRAC system (https://www.hracglobal.com/). Another mode of action was annotated for quinclorac (cellulose synthesis) and bromoxynil (uncouplers).

Figure S1. Annotation correction

The original annotations for Contig817_pilon.380 (a) and Contig222_pilon.21 (b) were shown as boxes with solid lines. The missing exons in the original annotations (Ye *et al.*, 2020) were shown as boxes with dotted lines. The new gene models were distinguished with the suffix of a and b. The comparison of Contig222_pilon.21.a and Contig222_pilon.21.b suggests that the latter carries a single nucleotide deletion in the first exon at the position of an asterisk (*), leading to a premature stop codon.

Figure S2. Detection of differentially expression genes by RNA-seq

(a) Multi-dimensional scaling (MDS) plot of the RNA-seq libraries of the sensitive (S) and multiple-herbicide resistant (MHR) lines. (b) Volcano plot of the S and MHR lines. Differentially expressed genes (fold-change>3, FDR<0.05) are shown in dark gray. Upregulated P450 genes (Iwakami *et al.*, 2014a)are shown in yellow.

Figure S3. Diclofop-methyl sensitivity of P450 genes in *Echinochloa phyllopogon*

Independent calli were grown on the media with diclofop-methyl for three weeks. (a)

Candidate P450 genes. (b) CYP709C genes.

Figure S4. Diclofop-methyl sensitivity of *Arabidopsis thaliana* transformed with P450 genes in *Echinochloa phyllopogon*

(a) Transcript levels of T3 homozygous lines generated in this study. (b) Response to 30

µM diclofop-methyl. Bars represent SEM. *CYP81A21*#6-1 was previously established

line (Iwakami *et al.*, 2014a). The line exhibited the highest mRNA level among the tested

lines, and showed resistance to many herbicides (Iwakami *et al.*, 2014a; Dimaano *et al.*,

2020).

Figure S5. [1]H-NMR spectra for OH-diclofop-acid of M1 (A) and M2 (B)

Asterisks indicate the signals related to impurities.

Figure S6. Herbicide metabolizing activity of CYP709C69

(a) Rice calli responses to herbicides inhibiting acetyl-CoA carboxylase. Independent

calli were grown on the media with herbicides for three weeks. (b) *Arabidopsis thaliana*

responses to herbicides. Two highly expressing lines (#31-5 and #17-2) (Fig. S4) were

grown on the media with herbicides for 12 days. The herbicide classifications follow the

HRAC system (https://www.hracglobal.com/).

Figure S7. Progeny association of the mRNA levels of *CYP709C69*

Shoot mRNA levels (four plants in bulk) were quantified by real-time PCR in 16 F6 lines

that were previously characterized (Iwakami *et al.*, 2019; Chayapakdee *et al.*, 2020).

R6-R/S is a segregating line.

Figure S8. Herbicide sensitivity of *Echinochloa phyllopogon* in Japan

(a) Herbicides and the growth stage of *E. phyllopogon* used for the assay. 1x and 1/3x

field rate was applied. The 1x rate of quinclorac was followed by that registered in the

USA (Yasuor *et al.*, 2012). (b) Herbicide response of resistant *E. phyllopogon* (Eoz1814)

found in Japan. Shoot dry weight of 13 days after herbicide applications were evaluated.

Bars, SEM (n=5). Line 401, originated from California, was used as a sensitive control.

(a)

Diclofop-methyl

Esterase

Diclofop-acid
Active form

P450

OH-diclofop-acid

Glucosyl transferase

Diclofop-acid-*O*-Gluc

β-Glucosidase

Further metabolism

(b)

-80°C ------------- 0 HAT

2.5-leaf stage   30 min

1 hour   -80°C ------------- 1 HAT

3 hours   -80°C ------------- 3 HAT

6 hours   -80°C --- 6 HAT

(c)

MHR

M1   M2

Retention time (min)

(d)

M1

S   MHR

Peak area

Hours after treatment

M2

Peak area

Hours after treatment

(e)

CYP81A12   M1   M3   M2

CYP81A21

Empty

Retention time (min)

(a)

Gene ID	Annotation	logFC[a]	q value[a]	TPM		Rank in transcriptome		Rank in CYPome		Protein sequence	Progeny association
				S	MHR	S	MHR	S	MHR		
Contig363_pilon.90	CYP72A122	1.64	4.11E-09	65.64	210.46	2407	651	4	1		Yes
Contig168_pilon.232	CYP81A12	2.98	3.58E-10	13.88	112.18	8983	1390	42	4		Yes[b,c]
Contig1257_pilon.113	CYP81A21	4.30	9.18E-11	4.13	83.76	14348	1890	111	5		Yes[b,c]
Contig168_pilon.233	CYP81A25P	3.12	7.18E-11	6.96	62.00	12465	2556	80	7	Frame shift	Not tested
Contig1257_pilon.115	CYP81A19P	2.41	1.63E-09	10.87	59.79	10345	2656	58	8	Frame shift	Not tested
Contig363_pilon.94	CYP72A252v1	1.91	7.30E-08	13.17	50.81	9288	3100	48	11		Yes
Contig817_pilon.380.b	CYP709C69	6.83	5.00E-12	0.38	46.06	17366	3413	256	12		Yes
Contig90_pilon.215	CYP72A252v2	3.05	4.51E-10	5.29	45.46	13611	3457	98	13		Yes
Contig1299_pilon.29	CYP71C35	3.10	4.62E-11	3.08	27.21	15169	5510	126	25		No
Contig12_pilon.29	NGS709C67	1.66	7.85E-06	0.55	5.65	17207	13492	233	93		No
Contig10_pilon.131	CYP71S	3.29	1.94E-08	0.55	5.62	17207	13492	233	93		No
Contig960_pilon.352	CYP709C70	3.43	1.40E-06	0.39	4.63	17351	14119	253	107		No
Contig1257_pilon.112	CYP81A22	2.19	6.57E-06	0.92	4.47	16869	14220	204	108		No[b]
Contig222_pillon21.b	CYP704A	2.33	1.68E-05	0.65	3.51	17127	14891	222	122	Frame shift	Not tested
Contig490_pilon.258	CYP92A198	6.10	6.69E-08	0.04	3.10	17675	15206	340	128		Not tested
Contig1138_pilon.9	CYP88A	3.23	2.78E-06	0.28	2.71	17455	15504	269	136		×
Contig1194_pilon.509	CYP78C	1.71	9.21E-03	0.54	1.79	17213	16195	234	165		Not tested
Contig774_pilon.23	CYP71E	1.71	2.44E-03	0.43	1.49	17319	16447	248	177	Frame shift	No

(b) S 337 · 65,723 · MHR 463

(c) CYP72A122, CYP81A12, CYP72A252v1, CYP72A252v2, CYP709C69, CYP81A21, CYP92A198, CYP78C
Transcripts per million vs Rank in MHR line

(d) S × MHR → F1 → F2 F2 … F2 F2 → Self-pollination → F6 F6 … F6 F6 — Sensitive lines / Resistant lines

(e) CYP71C35, CYP71E, CYP71S, CYP72A122, CYP72A252v1, CYP72A252v2, CYP88A, CYP709C67, CYP709C69, CYP709C70 — Relative transcript level (S, R, F6–S, F6–R)

(a)

| | 0 μM | 0.2 μM | 2 μM | 4 μM | 6 μM |

GFP
CYP81A12
CYP81A21
CYP709C69

(b)

M1 M3
CYP81A12
CYP709C69 M2
Empty

Intensity
Retention time (min)

(c)

CYP81A12
CYP81A21

CYP709C69

Diclofop-acid

M1

M2

M3? (not identified in this study)

(a)

(b)

(a)

#15-1   #17-2

WT   #31-5

Clomazone (0.3 μM)

(b)

Chlorophyll content (ng/μl)

Relative transcript level

(c)

CYP709C69

?

Clomazone

5-OH clomazone

5-keto clomazone
Active form

(a)

(b)

(c)

(a)

- Eoz1804: S
- Eoz1814: MHR

- 401: S
- 511: MHR

(b)

401 Eoz 1804 511 Eoz 1814

(c)

CYP81A12

CYP81A21

CYP709C69