The ESKAPE mobilome contributes to the spread of antimicrobial resistance and CRISPR-

mediated conflict between mobile genetic elements

João Botelho[1,2*], Adrian Cazares[3,4], Hinrich Schulenburg[1,2]

[1]Antibiotic Resistance Evolution Group, Max Planck Institute for Evolutionary Biology, Plön,

Germany

[2]Department of Evolutionary Ecology and Genetics, Zoological Institute, Christian Albrechts

University, Kiel, Germany

[3]EMBL's European Bioinformatics Institute (EMBL-EBI), Wellcome Genome Campus, Cambridge,

United Kingdom

[4]Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge, United Kingdom

*Correspondence: botelho@evolbio.mpg.de

Abstract

Mobile genetic elements (MGEs) mediate the shuffling of genes among organisms. They

contribute to the spread of virulence and antibiotic resistance genes in human pathogens,

including the particularly problematic group of ESKAPE pathogens, such as *Enterococcus*

faecium, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*,

Pseudomonas aeruginosa, and *Enterobacter* sp. Here, we performed the first systematic analysis

of MGEs, including plasmids, prophages, and integrative and conjugative/mobilizable elements

(ICEs/IMEs), in the ESKAPE pathogens. We characterized over 1700 complete ESKAPE genomes

and found that different MGE types are asymmetrically distributed across these pathogens.

While some MGEs are capable of exchanging DNA beyond the genus (and phylum) barrier,

horizontal gene transfer (HGT) is mainly restricted by phylum or genus. We further observed

that most genes on MGEs have unknown functions and show intricate distribution patterns.

Moreover, AMR genes and anti-CRISPRs are overrepresented in the ESKAPE mobilome. Our

results also underscored species-specific trends shaping the number of MGEs, AMR, and

virulence genes across pairs of conspecific ESKAPE genomes with and without CRISPR-Cas

systems. Finally, we found that CRISPR targets vary according to MGE type: while plasmid

CRISPRs almost exclusively target other plasmids, ICEs/IME CRISPRs preferentially target

ICEs/IMEs and prophages. Overall, our study highlights the general importance of the ESKAPE

mobilome in contributing to the spread of AMR and mediating conflict among MGEs.

Introduction

Mobile genetic elements (MGEs) are DNA entities that are capable of capturing and shuffling genes intra- and intercellularly [1]. Coevolution of bacterial hosts with these MGEs has driven the evolution of complexity [2]. Movement within the genome is often mediated by specific MGEs, such as insertion sequences and transposons [3]. Others like plasmids, prophages, and integrative and conjugative/mobilizable elements (ICEs/IMEs) are key vectors for intercellular mobility, being responsible for a large fraction of the variability observed between bacterial species [4–6]. Bacteria undergo extensive horizontal gene transfer (HGT), and some estimates suggest that more than 80% of bacterial genes were laterally transferred at some point in their evolutionary history [7]. These events are largely shaped by ecological niches, by the difference in the GC content between pairs of bacteria exchanging material, and by phylogenetic barriers [8–10]. Network-based methods are useful to trace HGT events and recover shared content between bacterial genomes [10,11], and have been recently applied to explore the population structure of thousands of plasmids [12,13]. Even though this approach has been useful to explore population structure of plasmids, network analyses of other MGEs (such as prophages and ICEs/IMEs) is understudied. More importantly, only a few studies have used network-based approaches to explore the co-evolutionary dynamics of different MGE types [14,15].

MGEs carry non-essential genes that can provide their bacterial host with adaptive traits and alter their fitness, such as antimicrobial resistance (AMR) and virulence genes. These elements apply a myriad of ecological and evolutionary strategies to promote their own replication and transmission, which allow them to persist even in the absence of positive selection for the beneficial genes they carry [16,17]. As a predicted immune response against invading MGEs, bacteria have evolved different, often complex, mechanisms such as restriction-modification and clustered, regularly interspaced short palindromic repeats (CRISPR) and CRISPR-associated (Cas) genes [18]. These systems are usually clustered in 'defence islands' and are widespread in bacteria and archaea [19,20]. Spacer sequences, derived from fragments of invading MGEs (termed protospacers), are continuously incorporated at CRISPR loci, enabling partial reconstruction of recent HGT events. Curtailment of HGT through CRISPR-Cas immunity can be favourable when dealing with obligate parasites, but can be detrimental by hindering the acquisition of genetic novelty and/or beneficial traits carried by MGEs. Invasion by these MGEs can, however, be associated with fitness costs that may lead to selection against carriage [5,21]. Hence, bacteria often face a trade-off between immunity and acquisition of novel elements, which favour adaptation to different ecological niches and stressors, such as antibiotic pressure. MGEs can be equipped with inhibitors of CRISPR-Cas systems, called anti-CRISPR (Acr) proteins, which have

been reported mostly in prophages [22–24]. Recently, Acr proteins were identified in non-phage

MGEs, including plasmids and ICEs [25].

Bacterial pathogens belonging to the ESKAPE panel consist of five species (*Enterococcus faecium*,

Staphylococcus aureus, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, and *Pseudomonas*

aeruginosa) and one genus (*Enterobacter* sp.) [26,27]. These pathogens are frequently involved in

nosocomial infections, due to their multi-drug resistance and/or invasive phenotypes [28–33]. The

WHO recently published a list of pathogens for which new antibiotic development is urgently

required, and the ESKAPE pathogens were designated "priority status" [34]. AMR and virulence

genes are broadly distributed in plasmids across the ESKAPE pathogens [28,35], and also in ICEs [36,37].

Recently, CRISPR-Cas systems have been identified in plasmids and ICEs from several bacterial

species (including representatives of the ESKAPE pathogens), and may be involved in conflict

between MGEs [38–40].

In this study, we performed the first systematic analysis focusing on the ESKAPE pathogens

mobilome. We asked i) how prevalent are different MGEs (plasmids, ICEs/IMEs, and prophages)

across the ESKAPE pathogens; ii) how broad or constrained is the combined MGEs' network; iii)

which traits are overrepresented in these MGEs, and if AMR and virulence genes are differently

distributed in pairs of conspecific ESKAPE pathogens with and without CRISPR-Cas systems; iv)

whether the CRISPR spacers have a targeting bias towards different MGE types. We found that

plasmids, ICEs/IMEs, and prophages are unequally distributed across these pathogens, and

found signatures of DNA sharing events between different species. Uncovering the structure of

MGEs and masked (i.e., MGE-free) genomes allowed us to discover an overrepresentation of

AMR genes and anti-CRISPRs in the ESKAPE mobilome. Our results also unveiled ESKAPE-specific

trends of MGEs, AMR, and virulence genes promoted by the presence of CRISPR-Cas systems.

Finally, our work shows that CRISPR spacers found on ICEs/IMEs and plasmids across the ESKAPE

pathogens have different targeting biases.

Results

1. **MGEs are unevenly distributed among the ESKAPE pathogens**

We downloaded 1782 ESKAPE complete genomes from NCBI's RefSeq database. To correct for

species taxonomy, genomes with <95% average nucleotide identity (ANI) were removed for each

ESKAPE species (**Table S1**). Since this parameter is only applied for species delineation, we also

built a phylogenomic tree with *Enterobacter* sp. genomes and type strains belonging to the

Enterobacteriaceae family (**Figure S1**). Our curated dataset included 1746 complete genomes,

belonging to multiple MLST profiles (**Table S2**) and consisting of the same number of

chromosomes plus 2640 plasmids (**Figures 1A and 1D**). We confirmed that plasmids were

abundant in *K. pneumoniae* (~3 plasmids/genome). These replicons were also prevalent in every

ESKAPE except *P. aeruginosa* and *S. aureus* (**Figure 1A**). The majority of plasmids carried a

relaxase (62.5%, 1651/2640), and were classified as mobilizable (either self-conjugative or not)

[41].

To look for regions of genome plasticity (RGPs) exclusively integrated in the chromosome, we

used the 1746 chromosomal replicons to generate plasmid-free pangenomes for each ESKAPE

taxon. We identified a total of 50482 plasmid-free RGPs in chromosomal replicons (**Figure 1E**).

Of these, 2685 were classified as ICEs/IMEs due to the presence of relaxase and integrase

domains (**Figures 1D and 1E**). At least one ICE/IME was detected in >50% of genomes for all

ESKAPE pathogens, and was abundant in *E. faecium* and *P. aeruginosa* (~3 elements/genome)

(**Figures 1B and 1D**). After masking the ICEs/IMEs identified in the ESKAPE chromosomes, we

performed a search for prophages. Prophages were less frequently found than ICEs/IMEs and

plasmids (**Figures 1C and 1D**). Prophages were however abundant in *P. aeruginosa*, *Enterobacter*

sp., and *K. pneumoniae*. We found ~2 elements/genome on *K. pneumoniae*, whereas these

elements were particularly rare in *S. aureus*.

When looking into the presence/absence combination of co-occurring MGEs across the ESKAPE

pathogens, we noticed that the most frequent combination involved the presence of the three

MGEs (in 387 out of the 1746 genomes, **Table S2**). Only 144 strains carried none of these

elements. Examples of these contrasting combinations (i.e., all MGEs vs. no MGE at all) were

found in all representatives of the ESKAPE pathogens. We noticed that the majority of the strains

with the three MGEs co-occurring in the same genome belonged to *K. pneumoniae* (268/387),

while the no MGEs combination was mostly represented by *S. aureus* (118/144). Non-

surprisingly, we observed that *S. aureus* was the only ESKAPE representative with a proportion

of MGE/strain lower than one. Our results show that different MGEs are asymmetrically

distributed across the ESKAPE pathogens, with *K. pneumoniae* genomes taking the lead for the

co-occurrence of ICEs/IMEs, plasmids, and prophages (**Figure 1**).

2. DNA sharing between different vectors is common across the ESKAPE mobilome

MGEs tend to have a GC content lower than the one from its host [42–44]. However, how conserved

is this difference when considering different MGE types is poorly studied. We confirmed that

the arithmetic mean GC content of plasmids, ICEs, and prophages is significantly lower when

compared to masked genomes across the ESKAPE pathogens (**Figure S2**, p-value < 2.2e-16). With

the exception of *S. aureus*, we observed that plasmids across the ESKAPE pathogens have a wider

variation in their size, when compared with ICEs/IMEs and prophages (**Figure S3**). Across all

ESKAPE pathogens, we observed a weak positive correlation between the ICEs/IMEs and

plasmids' GC content and sequence length (R = 0.38 and 0.35, respectively, p<2.2e-16), and a

weak negative correlation between the prophages' GC content and sequence length (R = -0.074,

p = 0.0031, **Figure S4**).

We used an alignment-free sequence similarity comparison of the ESKAPE mobilome to infer an

undirected network (**Figures 2A and 2B**). The sparse network assigned 96.7% (6715) of the MGEs

into 150 clusters. The majority of MGE pairs shared little similarity, with a Jaccard Index (JI) value

below 0.25 (**Figure S5A**), in accordance with the high diversity frequently observed across MGEs.

The network revealed clear structural differentiation, where the majority of clusters were

homogeneous for a given ESKAPE/MGE pair (**Figures 2A and 2B**). The absence of pairwise

distance similarities with intermediate JI (**Figure S5A**) helps to explain this clustering in discrete

groups, instead of a continuous genetic structure. However, the two largest clusters challenge

interspecies and MGE type barriers and correspond to multiple MGEs with the four

Proteobacteria representatives in the first (i.e., *K. pneumoniae*, *Enterobacter* sp., *P. aeruginosa*,

and *A. baumannii*), and *S. aureus* and *E. faecium* in the second cluster. MGEs within these

promiscuous clusters tend to be more dissimilar than those assigned to densely connected and

ESKAPE/MGE pair-restricted clusters. To plot this network, we used as a threshold the mean

value (0.07612963) of the estimated pairwise distances between the 6941 MGEs identified in

this study. We also plotted the network using the median value (0.009765625) and confirmed a

lower sparsity of the network and the two larger clusters physically linked to one another (**Figure

S6**). In parallel, we built a network using the masked genomes as nodes connected by edges

indicating pairwise distances, and we exclusively observed species-specific clustering (**Figures

S5B and S7**), which is in agreement with cohesive evolutionary forces that maintain the genomic

coherence of bacterial species [45,46]. Interestingly, we verified that ICEs/IMEs from *P. aeruginosa*

are the MGE/ESKAPE pair most frequently associated with other bacteria, sharing DNA segments

with plasmids from *Enterobacter* sp., *K. pneumoniae*, and *A. baumannii*, as well as ICEs/IMEs

from *Enterobacter* sp. and *K. pneumoniae*, and prophages from *K. pneumoniae* (**Figure S8** and

Table S3). Even though rare, we observed cross-phylum interactions, between plasmids from *P.*

aeruginosa and plasmids from *S. aureus*, as well as between an ICEs/IME from *E. faecium* and a

plasmid from *K. pneumoniae* (**Figure S8** and **Table S3**). Altogether, our results highlight that

while some elements are capable of exchanging DNA beyond the genus (and phylum) barrier,

for the majority of clusters host similarity and MGE type restrain DNA sharing between different

ESKAPE MGEs.

3. The proteome of the ESKAPE mobilome is highly diverse in sequence and functions

To gain functional insights into the proteome of the ESKAPE mobilome, we investigated the

diversity of clusters of orthologous groups (COGs) encoded by the MGEs identified in this study.

COGs are protein sets conserved across different lineages that typically share functions, and are

therefore used for systematic function prediction in poorly characterised genomes [47].

Furthermore, COGs have been assigned to curated and uniform functional categories, enabling

the comparison of their distribution among genome sequences [48]. We distinguished 1183

different COGs in ESKAPE MGEs (**Figure S9**) encompassing most of the functional categories

reported in the COGs scheme (**Figure 3**), thereby highlighting the diversity of functions

associated with the ESKAPE mobilome. Plasmids, prophages, and ICEs/IMEs contain unique

COGs, consistent with their distinctive features as mobile elements (**Figure S9**). However, they

also shared 113 COGs, indicative of a common pool of proteins and functions carried by MGEs

in ESKAPE pathogens. We detected some variation in the relative contribution of MGE

proteomes to different COG functional categories among the ESKAPE pathogens (**Figure 3**). For

example, proteins associated with "carbohydrate transport and metabolism" are more frequent

in MGE proteomes of *E. faecium* (particularly in plasmids and ICEs/IMEs) than in other ESKAPE

(**Figure 3**). On the other hand, proteins of the "replication, recombination and repair" category

are similarly frequent amongst MGE proteomes of most ESKAPE. Another interesting example is

the relative high frequency of "translation, ribosomal structure and biogenesis" proteins in the

proteome of *K. pneumoniae* prophages, which appears to be distinct across MGE/ESKAPE pairs.

To explore the diversity of the ESKAPE MGEs proteome further, we clustered their 474223

proteins based on sequence similarity, resulting in 51960 groups (**Table S4**). Around 78% of the

representatives of these protein groups were assigned the tag "hypothetical protein" by Prokka,

underlining the large proportion of uncharacterised proteins encoded by ESKAPE MGEs. Among

the representatives with an assigned function, transposases and integrases were the most

frequent protein product (1829 cluster representatives) (**Table S4**). Recombinase, repressor and

resistance, were also common terms across the representative products, with more than 200

occurrences each; the latter being mostly associated with metal or drug resistance. We looked

for widespread protein clusters in the MGEs proteome, i.e. those observed in the three MGE

types or in at least three of the ESKAPE pathogens. Our search resulted in the identification of

466 protein clusters widespread across MGEs and 383 present in at least three different ESKAPE

(**Table S4**). Interestingly, only 38 clusters were identified in common between the two

widespread categories. Although hypothetical proteins dominated both widespread categories

(79% of MGEs and 52% of ESKAPE widespread protein clusters), various protein clusters with

functions associated with transposition and antimicrobial resistance were also identified (**Table**

S4). Hierarchical clustering of the MGE/ESKAPE pairs and protein clusters based on the

distribution and relative frequency of widespread proteins uncovered structured patterns of

sharing (**Figure 4**). For example, we detected a large component of protein clusters present in

plasmids of *Enterobacter* sp., *K. pneumoniae*, and *A. baumannii*. When it comes to protein

clusters present in different MGE types, we observed sets predominantly occurring in ICEs/IMEs

and prophages of *A. baumannii* and *S. aureus*. Overall, the distribution of ESKAPE widespread

proteins clustered MGE/ESKAPE pairs by MGE type (**Figure 4**). The clustering observed from the

distribution of MGE widespread proteins was more intricate but tended towards the same

ESKAPE pathogen, with the exception of *Enterobacter* sp. and *K. pneumoniae*, which were

typically clustered together.

4. AMR genes are overrepresented in the ESKAPE mobilome

In order to explore the AMR repertoire of the ESKAPE mobilome, we only focused on genes that

are horizontally transferred, such as beta-lactamases and aminoglycoside-modifying enzymes

(that lead to antimicrobial inactivation), and those that promote target site modification (such

as rRNA methyltransferases, the *vanA* and *vanB* gene clusters, and the staphylococcal cassette

chromosome *mec*). We observed that AMR genes are broadly distributed in plasmids and

ICEs/IMEs across the ESKAPE pathogens (**Figure S10**). All ESKAPE pathogens have a large

proportion of AMR-carrying plasmids (>35% of plasmids across the different ESKAPE carry at

least one AMR gene), while a high proportion of AMR-harbouring ICEs (>25%) was only observed

for *S. aureus* and *P. aeruginosa* (**Figures S11A and S11B**). As expected from the vast repertoire

of MGEs present in *K. pneumoniae* (**Figure 1D**), this species presented a wider selection of

different AMR genes. Some AMR genes were restricted to specific ESKAPE pathogens, while

others were more promiscuous. For example, different beta-lactamases (*bla* genes) were

prevalent among proteobacterial representatives of the ESKAPE pathogens, but were mostly

absent from *S. aureus* and *E. faecium* (**Figure S10**). The only exception was the *blaZ* gene, which

was exclusively identified in plasmids and ICEs/IMEs from *S. aureus*. This gene is typically

embedded within the SCCmec elements of this species and may have been acquired from

distantly related non-*Staphylococcus* species [49]. Genes encoding resistance to aminoglycosides

(*aac*, *ant*, and *aph* genes), chloramphenicol (*cat* genes), trimethoprim (*dfr* genes), and

tetracyclines (*tet* genes) were found in all representatives of the ESKAPE pathogens. Genes

involved in resistance to vancomycin (the *vanHAX* and *vanHBX* gene clusters) were exclusively

found in *S. aureus* and *E. faecium*, while genes coding resistance for quinolones (*qnr* genes) and

colistin (*mcr* genes) were only found in the proteobacterial representatives (**Figure S10**).

We next assessed the distribution of virulence genes. These genes are more prevalent in

ICEs/IMEs than in plasmids (**Figure S12**). In fact, we identified no virulence genes in *E. faecium*

plasmids, and only 0.6% of *A. baumannii* plasmids carry these genes. More than 25% ICEs/IMEs

in *S. aureus*, *K. pneumoniae*, and *E. faecium* carried at least one virulence gene (**Figures S11C**

and S11D). *K. pneumoniae* is again the ESKAPE pathogen carrying a wider variety of genes in

both ICEs/IMEs and plasmids. As previously reported [50], we noticed that virulence and AMR

genes are rarely encoded by prophages. An important exception is the enterotoxins found in

prophages from *S. aureus* (**Figure S12**), which are also known as pyrogenic toxin superantigens

(PTSAgs) [51]. Polyketide synthesis loci *ybt* and *clb* encoding the iron-scavenging siderophore

yersiniabactin and genotoxin colibactin, respectively, were solely identified in

Enterobacteriaceae representatives of the ESKAPE pathogens (i.e., *K. pneumoniae* and

Enterobacter sp.). These virulence loci were mostly present in ICEs/IMEs, as previously reported

[52], but we also found these genes on plasmids (**Figure S12**). Interestingly, *S. aureus* was the

ESKAPE pathogen with a higher proportion of both plasmids and ICEs/IMEs carrying at least one

AMR or virulence genes (**Figure S11**).

Since chromosomes are substantially larger than MGEs and consequentially have more genes,

we corrected the prevalence of AMR and virulence genes to the total number of genes present

in MGEs and masked genomes across the different ESKAPE pathogens. Overall, we noticed that

AMR genes were largely overrepresented in MGEs (~14x), when compared with masked

genomes (**Figure 5 and Table S5**). On the other hand, virulence genes were ~2x more likely to

be located on masked genomes. Taken together, our results show that plasmids are important

vectors for AMR genes across the ESKAPE pathogens, while ICEs/IMEs play a more important

role for the distribution of virulence genes. When compared with masked genomes, we

confirmed that AMR genes are preferentially distributed in the ESKAPE mobilome.

5. CRISPR-Cas systems shape the number of MGEs, AMR, and virulence genes across ESKAPE pathogens

CRISPR-Cas systems were identified in every ESKAPE pathogen except *E. faecium*, which was therefore excluded from subsequent analyses. The proportion of genomes with CRISPR-Cas systems varied across the ESKAPE pathogens, from around 45.7% for *P. aeruginosa* to around 0.7% for *S. aureus* (**Figure 6A** and **Table S2**). We then explored the prevalence of CRISPR-Cas systems across closely related strains belonging to the same MLST profile. Since *Enterobacter* sp. consists of multiple species, this ESKAPE pathogen was excluded from subsequent analyses. Given the low prevalence of CRISPR-Cas systems in *S. aureus*, this species was also excluded, and we focused exclusively on *P. aeruginosa*, *K. pneumoniae*, and *A. baumannii*. Interestingly, some sequence types (ST) consisted entirely of either CRISPR-Cas positive or negative genomes (**Table S2**). For example, the most frequent MLST profile in *A. baumannii* from our dataset was ST2 (n=101), which only included strains with no CRISPR-Cas systems. On the other hand, the second most prevalent MLST profile in this species (ST1, n=14), only consisted of strains with I-F CRISPR-Cas systems. The most frequently observed MLST profile from *K. pneumoniae* (ST11, n=105), consists mostly of CRISPR-Cas negative strains (96.2%, 101/105). The four strains with positive hits carried IV-A3 CRISPR-Cas systems on plasmids. ST258 (n=47) was the second most common *K. pneumoniae* MLST profile identified in our dataset, and again, consisted entirely of strains with no CRISPR-Cas systems. However, ST147 and ST15 (n=31 and n=23, respectively) carried I-E CRISPR-Cas systems in all strains. Finally, looking at *P. aeruginosa*, we found that the most prevalent MLST profiles in our dataset (ST235 and ST549, n=16 and n=11, respectively) carried no CRISPR-Cas systems. The only exception was found in a ST235 strain, which carried an I-C CRISPR-Cas system on an ICE/IME. In contrast, ST233 and ST1971 (n=8 for both) consisted exclusively of strains carrying the I-F CRISPR-Cas system on masked genomes (**Table S2**). These findings suggest that the presence or absence of CRISPR-Cas systems across the ESKAPE pathogens is largely driven by the expansion of specific clones.

Our analysis revealed a large variety of MGE-encoded CRISPR-Cas subtypes, with I-C, I-E, I-F, III-A, IV-A1, and IV-A3 represented across the dataset (**Figure 6B**). We found CRISPR-Cas systems on plasmids (n=28 IV-A3 subtype in *K. pneumoniae* and n=1 IV-A1 in *P. aeruginosa*) and ICEs/IMEs (n=7 I-C in *P. aeruginosa* and n=1 III-A in *S. aureus*) (**Figure 6B** and **Table S6**). The MGEs carrying these systems were large, ranging from 102-430 kb. Additionally, all CRISPR-Cas-carrying plasmids have a MOB relaxase. This is in agreement with previous findings [53], which

reported an enrichment of CRISPR-Cas systems across plasmids with conjugative functions and

of larger sizes. We also found AMR and virulence genes on these CRISPR-Cas positive MGEs, but

no anti-CRISPRs, suggesting that the CRISPR-Cas systems are functional.

When looking into the influence of GC content and sequence length in pairs of conspecific

ESKAPE pathogens with and without CRISPR-Cas systems, we would expect to observe smaller

and GC richer strains on those carrying these systems. These expectations could only be

confirmed for *P. aeruginosa*, for which CRISPR-Cas positive genomes were significantly smaller

than their counterparts (**Figure S13**, p-value 0.00037), as observed before [54–56]. Still, the non-

significant differences observed for *A. baumannii*, *Enterobacter* sp., *Klebsiella pneumoniae*, and

S. aureus could in part be explained by the low sample size of CRISPR-Cas positive genomes

(**Figure 6A**). Regarding the GC content, we observed significant differences in *A. baumannii*, *K.*

pneumoniae, and *P. aeruginosa*. CRISPR-Cas positive genomes were GC richer in *A. baumannii*

and *P. aeruginosa* (**Figure S14**, p-values 0.00098 and 0.00673, respectively). Curiously, we

noticed that CRISPR-Cas positive genomes in *K. pneumoniae* were GC poorer (**Figure S14**, p-value

2.5e-05). Given the known association between GC content and genome size [44], these

differences in CRISPR-Cas positive and negative *P. aeruginosa* genomes may be a spurious

correlation driven by small size of CRISPR-Cas positive genomes. So, we corrected the GC content

for variation in genome size, and we observed that the association was maintained (**Figure S15**,

p-value 0.00042), in accordance to a previous study [55].

Since virulence genes are overrepresented in the chromosome (**Figure 5**), we assessed the

distribution of these genes in pairs of conspecific ESKAPE pathogens with and without CRISPR-

Cas systems. Virulence genes were significantly less abundant in CRISPR-Cas positive genomes

from *P. aeruginosa* and *A. baumannii* (**Table S7** and **Figure S16**, p-values 1.7e-06 and 0.0016,

respectively). Given that *P. aeruginosa* genomes positive for these systems are significantly

smaller than their CRISPR-Cas negative counterparts (**Figure S13**), the lower prevalence of these

genes in CRISPR-Cas positive *P. aeruginosa* genomes may again be driven by a spurious

correlation. As so, we corrected the number of virulence genes for variation in genome size, and

we observed that indeed the difference was no longer significant (**Figure S17**, p-value 0.73),

confirming our prediction that the genome size was a confounding variable obscuring the effect

of CRISPR-Cas systems on the prevalence of virulence genes in *P. aeruginosa*.

We then explored whether CRISPR-Cas presence or absence reduced the number of MGEs

acquired in pairs of conspecific ESKAPE pathogens. We would expect to detect a smaller number

of MGEs in genomes harbouring these immune systems. Indeed, this trend was observed in *K.*

pneumoniae and *P. aeruginosa* (**Figure 7**). Conversely, CRISPR-Cas positive *A. baumannii*

genomes were significantly larger than genomes of the same species lacking these systems (p-

value 0.047). Finally, we focused on AMR and virulence genes carried exclusively by plasmids

and ICEs, as these were the most important vectors (**Figure S10**). For most MGE/ESKAPE pairs,

we observed no significant difference between pairs of conspecific genomes with and without

CRISPR-Cas systems. When it comes to AMR genes, we only observed significant differences in

MGEs from *P. aeruginosa* (**Figure S18A**, p-values 0.037). Indeed, AMR genes were more

prevalent on ICEs/IMEs from *P. aeruginosa* genomes with CRISPR-Cas systems (**Table S8**). A

similar correlation was previously reported [56]. Curiously, the less prevalent I-C CRISPR-Cas

subtype, which was exclusively identified in *P. aeruginosa* and mostly on ICEs/IMEs (**Table S2**),

was found to be positively correlated with certain AMR genes [54]. Regarding virulence genes, we

observed significant differences in MGEs from *K. pneumoniae*, where CRISPR-Cas-carrying

elements were associated with more virulence genes than their CRISPR-Cas negative

counterparts (**Figure S18B**, p-values 0.0054). Taken together, we observed species-specific

trends shaping the number of MGEs, AMR, and virulence genes across pairs of conspecific

ESKAPE genomes with and without CRISPR-Cas systems.

6. Anti-CRISPRs are overrepresented in the ESKAPE mobilome

Anti-CRISPR proteins (n=161) antagonising CRISPR-Cas subtypes I-E, I-F, and II-A were identified

in a total of 90 *P. aeruginosa*, *K. pneumoniae*, and *A. baumannii* MGEs (**Figure 6C** and **Table S9**).

ICEs/IMEs were the most important vectors for these counter-defence systems in *P. aeruginosa*,

while prophages were more frequent in *K. pneumoniae*. Even though *P. aeruginosa* was the

ESKAPE pathogen with a higher proportion of genomes with CRISPR-Cas systems (**Figure 6A**), we

observed Anti-CRISPR proteins on 28.3 % (n=34/120) CRIPR-Cas positive genomes from this

species (**Table S2**). We also looked for these proteins across the masked genomes, and found

hits in all ESKAPE except *S. aureus* (n=252, **Table S10**). After correcting their prevalence to total

number of genes, we verified that anti-CRISPRs are largely overrepresented in MGEs (~9x) when

compared with masked genomes (**Figure 5**). When compared with masked genomes, our results

show that Anti-CRISPR proteins are preferentially encoded in MGEs.

7. CRISPR spacers in ICEs/IMEs and plasmids have different targeting biases

We explored the targets for all CRISPR spacers, retrieved from complete CRISPR-Cas systems, but also orphan CRISPRs, identified in our collection of ESKAPE genomes. Since we provided here a representative dataset of plasmids, ICEs/IMEs, and prophages, we used this collection as a database and took the CRISPR spacers identified in masked genomes as a query. In parallel, we used the CRISPR spacers identified in MGEs as a query and the MGEs collection as a database. For the latter, and to avoid self-targeting hits, we masked all CRISPR spacers from the MGEs collection used as database. We found a total of 897 CRISPR spacers across the MGEs (n=554 on plasmids and n=343 on ICEs/IMEs, **Table S11**). Only a small fraction of plasmids and ICEs/IMEs carried these spacers (1.3%, 34/2640 plasmids and 0.6%, 17/2685 ICEs/IMEs, respectively). No CRISPR spacers were found on prophages, which is in agreement with the lack of CRISPR-Cas systems in these elements from our collection. Given the large number of MGEs and CRISPR-Cas-encoding plasmids in *K. pneumoniae* (**Figure 1** and **Figure 6B**), most spacers were found in this species (522/897). The large majority of plasmid spacers were identified on mobilizable plasmids (99.4%, 551/554).

We then looked for protospacers, and identified invading DNA fragments on 9.2% of total MGEs (n=641/6941, **Table S12**). A substantially larger fraction of CRISPR spacers from plasmids targeted mobilizable plasmids from our ESKAPE collection (92.6%, 21488/23200 of total plasmid spacer's interactions). Only a small fraction of plasmid spacers targeted protospacers in non-transmissible plasmids (5.2%), prophages (1.6%), and ICEs/IMEs (0.6%) (**Figure 8A**). Moreover, ICEs/IMEs spacers were also biased towards other ICEs/IMEs (70.7%, 795/1124 of total ICEs/IMEs interactions). However, unlike plasmids, a larger proportion of ICEs/IMEs spacers targeted prophages (27.7%) (**Figure 8A**). Only a small fraction of ICEs/IMEs spacers targeted plasmids (1.6%).

When it comes to CRISPR arrays on masked genomes, we observed a total of 13709 spacers in 30.8% ESKAPE genomes (n=538/1746, **Table S13**). Even though the prevalence of CRISPR-Cas systems in *S. aureus* was considerably low (< 1%, **Figure 6A**), we found spacers in 31.4% of the total *S. aureus* strains (n=174/553). Consistent with the absence of CRISPR-Cas systems in the *E. faecium* strains from our dataset, no spacers were identified in masked genomes from this species. The number of CRISPR spacers per genome varied considerably between the masked genomes of the ESKAPE pathogens (**Figure S19**), reaching as high as 189 in *A. baumannii*. In fact, only strains from this species carried more than 100 spacers per genome. We observed that 27.8% (3815/13709) of CRISPR spacers in ESKAPE masked genomes yielded significant matches to protospacer MGE sequences, found on 25.0% MGEs (n=1735/6941, **Table S14**). Most CRISPR spacers targeted plasmids (72.8%, 84256/115743 total interactions), followed by ICEs/IMEs

(16.9%) and prophages (10.3%). As observed for CRISPR spacers in MGEs, most plasmid targets

correspond to mobilizable plasmids (88.5%, 74567/84256). We then tested if MGE or masked

genome spacers preferentially targeted ESKAPE MGEs of variable size. We found that the CRISPR

spacer targets were significantly larger than those with no protospacers (p-value <2.2e-16,

Figures 8B and 8C). Altogether, our results show that ICEs/IMEs preferentially target ICEs/IMEs

and also prophages, while plasmids almost exclusively target other plasmids. Our data also

shows that CRISPR spacers either on MGEs or masked genomes consistently target larger MGEs.

Discussion

In this work, we performed a systematic analysis of ICEs/IMEs, plasmids, and prophages across

all ESKAPE pathogens. We focused on this panel because the ESKAPE group of pathogens consist

of clinically-relevant bacteria, for which many genomes are completely sequenced (an important

parameter when delineating MGEs), and which include representatives of both Proteobacteria

and Firmicutes, and also phylogenetically divergent bacteria (with exception of *K. pneumoniae*

and *Enterobacter* sp., the remaining representatives belong to different bacterial families).

Studying MGEs in parallel allowed us to explore their uneven distribution across a collection of

complete genomes from important pathogens, and to explore DNA sharing events between

different MGE types. By separating these elements from masked (MGE-free) genomes, we were

able to observe an overrepresentation of AMR genes and anti-CRISPRs across the ESKAPE

mobilome. Furthermore, we assessed the influence of CRISPR-Cas defence systems in shaping

the acquisition of MGEs and beneficial genes, and we unveiled different targeting biases for

CRISPR spacers identified in plasmids and ICEs/IMEs.

The network-based approach used here to study the ESKAPE MGEs revealed a clear structural

differentiation, where the majority of clusters were homogeneous for a given ESKAPE/MGE pair.

Using pairwise genetic distances of alignment-free *k*-mer sequences has circumvented the

exclusion of non-coding elements that was observed in gene content similarity networks from

previous work [57], providing a more comprehensive picture of plasmid population and dynamics

[12]. Other groups have shown that plasmids form coherent clusters [12,13], similar in concept to

what was observed for bacterial genomes [45,46,58]. Focusing on the ESKAPE pathogens, we

demonstrated here that the same happens for ICEs/IMEs and prophages for the majority of the

clusters. However, two large heterogeneous clusters were also observed. Unlike bacterial

genomes, where recombination between closely related replicons is the main force promoting

genomic cohesiveness [45,58], MGEs such as plasmids, ICEs/IMEs and prophages are capable of

exchanging DNA beyond the genus barrier (and even occasionally across phylum) [59]. Based on

our network-based approach, we observed that the ESKAPE mobilome appears to follow a

bipartite structure, with some elements being capable of shuffling DNA between distantly

related species.

The abundance of MGEs is strongly associated with the prevalence of AMR genes. This is

particularly evident for *K. pneumoniae*, which carries a high proportion of important vectors of

AMR genes, such as plasmids and ICEs/IMEs. After correcting the prevalence to total number of

genes, we observed that AMR genes are nearly 15 times more likely to be found on ESKAPE

MGEs than on masked genomes. Most likely the use of different antibiotics targeting either

Gram negative or positive infections may have selected for the emergence of different AMR

genes across the Proteobacteria/Firmicutes divide. For example, vancomycin is used as last

resort for the treatment of sepsis and other infections caused by Gram-positive bacteria, while

colistin mainly serves to target multi-drug resistant Gram-negative infections [26]. Anti-CRISPRs

are nearly 10 times more likely to be detected on these MGEs than on masked genomes. Unlike

AMR genes, where genes conferring resistance to similar antibiotics where identified in different

ESKAPE pathogens, the distribution of virulence genes across the ESKAPE mobilome was mostly

species-specific. This may be explained by different mechanisms of virulence and toxicity across

bacterial species. Even considering the relative proportion of these genes, we confirmed that

these genes are twice more likely to be identified in masked genomes than in MGEs.

Curiously, we found no CRISPR-Cas systems in our curated *E. faecium* dataset. In addition to our

main analysis, we specifically searched for the presence of these systems on the excluded

genomes, based on <95% genome similarity threshold defined for species delineation [58], and

found three strains with these systems (**Table S15**). Since the majority of the most

representative MLST profiles in our collection consists of genomes either with or without

CRISPR-Cas systems (**Table S2**), analysis of intra-ST CRISPR variability between pairs of

conspecific ESKAPE genomes was not performed in this study. Except for *P. aeruginosa* [54,55], we

found no evidence for genome length as a marker for HGT inhibition by CRISPR-Cas systems. A

similar observation was made before, for a different collection of bacterial pathogens [56].

Defence systems such as CRISPR-Cas are inherently costly to bacterial hosts, mainly due to

different forms of autoimmunity [60]. To offset the short-term deleterious effects, these systems

benefit from associating with MGEs, such as the examples observed here and elsewhere [38].

Although these and other defence mechanisms such as restriction-modification systems do not

qualify as bona fide MGEs (the systems lack mechanisms controlling their own replication), these

quasi-autonomous systems take advantage of MGEs to promote their own dissemination and
maintenance across bacterial hosts. Conversely, MGEs benefit from these systems and may
pervasively repurpose them for inter-MGE competition [53]. In fact, we found that spacers in the
ICE/IME and plasmid CRISPR arrays target competing MGEs, underscoring the genetic
independence of CRISPR-Cas systems in ICEs/IMEs and plasmids in MGE-MGE conflicts. Even
though we found no CRISPR-Cas systems in prophages, a phage-encoded system was identified
in *Vibrio cholerae*, and targeted phage-inducible chromosomal islands [61]. Importantly, the
presence of CRISPR-Cas subtypes preferentially distributed in MGEs (I-C and IV-A3) points to the
existence of distinct selective pressure that promote the maintenance of specific subtypes on
plasmids and ICEs/IMEs versus masked genomes. Unsurprisingly given the large sizes of these
systems, we observed a bias towards larger MGEs (both plasmids and ICEs/IMEs >100kb). Since
plasmids with an identifiable relaxase (conjugative and mobilizable) are larger than the so-called
non-transmissible plasmids [41], we also found a relaxase in all CRISPR-Cas positive plasmids.
Prophages seem to follow similar streamlining dynamics [62]. The high proportion of *P. aeruginosa*
genomes carrying both CRISPR-Cas systems and anti-CRISPRs (28.3%, **Table S2**) suggests a large
fraction of these systems may be non-functional in this species, as hypothesized before [54]. Since
most Anti-CRISPRs in *P. aeruginosa* were identified in ICES/IMEs, these results underscore a
major role played by the overlooked ICEs/IMEs in potentially silencing immune systems in this
species.

Our results shows that only a restricted fraction of CRISPR spacers matched protospacer MGE
sequences, which is in agreement with previous findings [53,55]. While the large majority of
plasmid-encoded spacers were predicted to target other plasmids, CRISPR arrays in ICEs/IMEs
preferentially targeted other ICEs/IMEs, but also a large proportion of prophages. This
complementary targeting preference can be explained by the different lifestyle of these MGEs.
Since plasmids are maintained as extra-chromosomal elements, and ICEs/IMEs and prophages
are integrated in the chromosome, we hypothesize that while plasmids preferentially target
plasmid competitors [38], ICEs/IMEs exploit CRISPR-Cas systems to protect their host against other
ICEs/IMEs and viral predators.

Several bioinformatic tools exist to look for plasmids and prophages, but currently the options
for ICEs/IMEs are scarce [63]. We provide here an accurate identification of these elements,
building upon a recently reported tool to scan RGPs [64]. However, our approach depends on the
availability of a pangenome for the considered taxa, which is an important limitation for species
with an insufficient number of completely sequenced genomes. Moreover, our work only
focused on three MGE types, which are likely to be of main importance for the dissemination of

genes involved in pathogenesis and AMR and thus critical for our understanding of the evolution

of the ESKAPE pathogens. Nevertheless, other MGEs should be considered in future work, for

example those that contribute to communication between cells (such as phage-inducible

chromosomal islands [65]), or intracellular transfer events (e.g., transposons and insertions

sequences [35]).

To conclude, our results indicate that ICEs/IMEs, plasmids, and prophages are asymmetrically

distributed across the ESKAPE pathogens. We found that these MGEs can exchange DNA beyond

the genus (and phylum) barrier, even though most clusters are constrained by host similarity

and/or the type of MGE. We observed that most ESKAPE MGE proteins have unknown functions

and have convoluted distribution patterns. When comparing against masked (MGE-free)

genomes, we observed the pervasive association of AMR genes and anti-CRISPRs with the

ESKAPE mobilome. We also found different targeting biases to CRISPR spacers found on plasmids

and ICEs/IMEs, highlighting their genetic independence. Taken together, our results illustrate

the potential of network-based approaches and comparative genomics to underscore the

composition and dynamics of gene flow across different MGEs, and sheds a new light in the role

of the overlooked ICEs/IMEs as important players in the MGE-MGE warfare in the ESKAPE

pathogens and thus the main groups of highly problematic human pathogens.

Material and methods

ESKAPE pathogens collection

We retrieved all complete ESKAPE genomes available in the NCBI Reference Sequence Database

(RefSeq,   accessed   on   12-11-2020,   using   ncbi-genome-download   v0.3.0

(https://github.com/kblin/ncbi-genome-download). Genomes listed as 'unverified' were

removed from our dataset. We also excluded genomes classified as 'Enterobacteriaeceae'.

Finally, we used pyANI v0.2.10 (https://github.com/widdowquinn/pyani) to calculate the

average nucleotide identity based on MUMmer (ANIm), and removed genomes with an ANIm

value below the 95% threshold for species delineation [58,66]. To evaluate the taxonomy of the

Enterobacter species, we retrieved genomes for *Enterobacteriaceae* type strains and used them

together with the *Enterobacter* genomes to create a phylogenetic tree using GToTree v1.5.22

(https://github.com/AstrobioMike/GToTree)[67] and the IQ-TREE algorithm to estimate maximum

likelihood [68]. We used the pre-built set of 74 single copy gene bacterial Hidden Markov Models

(HMM) available in GToTree. Genomes labelled as belonging to the *Enterobacter* genus, but

clustered in the phylogenomic tree with type strains other than those from the *Enterobacter*

genus, were removed from subsequent analyses. MLST profiles were determined with mlst

v2.19.0 (https://github.com/tseemann/mlst). The curated genomes were automatically

annotated using Prokka v1.14.6 (https://github.com/tseemann/prokka) [69].

Extraction of plasmids, ICEs/IMEs, and prophages

Since the ESKAPE pathogens (as most bacteria) are haploid, we separated the large replicon (i.e.,

the chromosome) from the extrachromosomal replicons. For the latter, only accessions with

'plasmid' and 'complete sequence' on their description were kept and were used for further

plasmid analyses.

To extract ICEs from chromosomal replicons, we used the chromosomal genbank files created

with Prokka as input to build a pangenome for each ESKAPE pathogen. We used ppanggolin

v1.1.96 (https://github.com/labgem/PPanGGOLiN)[70], which uses a graphical model and a

statistical method to partition the pangenome in persistent, shell and cloud genomes. We used

the panRGP method to build the pangenomes [64]. This method predicts RGPs, which are clusters

of genes made of accessory genes (shell and cloud genomes) in the pangenome graph. We then

used bedtools v2.30.0 (https://bedtools.readthedocs.io/en/latest/) [71] to extract RGPs from the

chromosomal replicons. The proteomes of these extracted RGPs were scanned for relaxases

with hmmer v.3.3.1 (http://hmmer.org/) [72] against MOBfamDB, a curated relaxase profile HMM

database [73]. Simultaneously, the proteomes were screened with hmmer against integrases

(Pfam accession PF00589) and recombinases (PF07508). RGPs with hits both for integrases

(phage integrase or recombinase) and relaxases were classified as putative ICEs/IMEs and were

kept for further analysis.

To look for prophages, we masked the ICEs/IMEs locations in the chromosomal replicons using

bedtools. The ICE-masked chromosomal genbank files were then used as input in phispy v4.2.6

(https://github.com/linsalrob/PhiSpy), which combines similarity- and composition-based

strategies to look for prophages [74]. We also masked the prophage regions in the chromosomal

replicons, to build the final masked genomes, that are ICE- and prophage-masked (these masked

replicons are also free of plasmids, since these are part of the extrachromosomal replicons).

Network-based approach and functional annotation

To estimate pairwise distances between all ESKAPE MGEs (plasmids, ICEs and prophages), we

reduced the MGEs into sketches and compared the Jaccard index (JI) and mutation distances

between pairs of MGEs using BinDash v 0.2.1 (https://github.com/zhaoxiaofei/bindash)[75]. Each

MGE sequence was converted to a set of 21-bp *k*-mers. We used the mean() and median()

functions in R to calculate the arithmetic mean and median of the JI, respectively. Only JI equal

to or above the mean and median were considered, and the mutation distances were used as

edge attributes to plot the network with Cytoscape v3.9.0 under the prefuse force directed

layout (https://cytoscape.org/). We used the Analyzer function in Cytoscape to compute a

comprehensive set of topological parameters, such as the clustering coefficient, the network

density, the centralization, and the heterogeneity.

MGE proteins assignment to COGs was retrieved from the tab-separated annotation files

produced by Prokka. Lists of non-redundant COGs were extracted separately for prophages,

plasmids   and   ICEs/IMEs   and   compared   with   venny   v2.1.0

(https://bioinfogp.cnb.csic.es/tools/venny/index.html) to identify unique and shared COGs.

Likewise, COGs occurrence was determined separately for the proteomes of the three MGE

types in the different ESKAPE. Information on the COGs classification into functional categories

was retrieved from https://ftp.ncbi.nih.gov/pub/COG/COG2020/data/. The relative frequency

of the different COG functional categories per MGE/ESKAPE pair was calculated by summing up

the occurrences of COGs belonging to a given functional category and dividing the resulting

number by the total number of proteins observed in the corresponding MGE/ESKAPE pair.

To explore the diversity of MGE-encoded proteins we combined their proteomes (474223

proteins) and clustered them using the cluster algorithm from the MMSEQseqs2 v13.45111

package [76]. The proteins were clustered at 80% sequence identity and 80% coverage, with

otherwise default settings to match the parameters used by ppanggolin when generating the

ESKAPE pangenomes. The relative frequency of the different protein clusters identified per

MGE/ESKAPE pair was calculated following the same approach used to estimate the relative

frequency of COG functional categories, but using the occurrence of proteins belonging to a

given cluster instead. Representatives of the 51960 clusters identified were annotated with

eggNOG-mapper v2 [77], with default settings to further explore the functions of the MGE-

encoded proteins.

We used abricate v1.0.1 (https://github.com/tseemann/abricate) to scan extracted MGEs and

masked genomes against antimicrobial resistance and virulence genes (using pre-downloaded

databases from Resfinder[78] and VFDB[79] containing 3108 and 3686 sequences, respectively, and

both updated on the 05-02-2021). To identify and classify CRISPR-Cas systems, we used

CRISPRCasTyper v1.2.3 (https://github.com/Russel88/CRISPRCasTyper)[80]. We also used this tool

to look for CRISPR spacers. The entire CRISPR arrays identified on MGEs were then masked using

bedtools, and these masked MGEs served as a local blast database, when using MGE CRISPR

spacers as a query. CRISPR spacers from masked genomes were also blasted against a local

database of our extracted (non-masked) MGEs. Hits with at least 95% nucleotide identity and

95% sequence coverage were considered as protospacers [81]. While a representative collection

of plasmids and virus is publicly available at RefSeq's NCBI database (n=33269 and 13778,

respectively, accessed on the 21/05/2021), a substantially smaller collection of ICEs/IMEs is

available at ICEberg database (n=1325), and was last updated in September 2018. Due to this

limitation in the number of publicly available ICEs/IMEs sequences, and to have a representative

collection of these three different types of MGEs, we focused on the curated dataset presented

in this study to look for targets of CRISPR spacers.

We retrieved an Anti-CRISPR collection of non-redundant proteins from Anti-CRISPRdb

(http://guolab.whu.edu.cn/anti-CRISPRdb/, accessed on the 19-02-2021). This collection was

used to build a local database with diamond v2.0.9 (https://github.com/bbuchfink/diamond)[82].

We used the blastp command in diamond to scan the MGEs and masked proteomes against the

anti-CRISPR local database. We used an amino acid-based homology approach to find Anti-

CRISPRs encoded in the ESKAPE mobilome. Even though recent approaches have applied a guilt-

by-association method to identify new Anti-CRISPRs [25], currently there is no tool available to

apply this method in a large dataset of bacterial genomes.

Statistical analysis

Comparisons between MGEs' GC content and sequence length were performed using the

Kruskal-Wallis test, and the p-values adjusted using the Holm–Bonferroni method. Comparisons

between pairs of conspecific genomes with and without CRISPR-Cas systems, as well as between

MGE targets for CRISPR spacers, were performed using the Wilcoxon test, and the p-values

adjusted using the Holm–Bonferroni method. Values above 0.05 were considered as non-

significant (ns). We used the following convention for symbols indicating statistical significance:

* for p <= 0.05, ** for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001.

Code availability

All code used to create the main figures are available at

https://gitlab.gwdg.de/botelho/eskape_mobilome).

References

1.   Ghaly, T. M. & Gillings, M. R. Mobile DNAs as Ecologically and Evolutionarily
627   Independent Units of Life. *Trends Microbiol.* **26**, 904–912 (2018).

628   2.   Koonin, E. V. Viruses and mobile elements as drivers of evolutionary transitions. *Philos.*
629   *Trans. R. Soc. B Biol. Sci.* **371**, (2016).

630   3.   Kazazian, H. H. Mobile Elements: Drivers of Genome Evolution. *Science* **303**, 1626–1632
631   (2004).

632   4.   San Millan, A. & MacLean, R. C. Fitness Costs of Plasmids: a Limit to Plasmid
633   Transmission. *Microbiol. Spectr.* **5** (2017).

634   5.   Botelho, J. & Schulenburg, H. The Role of Integrative and Conjugative Elements in
635   Antibiotic Resistance Evolution. *Trends Microbiol.* **29**, 8-18 (2020).

636   6.   Chen, J. *et al.* Genome hypermobility by lateral transduction. *Science* **362**, 207–212
637   (2018).

638   7.   Dagan, T., Artzy-Randrup, Y. & Martin, W. Modular networks and cumulative impact of
639   lateral transfer in prokaryote genome evolution. *Proc. Natl. Acad. Sci. U. S. A.* **105**,
640   10039–10044 (2008).

641   8.   Smillie, C. S. *et al.* Ecology drives a global network of gene exchange connecting the
642   human microbiome. *Nature* **480**, 241–244 (2011).

643   9.   Ellabaan, M. M. H., Munck, C., Porse, A., Imamovic, L. & Sommer, M. O. A. Forecasting
644   the dissemination of antibiotic resistance genes across bacterial genomes. *Nat.*
645   *Commun.* **12**, 2435 (2021).

646   10.   Popa, O., Hazkani-Covo, E., Landan, G., Martin, W. & Dagan, T. Directed networks reveal
647   genomic barriers and DNA repair bypasses to lateral gene transfer among prokaryotes.
648   *Genome Res.* **21**, 599–609 (2011).

649   11.   Kloesges, T., Popa, O., Martin, W. & Dagan, T. Networks of Gene Sharing among 329
650   Proteobacterial Genomes Reveal Differences in Lateral Gene Transfer Frequency at
651   Different Phylogenetic Depths. *Mol. Biol. Evol.* **28**, 1057–1074 (2011).

652   12.   Acman, M., van Dorp, L., Santini, J. M. & Balloux, F. Large-scale network analysis
653   captures biological features of bacterial plasmids. *Nat. Commun.* **11**, 2452 (2020).

654   13.   Redondo-Salvo, S. *et al.* Pathways for horizontal gene transfer in bacteria revealed by a
655   global map of their plasmids. *Nat. Commun.* **11**, 3602 (2020).

656   14.   Campbell, D. E. *et al.* Interrogation of the integrated mobile genetic elements in gut-
657   associated Bacteroidaceae with a consensus prediction approach. *bioRxiv*
658   2021.09.02.458807 (2021) doi:10.1101/2021.09.02.458807.

659   15.   Croucher, N. J. *et al.* Diversification of bacterial genome content through distinct
660   mechanisms over different timescales. *Nat. Commun.* **5**, 1–12 (2014).

661   16.   Wein, T., Hülter, N. F., Mizrahi, I. & Dagan, T. Emergence of plasmid stability under non-

662   selective conditions maintains antibiotic resistance. *Nat. Commun.* **10**, 2595 (2019).

17. Rodriguez-Beltran, J. *et al.* Multicopy plasmids allow bacteria to escape from fitness trade-offs during evolutionary innovation. *Nat. Ecol. Evol.* **2**, 873–881 (2018).

18. Koonin, E. V., Makarova, K. S., Wolf, Y. I. & Krupovic, M. Evolutionary entanglement of mobile genetic elements and host defence systems: guns for hire. *Nat. Rev. Genet.* **21**, 119-131 (2019).

19. Roberts, R. J., Vincze, T., Posfai, J. & Macelis, D. REBASE—a database for DNA restriction and modification: enzymes, genes and genomes. *Nucleic Acids Res.* **43**, D298–D299 (2015).

20. Makarova, K. S. *et al.* Evolutionary classification of CRISPR–Cas systems: a burst of class 2 and derived variants. *Nat Rev Microbiol.* **18**, 67-83 (2020).

21. Millan, A. S. *et al.* Positive selection and compensatory adaptation interact to stabilize non-transmissible plasmids. *Nat. Commun.* **5**, 5208 (2014).

22. Bondy-Denomy, J., Pawluk, A., Maxwell, K. L. & Davidson, A. R. Bacteriophage genes that inactivate the CRISPR/Cas bacterial immune system. *Nature* **493**, 429–432 (2013).

23. Li, Y. & Bondy-Denomy, J. Anti-CRISPRs go viral: the infection biology of CRISPR-Cas inhibitors. *Cell Host Microbe.* **29**, 704–714 (2021).

24. Mahendra, C. *et al.* Broad-spectrum anti-CRISPR proteins facilitate horizontal gene transfer. *Nat. Microbiol.* **5**, 620–629 (2020).

25. Pinilla-Redondo, R. *et al.* Discovery of multiple anti-CRISPRs highlights anti-defense gene clustering in mobile genetic elements. *Nat. Commun.* **11**, 5652 (2020).

26. De Oliveira, D. M. P. *et al.* Antimicrobial resistance in ESKAPE pathogens. *Clin. Microbiol. Rev.* **33**, e00181-19 (2020).

27. Mortensen, K., Lam, T. J. & Ye, Y. Comparison of CRISPR–Cas Immune Systems in Healthcare-Related Pathogens. *Front. Microbiol.* **12**, 758782 (2021).

28. Wyres, K. L. *et al.* Genomic surveillance for hypervirulence and multi-drug resistance in invasive *Klebsiella pneumoniae* from South and Southeast Asia. *Genome Med.* **12**, 11 (2020).

29. Botelho, J., Grosso, F. & Peixe, L. Antibiotic resistance in *Pseudomonas aeruginosa* – Mechanisms, epidemiology and evolution. *Drug Resist. Updat.* **44**, 100640 (2019).

30. Davin-Regli, A., Lavigne, J.-P. & Pagès, J.-M. *Enterobacter* spp.: Update on Taxonomy, Clinical Aspects, and Emerging Antimicrobial Resistance. *Clin. Microbiol. Rev.* **32**, e00002-19 (2019).

31. Copin, R. *et al.* Sequential evolution of virulence and resistance during clonal spread of community-acquired methicillin-resistant *Staphylococcus aureus*. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 1745–1754 (2019).

32. Gao, W., Howden, B. P. & Stinear, T. P. Evolution of virulence in *Enterococcus faecium*, a hospital-adapted opportunistic pathogen. *Curr Opin Microbiol.* **41**, 76–82 (2018).

33. Harding, C. M., Hennon, S. W. & Feldman, M. F. Uncovering the mechanisms of *Acinetobacter baumannii* virulence. *Nat Rev Microbiol.* **16**, 91–102 (2018).

34. World Health Organization. Global priority list of antibiotic-resistant bacteria to guide

research, discovery, and development of new antibiotics. http://www.who.int/news-
room/detail/27-02-2017-who-publishes-list-of-bacteria-for-which-new-antibiotics-are-
urgently-needed.

35. Partridge, S. R., Kwong, S. M., Firth, N. & Jensen, S. O. Mobile Genetic Elements
Associated with Antimicrobial Resistance. *Clin. Microbiol. Rev.* **31**, e00088-17 (2018).

36. Botelho, J., Mourão, J., Roberts, A. P. & Peixe, L. Comprehensive genome data analysis
establishes a triple whammy of carbapenemases, ICEs and multiple clinically relevant
bacteria. *Microb. genomics* **6**, mgen000424 (2020).

37. Paauw, A., Leverstein-van Hall, M. A., Verhoef, J. & Fluit, A. C. Evolution in quantum
leaps: Multiple combinatorial transfers of HPI and other genetic modules in
Enterobacteriaceae. *PLoS One* **5**, 8662 (2010).

38. Pinilla-Redondo, R. *et al.* Type IV CRISPR-Cas systems are highly diverse and involved in
competition between plasmids. *Nucleic Acids Res.* **48**, 2000–2012 (2020).

39. León, L. M., Park, A. E., Borges, A. L., Zhang, J. Y. & Bondy-Denomy, J. Mobile element
warfare via CRISPR and anti-CRISPR in Pseudomonas aeruginosa. *Nucleic Acids Res.* **49**,
2114–2125 (2021).

40. Moya-Beltrán, A. *et al.* Evolution of Type IV CRISPR-Cas Systems: Insights from CRISPR
Loci in Integrative Conjugative Elements of *Acidithiobacillia*. *CRISPR J.* **4**, 656-672
(2021).

41. Smillie, C., Garcillán-Barcia, M. P., Francia, M. V., Rocha, E. P. C. & de la Cruz, F. Mobility
of plasmids. *Microbiol. Mol. Biol. Rev.* **74**, 434–52 (2010).

42. Nishida, H. Comparative Analyses of Base Compositions, DNA Sizes, and Dinucleotide
Frequency Profiles in Archaeal and Bacterial Chromosomes and Plasmids. *Int. J. Evol.*
Biol. **2012**, 342482 (2012).

43. Cury, J., Touchon, M. & Rocha, E. P. C. Integrative and conjugative elements and their
hosts: composition, distribution and organization. *Nucleic Acids Res.* **45**, 8943–8956
(2017).

44. Almpanis, A., Swain, M., Gatherer, D. & McEwan, N. Correlation between bacterial G+C
content, genome size and the G+C content of associated plasmids and bacteriophages.
Microb. genomics **4**, e000168 (2018).

45. Achtman, M. & Wagner, M. Microbial diversity and the genetic nature of microbial
species. *Nat Rev Microbiol*. **6**, 431–440 (2008).

46. Shapiro, B. J. *et al.* Population genomics of early events in the ecological differentiation
of bacteria. *Science* **335**, 48–51 (2012).

47. Galperin, M. Y. *et al.* COG database update: focus on microbial diversity, model
organisms, and widespread pathogens. *Nucleic Acids Res.* **49**, D274–D281 (2021).

48. Galperin, M. Y., Kristensen, D. M., Makarova, K. S., Wolf, Y. I. & Koonin, E. V. Microbial
genome analysis: the COG approach. *Brief. Bioinform.* **20**, 1063–1070 (2019).

49. John, J., George, S., Nori, S. R. C. & Nelson-Sathi, S. Evolutionary route of resistant genes
in *Staphylococcus aureus*. *Genome Biol. Evol.* **11**, 2917-2926 (2019).

50. Enault, F. *et al.* Phages rarely encode antibiotic resistance genes: a cautionary tale for
virome analyses. *ISME J.* **11**, 237–247 (2017).

51.   MM, D., PM, O. & PM, S. Exotoxins of *Staphylococcus aureus*. *Clin. Microbiol. Rev.* **13**,
16–34 (2000).

52.   Lam, M. M. C. *et al.* Genetic diversity, mobilisation and spread of the yersiniabactin-
encoding mobile element ICEKp in *Klebsiella pneumoniae* populations. *Microb.*
Genomics **4**, e000196 (2018).

53.   Pinilla-Redondo, R. *et al.* CRISPR-Cas systems are widespread accessory elements across
bacterial and archaeal plasmids. *Nucleic Acids Res.* **1**, gkab859 (2021).

54.   van Belkum, A. *et al.* Phylogenetic Distribution of CRISPR-Cas Systems in Antibiotic-
Resistant *Pseudomonas aeruginosa*. *mBio* **6**, e01796-15 (2015).

55.   Wheatley, R. M. & MacLean, R. C. CRISPR-Cas systems restrict horizontal gene transfer
in *Pseudomonas aeruginosa*. *ISME J.* **15**, 1420-1433 (2020).

56.   Pursey, E., Dimitriu, T., Paganelli, F. L., Westra, E. R. & Houte, S. van. CRISPR-Cas is
associated with fewer antibiotic resistance genes in bacterial pathogens. *Philos. Trans.*
R. Soc. B **377**, 20200464 (2022).

57.   Halary, S., Leigh, J. W., Cheaib, B., Lopez, P. & Bapteste, E. Network analyses structure
genetic diversity in independent genetic worlds. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 127–
132 (2010).

58.   Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.
Nat. Commun. **9**, 5114 (2018).

59.   Cury, J., Oliveira, P. H., de la Cruz, F. & Rocha, E. P. C. Host Range and Genetic Plasticity
Explain the Coexistence of Integrative and Extrachromosomal Mobile Genetic Elements.
Mol. Biol. Evol. **35**, 2230–2239 (2018).

60.   Rollie, C. *et al.* Targeting of temperate phages drives loss of type I CRISPR–Cas systems.
Nature **578**, 149-153 (2020).

61.   Seed, K. D., Lazinski, D. W., Calderwood, S. B. & Camilli, A. A bacteriophage encodes its
own CRISPR/Cas adaptive response to evade host innate immunity. *Nature* **494**, 489–
491 (2013).

62.   Al-Shayeb, B. *et al.* Clades of huge phages from across Earth's ecosystems. *Nature* **578**,
425-431 (2020).

63.   Liu, M. *et al.* ICEberg 2.0: an updated database of bacterial integrative and conjugative
elements. *Nucleic Acids Res.* **47**, D660-D665 (2018).

64.   Bazin, A., Gautreau, G., Médigue, C., Vallenet, D. & Calteau, A. panRGP: a pangenome-
based method to predict genomic islands and explore their diversity. *Bioinformatics* **36**,
i651–i658 (2020).

65.   Fillol-Salom, A. *et al.* Phage-inducible chromosomal islands are ubiquitous within the
bacterial universe. *ISME J.* **12**, 2114-2128 (2018).

66.   Richter, M. & Rosselló-Móra, R. Shifting the genomic gold standard for the prokaryotic
species definition. *Proc. Natl. Acad. Sci.* **106**, 19126–19131 (2009).

67.   GToTree: a user-friendly workflow for phylogenomics. Bioinformatics **35**, 4162-4164
(2019).

68.   Nguyen, L. T., Schmidt, H. A., Von Haeseler, A. & Minh, B. Q. IQ-TREE: A fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol.*
Biol. Evol. **32**, 268–274 (2015).

69. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–
2069 (2014).

70. Gautreau, G. *et al.* PPanGGOLiN: Depicting microbial diversity via a partitioned
pangenome graph. *PLOS Comput. Biol.* **16**, e1007732 (2020).

71. Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).

72. Eddy, S. R. Accelerated profile HMM searches. *PLoS Comput. Biol.* **7**, e1002195 (2011).

73. Garcillán-Barcia, M. P., Redondo-Salvo, S., Vielva, L. & de la Cruz, F. MOBscan:
Automated Annotation of MOB Relaxases. in *Methods in Molecular Biology* vol. 2075
295–308 (Humana Press Inc., 2020).

74. Akhter, S., Aziz, R. K. & Edwards, R. A. PhiSpy: A novel algorithm for finding prophages
in bacterial genomes that combines similarity-and composition-based strategies.
Nucleic Acids Res. **40**, e126–e126 (2012).

75. Zhao, X. BinDash, software for fast genome distance estimation on a typical personal
laptop. *Bioinformatics* **35**, 671–673 (2019).

76. Steinegger, M. & Söding, J. Clustering huge protein sequence sets in linear time. *Nat.*
Commun. 2018 91 **9**, 1–8 (2018).

77. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* **38**, 5825-5829 (2021).

78. Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J. Antimicrob.*
Chemother. **67**, 2640–2644 (2012).

79. Chen, L., Zheng, D., Liu, B., Yang, J. & Jin, Q. VFDB 2016: Hierarchical and refined dataset
for big data analysis - 10 years on. *Nucleic Acids Res.* **44**, D694–D697 (2016).

80. Russel, J., Pinilla-Redondo, R., Mayo-Muñoz, D., Shah, S. A. & Sørensen, S. J.
CRISPRCasTyper: Automated Identification, Annotation, and Classification of CRISPR-
Cas Loci. *Cris. J.* **3**, 462–469 (2020).

81. Shmakov, S. A. *et al.* The CRISPR spacer space is dominated by sequences from species-
specific mobilomes. *mBio* **8**, (2017).

82. Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life scale
using DIAMOND. *Nat. Methods* **18**, 366–368 (2021).

Fig 1. Distribution of MGEs across the ESKAPE pathogens. Proportion of genomes carrying at least one **A**) plasmid, **B)** ICE/IME, or **C)** prophage. **D**) Total number of MGEs and of considered genomes per ESKAPE pathogen. Size of the circles is proportional to the number of identified elements. **E)** Total number of RGPs and ICEs/IMEs per ESKAPE. The size of the green bars is proportional to the total number of ICEs/IMEs identified per ESKAPE pathogen, and the relative number of ICEs/IMEs per RGPs is shown in percentage next to the green bars. Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

A)

ICEs/IMEs

Prophages

Plasmids

B)

Acinetobacter baumannii

Enterococcus faecium

Enterobacter sp.

Klebsiella pneumoniae

Pseudomonas aeruginosa

Staphylococcus aureus

Figure 2. Network of clustered MGEs, using the mean JI as a threshold. Network grouped by **A)** MGE type; **B)** species. Each MGE is represented by a node, connected by edges according to the pairwise distances between all MGE pairs. The network has a clustering coefficient of 0.836, a density of 0.028, a centralization of 0.102, and a heterogeneity of 0.942.

Source

ICE_IME Plasmid Prophage

Category

A: RNA processing and modification
C: Energy production and conversion
D: Cell cycle control, cell division, chromosome partitioning
E: Amino acid transport and metabolism
F: Nucleotide transport and metabolism
G: Carbohydrate transport and metabolism
H: Coenzyme transport and metabolism
I: Lipid transport and metabolism
J: Translation, ribosomal structure and biogenesis
K: Transcription
L: Replication, recombination and repair
M: Cell wall/membrane/envelope biogenesis
N: Cell motility
P: Inorganic ion transport and metabolism
O: Posttranslational modification, protein turnover, chaperones
R: General function prediction only
S: Function unknown
Q: Secondary metabolites biosynthesis, transport and catabolism
T: Signal transduction mechanisms
V: Defense mechanisms
U: Intracellular trafficking, secretion, and vesicular transport
W: Extracellular structures
X: Mobilome: prophages, transposons

Figure 3. Relative contribution of the ESKAPE MGEs proteome to COG functional categories. Each ESKAPE pathogen is shown in a separate panel. The relative frequency of proteins in the different COG functional categories was calculated separately for each MGE type by dividing the number of proteins belonging to a given category by the total number of proteins observed in the corresponding MGE/ESKAPE pair. Hence, the bars illustrate the incidence of proteins of a given functional category in the proteomes of the different MGE types per ESKAPE taxon. Different COG functional categories are shown along the X axis, including: A - RNA processing and modification; C - Energy production and conversion; D - Cell cycle control, cell division, chromosome partitioning; E - Amino acid transport and metabolism; F - Nucleotide transport and metabolism; G - Carbohydrate transport and metabolism; H - Coenzyme transport and metabolism; I - Lipid transport and metabolism; J - Translation, ribosomal structure and biogenesis; K - Transcription; L - Replication, recombination and repair; M - Cell wall/membrane/envelope biogenesis; N - Cell motility; O - Posttranslational modification, protein turnover, chaperones; P - Inorganic ion transport and metabolism; Q - Secondary metabolites biosynthesis, transport and catabolism; R - General function prediction only; S - Function unknown; T - Signal transduction mechanisms; U - Intracellular trafficking, secretion,

and vesicular transport; V - Defense mechanisms; W - Extracellular structures; X - Mobilome: prophages, transposons. COG functional categories B, Y, and Z (standing for chromatin structure and dynamics, nuclear structure, and cytoskeleton, respectively) were not identified in the dataset. Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Fig 4. Distribution of widespread protein clusters across MGE/ESKAPE pairs. The heatmaps show the distribution and relative frequency of protein clusters (columns) identified as widespread in either ESKAPE pathogens (i.e. present in at least three different ESKAPE; left side) or MGEs (i.e. present at least once in the three MGE types; right side). MGE/ESKAPE pairs are indicated on the left side of the heatmaps. The relative frequency of the protein clusters in the different MGE/ESKAPE pairs was calculated by dividing the number of occurrences identified by the total number of proteins observed in a given pair. The trees displayed at the top and left side of the heatmaps illustrate the hierarchical clustering of protein clusters and MGE/ESKAPE pairs using the ward.D method with euclidean distance. A list of the protein clusters, their products and relative frequency values is provided in the (**Table S3**). Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure 5. Relative proportion of AMR, virulence, and Anti-CRISPRs between MGEs and masked genomes across the ESKAPE pathogens. The number of AMR, virulence, or Anti-CRISPRs proteins found for MGEs or masked genomes per ESKAPE was normalized to the total number of proteins found for each element per ESKAPE pathogen.

Figure 6. Distribution of CRISPR-Cas and Anti-CRISPR across the ESKAPE mobilome. **A)** Proportion of CRISPR-Cas positive genomes across the ESKAPE pathogens. **B)** Distribution of CRISPR-Cas systems within ICEs/IMEs, plasmids, and masked genomes. Prophages are not shown because we found no CRISPR-Cas system in these elements. **C)** Distribution of Anti-CRISPR proteins across ICEs/IMEs, plasmids, and prophages. Even though we found Anti-CRISPRs on masked genomes, we only plotted those found on MGEs. A complete list of Anti-CRISPRs is given in **Table S6**. Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure 7. Boxplots comparing the number of MGEs present in pairs of conspecific ESKAPE pathogens, with and without CRISPR-Cas systems. Values above 0.05 were considered as non-significant (ns). We used the following convention for symbols indicating statistical significance: * for p <= 0.05, ** for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001. Ab, *A. baumannii*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure 8. CRISPR spacers are involved in MGE-MGE conflict. **A)** CRISPR spacers found on ICEs/IMEs (left side) and plasmids (right side), and their interactions with protospacers identified in ESKAPE MGEs. **B)** Boxplots comparing the sequence length of MGEs targeted by CRISPR spacers found in MGEs. **C)** Boxplots comparing the sequence length of MGEs targeted by CRISPR spacers found in masked genomes. Values above 0.05 were considered as non-significant (ns). We used the following convention for symbols indicating statistical significance: * for p <= 0.05, ** for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001. Ab, *A. baumannii*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure S1. Phylogenomic tree, representing type strains belonging to the *Enterobacteriaceae* family and all complete genomes classified as *Enterobacter*.

Genomes that are positioned in the non-*Enterobacter* cluster, and poorly classified as belonging to the *Enterobacter* genus, were highlighted in green.

Figure S2. Variation in GC content between MGEs and masked genomes. Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*. Values above 0.05 were considered as non-significant (ns). We used the following convention for symbols indicating statistical significance: * for p <= 0.05, ** for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001.

Figure S3. Variation in sequence length between MGEs. Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*. Values above 0.05 were considered as non-significant (ns). We used the following convention for symbols indicating statistical significance: * for p <= 0.05, ** for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001.

Figure S4. Scatter plot between the different MGEs' sequence length and GC content. Regression lines for each MGE type are shown in different colours. Confidence interval is displayed around smooth. Level of confidence interval is 0.95.

A

B

Figure S5. Density plot showing the distribution of Jaccard index (JI) values for the **A)** MGE and **B)** masked genomes network. The mean distribution for the JI is shown in blue, while the median is shown in green. In A), density values from 0-10 were zoomed in and shown in the left.

Figure S6. Network of clustered MGEs, using the median JI as a threshold. Network grouped **A)** MGE type; **B)** species. Each MGE is represented by a node, connected by edges according to the pairwise distances between all MGE pairs. The network has a clustering coefficient of 0.783, a density of 0.042, a centralization of 0.168, and a heterogeneity of 1.088.

Figure S7. Network of clustered masked genomes grouped by species, using the mean JI as a threshold. Each masked genome is represented by a node, connected by edges according to the pairwise distances between all masked genome pairs.

Figure S8. Chord diagram showing DNA sharing events between different ESKAPE pathogens. The width of sectors represents total number of inferred interactions between two different MGE/ESKAPE pairs. The width of links are proportional to the number of inferred DNA sharing events. Interactions between the same ESKAPE/MGE pairs were excluded. Interactions between the same ESKAPE, but different MGEs, were also removed. Interactions between *K. pneumoniae* and *Enterobacter* sp. were also removed, since interactions between these species is very common, as both belong to the *Enterobacteriaceae* family. Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

ICEs_IMEs (713)   **Plasmids (876)**

Prophages (262)

Figure S9. COGs identified in ESKAPE MGEs. Venn diagram showing the distribution of 1183 different COGs identified in ESKAPE ICEs/IMEs, plasmids and phages. The number of different COGs found for each MGE type is indicated in parenthesis. Note that the numbers displayed in the figure do not correspond to the number of proteins with an identifiable COG in MGE sequences but to the distinct COGs detected in the MGEs proteome.

Figure S10. Distribution of antimicrobial resistance (AMR) genes across different ESKAPE pathogens and mobile elements. The size of the circles is proportional to total counts. Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure S11. Proportion of mobile genetic elements (MGEs) carrying antimicrobial resistance (AMR) genes, identified in **A)** plasmids and **B)** ICEs/IMEs, and virulence (Vir) genes identified in **C)** plasmids and **D)** ICEs/IMEs. A separate proportion is shown for the different ESKAPE pathogens. Each MGE was considered as positive for AMR or Vir if at least one gene was identified. Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure S12. Distribution of virulence genes across different ESKAPE pathogens and mobile elements. The size of the circles is proportional to total counts. Ab, *A. baumannii*; Ef, *E. faecium*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure S13. Boxplots comparing the sequence length in pairs of conspecific ESKAPE pathogens, with and without CRISPR-Cas systems. Values above 0.05 were considered as non-significant (ns). We used the following convention for symbols indicating statistical significance: * for p <= 0.05, ** for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001. Ab, *A. baumannii*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure S14. Boxplots comparing the GC content in pairs of conspecific ESKAPE pathogens, with and without CRISPR-Cas systems. Values above 0.05 were considered as non-significant (ns). We

used the following convention for symbols indicating statistical significance: * for p <= 0.05, **
for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001. Ab, *A. baumannii*; En, *Enterobacter*
sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure S15. Boxplots comparing CRISPR-Cas positive and negative *P. aeruginosa* genomes, with
GC content normalized to sequence length of each genome. Values above 0.05 were considered
as non-significant (ns). We used the following convention for symbols indicating statistical
significance: * for p <= 0.05, ** for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001.

Figure S16. Boxplots comparing the number of virulence genes present in pairs of conspecific ESKAPE pathogens, with and without CRISPR-Cas systems. Values above 0.05 were considered as non-significant (ns). We used the following convention for symbols indicating statistical significance: * for p <= 0.05, ** for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001. Ab, *A. baumannii*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure S17. Boxplots comparing CRISPR-Cas positive and negative *P. aeruginosa* genomes, with virulence genes normalized to sequence length of each genome. Values above 0.05 were considered as non-significant (ns).

Figure S18. Boxplots comparing pairs of conspecific ESKAPE pathogens, with and without CRISPR-Cas systems, regarding MGE-encoded **A**) antimicrobial resistance (AMR) and **B**) virulence genes. Values above 0.05 were considered as non-significant (ns). We used the following convention for symbols indicating statistical significance: * for p <= 0.05, ** for p <= 0.01, *** for p <= 0.001, and **** for p <= 0.0001. Ab, *A. baumannii*; En, *Enterobacter* sp.; Kp, *K. pneumoniae*; Pa, *P. aeruginosa*; Sa, *S. aureus*.

Figure S19. Total number of CRISPR spacers per masked genome identified across the ESKAPE pathogens.