Got Milk? Maternal immune activation during the mid-lactational period affects nutritional milk quality and adolescent offspring sensory processing in male and female rats

Holly DeRosa[1], Salvatore G. Caradonna[2], Hieu Tran[1], Jordan Marrocco[2,3], Amanda C. Kentner[1]*

[1]School of Arts & Sciences, Health Psychology Program, Massachusetts College of Pharmacy and Health Sciences, Boston Massachusetts, United States 02115.
[2]Laboratory of Neuroendocrinology, The Rockefeller University, 1230 York Ave, New York, NY 10065.
[3]Department of Biology, Touro University, 227 W 60th St, New York, NY 10023.

*Corresponding author:
Amanda Kentner
amanda.kentner@mcphs.edu
Office #617-274-3360
Fax # 617-732-2959

Keywords
Maternal immune activation; lactation; nutrition; prepulse inhibition; mechanical allodynia; conditioned fear; microbiome; RNA sequencing; transcription; milk; breastfeeding; animal models; early life stress

Word Counts:
Abstract: 250
Main Text: 3995
Figures: 6
Supplementary Methods: 1
Supplementary Figures: 1
Supplementary Tables: 3 + Key Resources Table

Abstract

Background: Previous studies have underscored the importance of breastfeeding and parental care on offspring development and behavior. However, their contribution as dynamic variables in animal models of early life stress are often overlooked. **Methods:** In the present study, we investigated how lipopolysaccharide (LPS)-induced maternal immune activation (MIA) on postnatal day (P)10 affects maternal care, milk, and offspring development. **Results:** MIA was associated with elevated milk corticosterone concentrations on P10, which recovered by P11. In contrast, both milk triglyceride and percent creamatocrit values demonstrated a prolonged decrease following inflammatory challenge. Adolescent MIA offspring were heavier, which is often suggestive of poor early life nutrition. While MIA did not decrease maternal care quality, there was a significant compensatory increase in maternal licking and grooming the day following inflammatory challenge. However, this did not protect against disrupted neonatal huddling or later-life alterations in sensorimotor gating, conditioned fear, mechanical allodynia, or reductions in hippocampal parvalbumin interneuron expression in MIA offspring. MIA-associated changes in brain and behavior were likely driven by differences in milk nutritional values and not by direct exposure to LPS or inflammatory molecules as neither LPS binding protein nor interleukin-6 milk levels differed between groups. These findings reflected comparable microbiome and transcriptomic patterns at the genome-wide level. **Conclusions:** Animal models of early life stress can impact both parents and their offspring. One mechanism that can mediate the effects of such stressors is changes to maternal lactation quality which our data show can confer multifaceted and compounding effects on offspring physiology and behavior.

Introduction

Epidemiological studies have shown that prenatal exposure to maternal immune activation (MIA) is a leading risk factor for developing psychiatric disorders such as autism and schizophrenia (Mahic et al., 2017; Meyer, 2019). Administrating immunogens such as lipopolysaccharide (LPS) or polyinosinic:polycytidylic acid (poly I:C) to pregnant animals induces MIA by promoting a cascade of inflammatory cytokines and activating the hypothalamic-pituitary adrenal (HPA) axis of the mother (Kentner et al., 2019; Brown et al., 2009; Arsenault et al., 2014; Estes & McAllister, 2016). Together, these physiological responses are recognized to affect the long-term development of offspring (Kentner et al., 2019; Estes & McAllister, 2016). However, the impact on mothers and their offspring when MIA occurs during lactation is relatively unexplored. This is surprising because infections during the postnatal period are common, particularly in low resource settings, and can result in severe morbidity and mortality (Karolinski et al., 2010; Kutlesic et al., 2017; John et al., 2017; Hussein et al., 2011).

LPS administered to rodent dams during the lactational period can impact maternal care by reducing the amount of time spent in efficient arched back nursing postures and the number of pups retrieved and returned back to the nest when displaced (Vilela et al., 2013). In line with the behavioral alterations, MIA in lactating dams may influence maternal milk quality. For example, maternal LPS treatment on postnatal days 4, 11, and 18 reduced the mRNA expression of milk nutrient precursors such as glucose and fatty acid transporters in the rat mammary gland (Ling & Alcorn, 2010). Exposure to MIA during the lactational period can also impact offspring behavior. For example, Nascimento and colleagues (2015) observed a reduction in ultrasonic vocalizations in male neonates shortly after their dams were treated with LPS. Although this presumably involved changes in various forms of

maternal input signals (e.g., parental care and/or milk quality), the mechanisms of this MIA-derived effect on offspring behavior remain unclear.

Other exogenous factors have been shown to alter contents in breastmilk and subsequent offspring development in animals and humans (Lesorogol et al., 2018; Mäkelä et al., 2013; Chen et al., 2017; Grazia Di Benedetto et al., 2020; Edwards et al., 2021). For example, psychological stress in humans reduces milk microbial diversity (Browne et al., 2019). Further, cortisol in maternal milk was found to predict temperament and promote greater body weights in infant primates, although milk energy content was also associated with body weight (Hinde et al., 2014). Together, this evidence suggests that 1) disturbances to the maternal-neonatal environment alters the developmental trajectory of offspring, and 2) these alterations are mediated by changes in maternal care and/or nutritional milk quality. To explore these ideas, we assessed the effects of MIA challenge during the mid-lactational period on maternal care, maternal milk composition, and offspring brain development and behavior in Sprague-Dawley rats.

Methods and Experimental Overview

All animal procedures were conducted in accordance with the Association for Assessment and Accreditation of Laboratory Animal Care with protocols approved by the Massachusetts College of Pharmacy and Health Sciences (MCPHS) Institutional Animal Care and Use Committee. **Figure 1A** outlines the experimental procedures and groups evaluated in this study. Please see the **Supplementary Methods, Supplementary Table 1,** and **Key Resources Table** for a more detailed description of the methodological protocols and statistical analyses used in this study. Briefly, on postnatal day 10 (P10), female Sprague-Dawley rat mothers were removed from their litters and placed into a clean cage located in a separate procedure room. Dams were challenged intraperitoneally (i.p.) with either 100 µg/kg of the inflammatory endotoxin, lipopolysaccharide (LPS) or pyrogen-free saline. Milk samples were collected 2 hours later at the height of the inflammatory response and 24 hours later on P11 during the recovery period (see DeRosa et al., 2022).

Milk quality was determined through the use of several assays to measure levels of milk creamatocrit, lactose, triglycerides, protein, corticosterone, immunoglobulin (Ig) A, interleukin (IL)-6 and lipopolysaccharide binding protein (LBP). Milk quality was also more broadly examined using microbiome and RNA Sequencing (RNA-seq). Maternal behavior was assessed in the afternoon and evening of P10 and in the morning on P11. The offspring of saline and MIA treated dams completed an assessment of neonatal huddling, a form of social thermoregulation, 2 hours after being reunited with their mother on P10 and P11. Starting on P43, one male and one female offspring from each litter completed several behavioral measures including the open field and social preference tasks, as well as assessments of mechanical allodynia, sensorimotor gating (prepulse inhibition; PPI), and conditioned fear (see Yan & Kentner, 2017; Connors et al., 2014; Zhao et al., 2021a; Núñez Estevez et al., 2020). On P46, whole hippocampus was collected and later processed to assess the expression of parvalbumin (PV) interneurons using western blotting.

Two-way repeated measure ANOVAs (MIA x Time) were used to evaluate milk and behavioral measures across P10 and P11. Violations to the assumption of sphericity were addressed using the Huyndt Feldt correction. One-way ANOVAs were used as appropriate for all other measures unless there were violations to the assumption of normality (Shapiro-Wilk test) in which case Kruskal-Wallis tests were employed (expressed as X^2). The von Frey threshold data was assessed using body weight as a covariate. The partial eta-squared (n_p^2) is reported as an index of effect size for the ANOVAs (Miles and Shevlin, 2001). Because the dataset was not powered to evaluate sex-differences directly, male and female animals were

evaluated separately (Ordoñes Sanchez et al., 2021). Data are depicted as each sex separately and collapsed across males and females for display purposes.

For microbiome sequencing, samples underwent composition visualization, in addition to alpha-diversity and beta-diversity analyses using Qiime (v.1.9.1) and statistical comparisons were performed using Kruskal-Wallis (Caporaso et al., 2010). Linear discriminant analysis effect size (LEfSe; http://huttenhower.sph.harvard.edu/lefse/) was utilized to determine significant differences in taxonomy abundance between each group as previously described (Segata et al., 2011; Schellekens et al., 2021).

Results

MIA challenge affected maternal behavior and offspring huddling

Immune challenge during lactation significantly impacted the number of maternal care bouts directed towards pups. For example, a MIA by time interaction was identified for the number of licks pups received following maternal LPS exposure (**Figure 1B**). While the number of licks received was not significantly affected on P10, there was a general pattern of reduced care provided by MIA treated dams on the afternoon of P10. This was followed by a rebound of maternal pup licking by MIA dams on the morning of P11, possibly to compensate for overall reduced maternal care due to illness. The display of active and low crouch nursing behaviors and time on nest were affected by the circadian cycle (**Figure 1C,D,E**), but only time on nest was impacted by MIA experience. Maternal self-directed care was impacted as a function of sickness. There was a significant MIA by time interaction for total number of self-grooms (**Figure 1F**); MIA dams had fewer bouts of grooming on the evening of P10, which reversed substantially by the morning of P11. Similarly, the number of maternal eating bouts was elevated on P11, although there were no significant effects of time (**Figure 1G**) there was a main effect of MIA (**Figure 1G**). Additional maternal care data are presented in **Supplementary Figure 1A-D**.

Huddling behavior was significantly reduced in the offspring of MIA dams compared to saline (main effect of MIA; time spent huddling, **Figure 1H**; average number of clusters: **Figure 1I**). See **Figure 1J** for representative displays of huddling behavior in saline and MIA offspring.

MIA challenge impacted nutritional composition of milk and microbiome communities

For milk collection (see photographs in **Figure 2A** depicting the milk collection procedure), there was a significant MIA by time interaction for maternal milk levels of corticosterone (**Figure 2B**). This stress-associated hormone was elevated in milk samples 3-hours following MIA challenge (**Figure 2B)** and was recovered by P11. A MIA by time interaction was also present for % creamatocrit (**Figure 2C**), which is linearly related to the fat concentration and energy content of milk (Lucas et al., 1978; Paul et al., 2015; see **Supplementary Figure 2A, B**). These measures were significantly lower in the milk of MIA exposed mothers compared to saline on P11. Triglycerides were also reduced in the milk of MIA treated dams on P11 (**Figure 2D**).

In order to examine the possibility that MIA challenge increased expression inflammatory cytokines that could be transferred to nursing offspring, we analyzed the concentration of milk IL-6. While there were no significant differences in IL-6 concentration between saline and MIA treated dams, there was a main effect of time on this measure (**Figure 2E).** We also tested the idea that bound LPS could cross into maternal milk and be absorbed by offspring. LPS binds directly to LBP which triggers a downstream pro-inflammatory response and the release of acute phase proteins like IL-6 (Wurfel & Wright, 1997; Park & Lee, 2017). LBP

levels were measured in blood plasma and P10 milk samples. LPB levels were significantly greater in the blood plasma of MIA dams (**Figure 2F**). In milk, LPB was absent in both MIA and saline dams. Additional data on maternal milk composition (i.e., protein, lactose, and IgA concentrations) can be found in **Supplementary Figure 2C-E.**

Microbiome sequencing revealed no main effect of treatment group on alpha (**Figure 2G**) or beta diversity in milk samples (**Figure 2H, I**). Milk samples collected on P11 demonstrated significantly greater alpha diversity along the Shannon index compared to samples collected on P10 (**Figure 2G**). Between treatment groups, LEfSE analysis identified nine differently abundant taxa on P10 and six significantly abundant taxa on P11 (**Figure 2J, K**). These analyses revealed a greater abundance of *Pseudomonadaceae* and *Christensenellaceae* on P10, and a greater expression of *Stenotrophomonas maltophilia*, *Ruminococcaceae* and *Lachnospiraceae* on P11 in the milk from MIA dams. For additional data on the composition of the milk microbiome, see **Supplementary Figures 3-6.**

RNA-seq of the milk was performed to characterize the effects of mid-lactational MIA at the genome-wide level. A total of 829 genes were differently expressed on P10 between saline and MIA milk samples (**Figure 3A**). We analyzed clusters of gene networks that are involved in maternal milk nutrition, immune activation, and offspring brain development (Quintana et al., 2019; Oakley & Cidlowski, 2013; Polman et al., 2012; Zhu et al., 2016; Ozkan et al., 2020; Zhang et al., 2013; Bowen et al., 2013; Borish & Steinke, 2003; **Figure 3B).** Heatmap clustering revealed a general upregulation of genes related to nutrient transport and inflammation, while genes related to epigenetic modifications, parvalbumin interneuron development, and glucocorticoid signaling were largely downregulated by MIA on P10 (**Figure 3B**). In P11 samples (**Figure 4A,B),** a total of 893 genes were differentially affected between the milk of saline and MIA dams. Heat clustering analysis of these samples demonstrated a general downregulation in genes responsible for inflammation, glucocorticoid signaling, epigenetic modifiers, and prolactin in MIA samples, while genes related to oxytocin, nutrient transport, triglycerides, and parvalbumin interneuron development were largely upregulated. For more information on the genes indicated in the heatmap clustering, see **Supplementary Tables 2 and 3.**

The combination of MIA challenge and nutritional deficit affected adolescent offspring physiology and behavior.

Male, but not female MIA offspring were significantly heavier than saline controls on P43 (**Figure 5A**). Using body weight as a covariate, female offspring from MIA dams were found to have higher thresholds in the von Frey test compared to their saline counterparts (**Figure 5B**). This is suggestive of reduced sensitivity to tactile stimulation.

Although there were no differences in distance traveled (cm), percent time spent in the center, duration of time (seconds) spent in the center, or perimeter of the open field arena (see **Supplementary Figure 7A-D**), male offspring of MIA exposed mothers displayed an increased number of crosses into both the center (**Figure 5C**) and perimeter (**Figure 5D**) of the arena. Social preference was not affected by the early life experience for either sex (**Figure 5E**).

Sensorimotor gating was evaluated using PPI of the acoustic startle reflex. A significant MIA by time interaction (**Figure 5F**) and a significant main effect of time (**Figure 3G, H**, collapsed across males and females for display purposes) confirmed that all groups demonstrated an increased % PPI as the intensity was raised from 69 to 85 dB. While females were unaffected, male offspring originating from MIA exposed mothers had attenuated PPI values compared to saline controls at the 69dB, 73dB, and 81dB intensities. The mean percent prepulse inhibition across all five prepulse intensities was not affected as a function of maternal MIA exposure (**Figure 5I**).

With respect to conditioned fear, there was a main effect of MIA for % PPI during the trials without fear (**Figure 5J, K, L).** % PPI for trials containing the conditioned stimulus was not significantly different between saline and MIA animals for either sex (**Supplementary Figure 7E**, collapsed across males and females for display purposes). One-way ANOVA revealed a main effect of trial type (**Supplementary Figure 7E**), which was expected given that the added element of fear increases % PPI (Ishii et al., 2010; Balogh et al., 2002). Conditioned fear did not lead to significant changes in total mean % PPI when collapsed across all dB, regardless of MIA or sex (**Supplementary Figure 7F**, data collapsed across males and females for display purposes), suggesting that fear modulated % PPI similarly regardless of sex or MIA exposure.

In addition to these behavioral effects, PV/β-actin ratios were significantly reduced in the hippocampus of male MIA offspring compared to saline dams (**Figure 5M**).

Discussion

We demonstrate that variations in milk quality may be part of a mechanism that programs offspring physiology and behavior following stress exposure during the lactational period (see **Figure 6** for summary of proposed mechanisms). In the present study, MIA was associated with sustained changes in both the nutritional content and microbial profile of maternal milk. Moreover, the MIA-induced changes in milk fat and corticosterone levels were accompanied by physiological and behavioral alterations in offspring. For example, neonatal offspring from MIA-challenged dams spent less time huddling compared to controls. In adolescence, MIA offspring were heavier, demonstrated deficits in sensorimotor gating, threat detection, hippocampal PV expression, and had increased mechanical allodynia thresholds. Importantly, these MIA-associated effects in offspring were unlikely due to an overexposure to inflammatory cytokines. Instead, offspring behavior and bodyweight differences likely reflected the unique expression of the nutritional elements and metabolically relevant bacterial taxa found in P10 and P11 milk samples. While most of the MIA-related literature has been dedicated to uncovering the physiological and behavioral effects on offspring gestationally exposed to MIA, fewer studies have focused on the impact of MIA exposure after birth. The lactational period is an important time for neonates and here we underscore the value of maternal inputs (e.g., maternal milk quality) to offspring development.

MIA was associated with multiple changes in the composition of maternal milk content. We observed a significant increase in milk corticosterone 3-hours post LPS treatment that remitted by P11. Milk corticosterone is expressed linearly to plasma corticosterone (Patacchioli et al., 1992), validating our MIA model. Importantly, milk corticosterone can pass from a mother to her offspring where these glucocorticoids can remain active in the periphery and in the brain (Angelucci et al., 1985; Brummelte et al., 2010). Increased corticosterone exposure from breastmilk has previously contributed to HPA axis dysfunction as well as altered learning and memory abilities in rat pups of both sexes, independent of maternal care (Catalani et al., 1993; Catalani et al., 2002). In addition to its modulatory actions within the HPA axis, recent evidence has demonstrated the regulatory role of corticosterone in breastmilk, where the presence of this hormone (and the human equivalent, cortisol) can modulate nutritional elements including sodium, potassium (Zietek et al., 2021; Sullivan et al., 2011) and fat (Hinde et al., 2015). This evidence suggests that the temporary rise in corticosterone on P10 may have preceded downstream changes in milk nutritional quality. In line with this, MIA reduced % creamatocrit and milk triglycerides on P11. However, higher levels of glucocorticoids in milk alone are not a sufficient predictor of

offspring bodyweight (Brummelte et al., 2010; Hinde et al., 2015), suggesting that the actions of corticosterone in neonates are likely interwoven with milk nutrition.

Microbiome sequencing did not reveal significant differences in alpha or beta diversity between saline and MIA treated dams. While general diversity was not affected, LEfSe analysis uncovered significant differences in the expression of certain taxa on P10 and P11. Specifically, MIA dams exhibited a higher abundance of bacterial families including *Pseudomonadaceae*, which was previously shown to be over expressed in the ceca of pregnant mice with low serum triglycerides (Connor et al., 2018). Further, milk from MIA dams also exhibited elevated levels of *Stenotrophomonas maltofilia*, which has the ability to degrade hydrocarbon chains and therefore could prevent triglyceride synthesis (Singh & Fulekar et al., 2010). Given that the microbes in breast milk directly colonize the infant gut (Sheard & Walker, 1988; Pannaraj et al., 2017), it is likely that these differently expressed taxa are being passed to the offspring to influence future metabolic processes. Additional studies are needed to explore potential therapeutic interventions, such as targeting the microbiome (Holingue et al., 2020) which may rescue changes in milk composition and subsequent offspring developmental outcomes.

Differences in microbial and nutritional markers paralleled MIA-mediated changes found in the whole-transcriptome of the milk using RNA-seq. Notably, we observed a significant upregulation in aldosterone-promoting *Ace* and a downregulation of the mineralocorticoid receptor-coding *Nr3c2* in MIA milk on P10 (**Figure 3B, Supplementary Table 2**). These changes, coupled with the increase in P10 milk corticosterone, corroborated differences observed in milk nutrition, given that aldosterone is another crucial mediator of breastmilk nutrition and ion exchange (Zietek et al., 2021). These results were further supported by the differential genomic profiles between milk from MIA and saline dams for gene clusters related to triglycerides and nutrient transport. RNA-seq also revealed a general upregulation of genes involved in the inflammatory response in maternal milk on P10 in MIA-treated dams. Although LBP and IL-6 were equally regulated between saline and MIA dams, the possibility that other inflammatory cytokines may be active in milk and subsequently affect offspring development cannot be entirely ruled out. While there was an unexpected significant upregulation of several genes within the inflammation pathway on P11 (**Figure 4B, Supplementary Table 3**), this may be reflective of the simultaneous downregulation of epigenetic modifiers since cytokine release is often associated with aberrant epigenetic activity (Obata et al., 2015). Overall, these data demonstrate the extensive effects of MIA on maternal milk at the genome-wide level. Future investigations are needed to understand the action of these transcriptomic profiles within offspring after they are absorbed.

We used several behavioral measures to explore the role of lactational MIA exposure on offspring development. Neonatal 'huddling' is described as a form of social thermoregulation (Gilbert et al., 2009). One previous study found a reduction in nest seeking behavior in rat pups prenatally exposed to LPS (Baharnoori et al., 2012). Undernourishment has also been shown to reduce neonatal huddling in rat pups (Soriano et al., 2006), and huddling efficiency may be related to serum triglycerides in newborn rabbits (García-Torres et al., 2015). This evidence suggests that MIA-modulated huddling may emerge due to differences in the fat content of maternal milk. Indeed, pups from MIA-treated mothers displayed disorganized clumping behavior.

Prenatal MIA has been shown to modulate offspring performance on certain tasks, such as the open field, social preference, von Frey, PPI, and fear conditioning paradigms (Kentner et al., 2019; Fortier et al., 2007; Zhao et al., 2021a; Inceoglu et al., 2006; Howland et al., 2012). Here, we observed significant MIA induced reductions in % PPI that were exacerbated in male adult offspring while females demonstrated deficits in mechanical allodynia. MIA was also associated with a significant reduction in % PPI in trials without fear, following the foot

shock and conditioned stimulus (CS) pairing. These findings suggest MIA contributes to an impairment in the offspring's ability to discriminate between "safe" trials (without CS presentation) and "stressful" trials (with CS presentation). Early life stress has been shown to impair this form of contextual threat detection in adolescence, and this is especially true for females (Manzano-Nieves et al., 2018; Gerhard et al., 2020). These results, in addition to reduced neonatal huddling, point to generally impaired sensory processing and threat responses in the offspring of dams exposed to MIA during lactation.

We measured the expression of GABAergic PV interneurons in the offspring hippocampus since altered PV expression in this region is a common theme in prenatal MIA models and is associated with many MIA-associated behavioral and sensory processing changes (Zhao et al., 2021b; Nakamura et al., 2021; Vojtechova et al., 2021). Further, poor nutrition in early life can prevent the development of PV interneurons in rat hippocampus (Callahan et al., 2013) although this has only been explored in males. The emergence of neonatal huddling behavior is time-locked to the maturation of GABAergic signaling (Naskar et al., 2019). Therefore, mid-lactational MIA may disrupt this early form of social behavior by attenuating the development of PV cells. In adolescence, male MIA offspring demonstrated a reduced expression in hippocampal PV interneurons. In vivo inhibition of PV in the hippocampus has previously been shown to reduce PPI in male mice (Nguyen et al., 2014). Together, these results demonstrate one neuronal target by which MIA during mid-lactation contributes to lasting changes in offspring brain development in males. More work is needed to discern the neuronal targets of mid-lactational MIA in females.

In our study, MIA dams showed significant increases in pup-directed licking and grooming behavior the day following MIA challenge. The heightened maternal care demonstrated by MIA dams on P11 may have been a compensatory mechanism to buffer the effects of lactational MIA exposure. Tactile stimulation through the licking and grooming of pups is one form of maternal behavior that promotes brain development and modulates future offspring behavior (Champagne et al., 2003; Francis et al., 2002; Van Hasselt et al., 2012; Francis & Kuhar, 2008). Aside from a general pattern of decreased (but not significant) licking and grooming of pups on the afternoon of P10, we did not observe a drastic reduction in pup directed behaviors in MIA dams. The effects of LPS are highly dose dependent (Bison et al., 2009). While Vilela and colleagues (2013) observed more dramatic deficits in maternal care following MIA, they implemented a dose of LPS 5-times higher than the dose used in the present study. Moreover, lactational exposure to LPS using the same dose as that used in the current study also resulted in improved maternal care (Nascimento et al., 2015). Together, the results of our study suggest that stressors experienced during the neonatal period may interact with maternal care and lactation quality, affecting offspring physiology and behavior. Indeed, early life stress models affect both the parent(s) and the offspring; the dynamic impacts on each need to be considered as part of the mechanistic programming of the developmental trajectory.

Conclusions

Maternal care and milk quality are important components to offspring development, brain health, and behavior. Here, we demonstrate that an acute maternal immune response during mid-lactation is sufficient to trigger changes in breastmilk quality and its microbiome profile, in addition to the physiology and behavior of adolescent offspring. These results contribute to the broader literature by suggesting that the effects of MIA on offspring development are not restricted to the prenatal window. Moreover, this study ties in the characteristics of lactation and nutrition as part of a mechanism contributing to the trajectory of offspring development following early life stress. Given that animal models of early life stress can impact both

parents and their offspring, the quality of maternal milk should be considered among the variables investigated in future studies. Finally, this work underscores the importance of research focused on potential therapeutic interventions (e.g., nutritional lactation supplements) and necessitates a better representation of pregnant and nursing people to aim for increased equity and inclusivity in both basic and clinical research.

Funding and Disclosures
This project was funded by NIMH under Award Number R15MH114035 (to ACK). The authors wish to thank Mary Erickson, Sahith Kaki, Ada Cheng, Dr. Mattia Migliore, and Dr. Theresa Casey for their technical support in the early phases of this project. The authors would also like to thank the Massachusetts College of Pharmacy and Health Sciences Schools of Pharmacy and Arts & Sciences for their continual support, the Bioinformatic Resource Center at the Rockefeller University where the RNA-seq was performed, and the University of Massachusetts Boston where HD is a graduate student. Figure 6 made with BioRender.com. The content is solely the responsibility of the authors and does not necessarily represent the official views of any of the financial supporters.

Author Contributions
H.D., S.G.C., H.T., and A.C.K., ran the experiments; H.D., S.G.C., J.M., & A.C.K. analyzed and interpreted the data; H.D. and A.C.K. wrote the manuscript; A.C.K., designed and supervised the study.

Disclosure of Competing Interest
The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Figure Captions

Figure 1. Maternal care and offspring huddling behaviors following maternal immune activation (MIA) during the mid-lactational period. (A) Timeline of experimental procedures. Maternal care was evaluated in the afternoon and night of postnatal day (P) 10 and the morning of P11. (B) MIA x Time interaction for the total number of pup licks ($F_{(2, 26)} = 31.435$, $p = 0.001$, $n_p^2 = 0.701$). Pup licking and grooming were elevated in MIA dams on P11 ($p = 0.001$). The circadian cycle influenced the display of (C) active nursing bouts ($F_{(2, 26)} = 7.752$, $p = 0.0022$, $n_p^2 = 0.374$), which were elevated in both groups on P10. (D) Low crouch nursing bouts ($F_{(2, 26)} = 3.960$, $p = 0.032$, $n_p^2 = 0.233$), and (E) total time (seconds) that dam spent on nest were also affected by the circadian cycle, as dams spent less time in the low crouch nursing posture and overall less time on the nest during the evening on P10. Total number of (F) self-groom bouts demonstrated a MIA x time interaction ($F_{(1.322, 26)} = 12.33$, $p = 0.001$, $n_p^2 = 0.485$); Huyndt Feldt correction. MIA was associated with a reduction in self-grooming bouts on the evening of P10 ($p = 0.025$), which were increased on the morning of P11 ($p = 0.0001$). A main effect of MIA was observed for (G) the number of eating bouts ($F_{(1, 13)} = 10.64$, $p = 0.006$, $n_p^2 = 0.450$). (H) Total time pups spent huddling (seconds) was reduced by MIA ($F_{(1, 12)} = 30.079$, $p = 0.001$, $n_p^2 = 0.715$) as was the (I) average number of pup clusters across P10 and P11 ($F_{(1, 12)} = 8.33$, $p = 0.014$, $n_p^2 = 0.410$). (J) Shows representative photographs of the huddling behavior of saline and MIA litters. Data are expressed as mean ± SEM. n = 7-8 for all measures. *p < 0.05, ***p <0.001, MIA versus Saline; [a]p < 0.05, [aa]p < 0.01, main effect of time; [b]p < 0.05, [bb]p < 0.01, [bbb]p < 0.001, main effect of MIA.

Figure 2. Nutritional profile of milk and microbiome community distribution following maternal immune activation (MIA) during the mid-lactational period. (A) Photographs of maternal milk collection. A significant time interaction was observed for levels of maternal milk levels of (B) corticosterone (ng/mL) ($F(1, 13) = 8.496$, $p = 0.013$, $n_p^2 = 0.415$), which was elevated in the milk of MIA dams on P10 ($p = 0.009$). A MIA by time interaction was observed for (C) percent (%) creamatocrit ($F(1, 13) = 5.86$, $p = 0.032$, $n_p^2 = 0.858$), which was significantly lower in MIA samples on P11 ($p = 0.002$) along with (D) milk triglycerides (mg/dL) ($F(1, 13) = 6.496$, $p = 0.026$, $n_p^2 = 0.351$). There was a main effect of time on (E) milk interleukin (IL)-6 (pg/mL) concentrations ($F(1,13) = 33.01$, $p < 0.001$, $n_p^2 = 0.717$), in that IL-6 was increased in both groups on P11. Levels of (F) LPS binding protein (LPB) (ug/mL) were increased in the blood plasma of MIA dams ($X^2(1)=5.00$, $p=0.025$), but there were negligible levels in maternal milk ($p>0.05$). (G) Alpha diversity of P10 and P11 milk samples along the Shannon Index was increased in both groups on P11 compared to P10 ($X^2(1) = 5.33$, $p = 0.021$). (H) Beta diversity using principle coordinate analysis (PCoA) of P10 and (I) P11 milk samples were created using the matrix of pair-wise distance between samples determined by Bray-Curtis dissimilarity using unique amplicon variants. (J) Microbial composition of taxonomy at the family level for saline and MIA dams at P10 and P11 ($n = 6$-7). (K) LEfSe biomarkers plots. These figures display taxa that are significantly more abundant in the milk of saline-treated dams (blue bars) versus MIA-treated dams (red bars) on P10 and P11. These taxa were identified based on their significant distributions ($p < 0.05$) and effect sizes (LDA score) larger than 2 ($n = 6$-7). Among these taxa, MIA was associated with a higher level of *Pseudomonadaceae* (LDA score = 2.94, $p=0.02$) and *Christensenellaceae* (LDA score = 2.82, $p = 0.02$) on P10, and a greater expression of *Stenotrophomonas maltophilia* (LDA score= 3.45, $p=0.025$), *Ruminococcaceae* (LDA score = 3.35, $p = 0.02$) and *Lachnospiraceae* (LDA score= 3.85, $p = 0.01$) on P11. Data are expressed as mean ± SEM. *$p < 0.05$, ***$p <0.001$, MIA versus Saline; [a]$p < 0.05$, [aa]$p < 0.01$, main effect of time; [b]$p < 0.05$, [bb]$p < 0.01$, [bbb]$p < 0.001$, main effect of MIA. LPS – lipopolysaccharide.

Figure 3. Transcriptomic analyses of milk samples obtained on P10 from rat mothers exposed to either saline or maternal immune activation (MIA). (A) Volcano plot depicting the distribution of 829 genes based on log2 fold change and -log10 p values. Grey dots are genes, and the dots highlighted in blue represented genes that displayed the highest magnitude of significance ($padj<0.05$, $FC>1.3$). (B) Heatmaps of differentially expressed genes related specifically to milk nutrient transport, triglyceride concentration, parvalbumin interneuron development, aldosterone/MR signaling, inflammation, LPS binding, oxytocin, prolactin, glucocorticoid signaling (events following glucocorticoids binding to the GR receptor), glutamate/GABA, and epigenetic modifiers. Expression is represented with the log2 transformation of counts recorded with a z-score based on the average across experimental groups. Data are expressed as *$p <0.05$ or **adjusted $p <0.05$, or ***$FC>1.3$, MIA versus Saline. GR- glucocorticoid receptor. MR- mineralocorticoid receptor.

Figure 4. Transcriptomic analyses of milk samples obtained on P11 from rat mothers exposed to either saline or maternal immune activation (MIA). (A) Volcano plot depicting the distribution of 893 genes based on log2 fold change and -log10 p values. Grey dots are genes, and the dots highlighted in blue represented genes that displayed the highest magnitude of significance ($padj<0.05$, $FC>1.3$). (B) Heatmaps of differentially expressed genes related specifically to milk nutrient transport, triglyceride concentration, parvalbumin interneuron development, aldosterone/MR signaling, inflammation, LPS binding, oxytocin,

prolactin, glucocorticoid signaling (events following glucocorticoids binding to the GR receptor), glutamate/GABA, and epigenetic modifiers. Expression is represented with the log2 transformation of counts recorded with a z-score based on the average across experimental groups. Data are expressed as *p <0.05 or **adjusted p <0.05, or ***FC>1.3, MIA versus Saline. GR- glucocorticoid receptor. MR- mineralocorticoid receptor.

Figure 5. Adolescent offspring physiology and behavior following maternal immune activation (MIA) during the mid-lactational period. A main effect of MIA was observed in males for (A) postnatal day 43 body weights (grams) ($F(1,13) = 19.037$, p = 0.001, $n_p^2 = 0.594$. (B) Mechanical paw withdrawal thresholds (grams) on the von Frey test were significantly greater in female offspring ($F(1, 12) = 8.925$, p =0.011, $n_p^2 = 0.427$). Male offspring made a greater number of crosses into the (C) center ($F(1, 13) = 5.744$, p = 0.032, $n_p^2 = 0.306$); and (D) perimeter ($F(1, 13) = 6.513$, p = 0.024, $n_p^2 = 0.334$) of an open field arena. There were no effect of MIA on (E) social preference index. Line plots displaying percent prepulse inhibition (PPI) as a function of increasing prepulse intensities for (F) males (significant MIA by time interaction $F(4, 52) = 4.578$, p = 0.003, $n_p^2 = 0.260$). Male offspring from MIA dams demonstrated significant reductions in PPI at the 69dB ($F(1, 13) = 17.009$, p = 0.001, $n_p^2 = 0.743$), 73dB ($F(1, 13) = 5.927$, p =0.030, $n_p^2 = 0.581$), and 81dB ($F(1, 13) = 8.315$, p = 0.013, $n_p^2 = 0.633$) intensities. (G) PPI performance for females (main effect of dB $F(4, 52) = 80.357$, p = 0.001, $n_p^2 = 0.861$), and (H) male and female PPI combined for display purposes. (I) The bar plot shows the mean percent prepulse inhibition across all five prepulse intensities. Conditioned fear % PPI for (J) males (main effect of MIA for 73dB ($F(1, 13) = 6.787$, p = 0.022, $n_p^2 = 0.343$), 81dB ($F(1, 13) = 6.787$, p = 0.022, $n_p^2=0.343$) and 85 dB ($F(1, 13) = 13.124$, p = 0.003, $n_p^2=0.502$)). Conditioned fear % PPI for (K), females (main effect of MIA for 69dB ($F(1, 13) =12.969$, p = 0.003, $n_p^2=0.495$) and 81dB ($F(1, 13) = 25.117$, p = 0.001, $n_p^2=0.659$)). (L) Males and females combined for display purposes. (M) Hippocampal parvalbumin (PV) interneuron and densitometric ratios for male and female offspring from saline (black) and MIA (grey)-treated dams. PV/β-actin ratios were significantly reduced in male MIA offspring ($F(1, 13) =6.96$, p = 0.022, $n_p^2=.367$). Data are expressed as mean ± SEM, n = 7-8. *p <0.05, **p<0.01, ***p <0.001, MIA versus Saline. [a]p <0.05, [aa]p<0.01, [aaa]p <0.001, main effect of trial type.

Figure 6. Summary of mechanisms affecting milk quality and offspring brain and behavior following maternal immune activation (MIA) during the mid-lactational period. Mid-lactational MIA is associated with changes in the milk transcriptome and increased milk corticosterone concentrations, in addition to elevated microbial expression of *Christensenellaceae*. In concert, these effects may lead to reduced levels of milk LPS binding protein and fat content. We propose that this altered milk profile is one mechanism by which mid-lactational MIA contributes to the emergence of sex-specific behavioral and physiological changes in offspring. LPS – lipopolysaccharide; MR- mineralocorticoid receptor.

Supplementary Table 1. Maternal Immune Activation Model Reporting Guidelines Checklist.

Supplementary Table 2. Differentially expressed genes in P10 milk samples from the selected pathways. Genes with "NA" padj values indicate that FDR was not calculable due to fewer normalized counts than the optimal threshold.

Supplementary Table 3. Differentially expressed genes in P11 milk samples within the targeted pathways.

Key Resources Table.

Supplementary Methods.

Supplementary Figure 1. Maternal care behaviors following maternal immune activation (MIA) during the mid-lactational period. Total number of (A) pup retrievals, (B) passive nursing bouts, (C) nest building behaviors, and (D) maternal drinking bouts. Maternal care was evaluated in the afternoon and night of postnatal day (P) 10 and the morning of P11. Data are expressed as mean ± SEM, n = 7-8. ^{a}p < 0.05, main effect of time.

Supplementary Figure 2. Nutritional profile of milk following maternal immune activation (MIA) during the mid-lactational period. Maternal milk (A) fat concentration (g/L), (B) energy value (kcal/L), (C) protein concentration (mg/mL), (D) lactose concentration (ng/μL), and IgA concentration (mg/mL). Data are expressed as mean ± SEM, n = 7-8. *p < 0.05, ***p <0.001, MIA versus Saline; ^{a}p < 0.05, main effect of time.

Supplementary Figure 3. Cladogram of milk biomarkers associated with treatment group on P10 (determined by LEfSe). Diameter of the nodes indicates relative abundance of taxa for saline (green) and MIA (red) samples. Placement indicates the classification of taxa, where nodes decrease in rank the closer to the center of the diagram.

Supplementary Figure 4. Taxonomy heatmap demonstrating the top fifty most abundant species identified in samples on P10. Treatment group is indicated by the colored bar at the top of the figure (red = saline, blue = MIA). Each row represents the abundance for each taxon, with the taxonomy ID shown on the right. Each column represents the abundance for each sample.

Supplementary Figure 5. Cladogram of milk biomarkers associated with treatment group on P11 (determined by LEfSe). Diameter of the nodes indicates relative abundance of taxa for saline (green) and MIA (red) samples. Placement indicates the classification of taxa, where nodes decrease in rank the closer to the center of the diagram.

Supplementary Figure 6. Taxonomy heatmap demonstrating the top fifty most abundant species identified in samples on P11. Treatment group is indicated by the colored bar at the top of the figure (red = saline, blue = MIA). Each row represents the abundance for each taxon, with the taxonomy ID shown on the right. Each column represents the abundance for each sample.

Supplementary Figure 7. Adolescent offspring behavior following maternal immune activation (MIA) during the mid-lactational period. (A) Distance traveled (cm), (B) percent time spent in the center, and duration of time (seconds) spent in the (C) center and (D) perimeter of an open field arena. (E) Conditioned fear significantly increased % PPI regardless of drug or sex ($F_{(4, 52)}$= 8.19, p=0.008, n_p^2=.226). (F) % PPI for trials primed with conditioned fear collapsed across males and females for display purposes. Data are expressed as mean ± SEM, n = 7-8.

References

Anderson, G., Vaillancourt, C., Maes, M., & Reiter, R. J. (2017). Breastfeeding and the gut-brain axis: is there a role for melatonin?. Biomolecular concepts, 8(3-4), 185-195. https://doi.org/10.1515/bmc-2017-0009

Angelucci, L. (1985). A model for later-life effects of perinatal drug exposure: maternal hormone mediation. *Neurobehavioral Toxicology & Teratology*.

Arsenault, D., St-Amour, I., Cisbani, G., Rousseau, L.-S., & Cicchetti, F. (2014). The different effects of LPS and poly I:C prenatal immune challenges on the behavior, development and inflammatory responses in pregnant mice and their offspring. *Brain, Behavior, and Immunity*, *38*, 77–90. https://doi.org/10.1016/j.bbi.2013.12.016

Badillo-Suárez, P.A., Rodríguez-Cruz, M. & Nieves-Morales, X. Impact of Metabolic Hormones Secreted in Human Breast Milk on Nutritional Programming in Childhood Obesity. *Journal of Mammary Gland Biology Neoplasia, 22,* 171–191 (2017). https://doi.org/10.1007/s10911-017-9382-y

Baharnoori, M., Bhardwaj, S. K., & Srivastava, L. K. (2012). Neonatal behavioral changes in rats with gestational exposure to lipopolysaccharide: a prenatal infection model for developmental neuropsychiatric disorders. *Schizophrenia Bulletin, 38*(3), 444–456. https://doi.org/10.1093/schbul/sbq098

Balogh, S. A., Radcliffe, R. A., Logue, S. F., & Wehner, J. M. (2002). Contextual and cued fear conditioning in C57BL/6J and DBA/2J mice: context discrimination and the effects of retention interval. *Behavioral Neuroscience*, *116*(6), 947. doi: 10.1037//0735-7044.116.6.947.

Basolo, F., Fiore, L., Fontanini, G., Conaldi, P. G., Calvo, S., Falcone, V., & Toniolo, A. (1996). Expression of and response to interleukin 6 (IL6) in human mammary tumors. *Cancer Research, 56*(13), 3118-3122. 8674070

Bison, S., Carboni, L., Arban, R., Bate, S., Gerrard, P. A., & Razzoli, M. (2009). Differential behavioral, physiological, and hormonal sensitivity to LPS challenge in rats. *International Journal of Interferon, Cytokine and Mediator Research, 1*, 1-13. https://doi.org/10.2147/IJICMR.S4273

Borish, L.C. & Steinke, J.W. (2003). Cytokines and chemokines. *Journal of Allergy Clinical Immunology. 111*, 460-745. doi: 10.1067/mai.2003.108.

Bowen, W.S., Gandhapudi, S.K., Kolb, J.P., Mitchell, T.C. (2013). Chapter Three-Immunopharmacology of Lipid A Mimetics, *Advances in Pharmacology, 66,* 81-128. https://doi.org/10.1016/B978-0-12-404717-4.00003-2

Brown, R., Imran, S. A., & Wilkinson, M. (2009). Lipopolysaccharide (LPS) stimulates adipokine and socs3 gene expression in mouse brain and pituitary gland in vivo, and in N-1 hypothalamic neurons in vitro. Journal of *Neuroimmunology, 209*(1-2), 96-103. https://doi.org/10.1016/j.jneuroim.2009.02.001

Browne, P. D., Aparicio, M., Alba, C., Hechler, C., Beijers, R., Rodríguez, J. M., ... & de Weerth, C. (2019). Human milk microbiome and maternal postnatal psychosocial distress. *Frontiers in Microbiology, 10,* 2333. https://doi.org/10.3389/fmicb.2019.02333

Brummelte, S., Schmidt, K.L., Taves, M.D., Soma, K.K. & Galea, L. A. (2010). Chronic corticosterone during pregnancy and postpartum affects maternal care, cell proliferation and depressive-like behavior in the dam. *Hormones and Behavior, 58*(5), 769-779. https://doi.org/10.1016/j.yhbeh.2010.07.012

Callahan, L. S., Thibert, K. A., Wobken, J. D., & Georgieff, M. K. (2013). Early-life iron deficiency anemia alters the development and long-term expression of parvalbumin and perineuronal nets in the rat hippocampus. Developmental neuroscience, 35(5), 427-436. https://doi.org/10.1159/000354178

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K. et al. (2010) QIIME allows analysis of high-throughput community sequencing data. *Nature Methods, 7*: 335-336. https://doi.org/10.1038/nmeth.f.303

Catalani, A., Marinelli, M., Scaccianoce, S., Nicolai, R., Muscolo, L. A., Porcu, A., ... & Angelucci, L. (1993). Progeny of mothers drinking corticosterone during lactation has lower stress-induced corticosterone secretion and better cognitive performance. *Brain Research, 624*(1-2), 209-215. https://doi.org/10.1016/0006-8993(93)90079-3

Catalani, A., Casolini, P., Cigliana, G., Scaccianoce, S., Consoli, C., Cinque, C., ... & Angelucci, L. (2002). Maternal corticosterone influences behavior, stress response and corticosteroid receptors in the female rat. *Pharmacology Biochemistry and Behavior, 73*(1), 105-114. https://doi.org/10.1016/S0091-3057(02)00755-4

Champagne, F.A. & Curley, J.P. (2009). Epigenetic mechanisms mediating the long-term effects of maternal care on development. *Neuroscience Biobehavioral Reviews, 33*(4):593-600. doi: 10.1016/j.neubiorev.2007.10.009.

Champagne, F. A., Francis, D. D., Mar, A., & Meaney, M. J. (2003). Variations in maternal care in the rat as a mediating influence for the effects of environment on development. *Physiology & Behavior, 79*(3), 359-371. https://doi.org/10.1016/S0031-9384(03)00149-5

Chen, Y., Wang, J., Yang, S., Utturkar, S., Crodian, J., Cummings, S., Thimmapuram, J., San Miguel, P., Kuang, S., Gribskov, M., Plaut, K., Casey, T. (2017). Effect of high fat diet on secreted milk transcriptome in midlactation mice. *Physiology Genomics, 49,* 747-762, doi:10.1152/physiolgenomics.00080.2017.

Chen, J., Wu, Y., Sun, Y., Dong, X., Wang, Z., Zhang, Z., ... & Dong, G. (2019). Bacterial lipopolysaccharide induced alterations of genome-wide DNA methylation and promoter methylation of lactation-related genes in bovine mammary epithelial cells. *Toxins, 11*(5), 298. https://doi.org/10.3390/toxins11050298

Connor, K. L., Chehoud, C., Altrichter, A., Chan, L., DeSantis, T. Z., & Lye, S. J. (2018).

Maternal metabolic, immune, and microbial systems in late pregnancy vary with malnutrition in mice. *Biology of Reproduction, 98*(4), 579-592. https://doi.org/10.1093/biolre/ioy002

Connors, E. J., Shaik, A. N., Migliore, M. M., & Kentner, A. C. (2014). Environmental enrichment mitigates the sex-specific effects of gestational inflammation on social engagement and the hypothalamic pituitary adrenal axis-feedback system. *Brain, Behavior, and Immunity, 42,* 178-190. https://doi.org/10.1016/j.bbi.2014.06.020

DeRosa, H., Caradonna, S.G., Tran, H., Marrocco, J., & Kentner, A.C. (2022). Milking it for all it's worth: The effects of environmental enrichment on maternal nurturance, lactational quality, and offspring social behavior. *bioRxiv.* doi: https://doi.org/10.1101/2021.12.13.472509

Dockman, R. L., Carpenter, J. M., Diaz, A. N., Benbow, R. A., & Filipov, N. M. (2021). Sex differences in behavior, response to LPS, and glucose homeostasis in middle-aged mice. *Behavioural Brain Research, 113628*.https://doi.org/10.1016/j.bbr.2021.113628

Edwards, P.D., Lavergne, S.G., McCaw, L.K., Wijenayake, S., Boonstra, R., McGowan, P.O., Holmes, M.M. (2021). Maternal effects in animals: broadening our understanding of offspring programming. *Frontiers in Neuroendocrinology*, 62, 100924, doi: 10.1016/j.yfrne.2021.100924

Estes, M.L. & McAllister, A.K. (2016). Maternal immune activation: Implications for neuropsychiatric disorders. *Science, 353*(6301):772-7. doi: 10.1126/science.aag3194.

Fortier, M. E., Luheshi, G. N., & Boksa, P. (2007). Effects of prenatal infection on prepulse inhibition in the rat depend on the nature of the infectious agent and the stage of pregnancy. *Behavioural Brain Research, 181*(2), 270-277. https://doi.org/10.1016/j.bbr.2007.04.016

Francis, D. D., Young, L. J., Meaney, M. J., & Insel, T. R. (2002). Naturally occurring differences in maternal care are associated with the expression of oxytocin and vasopressin (V1a) receptors: gender differences. *Journal of Neuroendocrinology, 14*(5), 349-353.

Francis, D. D., & Kuhar, M. (2008). Frequency of maternal licking and grooming correlates negatively with vulnerability to cocaine and alcohol use in rats. *Pharmacology Biochemistry and Behavior, 90*(3), 497-500. https://doi.org/10.1016/j.pbb.2008.04.012

Franco, J. G., Fernandes, T. P., Rocha, C. P., Calviño, C., Pazos-Moura, C. C., Lisboa, P. C., Moura, E. G., & Trevenzoli, I. H. (2012). Maternal high-fat diet induces obesity and adrenal and thyroid dysfunction in male rat offspring at weaning. *The Journal of Physiology, 590*(21), 5503–5518. https://doi.org/10.1113/jphysiol.2012.240655

Garay, P. A., Hsiao, E. Y., Patterson, P. H., & McAllister, A. K. (2013). Maternal immune activation causes age-and region-specific changes in brain cytokines in offspring throughout development. *Brain, Behavior, and Immunity, 31*, 54-68. https://doi.org/10.1016/j.bbi.2012.07.008

García-Torres, E., Hudson, R., Castelán, F., Martínez-Gómez, M., & Bautista, A. (2015). Differential metabolism of brown adipose tissue in newborn rabbits in relation to position in the litter huddle. *Journal of Thermal Biology, 51*, 33-41. https://doi.org/10.1016/j.jtherbio.2015.03.003

Gilbert, C., McCafferty, D., Le Maho, Y., Martrette, J. M., Giroud, S., Blanc, S., & Ancel, A. (2010). One for all and all for one: the energetic benefits of huddling in endotherms. *Biological Reviews, 85*(3), 545-569. https://doi.org/10.1111/j.1469-185X.2009.00115.x

Grazia Di Benedetto, M., Bottanelli, C., Cattaneo, A., Pariante, C.M., Borsini, A. (2020). Nutritional and immunological factors in breast milk: a role in the intergenerational transmission from maternal psychopathology to child development. *Brain, Behavior, and Immunity*, 85, 57-68.

Hinde, K., Skibiel, A. L., Foster, A. B., Del Rosso, L., Mendoza, S. P., & Capitanio, J. P. (2015). Cortisol in mother's milk across lactation reflects maternal life history and predicts infant temperament. *Behavioral Ecology, 26*(1), 269-281. https://doi.org/10.1093/beheco/aru186

Holingue, C., Brucato, M., Ladd☐Acosta, C., Hong, X., Volk, H., Mueller, N. T., ... & Fallin, M. D. (2020). Interaction between Maternal Immune Activation and Antibiotic Use during Pregnancy and Child Risk of Autism Spectrum Disorder. *Autism Research, 13*(12), 2230-2241. https://doi.org/10.1002/aur.2411

Howland, J. G., Cazakoff, B. N., & Zhang, Y. (2012). Altered object-in-place recognition memory, prepulse inhibition, and locomotor activity in the offspring of rats exposed to a viral mimetic during pregnancy. *Neuroscience, 201,* 184-198. https://doi.org/10.1016/j.neuroscience.2011.11.011

Hsiao, E.Y. & Patterson, P.H. (2011). Activation of the maternal immune system induces endocrine changes in the placenta via IL-6. *Brain, Behavior, and Immunity, 25*(4), 604-15. doi: 10.1016/j.bbi.2010.12.017.

Hussein J, Mavalankar DV, Sharma S, D'Ambruoso L. (2011). A review of health system infection control measures in developing countries: what can be learned to reduce maternal mortality. *Global Health*, 7, 14.

Inceoglu, B., Jinks, S. L., Schmelzer, K. R., Waite, T., Kim, I. H., & Hammock, B. D. (2006). Inhibition of soluble epoxide hydrolase reduces LPS-induced thermal hyperalgesia and mechanical allodynia in a rat model of inflammatory pain. *Life Sciences, 79*(24), 2311–2319. https://doi.org/10.1016/j.lfs.2006.07.031

Ishii, D., Matsuzawa, D., Fujita, Y., Sutoh, C., Ohtsuka, H., Matsuda, S., ... & Shimizu, E. (2010). Enhancement of acoustic prepulse inhibition by contextual fear conditioning in mice is maintained even after contextual fear extinction. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *34*(1), 183-188.

John, C.C., Black, M.M., Nelson, C.A III. (2017). Neurodevelopment: the impact of nutrition

and inflammation during early to middle childhood in low-resource settings. *Pediatrics*, 139, e 20162828.

Kentner, A.C., Bilbo, S.D., Brown, A.S., Hsiao, E.Y., McAllister, A.K., Meyer, U., Pearce, B.D., Pletnikov, M.V., Yolken, R.H., Bauman, M.D. (2019). Maternal immune activation: reporting guidelines to improve the rigor, reproducibility, and transparency of the model. *Neuropsychopharmacology 44*, 245-258.

Karolinski A, Mazzoni A, Belizan JM, Althabe F, Bergel E, Buekens P (2010). Lost opportunities for effective management of obstetric conditions to reduce maternal mortality and severe maternal morbidity in Argentina and Uruguay. Int J Gynaecol Obstet, 110, 175–180.

Kutlesic, V., Breinski Isaacs, M., Freund, L.S., Hazra, R., Raiten, D.J. (2017). Executive summary: research gaps at the intersection of pediatric neurodevelopment, nutrition, and inflammation in low resource settings. Pediatrics, 139, S1-S11.

Lesorogol, C., Bond, C., Dulience, S. J. L., & Iannotti, L. (2018). Economic determinants of breastfeeding in Haiti: the effects of poverty, food insecurity, and employment on exclusive breastfeeding in an urban population. *Maternal & Child Nutrition, 14*(2), e12524. https://doi.org/10.1111/mcn.12524

Liu, L., Lin, Y., Liu, L., Bian, Y., Zhang, L., Gao, X., & Li, Q. (2015). 14-3-3γ regulates lipopolysaccharide-induced inflammatory responses and lactation in dairy cow mammary epithelial cells by inhibiting NF-κB and MAPKs and up-regulating mTOR signaling. *International Journal of Molecular Sciences, 16*(7), 16622-16641. https://doi.org/10.1016/j.aninu.2016.03.005

Ling B, Alcorn J. LPS-induced inflammation downregulates mammary gland glucose, fatty acid, and L-carnitine transporter expression at different lactation stages. (2010). *Research in Veterinary Science, 89*(2):200-2. doi: 10.1016/j.rvsc.2010.03.004

Lucas, A., Gibbs, J.A., Lyster, R.L., Baum, J.D. (1978). Creamatocrit: simple clinical technique for estimating fat concentration and energy value of human milk. Br Med J., 1 (6119):1018-1020, doi: 10.1136/bmj.1.6119.1018.

Mahic, M., Che, X., Susser, E., Levin, B., Reichborn-Kjennerud, T., Magnus, P., ... & Lipkin, W. I. (2017). Epidemiological and serological investigation into the role of gestational maternal influenza virus infection and autism spectrum disorders. *Msphere, 2*(3), e00159-17. https://doi.org/10.1128/mSphere.00159-17

Mäkelä, J., Linderborg, K., Niinikoski, H., Yang, B., & Lagström, H. (2013). Breast milk fatty acid composition differs between overweight and normal weight women: the STEPS Study. *European Journal of Nutrition, 52*(2), 727-735. https://doi.org/10.1007/s00394-012-0378-5

Meyer, U. (2019). Neurodevelopmental Resilience and Susceptibility to Maternal Immune Activation. *Trends in Neurosciences, 42*(11), 793–806. https://doi.org/10.1016/j.tins.2019.08.001

Miles, J., Shevlin, M., 2001. Applying regression and correlation: A guide for students and researchers. Sage.

Millett, C. E., Phillips, B. E., & Saunders, E. F. (2019). The sex-specific effects of LPS on depressive-like behavior and oxidative stress in the hippocampus of the mouse. *Neuroscience, 399*, 77-88. https://doi.org/10.1016/j.neuroscience.2018.12.008

Mueller, F. S., Richetto, J., Hayes, L. N., Zambon, A., Pollak, D. D., Sawa, A., ... & Weber-Stadlbauer, U. (2019). Influence of poly (I: C) variability on thermoregulation, immune responses and pregnancy outcomes in mouse models of maternal immune activation. *Brain, Behavior, and Immunity, 80*, 406-418. doi: 10.1016/j.bbi.2019.04.019

Nascimento, A. F., Alves, G. J., Massoco, C. O., Teodorov, E., Felicio, L. F., & Bernardi, M. M. (2015). Lipopolysaccharide-induced sickness behavior in lactating rats decreases ultrasonic vocalizations and exacerbates immune system activity in male offspring. *Neuroimmunomodulation, 22*(4), 213-221. https://doi.org/10.1159/000363350

Naskar, S., Narducci, R., Balzani, E. et al. (2019). The development of synaptic transmission is time-locked to early social behaviors in rats. *Nature Communications, 10*, 1195 https://doi.org/10.1038/s41467-019-09156-3

Nguyen, R., Morrissey, M.D., Mahadevan, V., Cajanding, J.D., Woodin, M.A., Yeomans, J.S. Takehara-Nishiuchi K, Kim JC. (2014). Parvalbumin and GAD65 interneuron inhibition in the ventral hippocampus induces distinct behavioral deficits relevant to schizophrenia. *Journal of Neuroscience. 34*(45):14948-60. doi: 10.1523/JNEUROSCI.2204-14.2014.

Núñez Estevez, K.J., Rondón-Ortiz, A.N., Nguyen, J.Q.T., & Kentner, A.C. (2020). Environmental influences on placental programming and offspring outcomes following maternal immune activation. *Brain, Behavior, and Immunity, 83*, 44-55. doi: 10.1016/j.bbi.2019.08.192.

Oakley, RH & Cidlowski, JA (2013). The biology of the glucocorticoid receptor: new signaling mechanisms in health and disease. J Allergy Clin Immunol., 132(5), 1033-1044. https://doi.org/10.1016/j.jaci.2013.09.007

Ordoñes Sanchez, E., Bavley, C.C., Deutschmann, A.U., Carpenter, R., Peterson, D.R., Karbalaei, R....Bangasser, D.A. (2021). Early life adversity promotes resilience to opioid addiction-related phenotypes in male rats and sex-specific transcriptional changes. *PNAS USA, 118*, e2020173118. doi: 10.1073/pnas.2020173118.

Ozkan, H, Tuzun, F, Taheri, S, Korhan, P, Akokay, P, Yılmaz, O, Duman, N, Özer, E, Tufan, E, Kumral, A, & Özkul, Y (2020). Epigenetic programming through breast milk and its impact on milk-siblings mating. Front. Genet., 11, 569232. https://doi.org/10.3389/fgene.2020.569232

Pannaraj, P. S., Li, F., Cerini, C., Bender, J. M., Yang, S., Rollie, A., ... & Aldrovandi, G. M.

(2017). Association between breast milk bacterial communities and establishment and development of the infant gut microbiome. JAMA Pediatrics, 171(7), 647-654.

Par Pharmaceutical, Inc, Pitocin [Label], Par Pharmaceutical, Chestnut Ridge, NY, (2020), https://dailymed.nlm.nih.gov/dailymed/fda/fdaDrugXsl.cfm?setid=6d4b2c25-2e5d-49b5-93bc-2ae8a20916d1&type=display

Park, B. & Lee, J.O. (2013). Recognition of lipopolysaccharide pattern by TLR4 complexes. Experimental Molecular Medicine 45, https://doi.org/10.1038/emm.2013.97

Patacchioli, F. R., Cigliana, G., Cilumbriello, A., Perrone, G., Capri, O., Alemà, S., ... & Angelucci, L. (1992). Maternal plasma and milk free cortisol during the first 3 days of breast-feeding following spontaneous delivery or elective cesarean section. *Gynecologic and Obstetric Investigation*, *34*(3), 159-163. https://doi.org/10.1159/000292751

Paul, H.A., Hallam, M.C., Reimer, R.A. (2015). Milk collection in the rat using capillary tubes and estimation of milk fat content by creamatocrit. *Journal of Visualized Experiments, 106*, 1-6, doi: 10.3791/53476.

Polman JA, Welten JE, Bosch DS, de Jonge RT, Balog J, van der Maarel SM, de Kloet ER, Datson NA. (2012). A genome-wide signature of glucocorticoid receptor binding in neuronal PC12 cells. BMC Neurosci.13:118. doi: 10.1186/1471-2202-13-118.

Quintana, DS, Rokicki, J, van der Meer, D, Alnæs, D, Kaufmann, T, Córdova-Palomera, A, Dieset, I, Andreassen, OA, & Westlye, LT. (2019). Oxytocin pathway gene networks in the human brain. Nat Commun 10, 668. https://doi.org/10.1038/s41467-019-08503-8

Schatz, K. C., Kyne, R. F., Parmeter, S. L., & Paul, M. J. (2018). Investigation of social, affective, and locomotor behavior of adolescent Brattleboro rats reveals a link between vasopressin's actions on arousal and social behavior. *Hormones and Behavior, 106*, 1-9. https://doi.org/10.1016/j.yhbeh.2018.08.015

Schellekens, H., Torres-Fuentes, C., van de Wouw, M., Long-Smith, C. M., Mitchell, A., Strain, C., ... & Cryan, J. F. (2021). Bifidobacterium longum counters the effects of obesity: Partial successful translation from rodent to human. *EBioMedicine, 63,* 103176.

Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., and Huttenhower, C. (2011) Metagenomic biomarker discovery and explanation. *Genome Biology, 12*: R60. doi: 10.1186/gb-2011-12-6-r60.

Sheard, N. F., & Walker, W. A. (1988). The role of breast milk in the development of the gastrointestinal tract. *Nutrition Reviews (USA).* doi: 10.1111/j.1753-4887.1988.tb05343.x.

Singh, D., & Fulekar, M. H. (2010). Biodegradation of petroleum hydrocarbons by Pseudomonas putida strain MHF 7109. *CLEAN–Soil, Air, Water, 38*(8), 781-786. https://doi.org/10.1002/clen.200900239

Soriano, O., Regalado, M., Torrero, C., & Salas, M. (2006). Contributions of undernutrition and handling to huddling development of rats. *Physiology & Behavior, 89*(4), 543-551. https://doi.org/10.1016/j.physbeh.2006.07.021

Sullivan, E. C., Hinde, K., Mendoza, S. P., & Capitanio, J. P. (2011). Cortisol concentrations in the milk of rhesus monkey mothers are associated with confident temperament in sons, but not daughters. *Developmental Psychobiology, 53*(1), 96–104. https://doi.org/10.1002/dev.20483

Vacca, M., Celano, G., Calabrese, F. M., Portincasa, P., Gobbetti, M., & De Angelis, M. (2020). The controversial role of human gut lachnospiraceae. *Microorganisms, 8*(4), 573. doi: 10.3390/microorganisms8040573

van Hasselt, F. N., Cornelisse, S., Yuan Zhang, T., Meaney, M. J., Velzing, E. H., Krugers, H. J., & Joëls, M. (2012). Adult hippocampal glucocorticoid receptor expression and dentate synaptic plasticity correlate with maternal care received by individuals early in life. *Hippocampus, 22*(2), 255-266. https://doi.org/10.1002/hipo.20892

Vilela, F.C., Antunes-Rodrigues, J., Elias, L.L., & Giusti-Paiva, A. (2013). Corticosterone synthesis inhibitor metyrapone preserves changes in maternal behavior and neuroendocrine responses during immunological challenge in lactating rats. *Neuroendocrinology. 97*(4):322-30. doi: 10.1159/000346354

Waters, J. L., & Ley, R. E. (2019). The human gut bacteria Christensenellaceae are widespread, heritable, and associated with health. *BMC Biology, 17*(1), 1-11. https://doi.org/10.1186/s12915-019-0699-4

Wu, W.L., Hsiao, E.Y., Yan, Z., Mazmanian, S.K., & Patterson, P.H. (2017). The placental interleukin-6 signaling controls fetal brain development and behavior. *Brain, Behavior, and Immunity. 62,* 11-23. doi: 10.1016/j.bbi.2016.11.007.

Wurfel, M.M. & Wright, S.D. (1997). Lipopolysaccharide-binding protein and soluble CD14 transfer lipopolysaccharide to phospholipid bilayers: preferential interaction with particular classes of lipid. *Journal of Immunology. 15;158*(8):3925-34. PMID: 9103463.

Yan S. & Kentner A.C. (2017). Mechanical allodynia corresponds to Oprm1 downregulation within the descending pain network of male and female rats exposed to neonatal immune challenge. *Brain, Behavior, and Immunity*, *63,* 148-159. doi: 10.1016/j.bbi.2016.10.007.

Zhang, TY, Labonté, B, Wen, XL, Turecki, G, & Meaney, MJ (2013). Epigenetic mechanisms for the early environmental regulation of hippocampal glucocorticoid receptor gene expression in rodents and humans. NPP, 38(1), 111–123. https://doi.org/10.1038/npp.2012.149

Zhao, X., Mohammed, R., Tran, H., Erickson, M., & Kentner, A. C. (2021a). Poly (I: C)-

induced maternal immune activation modifies ventral hippocampal regulation of stress reactivity: prevention by environmental enrichment. *Brain, Behavior, and Immunity*. https://doi.org/10.1016/j.bbi.2021.03.018

Zhao, X., Tran, H., DeRosa, H., Roderick, R.C., & Kentner, A.C. (2021b) Hidden talents: Poly (I:C)-induced maternal immune activation improves mouse visual discrimination performance and reversal learning in a sex-dependent manner. *Genes, Brain, and Behavior, 20*(7). doi: 10.1111/gbb.12755.

Zhu X, Girardo D, Govek EE, John K, Mellén M, Tamayo P, Mesirov JP, Hatten ME. Role of Tet1/3 Genes and Chromatin Remodeling Genes in Cerebellar Circuit Formation. (2016). *Neuron*. 89(1):100-12. doi: 10.1016/j.neuron.2015.11.030

Zietek, M., Sochaczewska, D., Swiatkowska-Freund, M., Celewicz, Z., & Szczuko, M. (2021). The possible role of corticosterone in regulating sodium and potassium concentrations in human milk. *Ginekologia Polska.* doi: 10.5603/GP.a2021.0072.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.