Got Milk? Maternal immune activation during the mid-lactational period affects nutritional milk quality and adolescent offspring sensory processing in male and female rats

Holly DeRosa[1], Hieu Tran[1], Amanda C. Kentner[1]*

School of Arts & Sciences, Health Psychology Program, Massachusetts College of Pharmacy and Health Sciences, Boston Massachusetts, United States 02115

*Corresponding author:
Amanda Kentner
amanda.kentner@mcphs.edu
Office #617-274-3360
Fax # 617-732-2959

Keywords
Maternal immune activation; lactation, nutrition; prepulse inhibition; mechanical allodynia; conditioned fear; microbiome; parvalbumin; milk, breastfeeding; animal models; early life stress

Abstract

The neonatal environment requires a high level of maternal demand in terms of both breastfeeding and other forms of maternal care. Previous studies have underscored the importance of these maternal factors on offspring development and behavior. However, their contribution as dynamic variables in animal models of early life stress are often overlooked. In the present study, we show that lipopolysaccharide (LPS)-induced maternal immune activation (MIA) on postnatal day (P)10 elevated milk corticosterone concentrations, which recovered by P11. In contrast, both milk triglyceride and percent creamatocrit values demonstrated a prolonged decrease following MIA. Sustained changes to the nutritional quality of milk were also evidenced by an increase of microbial communities associated with inefficient energy and lipid metabolism. Nutritional deficits in early development have been associated with metabolic dysfunction later in life. Indeed, these nutritional milk changes were reflected by increased adolescent offspring bodyweights. While MIA did not decrease maternal care quality, there was a significant compensatory increase in maternal licking and grooming the day that followed the inflammatory challenge. However, this did not protect against disrupted neonatal huddling or later-life alterations in sensorimotor gating, mechanical allodynia, or hippocampal parvalbumin interneuron expression in MIA offspring. Since milk concentrations of LPS binding protein and interleukin-6 were not different between MIA and control dams, these offspring effects were likely driven by nutritional factors and not by a direct exposure to LPS or inflammatory molecules. Animal models of early life stress can impact both parents and their progeny; our data show that inflammatory stress affects maternal lactation quality which acts as a mechanism to confer multifaceted and compounding effects on offspring.

1. Introduction

Epidemiological studies have shown that prenatal exposure to maternal immune activation (MIA) is a leading risk factor for developing psychiatric disorders such as autism and schizophrenia (Mahic et al., 2017; Meyer, 2019). Animal models of MIA often employ the bacterial endotoxin lipopolysaccharide (LPS) or the toll-like 3 receptor agonist polyinosinic:polycytidylic acid (Poly I:C) to elicit a maternal immune response (Mueller et al., 2019). Administration of either of these immunogens promotes a cascade of inflammatory cytokines in the host, such as tumor necrosis factor (TNF)-α, interleukins, and T cells, in addition to activation of the hypothalamic-pituitary-adrenal (HPA) axis (Kentner et al., 2018; Brown et al., 2009; Arsenault et al., 2014; Estes & McAllister, 2016). In concert, these physiological responses contribute to a variety of sickness behaviors, including ptosis (eye drooping), piloerection (raised and ruffled fur), lethargy, and reduced food intake (Connors et al., 2014; Arsenault et al., 2014). In addition to eliciting these acute sickness behaviors, LPS exposure can lead to other changes in both pregnant and non-pregnant animals such as reduced locomotion (Millett et al., 2019), anhedonia, and improved performance in the Barnes Maze and novel object recognition task (Dockman et al., 2021). While the sickness-inducing effects of immunogens are also recognized to affect the long-term development of offspring when exposed prenatally (Kentner et al., 2018; Estes & McAllister, 2016), the impact of maternal infection during lactation on both mothers and their offspring is relatively unexplored. This is surprising because infections during the postnatal period are common, particularly in low resource settings, and can result in severe morbidity and mortality (Karolinski et al., 2010; Kutlesic et al., 2017; John et al., 2017; Hussein et al., 2011).

When administered to rodent dams during the lactational period, LPS can contribute to deficits in maternal care. This has been evidenced by reductions in the efficient arched back nursing posture and the number of rat pups retrieved and returned back to the nest (Vilela et al., 2013). In line with the behavioral alterations, MIA in lactating dams may influence maternal milk quality. For example, maternal LPS treatment on postnatal days 4, 11, and 18 reduced the mRNA expression of milk nutrient precursors such as glucose and fatty acid transporters in the rat mammary gland (Ling & Alcorn, 2010). Exposure to MIA during the lactational period can also impact offspring behavior. For example, Nascimento and colleagues (2015) observed a reduction in ultrasonic vocalizations in male neonates shortly after their dams were treated with LPS. Although this presumably involved changes in various forms of maternal input signals (e.g., parental care and/or milk quality), the mechanisms of this MIA-derived effect on offspring behavior remain unclear.

Other exogenous factors have been shown to alter contents in breastmilk and subsequent offspring development (Lesorogol et al., 2018; Mäkelä et al., 2013; Chen et al., 2017; Grazia Di Benedetto et al., 2020; Edwards et al., 2021). For example, maternal high fat diet has been associated with offspring obesity alongside greater triglyceride and protein levels in milk (Franco et al., 2012). Psychological stress in humans reduces milk microbial diversity (Browne et al., 2019), which may have clinical implications for infant development. Indeed, maternal milk quality and the microbiota are critical to programming infant metabolism, gut microbiome, and the gut-brain axis (Badillo-Suárez et al., 2017; Sheard & Walker, 1988; Pannaraj et al., 2017; Anderson et al., 2014). Further, cortisol in maternal milk was found to predict temperament and promote greater body weights in infant primates, although milk energy content was also associated with body weight (Hinde et al., 2014). Together, this evidence suggests that 1) disturbances to the neonatal environment, such as MIA, alters the developmental trajectory of offspring, and 2) these alterations are mediated by changes in maternal care and/or nutritional milk quality. To explore these ideas further, we assessed the effects of MIA challenge during the mid-lactational period on maternal care, maternal milk composition, and offspring brain development and behavior in Sprague-Dawley rats.

2. Materials and Methods

2.1. Animals and Housing

Male and female Sprague-Dawley rats (Charles River, Wilmington, MA) were housed in same-sex pairs and maintained at 20°C on a 12 h light/dark cycle (0700-1900 light) in standard sized laboratory cages (27×48×22 cm). Food and water were available *ad libitum* throughout the study. Animal procedures were conducted in accordance with the Association for Assessment and Accreditation of Laboratory Animal Care with protocols approved by the Massachusetts College of Pharmacy and Health Sciences (MCPHS) Institutional Animal Care and Use Committee. **Figure 1A** outlines the experimental procedures and groups evaluated in this study. During breeding, animals were placed into larger cages (51 x 41 x 22 cm) and bred using a 1 male:2 female design. Pregnancy was confirmed by continued weight gain and visible teats during the later phase of gestation. Approximately two days prior to parturition, females were individually housed in fresh clean standard sized cages. Day of birth was designated as postnatal day (P)0 and litters were standardized to 10 pups per litter on P1. Wherever possible, litters were balanced with equal numbers of male and female pups. Animals stayed in this condition until they were weaned into same-sex pairs on P22, again using standard sized laboratory cages. All cages maintained a tube, a Nylabone chew toy®, and Nestlets® throughout the study, except during the two-day period prior to parturition

through until P14 when toys were removed to protect pups from injury. Additional Nestlets®
were available to litters during this time. One male and one female offspring from each litter
were evaluated on behavioral tasks beginning on P43. The behavioral monitoring equipment
was cleaned thoroughly with the disinfectant cleaner Quatriside TB (Pharmacal Research
Laboratories, Inc) between each animal and test. The inter-rater reliability between blinded
scorers was found to be greater than r=0.80 for each observational measure reported in this
study.

2.2. Maternal Immune Activation (MIA)

On the morning of P10, dams were removed from their litter and placed into a fresh clean
cage located in a separate procedure room. To model MIA during the lactational period, dams
were weighed and intraperitoneally (i.p.) administered either 100 µg/kg of the inflammatory
endotoxin, lipopolysaccharide (LPS; Escherichia coli, serotype 026:B6; L-3755, Sigma, St.
Louis, MO, USA; n = 7) or pyrogen-free saline (saline; n = 8) at the start of the separation
period. Animals remained in this cage for 2 hours until milking. The purpose of the
separation period was to allow for maternal milk accumulation so samples could be collected
at the height of the inflammatory response. In their regular holding room, pups were placed
into a smaller clean cage positioned on top of a heating pad to maintain their body
temperature. Pups were weighed immediately prior to being returned to their dams and again
2 and 24 hours later, alongside the inspection of milk bands, to monitor their health. Twenty-
four hours following the P10 inflammatory challenge, milk collection procedures were
repeated to evaluate sustained changes in milk quality. Additional details can be found in the
methodological reporting table from Kentner et al. (2019), provided as **Supplementary
Table S1**.

2.3. Milk Sample Collection and Composition

Milk was collected from each dam on P10 and P11 following a modified procedure from
Paul et al. (2015). Immediately following each 2-hour separation period, dams were lightly
anesthetized with isoflurane in O2, followed by the administration of 0.2 mL of oxytocin (20
USP/mL i.p.). The pharmacokinetic profile of isoflurane suggests that the drug is not
absorbed by offspring and that breastfeeding can resume immediately after anesthesia (Drugs
and Lactation Database, 2020; Lee & Rubin, 1993). Similarly, oxytocin is not expected to
affect offspring given its short plasma half-life of 1-6 minutes which is reduced even further
during lactation (Par Pharmaceutical Inc, 2020). Teats were prepared by moistening the
collection areas with distilled water. Milk was obtained from each dam by gently squeezing
the base of the teat and manually expelling the milk for collection. At the time of milk
collection, ~20 ul of sample was collected into a microhematocrit tube, which was then
sealed and placed into a hematocrit spinner (2 minutes at 13, 700 g; StatSpin CritSpin
Microhematocrit Centrifuge, Beckman Coulter, Inc). Measurements were taken to calculate
percent (%) creamatocrit by evaluating the sample separation into each of the cream and clear
layers by following the procedures outlined in Paul et al. (2015). The remaining milk that was
collected (~500 ul per animal) was aliquoted across microcentrifuge tubes and stored at -
80°C until further processing. Collection time took about 10-25 minutes per rat, and dams
were reunited with their litters in their home cage as soon as they fully awoke from
anesthesia.

Milk samples were homogenized by overnight rotation on a Mini Tube Rotator (Fisher
Scientific Cat. #88861051) at 4°C before analysis. A lactose assay kit (Sigma-Aldrich Cat.

#MAK017) was used to measure lactose content in milk diluted 1:500 with lactose assay buffer (as outlined by Chen et al., 2017). Triglycerides were evaluated using a colorimetric assay kit (Abcam, Cat. #ab65336) at 1:1000 dilution. The Pierce™ BCA Protein Assay Kit (Cat. #23227) was run at a 1:50 dilution to measure protein levels in milk. The small sample assay protocol of the corticosterone ELISA kit (#ADI-900–097, Enzo Life Sciences, Farmingdale, NY) was followed, as recommended by the manufacturer, using a 1:40 dilution. Milk levels of immunoglobulin (Ig) A were assessed with an assay kit (Bethyl Laboratories, Cat. # E111-102) using samples diluted to 1:1000 while interleukin (IL)-6 (Thermo Fisher Scientific, Cat. #BMS625) was evaluated using samples diluted 1:2. Undiluted milk samples and blood plasma (diluted 1:2) were processed using the LPS binding protein ELISA (Abcam, Cat. #ab269542). Manufacturer's instructions were followed for all ELISA kits (n = 7).

P10 (n = 6) and P11 (n = 6-7) milk samples underwent microbiome sequencing. DNA extraction was performed using the ZymoBIOMICS ®-96 MagBead DNA Kit (Zymo Research, Irvine, CA) and 16S targeted sequencing was completed using the *Quick-*16S™ NGS Library Prep Kit (Zymo Research, Irvine, CA) and V3-V4 16S primers (Zymo Research, Irvine, CA). The sequencing library was prepared using real-time PCR. Final PCR products were quantified with qPCR fluorescence readings and pooled together based on equal molarity. The final pooled library was cleaned with the Select-a-Size DNA Clean & Concentrator™ (Zymo Research, Irvine, CA) which was then quantified with TapeStation® (Agilent Technologies, Santa Clara, CA) and Qubit® (Thermo Fisher Scientific, Waltham, WA). The final library was sequenced on Illumina® MiSeq™ with a v3 reagent kit (600 cycles) and sequencing was performed with 10% PhiX spike-in. Unique amplicon sequence variants were inferred from raw reads using the DADA2 pipeline (Callahan et al., 2016). Potential sequencing errors and chimeric sequences were also removed with the DADA2 pipeline. Taxonomy assignment was performed using Uclust from Qiime (v.1.9.1) and referenced with the Zymo Research Database (Zymo Research, Irvine, CA).

2.4. Neonatal Huddling Behavior

Two-hours after being reunited with their dams on P10 and P11, male and female offspring were quickly weighed and evaluated on their huddling behavior (n = 7-8 full litters). This paradigm was adapted from Naskar and colleagues (2019). Briefly, litters were uniformly placed along the perimeter of a 40 cm x 40 cm area and video recorded for 10 minutes. The average number of pup clusters (2 or more pups in physical contact) was determined by extracting one video frame every 30 seconds and averaging the number of clusters observed across 20 frames. The average time spent together was determined by calculating the average time (in seconds) each pup spent in a clump for the entire 10-minute video.

2.5. Maternal Behavior

To evaluate whether lactational MIA exposure affected maternal behavior, we monitored passive maternal care between P10 and P11 (n = 7-8). The first observation took place on the afternoon of P10 (15:00hrs), following the inflammatory challenge. Maternal care observations took place again at 20:00hrs, and then at 07:30hrs on P11. Following the procedures of Strzelewicz et al (2021; 2019), each session consisted of six observations and a total composite score was calculated for each of the morning, afternoon, and evening time points. Dams were evaluated for 1-minute intervals per observation, with at least 5 minutes of no observations occurring between each of the 1-minute bins. Maternal care observations

recorded included the frequency of pup-directed behaviors (i.e., dam licking/grooming pup, active/high crouch nursing, passive/low crouch nursing, pup retrieval), self-directed behaviors (i.e., dam eating/drinking, dam self-grooming), and nest building/digging behavior. Total time the dam spent on nest (seconds) was also recorded.

2.6. Adolescent Open Field and Social Behavior

On P43, male and female offspring were habituated to an open field arena for five-minutes (40 cm × 40 cm × 28 cm; Duque-Wilckens et al., 2020; Williams et al., 2020; n = 7-8). An automated behavioral monitoring software program (Cleversys TopScan, Reston, VA) was used to evaluate animals on their duration spent (seconds) and frequency of crosses made in each of the center and perimeter of the arena. Percent time spent in the center of the arena and total distance traveled (cm) were also evaluated.

Immediately following the open field habituation period, two clean wire cups were placed on opposite ends of the arena to evaluate social preference (adapted from Crawley, 2007). One cup contained a novel object, and the other cup contained a novel rat of the same sex, age, and strain. Placement of the cups were counterbalanced between trials. Using a manual behavioral monitoring program (ODLog™ 2.0, http://www.macropodsoftware. com/), active investigation was recorded when an experimental rat directed its nose within 2 cm of a wire cup, or it was touching the cup. A social preference index was calculated by the formula ([time spent with the rat] / [time spent with the inanimate object + time spent with the rat]) - 0.5 (Scarborough et al., 2020).

2.7. Mechanical Allodynia

The next day, each animal was acclimatized for 30 minutes to an acrylic cage, with a wire grid floor. Mechanical allodynia was assessed using a pressure-meter which consisted of a hand-held force transducer fitted with a polypropylene rigid tip (Electronic von Frey Aesthesiometer, IITC, Inc, Life Science Instruments, Woodland Hills, CA, USA). The polypropylene tip was applied vertically to the central area of the animal's left hind paw with increasing force. The trial ended when the rat withdrew their paw from the tip, at which point the intensity of the stimulus was automatically recorded by the electronic pressure-meter. The average of four test trials was calculated as the mechanical withdrawal threshold (grams; Yan & Kentner, 2017).

2.8. Prepulse Inhibition of the Acoustic Startle Reflex

Three hours following the evaluation of mechanical allodynia, animals were placed into startle chambers and evaluated on prepulse inhibition (PPI) of the acoustic startle reflex (San Diego Instruments, San Diego, CA, USA). Sessions each consisted of pulse-alone, prepulse-plus-pulse and prepulse-alone trials, as well as no-stimulus trials in which only a background noise of 65 dB was presented Giovanoli et al., (2013). The acoustic startle software administered one 40-ms pulse of white noise (120 dB) in combination with one of five different prepulses. The prepulses were made up of a 20-ms burst of white noise at five different intensities (69, 73, 77, 81, 85 dB), corresponding to 4, 8, 12, 16, and 20 dB above the background noise). After a startle habituation phase, each trial stimulus was pseudorandomly presented 12 times. The average interval between successive trials (ITI) was 15 ± 5 sec. Sessions terminated with 6 consecutive pulse-alone trials. PPI was calculated as percent inhibition of the startle response obtained in the prepulse-plus-trials compared to pulse-alone trials: [1-(mean reactivity on prepulse-plus-pulse trials / mean reactivity on pulse-

alone trials) $\times$ 1/100] and expressed as % PPI for each animal at each of the five possible prepulse intensities.

2.9. Conditioned Fear

Fear conditioning has been shown to modulate PPI of the acoustic startle reflex (Ishii et al., 2010; Balogh et al., 2002). At the conclusion of the PPI session, animals were allowed to rest for 2 hours before commencing Day 1 of fear conditioning trials. On Day 1, animals received 10 trials of a light stimulus (CS) paired with a 0.6 mA foot shock (US). The next day (Day 2), animals completed the PPI task for a second time, with half of the pseudorandom trials beginning with the presentation of the CS in order to assess how the element of fear influences PPI of the acoustic startle reflex. Percent PPI was calculated (described above) for trials with and without the CS.

2.10. Tissue Collection

On P46, a mixture of ketamine/xylazine (150 mg/kg, i.p./50 mg/kg, i.p.) was used to anesthetize animals. Whole hippocampus was dissected, frozen on dry ice, and stored at −80°C for future processing. Maternal blood was collected at weaning in EDTA coated tubes following a cardiac puncture and spun at 1,000g for 10 minutes to obtain plasma.

2.11. Western Blotting

Offspring hippocampus was homogenized in a radioimmunoprecipitation assay buffer (RIPA) with protease inhibitor added (Thermo Fisher Scientific, Cat. #87786). Protein values were determined using the BCA assay (Pierce™, Cat. #23227). Samples were mixed with an equal volume of 2x Laemmli sample buffer (Bio Rad Laboratories Cat. #1610737) and denatured at 100°C for 5 minutes. 20µg of protein was loaded into each well of Mini-Protean® gels (Bio Rad Laboratories, Cat. #4568101) and transferred onto nitrocellulose membranes (Bio Rad Laboratories, Cat. #1620147). Membranes were blocked in 5% nonfat milk with TBS + 0.05% Tween 20 (TBST) for 1 hour at room temperature, washed with TBST, and incubated in a 1:1000 dilution of parvalbumin (PV) antibody (RnD Systems, Cat. # AF5058) plus TBS overnight at 4°C. The next morning, membranes were washed with TBST and incubated in an HRP-conjugated secondary antibody (1:1000, RnD Systems, Cat., #HAF016) made in 1% nonfat milk with TBS for 1 hour at room temperature. Membranes were then washed with TBST and exposed to a chemiluminescent substrate (Thermo Fisher Scientific, Cat. #34580) for 5 minutes prior to being scanned. After imaging, membranes were stripped (Thermo Fisher, Cat. # 21062) for 15 minutes at 37°C, blocked in 5% nonfat milk with TBST for 1 hour at room temperature, washed, and incubated in beta actin primary antibody (1:1000, Thermo Fisher Scientific, Cat. #MA515739) for 1 hour at room temperature. Membranes were exposed again to the chemiluminescence substrate for 5 minutes and imaged. Densitometry was used to obtain a ratio of PV/actin in order to quantify differences between groups.

2.12. Statistical analyses

Statistics were performed using the software package Statistical Software for the Social Sciences (SPSS) version 26.0 (IBM, Armonk, NY) or GraphPad Prism (version 9.0). Because the dataset was not powered to evaluate sex-differences directly, male and female animals were evaluated separately (Ordoñes Sanchez et al., 2021). Data are depicted as each sex separately and also collapsed across sex for display purposes. Two-way repeated measure

ANOVAs (MIA x Time) were used to evaluate milk and behavioral measures across P10 and P11. Violations to the assumption of sphericity were addressed using the Huyndt Feldt correction.

One-way ANOVAs were used as appropriate for all other measures unless there were violations to the assumption of normality (Shapiro-Wilk test) in which case Kruskal-Wallis tests were employed (expressed as X^2). The von Frey threshold data was assessed using body weight as a covariate. Data are graphically expressed as mean ± SEM. The partial eta-squared (n_p^2) is also reported as an index of effect size for the ANOVAs (the range of values being 0.02 = small effect, 0.13 = moderate effect, 0.26 = large effect; Miles and Shevlin, 2001).

For microbiome sequencing, samples underwent composition visualization, in addition to alpha-diversity and beta-diversity analyses using Qiime (v.1.9.1) and statistical comparisons were performed using Kruskal-Wallis (Caporaso et al., 2010). Linear discriminant analysis effect size (LEfSe; http://huttenhower.sph.harvard.edu/lefse/) was utilized to determine significant differences in taxonomy abundance between each group as previously described (Segata et al., 2011; Schellekens et al., 2021). Briefly, LEfSe uses a series of nonparametric tests to create a model that identifies taxa that are most likely to explain differences between experimental groups (Segata et a., 2011).

3. Results

3.1. MIA challenge affected maternal care and offspring huddling behavior

Immune challenge during lactation significantly impacted the number of maternal care bouts directed towards pups. For example, a MIA by time interaction was identified for the number of licks pups received following maternal LPS exposure (F(2, 26) = 31.435, p = 0.001, n_p^2 = 0.701; **Figure 1B**). While the number of licks received was not significantly affected on P10 (p >0.05), there was a general pattern of reduced care provided by MIA treated dams on the afternoon of P10 (p = 0.063). This was followed by a rebound of maternal pup licking by MIA dams on the morning of P11 (p = 0.001), possibly to compensate for overall reduced maternal care due to illness. The display of both active and low crouch nursing behaviors was affected by the circadian cycle (active: F(2, 26) = 7.752, p = 0.0022, n_p^2 = 0.374; low: F(2, 26) = 3.960, p = 0.032, n_p^2 = 0.233: **Figure 1C,D**), but not MIA experience (p>0.05). Neither time of day nor MIA affected the duration spent on the nest (p>0.05; **Figure 1E**). Maternal self-directed care was impacted as a function of sickness. There was a significant MIA by time interaction for the total number of self-grooms (F (1.322, 26) = 12.33, p = 0.001, n_p^2 = 0.485; Huyndt Feldt correction; **Figure 1F**). MIA dams had fewer bouts of grooming on the evening of P10 (p = 0.025), which reversed substantially by the morning of P11 (p = 0.0001). The number of maternal eating bouts also seemed to rebound in a compensatory manner following MIA challenge, although there was no significant effect of time (p>0.05; Figure 1G). There was however, a main effect of MIA (F(1, 13) = 10.64, p = 0.006, n_p^2 = 0.450; **Figure 1G**)). Additional maternal care data are presented in **Supplementary Figure 1A-D**.

Huddling behavior was significantly reduced in the offspring of LPS treated dams compared to saline (main effect of MIA; time spent huddling: F(1, 12) = 30.079, p = 0.001, n_p^2 = 0.715; **Figure 1H**; average number of clusters: F(1, 12) = 8.33, p = 0.014, n_p^2 = 0.410; **Figure 1I**). See **Figure 1J** for representative displays of huddling behavior in saline and LPS offspring.

3.2. MIA challenge impacted nutritional composition of milk and microbiome communities

For milk collection (see photographs in **Figure 2A** depicting this procedure), there was a significant MIA by time interaction for maternal milk levels of corticosterone (F(1, 13) = 8.496, p = 0.013, n_p^2 = 0.415; **Figure 2B**). This stress-associated hormone was elevated in milk samples 3-hours following MIA challenge ((F(1, 13) = 9.512, p = 0.009), and was recovered by P11 (p>0.05). A MIA by time interaction was also present for % creamatocrit (F(1, 13) = 5.86, p = 0.032, n_p^2 = 0.858; **Figure 2C**), which is linearly related to the fat concentration and energy content of milk (Lucas et al., 1978; Paul et al., 2015; see **Supplementary Figure 2A, B**). These measures were significantly lower in the milk of MIA exposed mothers compared to saline on P11 (p = 0.002). Triglycerides were also reduced in the milk of MIA treated dams on P11 (main effect of MIA: F(1, 13) = 6.496, p = 0.026, n_p^2 = 0.351; **Figure 2D**).

In order to examine the possibility that MIA induced expression of inflammatory cytokines, that could be transferred to nursing offspring, we analyzed the concentration of milk IL-6. Although cytokine expression in milk does not reflect levels found in the blood, chronic in vitro stimulation of human breast milk cells with LPS can induce the expression of this cytokine (Hawkes et al., 2002). While there were no significant differences in IL-6 concentration between saline and MIA treated dams (p > 0.05), there was a main effect of time on this measure (F(1,13) = 33.01, p < 0.001, n_p^2 = 0.717; **Figure 2E**). We also tested the idea that bound LPS could cross into maternal milk and be absorbed by offspring. LPS binds directly to LPS binding protein (LBP) forming a complex which can trigger a downstream pro-inflammatory response and the release of acute phase proteins like IL-6 (Wurfel & Wright, 1997; Park & Lee, 2017). LPB was significantly elevated in the blood plasma of LPS-exposed dams on P10 (X^2(1)=5.00, p=0.025; **Figure 2F**). In milk, LPB was absent in both MIA and saline mothers at this time point (p>0.05). Additional data on maternal milk composition (i.e., protein, lactose, and IgA concentrations) can be found in **Supplementary Figure 2C-E.**

Microbiome sequencing of milk samples did not reveal an effect of MIA on either alpha (**Figure 2G**) or beta diversity (main effect of MIA: p > 0.05; **Figure 2H, I**). However, milk samples collected on P11 demonstrated significantly greater alpha diversity along the Shannon index compared to samples collected on P10 (main effect of time: X^2(1) = 5.33, p = 0.021; **Figure 2G**)). Between treatment groups, LEfSE analysis identified 9 differently abundant taxa on P10 and 6 significantly abundant taxa on P11 **(Figure 2J, K)**. These analyses revealed a greater abundance of *Pseudomonadaceae* (LDA score = 2.94, p=0.02) and *Christensenellaceae* (LDA score = 2.82, p = 0.02) on P10, and a greater expression of *Stenotrophomonas maltophilia* (LDA score= 3.45, p=0.025), *Ruminococcaceae* (LDA score = 3.35, p = 0.02) and *Lachnospiraceae* (LDA score= 3.85, p = 0.01) on P11 in the milk from MIA dams. For additional data on the composition of the milk microbiome, see **Supplementary Figures 3-6.**

3.4. The combination of MIA challenge and nutritional deficit affected adolescent offspring physiology and behavior.

Male, but not female, MIA offspring were significantly heavier than saline controls on P43 (one-way ANOVA main effect of MIA: males: F(1,13) = 19.037, p = 0.001, n_p^2 = 0.594; females: p > 0.05; **Figure 3A**). Using body weight as a covariate, female offspring from LPS treated dams were found to have higher thresholds in the von Frey test compared to their

saline counterparts (females: $F(1, 12) = 8.925$, p $=0.011$, $n_p^2 = 0.427$; males: p > 0.05; **Figure 3B**). This is suggestive of reduced sensitivity to tactile stimulation.

Although there were no differences in distance travelled (cm), percent time spent in the center, nor in the duration of time (seconds) spent in the center or perimeter of the open field arena (p>0.05; see **Supplementary Figure 7A-D**), male offspring of MIA exposed mothers displayed an increased number of crosses into both the center ($F(1, 13) = 5.744$, p $= 0.032$, $n_p^2 = 0.306$; **Figure 3C**) and perimeter ($F(1, 13) = 6.513$, p $= 0.024$, $n_p^2 = 0.334$; **Figure 3D**) of the arena (females: p > 0.05). Social preference was not affected by the early life experience for either sex (p > 0.05; **Figure 3E**).

Sensorimotor gating was evaluated using PPI of the acoustic startle reflex. A significant interaction between MIA and prepulse intensity (males: $F(4, 52) = 4.578$, p $= 0.003$, $n_p^2 = 0.260$; **Figure 3F**) and a significant main effect of prepulse intensity (females: $F(4, 52) = 80.357$, p $= 0.001$, $n_p^2 = 0.861$; **Figure 3G, H** collapsed across sex for display purposes) confirmed that all groups demonstrated an increased % PPI as the intensity was raised from 69 to 85 dB. While females were unaffected (p > 0.05), male offspring originating from MIA exposed mothers had attenuated PPI values compared to saline controls at the 69dB ($F(1, 13) = 17.009$, p $= 0.001$, $n_p^2 = 0.743$), 73dB ($F(1, 13) = 5.927$, p $=0.030$, $n_p^2 = 0.581$), and 81dB ($F(1, 13) = 8.315$, p $= 0.013$, $n_p^2 = 0.633$) intensities. The mean percent prepulse inhibition across all five prepulse intensities was not affected as a function of maternal MIA exposure (p > 0.05; **Figure 3I**).

With respect to conditioned fear, % PPI for trials containing the conditioned stimulus was not significantly different between saline and MIA animals for either sex (p>0.05; **Supplementary Figure 7E** collapsed across sex for display purposes). One-way ANOVA revealed a main effect of trial type ($F(4, 52)= 8.19$, p$=0.008$, $n_p^2=.226$; **Supplementary Figure 7F**), which was expected given that the added element of fear increases % PPI (Ishii et al., 2010; Balogh et al., 2002). Conditioned fear did not lead to significant changes in total mean % PPI when collapsed across all dB, regardless of MIA or sex (p > 0.05; **Supplementary Figure 7E** data collapsed across sex for display purposes), suggesting that fear modulated % PPI similarly regardless of sex or drug treatment. There was however, a main effect of MIA for % PPI during the trials without fear for 69dB (females: $F(1, 13) =12.969$, p $= 0.003$, $n_p^2=0.495$), 73dB (males: $F(1, 13) = 6.787$, p $= 0.022$, $n_p^2 = 0.343$), 81dB (males: $F(1, 13) = 6.787$, p $= 0.022$, $n_p^2=0.343$; females: $F(1, 13) = 25.117$, p $= 0.001$, $n_p^2=0.659$) and 85 dB (males: $F(1, 13) = 13.124$, p $= 0.003$, $n_p^2=0.502$; **Figure 3J, K, L** collapsed across sex for display purposes).

In addition to these behavioral effects, PV/actin ratios were significantly reduced in the hippocampus of male MIA offspring compared to their control counterparts ($F(1, 13) =6.96$, p $= 0.022$, $n_p^2=.367$; **Figure 3M**).

Figure 4 outlines the proposed mechanism driving changes in milk quality that affect offspring developmental outcomes.

4. Discussion

In the present study, we demonstrate that variations in milk quality may serve as part of a mechanism programming offspring physiology and behavior following stressors experienced during the mid-lactational period. For example, we show that MIA was associated with sustained changes in both the nutritional content and microbial profile of maternal milk. Moreover, the MIA-induced changes in milk fat and corticosterone levels were accompanied by physiological and behavioral alterations in offspring. Neonatal offspring from MIA-challenged dams spent less time huddling compared to controls. In adolescence, these offspring were heavier, demonstrated deficits in sensorimotor gating, hippocampal PV

expression, and had increased mechanical allodynia thresholds. Importantly, these MIA-associated effects in offspring were not due to an overexposure to inflammatory cytokines, as milk concentrations of IL-6 and LBP were comparable between MIA and saline dams. Moreover, though a general pattern of reduced licking/grooming could be seen in MIA dams on P10, this effect was non-significant and care quality was overcompensated by P11. Rather, offspring behavior and bodyweight differences likely reflect the unique expression of the nutritional elements and metabolically relevant bacterial taxa found in P10 and P11 milk samples. While most of the MIA-related literature has been dedicated to uncovering the physiological and behavioral effects on offspring gestationally exposed to MIA, fewer studies have focused on the impact of MIA exposure after birth. The lactational period is an important time for neonates and here we highlight parental inputs (e.g., maternal milk quality) that can be altered by stressors in the shared environment, affecting offspring development.

MIA was associated with multiple changes in the composition of maternal milk content. We observed a significant increase in milk corticosterone 3-hours post LPS treatment that remitted by P11. Milk corticosterone is expressed linearly to plasma corticosterone (Patacchioli et al., 1992), validating our MIA model. Importantly, milk corticosterone can pass from a mother to her offspring where these glucocorticoids can remain active in the periphery and in the brain (Angelucci et al., 1985; Brummelte et al., 2010). Increased corticosterone exposure from breastmilk has previously contributed to HPA axis dysfunction as well as altered learning and memory abilities in rat pups of both sexes, independent of maternal care (Catalani et al., 1993; Catalani et al., 2002). In addition to its modulatory actions within the HPA axis, recent evidence has demonstrated the regulatory role of corticosterone in breastmilk, where the presence of this hormone (and the human equivalent, cortisol) can modulate nutritional elements including sodium, potassium, and fat (Zietek et al., 2021; Sullivan et al., 2011; Hinde et al., 2015). This evidence suggests that the temporary rise in corticosterone on P10 may have preceded downstream changes in milk nutritional quality. In line with this, MIA reduced % creamatocrit and milk triglycerides on P11. However, higher levels of glucocorticoids in milk alone are not a sufficient predictor of offspring bodyweight (Brummelte et al., 2010; Hinde et al., 2014), suggesting that the actions of corticosterone in neonates are likely interwoven with milk nutrition. Reductions in milk creamatocrit or triglycerides may instead have been due to direct actions of LPS. Indeed, previous studies have shown LPS to inhibit fat production in the mammary epithelial cells of cows (Liu et al., 2015; Chen et al., 2019) while LPS reduced fatty-acid transporter mRNA in the mammary gland of rats (Ling & Alcorn, 2010). It is unclear however, if inflammatory markers that follow LPS administration into the periphery are capable of crossing into maternal milk in vivo.

We measured levels of LBP and IL-6 in milk to investigate the possibility that maternally derived cytokines, or LPS itself, were being absorbed by offspring and affecting neurodevelopment. LBP was significantly elevated in the plasma of MIA dams compared to saline dams confirming inflammation. However, LBP levels were absent in the milk samples of both groups, suggesting that none, or negligible, amounts of LPS were present in milk. We also measured milk IL-6 levels given that this cytokine is a main component of fetal programming in prenatal MIA models (Wu et al., 2017; Hsiao & Patterson, 2011). Moreover, in the presence of LPS, maternal milk cells can produce high amounts of IL-6 in vitro (Hawkes et al., 2002). The detection of this cytokine in our control samples was not surprising given that IL-6 is involved in typical mammary gland functions such as involution and milk storage (Basolo et al., 1996). Overall, comparable milk IL-6 concentrations between the saline and MIA treatment groups indicate that offspring were not affected by maternal IL-6 levels. Future studies must rule out the possibility that other inflammatory markers (e.g.,

TNF-α and other interleukins) may cross from maternal blood to milk, although it is unlikely (Hawkes et al., 2002). This evidence supports the idea that differences in nutritional quality, and not inflammatory profile, may be more influential in modulating offspring development and behavior in this postnatal MIA model.

Microbiome sequencing did not reveal significant differences in alpha or beta diversity between our saline and MIA treated dams. While general diversity was not affected, LEfSe analysis uncovered significant differences in the expression of certain taxa on P10 and P11. MIA dams exhibited higher levels of *Christensenellaceae,* which is negatively correlated with serum triglycerides (Waters & Ley, 2019). This relationship may reflect the lower levels of triglycerides found in the milk of our MIA dams. Moreover, *Christensenellaceae* are also associated with reduced inflammation and a lower expression of the proinflammatory lipopolysaccharide binding protein (LBP) in the gut (Citronberg et al., 2018; Citronberg et al., 2016). Significant increases in *Christensenellaceae* may have been compensatory to attenuate LBP-signaling and subsequent inflammatory responses. Further, milk from MIA dams exhibited elevated levels of *Stenotrophomonas maltofilia*, which has the ability to degrade hydrocarbon chains and therefore could prevent triglyceride synthesis (Singh & Fulekar et al., 2010). The increased expression of *Lachnospiraceae* in the milk of LPS-treated MIA dams may in part explain the offspring bodyweight differences at P43 as this bacterium is associated with inefficient lipid and energy metabolism (Vacca et al., 2020; Palmas et al., 2021). Given that the microbes in breast milk directly colonize the infant gut (Sheard & Walker, 1988; Pannaraj et al., 2017), it is likely that these differently expressed taxa are being passed to the offspring to influence future metabolic processes. Additional studies are needed to explore potential therapeutic interventions, such as targeting the microbiome (Holingue et al., 2020) and the use of nutritional supplements, which may rescue changes in milk composition and subsequent offspring developmental outcomes.

We used several behavioral measures to explore the role of lactational MIA exposure on offspring development. Neonatal 'huddling' or 'clumping' is described as a form of social thermoregulation (Gilbert et al., 2009). One previous study found a reduction in nest seeking behavior in rat pups prenatally exposed to LPS (Baharnoori et al., 2012). Undernourishment has also been shown to reduce neonatal huddling in rat pups (Soriano et al., 2006), and huddling efficiency may be related to serum triglycerides in newborn rabbits (García-Torres et al., 2015). This evidence suggests that MIA-modulated huddling may emerge due to differences in the fat content of maternal milk. Here, pups from MIA treated mothers displayed disorganized clumping behavior. Impairments in huddling have also been associated with delayed development of glutamatergic and GABAergic signaling in the prefrontal cortex and hippocampus (Naskar et al., 2019; Baharnoori et al., 2009), which may have long-term implications for offspring in adulthood. Indeed, deficits in neonatal huddling may be predictive of other behavioral alterations later in life (Schatz et al., 2018).

Prenatal MIA has been shown to modulate offspring performance on certain tasks, such as the open field, social preference, von Frey, PPI, and fear conditioning paradigms (Kentner et al., 2018; Fortier et al., 2007; Zhao et al., 2021a; Inceoglu et al., 2006; Howland et al., 2012). Here, we observed significant MIA induced reductions in % PPI that were exacerbated in male adult offspring while females demonstrated deficits in mechanical allodynia. MIA was also associated with a significant reduction in % PPI in trials without fear, following the foot shock and conditioned stimulus (CS) pairing. These findings suggest MIA contributes to an impairment in the offspring's ability to discriminate between "safe" trials (without CS presentation) and "stressful" trials (with CS presentation). Early life stress has been shown to impair this form of contextual threat detection in adolescence, and this is especially true for females (Manzano-Nieves et al., 2018; Gerhard et al., 2020). These results, in addition to

reduced neonatal huddling, point to generally impaired sensory processing and threat response in the offspring of dams exposed to MIA during lactation.

We measured the expression of GABAergic PV interneurons in the offspring hippocampus since altered PV expression in this region is a common theme in prenatal MIA models and is associated with many MIA-associated behavioral and sensory processing changes (Zhao et al., 2021b; Nakamura et al., 2021; Vojtechova et al., 2021). Further, poor nutrition in early life can prevent the development of PV interneurons in rat hippocampus (Callahan et al., 2013) although this has only been explored in males. The emergence of neonatal huddling behavior is time-locked to the maturation of GABAergic signaling (Naskar et al., 2019). Therefore, mid-lactational MIA may disrupt this early form of social behavior by attenuating the development of PV cells. In adolescence, male MIA offspring demonstrated a reduced expression in hippocampal PV interneurons. In vivo inhibition of PV in the hippocampus has previously been shown to reduce PPI in male mice (Nguyen et al., 2014). Together, these results demonstrate one neuronal target by which MIA during mid-lactation contributes to lasting changes in offspring brain development in males. More work is needed to discern the neuronal targets of mid-lactational MIA in females.

We found no differences between saline and MIA offspring performance in the social preference paradigm, and MIA males exhibited less anxiolytic behavior as indicated by their increased number of crosses into the center of the open field task. While in sharp contrast to studies that have assessed gestational MIA, these results are not entirely surprising given the physiological differences between the pre and postnatal actions of MIA. For example, MIA in utero reorganizes the molecular construction of the placenta (Núñez Estevez et al., 2020; Hsaio & Patterson, 2011; Garay et al., 2013) and the results of the present study suggest lactational MIA targets milk quality and offspring nutrition. While complimentary, our results highlight the importance of the timing of MIA challenge to offspring developmental and behavioral programming.

Moreover, our MIA dams showed significant increases in pup-directed licking and grooming behavior the day following MIA challenge. The heightened maternal care demonstrated by MIA dams on P11 may have been a compensatory mechanism to buffer the effects of lactational MIA exposure. Tactile stimulation through the licking and grooming of pups is one form of maternal behavior that promotes brain development and modulates future offspring behavior (Champagne et al., 2003; Francis et al., 2002; Van Hasselt et al., 2012; Francis & Kuhar, 2008). Aside from a general pattern of decreased (but not significant) licking and grooming of pups on the afternoon of P10, we did not observe a drastic reduction in pup directed behaviors in MIA dams. The effects of LPS are highly dose dependent (Bison et al., 2009). While Vilela and colleagues (2013) observed more dramatic deficits in maternal care following MIA, they implemented a dose of LPS 5-times higher than the dose used in the present study. Moreover, lactational exposure to LPS using the same dose as that used in the current study also resulted in improved maternal care (Nascimento et al., 2015). Together, our results suggest that stressors experienced during the neonatal period may interact with maternal care and lactation quality, affecting offspring physiology and behavior. Indeed, early life stress models affect both the parent(s) and the offspring; the dynamic impacts on each need to be considered as part of the mechanistic programming of the developmental trajectory.

Conclusions

Maternal care and milk quality are important components to offspring development, brain health, and behavior. Here, we demonstrate that an acute maternal immune response during mid lactation is sufficient to trigger changes in breastmilk quality and its microbiome profile,

in addition to the physiology and behavior of adolescent offspring. These results contribute to the broader literature by suggesting that the effects of MIA on offspring development are not restricted to the prenatal window. Moreover, this study ties in the characteristics of lactation and nutrition as part of a mechanism contributing to the trajectory of offspring development following early life stress. Given that animal models of early life stress can impact both parents and their offspring, the quality of maternal milk should be considered among the variables investigated in future studies. Finally, this work underscores the importance of research focused on potential therapeutic interventions (e.g., nutritional lactation supplements) and necessitates a better representation of pregnant and nursing people in both basic and clinical research.

Funding and Disclosures

This project was funded by NIMH under Award Number R15MH114035 (to ACK). The authors wish to thank Mary Erickson, Sahith Kaki, Ada Cheng, and Dr. Theresa Casey for their technical support. The authors would also like to thank the MCPHS University School of Pharmacy and School of Arts & Sciences for their continual support. The content is solely the responsibility of the authors and does not necessarily represent the official views of any of the financial supporters.

Author Contributions
H.D., H.T., A.C.K., ran the experiments; H.D., & A.C.K. analyzed and interpreted the data, and wrote the manuscript; A.C.K., designed and supervised the study.

Declaration of Competing Interest
The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Figure Captions

Figure 1. Maternal care and offspring huddling behaviors following maternal immune activation (MIA) during the mid-lactational period. (A) Timeline of experimental procedures. Total number of maternal pup-directed behaviors (B) pup licks, (C) active nursing bouts, (D) low crouch nursing bouts, and (E) total time (seconds) that dam spent on nest. Total number of maternal self-directed behaviors (F) self-groom bouts and (G) eating bouts (n = 7-8). Maternal care was evaluated in the afternoon and night of postnatal day (P) 10 and the morning of P11. Pup huddling such as (H) total time spent huddling (seconds) and the (I) average number of clusters was evaluated across P10 and P11 following MIA (n = 7-8 full litters). (J) Shows representative photographs of the huddling behavior of saline and LPS litters. Data are expressed as mean ± SEM. *$p < 0.05$, ***$p < 0.001$, LPS versus Saline; [a]$p < 0.05$, [aa]$p < 0.01$, main effect of time; [b]$p < 0.05$, [bb]$p < 0.01$, [bbb]$p < 0.001$, main effect of MIA.

Figure 2. Nutritional profile of milk and microbiome community distribution following maternal immune activation (MIA) during the mid-lactational period. (A) Photographs of maternal milk collection. Maternal milk levels of (B) corticosterone (ng/mL), (C) percent creamatocrit, (D) triglycerides (mg/dL), (E) interleukin (IL)-6 (pg/mL), and (F) LPS binding

protein (LPB;ug/mL) (n = 7). (G) Alpha diversity of P10 and P11 milk samples along the Shannon Index. Beta diversity plots using principal coordinate analysis (PCoA) of (H) P10 and (I) P11 milk samples were created using the matrix of pair-wise distances between samples determined by Bray-Curtis dissimilarity using unique amplicon variants. (J) Microbial composition of taxonomy at the family level for saline and LPS dams at P10 and P11 (n = 6-7). (K) LEfSe biomarkers plots. These figures display taxa that are significantly more abundant in the milk of saline-treated dams (blue bars) versus LPS treated dams (red bars) on P10 and P11. These taxa were identified based on their significant distributions ($p < 0.05$) and effect sizes (LDA score) larger than 2 (n = 6-7). Data are expressed as mean $\pm$ SEM. *$p < 0.05$, ***$p < 0.001$, LPS versus Saline; [a]$p < 0.05$, [aa]$p < 0.01$, main effect of time; [b]$p < 0.05$, [bb]$p < 0.01$, [bbb]$p < 0.001$, main effect of MIA.

Figure 3. Adolescent offspring physiology and behavior following maternal immune activation (MIA) during the mid-lactational period. (A) P43 body weights (grams), (B) mechanical paw withdrawal thresholds (grams) on the von Frey test, number of crosses into the (C) center and (D) perimeter of an open field arena, and (E) social preference index. Line plots displaying percent prepulse inhibition (PPI) as a function of increasing prepulse intensities for (F) males, (G) females, and (H) males and females combined. (I) The bar plot shows the mean percent prepulse inhibition across all five prepulse intensities. Conditioned fear % PPI for (J) males, (K), females, and (L) males and females combined. (M) Hippocampal parvalbumin interneuron and densitometric ratios for male and female offspring from saline (black) and LPS (grey) treated dams. Data are expressed as mean $\pm$ SEM, n = 7-8. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, LPS versus Saline. [a]$p < 0.05$, [aa]$p < 0.01$, [aaa]$p < 0.001$, main effect of trial type.

Figure 4. proposed mechanism driving changes in milk quality that affect offspring developmental outcomes. Maternal immune activation during the mid-lactational period elevates *Christensenellaceae* in a compensatory fashion, resulting in reduced lipopolysaccharide binding protein, triglycerides, and creamatocrit concentrations in maternal milk. This causes nutritional deficits which impacts the physiological and behavioral developmental trajectory of offspring.

Supplementary Table S1. Maternal Immune Activation Model Reporting Guidelines Checklist.

Supplementary Figure 1. Maternal care behaviors following maternal immune activation (MIA) during the mid-lactational period. Total number of (A) pup retrievals, (B) passive nursing bouts, (C) nest building behaviors, and (D) maternal drinking bouts. Maternal care was evaluated in the afternoon and night of postnatal day (P) 10 and the morning of P11. Data are expressed as mean $\pm$ SEM, n = 7-8. [a]$p < 0.05$, main effect of time.

Supplementary Figure 2. Nutritional profile of milk following maternal immune activation (MIA) during the mid-lactational period. Maternal milk (A) fat concentration (g/L), (B) energy value (kcal/L), (C) protein concentration (mg/mL), (D) lactose concentration (ng/μL), and IgA concentration (mg/mL). Data are expressed as mean $\pm$ SEM, n = 7-8. *$p < 0.05$, ***$p < 0.001$, LPS versus Saline; [a]$p < 0.05$, main effect of time.

Supplementary Figure 3. Cladogram of milk biomarkers associated with treatment group on P10 (determined by LEfSe). Diameter of the nodes indicates relative abundance of taxa for

saline (green) and LPS (red) samples. Placement indicates the classification of taxa, where nodes decrease in rank the closer to the center of the diagram.

Supplementary Figure 4. Taxonomy heatmap demonstrating the top fifty most abundant species identified in samples on P10. Treatment group is indicated by the colored bar at the top of the figure (red = saline, blue = LPS). Each row represents the abundance for each taxon, with the taxonomy ID shown on the right. Each column represents the abundance for each sample.

Supplementary Figure 5. Cladogram of milk biomarkers associated with treatment group on P11 (determined by LEfSe). Diameter of the nodes indicates relative abundance of taxa for saline (green) and LPS (red) samples. Placement indicates the classification of taxa, where nodes decrease in rank the closer to the center of the diagram.

Supplementary Figure 6. Taxonomy heatmap demonstrating the top fifty most abundant species identified in samples on P11. Treatment group is indicated by the colored bar at the top of the figure (red = saline, blue = LPS). Each row represents the abundance for each taxon, with the taxonomy ID shown on the right. Each column represents the abundance for each sample.

Supplementary Figure 7. Adolescent offspring behavior following maternal immune activation (MIA) during the mid-lactational period. (A) Distance traveled (cm), (B) percent time spent in the center, and duration of time (seconds) spent in the (C) center and (D) perimeter of an open field arena. (E) % PPI collapsed across both sex and drug for all five prepulse intensities. (F) Total % PPI collapsed across sex and all prepulse intensities to show trials without fear (black line) and trials with fear (dotted line). Data are expressed as mean $\pm$ SEM, n = 7-8. *p <0.05, LPS versus Saline.

References

Anderson, G., Vaillancourt, C., Maes, M., & Reiter, R. J. (2017). Breastfeeding and the gut-brain axis: is there a role for melatonin? Biomolecular concepts, 8(3-4), 185-195. https://doi.org/10.1515/bmc-2017-0009

Angelucci, L. (1985). A model for later-life effects of perinatal drug exposure: maternal hormone mediation. *Neurobehavioral Toxicology & Teratology*.

Arsenault, D., St-Amour, I., Cisbani, G., Rousseau, L.-S., & Cicchetti, F. (2014). The different effects of LPS and poly I:C prenatal immune challenges on the behavior, development and inflammatory responses in pregnant mice and their offspring. *Brain, Behavior, and Immunity*, *38*, 77–90. https://doi.org/10.1016/j.bbi.2013.12.016

Badillo-Suárez, P.A., Rodríguez-Cruz, M. & Nieves-Morales, X. Impact of Metabolic Hormones Secreted in Human Breast Milk on Nutritional Programming in Childhood Obesity. *Journal of Mammary Gland Biology Neoplasia, 22,* 171–191 (2017). https://doi.org/10.1007/s10911-017-9382-y

Baharnoori, M., Bhardwaj, S. K., & Srivastava, L. K. (2012). Neonatal behavioral changes in rats with gestational exposure to lipopolysaccharide: a prenatal infection model for developmental neuropsychiatric disorders. *Schizophrenia Bulletin, 38*(3), 444–456. https://doi.org/10.1093/schbul/sbq098

Balogh, S. A., Radcliffe, R. A., Logue, S. F., & Wehner, J. M. (2002). Contextual and cued fear conditioning in C57BL/6J and DBA/2J mice: context discrimination and the effects of retention interval. *Behavioral Neuroscience*, *116*(6), 947. doi: 10.1037//0735-7044.116.6.947.

Basolo, F., Fiore, L., Fontanini, G., Conaldi, P. G., Calvo, S., Falcone, V., & Toniolo, A. (1996). Expression of and response to interleukin 6 (IL6) in human mammary tumors. *Cancer Research, 56*(13), 3118-3122. 8674070

Bison, S., Carboni, L., Arban, R., Bate, S., Gerrard, P. A., & Razzoli, M. (2009). Differential behavioral, physiological, and hormonal sensitivity to LPS challenge in rats. *International Journal of Interferon, Cytokine and Mediator Research, 1*, 1-13. https://doi.org/10.2147/IJICMR.S4273

Brown, R., Imran, S. A., & Wilkinson, M. (2009). Lipopolysaccharide (LPS) stimulates adipokine and socs3 gene expression in mouse brain and pituitary gland in vivo, and in N-1 hypothalamic neurons in vitro. Journal of *Neuroimmunology, 209*(1-2), 96-103. https://doi.org/10.1016/j.jneuroim.2009.02.001

Browne, P. D., Aparicio, M., Alba, C., Hechler, C., Beijers, R., Rodríguez, J. M., ... & de Weerth, C. (2019). Human milk microbiome and maternal postnatal psychosocial distress. *Frontiers in Microbiology, 10,* 2333. https://doi.org/10.3389/fmicb.2019.02333

Brummelte, S., Schmidt, K.L., Taves, M.D., Soma, K.K. & Galea, L. A. (2010). Chronic corticosterone during pregnancy and postpartum affects maternal care, cell proliferation and depressive-like behavior in the dam. *Hormones and Behavior, 58*(5), 769-779. https://doi.org/10.1016/j.yhbeh.2010.07.012

Callahan B.J., McMurdie P.J., Rosen M.J., Han A.W., Johnson A.J., Holmes S.P. (2016). DADA2: High resolution sample inference from Illumina amplicon data. *Nature Methods, 13*(7):581-3. https://doi.org/10.1038/nmeth.3869

Callahan, L. S., Thibert, K. A., Wobken, J. D., & Georgieff, M. K. (2013). Early-life iron deficiency anemia alters the development and long-term expression of parvalbumin and perineuronal nets in the rat hippocampus. Developmental neuroscience, 35(5), 427-436. https://doi.org/10.1159/000354178

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K. et al. (2010) QIIME allows analysis of high-throughput community sequencing data. *Nature Methods, 7*: 335-336. https://doi.org/10.1038/nmeth.f.303

Catalani, A., Marinelli, M., Scaccianoce, S., Nicolai, R., Muscolo, L. A., Porcu, A., ... &

Angelucci, L. (1993). Progeny of mothers drinking corticosterone during lactation has lower stress-induced corticosterone secretion and better cognitive performance. *Brain Research, 624*(1-2), 209-215. https://doi.org/10.1016/0006-8993(93)90079-3

Catalani, A., Casolini, P., Cigliana, G., Scaccianoce, S., Consoli, C., Cinque, C., ... & Angelucci, L. (2002). Maternal corticosterone influences behavior, stress response and corticosteroid receptors in the female rat. *Pharmacology Biochemistry and Behavior, 73*(1), 105-114. https://doi.org/10.1016/S0091-3057(02)00755-4

Champagne, F.A. & Curley, J.P. (2009). Epigenetic mechanisms mediating the long-term effects of maternal care on development. *Neuroscience Biobehavioral Reviews, 33*(4):593-600. doi: 10.1016/j.neubiorev.2007.10.009.

Champagne, F. A., Francis, D. D., Mar, A., & Meaney, M. J. (2003). Variations in maternal care in the rat as a mediating influence for the effects of environment on development. *Physiology & Behavior, 79*(3), 359-371. https://doi.org/10.1016/S0031-9384(03)00149-5

Chen, Y., Wang, J., Yang, S., Utturkar, S., Crodian, J., Cummings, S., Thimmapuram, J., San Miguel, P., Kuang, S., Gribskov, M., Plaut, K., Casey, T. (2017). Effect of high fat diet on secreted milk transcriptome in midlactation mice. *Physiology Genomics, 49*, 747-762, doi:10.1152/physiolgenomics.00080.2017.

Chen, J., Wu, Y., Sun, Y., Dong, X., Wang, Z., Zhang, Z., ... & Dong, G. (2019). Bacterial lipopolysaccharide induced alterations of genome-wide DNA methylation and promoter methylation of lactation-related genes in bovine mammary epithelial cells. *Toxins, 11*(5), 298. https://doi.org/10.3390/toxins11050298

Citronberg, J.S., Curtis, K.R., White, E. et al. (2018). Association of gut microbial communities with plasma lipopolysaccharide-binding protein (LBP) in premenopausal women. ISME J 12, 1631–1641 (2018). https://doi.org/10.1038/s41396-018-0064-6.

Citronberg, J. S., Wilkens, L. R., Lim, U., Hullar, M. A., White, E., Newcomb, P. A., ... & Lampe, J. W. (2016). Reliability of plasma lipopolysaccharide-binding protein (LBP) from repeated measures in healthy adults. Cancer Causes & Control, 27(9), 1163-1166. https://doi.org/10.1007/s10552-016-0783-9.

Connor, K. L., Chehoud, C., Altrichter, A., Chan, L., DeSantis, T. Z., & Lye, S. J. (2018). Maternal metabolic, immune, and microbial systems in late pregnancy vary with malnutrition in mice. *Biology of Reproduction, 98*(4), 579-592. https://doi.org/10.1093/biolre/ioy002

Connors, E. J., Shaik, A. N., Migliore, M. M., & Kentner, A. C. (2014). Environmental enrichment mitigates the sex-specific effects of gestational inflammation on social engagement and the hypothalamic pituitary adrenal axis-feedback system. *Brain, Behavior, and Immunity, 42,* 178-190. https://doi.org/10.1016/j.bbi.2014.06.020

Crawley, J.N., 2007. Social behavior tests for mice. In: Crawley, J.N. (Ed.), What's Wrong

with My Mouse? Strategies for Rodent Behavior Phenotyping. Society for Neuroscience, San Diego, pp. 63–70. doi:10.1002/9780470119051

Dockman, R. L., Carpenter, J. M., Diaz, A. N., Benbow, R. A., & Filipov, N. M. (2021). Sex differences in behavior, response to LPS, and glucose homeostasis in middle-aged mice. *Behavioural Brain Research, 113628*.https://doi.org/10.1016/j.bbr.2021.113628

Drugs and Lactation Database (LactMed) [Internet]. Bethesda (MD): National Library of Medicine (US); 2006-. Isoflurane. [Updated 2020 Nov 16]. Bookshelf URL: https://www.ncbi.nlm.nih.gov/books/

Duque-Wilckens, N., Torres, L.Y., Yokoyama, S., Minie, V.A., Tran, A.M., Petkova, S.P., Hao, R., Ramos-Maciel, S., Rios, R.A., Jackson, K., Flores-Ramirez, F.J., Garcia-Carachure, I., Pesavento, P.A., Iniguez, ~ S.D., Grinevich, V., & Trainor, B.C. (2020). Extrahypothalamicoxytocin neurons drive stress-induced social vigilance and avoidance. *PNAS* U S A. 5:202011890. doi: 10.1073/pnas.2011890117.

Edwards, P.D., Lavergne, S.G., McCaw, L.K., Wijenayake, S., Boonstra, R., McGowan, P.O., Holmes, M.M. (2021). Maternal effects in animals: broadening our understanding of offspring programming. *Frontiers in Neuroendocrinology*, 62, 100924, doi: 10.1016/j.yfrne.2021.100924

Estes, M.L. & McAllister, A.K. (2016). Maternal immune activation: Implications for neuropsychiatric disorders. *Science, 353*(6301):772-7. doi: 10.1126/science.aag3194.

Fortier, M. E., Luheshi, G. N., & Boksa, P. (2007). Effects of prenatal infection on prepulse inhibition in the rat depend on the nature of the infectious agent and the stage of pregnancy. *Behavioural Brain Research, 181*(2), 270-277. https://doi.org/10.1016/j.bbr.2007.04.016

Francis, D. D., Young, L. J., Meaney, M. J., & Insel, T. R. (2002). Naturally occurring differences in maternal care are associated with the expression of oxytocin and vasopressin (V1a) receptors: gender differences. *Journal of Neuroendocrinology, 14*(5), 349-353.

Francis, D. D., & Kuhar, M. (2008). Frequency of maternal licking and grooming correlates negatively with vulnerability to cocaine and alcohol use in rats. *Pharmacology Biochemistry and Behavior, 90*(3), 497-500. https://doi.org/10.1016/j.pbb.2008.04.012

Franco, J. G., Fernandes, T. P., Rocha, C. P., Calviño, C., Pazos-Moura, C. C., Lisboa, P. C., Moura, E. G., & Trevenzoli, I. H. (2012). Maternal high-fat diet induces obesity and adrenal and thyroid dysfunction in male rat offspring at weaning. *The Journal of Physiology, 590*(21), 5503–5518. https://doi.org/10.1113/jphysiol.2012.240655

Garay, P. A., Hsiao, E. Y., Patterson, P. H., & McAllister, A. K. (2013). Maternal immune activation causes age-and region-specific changes in brain cytokines in offspring throughout development. *Brain, Behavior, and Immunity, 31*, 54-68. https://doi.org/10.1016/j.bbi.2012.07.008

García-Torres, E., Hudson, R., Castelán, F., Martínez-Gómez, M., & Bautista, A. (2015). Differential metabolism of brown adipose tissue in newborn rabbits in relation to position in the litter huddle. *Journal of Thermal Biology, 51*, 33-41. https://doi.org/10.1016/j.jtherbio.2015.03.003

Gerhard, D. M., Meyer, H. C., & Lee, F. S. (2020). An adolescent sensitive period for threat responding: impacts of stress and sex. Biological Psychiatry. doi:10.1016/j.biopsych.2020.10.003

Gilbert, C., McCafferty, D., Le Maho, Y., Martrette, J. M., Giroud, S., Blanc, S., & Ancel, A. (2010). One for all and all for one: the energetic benefits of huddling in endotherms. *Biological Reviews, 85*(3), 545-569. https://doi.org/10.1111/j.1469-185X.2009.00115.x

Giovanoli, S., Engler, H., Engler, A., Richetto, J., Voget, M., Willi, R., Winter, C., Riva, M.A., Mortensen, P.B., Feldon, J., 2013. Stress in puberty unmasks latent neuropathological consequences of prenatal immune activation in mice. *Science, 339*, 1095-1099. doi: 10.1126/science.1228261

Grazia Di Benedetto, M., Bottanelli, C., Cattaneo, A., Pariante, C.M., Borsini, A. (2020). Nutritional and immunological factors in breast milk: a role in the intergenerational transmission from maternal psychopathology to child development. *Brain, Behavior, and Immunity*, 85, 57-68.

Hawkes, J.S., Bryan, DL. & Gibson, R.A. (2002). Cytokine production by human milk cells and peripheral blood mononuclear cells from the same mothers. *Journal of Clinical Immunology, 22*, 338–344. https://doi.org/10.1023/A:1020652215048

Hinde, K., Skibiel, A. L., Foster, A. B., Del Rosso, L., Mendoza, S. P., & Capitanio, J. P. (2015). Cortisol in mother's milk across lactation reflects maternal life history and predicts infant temperament. *Behavioral Ecology, 26*(1), 269-281. https://doi.org/10.1093/beheco/aru186

Holingue, C., Brucato, M., Ladd Acosta, C., Hong, X., Volk, H., Mueller, N. T., ... & Fallin, M. D. (2020). Interaction between Maternal Immune Activation and Antibiotic Use during Pregnancy and Child Risk of Autism Spectrum Disorder. *Autism Research, 13*(12), 2230-2241. https://doi.org/10.1002/aur.2411

Howland, J. G., Cazakoff, B. N., & Zhang, Y. (2012). Altered object-in-place recognition memory, prepulse inhibition, and locomotor activity in the offspring of rats exposed to a viral mimetic during pregnancy. *Neuroscience, 201,* 184-198. https://doi.org/10.1016/j.neuroscience.2011.11.011

Hsiao, E.Y. & Patterson, P.H. (2011). Activation of the maternal immune system induces endocrine changes in the placenta via IL-6. *Brain, Behavior, and Immunity, 25*(4), 604-15. doi: 10.1016/j.bbi.2010.12.017.

Hussein J, Mavalankar DV, Sharma S, D'Ambruoso L. (2011). A review of health system infection control measures in developing countries: what can be learned to reduce maternal mortality. *Global Health*, 7, 14.

Inceoglu, B., Jinks, S. L., Schmelzer, K. R., Waite, T., Kim, I. H., & Hammock, B. D. (2006). Inhibition of soluble epoxide hydrolase reduces LPS-induced thermal hyperalgesia and mechanical allodynia in a rat model of inflammatory pain. *Life Sciences, 79*(24), 2311–2319. https://doi.org/10.1016/j.lfs.2006.07.031

Ishii, D., Matsuzawa, D., Fujita, Y., Sutoh, C., Ohtsuka, H., Matsuda, S., ... & Shimizu, E. (2010). Enhancement of acoustic prepulse inhibition by contextual fear conditioning in mice is maintained even after contextual fear extinction. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *34*(1), 183-188.

John, C.C., Black, M.M., Nelson, C.A III. (2017). Neurodevelopment: the impact of nutrition and inflammation during early to middle childhood in low-resource settings. *Pediatrics*, 139, e 20162828.

Kentner, A.C., Bilbo, S.D., Brown, A.S., Hsiao, E.Y., McAllister, A.K., Meyer, U., Pearce, B.D., Pletnikov, M.V., Yolken, R.H., Bauman, M.D. (2019). Maternal immune activation: reporting guidelines to improve the rigor, reproducibility, and transparency of the model. *Neuropsychopharmacology 44*, 245-258.

Kentner, A.C., Scalia, S., Shin, J., Migliore, M.M., z7 Rondón-Ortiz, A.N. (2018). Targeted sensory enrichment interventions protect against behavioral and neuroendocrine consequences of early life stress. *Psychoneuroendocrinology, 98*:74-85. doi: 10.1016/j.psyneuen.2018.07.029.

Karolinski A, Mazzoni A, Belizan JM, Althabe F, Bergel E, Buekens P (2010). Lost opportunities for effective management of obstetric conditions to reduce maternal mortality and severe maternal morbidity in Argentina and Uruguay. Int J Gynaecol Obstet, 110, 175–180.

Kutlesic, V., Breinski Isaacs, M., Freund, L.S., Hazra, R., Raiten, D.J. (2017). Executive summary: research gaps at the intersection of pediatric neurodevelopment, nutrition, and inflammation in low resource settings. Pediatrics, 139, S1-S11.

Lee, J.J. & Rubin, A.P. (2993). Breast feeding and anesthesia. Anaesthesia, 48, 616-625, doi: 10.1111/j.1365-2044.1993.tb07130.x.

Lesorogol, C., Bond, C., Dulience, S. J. L., & Iannotti, L. (2018). Economic determinants of breastfeeding in Haiti: the effects of poverty, food insecurity, and employment on exclusive breastfeeding in an urban population. *Maternal & Child Nutrition, 14*(2), e12524. https://doi.org/10.1111/mcn.12524

Liu, L., Lin, Y., Liu, L., Bian, Y., Zhang, L., Gao, X., & Li, Q. (2015). 14-3-3γ regulates lipopolysaccharide-induced inflammatory responses and lactation in dairy cow mammary epithelial cells by inhibiting NF-κB and MAPKs and up-regulating mTOR signaling. *International Journal of Molecular Sciences, 16*(7), 16622-16641. https://doi.org/10.1016/j.aninu.2016.03.005

Ling B, Alcorn J. LPS-induced inflammation downregulates mammary gland glucose, fatty

acid, and L-carnitine transporter expression at different lactation stages. (2010). *Research in Veterinary Science, 89*(2):200-2. doi: 10.1016/j.rvsc.2010.03.004

Lucas, A., Gibbs, J.A., Lyster, R.L., Baum, J.D. (1978). Creamatocrit: simple clinical technique for estimating fat concentration and energy value of human milk. Br Med J., 1 (6119):1018-1020, doi: 10.1136/bmj.1.6119.1018.

Mahic, M., Che, X., Susser, E., Levin, B., Reichborn-Kjennerud, T., Magnus, P., ... & Lipkin, W. I. (2017). Epidemiological and serological investigation into the role of gestational maternal influenza virus infection and autism spectrum disorders. *Msphere, 2*(3), e00159-17. https://doi.org/10.1128/mSphere.00159-17

Mäkelä, J., Linderborg, K., Niinikoski, H., Yang, B., & Lagström, H. (2013). Breast milk fatty acid composition differs between overweight and normal weight women: the STEPS Study. *European Journal of Nutrition, 52*(2), 727-735. https://doi.org/10.1007/s00394-012-0378-5

Manzano-Nieves, G., Gaillard, M., Gallo, M., & Bath, K. G. (2018). Early life stress impairs contextual threat expression in female, but not male, mice. Behavioral Neuroscience, 132(4), 247. https://doi.org/10.1037/bne0000248

Meyer, U. (2019). Neurodevelopmental Resilience and Susceptibility to Maternal Immune Activation. *Trends in Neurosciences*, *42*(11), 793–806. https://doi.org/10.1016/j.tins.2019.08.001

Miles, J., Shevlin, M., 2001. Applying regression and correlation: A guide for students and researchers. Sage.

Millett, C. E., Phillips, B. E., & Saunders, E. F. (2019). The sex-specific effects of LPS on depressive-like behavior and oxidative stress in the hippocampus of the mouse. *Neuroscience, 399*, 77-88. https://doi.org/10.1016/j.neuroscience.2018.12.008

Mueller, F. S., Richetto, J., Hayes, L. N., Zambon, A., Pollak, D. D., Sawa, A., ... & Weber-Stadlbauer, U. (2019). Influence of poly (I: C) variability on thermoregulation, immune responses and pregnancy outcomes in mouse models of maternal immune activation. *Brain, Behavior, and Immunity, 80*, 406-418. doi: 10.1016/j.bbi.2019.04.019

Nascimento, A. F., Alves, G. J., Massoco, C. O., Teodorov, E., Felicio, L. F., & Bernardi, M. M. (2015). Lipopolysaccharide-induced sickness behavior in lactating rats decreases ultrasonic vocalizations and exacerbates immune system activity in male offspring. *Neuroimmunomodulation, 22*(4), 213-221. https://doi.org/10.1159/000363350

Naskar, S., Narducci, R., Balzani, E. et al. (2019). The development of synaptic transmission is time-locked to early social behaviors in rats. *Nature Communications, 10*, 1195 https://doi.org/10.1038/s41467-019-09156-3

Núñez Estevez, K.J., Rondón-Ortiz, A.N., Nguyen, J.Q.T., & Kentner, A.C. (2020).

Environmental influences on placental programming and offspring outcomes following maternal immune activation. *Brain, Behavior, and Immunity, 83*, 44-55. doi: 10.1016/j.bbi.2019.08.192.

Pannaraj, P. S., Li, F., Cerini, C., Bender, J. M., Yang, S., Rollie, A., ... & Aldrovandi, G. M. (2017). Association between breast milk bacterial communities and establishment and development of the infant gut microbiome. JAMA Pediatrics, 171(7), 647-654.

Par Pharmaceutical, Inc, Pitocin [Label], Par Pharmaceutical, Chestnut Ridge, NY, (2020), https://dailymed.nlm.nih.gov/dailymed/fda/fdaDrugXsl.cfm?setid=6d4b2c25-2e5d-49b5-93bc-2ae8a20916d1&type=display

Park, B. & Lee, J.O. (2013). Recognition of lipopolysaccharide pattern by TLR4 complexes. Experimental Molecular Medicine 45, https://doi.org/10.1038/emm.2013.97

Patacchioli, F. R., Cigliana, G., Cilumbriello, A., Perrone, G., Capri, O., Alemà, S., ... & Angelucci, L. (1992). Maternal plasma and milk free cortisol during the first 3 days of breast-feeding following spontaneous delivery or elective cesarean section. *Gynecologic and Obstetric Investigation*, *34*(3), 159-163. https://doi.org/10.1159/000292751

Paul, H.A., Hallam, M.C., Reimer, R.A. (2015). Milk collection in the rat using capillary tubes and estimation of milk fat content by creamatocrit. *Journal of Visualized Experiments, 106*, 1-6, doi: 10.3791/53476.

Ordoñes Sanchez, E., Bavley, C.C., Deutschmann, A.U., Carpenter, R., Peterson, D.R., Karbalaei, R….Bangasser, D.A. (2021). Early life adversity promotes resilience to opioid addiction-related phenotypes in male rats and sex-specific transcriptional changes. *PNAS USA, 118*, e2020173118. doi: 10.1073/pnas.2020173118.

Scarborough, J., Mueller, F., Arban, R., Dorner-Ciossek, C., Weber-Stadlbauer, U., Rosenbrock, H., Meyer, U., Richetto, J., 2020. Preclinical validation of the micropipette-guided drug administration (MDA) method in the maternal immune activation model of neurodevelopmental disorders. *Brain Behavior and Immunity 88*, 461–470. doi: https://doi.org/10.1016/j.bbi.2020.04.015

Schatz, K. C., Kyne, R. F., Parmeter, S. L., & Paul, M. J. (2018). Investigation of social, affective, and locomotor behavior of adolescent Brattleboro rats reveals a link between vasopressin's actions on arousal and social behavior. *Hormones and Behavior, 106*, 1-9. https://doi.org/10.1016/j.yhbeh.2018.08.015

Schellekens, H., Torres-Fuentes, C., van de Wouw, M., Long-Smith, C. M., Mitchell, A., Strain, C., ... & Cryan, J. F. (2021). Bifidobacterium longum counters the effects of obesity: Partial successful translation from rodent to human. *EBioMedicine, 63,* 103176.

Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., and Huttenhower, C. (2011) Metagenomic biomarker discovery and explanation. *Genome Biology, 12*: R60. doi: 10.1186/gb-2011-12-6-r60.

Sheard, N. F., & Walker, W. A. (1988). The role of breast milk in the development of the gastrointestinal tract. *Nutrition Reviews (USA)*. doi: 10.1111/j.1753-4887.1988.tb05343.x.

Singh, D., & Fulekar, M. H. (2010). Biodegradation of petroleum hydrocarbons by Pseudomonas putida strain MHF 7109. *CLEAN–Soil, Air, Water, 38*(8), 781-786. https://doi.org/10.1002/clen.200900239

Soriano, O., Regalado, M., Torrero, C., & Salas, M. (2006). Contributions of undernutrition and handling to huddling development of rats. *Physiology & Behavior, 89*(4), 543-551. https://doi.org/10.1016/j.physbeh.2006.07.021

Strzelewicz, A.R., Vecciarelli, H.A., Rondón-Ortiz, Raneri, A., Hill., M.J., Kentner, A.C. (2021). Interactive effects of compounding multidimensional stressors on maternal and male and female rat offspring outcomes. *Hormones and Behavior, 134*, 105013, doi: 10.1016/j.yhbeh.2021.105013.

Strzelewicz, A. R., Ordoñes Sanchez, E., Rondón-Ortiz, A. N., Raneri, A., Famularo, S. T., Bangasser, D. A., & Kentner, A. C. (2019). Access to a high resource environment protects against accelerated maturation following early life stress: A translational animal model of high, medium and low security settings. *Hormones and Behavior, 111*, 46–59. https://doi.org/10.1016/j.yhbeh.2019.01.003

Sullivan, E. C., Hinde, K., Mendoza, S. P., & Capitanio, J. P. (2011). Cortisol concentrations in the milk of rhesus monkey mothers are associated with confident temperament in sons, but not daughters. *Developmental Psychobiology, 53*(1), 96–104. https://doi.org/10.1002/dev.20483

Vacca, M., Celano, G., Calabrese, F. M., Portincasa, P., Gobbetti, M., & De Angelis, M. (2020). The controversial role of human gut lachnospiraceae. *Microorganisms, 8*(4), 573. doi: 10.3390/microorganisms8040573

van Hasselt, F. N., Cornelisse, S., Yuan Zhang, T., Meaney, M. J., Velzing, E. H., Krugers, H. J., & Joëls, M. (2012). Adult hippocampal glucocorticoid receptor expression and dentate synaptic plasticity correlate with maternal care received by individuals early in life. *Hippocampus, 22*(2), 255-266. https://doi.org/10.1002/hipo.20892

Vilela, F.C., Antunes-Rodrigues, J., Elias, L.L., & Giusti-Paiva, A. (2013). Corticosterone synthesis inhibitor metyrapone preserves changes in maternal behavior and neuroendocrine responses during immunological challenge in lactating rats. *Neuroendocrinology. 97*(4):322-30. doi: 10.1159/000346354

Waters, J. L., & Ley, R. E. (2019). The human gut bacteria Christensenellaceae are widespread, heritable, and associated with health. *BMC Biology, 17*(1), 1-11. https://doi.org/10.1186/s12915-019-0699-4

Williams, A.V., Duque-Wilckens, N., Ramos-Maciel, S., Campi, K.L., Bhela, S.K., Xu, C.K., Jackson, K., Chini, B., Pesavento, P.A., Trainor, B.C., 2020. Social approach and social vigilance are differentially regulated by oxytocin receptors in the nucleus accumbens. *Neuropsychopharmacology, 45*, 1423–1430. https://doi.org/10.1038/

s41386-020-0657-4.

Wu, W.L., Hsiao, E.Y., Yan, Z., Mazmanian, S.K., & Patterson, P.H. (2017). The placental interleukin-6 signaling controls fetal brain development and behavior. *Brain, Behavior, and Immunity. 62,* 11-23. doi: 10.1016/j.bbi.2016.11.007.

Wurfel, M.M. & Wright, S.D. (1997). Lipopolysaccharide-binding protein and soluble CD14 transfer lipopolysaccharide to phospholipid bilayers: preferential interaction with particular classes of lipid. *Journal of Immunology. 15;158*(8):3925-34. PMID: 9103463.

Zhao, X., Mohammed, R., Tran, H., Erickson, M., & Kentner, A. C. (2021a). Poly (I: C)-induced maternal immune activation modifies ventral hippocampal regulation of stress reactivity: prevention by environmental enrichment. *Brain, Behavior, and Immunity.* https://doi.org/10.1016/j.bbi.2021.03.018

Zhao, X., Tran, H., DeRosa, H., Roderick, R.C., & Kentner, A.C. (2021b) Hidden talents: Poly (I:C)-induced maternal immune activation improves mouse visual discrimination performance and reversal learning in a sex-dependent manner. *Genes, Brain, and Behavior, 20*(7). doi: 10.1111/gbb.12755.

Zietek, M., Sochaczewska, D., Swiatkowska-Freund, M., Celewicz, Z., & Szczuko, M. (2021). The possible role of corticosterone in regulating sodium and potassium concentrations in human milk. *Ginekologia Polska.* doi: 10.5603/GP.a2021.0072.

Figure 1

Figure 2

Figure 3

Figure 4