1

Regulation of PDF receptor signaling controlling

daily locomotor rhythms in *Drosophila*.

Weihua LI*, Jennifer S. Trigg* & Paul H. Taghert**

Department of Neuroscience

Washington University School of Medicine

St Louis, MO 63110

USA

* co-first authors

** for correspondence: taghertp@wustl.edu

Number of Figures: 7

Number of Tables: 1

Number of Supplemental Figures: 15

Number of Supplemental Tables: 5

Key Words:

Drosophila, Neuropeptide PDF, PDF Receptor, Circadian Rhythms, G protein-coupled Receptor,

Desensitization

Short title: PDF Receptor Desensitization

ABSTRACT

Each day and in conjunction with ambient daylight conditions, neuropeptide PDF regulates the phase and amplitude of locomotor activity rhythms in *Drosophila* through its receptor, PDFR, a Family B G protein-coupled receptor (GPCR). We studied the *in vivo* process by which PDFR signaling turns off, by converting as many as half of the 28 potential sites of phosphorylation in its C terminal tail to a non-phosphorylatable residue (alanine). We report that many such sites are conserved evolutionarily, and their conversion creates a specific behavioral syndrome opposite to loss-of-function phenotypes previously described for *pdfr*. That syndrome includes increases in the amplitudes of both Morning and Evening behavioral peaks, as well as multi-hour delays of the Evening phase. The precise behavioral effects were dependent on day-length, and most effects mapped to conversion of only a few, specific serine residues near the very end of the protein and specific to its A isoform. Behavioral phase delays under entraining conditions were often, but not always correlated with increases in circadian period. The behavioral phenotypes produced by the most severe PDFR variant were ligand-dependent *in vivo*, and not a consequence of changes to their pharmacological properties, nor of changes in their surface expression, as measured *in vitro*. The mechanisms underlying termination of PDFR signaling are complex, subject to regulation that is modified by season, and central to a better understanding of the peptidergic modulation of behavior.

AUTHOR SUMMARY

In multi-cellular organisms, circadian pacemakers create output as a series of phase markers across the 24 hour day to allow other cells to pattern diverse aspects of daily rhythmic physiology and behavior. Within circadian pacemaker circuits, neuropeptide signaling is essential to help promote coherent

circadian outputs. In the fruit fly *Drosophila* 150 neurons are dedicated circadian clocks and they all tell

the same time. In spite of such strong synchronization, they provide diverse phasic outputs in the form

of their discrete, asynchronous neuronal activity patterns. Neuropeptide signaling breaks the clock-

generated symmetry and drives many pacemakers away from their preferred activity period in the

morning. Each day, neuropeptide PDF is released by Morning pacemakers and delays the phase of

activity of specific other pacemakers to later parts of the day or night.   When and how the PDF that is

released in the morning stops acting is unknown. Furthermore, timing of signal termination is not fixed

because day length changes each day, hence the modulatory delay exerted by PDF must itself be

regulated. Here we test a canonical model of G protein-coupled receptor physiology to ask how PDF

receptor signaling is normally de-activated. We use behavioral measures to define sequence elements

of the receptor whose post-translational modifications (e.g., phosphorylation) may define the duration

of receptor signaling.

63   **INTRODUCTION**

64

65   In *Drosophila*, neuropeptide PDF signaling helps pattern the output of the fly circadian

66   pacemaker network that controls rhythmic daily locomotor activity [1-3]. Its functions have been

67   compared to those of neuropeptide VIP in regulating mammalian circadian physiology [4]. Historically

68   PDF was first isolated as an active principle (Pigment Dispersing Hormone) that mediates light-

69   dependent dispersion of pigment granules in diverse chromatophores of crustacea [5, 6]. In the insect

70   circadian system, PDF acts for a specified period within each 24 hr cycle, and works in conjunction with

71   environmental light to set phase and amplitude for locomotor activity rhythms that normally occur

72   around dawn and dusk [7-9]. Each day, the precise times of dusk and of dawn change, which alters the

73   time interval between them. These facts require that the time of PDF signaling, the point when it starts

74   and the point when it stops, must also be adjusted each day to appropriately follow and reflect these

75   daily variations in the light:dark transitions. PDF signaling starts following its release by specific

76   pacemaker neurons, whose period of activity *in vivo* tracks the dawn in a variety of photoperiodic

77   conditions [10]. We lack a comparable understanding of how PDF receptor signaling normally stops: this

78   work addresses that mechanism.

79

80   The PDF receptor (PDFR) is a member of the Family B (secretin receptor-like) GPCR group [11-

81   13]: it is G_s-coupled and its activation elevates cAMP levels *in vivo* [14]. It regulates different adenylate

82   cyclases (AC) in diverse target pacemakers (15, 16), which, through PKA activation, ultimately regulate

83   the pace of the molecular clock through regulation of Timeless [17]. PDFR autoreceptor signaling

84   promotes dramatic, daily morphological changes in the axonal terminals of sLNv pacemakers [18]. In

85   addition to its effects on the pace of the molecular pacemaker, PDFR activation also regulates calcium

86   dynamics in subsets of pacemaker neuron groups to help dictate their group-specific, daily phases of

activation (in the sLNv, in the 5th sLNv and in subsets of LNd, and DN3 groups – [10]). Such target cell-

specific delays of PER-dependent neuronal activity illustrate the basis by which the circadian network

produces a daily series of staggered phasic, neuronal outputs [19, 20]. Finally, PDF/PDFR signaling is

long-lasting: its depression of basal calcium levels in target neurons persists without abatement over

many hours [19]. These observations raise fundamental questions regarding the mechanism and the

time course by which PDFR signaling diminishes in anticipation of the next day's cycle of signaling.

93
The canonical model of GPCR phosphorylation and homologous desensitization features G

protein-coupled receptor kinases (GRKs) which associate with activated GPCRs and phosphorylate

cytosolic segments, thereby recruiting β-arrestins [21, 22]. β-arrestins uncouple the receptors from G

proteins [23, 24]. or enhance receptor endocytosis [25]; they can also serve as signal transducers by

recruiting distinct signaling molecules [26]. A second major regulatory mechanism to reduce GPCR

signaling is heterologous desensitization, whereby second-messenger-dependent kinases (PKA or PKC)

phosphorylate GPCRs [27].   Thus, we designed experiments to modify evolutionarily-conserved residues

in the C terminal tail of PDFR that could conceivably serve as substrates for phosphorylation and

subsequent signal termination. Our working hypothesis was that, by their actions *in vivo*, such modified

PDF receptors would reveal extended lifetimes of activation.

We know very little about the mechanisms that underlie normal termination of PDFR signaling.

Levels of *pdfr* cycle in many pacemaker groups in the *Drosophila* brain, but they peak at different times,

in different groups [28]. Sensitivity to PDF *in vivo* peaks in the early day and is regulated by the PER-

dependent clock, via post-transcriptional mechanisms: the EC50 for PDF responses in identified neurons

varies systematically 5-10 fold: as a consequence of RalA action, as a function of time of day, and as a

function of seasonality [29]. PDFR signaling is long-lasting: it persists for many hours *in vivo* [19], for as

long as free peptide ligand is available in the bath [14]. In addition, β-arrestin2-GFP is not efficiently

recruited to activated PDFR when the receptor is functionally-expressed in *hEK-293T* cells [29]. In

contrast, each of 13 other *Drosophila* neuropeptide GPCRs (including two Family B GPCRs, CG8422

and CG17415) efficiently recruit β-arrestin2-GFP, when they are activated by their cognate ligands in that

cellular environment [30-32].

Here we report (i) that PDFR is normally phosphorylated *in vivo* at conserved C terminal

residues; (ii) that loss of conserved PDFR phosphorylatable sites leads to a behavioral syndrome

opposite to loss-of-function *pdf* and *pdfr* phenotypes, with effects on both the amplitude and phase of

the daily locomotor peaks. This 'gain of function' approach reveals a multi-hour range of potential

phases for both the Morning and Evening activity peaks, within which neuropeptide PDF:PDFR signaling

normally specifies rhythmic behavior, according to season. In addition, using a structure-function

approach, this work identifies specific PDFR sequence elements that are major points at which the

duration of receptor activity is regulated, and through which behavior is modulated in season-specific

fashion.

RESULTS

PDFR C-Terminal sequences. Based on alternative splicing, the *pdfr* locus in *Drosophila*

melanogaster (CG13758) encodes very similar GPCRs which differ in their extreme C terminal sequences,

for which the PA and PD isoforms are representative (flybase.org/reports/FBgn0260753).   The PD

isoform is slightly longer and lacks the final ~20 AAs of the PA isoform. As described in the Supplemental

Information (*PDFR Isoforms*), we focused on the PA protein isoform of PDFR, as it has been used in the

majority of genetic studies in the field. To identify residues for mutation, we first used comparative

genomic analyses to assess how well specific sequences in the C-terminal region of the PA receptor are

conserved.   We obtained annotated *pdfr* genomic sequences from 16 additional species of *Drosophila*,

in both the *Sophophora* and *Drosophila* sub-families (**Suppl. Table 1**). Together this species collection

represents an estimated 40-60 MYr of *Drosophalid* evolution [33]. We defined residue V505 of the

melanogaster protein as the start of the C terminal sequence, following the consensus 7^th

transmembrane TM domain (TM7) (**Suppl. Figure 1**). These 17 different PDFRs all contain C-termini of

considerable length, and vary between 189 to 215 amino acids (AAs). The *melanogaster* PDFR-A C-

terminal contains 28 Ser, Thr, or Tyr residues (these may be subject to post-translational

phosphorylation and de-phosphorylation, and/ or other modifications). To survey putative functionality

among these, we chose 14 residues that are distributed across the length of the C-terminal and which

display high evolutionary conservation (**Suppl. Figure 1** and **Suppl. Table 3)**. For naming purposes, we

grouped them into arbitrary clusters (CL) numbered #1 to #7, with positions shown in **Figure 1.**

Following a common paradigm in study of GPCR physiology [e.g., 34-36], we performed an "Alanine

Scan": testing the consequences of their mutation to Alanine, which is a non-phosphorylatable analog

[41]. Some clusters have only a single modified AA (e.g. CL4 and CL5), while in others we concurrently

modified two AAs (e.g., CL1, CL6 and CL7), and in others, as many as six closely-positioned residues (e.g.,

CL2-3).

Study Overview. This series of variant receptors contains 10 different mutated versions of PDFR

(not including the WT 'parent' PDFR-A) and is arbitrarily divided into two broad categories as indicated

in **Figure 1**. The first group (termed "Simple Variants") targeted one or two clusters of conserved

Ser/Thr/Tyr AAs (i.e., CL 2-3A; 4A; 5A; 6A; 7A). The second category (termed "Multiple Variants")

targeted three or more of the AA clusters in various combinations (i.e., CL 1-4A; 1-5A; 1-6A; 1-7A and 5-

7A). The CL1-7A variant is the most severe as it mutates all 14 of the targeted AA residues. We studied

these ten PDFR variants, as well as the WT receptor, following expression in a *pdfr* mutant strain (*han^5537*

[11]; we measured the ability of individual PDFR variants to rescue and to shape the phases and

periodicity of rhythmic locomotor behavior. For the most part, we analyzed PDFR variants in the *pdfr*

mutant background to permit evaluation of their properties, without competition from endogenous

PDFR. Normally, PDFR-A is expressed throughout the ~150 cell pacemaker network, typically in subsets

of each clustered group, as revealed by its expression from an epitope-marked BAC transgene [37].

However, no simple *Pdfr*-Gal4 driver element recapitulates a majority of normal expression sites [37].

Therefore to effect broad PDFR-A expression in the pacemaker system, we used a *timeless*-Gal4 driver

element. Because PDF sensitivity varies with seasons [37], we tested this PDFR series in different

photoperiodic conditions, as well as in constant darkness. *pdfr* loss-of-function mutants display

advanced behavioral peaks under Light:Dark conditions, as well as weak and shortened free-running

periods under constant darkness [11].

The behavioral actions of the Ala-mutated PDFRs either resembled those of WT receptor, or

exhibited gain-of-function properties (behavioral actions opposite to those seen in loss-of-function *pdfr*

mutant flies). For purpose of clarity, the Main Figures report a comparison of properties for the WT

PDFR with those of the most extensively-mutated version of PDFR (called 1-7A), as indicated in **Fig. 1**.

Comparable data on the properties of the other variant receptors in the series is described in

Supplemental Figures. To better interpret the results produced by the different sequence variants, we

also present experiments that test assumptions used in the experimental design. These tested the

following hypotheses: i) that the behavioral effects of receptor variants are independent of activation by

the endogenous ligand, neuropeptide PDF; and ii) that the bulky C-terminal fusion of GFP (present in all

variants tested) strongly influenced the results. Both hypotheses were dis-proved.

Effects of WT vs 1-7A PDFR on locomotor behavior in Short Day (winter-like) conditions (Figure

2). In 8L:16D, *pdfr^han* mutant flies lack a prominent morning peak and their evening peak of activity

begins 1-2 hr earlier than controls; the example shown in **Fig. 2A-1** (*tim>no transgene,* boxed in yellow)

is heterozygous for the *tim*-Gal4 element.   Rescue by a WT-*pdfr* cDNA (**Fig. 2B-1** (*tim>pdfr*, boxed in

black)) did not produce obvious effects on activity in the time domain preceding or just past Lights-On

(Morning). However, it strongly affected the Evening peak: not its amplitude (**Fig. 2B-3 and B-4**) but

significantly delaying its phase by about 1 h (**Fig. 2E**) such that the Evening peak now extends past after

Lights-off (**Fig. 2B-4**). In contrast, expression of the 1-7A Multiple Variant driven by *tim*-Gal4 (*tim>*

pdfr1-7A, boxed in red) elevated the amplitude of activity during the period of ZT 18-21 prior to lights-

ON (**Fig. 2C-1 and C-2, orange arrow**): we speculate this is promotion of a "Morning" peak of activity. It

has the same phase as that produced by expression of the WT receptor (**Fig. 2D**), but significantly larger

amplitude (**Fig. 2C-2, orange arrow**). The 1-7A variant also delayed the Evening peak but to a much

greater extent than did the WT PDFR (**Fig. 2C-1**, **C-4**, blue arrow), producing a conspicuous, large

amplitude peak that occurred on average as late as ~3 h after lights-off, at ZT12, significantly more

delayed than the peak produced by WT PDFR (**Fig. 2E**). Following short day light entrainment, flies were

released into constant darkness for ~7-8 days (DD - constant conditions): resulting locomotor activity is

displayed in the double-plotted group actograms (**Fig. 2F -** yellow box**, G -** black box**,** and **H -** red box).

The properties of the persistent circadian rhythmic behavior are provided in **Table 1**. The period (tau) of

the 1-7A variant was considerably lengthened. The phase of the major DD rhythm matched that of the

much-delayed evening peak in the 1-7A flies (**Fig. 2H**, green line).   In sum, under winter-like

photoperiods, the PDFR 1-7A variant significantly increased the amplitude of the morning activity peak

and significantly delayed the phase of the evening peak; the latter phase was also reflected in persistent

rhythmic activity under subsequent DD.

Behavioral effects of the other PDFR Variants under Short Day (winter-like) conditions (Suppl.

Fig. 2 & 3). We tested each of the nine other PDFR Ala-variants by comparing their effects on locomotor

behavior as we had done for the PDFR 1-7A variant. **Suppl. Figure 2** presents these collected data with

Group eductions, Actograms and Phase analysis; **Suppl. Fig. 3** presents these same collected data with

bin-by-bin comparisons to WT-PDFR to consider amplitude changes. The phase of the Morning peak

was not obviously or systematically effected (**Suppl. Fig. 2L**), while that of the Evening peak was

significantly delayed by several (**Suppl. Fig. 2M**). We note two features – first the progressive addition

of more Ala substitutions in the series 1-4A, 1-5A, 1-6A (**Suppl. Fig. 2I-K**) tended to produce large and

delayed Evening peaks in the time period ZT11.5-12.5 (3-4 h after lights-off), with the variant 1-6A

producing the most pronounced delay. The delayed Evening peak activity at ZT12 sometimes appeared

at the expense of, the normal Evening peak activity that occurred prior to Lights-off (e.g., **Suppl**. **Fig.2E-1**

and F-1), although when averaged across all days in LD, that effect for specific variants was not

significant (**Supp**l. **Fig. 3E-2 and F-2**). Second, a delayed Evening phase was also produced by two of the

Simple PDFR variants (**Suppl. Fig. 2M**), the 6A and 7A variants, each of which contain only a pair of Ser-

to-Ala substitutions. The amplitude of the Morning peak (ZT18-20) was increased modestly by variants

2-3A,_7A and 1-5A (**Suppl. Fig 3C-1, G-1** and **J-1, red arrows**) or strongly by 6A (**Suppl. Fig. 3F-1, red**

arrow). The amplitude of the Evening peak was also increased by others in the PDFR variant series

(**Suppl. Fig. 3, blue arrows**), of which the 6A, 7A, 1-5A and 1-6A variants had effect sizes most similar to

that of 1-7A (**Supp. Fig. 3F-3. G-3. J-3 and K-3).**_ We also note_that_combining the Single 5A, 6A and 7A

variants into a Multiple Variant (5-7A) did not produce the anticipated additive effects on increasing

Morning peak amplitude at ~ZT19 (**Suppl. Fig. 3H-1),** nor on increasing Evening peak amplitude **(Suppl.**

Fig. 3H-3). Such anticipation was predicated on the effects seen individually with the 1-5A, 1-6A and 1-

7A variants (versus the 1-4A); instead the 5-7A produced only a modest increase in the size of the

delayed Evening peak (**Suppl**. **Fig. 3H-3**).

Effects of WT versus 1-7A PDFR on locomotor behavior under equinox conditions (Fig. 3). Under

12:12 conditions, *han* mutant flies (lacking *pdfr* function) typically display elevated nocturnal activity, a

lack of morning anticipation prior to Lights-on, and a pronounced advance in the peak of the Evening

behavior [11]. Although we note that some reports have observed remnants or a full bout of Morning

activity in *han* mutant flies [e.g., 37]. **Fig. 3A-1** displays activity patterns of *han* mutants that are

heterozygous for *tim*-Gal4; they generally matched prior descriptions of *pdfr* mutant behavior.

Restoration of *pdfr* function by *han*; *tim>* WT-*pdfr*, restored a Morning peak (anticipatory activity prior

to lights-on – **Fig. 3B-1 and -2**) and delayed the evening peak by 1-2 hrs (**Fig. 3B-1 and B-4**) [7,8,9].

Restoration of *pdfr* function by the 1-7A PDFR variant under these conditions (**Fig. 3C-1**) increased the

amplitude of the Morning activity peak but not its phase (**Fig. 3C-2, red arrow**) but not its phase (**Fig.**

3D). 1-7A expression in equinox conditions rescued the advanced phase of the Evening activity peak,

but did not further delay it past Lights-OFF (**Fig. 3C-4** and **3E**), unlike what we observed under short-day

conditions. The amplitude of the Evening peak was only modestly elevated following expression of WT

or 1-7A PDFRs. The period (tau) of the 1-7A variant was considerably lengthened, but see discussion

below about the influence of the GFP fusion. The phase of the major DD rhythm was delayed relative to

the evening peak under entraining conditions (**Fig. 3H**, green line).   In sum under equinox-like

photoperiods, the PDFR 1-7A variant significantly increased the amplitude of the morning activity peak

compared to WT PDFR, but did not differentially affect its phase, or the amplitude or phase of the

Evening peak; under subsequent DD, the rhythmic activity displayed a delayed phase starting within the

very first cycle.

Behavioral effects of the other PDFR Variants under equinox conditions (Suppl. Fig. 4 & 5). We

tested each of the nine other PDFR Ala-variants by comparing their effects on locomotor behavior as we

had done for the PDFR 1-7A variant. The phases of the Morning and Evening peaks were not obviously

or systematically effected (**Suppl**. **Fig. 4L** and **4M**). Relative to WT PDFR expression, the Morning Activity

peak amplitude was increased by expression of the 6A, 7A and 1-5A variants (**Suppl. Fig. 5F-1, G-1**, and **J-**

1, red arrows). Relative to WT PDFR expression, the Evening Activity peak amplitude was increased by

expression of the 4A, 6A, 7A, 5-7A and 1-5A variants (**Suppl. Fig. 5D-3, F-3, G-3**, **H-3** and **J-3**, blue

arrows). We also note that combining the Single 5A, 6A and 7A variants into a Multiple Variant (5-7A)

did not produce the anticipated additive effects on increasing Morning peak amplitude at ~ZT21 (**Suppl.**

Fig. 5H-1, versus 5F-1 and G-1).

Behavior under Long Day (summer) conditions (Fig. 4). Flies lacking *pdfr* function in 16L:8D

conditions display locomotor patterns similar to those in 12L:12D – lack of a clear Morning peak and

display of a broad Evening activity period that peaks ~2-3hr before Lights-off: the example in **Figure 4A-1**

(yellow box) displays behavior by *han* mutants that are also heterozygous for the *tim*-Gal4 element.

Rescue (*han*; *tim*> WT-*pdfr*, **Figure 4B-1, black box**) typically delayed Morning activity offset (**Fig. 4B-3**

and 4D) and delayed the Evening peak onset each by 1.5 h (**Fig. 4B-4 and 4E).**   Relative to that of WT

PDFR, expression of the 1-7A variant (**Figure 4C-1**) increased the amplitude of both the morning activity

peak (**Figure 4C-3**) and the evening activity peak (**Figure 4C-4**). It did not significantly alter the Morning

phase relative to effects of WT PDFR (**Figure 4D),** although there was a pronounced tendency to

broaden the duration of the Morning activity peak.   1-7A expression did significantly extend the delay

the Evening phase of activity relative the effect of Wt PDFR **(Figure 4E)**. In DD, the 1-7A PDFR variant

significantly lengthened circadian period and the dominant rhythmic activity displayed a phase several

hours delayed from the evening activity peak phase under light entraining conditions within the first

cycle (**Fig. 4H** – green bar). In summary the 1-7A PDFR variant significantly increased the amplitudes of

the Morning and Evening activity peaks under long days, and also slightly delayed the Evening activity

phase.

Behavioral effects of the other PDFR Variants under long day (summer-like) conditions (Suppl. Fig.

6 & 7). We tested each of the nine other PDFR Ala-variants by comparing their effects on locomotor

behavior as we had done for the PDFR 1-7A variant. There was no systematic effect of the variants on

the phase of the Morning peak (**Suppl. Fig. 6L**), but several did delay the Evening activity phase (**Suppl.**

Fig 6M), especially the 1-4A, 1-5A and 1-6A variants. A few variants affected the Morning peak

amplitude – 5A, 6A and 1-6A (**Suppl. Fig. 7E-2, F-2 and K-2**). Several increased the amplitude of the

Evening peak under these conditions, namely 2-3A, 4A, 5A, 6A, 7A, 1-5A and 1-6A (**Suppl. Fig. 7 C-3, D-3,**

E-3, F-3, G-3, J-3 and **K-3**). As observed in the other photoperiodic conditions, when we combining the

5A, 6A and 7A variants into a single 5-7A variant, it did not produce the expected additive effects on

behavior in the morning (**Suppl. Fig 7H-2**) or in the evening (**Suppl. Figure 7H-3**): instead these activities

appeared similar to effects displayed by the 1-4A variant.

Effects of PDFR variants on period and phase of rhythmic activity in DD. In DD, following Short Day

conditions, the PDFR variants that produced 3-4 h delays in the evening peak often generated periods ~

1-2 hr longer than the controls and also lowered % arrhythmicity (**Table 1**). This was especially true for

the Multiple Variant series (e.g., PDFR 1-5A, 1-6A and 1-7A (**Table 1**). However the correlation between

delayed evening peaks and a longer tau in DD was not absolute. For example, the 6A, 7A and 5-7A

variants all had strong and significant delaying effects on the Evening phase of activity under Short Day

conditions (compared to the effect of the WT PDFR), but they did not lengthen tau values in subsequent

DD conditions (**Table 1**). These observations suggest that delays of activity phases produced by PDFR

modulation in LD may not reflect a direct consequence of PDFR effects on the PER and TIM--dependent

clock [2, 3, 17]. In addition, the dominant activity periods in DD typically reflected the delayed (~ZT12)

Evening peak (**Suppl**. **Fig. 2F-2, 2G-2** and **2H-2**). Notably, the early Morning peak (~ZT18-19) promoted

by PDFR 6A and 7A variants clearly persisted in DD (**Suppl**. **Fig. 2F-2, 2G-2**). In DD following Equinox or

Long Day entrainment (with photophases > 8 hr in duration), we noted that several variants produced

dominant activity phases which were phase-delayed relative to the phase of the Evening peak displayed

in LD (e.g., 6A and 7A - **Suppl**. **Fig. 4F-2 and 4G-2**; **Suppl**. **Fig. 6F-2 and 6G-2).** Again, several of the

variants significantly increased Tau in DD (**Table 1**), yet had no delaying effects on phase in LD: examples

included 2-3A, 1-4A, and 1-5A, under equinox conditions (**Suppl. Fig. 4C-2, 4I-2, 4J-2**) and 4A and 1-5A in

long day conditions (**Suppl. Fig. 6D-2, 6J-2).**

Table 1.

Short Day Entrainment	N	n	%AR	tau	SEM	Average tau different from:		
						w[1118]*	pdfrEgfp**	pdf[01]
han[5304]; tim > w[1118]	5	72	22%	23.9	0.09	-	ns	nd
han[5304]; tim > pdfr	1	16	6%	23.7	0.13	ns	ns	nd
han[5304]; tim > pdfr1-7A No GFP	3	30	8%	23.6	0.07	ns	ns	nd
han[5304]; tim > pdfrEgfp	3	46	35%	24.0	0.09	ns	-	nd
han[5304]; tim > pdfrEgfp23A	2	31	13%	24.8	0.12	ns	ns	nd
han[5304]; tim > pdfrEgfp4A	2	31	26%	24.7	0.13	ns	ns	nd
han[5304]; tim > pdfrEgfp5A	2	43	26%	24.6	0.11	ns	ns	nd
han[5304]; tim > pdfrEgfp6A	3	50	38%	24.4	0.08	ns	ns	nd
han[5304]; tim > pdfrEgfp7A	3	45	27%	24.8	0.11	ns	ns	nd
han[5304]; tim > pdfrEgfp1-4A	3	40	12%	24.6	0.12	ns	ns	nd
han[5304]; tim > pdfrEgfp1-5A	3	31	16%	25.0	0.23	***	*	nd
han[5304]; tim > pdfrEgfp1-6A	3	42	14%	25.1	0.13	****	**	nd
han[5304]; tim > pdfrEgfp1-7A	4	59	16%	24.9	0.11	****	*	nd
han[5304]; tim > pdfrEgfp567A	2	37	16%	23.9	0.09	ns	ns	nd
tim > pdfrEgfp1-7A	1	15	0%	25.8	0.25	nd	nd	****
pdf[01]	1	12	67%	22.5	0.12	nd	nd	-
tim > pdfrEgfp1-7A; pdf[01]/pdf[01]	1	25	20%	23.6	0.12	nd	nd	ns

Equinox Entrainment	N	n	%AR	tau	SEM		
han[5304]; tim > w[1118]	6	68	34%	23.7	0.09	-	ns
han[5304]; tim > pdfr	3	16	25%	23.5	0.16	ns	ns
han[5304];; tim > pdfr1-7A No GFP	3	35	14%	23.7	0.08	ns	ns
han[5304]; tim > pdfrEgfp	4	48	13%	24.1	0.06	ns	-
han[5304]; tim > pdfrEgfp23A	2	54	48%	25.6	0.12	****	****
han[5304]; tim > pdfrEgfp4A	2	44	32%	24.1	0.07	ns	ns
han[5304]; tim > pdfrEgfp5A	2	47	34%	24.4	0.06	ns	ns
han[5304]; tim > pdfrEgfp6A	3	35	23%	24.3	0.13	ns	ns
han[5304]; tim > pdfrEgfp7A	3	47	21%	24.1	0.09	ns	ns
han[5304]; tim > pdfrEgfp1-4A	3	45	18%	24.5	0.08	**	ns
han[5304]; tim > pdfrEgfp1-5A	2	32	6%	25.0	0.13	****	***
han[5304]; tim > pdfrEgfp1-6A	2	32	19%	25.3	0.14	****	****
han[5304]; tim > pdfrEgfp1-7A	6	88	6%	25.1	0.17	****	****
han[5304]; tim > pdfrEgfp567A	2	45	24%	24.1	0.08	ns	ns

Long Day Entrainment	N	n	%AR	tau	SEM			
han[5304]; tim > w[1118]	7	105	47%	23.8	0.06	-	ns	nd
han[5304]; tim > pdfr	3	47	17%	24.9	0.15	ns	ns	nd
han[5304]; tim > pdfr1-7A No GFP	3	40	48%	24.6	0.09	ns	ns	nd
han[5304]; tim > pdfrEgfp	4	82	17%	24.9	0.08	ns	-	nd
han[5304]; tim > pdfrEgfp23A	2	48	23%	25.4	0.12	****	****	nd
han[5304]; tim > pdfrEgfp4A	1	27	22%	25.5	0.16	****	****	nd
han[5304]; tim > pdfrEgfp5A	1	36	11%	24.7	0.16	ns	ns	nd
han[5304]; tim > pdfrEgfp6A	2	51	22%	25.0	0.09	ns	ns	nd
han[5304]; tim > pdfrEgfp7A	2	54	13%	25.5	0.09	****	****	nd
han[5304]; tim > pdfrEgfp1-4A	3	40	8%	25.2	0.13	****	ns	nd
han[5304]; tim > pdfrEgfp1-5A	3	33	6%	25.2	0.14	****	*	nd
han[5304]; tim > pdfrEgfp1-6A	3	48	2%	25.6	0.09	****	****	nd
han[5304]; tim > pdfrEgfp1-7A	4	63	14%	25.3	0.08	****	***	nd
han[5304]; tim > pdfrEgfp567A	1	38	18%	24.3	0.09	ns	ns	nd
tim > pdfrEgfp1-7A	1	14	14%	25.5	0.13	nd	nd	****
pdf[01]	1	32	88%	22.7	0.05	nd	nd	-
tim > pdfrEgfp1-7A; pdf[01]/pdf[01]	1	29	21%	23.3	0.09	nd	nd	ns

Table 1. Locomotor activity measures of various genotypes during constant dark conditions.

Behavioral activity records from days 3-9 in constant conditions following entrainment in the three

indicated photoperiods (Short Day, Equinox and Long Day). N = number of independent experiments

performed for each genotype::photoperiod combination. n = total number of flies tested for each

genotype::photoperiod combination. % AR: the percentage of flies judged arrhythmic by criteria; tau =

average circadian periods calculated according to χ2-periodogram analysis. * complete genotype =

han[5304]; tim > w[1118]. ** complete genotype = *han[5304]; tim > pdfrEgfp*. Statistical analysis

compared tau's using Tukey's multiple comparisons post hoc test following a one-way ANOVA: ns: not

significant; * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001.; nd: not determined.

Control experiments. The design of the *PDF*R variants contains assumptions and genotypic

constraints, distinct from simply substituting Alanine at potential sites of phosphorylation in the

receptor C terminal tail. To assess the potential of some assumptions to affect the results we report, we

performed the following two sets of experiments as controls. ***Control experiments (i) - dependence of***

GPCR PDFR variant effects on the presence of PDF ligand. We assumed that the altered behavioral

phenotypes produced by certain PDFR variants depended on activation by their endogenous cognate

ligand, the neuropeptide PDF. However, we could not *a priori* exclude the possibility that PDFR variants

may in fact produce novel constitutive activity (neomorphic properties). We therefore placed the 1-7A

PDFR variant in the *pdf^{01}* (null) background and re-tested its activity when driven by *tim*-Gal4 under both

short day and long day conditions, and then in constant darkness. In all conditions, the behavioral

phenotypes largely resembled those of the *pdf^{01}* background (lack of a morning activity peak, advanced

evening peak, and shorter tau/s in DD). The results are shown in **Fig. 5** (under short days) and **Fig. 6**

(under long days) and tabulated in **Table 1**. The ability of the 1-7A PDFR variant to increase the

amplitude of the Morning peak was completely dependent on WT *pdf* function (**Fig. 5B-2**). Likewise its

ability to significantly delay the Evening phase into the dark period was completely dependent on WT

pdf function (**Fig. 5B-4 and 5E**). Under Long days, the ability of the 1-7A variant to increase the

amplitudes of the Morning and Evening activity peaks were strongly diminished by lack of WT pdf

function (**Fig. 6B-3** and **B-4**). The 1-7A variant did display some activity in the *pdf* mutant background:

for example, it produced a significant delay of the Evening peak phase (akin to the action of the WT

receptor, **Fig. 5A-3** and **Fig.6 A-4**) and a reduction in the % arrhythmicity to nearly the same value as

found in a WT *pdf* background. These results suggest some degree of constitutive (ligand-independent)

activity. Apart from these, we conclude that constitutive activity explains at best a small proportion of

the behavioral effects that distinguish the 1-7A PDFR variant from the WT receptor.

(i) - ***The influence of epitope fusions.*** The second control experiment asked whether the 1-7A

epitope fusions (4xFLAG and eGFP), might confer some of the differences in behavioral effects that we

observed. In part, the concern regarding fused sequences is mitigated by the fact that all UAS-*PDFR*

isoforms in the experimental series contained both epitopes (in addition – all were genetically

transduced to the same integration site). Hence comparisons across different variants should largely

normalize for the effects of the non-receptor sequences, and instead highlight variant-specific

properties. However, we wished to test this assumption explicitly to define the extent to which a bulky

GFP fused to the C Terminal tail might confer some of the behavioral properties we might otherwise

ascribe to PDFR C terminal sequences. We therefore compared a 1-7A version of PDFR to a wild type

PDFR, both of which lacked GFP fusions. We found that the additional Evening phase delays in short day

and long day produced by 1-7A expression occurred despite the absence of a GFP fusion (**Supp. Fig. 8B-**

4, 8H and **8J-4 and 8P**); in addition, the increase of the Morning amplitude under Long days occurred

despite the absence of a GFP fusion (**Suppl. Fig. 8J-3**). Finally, the phase of the main DD rhythmic

activity, following expression of 1-7A lacking a GFP fusion, reflected the delayed evening phase in short

day (**Suppl. Fig. 8F**, green bar) and was delayed relative to that of the LD Evening activity peak in long

days (**Suppl. Fig. 8N**, green bar). We note however that the longer period produced by the 1-7A variant

was not exhibited by the 1-7A lacking GFP, suggesting that a lengthened period depends at least in part

on the GFP fusion.   In sum, these results support the hypothesis that most of the key alterations in daily

locomotor rhythmicity derived from sequence variation of the PDFR, and not from the properties of the

fused GFP.

PDFR expression and signaling *in vitro*. The PDFR is a G_s-coupled receptor and several reports

have documented the importance of the downstream cAMP pathway to mediate PDF behavioral

regulation [15, 17, 38]. In that context, we asked whether the *in vivo* properties of those PDFR variants

that affected behavioral phase and amplitude could be correlated with *in vitro* properties when

expressed in *hEK-293T* calls. We found no strong correlation. Significant differences in basal signaling

levels, in EC50 or in maximum values of signal transduction were rare (**Suppl. Fig. 9, 10 and 11**). When

noted, they were poorly correlated with behavioral effects (**Suppl. Table 5**). We also measured surface

expression of the PFDR variants in *hEK* cells using β-lactamase N terminal fusions [39] to determine if

PDFR variants tended to display longer surface lifetimes. The 1-4 and 1-7 variants displayed higher basal

levels, but no others were different from the WT levels (**Suppl. Fig. 12**). Following 20 min exposure to

PDF, neither the WT not any of the variants displayed a change in surface expression levels (**Suppl. Fig.**

13).

Measuring the phosphorylation state of over-expressed PDFR *in vivo*. To obtain direct

evidence that PDFR sequences are phosphorylated *in vivo*, we first over-expressed an epitope-tagged-

PDFR WT construct using *tim*-Gal4, which directs expression broadly in cells that feature the PER-

dependent molecular oscillator. We immunoprecipitated the receptor from head extracts, then

employed tandem mass spectroscopy to determine which if any specific PDFR peptide fragments are

phosphorylated. We performed eight biological replicates, with two collections in the morning (ZT2-3)

and six in the evening hours (ZT11). **Suppl.Table 4** reports the phosphorylated peptides detected from

the PDFR-GFP fusion protein.   Among the 28 conserved Ser/Thr and Tyr residues in the PDFR C terminal

tail, five were phosphorylated in one or more of these samples. In the two Mornings samples S563 was

phosphorylated in one sample; in the Evening samples, S531 (in CL2) was phosphorylated in four of

samples, S534 (in CL2) in one, T543 (in CL3) in one, and S560 in two of six samples. The spectra

documenting detection of these phosphopeptides are presented in **Suppl. Fig. 14**.

Testing GRKs and β-arrestin2 contributions to locomotor rhythmicity. Regarding which

potential PDFR phosphorylation sites might be β-arrestin binding sites, we considered a recent report by

Zhou *et al.* [40], who proposed a conserved GPCR sequence motif that promotes high affinity

interactions with β -arrestin. The motif includes three phosphorylation sites that align with three

conserved, positively-charged pockets in the arrestin N-domain. In *D. melanogaster* PDFR, the CL1

region contains a match with the proposed motif beginning with S512 (SLATQLS) and shows moderate

sequence conservation: of the 17 species we considered, 9 others retained this precise motif (**Suppl. Fig.**

1). In addition, the *D. melanogaster* PDFR sequence beginning with S629 (SRTRGS) also displays a match

for the motif, but is retained in only 6 of the other 16 species. The evidence that these two sites

represent high affinity β -arrestin2 binding domains is therefore equivocal. We previously reported that

β-arrestin2-GFP is not efficiently recruited to activated PDFR when functionally-expressed in *hEK-293T*

cells [29]. In contrast, each of 13 other *Drosophila* neuropeptide GPCRs do efficiently recruit β-

arrestin2-GFP when expressed and activated in that cellular environment [30-32].

We tested the potential to detect *in vivo* involvement by *Drosophila* orthologues of the

canonical desensitization effectors - mammalian G-protein coupled receptor kinases (GRKs – GPRK1

(CG40129) and GPRK2 (CG17998)) and of mammalian β-arrestin2 (βarr2 – *kurtz* (CG1487)) - in the

control of locomotor rhythms We drove specific over-expression of WT cDNAs and RNAi constructs

using *tim*-Gal4, to broadly affect signaling in the circadian pacemaker system. The PDFR-PA protein is

highly restricted in its expression to subsets of the pacemaker neural network [9]. We predicted that

elimination of a desensitizing component for PDFR signaling should produce behavioral phenotypes

opposite to that of *pdfr* loss of function phenotypes: these would potentially include (for example) a

delayed evening activity phase in LD conditions and a longer tau under DD. GPRK1- and *krz*-specific

RNAi's produced normal average locomotor profiles under 12:12, while one of two GPRK2-specific RNAi

constructs tested slightly advanced the evening peak (**Suppl. Fig. 15**). Over-expressing GPRK1 cDNAs did

not affect morning or evening phase; over-expressing GPRK2 broadened the evening peak. Under DD,

krz RNAi flies were uniformly arrhythmic, while GPRK RNAi's were normal or slightly lengthened the

circadian period (**Suppl. Table 5**).   These results do not support the hypothesis, and suggest that the

kinetics by which PDFR signaling terminates do not depend exclusively on the activities of either dGPRK-

1 or -2, or on that of the *Drosophila* β-arrestin2 ortholog, *krz*.

DISCUSSION

We employed a gain-of-function approach to measuring neuropeptide GPCR function *in vivo*. In

doing so, we sought to learn about the maximal consequences of PDFR GPCR signaling and also learn

something about the phosphorylation events that normally control the duration of PDFR's signaling

lifetime. We found that expression of a PDF receptor variant, containing numerous Ala-substitutions of

conserved phosphorylatable residues in the C terminal domain (1-7A), fundamentally altered rhythmic

locomotor behavior in *Drosophila*. The changes included increases in the amplitudes of the Morning

and/or Evening activity peaks, delays in their phases and/ or lengthening of the free-running period.

Such results are generally consistent with a prediction whereby non-phosphorylatable PDFR variants –

those with a potential to increase the duration of PDFR signaling - would produce behavioral actions

opposite to those seen in loss-of-function *pdf* [1], or *pdfr* [11] mutant stocks. Accordingly, we speculate

that phosphorylation of the PDFR at these conserved residues normally defines the duration of its active

signaling state each day. The longer duration that follows the substitutions of Ala may allow receptor

signal strength to also increase by accrual: such an effect could explain the increased amplitude of the

Morning and Evening peaks, as seen with expression of some of the variants. In this context, we

acknowledge that modification of GPCRs by phosphorylation does not exclusively lead to signal

termination. GRK2-dependent phosphorylation of the smoothened (smo) GPCR follows reception of the

Hh signal and helps mediate its signal: phosphorylation leads to smo <u>activation</u> in both *Drosophila* and

mammalian systems [41]. Thus modification of putative phosphorylation sites on GPCRs (to preclude

phosphorylation) will not exclusively promote extended signaling. Therefore our interpretations must

correspondingly consider outcomes without *a priori* assumption of mechanisms.

GPCR signal termination. PDFR belongs to the Secretin Receptor Family (Family B) of

neuropeptide receptors [11-13]: there is no clear consensus regarding mechanisms of desensitization

and internalization for this receptor family. For VPAC2 receptors, phosphorylation and internalization is

mediated exclusively by GRK2 [42]. Likewise, *Drosophila* orthologues of Cortocotrophin Releasing Factor

receptors (CG8422, DH44-R1) and Calcitonin receptors (CG17415, DH31-R1) are internalized in *hEK* cells

following recruitment of β-arrestin2 [31-32]. In contrast, PDFR, which is also related to the mammalian

Calcitonin receptor, is not internalized following exposure to PDF [29]. VPAC2 receptors are also

regulated by heterologous receptor signaling: M3 cholinergic receptors via PKC signaling can block

VPAC2 phosphorylation, desensitization and internalization [43]. Furthermore, secretin receptors and

VPAC1 receptors undergo phosphorylation by GRKs and β-arrestin2-dependent desensitization, but

these are not sufficient to facilitate or mediate internalization [44-46]. Finally, GLP2-Receptor associates

with β-arrestin2 via its distal C terminal sequences, but that receptor domain is required neither for

GLP2-R desensitization nor its internalization [47]. Thus kinases other than GRKs and effectors other

than β-arrestin2 may regulate internalization of diverse Family B receptors.

Following activation of rhodopsin in the mammalian retina, visual arrestin is recruited with a

time constant of < 80 ms [48]; for many *Drosophila* neuropeptide receptors, β-arrestin2 is recruited

within a minute of exposure to ligand [30]. The PDFR GPCR signals over a time base of many hours [19]:

could phosphorylation/de-phosphorylation also regulate its activity? The mammalian blue-light

sensitive GPCR melanopsin (OPN4) mediates intrinsic light sensitivity in certain classes of retinal

ganglion cells (RGCs): melanopsin signaling is distinguished by long latencies, graded responses and

sustained RGC depolarization that can outlast the duration of the light stimulus [49]. Melanopsin-

expressing RGCs produce exceptionally long time-course integration because that GPCR behaves

differently from other opsins in two fundamental ways. Firstly, it undergoes photoequilibration

between signaling and silent states: a property that maintains the availability of pigment molecules for

activation, termed 'tristability' by Emmanual and Do [50]. Secondly, phosphorylation and β-arrestin

recruitment do contribute to the kinetics of melanopsin signaling, but over a long time-course [51-52].

Mure *et al.* [34] proposed that the distal portion of the melanopsin C terminal tail creates a steric

blockade covering a cluster of specific serine residues situated in more proximal regions. The time

required for relief of that blockade dictates the time course of GPCR phosphorylation and therefore

delays subsequent desensitization.

Specific versus non-specific phosphorylation regulating GPCR activity. The 1-7A PDFR variant

modified the greatest number of phosphorylatable residues in the series we tested (**Fig. 1**), and typically

produced the strongest behavioral effects. Notably, the 6A variant – which modified only two specific

Ser residues – produced effects that were nearly identical to those exhibited by flies expressing the 1-7A

form. This suggests that much PDFR post-translational modification, that which is capable of restricting

its signaling time-course, may be directed to specific phosphorylation sites near the extreme C terminal

end. There are two broadly divergent hypotheses to describe the mechanisms by which GPCR

phosphorylation promotes desensitization. The first proposes that modification of specific residues

have the greatest significance for downstream desensitizing mechanisms: such a mechanism can explain

the effects we have seen with the highly limited 6A PDFR variant. The second hypothesis invokes

triggering of a termination processes by the aggregate negative charge accumulated with bulk

phosphorylation, regardless of where it might occur along a GPCR's intracellular sequences. For some

GPCRs (e.g., OPN4), the evidence supports both models [34, 35]. Our data concerning PDFR

desensitization also support both viewpoints and we present a model in **Figure 7**. In particular we

regard CL6 and CL7 to be the specific residues especially critical in terminating the time course of PDFR

signaling. Notably, converting just the single pair of CL6 AAs, or the single pair of CL7 AAs, is enough to

generate hours-long delays in the peaks of locomotor rhythms under Light:Dark conditions, as well as

generating a significant increase in tau under Constant Dark conditions. We note that the splicing event

that distinguishes the PDFR-A and PDFR-D isoforms occurs just prior to the position of sequences

encoding CL6 and 7, such that the D form lacks these two highly conserved domains

(flybase.org/download/sequence/FBgn0260753/FBpp). Other clusters (like CL2-3) also appear to have

potential for 'specific' contributions to PDFR regulation. Together, these results speak to the potency of

the inferred specific termination mechanisms for PDFR GPCR signaling. In contrast, the evidence for the

bulk phosphorylation hypothesis comes from comparison of the 1-4A, versus the 1-5A, 1-6A and 1-7A

variants. This series inactivates increasing numbers of phosphorylatable residues, and with it we

observed increasingly delayed evening phases in short day conditions (e.g., **Suppl**. **Fig. 2I-2, 2J-2 and 2K-**

2). The tandem mass-spectroscopy results, indicating endogenous phosphorylation of certain PDFR

residues (**Supplemental Table 4**) demonstrates such post-translational modifications can occur.

However, we caution that these phospho-peptide measurements derive from whole head extracts and

may not provide a complete or relevant accounting of those sites that are modified in critical E

pacemaker neurons.

515   Our results also point to what we propose as "context-dependent" effects of modifying

specific GPCR residues – ones that may produce opposing behavioral effects, depending on the

phosphorylation status of neighboring sites. In particular we point to the opposing results we observed

with Ala-variants of the CL5 site.   In comparing results from the series 1-4A, 1-5A, and 1-6A, we found

that under Short Day conditions, the 1-4A had only mild effects on the evening peak phase compared to

over-expressing WT PDFR (e.g., **Suppl. Fig**. **2B-1** vs. **2I-1**), whereas 1-5A and 1-6A both produced

significant delays (**Suppl**. **Fig. 2M and Suppl. Fig. 3J-3 through 3K-3)**. The difference between the 1-4A

and 1-5A variants is a change in a single S residue at CL5 (S633A): Those observations suggest

modification of the CL5 residue normally promotes desensitization of PDFR and termination of PDFR

signaling. However, the PDFR variant 5-7A was constructed with the expectation that it would be as

effective as 1-5A, 1-6A or 1-7A in delaying the evening phase. Instead it proved only weakly effective.

This same mis-match of expected behavioral effects for 5-7A was also seen in other photoperiodic

conditions (**Suppl**. **Fig. 6M**). Likewise, Langlet *et al*. [36] reported that effects on mutating Serine

residues in the carboxy terminus of VPAC1 did not produce additive effects. We propose that

phosphorylation of CL5 will have either positive or negative consequences on PDFR desensitization

depending on which other neighboring residues are also modified. Thus we speculate that CL5 takes on

outsize importance in determining desensitization rates for PDFR and so may itself be subject to

exceptional regulation. Such complex interactions between phosphorylation sites is reminiscent of

interactions documented between diverse phosphorylation sites in the circadian clock protein PERIOD

[53-54]. Better resolution of these two paradoxical mechanisms ((i) specific phosphorylation versus

bulk negative charge, and (ii) context-dependent effects of CL5 site phosphorylation, awaits precise

molecular definition of where and when PDFR is modified in key pacemaker neurons *in vivo*, and by

which post-translational modifications.

Validating the behavioral phenotypes produced by PDFR variants. The actions of PDFR

variants we have described, following substitutions of Ala for various Ser/Thr or Tyr residues, are not

easily explained by a hypothesis invoking neomorphic or constitutive GPCR properties.   The evidence for

this conclusion is three-fold. First, the effects on both amplitude and phase of activity peaks by these

variants are not of a random assortment: rather they are all strictly opposite to that of loss-of-function

models for *pdf* and *pdfr*.   Second, the actions of PDFR variants display strong dependence on wild type

pdf gene function: this strongly argues that the actions of PDFR variants reflect responses to normal,

endogenous PDF signaling. Third, even the strongest actions of PDFR variants on the phases and

amplitudes of locomotor activity (that of the 1-7A variant) reflected PDFR sequence variation, and not

the properties of the GFP C-terminal fusion. Together these observations support the hypothesis that

phosphorylation of some or all of the C terminal residues we studied are normally modified to attenuate

the strength and duration of PDFR signaling during the 24-hr day.

PDF Signaling and seasonal adaptation. PDF neuropeptide actions were discovered in the

context of physiological adaptions to daylight in crustacea [5,6]. Helfrich-Förster [55] discovered PDF

expression within a defined subset of the insect *Drosophila* circadian pacemaker network. Renn *et al.*

[1] demonstrated that *pdf* makes a fundamental contribution by setting the normal behavioral phase of

Evening activity in *Drosophila* during light:dark entrainment and promoting normal rhythmicity during

constant darkness. PDF signaling coordinates with that driven by light: It works in parallel to direct

photosensitivity along with the CRY blue light photoreceptor. Flies doubly-mutant for *cry* and *pdf*

display pronounced deficits of locomotor rhythmicity [7, 8, 9], and can be as severe as measured in

clock-deficient flies. PDF and environmental light also work coordinately at the level of neuronal activity

patterns: in the case of the evening pacemakers (LNd and the 5th s-LNv), both light and PDF signaling

promote a delay in their PER-dependent activation period, and together help align it to a phase just

prior to dusk [19]. Based on the close association of PDF signaling and photoperiodic signaling, it is not

surprising that the duration of the photoperiod strongly influenced the effects of PDFR sequence

variants on the Morning versus the Evening activity peaks. We only observed delayed Evening activity

peak produced by PDFR gain-of-function variants under Short Day conditions: that delay appears

strongly inhibited by light durations > 8 hr. It remains unclear at what molecular level light exerts such

an inhibitory effect, however, we note a similarity of these observations with effects recently reported

from loss-of-function states for the phosphatase *PRL-1* [56]. Like the most active PDFR variants we

herein describe (e.g., 6A, 7A, 1-5A 1-6A and 1-7A), *PRL-1* mutants produce a 3-4 h delay in the phase of

the evening activity peak, but only under winter-like (short day) conditions, and not under equinox or

summer-like (long day) conditions. Kula-Eversole *et al.* [56] have shown that TIM phosphorylation is

affected by PRL-1 activity and suggest the seasonal action of PRL-1 to advance the evening locomotor

activity phase is mediated by modification of TIM levels within the transcription-translation feedback

loop. The extensive similarity of these two sets of behavioral phenotypes reveals either serial or parallel

pathways to effect comparable outcomes on locomotor activity. If serial, then according to the simplest

model, PRL1 acts downstream of PDFR, and the combined results to-date suggest PRL-1 would be

inhibited by PDFR activation. Further genetic and biochemical experiments are needed to evaluate if and

how these pathways converge.

Relations between in vitro and in vivo PDFR signaling measures. Using an *in vitro* assay for

cyclic AMP generation, we found that modifying phosphorylation properties of PDFR does not affect the

strength of signaling. That conclusion suggests that *in vivo* other GPCR features are normally affected to

regulate the extent/duration of PDFR signaling according to season. This speculation corresponds to

findings described with other Family B GPCRs like VPAC1 [46]: Mutation of all the Ser and Thr residues of

the C terminal tail, and of Ser250 to Ala, led to a receptor with binding properties and adenylate cyclase

activity similar to the wild type receptor; however that variant receptor was neither phosphorylated nor

internalized. We propose that the PDFR variants we have described do not modify locomotor behavior

by virtue of greater second messenger signaling. Rather they do so by generating cAMP (and perhaps

other second messengers) over time periods longer than that normal sustained by a WT receptor over a

portion of each 24-hr cycle.

Is PDFR modulation of diurnal phase independent of its modulation of circadian period? Many

PDFR variants strongly delayed the peak of evening activity (past lights-off) and also significantly

lengthened the circadian period (measured under constant darkness). A conventional interpretation of

these linked phenotypes is that PDFR signaling affects the pace of the molecular clock (re-setting the

period) and consequently affects behavioral phase indirectly, which is downstream of the clock. This

explanation holds for all the variants that both increased tau and delayed the evening phase (e.g., 1-5A,

1-6A, and 1-7A). However, the correlation was not always observed. For example, under winter-like

conditions, the 5A, 6A, 7A and 5-7A variants all significantly delayed the offsets of Evening activity peaks

versus that seen with WT PDFR, but these variants did not significantly lengthen circadian period.

Likewise, under long day, 6A produced a significant increase in tau versus that seen with WT PDFR, but

did not significantly delay the onset of the Evening activity peak. Therefore, we cannot rule out an

alternative hypothesis: that the duration of PDF>PDFR signaling to target pacemakers in the *Drosophila*

brain is modulated each day to effect (i) entrainment of the molecular clock, and independently (ii) to

directly delay to Evening pacemaker neuronal activity. In fact, the *pdfr han* mutant flies exhibited quasi-

normal tau values in this report (Table 1), although they more traditionally exhibit shortened ones in our

experience [37]. Evening locomotor activity derives in large part from the activity of the primary

Evening oscillators, termed E cells, the LNd and the 5[th] small LNv [10, 20, 57 – 59]. However, numerous

network interactions are also known to provide critical contributions to the accuracy, precision and

adaptability of that timing system [e.g., 59 – 64]. The Evening oscillator group itself is known to be

heterogeneous and to constitute at least three separate, functionally distinct oscillators [60], only some

of which express PDFR-A and respond to PDF [37, 60]. Vaze and Helfrich-Förster [65] reported that the

phase of Period protein accumulation in E neurons is phase-locked to the previous lights-off transition

and also sensitive to PDF signaling. They conclude that the peak of PER accumulation is key to

determining the phase of the evening activity peak, but also note that the correlation between the

Period-clock timing cue and the Evening activity peak is not perfect. They suggest the Evening

behavioral phase is better described by also factoring in the delay in E cell neuronal activity driven by

PDF neuromodulation [10, 19]. We speculate that it is the combination of two separate PDFR signaling

effects that is essential for proper rhythmic behavioral outcomes across seasons. If valid, the hypothesis

predicts a bifurcation of signaling pathways downstream of PDFR, to independently regulate diurnal

phase and also circadian period within E pacemaker neurons.

MATERIALS and METHODS

Fly Rearing and Stocks. *Drosophila* were raised on a cornmeal agar diet supplemented with

yeast at 25°C in 12 hr:12 hr LD cycles. The UAS-*pdfr* mutant transgenic series was created by injecting

yw P{nos phiC31\int.NLS}X;P{CaryP}attP40 embryos (Rainbow Transgenic Flies, Inc, Camarillo, CA).

pdfr mutant flies are described in [11]; *tim*(UAS)-Gal4 flies are described in [66] – *BL80941*). The

recombinant fly stock - *han^{5304}*; *tim* (UAS)-Gal4 - was confirmed by PCR and sequencing. The UAS-*pdfr*-

tandem construct was injected into *y[1] w[*] P{y[+t7.7]=nos-phiC31\int.NLS}X;*

P{y[+t7.7]=CaryIP}su(Hw)attP6 embryos. For PDFR-tandem fusion protein expression, we made a stable

yw; UAS-*pdfr*-tandem; *tim*-Gal4 stock.

hEK-293 Cell Culture. *hEK-293* cells were maintained in DMEM, 10% FBS and 100U/mL penicillin

and streptomycin in 5% CO_2 atmosphere at 37°C. For all transient transfections, 1.5 x 10^6 cells were used

to inoculate T25 flasks, incubated overnight, then transfected with 10ug plasmid DNA and 20 uL

lipofectamine 2000 reagent (Invitrogen Life Technologies). Five hours after transfection, cells were split

4 x 10^4 cells/well into a 96-well assay plate. We created a series of stably-transfected *hEK-293* cells

expressing WT and sequence-variant PDFRs, using the Flp-In System (Invitrogen Life Technologies,

Waltham, Massachusetts) per manufacturers recommendations, and maintained them in DMEM

supplemented with 150 µg/ml hygromycin B.

cAMP Assays. We measured PDF Receptor signaling activity using a CRE-*Luciferase* reporter

gene, following methods described by Johnson *et al.* [30]. The reporter gene construct was transiently

transfected to each stable cell line and luminescence measured using Firefly Luciferase Assay Kit

(Biotium, Inc., Fremont, California) and a Wallac 1420 VICTOR2 microplate reader (PerkinElmer, Inc.,

Waltham, Massachusetts). Concentration-effect curves, EC_{50}, top values and p-values were calculated

using the dose response, in a nonlinear regression using GraphPad Prism 8.0 software (San Diego,

California).

Locomotor activity Measures. All locomotor activity experiments were conducted with 2-5

days-old male flies at 25°C using Trikinetics Activity Monitors as previously described [9]. We crossed

Gal4 lines to w^{1118} to create control progeny. Locomotor activities were monitored for 6 days under

different photoperiodic conditions, and then for 9 days under constant dark (DD). To analyze rhythmicity

under constant conditions, we normalized the activity of flies from DD day 3 to day 9 and used χ2-

periodogram analysis with a 95% confidence cut-off, as well as SNR analysis [67] Arrhythmic flies were

defined by a power value $\leq$ 10 and width value $\leq$ 2, and period outside the range, 18 to 30 hours. To

analyze periods, we used Graphpad Instat (v. 8) software to run one-way ANOVA measures followed by

the Tukey-Kramer Multiple Comparisons Test. We used Clocklab (Actimetrics) software to produce

actograms and the Brandeis Rhythms Package [73] to produce average activity plots (group eductions).

To analyze onset and offset phases of Morning and Evening activity bouts on the final two days of

light:dark entrainment (LD 5&6), we followed the method of Kula-Eversole *et al.* [56]: $[(A_{n+2} + A_{n+1}) - (A_{n-1} + A_{n-2})] = \Delta Activity$

$_1 + A_{n-2}) = \Delta Activity$]. Experimental genotypes were tested in the *han* mutant background*, pdfr^{5304}* [11]*,* or

in the *pdf^{01}* (null) mutant background [1], or in the w^{1118} background, as noted.

Presentation of behavioral data with expression of PDFR variants. Figures 2-4 all describe

rhythmic locomotor behavior of adult *pdfr^han^* mutant flies that are expressing either a WT *pdfr* cDNA or

the 1-7A variant. Figures 5 and 6 use a similar format but they introduce behavioral experiments

performed in *pdfr^+^* or *pdf* mutant backgrounds, as described. All behavioral experiments extend across

~6 days of light entrainment (LD), followed by activity during nine or more days of constant darkness

(DD). For each genotype, we display locomotor activity during light entrainment in four ways: (i) group

eductions (in panels marked **A, B and C**-**1** in these Figures), (ii) direct comparisons between two

genotypes of the amplitudes of behavioral peaks using bin-by-bin analyses during the last two days of

entrainment (days 5-6) (in panels marked **A, B and C-2-4**); (iii) derivations of Morning and Evening

behavioral phases (in Panels **D and E**); and (iv) double-plotted group actograms (in Panels **F, G and H**).

and The Bin-by-Bin analyses directly compare experimental genotypes to a control one [cf. 42]. They

present data averaged across 2 to 5 independent experiments, to help identify the differences that most

consistently correlate with genotype. Finally, Table 1 compiles measures of rhythmic activity displayed

by the different genotypes in days 3-9 of constant darkness.

Additional details on Methods and procedures are found in **Supplemental Information**.

ACKNOWLEDGEMENTS

We thank many colleagues for generously sharing reagents, fly strains and providing technical advice, all

of which helped us pursue these studies. These include Xuemei Si, and Drs. Ali Salahpour, Dmitri

Nusinow, Johanna Chiu, Xitong Liang, Paul Hardin, Zsolt Lenkei, Ken Blumer and David Wheeler. We

thank the staff that manages the Flybase resource which provides valuable information, and the

Danforth Plant Science Center Mass Spectroscopy Facility for expert assistance, and the Bloomington

Drosophila Stock Center, and the Vienna *Drosophila* Research Center for providing many valuable fly

stocks. We also thank Joel Levine and Orie Shafer for kindly providing comments on the manuscript.

AUTHOR CONTRIBUTIONS

W.L., J.T. and P.H.T designed experiments. W.L. and J.T. performed experiments. W.L., J.T. and P.H.T

analyzed experiments. P.H.T wrote the manuscript.

LITERATURE CITED

1. Renn SC, Park JH, Rosbash M, Hall JC, Taghert PH. A *pdf* neuropeptide gene mutation and ablation of

PDF neurons each cause severe abnormalities of behavioral circadian rhythms in *Drosophila*.

Cell. 1999; 99:791-802.   PMID: 10619432.

2. Peng Y, Stoleru D, Levine JD, Hall JC, Rosbash M. *Drosophila* free-running rhythms require intercellular

communication. PLoS Biol. 2003; 1:E13. PMID: 12975658.

3. Lin Y, Stormo GD, Taghert PH. The neuropeptide pigment-dispersing factor coordinates pacemaker

interactions in the *Drosophila* circadian system. J Neurosci. 2004; 24:7951-7. PMID: 15356209.

4. Vosko AM, Schroeder A, Loh DH, Colwell CS. Vasoactive intestinal peptide and the mammalian

circadian system. Gen Comp Endocrinol. 2007; 152:165-75. PMID: 17572414.

5. Fernlund, P. Structure of a light-adapting hormone from the shrimp, *Pandalus borealis.* Biochim

Biophysica Acta 1976; 439: 17-25. PMID: 95295.

6. Rao KR, Riehm JP, Zahnow CA, Kleinholz LH, Tarr GE, Johnson L, Norton S, Landau M, Semmes OJ,

Sattelberg RM, Jorenby WH, Hintz MF. Characterization of a pigment-dispersing hormone in

eyestalks of the fiddler crab *Uca pugilator*. Proc Nat'l Acad Sci (Wash) U S A. 1985; 82:5319-22.

PMID: 16593589.

7. Cusumano P, Klarsfeld A, Chélot E, Picot M, Richier B, Rouyer F. PDF-modulated visual inputs and

cryptochrome define diurnal behavior in *Drosophila*. Nat Neurosci. 2009; 12:1431-7. PMID:

19820704.

8. Zhang L, Lear BC, Seluzicki A, Allada R. The CRYPTOCHROME photoreceptor gates PDF neuropeptide

signaling to set circadian network hierarchy in *Drosophila*. Curr Biol. 2009; 19:2050-5. PMID:

19913424.

9. Im SH, Li W, Taghert PH. PDFR and CRY signaling converge in a subset of clock neurons to modulate

the amplitude and phase of circadian behavior in *Drosophila*. PLoS One. 2011; 6:e18974. PMID:

21559487.

10. Liang X, Holy TE, Taghert PH. Synchronous *Drosophila* circadian pacemakers display nonsynchronous

Ca^{2+} rhythms. Science (Wash). 2016; 351:976-81. PMID: 26917772.

11. Hyun S, Lee Y, Hong ST, Bang S, Paik D, Kang J, Shin J, Lee J, Jeon K, Hwang S, Bae E, Kim J. *Drosophila*

GPCR Han is a receptor for the circadian clock neuropeptide PDF. Neuron. 2005; 48:267-78.

PMID: 16242407.

12. Lear BC, Merrill CE, Lin JM, Schroeder A, Zhang L, Allada R. A G protein-coupled receptor, groom-of-

PDF, is required for PDF neuron action in circadian behavior. Neuron. 2005; 48:221-7. PMID:

16242403.

13. Mertens I, Vandingenen A, Johnson EC, Shafer OT, Li W, Trigg JS, De Loof A, Schoofs L, Taghert PH.

PDF receptor signaling in *Drosophila* contributes to both circadian and geotactic behaviors.

Neuron. 2005; 48:213-9. PMID: 16242402.

14. Shafer OT, Kim DJ, Dunbar-Yaffe R, Nikolaev VO, Lohse MJ, Taghert PH. Widespread receptivity to

neuropeptide PDF throughout the neuronal circadian clock network of *Drosophila* revealed by

real-time cyclic AMP imaging. Neuron. 2008; 58:223-37. PMID: 18439407.

15. Duvall LB, Taghert PH. The circadian neuropeptide PDF signals preferentially through a specific

adenylate cyclase isoform AC3 in M pacemakers of *Drosophila*. PLoS Biol. 2012; 10:e1001337.

PMID: 22679392.

16. Duvall LB, Taghert PH. E and M circadian pacemaker neurons use different PDF receptor

signalosome components in *Drosophila*. J Biol Rhythms. 2013; 28:239-48. PMID: 23929551.

17. Seluzicki A, Flourakis M, Kula-Eversole E, Zhang L, Kilman V, Allada R. Dual PDF signaling pathways

reset clocks via TIMELESS and acutely excite target neurons to control circadian behavior. PLoS

Biol. 2014; 12(3):e1001810.. PMID: 24643294.

18. Depetris-Chauvin A, Fernández-Gamba A, Gorostiza EA, Herrero A, Castaño EM, Ceriani MF. Mmp1

processing of the PDF neuropeptide regulates circadian structural plasticity of pacemaker

neurons. *PLoS Genet*. 2014; doi:10.1371/journal.pgen.1004700. PMID: 25356918.

19. Liang X, Holy TE, Taghert PH. A Series of Suppressive Signals within the *Drosophila* Circadian Neural

Circuit Generates Sequential Daily Outputs. Neuron. 2017; 94:1173-1189. PMID: 28552314.

20. Liang X, Ho MCW, Zhang Y, Li Y, Wu MN, Holy TE, Taghert PH. Morning and Evening Circadian

Pacemakers Independently Drive Premotor Centers via a Specific Dopamine Relay. Neuron.

2019; 102:843-857. PMID: 30981533.

21. Pincas H, González-Maeso J, Ruf-Zamojski F, Sealfon SC. G Protein-Coupled Receptors. In:

Belfiore A, LeRoith D, editors. Principles of Endocrinology and Hormone Action.

Endocrinology. Springer, Cham. https://doi.org/10.1007/978-3-319-44675-2_6. 2018.

22. Lefkowitz RJ, Shenoy SK. Transduction of receptor signals by β-Arrestins. Science (Wash). 2005;

308:512-7. PMID: 15845844.

23. Benovic JL, Kühn H, Weyand I, Codina J, Caron MG, Lefkowitz RJ. Functional desensitization of the

isolated beta-adrenergic receptor by the beta-adrenergic receptor kinase: potential role of an

analog of the retinal protein arrestin (48-kDa protein). Proc Nat'l Acad Sci (Wash) U S A. 1987;

84:8879-82. PMID: 2827157.

24. Lohse MJ, Benovic JL, Codina J, Caron MG, Lefkowitz RJ. β-Arrestin: a protein that regulates beta-

adrenergic receptor function. Science (Wash). 1990; 248:1547-50. PMID: 2163110.

25. Ferguson SS, Downey WE 3rd, Colapietro AM, Barak LS, Ménard L, Caron MG. Role of β-arrestin in

mediating agonist-promoted G protein-coupled receptor internalization. Science (Wash). 1996;

271:363-6. PMID: 8553074.

26. Smith JS, Rajagopal S. The β-Arrestins: Multifunctional Regulators of G Protein-coupled Receptors. J

Biol Chem. 2016; 291:8969-77. PMID: 26984408.

27. Kelly E, Bailey CP, Henderson G. Agonist-selective mechanisms of GPCR desensitization. Brit J of

Pharm. 2008; 153 Suppl 1: S379–88. PMID 18059321.

28. Abruzzi, K. C., Zadina, A., Luo, W., Wiyanto, E., Rahman, R., Guo, F., Shafer, O., & Rosbash, M. RNA-

seq analysis of Drosophila clock and non-clock neurons reveals neuron-specific cycling and novel

candidate neuropeptides. PLoS Genetics, 2017; *13*(2), e1006613

doi.org/10.1371/journal.pgen.1006613

29. Klose M, Duvall L, Li W, Liang X, Ren C, Steinbach JH, Taghert PH. Functional PDF Signaling in the

Drosophila Circadian Neural Circuit Is Gated by Ral A-Dependent Modulation. Neuron. 2016;

90:781-794. PMID: 27161526.

30. Johnson EC, Bohn LM, Barak LS, Birse RT, Nässel DR, Caron MG, Taghert PH. Identification of

Drosophila neuropeptide receptors by G protein-coupled receptors-beta-arrestin2 interactions. J

Biol Chem. 2003; 278(52):52172-8. PMID: 14555656.

31. Johnson EC, Bohn LM, Taghert PH. *Drosophila* CG8422 encodes a functional diuretic hormone

receptor. J Exp Biol. 2004; 207:743-8. PMID: 14747406.

32. Johnson EC, Shafer OT, Trigg JS, Park J, Schooley DA, Dow JA, Taghert PH. A novel diuretic hormone

receptor in *Drosophila*: evidence for conservation of CGRP signaling. J Exp Biol. 2005; 208:1239-

46. PMID: 15781884.

33. Throckmorton, LH. The phylogeny, ecology and geography of *Drosophila*. In: King RC., editor.

Handbook of Genetics. Vol 3. New York: Plenum; 1975. pp 421-469.

34. Mure LS, Hatori M, Zhu Q, Demas J, Kim IM, Nayak SK, Panda S. Melanopsin-Encoded Response

Properties of Intrinsically Photosensitive Retinal Ganglion Cells. Neuron. 2016; 90:1016-27.

PMID: 27181062.

35. Valdez-Lopez JC, Gulati S, Ortiz EA, Palczewski K, Robinson PR. Melanopsin Carboxy-terminus

phosphorylation plasticity and bulk negative charge, not strict site specificity, achieves

phototransduction deactivation. PLoS One. 2020; 15:e0228121. PMID: 32236094.

36. Langlet C, Langer I, Vertongen P, Gaspard N, Vanderwinden JM, Robberecht P. Contribution of the

carboxyl terminus of the VPAC1 receptor to agonist-induced receptor phosphorylation,

internalization, and recycling. J Biol Chem. 2005; 280: 28034 −28043. PMID: 15932876.

37. Im SH, Taghert PH PDF receptor expression reveals direct interactions between circadian oscillators

in *Drosophila.* J Comp Neurol. 2010; 518:1925-45. PMID**:** 20394051.

38. Zhang Y, Emery P. GW182 controls *Drosophila* circadian behavior and PDF-receptor signaling.

Neuron. 2013; Apr 10;78(1):152-65. PMID: 23583112.

39. Beerepoot P, Lam VM, Salahpour A. Measurement of G protein-coupled receptor surface

expression. J Recept Signal Transduct Res. 2013; Jun;33(3):162-5. PMID: 23557016.

40. Zhou XE, He Y, de Waal PW, Gao X, Kang Y, Van Eps N, Yin Y, Pal K, Goswami D, White TA, Barty A,

Latorraca NR, Chapman HN, Hubbell WL, Dror RO, Stevens RC, Cherezov V, Gurevich VV, Griffin

PR, Ernst OP, Melcher K, Xu HE. Identification of Phosphorylation Codes for Arrestin Recruitment

by G Protein-Coupled Receptors. Cell. 2017; 170:457-469.e13. PMID: 28753425.

41. Li S, Li S, Han Y, Tong C, Wang B, Chen Y, Jiang, J. (2016). Regulation of Smoothened Phosphorylation

and High-Level Hedgehog Signaling Activity by a Plasma Membrane Associated Kinase. PLoS

Biology, 2016; *14*(6), e1002481. PMID: 27280464.

42. Murthy KS, Mahavadi S, Huang J, Zhou H, and Sriwai W. Phosphorylation of GRK2 by PKA augments

GRK2-mediated phosphorylation, internalization, and desensitization of VPAC2 receptors in

smooth muscle. Am J Physiol Cell Physiol. 2008; 294: C477–C487. PMID: 18077607.

43. Huang J, Mahavadi S, Sriwai W, Grider JR, Murthy KS. Cross-regulation of VPAC(2) receptor

desensitization by M(3) receptors via PKC-mediated phosphorylation of RKIP and inhibition of

GRK2. Am J Physiol Gastrointest Liver Physiol. 2007; 292:G867-74. PMID: 17170028.

44. Shetzline MA, Premont RT, Walker JK, Vigna SR, Caron MG. A role for receptor kinases in the

regulation of class II G protein-coupled receptors. Phosphorylation and desensitization of the

secretin receptor. J Biol Chem. 1998 Mar 20;273(12):6756-62. PMID: 9506976.

45. Walker JK, Premont RT, Barak LS, Caron MG, Shetzline MA. Properties of secretin receptor

internalization differ from those of the 2-adrenergic receptor. J Biol Chem 1999; 274: 31515–

31523. PMID: 10531354.

46. Marie JC, Rouyer-Fessard C, Couvineau A, Nicole P, Devaud H, El Benna J, Laburthe M. Serine 447 in

the carboxyl tail of human VPAC1 receptor is crucial for agonist-induced desensitization but not

internalization of the receptor. Mol Pharmacol. 2003; 64: 1565–1574. PMID: 14645688.

47. Estall JL, Koehler JA, Yusta B, Drucker DJ. The glucagon-like peptide-2 receptor C terminus

modulates beta-arrestin-2 association but is dispensable for ligand-induced desensitization,

endocytosis, and G-protein-dependent effector activation. J Biol Chem. 2005; 280:22124-34.

PMID: 15817468.

48. Krispel CM, Chen D, Melling N, Chen YJ, Martemyanov KA, Quillinan N, Arshavsky VY, Wensel TG,

Chen CK, Burns ME. RGS expression rate-limits recovery of rod photoresponses. Neuron. 2006;

51:409-16. PMID: 16908407.

49. Berson DM, Dunn FA, Takao M. Phototransduction by retinal ganglion cells that set the circadian

clock. Science (Wash). 2002; 295:1070-3. PMID: 11834835.

50. Emanuel AJ, Do MT. Melanopsin tristability for sustained and broadband phototransduction.

Neuron. 2015; 85:1043-55. PMID: 25741728.

51. Somasundaram P, Wyrick GR, Fernandez DC, *et al.* C-terminal phosphorylation regulates the kinetics

of a subset of melanopsin-mediated behaviors in mice. Proc Nat'l. Acad Sci (Wash) U S A. 2017;

114:2741-2746. PMID: 28223508.

52. Dephoure N, Gould KL, Gygi SP, Kellogg DR. Mapping and analysis of phosphorylation sites: a quick

guide for cell biologists. Mol Biol Cell. 2013; 24:535-42. PMID: 23447708.

53. Chiu JC, Ko HW, Edery I. (2011) NEMO/NLK phosphorylates PERIOD to initiate a time-delay

phosphorylation circuit that sets circadian clock speed. Cell. 2011; 145:357-70. PMID: 21514639.

54. Yu W, Houl JH, Hardin PE. NEMO kinase contributes to core period determination by slowing the

pace of the *Drosophila* circadian oscillator. Curr Biol. 2011; 21:756-61. PMID: 21514156.

55. Helfrich-Förster C. The period clock gene is expressed in central nervous system neurons which also

produce a neuropeptide that reveals the projections of circadian pacemaker cells within the

brain of *Drosophila melanogaster*. Proc Nat'l Acad Sci (Wash) U S A. 1995; 92:612-6. PMID:

7831339.

56. Kula-Eversole E, Lee DH, Samba I, Yildirim E, Levine DC, Hong HK, Lear BC, Bass J, Rosbash M, Allada

R. Phosphatase of Regenerating Liver-1 Selectively Times Circadian Behavior in Darkness via

Function in PDF Neurons and Dephosphorylation of TIMELESS. Curr Biol. 2021; 31:138-149.e5.

PMID: 33157022.

57. Grima B, Chélot E, Xia R, Rouyer F. Morning and evening peaks of activity rely on different clock

neurons of the *Drosophila* brain. Nature. 2004; Oct 14;431(7010):869-73. PMID: 15483616.

58. Stoleru D, Peng Y, Agosto J, Rosbash M. Coupled oscillators control morning and evening locomotor

behaviour of *Drosophila*. Nature. 2004; 431:862-8. PMID: 15483615.

59.. Rieger D, Shafer OT, Tomioka K, Helfrich-Förster C. Functional analysis of circadian pacemaker

neurons in Drosophila melanogaster. J Neurosci. 2006; Mar 1;26(9):2531-43. PMID: 16510731.

60. Zhang Y, Liu Y, Bilodeau-Wentworth D, Hardin PE, Emery P. Light and temperature control the

contribution of specific DN1 neurons to *Drosophila* circadian behavior. Curr Biol. 2010; Apr

13;20(7):600-5. PMID: 20362449.

61. Yao Z, Shafer OT. The *Drosophila* circadian clock is a variably coupled network of multiple peptidergic

units. Science (Wash). 2014; 343(6178):1516-20. PMID: 24675961.

62. Schlichting M, Díaz MM, Xin J, Rosbash M. Neuron-specific knockouts indicate the importance of

network communication to *Drosophila* rhythmicity. Elife. 2019 Oct 15;8:e48301. PMID:

31613223

63. Delventhal R, O'Connor RM, Pantalia MM, Ulgherait M, Kim HX, Basturk MK, Canman JC, Shirasu-Hiza

M. Dissection of central clock function in *Drosophila* through cell-specific CRISPR-mediated clock

gene disruption. Elife. 2019;8:e48308. PMID: 31613218.

64. Menegazzi P, Beer K, Grebler V, Schlichting M, Schubert FK, Helfrich-Förster C. A Functional Clock

Within the Main Morning and Evening Neurons of *D. melanogaster* Is Not Sufficient for Wild-

Type Locomotor Activity Under Changing Day Length. Front Physiol. 2020; Mar 26;11:229. PMID:

32273848.

65. Schlichting M, Menegazzi P, Lelito KR, Yao Z, Buhl E, Dalla Benetta E, Bahle A, Denike J, Hodge JJ,

Helfrich-Förster C, Shafer OT. A Neural Network Underlying Circadian Entrainment and

Photoperiodic Adjustment of Sleep and Activity in *Drosophila*. J Neurosci. 2016; 36(35):9084-96.

PMID: 27581451.

66. Vaze KM, Helfrich-Förster C. The Neuropeptide PDF Is Crucial for Delaying the Phase

of *Drosophila's* Evening Neurons Under Long Zeitgeber Periods. J Biol Rhythms. 2021;36(5):442-

460. PMID: 34428956.

67. Blau J, Young MW. Cycling *vrille* expression is required for a functional *Drosophila* clock. Cell. 1999;

Dec 10;99(6):661-71 PMID: 10612401.

68. Levine JD, Funes P, Dowse HB, Hall JC. Signal analysis of behavioral and molecular cycles. BMC

Neurosci. 2002; 3:1. PMID: 1182533.

69. Wheeler DA, Hamblen-Coyle MJ, Dushay MS, Hall JC. Behavior in light-dark cycles of *Drosophila*

mutants that are arrhythmic, blind, or both. J Biol Rhythms. 1993; 8:67-94. PMID: 8490212.

70. Shafer OT, Taghert PH. RNA-interference knockdown of *Drosophila pigment dispersing factor* in

neuronal subsets: the anatomical basis of a neuropeptide's circadian functions. PLoS One. 2009;

4:e8298. PMID: 20011537..

FIGURE LEGENDS

Figure 1. A map illustrating the series of PDFR sequence variants used to test a role for PDFR

phosphorylation in regulating the locomotor rhythms. The C Terminal of the PDFR-A isoform is

diagrammed (top) to the right of the 7[th] transmembrane domain (TM7, blue). Asterisks indicate the

positions of the 28 serine, threonine or tyrosine residues present within the PDFR C Terminal tail. The

residues marked by the white asterisks were those chosen for study by sequence alteration to encode

Alanine; they were given arbitrary designations as Clusters (CL) 1 through 7. Each cluster contains a

single residue (e.g., CL4 or CL5), or as many as four residues (e.g., CL2). The ten sequence variants are

named by the Cluster(s) that was altered followed by the letter 'A". Simple Variants include those

wherein residues in one or two Clusters were altered. Multiple Variants include those wherein residues

within three or more Clusters were altered. All 14 targeted residues were altered in the PDFR 1-7A

variant. The green arrows mark the WT and 1-7A PDFR variant: their behavioral effects are featured in

the Main Figures. The yellow arrows mark the nine other PDFR variants: their behavioral effects are

described in Supplemental Figures.

Figure 2. Locomotor Rhythms exhibited by *pdfr* mutant flies expressing the WT PDFR versus

the 1-7A PDFR Variant under Winter-like (Short Day) conditions. All behavioral records were recorded

from *han* (*pdfr* mutant) flies that expressed either no UAS transgene (A, yellow box), or a UAS-WT *pdfr*

transgene (B, black box) or a UAS-1-7A *pdfr* (C, red box)). Panels (A)-(C) contains sub-panels 1-4. Sub-

Panel (1) displays group eductions (activity averaged over days 5 and 6 of light entrainment) with open

bars indicating the 8-hr periods of Lights-on and filled bars the 16-hr periods of Lights-off. *TG*:

transgene. The red and blue arrows indicate Morning and Evening activity periods respectively in the 1-

7A records that have distinguished amplitude or phase. Amplitude measures are displayed in Panels 2-

4: 30 min bins for each genotype, color-coded and directly-compared (i.e., Panel C-2 through C-4

compares the amplitudes of the WT PDFR activity (black) with that of the 1-7A PDFR activity (red). The

bins representing the light-dark transitions were removed. Blue asterisks mark amplitudes that are

significantly different between genotypes by ANOVA followed by a Student's T-test ($p < 0.05$). Panel (D)

displays the average Phase Onset timepoint for the Morning activity for each genotype over the last two

days of entrainment (LD 5-6). Panel (E) displays the average Phase Offset timepoint for the Evening

activity for each genotype over the last two days of entrainment (LD 5-6). Analyses in (D) and (E)

represent ANOVA followed by Dunnett's post hoc multiple comparisons of all compared to WT: ns = not

significant; * = $p<0.05$; ** = $p<0.01$; *** = $p<0.005$; **** = $p<0.001$. Panels (F) – (H) present double

plotted actograms of group activity for each genotype (color-coded) over the ~6 days of light

entrainment, followed by ~9 days of constant darkness (DD). The green bars indicate the phases of the

dominant activity periods in DD.

Figure 3. Locomotor Rhythms exhibited by *pdfr* mutant flies expressing the WT PDFR versus the 1-7A

PDFR Variant under Equinox-like (12L:12D) conditions. All behavioral records were recorded from *han*

(*pdfr* mutant) flies that expressed either no UAS transgene (A, yellow box), or a UAS-WT *pdfr* transgene

(B, black box) or a UAS-1-7A *pdfr* (C, red box)). The Panels display behavioral activity with a format

similar to the one shown in Figure 2. Panels (A)-(C) display average group eductions (activity averaged

over days 5 and 6 of light entrainment, with open bars indicating the 12-hr periods of Lights-on and

filled bars the 12-hr periods of Lights-off) and bin-by-bin analyses to compare activity amplitudes

between genotypes. *TG*: transgene. The red arrow indicates Morning activity in flies expressing the 1-7A

variant with distinguished amplitude. Blue asterisks mark amplitudes that are significantly different

between genotypes by ANOVA followed by a Student's T-test ($p < 0.05$). Panels (D) and (E) display the

phases of morning Onsets and Evening Offsets respectively. Gold asterisks: ANOVA followed by

Dunnett's post hoc multiple comparisons of all compared to WT: ns = not significant; * = p<0.05; ** =

p<0.01; *** = p<0.005; **** = p<0.001. Panels (F) – (H) present double plotted actograms of group

activity for each genotype over the ~6 days of light entrainment, followed by ~9 days of constant

darkness (DD). The green bars indicate the phase of the dominant activity period in DD.

Figure 4. Locomotor Rhythms exhibited by *pdfr* mutant flies expressing the WT PDFR versus

the 1-7A PDFR Variant under Long-day (summer-like) conditions. All behavioral records were recorded

from *han* (*pdfr* mutant) flies that expressed either no UAS transgene (A, yellow box), or a UAS-WT *pdfr*

transgene (B, black box) or a UAS-1-7A *pdfr* (C, red box)). The Panels present behavioral activity with a

format similar to the one shown in Figure 2. Panels (A)-(C) display average group eductions (activity

averaged over days 5 and 6 of light entrainment, with open bars indicating the 16-hr periods of Lights-on

and filled bars the 8-hr periods of Lights-off) and bin-by-bin analyses to compare activity amplitudes

between genotypes. *TG*: transgene. The red and blue arrows indicate Morning and Evening activity

respectively, in flies expressing the 1-7A variant with distinguished amplitude and/or phase. Blue

asterisks mark amplitudes that are significantly different between genotypes by ANOVA followed by a

Student's T-test ($p < 0.05$). Panels (D) and (E) display the phases of Morning Offsets and Evening Onsets

respectively. Gold asterisks: ANOVA followed by Dunnett's post hoc multiple comparisons of all

compared to WT: ns = not significant; * = $p<0.05$; ** = $p<0.01$; *** = $p<0.005$; **** = $p<0.001$. Panels (F)

– (H) display double plotted actograms of activity during the ~6 days of light:dark conditions, followed

by activity during ~9 days of DD. The green bars indicate the phase of the dominant activity period in

DD.

Figure 5. Locomotor Rhythms exhibited by WT versus *pdf* mutant flies expressing the 1-7A PDFR

Variant under Short-day (winter-like) conditions. The Panels present behavioral activity with a format

similar to that described in Figure Legend 2. Behavioral records were recorded from pdf^{01} (null) flies that

expressed either no UAS transgene (A, yellow box), or a UAS-1-7A *pdfr* transgene (B, black box), or from

control flies (*pdf⁺)* that expressed the UAS-1-7A *pdfr* (C, red box)). *TG*: transgene. Panels (A)-(C) display

average group eductions (activity averaged over days 5 and 6 of light entrainment, with open bars

indicating the 8-hr periods of Lights-on and filled bars the 16-hr periods of Lights-off) and bin-by-bin

analyses to compare activity amplitudes between genotypes. The red and blue arrows indicate Morning

and Evening activity of control (red) flies with distinguished amplitude and/or phase, respectively. Blue

asterisks mark amplitudes that are significantly different between genotypes by ANOVA followed by a

Student's T-test ($p < 0.05$). Panels (D) and (E) display the phases of Morning activity Onsets and Evening

activity Offsets respectively. Gold asterisks: ANOVA followed by Dunnett's post hoc multiple

comparisons of all compared to WT: ns = not significant; * = p<0.05; ** = p<0.01; *** = p<0.005; **** =

p<0.001. Panels (F) – (H) display double plotted actograms of activity during the ~6 days of light:dark

conditions, followed by activity during ~9 days of DD. The green bars indicate the phase of the

dominant activity period in DD.

Figure 6. Locomotor Rhythms exhibited by WT versus *pdf* mutant flies expressing the 1-7A PDFR

Variant under Long-day (summer-like) conditions. The Panels present behavioral activity with a format

similar to that described in Figure Legend 5. Behavioral records were recorded from pdf^{01} (null) flies that

expressed either no UAS transgene (A, yellow box), or a UAS-1-7A *pdfr* transgene (B, black box), or from

control flies (*pdf⁺)* that expressed the UAS-1-7A *pdfr* (C, red box)). *TG*: transgene. Panels (A)-(C) display

average group eductions (activity averaged over days 5 and 6 of light entrainment, with open bars

indicating the 16-hr periods of Lights-on and filled bars the 8-hr periods of Lights-off) and bin-by-bin

analyses to compare activity amplitudes between genotypes. The red and blue arrows indicate Morning

and Evening activity of control (red) flies with distinguished amplitude and/or phase, respectively. Blue

asterisks mark amplitudes that are significantly different between genotypes by ANOVA followed by a

Student's T-test (p < 0.05). Panels (D) and (E) display the phases of Morning activity Offsets and Evening

activity Onsets respectively. Gold asterisks: ANOVA followed by Dunnett's post hoc multiple

comparisons of all compared to WT: ns = not significant; * = p<0.05; ** = p<0.01; *** = p<0.005; **** =

p<0.001. Panels (F) – (H) display double plotted actograms of activity during the ~6 days of light:dark

conditions, followed by activity during ~9 days of DD. The green bars indicate the phase of the

dominant activity period in DD.

1006

Figure 7. A model predicting the effects of phosphorylating different sites on the PDFR CT on

downstream signaling. The CT is diagrammed as a grey box running horizontally. Asterisks indicate the

positions Ser/ Thr/ and Tyr residues as described in Fig. 1 and Suppl. Fig. 1: white asterisks are the

subset of Ser/Thr/ and Tyr residues that were targets of mutational analysis in this study. Mutation of

CL6 and CL7 consistently demonstrated the greatest delaying effects on the phases of Evening

locomotor peaks; they were also among the sites with greatest influence on the amplitudes of the

Morning and Evening peaks. The results suggest their phosphorylation normally will have the greatest

effect to slow or terminate PDFR signaling. Other sites are also effective, although to lesser degrees (as

indicated by their font sizes), including CL2-3, CL4 and CL5. Primarily, such phosphorylation will

decrease the duration of PDFR signaling (red bars) and so reduce (for example) the delay that PDF

imposes on the period of neuronal activation displayed by PDFR-responsive pacemaker groups (like the

Evening cells and the DN3 [10, 19]). The model also predicts that the effects of phosphorylating CL5 will

depend on the phosphorylation status of neighboring sites. In some contexts (e.g., PDFR 1-5A), it will

help terminate PDFR signaling, but in others (e.g., PDFR 567A) it may promote the duration or extent of

PDFR signaling, perhaps by blocking the effects of phosphorylating CL6 and CL7 (green bar).

1022

1023

1024

1025

1026

1027

1028

1029

1030

Figure 1.

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7