1

Regulation of PDF receptor signaling controlling

daily locomotor rhythms in *Drosophila*.

Weihua LI*, Jennifer Trigg* & Paul H. Taghert**

Department of Neuroscience

Washington University School of Medicine

St Louis, MO 63110

USA

* co-first authors

** for correspondence: taghertp@wustl.edu

Number of Figures: 8

Number of Tables: 1

Number of Supplemental Figures: 12

Number of Supplemental Tables: 5

Key Words:

Drosophila, PDF, PDF Receptor, Circadian Rhythms, G protein-coupled Receptor

Short title: PDF Receptor Desensitization

ABSTRACT

G protein-coupled receptors (GPCRs) trigger second messenger signaling cascades following

activation by cognate ligands. GPCR signaling ceases following receptor desensitization or uncoupling

from G proteins. Each day and in conjunction with ambient daylight conditions, neuropeptide PDF

regulates the phase and amplitude of locomotor activity rhythms in *Drosophila* through its receptor, a

Family B GPCR. Its time of action – when it starts signaling and when it stops – much change every day

to following changing day lengths. We studied the process by which PDF Receptor (PDFR) signaling

turns off *in vivo*, by modifying as many as half of the 28 potential sites of phosphorylation in its C

terminal tail. We report that many such sites are conserved evolutionarily, and that in general their

conversion to a non-phosphorylatable residue (alanine) creates a specific behavioral syndrome opposite

to loss of function phenotypes previously described for *pdfr*. Such "gain of function" *pdfr* phenotypes

include increases in the amplitudes of both Morning and Evening behavioral peaks as well as multi-hour

delays of their phases. Such effects were most clearly associated with a few specific serine residues, and

were seen following alanine-conversion of as few as one or two residues. The behavioral phenotypes

produced by these PDFR sequence variants are not a consequence of changes to the pharmacological

properties or of changes in their surface expression, as measured *in vitro*. We conclude that the

mechanisms underlying termination of PDFR signaling are complex and central to an understanding of

how this critical neuropeptide modulates daily rhythmic behavior.

AUTHOR SUMMARY

In multi-cellular organisms, circadian pacemakers create output as a series of phase markers across the 24

hr day to allow other cells to pattern diverse aspects of daily rhythmic physiology and behavior.   Within

circadian pacemaker circuits, neuropeptide signaling is essential to help promote coherent circadian

outputs. In the fruit fly *Drosophila* 150 neurons are dedicated circadian clocks and are heavily

synchronized – they all tell the same time. Yet they provide diverse phasic outputs in the form of their

discrete, asynchronous neuronal activity patterns. Neuropeptide signaling breaks that symmetry and

drives many pacemakers away from their preferred activity period in the morning. Each day,

neuropeptide PDF is released by Morning pacemakers and delays activity by specific other pacemakers to

later parts of the day.   When and how PDF stops acting each day is unknown. Further, timing of signal

termination is not fixed because day length changes each day, hence the modulatory delay exerted by

PDF must itself be regulated. Here we consider canonical models of G protein-coupled receptor

physiology to ask how PDF receptor signaling is normally de-activated. We use behavioral measures to

define sequence elements of the receptor whose post-translational modifications (e.g., phosphorylation)

regulate receptor signaling duration.

INTRODUCTION

In *Drosophila*, neuropeptide PDF signaling helps pattern the output of the fly circadian

pacemaker network in controlling rhythmic daily locomotor activity [1-3]. Its functions have been

compared to those of neuropeptide VIP in regulating mammalian circadian physiology [4]. Historically

PDF was first isolated as an active principle (Pigment Dispersing Hormone) that mediates light-

dependent dispersion of pigment granules in diverse chromatophores of crustacea [5, 6]. In the insect

circadian system, PDF acts for a specified period within each 24 hr cycle, and works in conjunction with

environmental light to set phase and amplitude for locomotor activity rhythms that normally occur

around dawn and dusk [7-9]. Each day, the precise time of dusk and of dawn change, as does the time

interval between them. These facts require that the time of PDF signaling, the point when it starts and

the point when it stops, must also be adjusted each day to appropriately follow and reflect these daily

variations in the light:dark transitions. PDF signaling starts following its release by specific pacemaker

neurons, whose period of activity *in vivo* tracks the dawn in a variety of photoperiodic conditions [10].

Here we address molecular mechanisms by which PDF receptor signaling normally stops.

The PDF receptor (PDFR) is a member of the Family B (secretin receptor-like) GPCR group [11-

13]: it is G_s-coupled and its activation elevates cAMP levels *in vivo* [14]. It regulates different adenylate

cyclases (AC) in diverse target pacemakers (15, 16, Duvall and Taghert, 2012; 2013), which, through PKA

activation, ultimately regulate the pace of the molecular clock through regulation of Timeless [17]. PDFR

autoreceptor signaling promotes dramatic, daily morphological changes in the axonal terminals of sLNv

pacemakers [18]. PDFR activation also regulates calcium dynamics in subsets of pacemaker neuron

groups to help dictate their group-specific, daily phases of activation (in the sLNv, in the 5th sLNv and in

subsets of LNd, and DN3 groups – [10]). Such target cell-specific delays of PER-dependent neuronal

activity illustrates the basis by which the circadian network produces a daily series of phasic, neuronal

outputs to temporally bias diverse downstream rhythmic events [19, 20]. Finally, PDF/PDFR signaling is

long-lasting: its depression of basal calcium levels in target neurons persists without abatement over

many hours. These observations raise fundamental questions regarding the mechanism and the time

course by which PDFR signaling diminishes in anticipation of the next day's cycle of signaling.

92
The canonical model of GPCR phosphorylation and homologous desensitization features G

protein-coupled receptor kinases (GRKs) which associate with activated GPCRs and phosphorylate

cytosolic segments, thereby recruiting β-arrestins [21, 22]. β-arrestins uncouple the receptors from G

proteins [23, 24]. or enhance receptor endocytosis [25]; they can also serve as signal transducers by

recruiting distinct signaling molecules [26]. A second major regulatory mechanism to reduce GPCR

signaling is heterologous desensitization, whereby second-messenger-dependent kinases (PKA or PKC)

phosphorylate GPCRs [27]. Thus, we designed experiments to modify evolutionarily-conserved residues

in the C terminal tail of PDFR that could conceivably serve as substrates for phosphorylation and

subsequent signal termination. Our working hypothesis is that such modifications will reveal an

extension of the lifetime of PDFR activation *in vivo*.

Following activation of rhodopsin in the mammalian retina, visual arrestin is recruited with a

time constant of < 80 ms [28]; for many *Drosophila* neuropeptide receptors, β-arrestin2 is recruited

within a minute of exposure to ligand [29]. Does phosphorylation/de-phopsphorylation also regulate

the activity of a GPCR like PDFR, that signals over a much longer time base? The mammalian blue-light

sensitive GPCR melanopsin (OPN4) mediates intrinsic light sensitivity in certain classes of retinal

ganglion cells (RGCs): melanopsin signaling is distinguished by long latencies, graded responses and

sustained RGC depolarization that can outlast the duration of the light stimulus [30]. Melanopsin-

expressing RGCs produce exceptionally long timecourse integration because that GPCR behaves

differently from other opsins in two fundamental ways. Firstly, it undergoes photoequilibration

between signaling and silent states: a property that maintains the availability of pigment molecules for

activation, termed 'tristability' by Emmanual and Do [31]. Secondly, phosphorylation and β-arrestin

recruitment do contribute to the kinetics of melanopsin signaling, but over a long time course[32, 33].

Mure *et al.* [33] proposed that the distal aspect of the melanopsin C terminal tail creates a steric

blockade covering a cluster of specific serine residues situated in more proximal regions. The time

required for relief of that blockade therefore dictates a delayed time course of phosphorylation.

We know very little about the mechanisms that underlie termination of PDFR signaling. The *pdfr*

gene displays a transcript rhythm that peaks in the late day [34]. Sensitivity to PDF *in vivo* peaks in the

early day and is regulated by the PER-dependent clock, via post-transcriptional mechanisms*:* the EC50

for PDF responses in identified neurons varies systematically 5-10 fold: as a consequence of Ral A action,

as a function of time of day, and as a function of seasonality [35]. PDFR signaling is long-lasting: it

persists for many hours *in vivo* [19], for as long as free peptide ligand is available in the bath [14]. In

addition, β-arrestin2-GFP is not efficiently recruited to activated PDFR when the receptor is functionally-

expressed in *hEK-293T* cells [35]. In contrast, each of 13 other *Drosophila* neuropeptide GPCRs (including

two Family B GPCRs, CG8422 and CG17415) efficiently recruit β-arrestin2-GFP when they are activated

by their cognate ligands in that cellular environment 29, 36, 37].

Here we report (i) that PDFR is normally phosphorylated *in vivo* at conserved C terminal

residues; and (ii) that loss of conserved PDFR phosphorylatable sites leads to a behavioral syndrome

opposite to loss-of-function *pdf* and *pdfr* phenotypes. This 'gain of function' approach reveals a multi-

hour range of potential phases for both the Morning and Evening activity peaks, within which

neuropeptide PDF:PDFR signaling normally specifies rhythmic behavior, according to season.

Importantly, it shows that PDFR signaling affects the amplitude of the activity peaks and not just their

phases. Finally, this work identifies specific PDFR sequence elements that are major points at which the

duration of receptor activity is dictated, and through which behavior is modulated in season-specific

fashion.

RESULTS

PDFR C-Terminal sequences. Based on alternative splicing, the *pdfr* locus in *Drosophila*

melanogaster (CG13758) encodes GPCRs with two different C terminal sequences, for which the PA and

PD isoforms are representative (flybase.org/reports/FBgn0260753).   The PD isoform is slightly longer

and lacks the final ~20 AAs of the PA isoform. As described in the Supplemental Information (*PDFR*

Isoforms), we focused on the PA protein isoform of PDFR, as it has been used in the majority of genetic

studies in the field. To identify residues for mutation, we first used comparative genomic analyses to

assess how well specific sequences in the C-terminal region of the PA receptor are conserved.   We

obtained annotated *pdfr* genomic sequences from 16 additional species of *Drosophila*, in both the

Sophophora and *Drosophila* sub-families (**Supplemental Table 1**). Together this species collection

represents an estimated 40-60 MYr of *Drosophalid* evolution [38]. We defined residue V505 of the

melanogaster protein as the start of the C terminal sequence, following the consensus 7th

transmembrane TM domain (TM7) (**S Figure 1**). These 17 different C terminal sequences all display

considerable length, and vary between 189 to 215 amino acids (AAs). The *melanogaster* PDFR-A C-

terminal contains 28 Ser, Thr, or Tyr residues (these may be subject to post-translational

phosphorylation and de-phosphorylation). To survey putative functionality among these, we chose 14

residues that are distributed across the length of the C-terminal and which display high evolutionary

conservation (**Supplemental Figure 1** and **Supplemental Table 2)**. For naming purposes, we grouped

them into arbitrary clusters (CL) numbered #1 to #7, with positions shown in **Figure 1.** Following a

common paradigm in study of GPCR physiology [e.g., 33, 39, 40], we tested the consequences of their

mutation to Alanine, a non-phosphorylatable analog [41]. Some clusters have only a single modified AA

(e.g. CL4 and CL5), while in others we modified two AAs (e.g., CL1, CL6 and CL7), and in others, as many

as six closely-positioned residues (e.g., CL2-3).

This series of variant receptors contains 10 different mutated versions of PDFR (not including

the WT 'parent' PDFR-A) and is arbitrarily divided into two broad categories as indicated in **Figure 1**. The

first group (termed "Simple Variants") targeted one or two clusters of conserved Ser/Thr/Tyr AAs (i.e.,

CL 2-3A; 4A; 5A; 6A; 7A). The second category (termed "Multiple Variants") targeted three or more of

the AA clusters in various combinations (i.e., CL 1-4A; 1-5A; 1-6A; 1-7A and 5-7A). The CL1-7A variant

mutates all 14 of the targeted residues. We studied these ten variants, as well as the WT receptor, in a

pdfr mutant strain (*han[5537]* – [11]) for their ability to rescue and to shape the phases and periodicity of

rhythmic locomotor behavior. We analyzed PDFR variants in the *pdfr* mutant background to permit

evaluation of their properties without competition from endogenous PDFR [although we acknowledge

that such possible interactions warrant future study]. Further we tested this PDFR series in different

photoperiodic conditions reflecting different seasons, as well as in constant darkness. *pdfr* loss-of-

function mutants display advanced behavioral peaks under Light:Dark conditions, as well as weak and

shortened free-running periods under constant darkness [11]. In general, we found that several of these

Ala-mutant PDFRs delayed the phase of the Morning and/or Evening activity peaks (**Figures 2-7**) and that

some also lengthened the free-running period (**Table 1**). Such results are generally consistent with a

prediction whereby non-phosphorylatable PDFR variants – those having a potential to increase the

duration of PDFR signaling - will produce behavioral actions opposite to those seen in loss-of-function

pdfr mutant flies. To better interpret the results produced by the different sequence variants, we also

present experiments that test assumptions used in the experimental design. These tests consider: i)

whether the C-terminal fusion of GFP (present in all variants tested) influenced the results; and ii)

whether the behavioral effects of receptor variants are independent of activation by the endogenous

ligand, neuropeptide PDF.

Analysis of Behavior with PDFR variant expression. Figures 2 -7 describe rhythmic locomotor

behavior of adult *pdfr* mutant flies expressing diverse *pdfr* transgenes in the aggregate, during six days

of light entrainment (LD), followed by activity during nine or more days of constant darkness (DD). For

each genotype, we display locomotor activity during light entrainment in four ways: (i) group actograms,

(ii) group eductions, (iii) derivations of behavioral phase, and (iv) direct comparisons of the amplitudes

of behavioral peaks between genotypes. Thus in Figures 2, 4 and 6, a representative group actogram

displays all locomotor activity double-plotted across all days in LD and DD (Panels A-L, sub-panel **-1** in

each Figure). Second a representative group eduction displays a 24 h average of the activity in the final

two days in LD (days 5-6) (Panels A-L, sub-panel **-2** in each Figure). Next, we define the phase onsets and

offsets for Morning and Evening activity peaks during the same final two LD days across all experiments

(Panels M and N in Figures 2, 4 and 6). For the comparison of peak amplitudes in these three

photoperiodic conditions, we describe Bin-by-Bin analyses of locomotor activity during the entire LD

period (days 2-6) in Figures 3, 5 and 7. The Bin-by-Bin analyses directly compares experimental

genotypes to a control one [cf. 42]. They present data averaged across 2 to 5 independent experiments,

to help identify the differences that most consistently correlate with genotype. Finally, Table 1 compiles

measures of rhythmic activity displayed by the different genotypes in days 3-9 of constant darkness.

Behavior under Short Day (winter-like) conditions (Figures 2 and 3). Flies lacking *pdfr* function

in 8L:16D lack a prominent morning peak and their evening peak of activity begins earlier than controls;

the example shown in **Figure 2A-2** is heterozygous for the *tim*-Gal4 element.   Rescue by WT-*pdfr* using

tim-Gal4 significantly delayed (i.e., rescued) the Evening peak initiation by about 1 h (**Fig. 2B-2).**

Expression of each of the Single Variants driven by *tim*-Gal4 produced also delayed the Evening peak

(**Fig. 2C-2** to **G-2**), with the variants 6A and 7A producing conspicuous, large amplitude peaks that

occurred on average as late as ~3 h after lights-off, at ZT12 (**2F-2** and **2G-2**).   Variants 2-3A, 6A and 7A

also produced elevated activity during the period of ZT 18-19 (**Fig. 3C-1,2F-1** and **2G-1**): we speculate

this is promotion of a "Morning" peak of activity.   Among the five Multiple Variants, we noted three

main effects (**Fig. 3**).   First the progressive addition of more Ala substitutions in the series 1-5A, 1-6A and

1-7A (**Figure 3D-5** through **3G-5**) produced large and delayed Evening peaks in the time period ZT11.5-

12.5 (3-4 h after lights-off), with the variant 1-6A producing the most pronounced delay.   The delayed

Evening peak activity at ZT12 appeared often appeared at the expense of, the normal Evening peak

activity that occurred prior to Lights-off (e.g., **Fig.3G-4**), although when averaged across all days in LD,

that effect was not significant (**Figure 3 – panel 2**). Second, the PDFR 1-5A and 1-7A forms in the

Multiple Variant series produced significant "Morning" peaks around ZT19, (**Fig 3J-1** and **3L-1**), Finally,

we noted that combining the Single 5A, 6A and 7A variants into a Multiple Variant (5-7A) did not

produce the anticipated additive effects on promoting morning locomotor activity at ~ZT19 (**Fig. 3C-3).**

Such anticipation was predicated on the effects seen with 1-5A, 1-6A and 1-7A (versus 1-4A) (**Fig 3J-1**

and-3, K-3 and L-1 and -3).   Likewise the 5-7A did not produce the strong delay in the Evening activity

peak to ~ZT12 that was expected based on effects seen with 1-5A, 1-6A and 1-7A (versus 1-4A): instead

it produced only a minor delay (**Fig. 3H-3**).

Behavior under equinox conditions (Figures 4 and 5).   Under 12:12 conditions, *han* mutant flies

(lacking *pdfr* function) typically display elevated nocturnal activity, a lack of morning anticipation prior to

Lights-on, and a pronounced advance in the peak of the Evening behavior [11]. **Fig. 4A-1** to **A-2** display

activity patterns of *han* mutants that are heterozygous for *tim*-Gal4; they generally matched prior

descriptions of *pdfr* mutant behavior. Restoration of some *pdfr* function by *han*; *tim>* WT-*pdfr*,

produced a modest Morning peak (anticipatory activity prior to lights-on – **Fig. 4B-2**) and delayed the

evening peak by 1-2 hrs (**Figures 4N and 5B-2**) [7,8,9]. Among the five "Simple PDFR Variants" (**Fig 4C-G**),

the 6A and 7A PDFR isoforms produced significantly elevated morning activity, with a peak ~2 hr prior to

Lights-on in the period ZT22-23 (**Fig 5F-1** and **G-1**). This elevated pre-Lights-on activity peaked ~3-4 hr

later than that seen under Short Day conditions with these same PDFR variants. Separately, the 6A

isoform increased Morning activity after the lights-on signal (**Fig. 5F-2)** and the 4A, 6A and 7A isoforms

increased the amount of Evening activity, but without changing its phase (**Fig. 5D-5, F- 5** and **G-5**). In the

case of the five Multiple Variants, we noted three main effects. First the progressive addition of more

Ala substitutions in the series 1-5A, 1-6A and 1-7A all produced significant additional Morning activity in

the period ZT22-23 compared to that seen with 1-4A. Notably, all of that activity appeared to decline

prior to the Lights-on signal (**Fig 5E-3, F-3, G-3**). Those variants also produced increased activity

immediately after Lights-on: an effect seemingly distinct from that produced prior to Lights-on (**Figure 5**

J-42 K-2 and **L-2**). Second, combining the 5A, 6A and 7A variants in PDFR 5-7A did not produce additive

effects on behavior in the morning, either before or after Lights-on (**Fig 5H-1** and **-2**): instead the activity

at ZT22-23 and ZT 0.5-3.5 appeared similar only to that displayed by rescue with a WT PDFR. Finally, the

1-5A variant was notable among Multiple Variants in LD12:12 conditions by displaying elevated activity

in mid-day between the Morning and Evening Activity peaks (**Fig 5E-4** and **E-5**).

Behavior under Long Day (summer) conditions (Figures 6 and 7). Flies lacking *pdfr* function in

16L:8D display locomotor patterns similar to those in 12L:12D – lack of an obvious Morning peak and

produce a broad Evening activity period that peaks ~3hr before Lights-off: the example in **Figure 6A-2**

displays behavior by *han* mutants that are also heterozygous for the *tim*-Gal4 element. Rescue (*han*;

tim> WT-*pdfr*, **Figure 6B-2**) typically delayed Morning activity offset (**Fig. 6M)** and delayed the Evening

peak onset each by about 1 h (**Fig. 6N).** Expression of the 5A and 6A PDFR variants produced

significantly greater amounts of Morning activity after Lights-on (**Fig. 7C-3** through **G-3**), while all of the

Single Variants increased the amplitude of the Evening peak without affecting its phase (**Fig. 6 C-5** to **G-**

5). Among the five Multiple Variants, we noted three main effects (all comparable to those seen

previously under 12L:12D entraining conditions). First the 1-6A and 1-7A variants displayed significantly

increased Morning activity after Lights-on (**Fig. 7 K-2** and **L-4**). Second, the variants 1-5A, 1-6A and 1-7A

increased the amplitude of the evening peak without affecting its phase (**Fig. 7J-3, K-3** and **G-3**). Third,

as in the other photoperiodic conditions, when we combining the 5A, 6A and 7A variants into a single 5-

7A variant, it did not produce the expected additive effects on behavior in the morning (**Fig 7H-2**) or in

the evening (**Figure 7H-3**): instead these activities appeared similar to effects displayed by the 1-4 and 1-

5 variants.

Behavior under DD. In DD following Short Day conditions, the PDFR variants that produced 3-4 h

delays in the evening peak often generated periods ~ 1-2 hr longer than the controls and also lowered %

arrhythmicity (**Table 1**). This was especially true for the Multiple Variant series (e.g., PDFR 1-5A, 1-6A

and 1-7A (**Table 1**). However the correlation between delayed evening peaks and a longer tau in DD was

not absolute" for example, the 6A, 7A and 5-7A variants all had strong significant delaying effects on the

Evening phase of activity under Short Day conditions (compared to the effect of the WT PDFR), but they

did not lengthen tau values in subsequent DD conditions (Table 1). These observations suggest that

delays of activity phases produced by PDFR modulation in LD are not necessarily a simple or direct

consequence of its effects on the PER and TIM--dependent clock [2, 3, 17]. In addition, the dominant

activity periods in DD typically reflected the delayed (~ZT12) Evening peak (**Fig. 2K-1** and **L-1**). Notably,

the early Morning peak (~ZT18-19) promoted by PDFR 6A and 7A clearly persisted in DD (**Fig. 2F-1** and

G-1).

Table 1.

Short Day Entrainment	N	n	%AR	tau	SEM	Average tau different from: w[1118]	pdfrEgfp	pdf[01]
han[5304]; tim > w[1118]	5	72	22%	23.9	0.09	-	ns	nd
han[5304]; tim > pdfr	1	16	6%	23.7	0.13	ns	ns	nd
han[5304]; tim > pdfr1-7A No GFP	3	30	8%	23.6	0.07	ns	ns	nd
han[5304]; tim > pdfrEgfp	3	46	35%	24.00	0.09	ns	-	nd
han[5304]; tim > pdfrEgfp23A	2	31	13%	24.8	0.12	ns	ns	nd
han[5304]; tim > pdfrEgfp4A	2	31	26%	24.7	0.13	ns	ns	nd
han[5304]; tim > pdfrEgfp5A	2	43	26%	24.58	0.11	ns	ns	nd
han[5304]; tim > pdfrEgfp6A	3	50	38%	24.37	0.08	ns	ns	nd
han[5304]; tim > pdfrEgfp7A	3	45	27%	24.83	0.11	ns	ns	nd
han[5304]; tim > pdfrEgfp1-4A	3	40	12%	24.6	0.12	ns	ns	nd
han[5304]; tim > pdfrEgfp1-5A	3	31	16%	25	0.23	***	*	nd
han[5304]; tim > pdfrEgfp1-6A	3	42	14%	25.1	0.13	****	**	nd
han[5304]; tim > pdfrEgfp1-7A	4	59	16%	24.9	0.11	****	*	nd
han[5304]; tim > pdfrEgfp567A	2	37	16%	23.9	0.09	ns	ns	nd
tim > pdfrEgfp1-7A	1	15	0%	25.77	0.25	nd	nd	****
pdf[01]	1	12	67%	22.50	0.12	nd	nd	-
tim > pdfrEgfp1-7A; pdf[01]/pdf[01]	1	25	20%	23.63	0.12	nd	nd	ns

Equinox Entrainment	N	n	%AR	tau	SEM		
han[5304]; tim > w[1118]	6	68	34%	23.7	0.09	-	ns
han[5304]; tim > pdfr	3	16	25%	23.5	0.16	ns	ns
han[5304];; tim > pdfr1-7A No GFP	3	35	14%	23.67	0.08	ns	ns
han[5304]; tim > pdfrEgfp	4	48	13%	24.1	0.06	ns	-
han[5304]; tim > pdfrEgfp23A	2	54	48%	25.6	0.12	****	****
han[5304]; tim > pdfrEgfp4A	2	44	32%	24.1	0.07	ns	ns
han[5304]; tim > pdfrEgfp5A	2	47	34%	24.4	0.06	ns	ns
han[5304]; tim > pdfrEgfp6A	3	35	23%	24.3	0.13	ns	ns
han[5304]; tim > pdfrEgfp7A	3	47	21%	24.1	0.09	ns	ns
han[5304]; tim > pdfrEgfp1-4A	3	45	18%	24.5	0.08	**	ns
han[5304]; tim > pdfrEgfp1-5A	2	32	6%	25.00	0.13	****	***
han[5304]; tim > pdfrEgfp1-6A	2	32	19%	25.27	0.14	****	****
han[5304]; tim > pdfrEgfp1-7A	6	88	6%	25.1	0.17	****	****
han[5304]; tim > pdfrEgfp567A	2	45	24%	24.1	0.08	ns	ns

Long Day Entrainment	N	n	%AR	tau	SEM			
han[5304]; tim > w[1118]	7	###	47%	23.8	0.06	-	ns	nd
han[5304]; tim > pdfr	3	47	17%	24.88	0.15	ns	ns	nd
han[5304]; tim > pdfr1-7A No GFP	3	40	48%	24.57	0.09	ns	ns	nd
han[5304]; tim > pdfrEgfp	4	82	17%	24.87	0.08	ns	-	nd
han[5304]; tim > pdfrEgfp23A	2	48	23%	25.35	0.12	****	****	nd
han[5304]; tim > pdfrEgfp4A	1	27	22%	25.50	0.16	****	****	nd
han[5304]; tim > pdfrEgfp5A	1	36	11%	24.66	0.16	ns	ns	nd
han[5304]; tim > pdfrEgfp6A	2	51	22%	25.02	0.09	ns	ns	nd
han[5304]; tim > pdfrEgfp7A	2	54	13%	25.46	0.09	****	****	nd
han[5304]; tim > pdfrEgfp1-4A	3	40	8%	25.2	0.13	****	ns	nd
han[5304]; tim > pdfrEgfp1-5A	3	33	6%	25.18	0.14	****	*	nd
han[5304]; tim > pdfrEgfp1-6A	3	48	2%	25.59	0.09	****	****	nd
han[5304]; tim > pdfrEgfp1-7A	4	63	14%	25.3	0.08	****	***	nd
han[5304]; tim > pdfrEgfp567A	1	38	18%	24.31	0.09	ns	ns	nd
tim > pdfrEgfp1-7A	1	14	14%	25.50	0.13	nd	nd	****
pdf[01]	1	32	88%	22.75	0.05	nd	nd	-
tim > pdfrEgfp1-7A; pdf[01]/pdf[01]	1	29	21%	23.30	0.09	nd	nd	ns

Table 1. Locomotor activity measures of various genotypes during constant dark conditions.

Behavioral activity records from days 3-9 in constant conditions following entrainment in the three

indicated photoperiods (Short Day, Equinox and Long Day). N = number of independent experiments

performed for each genotype::photoperiod combination. n = total number of flies tested for each

genotype::photoperiod combination.   % AR: the percentage of flies judged arrhythmic by criteria; tau =

average circadian periods calculated according to χ2-periodogram analysis. Statistical analysis

compared tau's using Tukey's multiple comparisons post hoc test following a one-way ANOVA: ns: not

significant; * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001.; nd: not determined.

In DD following Equinox or Long Day entrainment, we noted that several variants produced

dominant activity phases which were phase-delayed relative to the Evening peak displayed in LD (e.g.,

Fig.4F-1, G-1, and **J-, K-** and **L-1; Fig.6F-1, G-1,** and **J-, K-** and **L-1**). Thus, photophases > 8 hr in duration

appear to repress expression of the delayed, post Lights-off Evening activity observed under SD

conditions: this was suggested by the emergence of a delayed phase activity peak once flies only after

release to DD. Once again, several of the variants significantly increased Tau in DD (**Table 1**), yet had

only slight effects on phase in LD.

Control experiments. The design of the *PDF*R variants contains numerous assumptions and

genotypic constraints, distinct from simply substituting Alanine at potential sites of phosphorylation in

the receptor C terminal tail. To determine their potential to affect the results we report, we performed

the following two sets of experiments as controls. (i) - ***The influence of epitope fusions.*** The PDFR

variants we have tested include epitope fusions (4xFLAG and eGFP), which in principle could confer

some of the properties observed. In part, the concern regarding fused sequences is mitigated by the

fact that all UAS-receptor isoforms described above (WT and sequence-variant) contained both epitopes

(in addition – all were genetically transduced to the same integration site). Hence comparisons across

different variants (e.g., **Figures 2 through 7**), should largely normalize for the effects of the non-receptor

sequences, and instead highlight variant-specific properties. However, we wished to test this

assumption explicitly to define the extent to which a bulky GFP fused to the C Terminal tail might confer

some of the behavioral properties we might otherwise ascribe to PDFR C terminal sequences. We

therefore compared a 1-7A version of PDFR to a wild type PDFR, both of which lacked GFP fusions. The

results in both Short Day and Long Day conditions, followed by release into DD, support the hypothesis

that alterations in daily locomotor rhythmicity were largely derived from sequence variation of the

PDFR, and not from the properties of the fused GFP (**Supplemental Figure 4** and tabulated in **Table 1**).

Control experiments (ii) - dependence of GPCR PDFR variant effects on the presence of PDF ligand.

Second, we assumed that the altered behavioral phenotypes produced by certain PDFR variants

depended on activation by their endogenous cognate ligand, the neuropeptide PDF. However, we could

not *a priori* exclude the possibility that PDFR variants may in fact produce novel constitutive activity

(neomorphic properties). We therefore placed the 1-7A PDFR variant in the *pdf^{01}* (null) background and

re-tested its activity when driven by *tim*(UAS)-Gal4 under both short day and long day conditions, and

then in constant darkness. In all conditions, the behavioral phenotypes largely resembled those of the

pdf^{01} background (lack of a morning activity peak, advanced evening peak, and shorter tau/s in DD). The

results are shown in **Supplemental Figure 5** and tabulated in **Table 1**. The 1-7A PDFR variant did reduce

the % arrhythmicity present in the *pdf* mutant background to nearly the same value as found in a WT

pdf background, which suggests some degree of constitutive (ligand-independent) activity. Apart from

that one aspect, we conclude that constitutive activity explains at best a small proportion of the

behavioral effects of the PDFR variants here tested.

PDFR expression and signaling *in vitro*. The PDFR is a G_s-coupled receptor and several reports

have documented the importance of the downstream cAMP pathway to mediate PDF behavioral

regulation [15, 17, 43]. In that context, we asked whether the *in vivo* properties of those PDFR variants

that affected behavioral phase and amplitude could be correlated with *in vitro* properties when

expressed in *hEK-293T* calls. We found no strong correlation. Significant differences in basal signaling

levels, in EC50 or in maximum values of signal transduction were rare (**Supplemental Figures 6-8**).

When noted, they were poorly correlated with behavioral effects (**Supplemental Table 5**). We also

measured surface expression of the PFDR variants in *hEK* cells using β-lactamase N terminal fusions [44]

to determine if PDFR variants tended to display longer surface lifetimes. The 1-4 and 1-7 variants

displayed higher basal levels, but no others were different from the WT levels (**Supplemental Figure 9**).

Following 20 min exposure to PDF, neither the WT not any of the variants displayed a change in surface

expression levels (**Supplemental Figure 10**).

Measuring the phosphorylation state of over-expressed PDFR *in vivo*. To obtain direct

evidence that PDFR sequences are phosphorylated *in vivo*, we first over-expressed an epitope-tagged-

PDFR WT construct using *tim*-Gal4, which directs expression broadly in cells that feature the PER-

dependent molecular oscillator. We immunoprecipitated the receptor from head extracts, then

employed tandem mass spectroscopy to determine which if any specific PDFR peptide fragments are

phosphorylated. We performed eight biological replicates, with two collections in the morning (ZT2-3)

and six in the evening hours (ZT11). **Table S3** reports the phosphorylated peptides detected from the

PDFR-GFP fusion protein.   Among the 28 conserved Ser/Thr and Tyr residues in the PDFR C terminal tail,

five were phosphorylated in one or more of these samples. In the two Mornings samples S563 was

phosphorylated in one sample; in the Evening samples, S531 (in CL2) was phosphorylated in four of

samples, S534 (in CL2) in one, T543 (in CL3) in one, and S560 in two of six samples. The spectra

documenting detection of these phosphopeptides are presented in **Supplemental Figure 2**.

Testing GRKs and β-arrestin2 contributions to locomotor rhythmicity. Regarding which

potential PDFR phosphorylation sites might be β-arrestin binding sites, we considered a recent report by

Zhou *et al.* [45] who proposed a conserved GPCR sequence motif that promotes high affinity

interactions with β-arrestin. The motif includes three phosphorylation sites that align with three

conserved, positively-charged pockets in the arrestin N-domain. In *D. melanogaster* PDFR, the CL1

region contains a match with the proposed motif beginning with S512 (SLATQLS) and shows moderate

sequence conservation: of the 17 species we considered, 9 others retained this precise motif (**S Figure**

1). In addition, the *D. melanogaster* PDFR sequence beginning with S629 (SRTRGS) also displays a match

for the motif, but is retained in only 6 of the other 16 species. The evidence that these two sites

represent high affinity β-arrestin2 binding domains is therefore equivocal. We previously reported that

β-arrestin2-GFP is not efficiently recruited to activated PDFR when functionally-expressed in *hEK-293T*

cells [35]. In contrast, each of 13 other *Drosophila* neuropeptide GPCRs do efficiently recruit β-

arrestin2-GFP when expressed and activated in that cellular environment [29, 36, 37].

We tested the potential to detect *in vivo* involvement by *Drosophila* orthologues of the

canonical desensitization effectors - mammalian G-protein coupled receptor kinases (GRKs – GPRK1

(CG40129) and GPRK2 (CG17998)) and of mammalian β-arrestin2 (βarr2 – *kurtz* (CG1487)) - in the

control of locomotor rhythms We drove specific over-expression of WT cDNAs and RNAi constructs

using *tim*-Gal4, to broadly affect signaling in the circadian pacemaker system. The PDFR-PA protein is

highly restricted in its expression to subsets of the pacemaker neural network [46]. We predicted that

elimination of a desensitizing component for PDFR signaling should produce behavioral phenotypes

opposite to that of *pdfr* loss of function phenotypes: these would potentially include (for example) a

delayed evening activity phase in LD conditions and a longer tau under DD. GPRK1- and *krz*-specific

RNAi's produced normal average locomotor profiles under 12:12, while one of two GPRK2-specific RNAi

constructs tested slightly advanced the evening peak (**Supplemental Figure 5**). Over-expressing GPRK1

cDNAs did not affect morning or evening phase; over-expressing GPRK2 broadened the evening peak.

Under DD, *krz* RNAi flies were uniformly arrhythmic, while GPRK RNAi's were normal or slightly

lengthened the circadian period (**Supplemental Table 10**).   These results do not support the hypothesis

and suggest that the kinetics by which PDFR signaling terminates do not depend exclusively on the

activities of either dGPRK-1 or -2, or on that of the *Drosophila* β-arrestin2 ortholog, *krz*.

DISCUSSION

We found that several PDF receptor variants, containing Ala-substitutions of phosphorylatable

residues, modulated the amplitudes of the Morning and/or or Evening activity peaks, or delayed their

phases and/ or lengthened the free-running period. Such results are generally consistent with a

prediction whereby non-phosphorylatable PDFR variants – those with a potential to increase the

duration of PDFR signaling - would produce behavioral actions opposite to those seen in loss-of-function

pdf [1], or *pdfr* [11] mutant stocks.   The loss of function phenotypes include advanced behavioral peaks,

as well as weaker and shortened free-running periods. In this context, we acknowledge that GPCR

modification by phosphorylation does not exclusively lead to signal termination. GRK2-dependent

phosphorylation of the smoothened (smo) GPCR (which helps mediate Hh signaling) follows reception of

the Hh signal: that leads to smo <u>activation</u> in both *Drosophila* and mammalian systems [47]. Thus

modification of putative phosphorylation sites on GPCRs (to preclude phosphorylation) will not

exclusively promote extended signaling and therefore our interpretations must correspondingly

consider outcomes without *a priori* assumption of mechanisms.

GPCR signal termination. PDFR belongs to the Secretin Receptor Family (Family B) of

neuropeptide receptors [11-13]: there is no clear consensus regarding mechanisms of desensitization

and internalization for this receptor family. For VPAC2 receptors, phosphorylation and internalization is

mediated exclusively by GRK2 (48. Murthy, 2008;). Likewise, *Drosophila* orthologues of Cortocotrophin

Releasing Factor receptors (CG8422, DH44-R1) and Calcitonin receptors (CG17415, DH31-R1) are

internalized in *hEK* cells following recruitment of β-arrestin2 [29]. In contrast, PDFR, which is also

related to the mammalian Calcitonin receptor, is not internalized following exposure to PDF [35]. VPAC2

receptors are also regulated by heterologous receptor signaling: M3 cholinergic receptors via PKC

signaling can block VPAC2 phosphorylation, desensitization and internalization [49]. Furthermore,

secretin receptors and VPAC1 receptors undergo phosphorylation by GRKs and β-arrestin2-dependent

desensitization, but these are not sufficient to facilitate or mediate internalization [50-52]. Finally, GLP2-

Receptor associates with β-arrestin2 via its distal C terminal sequences, but that receptor domain is

required neither for GLP2-R desensitization nor its internalization [53]. Thus kinases other than GRKs

and effectors other than β-arrestin2 may regulate internalization of diverse Family B receptors.

Validating the behavioral phenotypes produced by PDFR variants. The actions of PDFR

variants we have described, following substitutions of Ala for various Ser/Thr or Tyr residues, are not

easily explained by a hypothesis invoking neomorphic or constitutive GPCR properties.   The evidence for

this conclusion is three-fold. First, the effects on both amplitude and phase of activity peaks by these

variants are not of a random assortment: rather they are all strictly opposite to that of loss-of-function

models for *pdf* and *pdfr*.   Second, the actions of PDFR variants display strong dependence on wild type

pdf gene function: this strongly argues that the actions of PDFR variants reflect responses to normal,

endogenous PDF signaling. Third, the action of PDFR variants on the phases and amplitudes of

locomotor activity reflected PDFR sequence variation, and not the properties of the GFP C-terminal

fusion. Together these control observations support the hypothesis that phosphorylation of some or all

of the C terminal residues we studied are normally modified to attenuate the strength and duration of

PDFR signaling.

PDF Signaling and seasonal adaptation. PDF neuropeptide actions were discovered in the

context of physiological adaptions to daylight in crustacea [5,6]. Helfrich-Förster [54] discovered PDF

expression within a defined subset of the insect *Drosophila* circadian pacemaker network. Renn *et al.*

[1] demonstrated that *pdf* makes a fundamental contribution by setting the normal behavioral phase of

Evening activity in *Drosophila* during light:dark entrainment and promoting normal rhythmicity during

constant darkness. PDF signaling coordinates with that driven by light: It works in parallel to direct

photosensitivity along with the CRY blue light photoreceptor. Flies doubly mutant for *cry* and *pdf* display

severe deficits of locomotor rhythmicity [7, 8], and such effects can be as severe as measured in clock-

deficient flies [9]. PDF and environmental light also work coordinately at the level of neuronal activity

patterns: in the case of the evening pacemakers (LNd and the 5[th] s-LNv), both light and PDF signaling

promote a delay in their PER-dependent activation period, and help align it to a phase just prior to dusk

[19].

Based on the close association of PDF signaling and photoperiodic signaling, we considered it

essential to test the different variant PDF Receptors in a variety of photoperiodic conditions. The

consensus of many prior observations indicates that PDF neuropeptide signaling is necessary for the

display of a Morning peak (its amplitude) and for proper adjustment of phase of the Evening activity

peak [e.g., 1, 7, 8, 55-57]. Our present observations, concerning likely gain-of-function *pdfr* variants,

support a revised hypothesis regarding regulation of the Morning peak: we propose PDF/PDFR can

affect - and normally does affect - both the phase and amplitude of the both Morning and Evening

activity peaks. Thus the consensus of many prior descriptions – concluding there is "no Morning activity

peak" in *Pdf* and *Pdfr* mutants – beginning with the description by Renn *et al.* [1], may have failed to

discern the presence of a greatly-diminished Morning peak that was also advanced. There is some

precedent for this contrary conclusion: Yoshii *et al.* [58] described behavior of *pdf*^{01} mutants and noted

advanced morning peaks especially under longer photoperiods. Likewise, the duration of the

photoperiod strongly influenced the effects we observed on the Morning versus the Evening activity

peaks. We only observed delayed Evening activity peak, produced by PDFR gain-of-function variants,

under Short Day conditions: the delayed evening peak appears strongly inhibited by light durations > 8

hr: and it remains unclear at what molecular level light exerts this inhibitory effect. We note a striking

similarity of these observations with effects recently described in loss-of-function states for the

phosphatase *PRL-1* [51].   Like the most active PDFR variants we herein describe (e.g., 6A, 7A, 1-5A 1-6A

and 1-7A), *PRL-1* mutants produce a 3-4 h delay in the phase of the evening activity peak, but only under

winter-like (short day) conditions, not under equinox or summer-like (long day) conditions. Kula-

Eversole *et al.* [59] have shown that TIM phosphorylation is affected by PRL-1 activity and suggest the

seasonal action of PRL-1 to advance the evening locomotor activity phase is mediated by modification of

TIM levels within the transcription-translation feedback loop. The extensive similarity of these two sets

of behavioral phenotypes reveals either serial or parallel pathways to effect comparable outcomes on

locomotor activity. If serial, then according to the simplest model, PRL1 acts downstream of PDFR, and

the combined results to-date suggest PRL-1 is inhibited by PDFR activation. Further genetic and

biochemical experiments are needed to evaluate if and how these pathways converge.

PDFR signaling effects on phase and on period. The behavioral effects of the PDFR variants

tested were most robust in modulating the phase of evening activity bouts. Evening locomotor activity

derives in large part from the activity of the primary Evening oscillators, termed E cells, the LNd and the

5[th] small LNv [10, 20, 51, 60-63]. However, numerous network interactions are also known to provide

critical contributions to the accuracy, precision and adaptibility of that timing system [e.g., 64 -67]. The

Evening oscillator group itself is known to be heterogeneous and to constitute at least three separate,

functionally distinct oscillators [68], only some of which express PDFR and respond to PDF [57, 68]. Vaze

and Helfrich-Förster [57] reported that the phase of Period protein accumulation in E neurons is phase-

locked to the previous lights-off transition and also sensitive to PDF signaling. They conclude that the

peak of PER accumulation is key to determining the phase of the evening activity peak, but also note

that the correlation between the Period-clock timing cue and the Evening activity peak is not perfect.

They suggest the Evening behavioral phase is better described by also factoring in the delay in E cell

neuronal activity driven by PDF neuromodulation [10, 19]. In that context, we interpret the many

examples of delayed Evening peaks resulting from PDFR variants (e.g., Figure 2M) as products at least

partly derived from PDF modulation (delay) of the cell intrinsic molecular clocks in E cells: that

conclusion is supported by the larger tau values in DD evident in the behavior of many PDFR variants

(Table 1). However, among the different variants, the association between delayed Evening activity

peaks in LD and longer tau values in DD was not perfect. For example, under short days, the 5A, 6A, 7A

and 5-7A variants all significantly delayed the offsets of Evening activity peaks versus that seen with WT

PDFR, but these variants did not significantly lengthen circadian period. Likewise under long day, 4A and

7A produced significant increases in tau versus that seen with WT PDFR, but they did not significantly

delay the onset of the Evening activity peak. Our working hypothesis is therefore that the duration of

PDF>PDFR signaling to target pacemakers in the *Drosophila* brain is modulated each day to

independently effect (i) entrainment of clock period, and (ii) delays of pacemaker neuronal activity. It is

the combination of these two PDFR signaling effects that is essential for proper rhythmic behavioral

outcomes across seasons. The signaling pathways downstream of PDFR, that independently regulate

diurnal phase and circadian period within E pacemaker neurons, are largely unknown.

Relations between in vitro and in vivo PDFR signaling measures. Using an *in vitro* assay for

cyclic AMP generation, we found that modifying phosphorylation properties of PDFR does not affect the

strength of signaling. That conclusion suggests that *in vivo* other GPCR features are normally affected to

regulate the extent/duration of PDFR signaling according to season, which corresponds to findings

described with other Family B GPCRs like VPAC1 [40]. Mutation of all the Ser and Thr residues of the C

terminal tail and of Ser250 to Ala led to a receptor with binding properties and adenylate cyclase activity

not different from that of the wild type receptor; however that variant receptor was neither

phosphorylated nor internalized. We propose that the PDFR variants we have described do not modify

locomotor behavior by virtue of greater second messenger signaling. Rather they do so by generating

cAMP (and perhaps other second messengers) over time periods longer than normal sustained by a WT

receptor over a portion of a 24-hr cycle.

Specific versus non-specific phosphorylation regulating GPCR activity. There are two broadly

divergent hypotheses to describe the mechanisms by which GPCR phosphorylation promotes

desensitization. The first posits that modification of specific residues have greatest significance for

downstream desensitizing mechanisms. The second invokes the triggering of termination processes by

the aggregate negative charge accumulated with phosphorylation, regardless of where it might occur.

For some GPCRs (e.g., OPN4), the evidence supports both models [33, 39]. Our data concerning PDFR

desensitization also support both viewpoints and we present a model in **Figure 8**. The evidence for the

importance of PDFR sequence specificity rests on the data that modification of certain single Clusters is,

among all variants tested, the most effective in affecting locomotor activity rhythms. In particular we

regard CL6 and CL7 to be especially critical in terminating the time course of PDFR signaling. Notably,

converting just the single pair of CL6 AAs, or the single pair of CL7 AAs, is enough to generate hours-long

delays in the peaks of locomotor rhythms under Light:Dark conditions (**Figure 2 F-G, 4F-G; 6 F-G, 7 F-G**),

as well as generating a significant increase in tau under Constant Dark conditions (**Table 1**). We note

that the splicing event that distinguishes the PDFR-A and PDFR-D isoforms occurs just prior to the

position of sequences encoding CL6 and 7, such that the D form lacks these two highly conserved

domains. Other clusters (like CL2-3) also appear to have potential for 'specific' contributions to PDFR

regulation. Together, these results speak to the potency of the inferred specific termination

mechanisms for PDFR GPCR signaling. In contrast, the evidence for the bulk phosphorylation hypothesis

comes from comparison of the 1-4A, versus the 1-5A, 1-6A and 1-7A variants. By inactivating increased

numbers of phosphorylatable residues, we observed a graded series increasingly delayed evening

phases in short day conditions (e.g., **Figure 2I-2 though 2L-2**).   The tandem mass-spectroscopy results,

indicating endogenous phosphorylation of certain PDFR residues (**Supplemental Table 3**) demonstrates

such post-translational modifications can occur, but likely does not provide a complete accounting of

them.

Finally, our results point to what we propose as "context-dependent" effects of modifying

specific GPCR residues – ones that may produce opposite behavioral effects, depending on the

phosphorylation status of neighboring sites. In particular we point to the contradictory results we

observed with Ala-variants of the CL5 site.   In comparing results from the series 1-4A, 1-5A, 1-6A and 1-

7A, we found that the 1-4A had only mild effects on the evening peak compared to over-expressing WT

PDFR (e.g., Figure **2B-2** vs. **2I-2**), whereas the others produced significant delays (**Figures 2M and 3I-3**

through 3L-3) under Short Day conditions. Those observations suggest modification of the CL5 residue

(S6333) promotes desensitization of PDFR and termination of PDFR signaling. However, the PDFR

variant 5-7A was constructed with the expectation that it would be as effective as 1-5A, 1-6A or 1-7A in

delaying the evening phase. Instead it proved only weakly effective. This same mis-match of expected

behavioral effects for 5-7A was also seen in other photoperiodic conditions (**Figure 6N, and 7H-3 vs 7J-3**

through 7L-3). Likewise, Langlet *et al*. [40] reported that effects on mutating Serine residues in the

carboxy terminus of VPAC1 did not produce additive effects. We propose that phosphorylation of CL5

will have either positive or negative consequences on PDFR desensitization depending on which other

neighboring residues are also modified. Thus we speculate that CL5 takes on outsize importance in

determining desensitization rates for PDFR and so may itself be subject to exceptional regulation. Such

complex interactions between phosphorylation sites is reminiscent of interactions documented between

diverse phosphorylation sites in the circadian clock protein PERIOD [69, 70].   Better resolution of these

apparent conflicting mechanisms (specific phosphorylation versus bulk negative charge, and paradoxical

effects of the CL5 site) awaits precise molecular definition of where and when PDFR is modified *in vivo*,

and by which modifying pathways.

MATERIALS and METHODS

Fly Rearing and Stocks. *Drosophila* were raised on a cornmeal agar diet supplemented with

yeast at 25°C in 12 hr:12 hr LD cycles. The UAS-*pdfr* mutant transgenic series was created by injecting

yw P{nos phiC31\int.NLS}X;P{CaryP}attP40 embryos (Rainbow Transgenic Flies, Inc, Camarillo, CA). The

UAS-*pdfr*-tandem construct was injected into *y[1] w[*] P{y[+t7.7]=nos-phiC31\int.NLS}X;*

P{y[+t7.7]=CaryIP}su(Hw)attP6 embryos. For PDFR-tandem fusion protein expression, we made a stable

yw; *tim*(UAS)-Gal4; UAS-*pdfr*-tandem stock. *pdfr* mutant flies are described in [11]; *tim*(UAS)-Gal4 flies

are described in [71] – *BL80941*). The recombinant fly stock - *han^5304*; *tim* (UAS)-Gal4 - was confirmed by

PCR and sequencing.

hEK-293 Cell Culture. *hEK-293* cells were maintained in DMEM, 10% FBS and 100U/mL penicillin

and streptomycin in 5% CO_2 atmosphere at 37^{o}C. For all transient transfections, 1.5×10^6 cells were used

to inoculate T25 flasks, incubated overnight, then transfected with 10ug plasmid DNA and 20 uL

lipofectamine 2000 reagent (Invitrogen Life Technologies). Five hours after transfection, cells were split

4×10^4 cells/well into a 96-well assay plate. We created a series of stably-transfected *hEK-293* cells

expressing WT and sequence-variant PDFRs, using the Flp-In System (Invitrogen Life Technologies,

Waltham, Massachusetts) per manufacturers recommendations, and maintained them in DMEM

supplemented with 150 µg/ml hygromycin B.

cAMP Assays. We measured PDF Receptor signaling activity using a CRE-*Luciferase* reporter

gene, following methods described by Johnson *et al.* [29]. The reporter gene construct was transiently

transfected to each stable cell line and luminescence measured using Firefly Luciferase Assay Kit

(Biotium, Inc., Fremont, California) and a Wallac 1420 VICTOR2 microplate reader (PerkinElmer, Inc.,

Waltham, Massachusetts). Concentration-effect curves, EC_{50}, top values and p-values were calculated

using the dose response, in a nonlinear regression using GraphPad Prism 8.0 software (San Diego,

California).

Locomotor activity Measures. All locomotor activity experiments were conducted with 2-5

days-old male flies at 25°C using Trikinetics Activity Monitors as previously described [9]. We crossed

Gal4 lines to *w^1118* to create control progeny. Locomotor activities were monitored for 6 days under

different photoperiodic conditions, and then for 9 days under constant dark (DD). To analyze rhythmicity

under constant conditions, we normalized the activity of flies from DD day 3 to day 9 and used χ2-

periodogram analysis with a 95% confidence cut-off, as well as SNR analysis [72] Arrhythmic flies were

defined by a power value $\leq$ 10 and width value $\leq$ 2, and period outside the range, 18 to 30 hours. To

analyze periods, we used Graphpad Instat (v 8) software to run one-way ANOVA measures followed by

the Tukey-Kramer Multiple Comparisons Test. We used Clocklab (Actimetrics) software to produce

actograms and the Brandeis Rhythms Package [73] to produce average activity plots (group eductions).

To analyze onset and offset phases of Morning and Evening activity bouts on the final two days of

light:dark entrainment (LD 5&6), we followed the method of Kula-Eversole *et al.* (55, 2021): $[(A_{n+2} + A_{n+1})$

$- (A_{n-1} + A_{n-2}) = \Delta Activity]$. Experimental genotypes were tested in the *han* mutant background*, pdfr^{5304}*

[11]*,* or in the *pdf^{01}* (null) mutant background [1], or in the *w^{1118}* background, as noted.

Additional details on Methods and procedures are found in **Supplemental Information**.

ACKNOWLEDGEMENTS

We thank many colleagues for generously sharing reagents, fly strains and providing technical advice, all

of which helped us pursue these studies. These include Drs. Xuemei Si, Ali Salahpour, Dmitri Nusinow,

Johanna Chiu, Xitong Liang, Paul Hardin, Zsolt Lenkei, Ken Blumer and David Wheeler. We thank the

staff that manages the Flybase resource which provides valuable information, and the Danforth Plant

Science Center Mass Spectroscopy Facility for expert assistance, and the Bloomington *Drosophila* Stock

Center, and the Vienna *Drosophila* Research Center for providing many valuable fly stocks. We also

thank Joel Levine and Orie Shafer for kindly providing comments on the manuscript.

AUTHOR CONTRIBUTIONS

W.L., J.T. and P.H.T designed experiments. W.L. and J.T. performed experiments. W.L., J.T. and P.H.T

analyzed experiments. P.H.T wrote the manuscript.

LITERATURE CITED

1. Renn SC, Park JH, Rosbash M, Hall JC, Taghert PH. A *pdf* neuropeptide gene mutation and ablation of

PDF neurons each cause severe abnormalities of behavioral circadian rhythms in *Drosophila*.

Cell. 1999; 99:791-802.   PMID: 10619432.

2. Peng Y, Stoleru D, Levine JD, Hall JC, Rosbash M. *Drosophila* free-running rhythms require intercellular

communication. PLoS Biol. 2003; 1:E13. PMID: 12975658.

3. Lin Y, Stormo GD, Taghert PH. The neuropeptide pigment-dispersing factor coordinates pacemaker

interactions in the *Drosophila* circadian system. J Neurosci. 2004; 24:7951-7. PMID: 15356209.

4. Vosko AM, Schroeder A, Loh DH, Colwell CS. Vasoactive intestinal peptide and the mammalian

circadian system. Gen Comp Endocrinol. 2007; 152:165-75. PMID: 17572414.

5. Fernlund, P. Structure of a light-adapting hormone from the shrimp, *Pandalus borealis.* Biochim

Biophysica Acta 1976; 439: 17-25. PMID: 95295.

6. Rao KR, Riehm JP, Zahnow CA, Kleinholz LH, Tarr GE, Johnson L, Norton S, Landau M, Semmes OJ,

Sattelberg RM, Jorenby WH, Hintz MF. Characterization of a pigment-dispersing hormone in

eyestalks of the fiddler crab *Uca pugilator*. Proc Nat'l Acad Sci (Wash) U S A. 1985; 82:5319-22.

PMID: 16593589.

7. Cusumano P, Klarsfeld A, Chélot E, Picot M, Richier B, Rouyer F. PDF-modulated visual inputs and

cryptochrome define diurnal behavior in *Drosophila*. Nat Neurosci. 2009; 12:1431-7. PMID:

19820704.

8. Zhang L, Lear BC, Seluzicki A, Allada R. The CRYPTOCHROME photoreceptor gates PDF neuropeptide

signaling to set circadian network hierarchy in *Drosophila*. Curr Biol. 2009; 19:2050-5. PMID:

19913424.

9. Im SH, Li W, Taghert PH. PDFR and CRY signaling converge in a subset of clock neurons to modulate

the amplitude and phase of circadian behavior in *Drosophila*. PLoS One. 2011; 6:e18974. PMID:

21559487.

10. Liang X, Holy TE, Taghert PH. Synchronous *Drosophila* circadian pacemakers display nonsynchronous

Ca^{2+} rhythms. Science (Wash). 2016; 351:976-81. PMID: 26917772.

11. Hyun S, Lee Y, Hong ST, Bang S, Paik D, Kang J, Shin J, Lee J, Jeon K, Hwang S, Bae E, Kim J. *Drosophila*

GPCR Han is a receptor for the circadian clock neuropeptide PDF. Neuron. 2005; 48:267-78.

PMID: 16242407.

12. Lear BC, Merrill CE, Lin JM, Schroeder A, Zhang L, Allada R. A G protein-coupled receptor, groom-of-

PDF, is required for PDF neuron action in circadian behavior. Neuron. 2005; 48:221-7. PMID:

16242403.

13. Mertens I, Vandingenen A, Johnson EC, Shafer OT, Li W, Trigg JS, De Loof A, Schoofs L, Taghert PH.

PDF receptor signaling in *Drosophila* contributes to both circadian and geotactic behaviors.

Neuron. 2005; 48:213-9. PMID: 16242402.

14. Shafer OT, Kim DJ, Dunbar-Yaffe R, Nikolaev VO, Lohse MJ, Taghert PH. Widespread receptivity to

neuropeptide PDF throughout the neuronal circadian clock network of *Drosophila* revealed by

real-time cyclic AMP imaging. Neuron. 2008; 58:223-37. PMID: 18439407.

15. Duvall LB, Taghert PH. The circadian neuropeptide PDF signals preferentially through a specific

adenylate cyclase isoform AC3 in M pacemakers of *Drosophila*. PLoS Biol. 2012; 10:e1001337.

PMID: 22679392.

16. Duvall LB, Taghert PH. E and M circadian pacemaker neurons use different PDF receptor

signalosome components in *Drosophila*. J Biol Rhythms. 2013; 28:239-48. PMID: 23929551.

17. Seluzicki A, Flourakis M, Kula-Eversole E, Zhang L, Kilman V, Allada R. Dual PDF signaling pathways

reset clocks via TIMELESS and acutely excite target neurons to control circadian behavior. PLoS

Biol. 2014; 12(3):e1001810.. PMID: 24643294.

18. Depetris-Chauvin A, Fernández-Gamba A, Gorostiza EA, Herrero A, Castaño EM, Ceriani MF. Mmp1

processing of the PDF neuropeptide regulates circadian structural plasticity of pacemaker

neurons. *PLoS Genet*. 2014; doi:10.1371/journal.pgen.1004700. PMID: 25356918.

19. Liang X, Holy TE, Taghert PH. A Series of Suppressive Signals within the *Drosophila* Circadian Neural

Circuit Generates Sequential Daily Outputs. Neuron. 2017; 94:1173-1189. PMID: 28552314.

20. Liang X, Ho MCW, Zhang Y, Li Y, Wu MN, Holy TE, Taghert PH. Morning and Evening Circadian

Pacemakers Independently Drive Premotor Centers via a Specific Dopamine Relay. Neuron.

2019; 102:843-857. PMID: 30981533.

21. Pincas H, González-Maeso J, Ruf-Zamojski F, Sealfon SC. G Protein-Coupled Receptors. In:

Belfiore A, LeRoith D, editors. Principles of Endocrinology and Hormone Action.

Endocrinology. Springer, Cham. https://doi.org/10.1007/978-3-319-44675-2_6. 2018.

22. Lefkowitz RJ, Shenoy SK. Transduction of receptor signals by β-Arrestins. Science (Wash). 2005;

308:512-7. PMID: 15845844.

23. Benovic JL, Kühn H, Weyand I, Codina J, Caron MG, Lefkowitz RJ. Functional desensitization of the

isolated beta-adrenergic receptor by the beta-adrenergic receptor kinase: potential role of an

analog of the retinal protein arrestin (48-kDa protein). Proc Nat'l Acad Sci (Wash) U S A. 1987;

84:8879-82. PMID: 2827157.

24. Lohse MJ, Benovic JL, Codina J, Caron MG, Lefkowitz RJ. β-Arrestin: a protein that regulates beta-

adrenergic receptor function. Science (Wash). 1990; 248:1547-50. PMID: 2163110.

25. Ferguson SS, Downey WE 3rd, Colapietro AM, Barak LS, Ménard L, Caron MG. Role of β-arrestin in

696   mediating agonist-promoted G protein-coupled receptor internalization. Science (Wash). 1996;

697   271:363-6. PMID: 8553074.

698   26. Smith JS, Rajagopal S. The β-Arrestins: Multifunctional Regulators of G Protein-coupled Receptors. J

699   Biol Chem. 2016; 291:8969-77. PMID: 26984408.

700   27. Kelly E, Bailey CP, Henderson G. Agonist-selective mechanisms of GPCR desensitization. Brit J of

701   Pharm. 2008; 153 Suppl 1: S379–88. PMID 18059321.

702   28. Krispel CM, Chen D, Melling N, Chen YJ, Martemyanov KA, Quillinan N, Arshavsky VY, Wensel TG,

703   Chen CK, Burns ME. RGS expression rate-limits recovery of rod photoresponses. Neuron. 2006;

704   51:409-16. PMID: 16908407.

705   29. Johnson EC, Bohn LM, Barak LS, Birse RT, Nässel DR, Caron MG, Taghert PH. Identification of

706   *Drosophila* neuropeptide receptors by G protein-coupled receptors-beta-arrestin2 interactions. J

707   Biol Chem. 2003; 278(52):52172-8. PMID: 14555656.

708   30. Berson DM, Dunn FA, Takao M. Phototransduction by retinal ganglion cells that set the circadian

709   clock. Science (Wash). 2002; 295:1070-3. PMID: 11834835.

710   31. Emanuel AJ, Do MT. Melanopsin tristability for sustained and broadband phototransduction.

711   Neuron. 2015; 85:1043-55. PMID: 25741728.

712   32. Somasundaram P, Wyrick GR, Fernandez DC, *et al.* C-terminal phosphorylation regulates the kinetics

713   of a subset of melanopsin-mediated behaviors in mice. Proc Nat'l. Acad Sci (Wash) U S A. 2017;

714   114:2741-2746. PMID: 28223508.

715   33. Mure LS, Hatori M, Zhu Q, Demas J, Kim IM, Nayak SK, Panda S. Melanopsin-Encoded Response

716   Properties of Intrinsically Photosensitive Retinal Ganglion Cells. Neuron. 2016; 90:1016-27.

717   PMID: 27181062.

34. Abruzzi K, Chen X, Nagoshi E, Zadina A, Rosbash M. RNA-seq profiling of small numbers of *Drosophila*

neurons. Methods Enzymol. 2015; 551:369-86. PMID: 25662465.

35. Klose M, Duvall L, Li W, Liang X, Ren C, Steinbach JH, Taghert PH. Functional PDF Signaling in the

Drosophila Circadian Neural Circuit Is Gated by Ral A-Dependent Modulation. Neuron. 2016;

90:781-794. PMID: 27161526.

36. Johnson EC, Bohn LM, Taghert PH. *Drosophila* CG8422 encodes a functional diuretic hormone

receptor. J Exp Biol. 2004; 207:743-8. PMID: 14747406.

37. Johnson EC, Shafer OT, Trigg JS, Park J, Schooley DA, Dow JA, Taghert PH. A novel diuretic hormone

receptor in *Drosophila*: evidence for conservation of CGRP signaling. J Exp Biol. 2005; 208:1239-

46. PMID: 15781884.

38. Throckmorton, LH. The phylogeny, ecology and geography of *Drosophila*. In: King RC., editor.

Handbook of Genetics. Vol 3. New York: Plenum; 1975. pp 421-469.

39. Valdez-Lopez JC, Gulati S, Ortiz EA, Palczewski K, Robinson PR. Melanopsin Carboxy-terminus

phosphorylation plasticity and bulk negative charge, not strict site specificity, achieves

phototransduction deactivation. PLoS One. 2020; 15:e0228121. PMID: 32236094.

40. Langlet C, Langer I, Vertongen P, Gaspard N, Vanderwinden JM, Robberecht P. Contribution of the

carboxyl terminus of the VPAC1 receptor to agonist-induced receptor phosphorylation,

internalization, and recycling. J Biol Chem. 2005; 280: 28034 –28043. PMID: 15932876.

41. Dephoure N, Gould KL, Gygi SP, Kellogg DR. Mapping and analysis of phosphorylation sites: a quick

guide for cell biologists. Mol Biol Cell. 2013; 24:535-42. PMID: 23447708.

42. Shafer OT, Taghert PH. RNA-interference knockdown of *Drosophila pigment dispersing factor* in

neuronal subsets: the anatomical basis of a neuropeptide's circadian functions. PLoS One. 2009;

4:e8298. PMID: 20011537..

43. Zhang Y, Emery P. GW182 controls *Drosophila* circadian behavior and PDF-receptor signaling.

Neuron. 2013; Apr 10;78(1):152-65. PMID: 23583112.

44. Beerepoot P, Lam VM, Salahpour A. Measurement of G protein-coupled receptor surface expression.

J Recept Signal Transduct Res. 2013; Jun;33(3):162-5. PMID: 23557016.

45. Zhou XE, He Y, de Waal PW, Gao X, Kang Y, Van Eps N, Yin Y, Pal K, Goswami D, White TA, Barty A,

Latorraca NR, Chapman HN, Hubbell WL, Dror RO, Stevens RC, Cherezov V, Gurevich VV, Griffin

PR, Ernst OP, Melcher K, Xu HE. Identification of Phosphorylation Codes for Arrestin Recruitment

by G Protein-Coupled Receptors. Cell. 2017; 170:457-469.e13. PMID: 28753425.

46. Im SH, Taghert PH PDF receptor expression reveals direct interactions between circadian oscillators

in *Drosophila.* J Comp Neurol. 2010; 518:1925-45. PMID**:** 20394051.

47. Li S, Li S, Han Y, Tong C, Wang B, Chen Y, Jiang, J. (2016). Regulation of Smoothened Phosphorylation

and High-Level Hedgehog Signaling Activity by a Plasma Membrane Associated Kinase. PLoS

Biology, 2016; *14*(6), e1002481. PMID: 27280464.

48. Murthy KS, Mahavadi S, Huang J, Zhou H, and Sriwai W. Phosphorylation of GRK2 by PKA augments

GRK2-mediated phosphorylation, internalization, and desensitization of VPAC2 receptors in

smooth muscle. Am J Physiol Cell Physiol. 2008; 294: C477–C487. PMID: 18077607.

49. Huang J, Mahavadi S, Sriwai W, Grider JR, Murthy KS. Cross-regulation of VPAC(2) receptor

desensitization by M(3) receptors via PKC-mediated phosphorylation of RKIP and inhibition of

GRK2. Am J Physiol Gastrointest Liver Physiol. 2007; 292:G867-74. PMID: 17170028.

50. Shetzline MA, Premont RT, Walker JK, Vigna SR, Caron MG. A role for receptor kinases in the

regulation of class II G protein-coupled receptors. Phosphorylation and desensitization of the

secretin receptor. J Biol Chem. 1998 Mar 20;273(12):6756-62. PMID: 9506976.

51. Walker JK, Premont RT, Barak LS, Caron MG, Shetzline MA. Properties of secretin receptor

internalization differ from those of the 2-adrenergic receptor. J Biol Chem 1999; 274: 31515–

31523. PMID: 10531354.

52. Marie JC, Rouyer-Fessard C, Couvineau A, Nicole P, Devaud H, El Benna J, Laburthe M. Serine 447 in

the carboxyl tail of human VPAC1 receptor is crucial for agonist-induced desensitization but not

internalization of the receptor. Mol Pharmacol. 2003; 64: 1565–1574. PMID: 14645688.

53. Estall JL, Koehler JA, Yusta B, Drucker DJ. The glucagon-like peptide-2 receptor C terminus modulates

beta-arrestin-2 association but is dispensable for ligand-induced desensitization, endocytosis,

and G-protein-dependent effector activation. J Biol Chem. 2005; 280:22124-34. PMID:

15817468.

54. Helfrich-Förster C. The period clock gene is expressed in central nervous system neurons which also

produce a neuropeptide that reveals the projections of circadian pacemaker cells within the

brain of *Drosophila melanogaster*. Proc Nat'l Acad Sci (Wash) U S A. 1995; 92:612-6. PMID:

7831339.

55. Stoleru D, Peng Y, Agosto J, Rosbash M. Coupled oscillators control morning and evening locomotor

behaviour of *Drosophila*. Nature. 2004; 431:862-8. PMID: 15483615.

56. Schlichting M, Menegazzi P, Lelito KR, Yao Z, Buhl E, Dalla Benetta E, Bahle A, Denike J, Hodge JJ,

Helfrich-Förster C, Shafer OT. A Neural Network Underlying Circadian Entrainment and

Photoperiodic Adjustment of Sleep and Activity in *Drosophila*. J Neurosci. 2016; 36(35):9084-96.

PMID: 27581451.

57. Vaze KM, Helfrich-Förster C. The Neuropeptide PDF Is Crucial for Delaying the Phase

of *Drosophila's* Evening Neurons Under Long Zeitgeber Periods. J Biol Rhythms. 2021;36(5):442-

460. PMID: 34428956.

58. Yoshii T, Wülbeck C, Sehadova H, Veleri S, Bichler D, Stanewsky R, Helfrich-Förster C. The

neuropeptide pigment-dispersing factor adjusts period and phase of *Drosophila's* clock. J

Neurosci. 2009; Feb 25;29(8):2597-610. PMID: 19244536.

59. Kula-Eversole E, Lee DH, Samba I, Yildirim E, Levine DC, Hong HK, Lear BC, Bass J, Rosbash M, Allada

R. Phosphatase of Regenerating Liver-1 Selectively Times Circadian Behavior in Darkness via

Function in PDF Neurons and Dephosphorylation of TIMELESS. Curr Biol. 2021; 31:138-149.e5.

PMID: 33157022.

60. Grima B, Chélot E, Xia R, Rouyer F. Morning and evening peaks of activity rely on different clock

neurons of the *Drosophila* brain. Nature. 2004; Oct 14;431(7010):869-73. PMID: 15483616.

61. Stoleru D, Peng Y, Nawathean P, Rosbash M. A resetting signal between *Drosophila* pacemakers

synchronizes morning and evening activity. Nature. 2005; Nov 10;438(7065):238-42. PMID:

16281038.

62. Rieger D, Shafer OT, Tomioka K, Helfrich-Förster C. Functional analysis of circadian pacemaker

neurons in Drosophila melanogaster. J Neurosci. 2006; Mar 1;26(9):2531-43. PMID: 16510731

63. Picot M, Cusumano P, Klarsfeld A, Ueda R, Rouyer F. Light activates output from evening neurons

and inhibits output from morning neurons in the *Drosophila* circadian clock. PLoS Biol. 2007;

Nov;5(11):e315. PMID: 18044989;

64. Zhang Y, Liu Y, Bilodeau-Wentworth D, Hardin PE, Emery P. Light and temperature control the

contribution of specific DN1 neurons to *Drosophila* circadian behavior. Curr Biol. 2010; Apr

13;20(7):600-5. PMID: 20362449.

65. Delventhal R, O'Connor RM, Pantalia MM, Ulgherait M, Kim HX, Basturk MK, Canman JC, Shirasu-Hiza

M. Dissection of central clock function in *Drosophila* through cell-specific CRISPR-mediated clock

gene disruption. Elife. 2019;8:e48308. PMID: 31613218.

66. Menegazzi P, Beer K, Grebler V, Schlichting M, Schubert FK, Helfrich-Förster C. A Functional Clock

Within the Main Morning and Evening Neurons of *D. melanogaster* Is Not Sufficient for Wild-

Type Locomotor Activity Under Changing Day Length. Front Physiol. 2020; Mar 26;11:229. PMID:

32273848.

67. Schlichting M, Díaz MM, Xin J, Rosbash M. Neuron-specific knockouts indicate the importance of

network communication to *Drosophila* rhythmicity. Elife. 2019 Oct 15;8:e48301. PMID:

31613223

68. Yao Z, Shafer OT. The *Drosophila* circadian clock is a variably coupled network of multiple peptidergic

units. Science (Wash). 2014; 343(6178):1516-20. PMID: 24675961.

69. Chiu JC, Ko HW, Edery I. (2011) NEMO/NLK phosphorylates PERIOD to initiate a time-delay

phosphorylation circuit that sets circadian clock speed. Cell. 2011; 145:357-70. PMID: 21514639.

70. Yu W, Houl JH, Hardin PE. NEMO kinase contributes to core period determination by slowing the

pace of the *Drosophila* circadian oscillator. Curr Biol. 2011; 21:756-61. PMID: 21514156.

71. Blau J, Young MW. Cycling vrille expression is required for a functional *Drosophila* clock. Cell. 1999;

Dec 10;99(6):661-71 PMID: 10612401.

72. Levine JD, Funes P, Dowse HB, Hall JC. Signal analysis of behavioral and molecular cycles. BMC

Neurosci. 2002; 3:1. PMID: 1182533.

73. Wheeler DA, Hamblen-Coyle MJ, Dushay MS, Hall JC. Behavior in light-dark cycles of *Drosophila*

mutants that are arrhythmic, blind, or both. J Biol Rhythms. 1993; 8:67-94. PMID: 8490212.

Figure 1. A map illustrating the series of PDFR sequence variants used to test a role for PDFR phosphorylaytion in regulating the locomotor rhythms. The C Terminal of the PDFR-A isoform is diagrammed (top) immediately following the position of the 7^{th} transmembrane domain (TM7, blue). Asterisks indicate the positions of the 28 serine, threonine or tyrosine residues present within the PDFR C Terminal tail. The residues marked by the white asterisks were those chosen for study by sequence alteration to encode Alanine; they were given arbitrary designations as Clusters (CL) 1 through 7. Each cluster contains a single residue (e.g., CL4 or CL5), or as many as four residues (e.g., CL2). The ten sequence variants are named by the Cluster(s) that was altered followed by the letter 'A". Simple Variants include those wherein residues in one or two Clusters were altered. Multiple Variants include those wherein residues within three or more Clusters were altered. All 14 targeted residues were altered in the PDFR 1-7A variant.

Figure 2. Locomotor Rhythms exhibited by WT PDFR and by Simple PDFR Variants under Winter-like (Short Day) conditions. All behavioral records were recorded from *han* (*pdfr* mutant) flies that expressed either no UAS transgene (A), or a UAS-WT *pdfr* transgene (B) or a variety of Simple *PDFR* Variants, including 2-3A (C), 4A (D), 5A (E), 6A (F) or 7A (G), or Multiple PDFR variants, including 5-7A (H), 1-4A (I), 1-5A (J), 1-6A (K) and 1-7A (L). Each Panel (A)-(L) contains sub-panels (1) and (2). Sub-Panel (1) displays a double-plotted group actogram throughout the 6 days of Light: dark entrainment, followed

by ~9 days of constant darkness (DD, grey background). Sub-panel (2) displays a daily plot of locomotor

activity (a group eduction) averaged over the last two days of entrainment (LD 5-6). Open bars indicate

the periods of Lights-on and filled bars indicate periods of Lights-off. Panel M displays the average

Phase Offset timepoint for the Evening activity phase (marked by a Blue Arrow) for each genotype over

the last two days of entrainment (LD 5-6). Panel N displays the average Phase Onset timepoint for the

Morning activity phase (marked by a Red Arrow) for each genotype over the last two days of

entrainment (LD 5-6). The positions of the Blue and Red arrow in panels A-L are representative; panels

M and N presents their true values. Analyses represent ANOVA followed by Dunnett's post hoc multiple

comparisons of all compared to WT: ns = not significant; * = $p<0.05$; ** = $p<0.01$; *** = $p<0.005$; **** =

$p<0.001$.

Figure 3. Amplitude measures of Locomotor activity exhibited by WT PDFR and by diverse

PDFR Variants under Winter-like (Short Day) conditions. All behavioral records were recorded from

han (*pdfr* mutant) flies that expressed either no UAS transgene (A – marked in YELLOW), or a WT *pdfr*

cDNA (B – marked in BLACK) or a variety of Simple *pdfr* Variants (marked in RED) including 2-3A (C), 4A

(D), 5A (E), 6A (F), 7A (G), 5-7A (H), 1-4A (I), 1-5A (J), 1-6A (K) or 1-7A (L). These measures were

averaged across all experiments run for each individual genotype (see Table 1 for N and n values). Each

Panel (A)-(L) contains sub-panels (1) through (3), each of which displays Bin-by-Bin analyses of activity

levels sorted by 30 min bins, for three different time periods: (1) ZT17-23.5; (2) ZT 0.5 – 8; (3) ZT 8.5-16.

The missing bin at timepoint 0 contains the startle response that accompanies the sudden lights-on

signal. Asterisks indicate significantly-different activity levels according to a Student's T-test ($p< 0.05$).

Red arrows indicate elevated Morning activity levels displayed by different PDFR variants. Blue arrows

indicate elevated Evening activity levels produced by different PDFR variants.

Figure 4. Locomotor Rhythms exhibited by WT PDFR and by Simple PDFR Variants under Equinox (12:12) conditions. All behavioral records were recorded from *han* (*pdfr* mutant) flies that expressed either no UAS transgene (A), or a UAS-WT *pdfr* transgene (B) or a variety of Simple *pdfr* Variants, including 2-3A (C), 4A (D), 5A (E), 6A (F) or 7A (G), or Multiple pdfr variants, including 5-7A (H), 1-4A (I), 1-5A (J), 1-6A (K) and 1-7A (L). Each Panel (A)-(L) contains sub-panels (1) and (2). Sub-Panel (1) displays a double-plotted group actogram throughout the 6 days of Light: dark entrainment, followed by ~9 days of constant darkness (DD, grey background). Sub-panel (2) displays a daily plot of locomotor activity (a group eduction) averaged over the last two days of entrainment (LD 5-6). Open bars indicate the periods of Lights-on and filled bars indicate periods of Lights-off. Panel M displays the average Phase Offset timepoint for the Evening activity phase (marked by a Blue Arrow) for each genotype over the last two days of entrainment (LD 5-6). Panel N displays the average Phase Onset timepoint for the Morning activity phase (marked by a Red Arrow) for each genotype over the last two days of entrainment (LD 5-6). The positions of the Blue and Red arrow in panels A-L are representative; panels M and N presents their true values. Analyses represent ANOVA followed by Dunnett's post hoc multiple comparisons of all compared to WT: ns = not significant; * = $p<0.05$; ** = $p<0.01$; *** = $p<0.005$; **** = $p<0.001$.

Figure 5. Locomotor Rhythms exhibited by WT PDFR and by Multiple PDFR Variants under Equinox (12:12) conditions. All behavioral records were recorded from *han* (*pdfr* mutant) flies that expressed either no UAS transgene (A – marked in YELLOW), or a WT *pdfr* cdNA (B – marked in BLACK) or a variety of Simple *pdfr* Variants (marked in RED) including 2-3A (C), 4A (D), 5A (E), 6A (F), 7A (G), 5-7A (H), 1-4A (I), 1-5A (J), 1-6A (K) or 1-7A (L). These measures were averaged across all experiments run for each individual genotype (see Table 1 for N and n values). Each Panel (A)-(L) contains sub-panels (1)

through (3), each of which displays Bin-by-Bin analyses of activity levels sorted by 30 min bins, for three

different time periods: (1) ZT17-23.5; (2) ZT 0.5 – 8; (3) ZT 8.5-16. The missing bin at timepoint 0

contains the startle response that accompanies the sudden lights-on signal. Asterisks indicate

significantly-different activity levels according to a Student's T-test (p< 0.05). Red arrows indicate

elevated Morning activity levels displayed by different PDFR variants. Blue arrows indicate elevated

Evening activity levels produced by different PDFR variants.

Figure 6. Locomotor Rhythms exhibited by WT PDFR and by Simple PDFR Variants under

Summer-like (Long Day) conditions. All behavioral records were recorded from *han* (*pdfr* mutant) flies

that expressed either no UAS transgene (A), or a UAS-WT *pdfr* transgene (B) or a variety of Simple *pdfr*

Variants, including 2-3A (C), 4A (D), 5A (E), 6A (F) or 7A (G), or Multiple *pdfr* variants, including 5-7A (H),

1-4A (I), 1-5A (J), 1-6A (K) and 1-7A (L). Each Panel (A)-(L) contains sub-panels (1) and (2). Sub-Panel (1)

displays a double-plotted group actogram throughout the 6 days of Light: dark entrainment, followed by

~9 days of constant darkness (DD, grey background). Sub-panel (2) displays a daily plot of locomotor

activity (a group eduction) averaged over the last two days of entrainment (LD 5-6). Open bars indicate

the periods of Lights-on and filled bars indicate periods of Lights-off. Panel M displays the average

Phase Offset timepoints for the Evening activity periods (marked by a Blue Arrow) for each genotype

over the last two days of entrainment (LD 5-6). Panel N displays the average Phase Onset timepoints for

the Morning activity periods (marked by a Red Arrow) for each genotype over the last two days of

entrainment (LD 5-6). The positions of the Blue and Red arrows in panels A-L are representative; panels

M and N presents their true values. . Analyses represent ANOVA followed by Dunnett's post hoc

multiple comparisons of all compared to WT: ns = not significant; * = p<0.05; ** = p<0.01; *** =

p<0.005; **** = p<0.001.

Figure 7. Locomotor Rhythms exhibited by WT PDFR and by Multiple PDFR Variants under

Summer-like (Long Day) conditions. All behavioral records were recorded from *han* (*pdfr* mutant) flies

that expressed either no UAS transgene (A – marked in YELLOW), or a WT *pdfr* cdNA (B – marked in

BLACK) or a variety of Simple *pdfr* Variants (marked in RED) including 2-3A (C), 4A (D), 5A (E), 6A (F), 7A

(G), 5-7A (H), 1-4A (I), 1-5A (J), 1-6A (K) or 1-7A (L). These measures were averaged across all

experiments run for each individual genotype (see Table 1 for N and n values). Each Panel (A)-(L)

contains sub-panels (1) through (3), each of which displays Bin-by-Bin analyses of activity levels sorted

by 30 min bins, for three different time periods: (1) ZT17-23.5; (2) ZT 0.5 – 8; (3) ZT 8.5-16. The missing

bin at timepoint 0 contains the startle response that accompanies the sudden lights-on signal. Asterisks

indicate significantly-different activity levels according to a Student's T-test (p< 0.05). Red arrows

indicate elevated Morning activity levels displayed by different PDFR variants. Blue arrows indicate

elevated Evening activity levels produced by different PDFR variants.

Figure 8. A model predicting the effects of phosphorylating different sites on the PDFR CT on

downstream signaling. Asterisks indicate the positions Ser/ Thr/ and Tyr residues as described in Figure

1 and Supplemental Figure 1: white asterisks were targets of mutational analysis in this study. Mutation

of CL6 and CL7 consistently demonstrated the greatest delaying effects on the phases of locomotor

peaks, suggesting their phosphorylation normally will have the greatest effect to slow or terminate PDFR

signaling. Other sites are also effective, although to lesser degrees, including CL2-3, CL4 and CL5.

Primarily, such phosphorylation will decrease the extent of PDFR signaling (red bars) and so reduce the

delay that PDF imposes on the period of neuronal activation displayed by PDFR-responsive pacemaker

groups (like the Evening cells and the DN3: Liang *et al.*, 2016, 2017). The model also predicts that the

effects of phosphorylating CL5 will depend on the phosphorylation status of neighboring sites.   In some

contexts (e.g., PDFR 1-5A), it will help terminate PDFR signaling, but in others (e.g., PDFR 567A) it may

promote the duration or extent of PDFR signaling, perhaps by blocking the effects of phosphorylating

CL6 and CL7 (green bar).

Figure 1.

Figure 1

Figure 2

Figure 2

Figure 3 .

Figure 3

Figure 4

Figure 4

Figure 5

Figure 5

Figure 6

Figure 6

Figure 7.

Figure 7

Figure 8.

Figure 8