Putative host-derived insertions in the genomes of circulating SARS-CoV-2 variants

Yiyan Yang[1], Keith Dufault-Thompson[1], Rafaela Salgado Fontenele[1], Xiaofang Jiang[1,*]

[1]National Library of Medicine, National Institutes of Health, Bethesda, Maryland, USA

[*] Correspondence: Xiaofang Jiang (xiaofang.jiang@nih.gov)

ABSTRACT

Insertions in the SARS-CoV-2 genome have the potential to drive viral evolution, but

the source of the insertions is often unknown. Recent proposals have suggested that

human RNAs could be a source of some insertions, but the small size of many

insertions makes this difficult to confirm. Through an analysis of available direct

RNA sequencing data from SARS-CoV-2 infected cells, we show that viral-host

chimeric RNAs are formed through what are likely stochastic RNA-dependent RNA

polymerase template switching events. Through an analysis of the publicly available

GISAID SARS-CoV-2 genome collection, we identified two genomic insertions in

circulating SARS-CoV-2 variants that are identical to regions of the human 18S and

28S rRNAs. These results provide direct evidence of the formation of viral-host

chimeric sequences and the integration of host genetic material into the SARS-CoV-2

genome, highlighting the potential importance of host-derived insertions in viral

evolution.

IMPORTANCE

Throughout the COVID-19 pandemic, the sequencing of SARS-CoV-2 genomes has

revealed the presence of insertions in multiple globally circulating lineages of

SARS-CoV-2, including the Omicron variant. The human genome has been suggested

to be the source of some of the larger insertions, but evidence for this kind of event

occurring is still lacking. Here, we leverage direct RNA sequencing data and

SARS-CoV-2 genomes to show host-viral chimeric RNAs are generated in infected

cells and two large genomic insertions have likely been formed through the

incorporation of host rRNA fragments into the SARS-CoV-2 genome. These

host-derived insertions may increase the genetic diversity of SARS-CoV-2 and

expand its strategies to acquire genetic materials, potentially enhancing its

adaptability, virulence, and spread.

KEYWORDS

SARS-CoV-2, insertion, host-virus chimeric reads, ribosomal RNA

INTRODUCTION

During the COVID-19 pandemic, insertions have been frequently acquired in

SARS-CoV-2 lineages (1-4). Insertions have been associated with several globally

circulating lineages, including the insertion of one amino acid at position 146 of the S

protein (ins146N) of the variant of interest Mu (B.1.621) (4), insertions at the

recurrent insertion site 214 of the NTD region on the S protein that occurred in the

lineages B.1.214.2 (ins214TDR) and A.2.5 (ins214AAG) (1), and the insertion

ins214EPE in the recently-emerged variant of concern Omicron (5). Although there is

insufficient evidence to show the direct impact these insertions have on viral spread

and interference with immune responses, the fact that variants carrying those

insertions have circulated for long periods suggests that they might be advantageous

or neutral for the transmission. Results from a long-term *in vitro* experiment where

SARS-CoV-2 was co-incubated with highly neutralizing antibodies have also shown

that an 11 amino acid insertion (ins248KTRNKSTSRRE) at the NTD N5 loop of the

S protein was able to drive antibody escape suggesting a potential role of insertions in

enhancing infectivity and virulence (6). Taken together, insertions have the potential

to increase genetic diversity in SARS-CoV-2 and contribute to the continued

evolution of the virus.

Previous research has shown that most small insertions in the SARS-CoV-2 genome

likely originated from template sliding, local duplication, or template switching

between viruses (2). Longer insertions (equal or larger than nine nucleotides) have

been detected in multiple coronavirus genomes, including in variants of concern like

the Omicron variant, but their origin remains unknown. Host genetic material has

been suggested as a possible source for these insertions (5, 7). Venkatakrishnan et al.

suggested that the unique insertion (ins214EPE) in the Omicron variant could have

originated from the human common cold virus HCoV-229E or the human genome

based on BLAST search (5), and the human genome has been speculated to be the

source of multiple other small insertions (7). However, given that these insertion

sequences are typically short, sequence comparisons tend to be less informative, and

false-positive matches have a high chance of occurring. Additionally, coronavirus

replication occurs in modified endoplasmic reticulum-derived double-membrane

vesicles, providing a physical barrier between viral and host genetic material (8), and

coronavirus replication complexes are known to contain enzymes with proofreading

activity (9), both of which likely play roles in limiting the formation of host-virus

chimeric sequences.

Human-derived insertions in the SARS-CoV-2 genome would likely be generated

through RdRp-driven template switching events between SARS-CoV-2 and host

mRNA. While template switching events between coronaviruses are common (10-13)

and likely contribute to the emergence of SARS-CoV-2 lineages including the

deltacron variant (14), template switching events between coronaviruses and host

RNAs are rarely documented (15, 16). Chimeric reads between SARS-CoV-2 RNA

and human RNA have been detected but were interpreted as a signal of SARS-CoV-2

integration into the human genome in a previous controversial study (17). Others have

suggested that the chimeric reads were likely to be template switching artifacts

mediated by reverse transcriptase or PCR during library preparation (18-21). One

possible explanation that was largely omitted in these studies is that the

SARS-CoV-2-host chimeric RNA could be generated by RdRp-driven template

switching.

Here, to investigate the possible existence of SARS-CoV-2-host chimeric RNA, we

take advantage of the publically available Nanopore direct RNA sequencing data of

SARS-CoV-2. Direct RNA-seq sequences the individual polyadenylated RNAs

directly mitigating the possible formation of chimeric reads during library preparation

or amplification. We first identified SARS-CoV-2-host chimeric RNA from direct

RNA-seq data and showed that RdRp-driven template switching between

SARS-CoV-2 and host mRNA occurs, but it is infrequent and stochastic. We also

found that highly expressed host genes and structural RNA genes have a higher

chance to be observed in chimeric RNA reads. We then systematically analyzed the

SARS-CoV-2 genomes deposited in the GISAID database (22), resulting in the

identification of two insertions in functional SARS-CoV-2 genomes that likely

originated from the host 18S and 28S rRNAs.

RESULTS

Host-virus mRNA chimera are rare but do exist

We first analyzed direct RNA-seq data from SARS-CoV-2 infected cell lines to

identify sequences formed from chimeric host-viral RNAs. The direct RNA-seq data

were quality filtered and mapped to both the host and SARS-CoV-2 transcriptomes to

identify potential chimeric sequences. Out of the 30 samples that were analyzed,

host-viral chimeric reads were detected in 16 of the samples with an average of 0.029%

(standard deviation 0.048%) of the reads mapped to SARS-CoV-2 being chimeric

(Supplementary Table 1). Chimeric reads were typically rare, making up 0.206% of

one sample, but less than 0.06% of the other 15 samples, and these rates may be an

overestimation due to the cell lines used compared to what would be observed in *in*

vivo conditions. Additionally, chimeric reads detected in five samples were further

investigated using paired-end sequencing short reads from the same samples

(Supplementary Table 2). Approximately 1.4% (5 out of 357) of chimeric reads were

supported by at least five read pairs spanning the junctions. This finding implies that a

small fraction of the host-viral chimeric mRNA molecules could function as templates

for RNA replication.

We then analyzed the chimeric reads to identify trends in how the viral and host RNA

sequences were joined. All the viral-derived sequences in chimeric reads were

annotated as positive-sense RNA and a majority (92.49%) of the reads contained

host-derived positive-sense sequences. Upon further examination, the few host reads

that were identified as being negative-sense were largely long non-coding RNAs that

were present in the raw reads as the negative-sense sequences, making it likely that

they were mis-annotated rather than actually being derived from negative-sense RNA.

These results suggest that the host-viral chimeric sequences are not the result of the

integration of the viral genetic material into the host genome, which would have

resulted in a nearly equal mix of positive and negative sense viral sequences (17).

Most likely, these host-viral chimeric sequences were created from

positive-to-positive-strand template switching events (23, 24).

Viral-host chimeric read formation is likely a stochastic process

The chimeric reads were then analyzed to determine if there were any patterns in the

composition of the sequences and in which positions relative to the references they

were formed. Both viral to host and host to viral chimeric sequences were detected in

the direct RNA-seq data, but the chimeric reads did not show a preference for either

organization (Supplementary Table 1). Both types of sequences were seen in

approximately the same frequency, with viral to host reads making up 55% of the

chimeric sequences and host to viral reads making up 45%. This lack of strong

preference may indicate that host RNA can be readily recognized by viral RdRp, but

other factors like the exclusion of host RNA by the formation of the

double-membrane vesicles might prevent the formation of chimeric RNAs. When

examining the positions of the junctions on the viral RNA sequences, we found there

was a bias toward the junction sites being located in the dense coding region near the

three prime end of the sequence, with fewer junctions being identified in the ORF1ab

genes, the largest region of the genome (Fig.1). This is likely due to the ORF1ab

region not being retained in the canonical SARS-CoV-2 subgenomic RNAs resulting

in fewer viral RNAs being synthesized with these regions that could form chimeric

RNAs (25). It suggests that the process by which chimeric sequences are formed is

likely stochastic, depending on the availability of template RNA molecules.

Previous studies have also found that indel formation and template switching events

preferentially occur in the loops and stems formed in the RNA secondary structure (2,

3). First a permutation test was used to investigate if junction sites were commonly

located in stems (positions that form base-pairs) or non-stem regions (non-base-paired

positions) in the viral RNA. The results of this test showed a significant (P-values <

0.01) preference for the formation of junctions in non-base-paired regions of the RNA

secondary structure (Fig.1). One-sided Fisher's exact tests were performed to explore

if junction sites were enriched in specific types of RNA structures. Consistent with the

results of the permutation test, stems were under-represented at the junction sites

(Supplementary Table 3). We speculate that the non-base-paired regions of the

SARS-CoV-2 RNA may be more susceptible to stochastic template-switching events

due to their more "open" configurations, where the viral RdRp could easily attach or

detach as it moves along the RNA.

An examination of the types of human gene sequences found in the chimeric

sequences revealed an enrichment of non-coding RNAs and highly expressed genes.

We found that a disproportionate number of non-coding RNAs, mainly long

non-coding RNAs (lncRNAs), were forming parts of the chimeric reads compared to

their abundance in the human genomes. These non-coding RNA chimeric sequences

made up 8.8% and 10.5% of the chimeric reads detected in the Caco and Calu cell

lines, respectively, while non-coding sequences made up only 4% of the genes

annotated in the human genome. This enrichment of non-coding RNA chimeric

sequences was tested using Fisher's exact test confirming that the trend was

significant (Caco cells: odds ratio=2.2, P-value=0.043; Calu cells: odds ratio=2.8,

P-value=0.001). When analyzed in the context of the expression level of the host

genes in each sample, we also observed an enrichment for highly expressed genes

forming parts of the chimeric sequences (Fig.2). This enrichment was confirmed

through the Mann-Whitney U tests showing that the trend was significant in the two

human cell lines (P-value < 2.2e-16 for both) and the *Chlorocebus sabaeus* (green

monkey) cell line (P-value < 2.2e-16). These results appear to highlight two groups of

sequences that are forming chimeric RNAs, structural RNAs like lncRNAs, which

may be susceptible due to their secondary structures, and highly expressed genes,

which would have more RNA molecules present for template-switching events to

occur with. This suggests that the formation of chimeras is largely stochastic, with

factors like the abundance of RNAs playing a large role, but that certain RNA

molecules may be more susceptible to these events due to their structure.

Systematic search for host-derived insertions in SARS-CoV-2 genomes

We performed a survey of the GISAID SARS-CoV-2 genomes to identify insertions

with potential host origins. Insertions were detected based on alignments and

comparison to the Wuhan-Hu-1/2019 reference genome. Only insertions greater than

or equal to 21 nucleotides long and that were found outside of the 5' and 3'

untranslated regions were considered in subsequent analyses (Supplementary Table 4).

Of the 36 insertions that were found, 17 of them were found in multiple SARS-CoV-2

genomes but were not monophyletic. Upon further examination, the genomes

containing these insertions tended to be sequenced by the same labs around the same

times making it likely that these detected insertions are due to library preparation or

sequencing errors rather than the result of multiple independent insertion events in

different viral lineages. Of the 19 other insertions, 16 of them were only detected in a

single genome, and while many of these had plausible hits to human genes, it is

difficult to assess if these are true insertions or library preparation or sequencing

artifacts due to their limited presence.

The three remaining insertions were from monophyletic virus variants and were

further examined to determine if they had plausible homologous sequences in the

human genome. Two of the insertions were found to be identical to conserved

segments of the 28S and 18S rRNAs and were analyzed further. The remaining

insertion was 21 nucleotides long and was found in 6 SARS-CoV-2 genomes of the

Alpha B.1.1.7 lineage. These genomes were collected in early March of 2021 from

England, United Kingdom by two laboratories, and sequenced at the same location

using the same sequencing platform. The raw reads were available for two of the

genomes, namely England/ALDP-13C8C28/2021 (EPI_ISL_1331302) and

England/QEUH-13C1955/2021 (EPI_ISL_1332461), and were examined directly,

providing confirmation that the insertion was present and likely not an artifact.

Unfortunately, no plausible source for this insertion was able to be identified using a

BLAST search in the NCBI non-redundant nucleotide database and a collection of

coronavirus genomes with a cutoff E-value of 1e-2, and it was not analyzed further.

28S rRNA-derived insertion in SARS-CoV-2 genomes

We detected a 27-nucleotide long insertion in five SARS-CoV-2 genomes

(Supplementary Table 5 and Fig.3A) at position 7120 of the reference genome

(China/Wuhan-Hu-1/2019). The five genomes containing the 28S rRNA-derived

insertions were collected by different laboratories and were sequenced on different

sequencing platforms, making it extremely unlikely that laboratory error is

responsible for the presence of the insertions. The five genomes belong to a

monophyletic group. In this clade, there are three other variants whose assembled

genomes do not contain the insertion. We were able to obtain access to the raw

genome sequencing data of two of the three variants — USA/WA-PHL-005726/2021

(EPI_ISL_6259191) and USA/HI-H215617/2021 (EPI_ISL_6540096). We then did

further analysis on the raw sequencing to check if the insertion is indeed missing.

First, we generated consensus genome sequences based on the alignment of

sequencing reads to the SARS-CoV-2 reference genome and found the consensus

sequences did not contain the insertion. Next, we manually added the 28S

rRNA-derived insertion at position 7120 of the consensus genome and compared the

reads exclusively aligned to the consensus genome with the insertion and the reads

exclusively aligned to the consensus genome without the insertion. We found that

99.76% (8700/8721 for EPI_ISL_6259191) and 99.93% (1502/1503 for

EPI_ISL_6540096) of the exclusively-mapped reads support the presence of the

insertion in the genomes. The reason that the insertion is missing in the submitted

genomes (EPI_ISL_6259191 and EPI_ISL_6540096) is likely that the assembly was

generated using an insertion-unaware approach, such as reference-based consensus

calling. For the only one variant that was missing the insertion in the genomes, we are

not able to assess if it is due to failure to identify the insertion based on the consensus

caller or the subsequent loss of the inserted sequence.

By performing the BLAST search for this insertion against the human transcripts

(Release 109 RNAs), an exact match (E-value: 2e-06) of this insertion was found in

the nucleotide sequences of 28S ribosomal RNA (Fig.3B). We observed an extra three

overlapping bases in the pairwise alignment of SARS-CoV-2 variants containing the

insertion and the human 28S rRNA sequence, extending the length of identity

nucleotide bases from 27 nucleotides to 30 nucleotides. The identical region was

located at positions 4969-4998 of the human 28S rRNA (based on the structure of

PDB 5AJ0 Chain A2) and makes up part of the highly conserved loop 94 stem of

domain 7 of the rRNA molecule according to the Gorski et al.'s segmentation of

human 28S rRNA (26) (Fig.3B).

Due to the high level of sequence conservation of 28S rRNA, asserting the origin of

the insertion-related 30 nucleotide sequences is impossible based on sequence identity

alone. In the human genome (GRCh38 release 105), three 28S rRNA gene copies in

chromosome 21 and one copy in chromosome 12 contain the exact 30 nucleotide

sequences. When we searched the 30 nucleotide sequences in the LSU rRNA database

downloaded from SILVA (27), 98 organisms were found to contain the sequences.

The last common ancestor of these 98 organisms is *Euteleostomi* (bony vertebrates).

Given the fact that the insertion emerged from the SARS-CoV-2 variant circulating in

humans, the originating organism of the 28S rRNA-derived insertion is most likely

humans.

The nine amino acid insertion is located at position 1467 of the ectodomain (3Ecto) in

the Nsp3 protein, the only domain of this protein located on the lumenal side of the

endoplasmic reticulum (Fig.3C). Nsp3 along with Nsp4 and Nsp6 have been shown to

be involved in the formation of double membrane vesicles in coronavirus infected

cells (28, 29). The 3Ecto domain is specifically involved in the recruitment of Nsp4

and has been shown to be an essential component of Nsp3 for correct

double-membrane vesicle formation (28). At this point, it is unclear if this insertion

would have had an effect on viral fitness, but given its location in the 3Ecto domain, it

is possible that the insertion could have an effect on the interactions between Nsp3

and other proteins and on the membrane rearrangement process.

The monophyletic group with the 28S rRNA-derived insertion belonged to the

AY.103 group of the delta lineage (30) (Fig.3A). The AY.103 variant was first

detected worldwide on January 1st, 2021 and in the USA on January 2nd, 2021. The

clade containing the 28S rRNA-derived insertion is defined by five nucleotide

mutations (T7900C, A10420T, C18646T, C25721T, and C29668T). By September

2021, AY.103 had become the most common delta lineage in the United States and

has continued to be responsible for a significant fraction of cases until the recent

emergence of the Omicron variant (31). The five genomes containing the 28S

rRNA-derived insertion were collected between October 9th and November 10th in

2021 from the states of Washington, Idaho, Massachusetts, and California, indicating

that these variants were likely being transmitted over this timeframe, but the extent to

which it was being spread seems to be low as Idaho was the only state where multiple

genomes were collected from and no genomes containing the insertion have been

reported since. Based on the limited spread of the viruses containing the 28S

rRNA-derived insertion, it is likely that the insertion might not confer phenotypic

advantages or is possibly disadvantageous to the virus. Nonetheless, our data show

that AY.103 lineages containing this insertion were viable and were transmitted for a

short period of time.

18S rRNA-derived insertion in SARS-CoV-2 genomes

A 24-nucleotide insertion was detected in two genomes at position 27492 in the

genome of the reference genome (China/Wuhan-Hu-1/2019) (Supplementary Table 5).

A sequence search against human transcripts (Release 109) was performed using

BLAST (32), resulting in the identification of an exact match to a 24 nucleotide

stretch (E-value: 2e-5) of the 18S rRNA sequence. When aligned to the full 18S

rRNA sequence, it was found that the identical region extended one additional

nucleotide outside of the insertion region, bringing the identical stretch to 25

nucleotides (Fig.4A). The insertion was identical to a highly conserved region of the

18S rRNA (at positions 399-423 in 18S rRNA), consisting of a portion of the helix 12

of the 5' domain (33, 34). In the human genome alone there are five copies of the 18S

rRNA gene on chromosome 21 that contain identical matches for this 25 nucleotide

sequence. When compared to the SSU rRNA SILVA database (27), identical

sequences were found in the 18S sequences of 2289 organisms, which had a common

ancestor of *Opisthokonta* (Fungi/Metazoa group). Considering that the viral samples

were circulating in human populations, it is highly likely that the insertion was

derived from human 18S rRNA.

The insertion is in the SARS-CoV-2 ORF7a protein, encoding an eight amino acid

sequence that is located between the proline and cysteine at positions 34 and 35 in the

reference protein sequence (Fig.4B). The cysteine at position 35 is known to form a

disulfide bond with a cysteine at position 67 and is thought to help stabilize the

beta-sheet structure (35, 36) and the possible functions of the proline at position 34

are not known. The ORF7a protein has been shown to contain an

immunoglobulin-like ectodomain between residues 16 and 96 on the protein which is

thought to have a role of binding to human immune cells and modulating immune

response (35-37). Given the proximity of the insert to the disulfide bond forming

cysteine at position 34 and the size of the insert it is possible that this insert would

have an effect on the overall structure and immunoregulatory functions of ORF7a, but

without additional evidence, the effect of this insertion on the fitness of the virus

remains unknown.

The two genomes containing the 18S rRNA insertion were from the same clade in the

Alpha B.1.1.7 SARS-CoV-2 lineage, which was first identified in England, United

Kingdom in mid-December of 2020 (Fig. 4C). This variant was designated as a

variant of concern due to its transmissibility and large number of mutations and

quickly became the dominant variant in England while spreading to other countries

(38). The genomes containing the 18S rRNA-derived insertion, along with the other

four genes in the same clade, were collected in April and May of 2021 in Oregon,

United States. The genomes from the variants containing the insertion were collected

and sequenced by different labs using different sequencing platforms, making it

unlikely that the insertion was a sequencing or library preparation artifact. We did not

detect the insertion in any of the other four genomes from this clade, indicating that

either they do not have the insertion, they have it but it was not detected, or that the

insertion was only acquired in a sub-clade within this group. After May of 2021, no

new genomes containing this insertion were collected, indicating that the period

during which these lineages were circulating may have been brief. While these viral

variants seem to be viable and transmitted for a short period of time, the insertion

likely does not confer a significant advantage or may be disadvantageous for the virus

resulting in its limited spread.

DISCUSSION

Insertions in the SARS-CoV-2 genome can be introduced through multiple

mechanisms and have the potential to give rise to new variants with enhanced

infectivity, pathogenicity, and antibody escape (2, 6), but the source of these

insertions is often difficult to determine and has been hotly debated (5, 7). Leveraging

available direct RNA sequencing data and an analysis of SARS-CoV-2 genomes, we

have found evidence of the formation of viral-host chimeric RNA sequences and

described two novel human-derived genomic insertions present in circulating variants

of SARS-CoV-2.

Through our screening of direct RNA-seq data from SARS-CoV-2 infected cell lines,

we found that viral-host chimeric RNAs were rare but were present in approximately

half of the samples analyzed. The chimeric reads all contained positive-sense viral

RNA sequences, indicating that these chimeric sequences are not the result of the

integration of the viral genetic material into the host genome, which would have

resulted in a nearly equal mix of positive and negative sense viral sequences (17).

This process does appear to be stochastic in nature though, with no preference for

starting with host or viral sequences during chimera formation and a higher frequency

of chimeras being formed with highly expressed genes in the cells. The regions in the

RNA where these template switching events occur appears to be influenced by the

secondary structure of the viral RNA, possibly due to certain structures being more

susceptible to template switching events similar to what has been reported in previous

studies (2, 3). The accurate determination of the exact junction boundaries and

potential base-pairings were hindered by the high error rate of 14% in direct

RNA-sequencing data and the limited number of host-viral chimeras detected in this

study. The exact molecular basis for the viral-host chimera remains unclear and future

investigation with larger sets of error-corrected direct RNA-seq data of SARS-CoV-2

could be beneficial to address this question.

The formation of host-viral chimeric mRNAs or subgenomic RNAs could mostly be

transient events, not having a long-term impact on viral fitness, but the possibility of

human-derived insertions in the coronavirus genomes could have significant

implications considering the role that genomic insertions seem to have in the

evolution of new SARS-CoV-2 variants (5, 6). The putative 18S and 28S-derived

insertions were identified in circulating variants of the SARS-CoV-2, and while these

particular variants did not seem to spread widely, they do provide evidence that

human genetic material can be a source of genomic insertions in SARS-CoV-2.

Interestingly, rRNAs have been established to be a source of insertions in influenza

genomes, in some cases resulting in significantly more pathogenic viral variants (39,

40). It has been speculated that these recombination events often occur with host

rRNAs due to their abundance in the cells, the presence of recombination hotspots on

rRNA molecules, and the utilization of host rRNAs during viral replication (39).

Similar factors may play a role in the formation of these rRNA-derived insertions in

SARS-CoV-2, but the formation of double-membrane vesicles during SARS-CoV-2

would seemingly complicate this process. There may be accidental capture of host

RNAs inside of the double-membrane vesicles during their formation or some

crossover of host RNA from the cytosol, but evidence of this is lacking and warrants

further investigation.

CONCLUSIONS

Overall, our results suggest that viral-host chimeric sequences can be formed, likely

through stochastic RdRp template switching events. Furthermore, we have identified

two long insertions in SARS-CoV-2 genomes in previously circulating variants which

are likely derived from human ribosomal RNAs. While the source of smaller

insertions that are present in many SARS-CoV-2 genomes are still difficult to identify

due to their short lengths, these results provide evidence that bolsters the hypothesis

that some of them are derived from human genetic material. The mechanisms at work

in the formation of these chimeric RNAs and genomic insertions are still unclear but

warrant further study considering the potential importance of these processes in viral

evolution and the emergence of new variants.

METHODS

Identification of host-virus chimeric reads in SARS-CoV-2 direct-RNA seq data

The nanopore direct RNA-seq data from SARS-CoV-2 infected cell lines were

downloaded from the NCBI SRA database (Supplementary Table 1). All reads were

quality trimmed using NanoFilt v2.8.0 (41), to remove the first 50 nucleotides of each

read and require an average quality score of at least 10 over the length of the read.

The trimmed reads were then mapped using Minimap2 v2.23 (42) to the

SARS-CoV-2 reference genome (NCBI GenBank accession: NC_045512.2) (43), and

either a reference *Chlorocebus sabaeus* transcriptome

(ftp://ftp.ensembl.org/pub/release-105/fasta/chlorocebus_sabaeus/) or human

transcriptome (ftp://ftp.ensembl.org/pub/release-105/fasta/homo_sapiens/). The

mapping files were converted to the Pairwise mApping Format (PAF) using the

paftools script that is part of Minimap2 (42). Reads that mapped to both the host and

SARS-CoV-2 transcriptomes were extracted for analysis as potential chimeric

sequences. To avoid including chimeric reads that resulted from technical artifacts

such as those caused by misinterpretation of open-pore states by base-calling

softwares (19), additional quality filtering was applied to the chimeric reads. The

distance between the mapped regions of the virus and the host sequence on the

chimeric reads was required to be less than 15 nucleotides, the junction was required

to be formed in the middle of the genes (not within the last 50 nucleotides of the first

gene sequence, nor the first 50 nucleotides of the second gene sequence), and the

quality score within 20 bp of either side of the junction was required to be higher than

the 20th percentile quality score for that read.

Mapping short reads to direct-RNA seq chimeric reads

We collected paired-end sequencing data on five samples with corresponding direct

RNA-sequencing data. The short reads were first preprocessed with fastp v0.23.1 (44)

and then mapped to the chimeric reads from the same samples by using Minimap2

v2.23 (42) with options "-ax sr -w 5" to tolerate the high error rate of the Nanopore

direct RNA-sequencing reads (45). Read pairs spanning the junctions were detected

and counted with a custom script. The numbers of read pairs supporting the chimeric

reads are provided in Supplementary Table 2.

Analysis of junction positions in relation to viral RNA secondary structure

The RNA secondary structure of the SARS-CoV-2 reference genome was obtained

from previous studies (46, 47) and bpRNA (48) was used to assign each residue to

secondary structure elements. A junction site was considered in the stem if the two

flanking nucleotides were in the same stem. To investigate if junctions tend to happen

in non-stem regions, the number of junctions occurring in base-paired positions were

calculated and compared with a background distribution for the numbers of junctions

located in stems derived from a 1000-time random sampling of the same number of

sites along the viral RNA strand. To further examine which types of structural

elements are over- or under-presented at junction sites in virus-host chimeric reads

and in host-virus chimeric reads, one-sided Fisher's exact test was performed.

Analysis of the expression level of host genes observed in chimeric reads

Gene expression profiles for two SARS-CoV-2 infected Caco-2 cell line samples

(GSM4477888, GSM4477889), two SARS-CoV-2 infected Calu-3 cell line samples

(GSM4477962, GSM4477963), and three SARS-CoV-2 infected Vero-6 cell line

samples (GSM4916368, GSM4916369, GSM4916370) were downloaded from the

GEO database. The read counts of each gene were normalized by the total number of

reads in each sample and by the gene length (RPKM) to represent the gene expression

level. The background gene set was composed of all expressed protein-coding genes

in the cell line. To evaluate whether the expression level of the host protein-coding

genes in chimeric reads is significantly greater than the expression level of the

background gene set, a one-sided Mann-Whitney U test was performed for each

sample.

Identification of insertions in SARS-CoV-2 genomes

The SARS-CoV-2 genomes available at GISAID (https://www.gisaid.org/) on

2021-12-17 were downloaded for analysis (n=6,163,073). The sequences were then

processed by NextClade CLI v1.7.0 (49) which generated a multiple sequence

alignment against the reference genome (Wuhan-Hu-1/2019) and provided a list of

single nucleotide polymorphisms, insertions, and deletions associated with each

genome sequence. Only sequences that passed all quality controls and were assessed

as "good" applied by NextClade were used for further analysis (n=5,226,229).

Insertions greater than or equal to 21 nucleotides long and found outside of the 5' and

3' untranslated regions of the viral genomes were kept. They were searched in the

NCBI non-redundant nucleotide database and a collection of coronavirus genomes

with BLASTN (E-value≤1e-2) (32) to explore their possible origins.

Monophyletic test

To check if the insertions of interest formed a monophyletic group, all genomes that

contained the same insertion were analyzed using UShER: Ultrafast Sample

placement on Existing tRee v0.5.1 (50) against a phylogenetic tree with available

genomes (n=6,257,569) from GISAID, GenBank, COG-UK and CNCB generated by

sarscov2phylo pipeline v.13-11-20 (51). The sequences are placed within an updated

global subsampled SARS-CoV-2 phylogenetic tree and local subtrees are computed to

show more sequences with the same context of the ones being analyzed.

Verification of insertions with raw sequencing data

The raw genome sequencing data of USA/WA-PHL-005726/2021 (EPI_ISL_6259191)

and USA/HI-H215617/2021 (EPI_ISL_6540096) were analyzed to check if the

insertion is indeed missing. The raw sequencing reads were processed for quality

control using fastp v0.23.1 (44) with default parameters and mapped to the

SARS-CoV-2 reference genome using BWA mem v0.7.17 (52). Primer sequences in

reads of EPI_ISL_6259191 were soft clipped using ivar trim (parameters: -m 1 -q 0 -s

4 -e) and reads in amplicons with variants in primer binding sites were removed by

ivar removereads v1.3.1 (53). The sequencing data of EPI_ISL_6540096 were

preprocessed by the providing laboratory and the primers were removed. Consensus

genome sequences were generated based on the alignments and it was found that the

consensus sequences did not contain the insertion. The 28S rRNA-derived insertion

was manually added at position 7120 of the consensus genomes to generate consensus

genomes with the insertion. The alignment files were converted to FASTQ format

using samtools fastq command v1.14 (54) and re-aligned to the consensus genomes

with or without the insertion using bowtie v2.4.4 (45) (parameter: --xeq). Reads

exclusively aligned to the consensus genome with the insertion and exclusively

aligned to the consensus genome without the insertion were identified with a custom

script

(https://github.com/ncbi/SARS2_host_derived_insertions/blob/main/verify_insertion/i

nsertion_match_reads.py).

Data availability

The datasets generated in this study and scripts are available in the github repository,

https://github.com/ncbi/SARS2_host_derived_insertions.

Competing interests: The authors declare that they have no competing interests.

Funding: All authors are supported by the Intramural Research Program of the NIH,

National Library of Medicine.

Authors' contributions: YY was involved in the execution of the analyses,

interpretation of the results, and writing and revision of the manuscript. KD was

involved in the interpretation of the results and writing and revision of the manuscript.

RF was involved in the execution of the analyses and writing of the manuscript. XJ

was involved in the conceptualization, planning, interpretation of the results, and

revision of the manuscript. All authors read and approved the final manuscript.

Acknowledgments: This work utilized the computational resources of the NIH HPC

Biowulf cluster (http://hpc.nih.gov). We would like to thank Eugene V. Koonin and

Sofya K. Garushyants for their thoughtful comments on our manuscript and their code

on how to perform permutation tests and plot the results provided at

https://github.com/garushyants/covid_insertions_paper. We gratefully acknowledge

the researchers from the originating laboratories responsible for obtaining the

specimens and the submitting laboratories where genetic sequence data were

generated and shared via the GISAID Initiative, on which this research is based

(Supplementary Table 6). We want to particularly thank the Washington State Public

Health Laboratories and the State of Hawaii Laboratories Division for sharing the raw

sequencing data for the genomes USA/WA-PHL-005726/2021 and

USA/HI-H215617/2021.

REFERENCES

1. Gerdol M, Dishnica K, Giorgetti A. 2022. Emergence of a recurrent insertion in the N-terminal domain of the SARS-CoV-2 spike glycoprotein. Virus Res:198674.

2. Garushyants SK, Rogozin IB, Koonin EV. 2021. Template switching and duplications in SARS-CoV-2 genomes give rise to insertion variants that merit monitoring. Communications biology 4:1-9.

3. Chrisman BS, Paskov K, Stockham N, Tabatabaei K, Jung J-Y, Washington P, Varma M, Sun MW, Maleki S, Wall DP. 2021. Indels in SARS-CoV-2 occur at template-switching hotspots. BioData Min 14:1-16.

4. Laiton-Donato K, Franco-Muñoz C, Álvarez-Díaz DA, Ruiz-Moreno HA, Usme-Ciro JA, Prada DA, Reales-González J, Corchuelo S, Herrera-Sepúlveda MT, Naizaque J. 2021. Characterization of the emerging B. 1.621 variant of interest of SARS-CoV-2. Infect Genet Evol 95:105038.

5. Venkatakrishnan A, Anand P, Lenehan PJ, Suratekar R, Raghunathan B, Niesen MJ, Soundararajan V. 2021. Omicron variant of SARS-CoV-2 harbors a unique insertion mutation of putative viral or human genomic origin.

6. Andreano E, Piccini G, Licastro D, Casalino L, Johnson NV, Paciello I, Dal Monego S, Pantano E, Manganaro N, Manenti A. 2021. SARS-CoV-2 escape from a highly neutralizing COVID-19 convalescent plasma. Proceedings of the National Academy of Sciences 118.

7. Peacock TP, Bauer DL, Barclay WS. Putative host origins of RNA insertions in SARS-CoV-2 genomes. Image (IN) 556:30.4.

8. Knoops K, Kikkert M, Worm SHvd, Zevenhoven-Dobbe JC, Van Der Meer Y, Koster AJ, Mommaas AM, Snijder EJ. 2008. SARS-coronavirus replication is supported by a reticulovesicular network of modified endoplasmic reticulum. PLoS Biol 6:e226.

9. Robson F, Khan KS, Le TK, Paris C, Demirbag S, Barfuss P, Rocchi P, Ng W-L. 2020. Coronavirus RNA proofreading: molecular basis and therapeutic targeting. Mol Cell 79:710-727.

10. Banner LR, Mc Lai M. 1991. Random nature of coronavirus RNA recombination in the absence of selection pressure. Virology 185:441-445.

11. Liao C, Lai M. 1992. RNA recombination in a coronavirus: recombination between viral genomic RNA and transfected RNA fragments. J Virol 66:6117-6124.

12. Yang Y, Yan W, Hall AB, Jiang X. 2021. Characterizing transcriptional regulatory sequences in coronaviruses and their role in recombination. Mol Biol Evol 38:1241-1248.

13. Simon-Loriere E, Holmes EC. 2011. Why do RNA viruses recombine? Nature Reviews Microbiology 9:617-626.

550   14.   Bolze A, White S, Basler T, Dei Rossi A, Roychoudhury P, Greninger AL,
551   Hayashibara K, Wyman D, Kil E, Dai H. 2022. Evidence for SARS-CoV-2
552   Delta and Omicron co-infections and recombination. medRxiv.
553   15.   Forni D, Cagliani R, Clerici M, Sironi M. 2017. Molecular evolution of human
554   coronavirus genomes. Trends Microbiol 25:35-48.
555   16.   Yan B, Chakravorty S, Mirabelli C, Wang L, Trujillo-Ochoa JL, Chauss D,
556   Kumar D, Lionakis MS, Olson MR, Wobus CE. 2021. Host-virus chimeric
557   events in SARS-CoV-2-infected cells are infrequent and artifactual. J Virol
558   95:e00294-21.
559   17.   Zhang L, Richards A, Barrasa MI, Hughes SH, Young RA, Jaenisch R. 2021.
560   Reverse-transcribed SARS-CoV-2 RNA can integrate into the genome of
561   cultured human cells and can be expressed in patient-derived tissues.
562   Proceedings of the National Academy of Sciences 118.
563   18.   Briggs E, Ward W, Rey S, Law D, Nelson K, Bois M, Ostrov N, Lee HH,
564   Laurent JM, Mita P. 2021. Assessment of potential SARS-CoV-2 virus
565   integration into human genome reveals no significant impact on RT-qPCR
566   COVID-19 testing. Proceedings of the National Academy of Sciences 118.
567   19.   Parry R, Gifford RJ, Lytras S, Ray SC, Coin LJ. 2021. No evidence of
568   SARS-CoV-2 reverse transcription and integration as the origin of chimeric
569   transcripts in patient tissues. Proceedings of the National Academy of
570   Sciences 118.
571   20.   Smits N, Rasmussen J, Bodea GO, Amarilla AA, Gerdes P, Sanchez-Luque FJ,
572   Ajjikuttira P, Modhiran N, Liang B, Faivre J. 2021. No evidence of human
573   genome integration of SARS-CoV-2 found by long-read DNA sequencing.
574   Cell Rep 36:109530.
575   21.   Zhang L, Richards A, Barrasa MI, Hughes SH, Young RA, Jaenisch R. 2021.
576   Response to Parry et al.: Strong evidence for genomic integration of
577   SARS-CoV-2 sequences and expression in patient tissues. Proc Natl Acad Sci
578   U S A 118.
579   22.   Shu Y, McCauley J. 2017. GISAID: Global initiative on sharing all influenza
580   data–from vision to reality. Eurosurveillance 22:30494.
581   23.   Wang D, Jiang A, Feng J, Li G, Guo D, Sajid M, Wu K, Zhang Q, Ponty Y,
582   Will S. 2021. The SARS-CoV-2 subgenome landscape and its novel
583   regulatory features. Mol Cell 81:2135-2147. e5.
584   24.   Wu H-Y, Brian DA. 2007. 5′ -Proximal Hot Spot for an Inducible
585   Positive-to-Negative-Strand Template Switch byCoronavirus RNA-Dependent
586   RNA Polymerase. J Virol 81:3206-3215.
587   25.   Kim D, Lee J-Y, Yang J-S, Kim JW, Kim VN, Chang H. 2020. The
588   architecture of SARS-CoV-2 transcriptome. Cell 181:914-921. e10.
589   26.   Gorski JL, Gonzalez IL, Schmickel RD. 1987. The secondary structure of
590   human 28S rRNA: the structure and evolution of a mosaic rRNA gene. J Mol
591   Evol 24:236-251.

592   27.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J,
593   Glöckner FO. 2012. The SILVA ribosomal RNA gene database project:
594   improved data processing and web-based tools. Nucleic Acids Res
595   41:D590-D596.

596   28.   Hagemeijer MC, Monastyrska I, Griffith J, van der Sluijs P, Voortman J, en
597   Henegouwen PMvB, Vonk AM, Rottier PJ, Reggiori F, De Haan CA. 2014.
598   Membrane rearrangements mediated by coronavirus nonstructural proteins 3
599   and 4. Virology 458:125-135.

600   29.   Lei J, Kusov Y, Hilgenfeld R. 2018. Nsp3 of coronaviruses: Structures and
601   functions of a large multi-domain protein. Antiviral Res 149:58-74.

602   30.   Khare S, Gurry C, Freitas L, Schultz MB, Bach G, Diallo A, Akite N, Ho J,
603   Lee RT, Yeo W. 2021. GISAID's Role in Pandemic Response. China CDC
604   Weekly 3:1049.

605   31.   Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C,
606   Sagulenko P, Bedford T, Neher RA. 2018. Nextstrain: real-time tracking of
607   pathogen evolution. Bioinformatics 34:4121-4123.

608   32.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K,
609   Madden TL. 2009. BLAST+: architecture and applications. BMC
610   Bioinformatics 10:1-9.

611   33.   Gopanenko AV, Malygin AA, Karpova GG. 2015. Exploring human 40S
612   ribosomal proteins binding to the 18S rRNA fragment containing major $3'$
613   -terminal domain. Biochimica et Biophysica Acta (BBA)-Proteins and
614   Proteomics 1854:101-109.

615   34.   Granneman S, Petfalski E, Swiatkowska A, Tollervey D. 2010. Cracking pre
616   ‐ 40S ribosomal subunit structure by systematic analyses of RNA−protein
617   cross‐linking. The EMBO journal 29:2026-2036.

618   35.   Cao Z, Xia H, Rajsbaum R, Xia X, Wang H, Shi P-Y. 2021. Ubiquitination of
619   SARS-CoV-2 ORF7a promotes antagonism of interferon response. Cell Mol
620   Immunol 18:746-748.

621   36.   Zhou Z, Huang C, Zhou Z, Huang Z, Su L, Kang S, Chen X, Chen Q, He S,
622   Rong X. 2021. Structural insight reveals SARS-CoV-2 ORF7a as an
623   immunomodulating factor for human CD14+ monocytes. IScience 24:102187.

624   37.   Su C-M, Wang L, Yoo D. 2021. Activation of NF-κB and induction of
625   proinflammatory cytokine expressions mediated by ORF7a protein of
626   SARS-CoV-2. Sci Rep 11:1-12.

627   38.   Volz E, Mishra S, Chand M, Barrett JC, Johnson R, Geidelberg L, Hinsley
628   WR, Laydon DJ, Dabrera G, O'Toole Á. 2021. Assessing transmissibility of
629   SARS-CoV-2 lineage B. 1.1. 7 in England. Nature 593:266-269.

630   39.   Gultyaev AP, Spronken MI, Funk M, Fouchier RA, Richard M. 2021.
631   Insertions of codons encoding basic amino acids in H7 hemagglutinins of
632   influenza A viruses occur by recombination with RNA at hotspots near
633   snoRNA binding sites. RNA 27:123-132.

634   40.   Khatchikian D, Orlich M, Rott R. 1989. Increased viral pathogenicity after
635   insertion of a 28S ribosomal RNA sequence into the haemagglutinin gene of
636   an influenza virus. Nature 340:156-157.
637   41.   De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. 2018.
638   NanoPack: visualizing and processing long-read sequencing data.
639   Bioinformatics 34:2666-2669.
640   42.   Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences.
641   Bioinformatics 34:3094-3100.
642   43.   Wu F, Zhao S, Yu B, Chen Y-M, Wang W, Song Z-G, Hu Y, Tao Z-W, Tian
643   J-H, Pei Y-Y. 2020. A new coronavirus associated with human respiratory
644   disease in China. Nature 579:265-269.
645   44.   Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ
646   preprocessor. Bioinformatics 34:i884-i890.
647   45.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2.
648   Nature methods 9:357-359.
649   46.   Cao C, Cai Z, Xiao X, Rao J, Chen J, Hu N, Yang M, Xing X, Wang Y, Li M.
650   2021. The architecture of the SARS-CoV-2 RNA genome inside virion.
651   Nature communications 12:1-14.
652   47.   Huston NC, Wan H, Strine MS, Tavares RdCA, Wilen CB, Pyle AM. 2021.
653   Comprehensive in vivo secondary structure of the SARS-CoV-2 genome
654   reveals novel regulatory motifs and mechanisms. Mol Cell 81:584-598. e5.
655   48.   Danaee P, Rouches M, Wiley M, Deng D, Huang L, Hendrix D. 2018. bpRNA:
656   large-scale automated annotation and analysis of RNA secondary structure.
657   Nucleic Acids Res 46:5381-5394.
658   49.   Aksamentov I, Roemer C, Hodcroft EB, Neher RA. 2021. Nextclade: clade
659   assignment, mutation calling and quality control for viral genomes. Journal of
660   Open Source Software 6:3773.
661   50.   Turakhia Y, Thornlow B, Hinrichs AS, De Maio N, Gozashti L, Lanfear R,
662   Haussler D, Corbett-Detig R. 2021. Ultrafast Sample placement on Existing
663   tRees (UShER) enables real-time phylogenetics for the SARS-CoV-2
664   pandemic. Nat Genet 53:809-816.
665   51.   Lanfear R, Mansfield R. 2020. A global phylogeny of SARS-CoV-2
666   sequences from GISAID. Zenodo doi 10.
667   52.   Li H, Durbin R. 2009. Fast and accurate short read alignment with
668   Burrows–Wheeler transform. Bioinformatics 25:1754-1760.
669   53.   Grubaugh ND, Gangavarapu K, Quick J, Matteson NL, De Jesus JG, Main BJ,
670   Tan AL, Paul LM, Brackney DE, Grewal S. 2019. An amplicon-based
671   sequencing framework for accurately measuring intrahost virus diversity using
672   PrimalSeq and iVar. Genome Biol 20:1-19.
673   54.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G,
674   Abecasis G, Durbin R. 2009. The sequence alignment/map format and
675   SAMtools. Bioinformatics 25:2078-2079.

FIGURE LEGENDS

Fig.1 Locations of the chimeric read junction sites and permutation tests for the number of junction sites in stems. Diagrams show how frequently junction sites occur at each position on the SARS-CoV-2 genome for **(A)** 5'-human-SARS2-3', **(B)** 5'-SARS2-human-3', **(C)** 5'-monkey-SARS2-3', and **(D)** 5'-SARS2-monkey-3' chimeric reads. Positions are colored based on the secondary structure of the SARS-CoV-2 RNA, with red lines indicating that the position is in the non-stem region, while gray indicates that the position is located in the stem region. Histograms following each diagram show the corresponding results of permutation tests used to test if the junction sites of chimeric reads are within base-paired regions of the viral RNA. Each test consists of 1000 permutations and the actual frequency of junction sites occurring in the stem regions is marked with a vertical red line.

Fig.2 The expression level of host genes observed in chimeric reads. The expression level of host protein-coding genes observed in chimeric reads is significantly higher than the background protein-coding gene expression level based on studies on **(A)** *Homo sapiens* Caco-2 cell line, **(B)** *Homo sapiens* Calu-3 cell line, and **(C)** *Chlorocebus sabaeus* Vero-6 cell line.

Fig.3 The 28S rRNA-derived insertion in SARS-CoV-2 genomes. (A) The phylogeny tree shows the genomes containing the human 28S-derived insertion. The clade where the insertion was detected is highlighted with a red box and the genomes with the insertion are marked with red circles at the tips. The asterisk (*) indicates that the insertion should be present in the variant based on raw sequencing data. **(B)** The insertion in SARS-CoV-2 genomes potentially originate from the host 28S rRNA shown by the sequence alignment of SARS-CoV-2 reference genome (NCBI accession: NC_045512.2, GISAID accession: China/Wuhan-Hu-1/2019) (pink), USA/CA-CDC-FG-169171/2021 (NCBI accession: OL591909.1, GISAID accession: EPI_ISL_6624703) (pink) and human 28S rRNA (chain A2 of PDB 5AJ0) (blue). There are five possible alignments for mapping this insertion to the reference. Only the alignment with the sequence inserted after the 3rd position of 2285th codon in ORF1ab is shown. The putative insertion origin is colored in red. The numbers listed above and below the alignment indicate the positions of aligned bases in the original sequences. The insertion sequence (red) was mapped to the 28s rRNA (blue) in a human polysome 3D structure (PDB: 5AJ0). A zoom-in view of the RNA secondary structure shows that the insertion is located on the No. 94 stem of domain 7 (position: 4969-4998) 28S rRNA region (highlighted red). **(C)** Diagram shows the position of the human 28S rRNA-derived insertion in the ectodomain (3Ecto) of Nsp3 protein.

Fig.4 The 18S rRNA-derived insertion in SARS-CoV-2 genomes. (A) The insertion in SARS-CoV-2 genomes potentially originates from the host 18S rRNA shown by the sequence alignment of SARS-CoV-2 reference genome (NCBI

accession: NC_045512.2, GISAID accession: China/Wuhan-Hu-1/2019) (pink),

USA/OR-OSPHL00675/2021 (GISAID accession: EPI_ISL_2339305) (pink) and

human 18S rRNA (purple). The putative insertion origin is colored in red. The

numbers listed above and below the alignment indicate the positions of aligned bases

in the original sequences. The insertion sequence (red) was mapped to the 18S rRNA

(purple) in a human polysome 3D structure (PDB: 5AJ0). A zoom-in view of the

RNA secondary structure shows that the insertion covers parts of helices 11 and 12 of

the 5' domain of the 18S rRNA. The location of the putative insertion sequence is

highlighted red. **(B)** Diagram shows the position of the human 18S-derived insertion

on the structure of the SARS-CoV-2 ORF7a protein (PDB: 7CI3). **(C)** The phylogeny

tree shows the genomes containing the human 18S-derived insertion. The clade where

the insertion was detected is highlighted with a red box and the genomes with the

insertion are marked with red circles.

SUPPLEMENTAL MATERIAL

Supplementary Tables

Supplementary Table 1: Direct RNA-seq data analysis. Metadata associated with all 30 of the analyzed directed RNA-seq samples is provided along with the number of reads mapped to the SARS-CoV-2 transcriptome and the count and frequency of chimeric reads in each sample. The counts of chimeric reads in the host to virus and virus to host orientation are listed for each sample. The references DOI for each sample is also listed.

Supplementary Table 2: Chimeric reads supported by spanning junction short reads.

Supplementary Table 3: Secondary structure element enrichment analysis of nucleotide sequences at junction sites.

Supplementary Table 4: Long insertions identified in GISAID that are not derived from SARS-CoV-2. The location on the SARS-CoV-2 reference genome, insertion sequence, insertion length, and what gene they are located in are provided for each of the 36 detected insertions. SARS-CoV-2 genomes with the insertion, whether those genomes were monophyletic, and short descriptions of putative matches are also provided for each insertion.

Supplementary Table 5: Information on genomes related to the three verified insertions. Metadata associated with each of the SARS-CoV-2 genomes with putative insertions that were analyzed including the variant types, collection dates, collecting labs, and sequencing methods are provided.

Supplementary Table 6: GISAID acknowledgement table.

A

AY.103

USA/WA-CDC-IBX436522269722/2021-10-09
USA/CA-CDC-FG-169171/2021-11-10
USA/WA-S14864/2021-10-11
USA/MA-CDC-LC0362672/2021-11-02
USA/HI-H215617/2021-10-21 *
USA/WA-PHL-005726/2021-10-25 *
USA/ID-IBL-770307/2021-11-02
USA/ID-CDC-IBX205486403071/2021-10-30

Tree scale 5

B

7091   7120   7121   7152

SARS-CoV-2 Wuhan-Hu-1   GTCACTATTGCAACCTACTGTACTGGTTCT--------------------ATACCTTGTAGTGTTTGTCTTAGTGGTTTAGA

SARS-CoV-2 EPI_ISL_6624703   GTCACTATTGCAACCTACTGTACTGGTTCTTAGCAGAGCAGCTCCCTCGCTGCGATCTATATCTTGTAGTGTTTGTCTTAGTGGTTTAGA

Human 28S rRNA

4924   4969   4998   5029

AAACCATTCGTAGACGACCTGCTTCTGGGTCGGGGTTTCGTACGTAGCAGAGCAGCTCCCTCGCTGCGATCTATTGAAAGTCAGCCCTCGACACAAGGGTTTGTC

28S rRNA

5.8S rRNA

5S rRNA

18S rRNA

PDB: 5AJ0

C

Insertion at position 1467 **SRAAPSLRS**

1437   1518

3Ecto

ER lumen

TM   TM

Cytoplasm

N-terminal   Nsp3   C-terminal

1945

```
                27463                          27492              27493                       27523
SARS-CoV-2 Wuhan-Hu-1     TAGAGGTACAACAGTACTTTTAAAAGAACC------------------------TTGCTCTTCTGGAACATACGAGGGCAATTCA
                          ||||||||||||||||||||||||||||||                        |||||||||||||||||||||||||||||||
SARS-CoV-2 EPI_ISL_2339305 TAGAGGTACAACAGTACTTTTAAAAGAACCCACGGGTGACGGGGAATCAGGGTTTTGCTCTTCTGGAACATACGAGGGCAATTCA

Human 18S rRNA                                            |||||||||||||||||||||||||||
                    357                          CCTACCATGGTGACCCACGGGTGACGGGGAATCAGGGTTCGATTCCGGAGAGGGAGCCTGAGAAACGGCTACC  457
                    CCTATCAACTTTCGATGGTAGTCGCCGTGCCTA            399                   423
```

28S rRNA

5.8S rRNA

18S rRNA

PDB: 5AJ0

B

disulfide bond

CYS-67

CYS-35

Insertion at position 34

TGDGESGF

PRO-34

beta sheet C

beta sheet B

PDB: 7CI3

ORF7a

C

C7125T

    USA/OR-OSPHL00675/2021-05-06
    USA/OR-OHSU-211930214/2021-05-05
    USA/OR-CDC-ASC210075576/2021-04-19

C29095T
    USA/OR-CDC-LC0056173/2021-04-30
    USA/OR-OHSU-211740128/2021-04
    USA/OR-OHSU-211790160/2021-04

C12473T

G1458A
    USA/ND-NDDH-01971/2021-05-22
    USA/ND-NDDH-02292/2021-06-05
C29253T
    USA/ND-NDDH-02143/2021-05-24
    USA/ND-CDC-2-4524396/2021-05-25
    USA/ND-CDC-2-4524396/2021-05-25

C21597T
    USA/MN-MDH-4707/2021-03-26
    USA/MN-MDH-4858/2021-03-26
    USA/MN-UMGC-2120/2021-04-01

    USA/MN-CDC-IBX346610039626/2021-03-21
    USA/MN-CDC-IBX290234783599/2021-03-24

Tree scale: 1