Title: Improving cassava bacterial blight resistance by editing the epigenome

Authors: Kira M. Veley[1], Kiona Elliott[1,2], Greg Jensen[1], Zhenhui Zhong[3], Suhua Feng[3,4], Marisa Yoder[1], Kerrigan B. Gilbert[1], Jeffrey C. Berry[1], Zuh-Jyh Daniel Lin[1], Basudev Ghoshal[3]†, Javier Gallego-Bartolomé[3]‡, Joanna Norton[4], Sharon Motomura-Wages[4], James C. Carrington[1], Steven E. Jacobsen[3,5,6]*, Rebecca S. Bart[1]*

Affiliations:

[1]Donald Danforth Plant Science Center; Saint Louis, MO 63132, USA.

[2]Division of Biological and Biomedical Sciences, Washington University in Saint Louis, Saint Louis, MO, 63110, USA.

[3]Department of Molecular, Cell and Developmental Biology, University of California at Los Angeles; Los Angeles, CA, 90095, USA.

[4]College of Tropical Agriculture & Human Resources, University of Hawaii at Manoa, Hilo, HI, 96720, USA.

[5]Eli & Edythe Broad Center of Regenerative Medicine & Stem Cell Research, University of California at Los Angeles; Los Angeles, CA, 90095, USA.

[6]Howard Hughes Medical Institute, University of California at Los Angeles; Los Angeles, CA, 90095, USA.

*Co-corresponding authors. Email: SEJ: jacobsen@ucla.edu; RSB: rbart@danforthcenter.org

†Current address: Summerland Research and Development Centre, Agriculture and Agri-Food Canada, Summerland, BC, V0H 1Z0, Canada.

‡Current address: Instituto de Biología Molecular y Celular de Plantas (IBMCP), CSIC-Universidad Politécnica de Valencia, 46011 Valencia, Spain.

Abstract

Pathogens rely on expression of host susceptibility (S) genes to promote infection and disease.

As DNA methylation is an epigenetic modification that affects gene expression, blocking access

to S genes through targeted methylation could increase disease resistance. *Xanthomonas phaseoli*

pv. *manihotis*, the causal agent of cassava bacterial blight (CBB), uses transcription activator-

like20 (TAL20) to induce expression of the S gene *MeSWEET10a.* In this work, we direct

methylation to the TAL20 effector binding element within the *MeSWEET10a* promoter using a

synthetic zinc-finger DNA binding domain fused to a component of the RNA-directed DNA

methylation pathway. We demonstrate that this methylation prevents TAL20 binding, blocks

transcriptional activation of *MeSWEET10a in vivo* and that these plants display decreased CBB

symptoms while maintaining normal growth and development. This work therefore presents an

epigenome editing approach useful for crop improvement.

Introduction

Susceptibility (S) genes present a liability within a host genome, as their exploitation is a

common mechanism used by diverse classes of pathogens[1]. However, mutating these loci may

have negative consequences as these genes are often necessary for normal plant growth and

reproductive development[2–5]. In many cases, pathogens actively upregulate expression of S

genes during infection. Plant pathogenic bacteria within the *Xanthomonas* and *Ralstonia* genera

express transcription activator-like (TAL) effectors that activate transcription of S genes by

recognizing and binding specific, largely predictable, promoter sequences (effector binding

elements, EBEs) within the host genome[4,6–9]. Previous research demonstrated that gene editing

can be used to prevent the exploitation of S genes by pathogens and that this results in increased

resistance[10–14]. Epigenomic modifications can also influence gene expression. Because TAL

binding can be inhibited by DNA methylation[15,16], we hypothesized that targeting methylation to

a TAL effector binding site would block induction of the S gene and result in increased

resistance as measured by decreased disease symptoms.

In eukaryotes, cytosine DNA methylation is an important mechanism of epigenetic gene

regulation[17]. In plants, the RNA-directed DNA methylation (RdDM) pathway is responsible for

establishing *de novo* 5-methylcytosine methylation. Previous targeted methylation efforts

leveraged naturally occurring epialleles from Arabidopsis that affect flowering time[18,19]. These

studies used CRISPR-Cas9-based tools combined with components of the methylation pathways,

either the catalytic domain of the plant methyltransferase DOMAINS REARRANGED

METHYLTRANSFERASE 2 (DRM2) involved in RdDM or the bacterial CG-methyltransferase

MQ1[20,21]. In addition, a number of RdDM proteins are effective in targeting methylation when

fused with artificial zinc-fingers (ZFs), including DRM2 and DEFECTIVE IN MERISTEM

SILENCING 3 (DMS3), a protein that in Arabidopsis targets RNA polymerase V binding to

chromatin to trigger RdDM[22,23]. While CRISPR-based tools are highly specific, ZF-based

targeting systems are preferable for some applications, as they generally require smaller

transgenes and are not constrained by the location of PAM sites.

Cassava bacterial blight (CBB) is caused by *Xanthomonas phaseoli* pv. *manihotis*[24]

(*Xam*, aka. *Xanthomonas axonopodis* pv. *manihotis)* and occurs in all regions where cassava

(*Manihot esculenta* Crantz) is grown[25,26]. Recently, a resistance gene was described that may

provide some tolerance to certain strains of Xam[27], however, currently, there are no strong broad

spectrum resistance genes available to breeders. Symptoms of an infection begin with dark

green, angular, "water-soaked" lesions on leaves (Fig. 1A) after which *Xam* moves through the

vasculature causing leaf wilting and eventual death of the plant. *Xam* secretes 20 - 30 effectors,

including multiple TAL effectors, into cassava cells[28]. We previously demonstrated that one of

these TAL effectors, TAL20, is a major virulence determinant for this pathogen. TAL20 induces

ectopic S gene expression of a SWEET (Sugars Will Eventually be Exported Transporters)

family sugar transporter, *MeSWEET10a*[6,29]. Expression of *MeSWEET10a* during infection is

required for water-soaking phenotypes and full virulence[6]. In wild-type cassava, the promoter of

MeSWEET10a is not methylated[30]. Thus, we hypothesized that targeted methylation at the

TAL20 EBE would prevent ectopic expression of *MeSWEET10a* and provide resistance to CBB

(Fig. 1B).

Here, we present epigenetic editing as a useful approach toward crop improvement. We

show directed DNA methylation to the TAL20 EBE within the *MeSWEET10a* promoter using a

ZF fused to DMS3. We demonstrate that cytosine methylation inhibits TAL20 binding to the

EBE, preventing transcriptional activation of *MeSWEET10a* and decreasing bacterial blight

symptoms in cassava plants.

Results

Establishing methylation at the *MeSWEET10a* promoter

We first tested the effect of methylation on TAL20 binding to the EBE *in vitro* using an

Electrophoretic Mobility Shift Assay (EMSA) (Supplementary Fig. 1). As shown previously[6],

purified 6xHis-TAL20$_{Xam668}$ binds the unmethylated EBE oligo (lane 2) and, as expected,

providing an excess of unlabeled, unmethylated oligo outcompeted for binding (lane 5).

Conversely, 6xHis-TAL20$_{Xam668}$ does not efficiently bind the methylated oligo (lane 3), nor did

the methylated oligo compete for TAL20 binding, even when supplied in excess (lanes 6 and 7).

Therefore, the EBE of *MeSWEET10a* was deemed a good candidate for engineering epigenetic

disease resistance. Of the RdDM factors tested in *Arabidopsis thaliana*, DMS3 was shown to be

the most effective at delivering methylation to a given locus[23]. Transgenic cassava plants were

generated expressing an artificial ZF targeting the EBE of *MeSWEET10a* fused to Arabidopsis

DMS3 (Fig. 2A, Supplementary Figs. 2-7). The targeted region of the *MeSWEET10a* promoter

was assessed for methylation using amplicon-based bisulfite sequencing (ampBS-Seq) (Fig. 2B,

Supplementary Figs. 2-7). Plants expressing the ZF alone showed no methylation at or around

the EBE of *MeSWEET10a*. In contrast, all plants expressing DMS3-ZF contained high levels of

ectopic CpG methylation within and near the ZF target, including the only two CpG sites within

the EBE (Fig. 2B, Supplementary Figs. 2-7). The specific analysis that the plants were used for

in each Supplementary Fig. 2-7 are indicated in the title of the figure. Plant phenotypes were

evaluated under greenhouse and field conditions using our standard methods (see methods for

details). These analyses indicated that methylation did not grossly affect root and shoot growth

and development (Fig. 2C, D, Supplementary Fig. 8). Whole Genome Bisulfite Sequencing

(WGBS) was used to assess specificity of the DMS3-ZF. High levels of *de novo* DNA

methylation were observed within the target region, similar to the ampBS-seq result (Fig. 2B,

Fig. 3A), while DNA methylation within an extended region of 5 kb on either side was

comparable between the DMS3-ZF-expressing lines and the ZF alone control (Fig. 3A, B).

Additionally, CG, CHG, and CHH methylation across all chromosomes, as well as within genes,

was similar between the DMS3-ZF-expressing lines and ZF alone control (Fig. 3C, D,

Supplementary Fig. 9, and Supplementary Table 1). We investigated potential off-targets by

predicting ZF binding sites with up to 4 mismatches within the target sequence (5'-

CGCTTCTCGCCCATCCAT - 3'). This yielded 1,542 potential off-target regions genome-wide

(Supplementary Fig. 10A). We compared methylation levels of these potential off-targets, plus

100bp flanking sequences, in WT and the two DMS3-ZF-expressing lines 133 and 204. No

significant differences were detected (Supplementary Fig. 10B). As a complementary approach

to identify potential non-specific activity, differentially methylated regions (DMRs) (CG

variation > 0.4, CHG variation > 0.2, and CHH variation > 0.1, respectively) were obtained by

comparing the two DMS3-ZF lines with the WT control. This method successfully detected two

hypermethylated DMRs in the targeted MeSWEET10a region (Fig. 3A). However, most of the

other detected DMRs were hypoDMRs (i. e. regions showing decreased methylation in the

DMS3-ZF lines) rather than hyperDMRs (increased methylation in the DMS3-ZF lines). An

overall decrease in genome methylation has been attributed to the plant tissue culture process in

cassava[31] and in many other plant species including maize[32], rice[33] and oil palm[34]. Cross

referencing the observed DMRs with the 1,542 potential off-targets revealed that very few of

DMRs (particularly hyperDMRs) fell into off-target regions, and the chance was

indistinguishable from the random control regions (Supplementary Fig. 10C, Supplementary

Table 2). Together, these results demonstrate a reasonable level of specificity for targeted

DMS3-ZF-mediated methylation in cassava.

131   Cytosine methylation prevents *MeSWEET10a* transcriptional activation by TAL20 *in vivo*

We next tested the hypothesis that CpG methylation at the TAL20 EBE would prevent

activation of *MeSWEET10a in vivo*. Compared to controls, plants from independent DMS3-ZF-

expressing lines showed a significant lack of *MeSWEET10a* induction in response to *Xam*

infection (Fig. 4A). The suppression of *MeSWEET10a* induction occurred in independently

derived transgenic lines and across independent experiments. This suppression did not correlate

with increased expression of DMS3-ZF (Supplementary Fig. 11). These data indicate that

methylation prevents TAL20-mediated activation of *MeSWEET10* expression.

139   Epigenomic editing at *MeSWEET10a* decreases cassava bacterial blight symptoms

To quantify the effect of methylation at the *MeSWEET10a* EBE on CBB disease *in vivo*,

we measured "water-soaked" lesions (Fig. 1A, Fig. 4B, Supplementary Fig. 12), which are the

most obvious early disease symptom and requires induction of *MeSWEET10a*[6,35]. Quantitative

analysis of images from *Xam*-infiltrated leaves revealed that plants with methylation at the EBE

had a reduced infected area compared to wild-type (Fig. 4C). Further, the area of infection in

DMS3-ZF leaves infiltrated with wild-type *Xam* was comparable to leaves infected with *Xam*

lacking TAL20 (XamΔTAL20). No significant difference in bacterial population was observed in

the DMS3-ZF lines (Supplementary Fig. 13). In addition to a decrease in lesion size, the intensity

of water-soaking at and around the infiltration site was also reduced in EBE methylated plants.

Grey-scale values from images of lesions were compared to mock-inoculated sites on the same

leaf, where more water-soaking equates to a darker appearance. Water-soaking intensity was

reduced in plants with methylation at the EBE (Fig. 4D). These results demonstrate that

methylation of the EBE site within the promoter of the S gene *MeSWEET10a* results in a

reduction in CBB-related symptoms.

Discussion

In summary, we show that DMS3 paired with an artificial ZF caused targeted, *de novo*

establishment of methylation at the EBE of *MeSWEET10a,* blocking TAL20-induced ectopic

expression of *MeSWEET10a* and resulted in decreased CBB disease symptoms. Cassava is

typically clonally propagated, and we have monitored methylation over time throughout the

course of this study (three-four years maintenance in tissue culture and in the greenhouse). We

conclude that the methylation is stable in the presence of the transgene through clonal

propagation. Because the components of the technology presented here do not derive from

cassava (Arabidopsis DMS3 paired with a synthetic zinc finger), it is reasonable to hypothesize

that this approach may be similarly successful in other crop species. Gene editing represents an

existing, powerful strategy that can achieve similar outcomes to the work presented here, and we

expect this will continue to be the favored approach for many applications[5,10–14,36–40]. At the same

time, as targeted methylation technologies continue to mature, there will likely be cases where it

may be preferred. For example, DNA methylation may be applied as a gene expression dial,

opposed to an on/off switch. Similarly, as DNA methylation does not change the DNA sequence,

it may be possible to use DNA methylation changes to achieve expression patterns in a tissue- or

environment-specific manner. This might be achieved in transgenic plants by driving the

methylation reagents using a tissue- or environment-specific promoter. Additionally, a key near-

term future goal will be to study the heritability of DNA methylation in the absence of the

targeting transgene[41,42]. It seems likely that the size of the region with de novo methylation, the

type of methylation (CG, CHG, CHH), and the specific location in the genome may all affect

heritability. These factors will each need to be investigated, and while there is value in pursuing

this work in the crop of interest, we expect that much can be learned using model systems.

Ultimately, the practicality of this strategy will be determined by the heritability and stability of

epigenetic edits, and this is likely to be application- and target-specific. Nonetheless, the

transgenic epigenetic approach shown here may be useful in a variety of other crop improvement

programs.

Methods

<u>Construct design and cloning</u>

The genomic coding sequence of Arabidopsis *DMS3* (AT3G49250), including promoter,

was cloned into the pEG302 backbone as described[23,44]. The ZF targeting the EBE of

MeSWEET10a was designed as previously described[22] to target the sequence 5'-

CGCTTCTCGCCCATCCAT-3'. CodeIT (zinc.genomecenter.ucdavis.edu:8080/Plone/codeit)

was used to generate the amino acid in silico (linker type "normal"). The resulting amino acid

sequence was plant codon optimized, synthesized by IDT and cloned at the 3' end of *DMS3*

(AT3G49250) in pEG302. A hygromycin resistance cassette was cloned into an available PmeI

site for selection in cassava. Vector sequences and maps are available in supplementary

materials.

<u>Production of plants and growth conditions</u>

The transgenic cassava line expressing the ZF with no methylating factor was generated for

a different project, and thus a different background was used: 60444 (ZFΔDMS3) versus

TME419 (DMS3-ZF). The promoter region of interest is identical in both genetic backgrounds,

as verified by Sanger sequencing (Extended Data Fig. 14). Transgenic cassava lines of cultivar

TME419 and 60444 were generated and maintained *in vitro* as described previously[45] with some

modifications, as follows: transgenic FEC cells were selected for resistance using hygromycin

(20 mg / ml) on spread plates (also containing 125 mg / L cefotaxime). On stage 1, 2, and 3

plates, 20 mg / ml hygromycin was used as well. Growth chamber settings used for tissue

culture: 28°C +/- 1°C, 75 μmol·m^{-2}·s^{-1} light; 16 hrs light / 8 hrs dark. *In vitro* plantlets were

propagated and established in soil on a misting bench with domes. Once established, plants were

removed from the misting bench and acclimated to greenhouse conditions (28°C; humidity

>50%; 16 hrs light / 8 hrs dark; 1000 W light fixtures supplementing natural light levels below

400 W / m^2). 1000 W light fixtures supplementing natural light levels below 400 W / m^2). For

field-grown plants, individuals were first established in the greenhouse (natural light and

temperature, 17-28°C, humidity >50%) for 29 weeks then transitioned outside and planted 3 feet

apart (field location 19.640, -155.083).

<u>PCR bisulfite sequencing (ampBS-seq) library preparation and analysis</u>

PCR-based BS-seq (ampBS-seq) of the *MeSWEET10a* promoter was performed as

described[21]. Leaf tissue and DNA extracted using the GenElute™ Plant Genomic DNA Miniprep

Kit (Sigma) and sodium bisulfite treated using the Epitect Bisulfite Conversion kit (QIAGEN).

The region of interest within the *MeSWEET10a* promoter was amplified using the primers in

Extended Data Table 3, giving a 233bp product. Libraries were generated from purified PCR

products amplified from the bisulfite-treated DNA, using an Ovation Ultralow V2 kit (NuGEN)

or a Kapa DNA HyperPrep Kit (Roche) in combination with IDT UD barcodes (Illumina). iSeq

100 (Illumina) was used to sequence the libraries. Raw sequencing reads were aligned to the

region of interest using the epigenomic analysis plugin for CLC Genomics Workbench 20.0

(QIAGEN). Reads shorter than 125 bp or longer than 152 bp were discarded. Reads were

mapped to the following sequence:

ATATTGTTTTTTTTTTTTTAAAAAAAATAATAAAAGAAATAAGGTTACTGTTACATT

GACATATTTTATTCACTTTAATCATGCATGCAACTTGACTTCATTCCGTTCCCTGGAT

TCCTCCCCTATATAAACGCTTCTCGCCCATCCATCATTGCACAACATAGCTAGAGTTT

CCTCTTGAGAAAGAGAGTTTCCTCTGCATAAGGGAAAGAGAGTTTTTATTATAGTYG

GA. For mapping, the following parameters were used: non-directional, no masking, match score

= 1, mismatch cost = 2, affine gap cost, insertion open cost = 6, insertion extend cost = 1,

deletion open cost = 6, deletion extend cost = 1, length fraction = 0.9, similarity fraction = 0.9.

Methylation was called for cytosines with greater than 500 reads mapped and the resulting

methylation level was plotted using the provided script (MeSWEET10a_07272021-

for_publication.0920.R).

<u>Whole Genome Bisulfite sequencing (WGBS) library preparation and analysis</u>

Genomic DNA from samples were end-repaired and ligated with TruSeq LT DNA single

adapters (Illumina) using Kapa DNA HyperPrep kit (Roche). Adapter-ligated DNA was

converted with EpiTect Bisulfite Conversion Kit (Qiagen). Converted DNA was PCR-amplified

by MyTaq polymerase (Bioline) for 12 cycles. Library quality and size were determined by

D1000 ScreenTape (Agilent) and libraries were subsequently purified by AMPure XP beads

(Beckman Coulter). Library concentrations were measured with Qubit dsDNA Broad-Range

Assay kit (ThermoFisher). Libraries were sequenced on NovaSeq 6000 sequencer (Illumina).

WGBS reads were mapped to the TME7 phase0 genome with BSMAP (v2.90) allowing 1

mismatch and 1 best hit (-v 1 -w 1)[46,47]. Duplicated reads were removed with SAMtools

(v1.3.1)[48]. Reads with three or more consecutive methylated CHH sites were considered as

unconverted reads and removed in the following analysis. Conversion rate was estimated by

calculating methylation level of the chloroplast genome. DNA methylation level at each cytosine

was calculated by the number of methylated C vs. total C and T account. Methylation levels of

each chromosome were determined by calculating CpG, CHG, and CHH level of 100 kb bins.

Differentially Methylated Regions (DMRs) were called using BSMAP (v2.90) with $p < 0.01$,

where the differences in CG, CHG, and CHH methylation were at least 0.4, 0.2, and 0.1,

respectively. Methylation track files were visualized with Integrative Genomics Viewer (IGV,

v3.0).

<u>Off-target analysis</u>

ZF off-target sites were predicted by Cas-OFFinder by searching cassava genome with target site

(5'-CGCTTCTCGCCCATCCAT-3') allowing 4 mismatches and no PAM site[49]. Methylation

levels of off-target sites and 100 bp flanking sequences (50 bp upstream and 50 bp downstream)

were compared between samples. Sequence consensus analysis of potential off-target sites was

performed with WebLogo (v3.6.0)[50].

<u>SDS-PAGE gel and western blotting</u>

All individuals used for qPCR and disease assay experiments were checked for construct

expression (Extended Data Figs. 3 (qPCR1), 4 (qPCR2), 5 (Water-soaking), 6 (qPCR3), 7

(Water-soaking, qPCR4), 8 (Water-soaking)). Tissue (100 mg) was collected from each

individual plant to be used in an experiment in a 1.5 ml microcentrifuge tube and frozen in liquid

nitrogen along with four 3mm glass beads. A tissue lyser (TissueLyser II, QIAGEN) was used to

homogenize the sample (30/second frequency, 2 minutes, performed twice with sample rotation

between runs). 400 ul of 1x Laemmli Sample buffer (.0625M Tris-HCl pH 6.8, 2% SDS, 10%

glycerol, 100 mM DTT) was added to each sample, then samples were boiled 90-100°C for 4

min, centrifuged for 2 min and then either run immediately on a gel or frozen for storage at -

80°C. For SDS-PAGE, 10ul of each sample was run on a 12% TEO-Tricine Precast Gel

(RunBlue™, AbCam, ab119203) then transferred to PVDF membrane (ThermoFisher

Scientific). For western blotting, monoclonal ANTI-FLAG® M2-Peroxidase (HRP) antibody

(A8592, Sigma) was used (1:2000 dilution in 5% milk in TBST). HRP activity was assessed

using SuperSignal™ West Femto substrate (ThermoFisher Scientific). Chemiluminescence was

detected using the ChemiDoc XRS+ System equipped with Image Lab 6.1 software (Bio-Rad).

The molecular weight of protein bands was estimated by comparison with the BenchMark™ Pre-

stained Protein Ladder (Thermo Fisher Scientific). The abundance of the 3xFLAG-ZF proteins

were quantified from the band intensities also using Image Lab Software. Coomassie staining

was performed on membranes using Coomassie brilliant blue (0.025% (w/v) Coomassie brilliant

blue R-250 (ThermoFisher Scientific) in 40% methanol/7% acetic acid (v/v)). Staining was

performed for 3 min followed by destain (solution: 50% methanol/7% acetic acid (v/v)) for 30

min. Then membranes were rinsed with water and allowed to dry overnight before imaging.

Electrophoretic mobility shift assay (EMSA)

For protein purification, $TAL20_{Xam668}$ was cloned into pDEST17 (Thermo Fisher Scientific)

using Gateway^{TM} technology, resulting in the addition of a 6xHis-tag at the N-terminus of the

protein upon expression. 6xHis-TAL20$_{Xam668}$ was expressed and purified from *E. coli* BL21 cells

using HisPur™ Ni-NTA Spin Columns (Thermo Fisher Scientific). Protein size and purity was

assessed by SDS-PAGE (4-12% gradient gel) followed by western blot (Extended Data Fig. 1A)

using an anti-His antibody (His-Tag (27E8) Mouse mAb (HRP Conjugate), Catalog #9991, Cell

Signaling Technology). For the EMSA, the sequence previously shown to be able to be bound by

TAL20$_{Xam668}$ [6], with and without methylation, was synthesized (Integrated DNA Technologies)

and biotin-labeled (Pierce™ Biotin 3' End DNA Labeling Kit, Thermo Fisher Scientific). The

sequence used is as follows, with methylated cytosines indicated by underlining: sense 5'-

CCCTATATAAA<u>C</u>G<u>C</u>TT<u>C</u>T<u>C</u>G<u>CCC</u>AT<u>C</u>CATCATT-3', anti-sense: 3'-

GGGATATATTTG<u>C</u>GAAGAG<u>C</u>GGGTAGGTAGTAA-5'. EMSA was performed with the

Light Shift® Chemiluminescent EMSA Kit (Thermo Fisher Scientific) according to the

manufacturer's protocol. Biotin-labeled oligonucleotides were mixed with 0, 20, 200, 300 and

1000 fmol of purified TAL20$_{Xam668}$ effector protein as indicated in the chart in Extended Data

Fig. 1B. All binding reactions were incubated at room temperature for 20 min. For competition

assays, unlabeled oligonucleotides were pre-incubated with other reaction components for an

additional 20 min. Gel electrophoresis was performed on a 6% native polyacrylamide gel.

<u>Bacterial inoculations</u>

Xanthomonas strains were grown on plates containing necessary antibiotics for 2-3 days at

30°C. The *Xanthomonas* strains used and their antibiotic resistances are as follows: 1. *Xam668*

(rifampicin 50 µg / ml) and 2. *Xam668ΔTAL20* (suicide vector knockout[6], tetracycline 5 µg / ml,

rifampicin 50 µg / ml). Bacteria were scraped from the plates into 10 mM MgCl$_2$ to either OD$_{600}$

= 1.0 (RT-qPCR) or OD$_{600}$ = 0.01 (water-soaking) concentrations according to established lab

protocols. Leaves (2-3 weeks after plants were transferred to a greenhouse) were inoculated with

a 1.0 ml needleless syringe using one bacterial strain per lobe, three injection points on each of

2-3 leaves. After inoculation, plants were kept under fluorescent light (12 hr day / night light

cycle) at 27°C for either 48 hours (RT-qPCR and ampBS-seq) or 4 days (water-soaking).

RNA extraction and RT–qPCR analysis

For RNA, 48 hours post-infiltration, samples were taken using an 8 mm cork borer at each

infiltration site, frozen in liquid nitrogen, ground to a fine powder (3 mm glass beads added to

tissue, ground using tissue lyser (TissueLyser II, QIAGEN), 30/second frequency, 2 minutes,

performed twice with sample rotation between runs) and total RNA was extracted (Spectrum™

Plant Total RNA Kit, Sigma). Technical replicates were pooled, thus each RNA sample

contained three injection sites from one leaf. 1.0 µg DNase-treated RNA was reverse-transcribed

to cDNA using SuperScript™ III Reverse Transcriptase (Thermo Fisher Scientific) according to

the manufacturer's instructions.

Quantitative reverse transcription PCR (RT-qPCR) was done with SYBR Select Master Mix

(Thermo Fisher Scientific) on a CFX384 Touch Real-Time PCR Detection System (Bio-Rad).

Primers specific for cassava GTPb (Manes.09G086600) and PP2A4 (Manes.09G039900) genes

were used as the internal controls to calculate relative expression levels (Table S1)[51,52].

Normalized relative expression was calculated using formulas described[53].

Water-soaking area and intensity quantification, image analysis

Cassava leaves were infiltrated with either 10 mM $MgCl_2$ alone (mock) or containing

Xanthomonas (*Xam668* strains with and without TAL20) as described above and imaged using a

Canon EOS Rebel T5i camera with a 15-85mm lens at 0- and 4-days post-inoculation. Images

were grey balanced using a X-Rite Passport by estimating the saturation of the six grey chips and

lowering the brightness accordingly. Gray corrected images were uploaded to FIJI[54] and

duplicated. In order to define the infected area, visible water-soaking spots were manually

outlined on the duplicate image using the pencil tool (color: #ff00b6 and size 2). The outlined

images were converted from RGB to LAB and split to obtain the A color channel. The A channel

images were thresholded, converted to a mask and the mask for each spot was added to the ROI

manager using the analyze particle tool. The ROI masks were applied to the original RGB grey

corrected images. In order to define the background color of a given leaf, mock-infiltrated spots

(no water-soaking) were added to the ROI manager using an arbitrarily sized rectangle selection

tool consistently set to a W = 51 and H = 61. Area and grey scale mean data were obtained for

each infiltrated spot using the FIJI measure tool. Grey balancing using the X-Rite Passport is a

coarse adjustment and finer standardization is performed post image processing using statistical

methods. Grey value standardization was achieved by estimating the grand mean of all grey

values in each image and centering those values to the grand mean of all images by using model

residuals. Further, the residuals within each image were re-centered with respect to the mock

treatment by subtracting the mean grey value of mock from all other features.

Bacterial growth assay

Cassava leaves were infiltrated with either 10 mM $MgCl_2$ alone (mock) or containing

Xanthomonas (*Xam668* strains with and without TAL20) as described above. Leaf punch

samples were taken at the site of infiltration using a 5 mm cork borer at 0- and 4-days post-

inoculation. For day-0 samples, infiltrated spots were allowed to dry down prior to processing.

Individual leaf punches were transferred to 2 mL Eppendorf Safelock tubes with 200 ul of 10

mM $MgCl_2$ and three disposable 3 mm Propper solid glass beads. Samples were ground with a

Qiagen Tissuelyzer at 28 hZ for 3 minutes. 200 ul of the ground sample were transferred to the

first column of a labelled 96-well plate. Serial dilutions were performed by transferring 20 ul of

the non-dilute sample (10^1) to the next well containing 180 ul of 10 mM $MgCl_2$. Samples were

serially diluted to 10^4 for day-0 and 10^6 for day-4. 10 ul of each serial dilution was spread onto

labelled quadrants of a NYG plate with cycloheximide and the appropriate antibiotics for the

infiltrated bacterial strain. Plates were incubated at 30°C for 2-3 days, and the number of

colonies were counted. Colony forming Units (CFU) reported in this manuscript were

transformed by sample area (CFU/cm^2 where $cm^2 = 0.52$).

<u>Storage root production and measurements</u>

Storage root production and analysis was conducted with some modification as

described[43]. Briefly, plants were maintained in small, 2-inch pots for 11 weeks in order to induce

storage root production. The same greenhouse conditions were used as described above. After 11

weeks, roots were removed from pots, washed to remove soil, and storage roots isolated from the

root mass and weighed for fresh weight measurements. For starch content estimation

(approximately 80% of storage root weight), the storage roots were peeled, cut into

approximately 2 cm cubes, placed into a 50 ml conical tube, and frozen in dry ice and stored at -

80°C prior to lyophilization. Samples were lyophilized for two days, then weighed to obtain dry

weight.

Data availability

All data and constructs are available on request. All amplicon-based bisulfite sequencing data,

western blots, and images used for imaged-based analysis of water-soaking disease symptoms

are available in a FigShare repository (DOI: 10.6084/m9.figshare.16887934,

https://figshare.com/s/9b2cf4f3cdf894e67baf). Whole genome bisulfite sequencing (WGBS) data

is available at GEO (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE187022):

GSM5667182 (WGBS data of wild type TME419), GSM5667183 (WGBS data of DMS3-ZF

line 133), GSM5667184 (WGBS data of DMS3-ZF line 204). Source data are provided with this

paper.

Code availability

388   R code for generating plots are available in a FigShare repository

389   (DOI:10.6084/m9.figshare.16887934, https://figshare.com/s/9b2cf4f3cdf894e67baf).

References

1.   van Schie, C. C. N. & Takken, F. L. W. Susceptibility genes 101: how to be a good host. *Annu Rev*
Phytopathol **52**, 551–581 (2014).

2.   Chu, Z. *et al.* Promoter mutations of an essential gene for pollen development result in disease resistance in
rice. *Genes Dev.* **20**, 1250–1255 (2006).

3.   Yang, B., Sugio, A. & White, F. F. Os8N3 is a host disease-susceptibility gene for bacterial blight of rice.
PNAS **103**, 10503–10508 (2006).

4.   Antony, G. *et al.* Rice xa13 Recessive Resistance to Bacterial Blight Is Defeated by Induction of the Disease
Susceptibility Gene Os-11N3. *The Plant Cell* **22**, 3864–3876 (2010).

5.   Oliva, R. *et al.* Broad-spectrum resistance to bacterial blight in rice using genome editing. *Nature*
Biotechnology **37**, 1344–1350 (2019).

6.   Cohn, M. *et al.* Xanthomonas axonopodis virulence is promoted by a transcription activator-like effector-
mediated induction of a SWEET sugar transporter in cassava. *Mol. Plant Microbe Interact.* **27**, 1186–1198
(2014).

7.   Hu, Y. *et al.* Lateral organ boundaries 1 is a disease susceptibility gene for citrus bacterial canker disease.
PNAS **111**, E521–E529 (2014).

8.   Cox, K. L. *et al.* TAL effector driven induction of a SWEET gene confers susceptibility to bacterial blight of
cotton. *Nature Communications* **8**, 1–14 (2017).

9.   Phillips, A. Z. *et al.* Genomics-enabled analysis of the emergent disease cotton bacterial blight. *PLOS*
Genetics **13**, e1007003 (2017).

10. Nekrasov, V. *et al.* Rapid generation of a transgene-free powdery mildew resistant tomato by genome
deletion. *Sci Rep* **7**, 482 (2017).

11. Peng, A. *et al.* Engineering canker-resistant plants through CRISPR/Cas9-targeted editing of the susceptibility
gene CsLOB1 promoter in citrus. *Plant Biotechnol. J.* **15**, 1509–1519 (2017).

12. Zhang, Y. *et al.* Simultaneous modification of three homoeologs of TaEDR1 by genome editing enhances
powdery mildew resistance in wheat. *The Plant Journal* **91**, 714–724 (2017).

13. Gomez, M. A. *et al.* Simultaneous CRISPR/Cas9-mediated editing of cassava eIF4E isoforms nCBP-1 and
nCBP-2 reduces cassava brown streak disease symptom severity and incidence. *Plant Biotechnol. J.* **17**, 421–
434 (2019).

14. Xu, Z. *et al.* Engineering Broad-Spectrum Bacterial Blight Resistance by Simultaneously Disrupting Variable
TALE-Binding Elements of Multiple Susceptibility Genes in Rice. *Molecular Plant* **12**, 1434–1446 (2019).

15. Bultmann, S. *et al.* Targeted transcriptional activation of silent oct4 pluripotency gene by combining designer
TALEs and inhibition of epigenetic modifiers. *Nucleic Acids Res* **40**, 5368–5377 (2012).

16. Valton, J. *et al.* Overcoming Transcription Activator-like Effector (TALE) DNA Binding Domain Sensitivity
to Cytosine Methylation*♦. *Journal of Biological Chemistry* **287**, 38427–38432 (2012).

17. O'Malley, R. C. *et al.* Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. *Cell* **165**,
1280–1292 (2016).

18. Law, J. A. & Jacobsen, S. E. Establishing, maintaining and modifying DNA methylation patterns in plants and
animals. *Nat Rev Genet* **11**, 204–220 (2010).

19. Zhang, H., Lang, Z. & Zhu, J.-K. Dynamics and function of DNA methylation in plants. *Nat Rev Mol Cell
Biol* **19**, 489–506 (2018).

20. Papikian, A., Liu, W., Gallego-Bartolomé, J. & Jacobsen, S. E. Site-specific manipulation of Arabidopsis loci
using CRISPR-Cas9 SunTag systems. *Nature Communications* **10**, 1–11 (2019).

21. Ghoshal, B., Picard, C. L., Vong, B., Feng, S. & Jacobsen, S. E. CRISPR-based targeting of DNA methylation
in Arabidopsis thaliana by a bacterial CG-specific DNA methyltransferase. *PNAS* **118**, (2021).

22. Johnson, L. M. *et al.* SRA- and SET-domain-containing proteins link RNA polymerase V occupancy to DNA
methylation. *Nature* **507**, 124–128 (2014).

23. Gallego-Bartolomé, J. *et al.* Co-targeting RNA Polymerases IV and V Promotes Efficient De Novo DNA
Methylation in Arabidopsis. *Cell* **176**, 1068-1082.e19 (2019).

24. Constantin, E. C. *et al.* Genetic characterization of strains named as Xanthomonas axonopodis pv.
dieffenbachiae leads to a taxonomic revision of the X. axonopodis species complex. *Plant Pathology* **65**, 792–
806 (2016).

25. López, C. E. & Bernal, A. J. Cassava Bacterial Blight: Using Genomics for the Elucidation and Management
of an Old Problem. *Tropical Plant Biology* **5**, 117–126 (2012).

26. Lin, Z. J. D., Taylor, N. J. & Bart, R. Engineering Disease-Resistant Cassava. *Cold Spring Harb Perspect Biol*
11, a034595 (2019).

27. Bickhart, D. M. *et al.* Generating lineage-resolved, complete metagenome-assembled genomes from complex
microbial communities. *Nat Biotechnol* 1–9 (2022) doi:10.1038/s41587-021-01130-z.

28. Bart, R. *et al.* High-throughput genomic sequencing of cassava bacterial blight strains identifies conserved
effectors to target for durable resistance. *PNAS* **109**, E1972–E1979 (2012).

29. Cohn, M., Morbitzer, R., Lahaye, T. & Staskawicz, B. J. Comparison of gene activation by two TAL effectors
from Xanthomonas axonopodis pv. manihotis reveals candidate host susceptibility genes in cassava. *Mol.*
Plant Pathol. **17**, 875–889 (2016).

30. Wang, H. *et al.* CG gene body DNA methylation changes and evolution of duplicated genes in cassava. *PNAS*
112, 13729–13734 (2015).

31. Lim, Y.-W. *et al.* Mutations in DNA polymerase δ subunit 1 co-segregate with CMD2-type resistance to
Cassava Mosaic Geminiviruses. *Nat Commun* **13**, 1–11 (2022).

32. Stelpflug, S. C., Eichten, S. R., Hermanson, P. J., Springer, N. M. & Kaeppler, S. M. Consistent and heritable
alterations of DNA methylation are induced by tissue culture in maize. *Genetics* **198**, 209–218 (2014).

33. Stroud, H. *et al.* Plants regenerated from tissue culture contain stable epigenome changes in rice. *eLife*
https://elifesciences.org/articles/00354 (2013) doi:10.7554/eLife.00354.

34. Ong-Abdullah, M. *et al.* Loss of Karma transposon methylation underlies the mantled somaclonal variant of
oil palm. *Nature* **525**, 533–537 (2015).

35. Aung, K., Jiang, Y. & He, S. Y. The role of water in plant-microbe interactions. *Plant J* **93**, 771–780 (2018).

36. Li, T., Liu, B., Spalding, M. H., Weeks, D. P. & Yang, B. High-efficiency TALEN-based gene editing
produces disease-resistant rice. *Nature Biotechnology* **30**, 390–392 (2012).

37. Wang, F. *et al.* Enhanced Rice Blast Resistance by CRISPR/Cas9-Targeted Mutagenesis of the ERF
Transcription Factor Gene OsERF922. *PLoS One* **11**, (2016).

38. Blanvillain-Baufumé, S. *et al.* Targeted promoter editing for rice resistance to Xanthomonas oryzae pv. oryzae
reveals differential activities for SWEET14-inducing TAL effectors. *Plant Biotechnol. J.* **15**, 306–317 (2017).

39. Chandrasekaran, J. *et al.* Development of broad virus resistance in non-transgenic cucumber using
CRISPR/Cas9 technology. *Molecular Plant Pathology* **17**, 1140–1153 (2016).

40. Wang, Y. *et al.* Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable
resistance to powdery mildew. *Nature Biotechnology* **32**, 947–951 (2014).

41. Schmitz, R. J. *et al.* Transgenerational Epigenetic Instability Is a Source of Novel Methylation Variants.
Science **334**, 369–373 (2011).

42. Zheng, X. *et al.* Transgenerational Variations in DNA Methylation Induced by Drought Stress in Two Rice
Varieties with Distinguished Difference to Drought Resistance. *PLoS One* **8**, e80253 (2013).

43. Beyene, G., Chauhan, R. D., Gehan, J., Siritunga, D. & Taylor, N. Cassava shrunken-2 homolog MeAPL3
determines storage root starch and dry matter content and modulates storage root postharvest physiological
deterioration. *Plant Mol Biol* (2020) doi:10.1007/s11103-020-00995-z.

44. Law, J. A. *et al.* A Protein Complex Required for Polymerase V Transcripts and RNA- Directed DNA
Methylation in Arabidopsis. *Current Biology* **20**, 951–956 (2010).

45. Taylor, N. *et al.* A High-throughput Platform for the Production and Analysis of Transgenic Cassava
(Manihot esculenta) Plants. *Tropical Plant Biol.* **5**, 127–139 (2012).

46. Xi, Y. & Li, W. BSMAP: whole genome bisulfite sequence MAPping program. *BMC Bioinformatics* **10**, 232
(2009).

47. Mansfeld, B. N. *et al.* Large structural variations in the haplotype-resolved African cassava genome. *Plant J*
(2021) doi:10.1111/tpj.15543.

48. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

49. Bae, S., Park, J. & Kim, J.-S. Cas-OFFinder: a fast and versatile algorithm that searches for potential off-
target sites of Cas9 RNA-guided endonucleases. *Bioinformatics* **30**, 1473–1475 (2014).

50. Crooks, G. E., Hon, G., Chandonia, J.-M. & Brenner, S. E. WebLogo: A Sequence Logo Generator. *Genome*
Res. **14**, 1188–1190 (2004).

51. Moreno, I., Gruissem, W. & Vanderschuren, H. Reference genes for reliable potyvirus quantitation in cassava
and analysis of Cassava brown streak virus load in host varieties. *Journal of Virological Methods* **177**, 49–54
(2011).

52. Hu, M., Hu, W., Xia, Z., Zhou, X. & Wang, W. Validation of Reference Genes for Relative Quantitative Gene
Expression Studies in Cassava (Manihot esculenta Crantz) by Using Quantitative Real-Time PCR. *Front.*
Plant Sci. **7**, (2016).

53. Hellemans, J., Mortier, G., De Paepe, A., Speleman, F. & Vandesompele, J. qBase relative quantification
framework and software for management and automated analysis of real-time quantitative PCR data. *Genome*
Biology **8**, R19 (2007).

54. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676–682
(2012).

505   43

Acknowledgments: We are grateful to our scientific advisory board, including Drs. Morag

Ferguson, Nathan Springer, and Nigel Taylor. We also acknowledge Dr. Noah Fahlgren,

Director of the Data Science lab at DDPSC, and the additional members of our labs who

contributed to this work in various ways, especially Ke Ke, Molly Kuhs, Joshua Sumner, and Dr.

Xinggou Zheng. We also thank David Segal for assistance with the design of the artificial zinc

finger. This work was funded by the Bill and Melinda Gates Foundation (OPP1125410, RBS,

SEJ, JCC) and the National Science Foundation (GRFP DGE-2139839 and DGE-1745038, KE).

Author contributions:

Conceptualization: SMW, JCC, SEJ, RSB

Methodology: KMV, KE, GJ, ZZ, SF, MY, KBG, JCB, ZJDL, BG, JGB, JCC, SEJ, RSB

Investigation: KMV, KE, GJ, ZZ, SF, MY, KBG, JCB, ZJDL, BG, JGB, JN

Visualization: KMV, KE, ZZ, MY, KBG, JCB

Funding acquisition: SMW, JCC, SEJ, RSB

Project administration: KBG, JN, SMW, JCC, SEJ, RSB

Supervision: SMW, JCC, SEJ, RSB

Writing – original draft: KMV, ZZ, RSB

Writing – review & editing: KMV, KE, GJ, ZZ, SF, MY, KBG, JCB, ZJDL, BG, JGB,

JN, SMW, JCC, SEJ, RSB

Competing interests: Authors declare that they have no competing interests.

Fig. 1. Overview of strategy to improve disease resistance using epigenetic modification. (**A**)

Left: Example of CBB-infected cassava in a field (Uganda, 2019). Right: selected region from

leaf on the left. Water-soaked regions are indicated with white arrows. (**B**) Graphical

representation of epigenetic CBB disease resistance strategy. Top: In WT plants, TAL20 from

Xam binds a specific sequence (EBE) present in the promoter of the S gene *MeSWEET10a*. Upon

binding, TAL20 induces the ectopic expression of MeSWEET10a, a sugar transporter, which is

required for establishment of disease. A typical 'water-soaked' lesion is shown to the right, an

early indicator of CBB. Bottom: DNA methylation prevents TAL20 from binding the EBE.

MeSWEET10a expression is not induced, and disease symptoms are reduced.

Fig. 2. DMS3-ZF expression results in CpG methylation at the EBE *in vivo*. (**A**) Expression

of transgenes in individual plants from two independent DMS3-expressing transgenic lines (133

and 204) as well as a ZF-only negative control line (216). Cassava variety information for each

sample is given above the lanes. First two rows: representative western blots (anti-FLAG)

showing expression of the ZF (ZF-3xFLAG) protein with (top) and without (middle) DMS3.

Relevant size standards are shown to the right (kD). Bottom: Coomassie Brilliant Blue stained

Rubisco, loading control. Uncropped images of the western and Coomassie are shown in

Supplementary Fig. 5A. (**B**) Representative PCR-based bisulfite sequencing (ampBS-seq)

results from samples shown in A. Details results for these samples are shown in **Supplementary**

Fig. 5A. Top: Graphical depiction of *MeSWEET10a* promoter region assessed for methylation.

The EBE (grey), a presumed TATA box (blue) and the ZF binding site (orange) are indicated.

The predicted 5' UTR and *MeSWEET10a* transcriptional start site are shown in green. The area

within the dotted lined box (233 bp) was subjected to ampBS-seq. Bottom: CpG, CHG, and CHH

DNA methylation levels (percent, y-axis) of the *MeSWEET10a* promoter (EBE, grey) measured

by ampBS-seq with and without DMS3-ZF. Background of tissue for each plot is indicated to the

right. Methylation was called for cytosines with greater than 500 reads mapped. The results in

panels A and B are representative examples of the six independent experiments presented in

Supplementary Figs. 2-7. (C) Representative wild-type (TME419) and (**D**) DMS3-ZF-

expressing (line #133) plants. Scale bar = 14 cm. Additional data quantifying growth in the field

are shown in **Supplementary Fig. 8**.

Fig. 3. Targeted methylation at MeSWEET10a shows high specificity. (A) Distribution of

CpG, CHG, and CHH methylation over the targeted and flanking 5 kb regions of the

MeSWEET10a promoter in wild-type TME419 and two independent DMS3-ZF lines (transgenic

lines 133 and 204). Grey box indicates the targeted region within the *MeSWEET10a* promoter.

ZF binding site is shaded within the pink box. Two detected hypermethylated differentially

methylated regions (DMR) are also indicated. (**B**) Summary of CpG, CHG, and CHH

methylation level of targeted and flanking 5 kb regions of the *MeSWEET10a* promoter in wild-

type TME419 and the two independent DMS3-ZF lines. (**C**) Whole genome CpG, CHG, and

CHH methylation level of wild-type TME419 and the two representative DMS3-ZF lines. (**D**)

Metaplot of CpG, CHG, and CHH methylation over the gene of protein coding genes (n =

33,653) and flanking 1 kb regions in wild-type TME419 and the two independent DMS3-ZF

lines. Transcriptional start sites (TSS) and transcriptional termination sites (TTS) are indicated.

Fig. 4. Effect of methylation on CBB disease phenotypes in cassava. (**A**) Normalized

MeSWEET10a expression (y-axis, log10 scale) in WT and transgenic lines as determined by RT-

qPCR. The cassava genes GTPb (Manes.09G086600) and PP2A4 (Manes.09G039900) were

used as internal controls. Boxes are colored according to Xanthomonas treatment. Biological

replicates (black dots) included in each background (x-axis) are as follows: n = 4, 9, 4, 5, 10

examined over 4 independent experiments. The n included in each treatment group for each

background are consistent. Horizontal black line within boxes indicates the value of the median

while the box limits indicate the 25th and 75th percentiles as determined by R software; whiskers

extend 1.5 times the interquartile range (1.5xIQR) from the 25th and 75th percentiles. Two-sided

Welch's *t*-test *p*-values are noted above brackets within plot. (**B**) Representative images of water-

soaking phenotype of leaves from TME419 WT and DMS3-ZF-expressing plants. Images were

taken 4 days post-infection with either *Xam668* (XamWT) or a *Xam668* TAL20 deletion mutant

(XamΔTAL20). Scale bar = 0.5 cm. (**C**) Observed area (pixels, y-axis) of water-soaking from

images of Xam-infiltrated leaves (backgrounds, x-axis) 4 days post-infiltration. Images from (**B**)

and **Supplementary Fig. 11** are included in dataset. Calculated *p*-values (Kolmogorov-Smirnov

test) are shown above brackets within plot. (**D**) Intensity of water-soaking phenotype (y-axis) of

region measured in panel **C**. The negative mean grey-scale value for the water-soaked region

relative to the average of the mock-treated samples within the same leaf is reported (see methods

for details). Calculated *p*-values (two-sided Kolmogorov-Smirnov test) are shown above brackets

within plot. Both box plots: Biological replicates (black dots) included in each background (x-

axis) are as follows: n = 24, 30, 24, 24, 24 examined over three independent experiments.

Horizontal black line within boxes indicates the value of the median while the box limits indicate

the 25th and 75th percentiles as determined by R software; whiskers extend 1.5 times the

interquartile range (1.5xIQR) from the 25th and 75th percentiles.

A

	60444			TME 419		
		ZF			DMS3-ZF	
	WT	216		WT	204	133

DMS3-ZF — 115 kD

ZF — 26 kD

Coomassie

B

TATA Box   ZF Binding Site

233 bp   EBE   5' UTR   MeSWEET10a

Methylation (%)

[0–100] — 60444 WT
[0–100] — ZF 216
[0–100] — TME419 WT
[0–100] — 204 DMS3-ZF
[0–100] — 133

Methylation context: ■ CpG ■ CHG ■ CHH

C

D

A

Zoomed in
Chr.6: 24.084-24.096 Mb

Target region ZF

TME419 — mCG / mCHG / mCHH
DMS3-ZF 133 — mCG / mCHG / mCHH
DMS3-ZF 204 — mCG / mCHG / mCHH

TE   MeSWEET10a

DMR   DMR

1 Kb   100 bp

B

Methylation level of target and flanking 5 Kb

TME419   133   204

CG CHG CHH   CG CHG CHH   CG CHG CHH
Up 5 Kb   Target region   Down 5 Kb

C

Whole genome methylation level

TME419   133   204

CG   CHG   CHH

D

Genebody methylation (n=33653)

TME419
133
204

CG
CHG
CHH

-1 Kb   TSS   TTS   1 Kb