Hemato-vascular specification requires *arnt1* and *arnt2* genes in zebrafish embryos

Authors: Hailey E. Edwards, Jaclyn P. Souder, Daniel A. Gorelick

Center for Precision Environmental Health, Department of Molecular and Cellular Biology,

Baylor College of Medicine, Houston, Texas, USA

Correspondance to DAG gorelick@bcm.edu

ABSTRACT

During embryonic development, a subset of cells in the mesoderm germ layer are specified as

hemato-vascular progenitor cells, which then differentiate into endothelial cells and

hematopoietic stem and progenitor cells. In zebrafish, the transcription factor *npas4l*, also

known as *cloche*, is required for the specification of hemato-vascular progenitor cells.

However, it is unclear if *npas4l* is the sole factor at the top of the hemato-vascular specification

cascade. Here we show that *arnt1* and *arnt2* genes are required for hemato-vascular

specification. We found that *arnt1;arnt2* double homozygous mutant zebrafish embryos (herein

called *arnt1/2* mutants), but not *arnt1* or *arnt2* single mutants, lack blood cells and most

vascular endothelial cells. *arnt1/2* mutants have reduced or absent expression of *etv2* and *tal1*,

the earliest known endothelial and hematopoietic transcription factor genes. *npas4l* and *arnt*

genes are PAS domain-containing bHLH transcription factors that function as dimers. We

found that Npas4l binds both Arnt1 and Arnt2 proteins *in vitro*, consistent with the idea that

PAS domain-containing bHLH transcription factors act in a multimeric complex to regulate

gene expression. Our results demonstrate that *npas4l*, *arnt1* and *arnt2* act together as master

regulators of endothelial and hematopoietic cell fate. Our results also demonstrate that *arnt1*

and *arnt2* act redundantly in a transcriptional complex containing *npas4l*, but do not act

redundantly when interacting with another PAS domain-containing bHLH transcription factor,

the aryl hydrocarbon receptor. Altogether, our data enhance our understanding of hemato-

vascular specification and the function of PAS domain-containing bHLH transcription factors.

INTRODUCTION

Endothelial and hematopoietic cells are derived from the mesoderm during vertebrate

embryogenesis. These two cell types are closely linked and arise from a common progenitor

during development (Huber et al., 2004; Kataoka et al., 2011; Liao et al., 2000). The

differentiation and specification of endothelial and hematopoietic cells occurs in two waves—

primitive and definitive hematopoiesis (Costa et al., 2012; Paik and Zon, 2010; Palis et al.,

1999). During primitive hematopoiesis the first hemato-vascular progenitors are observed in

the lateral plate mesoderm of zebrafish and in the extraembryonic yolk sac mesoderm of mice

(Lee et al., 1994; Palis et al., 1999; Stainier et al., 1993). These progenitors give rise to blood

islands which contain primitive erythroid cells (Palis et al., 1999). During definitive

hematopoiesis in mice and zebrafish, fully multipotent hematopoietic stem cells (HSCs) are

derived from specialized epithelial cells, hemogenic endothelium (Hirschi, 2012; Kobayashi et

al., 2014).

In humans, the genes that drive the specification of hemato-vascular progenitors from

the multipotent mesoderm are not completely understood. We have a more complete

understanding of this process in other vertebrates. In zebrafish and chicken, Neuronal PAS 4-

like (*npas4l)* is essential for the specification of hemato-vascular progenitors from multipotent

mesoderm (Reischauer et al., 2016; Weng et al., 2020). Mammals do not appear to have a

npas4l homolog in their genomes and the mammalian functional equivalent to zebrafish *npas4l*

is unknown. Many genes involved in embryonic hematopoiesis are conserved between

zebrafish, mice, and humans (Aplan et al., 1992; Brownlie et al., 1998; Casie Chetty et al.,

2017; Dooley et al., 2005; Ferrara et al., 1996; Gong et al., 2004; Kalev-Zylinska et al., 2002b;

Kataoka et al., 2011; Okuda et al., 1996; Wang et al., 1996), but *npas4l* appears to be an

exception. Understanding the specific mechanisms of hemato-vascular specification in

zebrafish may help identify functional homologs in humans and other mammals.

Understanding the specific mechanisms of hemato-vascular specification may prove useful in

regenerative therapies for blood and vascular diseases. Recipients of embryonic stem cell-

based regenerative therapies have an increased risk of tumor formation. A promising

alternative, which ameliorates the risk of tumor formation, is to produce specific multipotent

progenitors from embryonic stem cells prior to transplantation (Doss et al., 2012; Gunaseeli et

al., 2010).

Zebrafish embryos and larvae are an ideal organism in which to study hemato-vascular

development *in vivo*. Zebrafish oocytes are fertilized externally and embryos are optically

transparent. Zebrafish embryos can survive up to 7 days post fertilization without circulating

red blood cells because oxygen can freely diffuse through the skin (Hughes et al., 2019;

Pelster and Burggren, 1996). This trait allows for characterization of cardiovascular

phenotypes that would be lethal in embryos from other organisms.

In zebrafish, endothelial cells and HSCs can be derived from a single hemato-vascular

progenitor cell population called the hemangioblast, which is hypothesized to exist in mammals

but has not been definitively confirmed (Huber et al., 2004; Moore et al., 2018; Shu et al.,

2015; Zhang and Featherstone, 2020). It has been previously reported that the basic helix-

loop-helix Per-Arnt-Sim (bHLH-PAS) transcription factor, *npas4l,* is the master regulator of

hemato-vascular specification in zebrafish (Reischauer et al., 2016) and in chicken (Weng et

al., 2020).

bHLH-PAS transcription factors are classified as Class I or Class II, which must

dimerize to regulate transcription (Erbel et al., 2003; Pongratz et al., 1998)(Figure 1A). Npas4l

is a Class I bHLH-PAS protein, but its binding partner during hemato-vascular progenitor

specification is not known. We hypothesized that Arnt1 or Arnt2 is the dimerization partner of

Npas4l in zebrafish because there are only 4 bHLH-PAS Class II proteins encoded in the

zebrafish genome and *arnt1* and *arnt2* genes are expressed at the the same stages of embryo

development as *npas4l* (Prasch et al., 2006; Reischauer et al., 2016; Tanguay et al., 2000).

To test this hypothesis, we examined hemato-vascular development in zebrafish with

loss-of-function mutations in *arnt1* and *arnt2* genes. We find that Arnt1 and Arnt2 are essential

for hemato-vascular specification. Double homozygous *arnt1;arnt2* zebrafish embryos

phenocopy *npas4l* mutants (also called *cloche*), showing lack of circulating blood, absence of

hematopoietic stem and progenitor cells, and dramatic reduction in the number of vascular

endothelial cells. Additionally, we show that zebrafish Npas4l protein interacts with both

zebrafish Arnt1 and zebrafish Arnt2 proteins *in vitro*. Together, our results suggest that Arnt1

and Arnt2 act redundantly and form a transcriptional complex with Npas4l that sits atop the

hemato-vascular specification cascade.

MATERIALS AND METHODS
91

Zebrafish

Adult zebrafish were raised at 28.5°C on a 14-h light, 10-h dark cycle in the UAB Zebrafish

Research Facility or the BCM Zebrafish Research Facility in an Aquaneering recirculating

water system (Aquaneering, Inc., San Diego, CA) and a Tecniplast recirculating water system

(Tecniplast USA, Inc., West Chester, PA). All zebrafish used for experiments were wild-type

AB strain (Westerfield, 2000) and all mutant lines were generated on the AB strain. All

procedures were performed in accordance with Institutional Animal Care and Use Committees

(IACUC) and BCM guidelines.

Embryo collection

Adult zebrafish were allowed to spawn naturally in pairs or in groups. Embryos were collected

in intervals of 20 minutes to ensure precise developmental timing or staged following

collection, placed in 60 cm^2 Petri dishes at a density of no more than 100 per dish in E3B

media (60X E3B: 17.2g NaCl, 0.76g KCl, 2.9g CaCl$_2$-2H$_2$O, 2.39g MgSO$_4$ dissolved in 1L Milli-

Q water; diluted to 1X in 9L Milli-Q water plus 100 µL 0.02% methylene blue), and then stored

in an incubator at 28.5°C on a 14-h light, 10-h dark cycle.

Genotyping adult zebrafish

Genomic DNA was isolated from tail biopsies from individual adult zebrafish by incubation in

50 µL ELB (10 mM Tris pH 8.3, 50 mM KCl, 0.3% Tween 20) with 0.5 µL proteinase K (800

U/ml, NEB) in 96 well plates, one sample per well, at 55°C for 8 hours (tail biopsies).

Proteinase K was inactivated by incubation at 98°C for 10 minutes and DNA was stored at -

20°C. Genotyping was performed by PCR and high-resolution melting curve analysis as

described (Parant et al., 2009; Romano et al., 2017). All melting curves were generated with a

Bio-Rad CFX96 Real-Time System over a 70-95°C range and analyzed with the Bio-Rad CFX

Manager 3.1 or the Bio-Rad CFX Maestro 4.1 software. All mutations were confirmed by TA

cloning and sequencing.

CRISPR-Cas9 Mutant Generation

Cas9 mRNA and gRNAs for *arnt1,* and *arnt2* mutants were generated as previously described

(Romano et al., 2017; Souder and Gorelick, 2019). Cas9 mRNA was transcribed from a

linearized pT3TS-nCas9n plasmid, Addgene #46757 (Jao et al., 2013), and purified. The target

sequences were identified using CHOPCHOP (Labun et al., 2016; Labun et al., 2019).

Oligonucleotides were annealed to each other and cloned into a pT7-gRNA plasmid, Addgene

#46759 (Jao et al., 2013). Oligonucleotides utilized are shown in Table 1. gRNAs were

synthesized from plasmids using a MEGAshortscript T7 Transcription Kit (Invitrogen AM1354)

and purified.

One-cell-stage embryos were injected using glass needles pulled on a Sutter

Instruments Fleming/Brown Micropipette Puller, model P-97 and a regulated air-pressure

micro-injector (Harvard Apparatus, NY, PL1–90). Each embryo was injected with a 1nL

solution containing one gRNA for *arnt1* and one gRNA for *arnt2* (30 ng/µL per target), Cas9

mRNA (150 ng/µL), and 0.1% phenol red. Mixtures were injected into the yolk of each embryo.

Approximately, 100 injected embryos per gRNA pair were raised to adulthood and crossed to

AB zebrafish to generate F1 embryos. F1 offspring with heritable mutations were sequenced to

identify mutations predicted to cause loss of function.

Live Imaging

Embryos were imaged with a Nikon SMZ25 microscope equipped with a Hamamatsu ORCA-

Flash4.0 digital CMOS camera. Images were equally adjusted for brightness and contrast in

Adobe Photoshop CC 2020. Embryos and larvae were anesthetized with 0.04% tricaine and

imaged in Petri dish containing E3B. All embryos were genotyped after imaging.

Whole mount *in situ* hybridization

etv2, tal1, runx1, and *flk1* antisense RNA probes were used as described (Moore et al., 2013;

Thisse and Thisse, 2008). All DNA clones were verified by sequencing. Digoxigenin-labeled

antisense RNA probes were synthesized using T7 for *runx1* and *flk1*, or T3 for *etv2*.

Colorimetric *in situ* hybridization (ISH) was performed on fixed zebrafish embryos and larvae

as previously described, with 5% dextran in the hybridization buffer (Lauter et al., 2011; Thisse

and Thisse, 2008). Following ISH, embryos were washed several times in 1x PBST

(phosphate buffered saline, 0.01% Tween), cleared with glycerol or mounted in 3%

methylcellulose on a class coverslip, and imaged on either a Nikon SMZ18 stereoscopic

microscope or an inverted Nikon ECLIPSE Ti2-E microscope. Embryos were then genotyped.

Genomic DNA was extracted using the HotSHOT protocol (Dobrzycki et al., 2018). Briefly,

embryos were suspended in 20 µL of lysis buffer (25 mM NaOH, 0.2 mM EDTA) and incubated

at 95°C for 30 minutes, cooled to 4°C, then an equal volume of neutralization buffer (40 mM

Tris-HCl) was added. DNA regions of interest were amplified with JumpStart REDTaq

ReadyMix (Sigma-Milipore, P0982) PCR according to manufacturer's protocols.

Cell Culture and Transient Transfection

HEK293T cells (gift of Dr. Charles Foulds), were thawed and maintained at 37°C in humidified

5% CO_2 atmosphere incubator. Cells were cultured in DMEM supplemented with 10% fetal

bovine serum (FBS) (Gibco, 2062564), and 1% penicillin-streptomycin (Gibco, 15140-122).

After 24 hours passaging in a 100mm plate at a density of 2.2 x 10^6 cells per dish and reaching

at least 70 confluency, the media was changed to DMEM supplemented with 1% dialyzed

FBS. Cells were transiently transfected with 12 µg of total DNA and Lipofectamine 2000

according to manufacturer's instructions (Invitrogen, 2400439) with *npas4l-v5* together with

either *arnt1-myc* or *arnt2-HA* plasmid DNA.

Cloning of plasmids for expression in cultured cells was performed as follows. For

npas4l-V5, we took pCS2+npas4lORF (gift of Dr. Leonard Zon) that contains the open reading

frame (ORF) of zebrafish *npas4l* downstream of the CMV promoter. This plasmid was modified

to include a C-terminal V5 tag using Q5 Site-Directed Mutagenesis Kit (New England Biolabs,

#E0554S). For arnt1-myc, we synthesized a plasmid containing CMV promoter driving a

codon-optimized zebrafish *arnt1* ORF (ENSDART00000081852.5) with a C-terminal myc tag

(Twist Bioscience, San Francisco, CA). For arnt2-myc, we synthesized a gene block of the

zebrafish *arnt2* ORF (ENSDART00000158162.2, Integrated DNA Technologies). The gene

block was integrated into a plasmid containing a CMV promoter using pcDNA3.3-TOPO TA

cloning kit (Invitrogen, K8300-01) in accordance with manufacturer's instructions. A C-terminal

HA tag was added using Q5 Site-Directed Mutagenesis Kit (New England Biolabs, #E0554S).

Co-immunoprecipitation and Western blot

Transiently transfected cells were washed with ice cold PBS and lysed in 500 µL of 1x RIPA

buffer containing Pierce Protease (ThermoScientific, A32955). Cells were sonicated in 6 x 30

second intervals using a Bioruptor 300 Machine (Diagenode), then centrifuged for 10 minutes

at 10,000g at 4°C. The lysate (supernatant) was removed from the pellet and placed in fresh

1.5mL Eppendorf tube. Cell lysate was precleared with Protein A beads (Invitrogen #159 18-

014) for 30 minutes 4°C, incubated with either 5 μg of V5 antibody (ThermoFisher Scientific

R960-25) or myc antibody (SantaCruz, SC-40) diluted 1:200 for 1 hour, then

immunoprecipitated with 20 μL of Protein A beads for 1 hour at 4°C. The beads were washed

three times with RIPA buffer and once with PBS and finally suspended in 30μL of PBS. Equal

volume of 2x Laemmli Sample Buffer (BioRad, cat# 1610737) was added to each sample and

boiled for 5 minutes at 95°C then placed on ice. Samples were run on 4-20% Mini-PROTEAN

TGX gels (BioRad, cat# 4561093) and transferred onto PVDF membrane using Trans-Blot

Turbo Transfer System (BioRad) using the pre-defined Mixed MW program. The membranes

were washed several times with 1x Tris-Buffered Saline, 0.1% Tween 20 (TBST)

(GBiosciences, R043) and incubated for 1 hour with agitation in 5% milk in 1x TBST.

Membranes were probed with 1:1000 V5-HRP (ThermoFisher Scientific, R960-25), 1:1000

myc-HRP (SantaCruz, SC-40), or 1:200 HA (Cell Signaling, 3724) antibodies rocking overnight

at 4°C. Membranes were washed 5 times with 1xTBST on orbital rocker for 15 minutes at room

temperature. Membranes probed with the anti-HA antibody were then probed with 1:10,000

IRDye 800CW donkey anti-rabbit (Licor, 926-32213) in 2.5%-5% milk in 1x TBST for 1 hour

and then washed multiple times with 1xTBST on orbital rocker for 5 minutes prior to imaging.

Membranes probed with HRP conjugated antibodies were developed using Clarity™ Western

ECL Substrate (BioRad, 170-5061). Membranes were imaged using BioRad ChemiDoc™ MP

Imaging System.

TCDD exposure

At 1-day post fertilization (dpf) embryos were exposed to 10 ng/mL 2,3,7,8-

tetrachlorodibenzodioxin (TCDD, AccuStandard Inc, D404N) or vehicle (0.1% DMSO). All

treatments were performed on a single clutch of at least 20 embryos; embryos treated with

vehicle were from the same clutches as embryos treated with TCDD. Chemical exposures

were performed on at least 2 clutches per genotype. Embryos were raised at 28.5°C on a 14-h

light, 10-h dark cycle until 3 dpf, when they were imaged and then genotyped.

RESULTS

arnt1/2 homozygous mutants lack circulating blood

To determine if *arnt1* or *arnt2* is required for hemato-vascular development, we generated heritable mutant alleles for *arnt1 (arnt1^{bcm1}* and *arnt1^{bcm2}*) and *arnt2 (arnt2^{bcm3}*) using CRISPR/Cas9. For *arnt1^{bcm1}*, we generated zebrafish with a 5 bp insertion in exon 5, which results in a frameshift at amino acid (AA) 126 and a premature stop codon in the first PAS domain at AA139 (Fig. 1B). For *arnt1^{bcm2}*, we generated zebrafish with an 11 bp deletion in exon 10, which results in a frameshift at AA230 and a premature stop codon between the two PAS domains at AA250 (Fig. 1B). We did not observe any phenotypic differences between homozygous *arnt1^{bcm/bcm1}* and *arnt1^{bcm2/bcm2}* mutant embryos. These mutants are morphologically indistinct from wildtype during development and homozygous adults are viable and fertile, which is consistent with previously published *arnt1* mutants (Marchi et al., 2020). We also generated maternal zygotic *arnt1* mutants (derived from homozygous *bcm1* or *bcm2* parents) and these were morphologically indistinct from wildtype. For *arnt2^{bcm3}*, we generated zebrafish with a 3 bp deletion and 1 bp insertion in exon 9, which resulted in a missense mutation at AA286 and a premature stop codon between the two PAS domains at AA287 (Fig. 1C). Homozygous *arnt2* mutants *(arnt2^{bcm3/bcm3}*) were viable at 5 days post fertilization (dpf) and grossly morphologically normal, however they did not survive to adulthood. Our observations are consistent with independently generated *arnt2* mutant zebrafish lines (Hill et al., 2009; Lohr et al., 2009).

We crossed *arnt1^{bcm2/+}; arnt2^{bcm3/+}* adult fish to each other and analyzed the phenotype and genotype of the resulting embryos. We observed all genotypes at the expected Mendelian ratios (**Table 1**). The single mutant larvae, *arnt1^{bcm2/bcm2}*, herein called *arnt1* mutants (**Fig. 1D and D'**)*,* and *arnt2^{bcm3/bcm3}*, herein called *arnt2* mutants (**Fig. 1E and E'**), are phenotypically indistinguishable from their wildtype siblings by gross morphology (**Fig. 1C and C'**). However, *arnt1^{bcm2/bcm2};arnt2^{bcm3/bcm3}* larvae, herein called *arnt1/2* mutants, showed a clear phenotype (**Fig. 1F**). These larvae lack all circulating blood (**Fig. 1H , M3 and M4**). All other intermediate *arnt1;arnt2* genotypes, e.g.,

arnt1^{bcm2/+};arnt2^{bcm3/bcm3}, appeared phenotypically wildtype (Fig. 1G, and data not shown). These results

suggest that *arnt1* and *arnt2* function redundantly, because only the *arnt1/2* double homozygous

mutants show a phenotype. Additionally, the *arnt1/2* mutant phenotype appears identical to the

previously described *npas4l* mutant phenotype (also known as *cloche*) (Reischauer et al., 2016;

Stainier et al., 1995a). Like *npas4l* mutants, *arnt1/2* mutants exhibit cardiac edema with enlarged heart

atrium, likely a secondary effect of the lack of blood and endocardial cells (Fig 1G). Taken together,

these data suggest that *arnt1* and *arnt2* have a similar function as *npas4l*.

arnt1/2 mutants have reduced expression of hematopoietic and endothelial cell markers at 2 dpf

We hypothesized that the Arnt proteins are dimerization partners with Npas4l and that *arnt1/2*

mutants would exhibit the same phenotypes as *npas4l* mutants. Therefore, we examined *arnt1/2*

mutant embryos for additional phenotypes found in *npas4l* mutants. Previous studies determined that

npas4l mutants lack almost all endothelial cells, except some cells located in the lower trunk and caudal

tail of the larvae (Liao et al., 1997; Stainier et al., 1995a). To determine if our *arnt1/2* mutants showed a

similar reduction in endothelial cells, we performed whole-mount *in situ* hybridization (WISH) on 2 dpf

embryos for *flk-1* (also known as *kdrl*). *flk-1*, a widely used marker of endothelial cells in blood vessels,

is essential for the development of blood vessels in mice and zebrafish (Dumont et al., 1995; Liao et al.,

1997; Shalaby et al., 1995). We found that *maternal zygotic arnt1* mutants (*MZarnt1*) and *arnt2*

mutants showed clear expression of *flk-1* in the vasculature, similar to wild type (**Fig 2B-C**). In contrast,

arnt1/2 mutants had a reduced number of *flk-1*-positive cells (**Fig. 2D and D'**). The *flk-1*-positive cells

present in the *arnt1/2* mutant were in the caudal tail and likely comprised a portion of the caudal vein

(**Fig. 2D'**), similar to the *flk-1* expression pattern seen in *npas4l* mutants (Liao et al., 1997). This

suggests that *npas4l* and *arnt1* or *arnt2* are required for the development of most blood vessels.

In addition to a significant reduction of endothelial cells, *npas4l* mutants show a reduction in the

number of hematopoietic stem cells (HSC) (Stainier et al., 1995a; Thompson et al., 1998). To

determine if *arnt1/2* mutants contain HSCs, we examined *runx1* expression by WISH. *runx1* is a marker

of HSCs and is required for definitive hematopoiesis in zebrafish (Kalev-Zylinska et al., 2002a). By 48

hpf, *runx1*-positive HSCs are localized to the area between the dorsal aorta and the posterior caudal

vein (Lam et al., 2009). At 2 dpf, wildtype, *arnt1*, and *arnt2* single mutants all contain *runx1*-positive

cells in the blood island region (**Fig 2E-G)**, indicating the presence of HSCs. However, *arnt1/2* mutants

lack *runx1*-positive cells in this region (**Fig. 2H**). This suggests that *arnt1* or *arnt2* is necessary for

hematopoietic cell development during embryogenesis, similar to *npas4l* (Liao et al., 1998; Stainier et

al., 1995a; Thompson et al., 1998).

Hematopoietic and endothelial progenitor populations are reduced in *arnt1/2* mutants

Hematopoietic cells and endothelial cells arise from a common progenitor population in

zebrafish (Vogeli et al., 2006). To determine if *arnt1/2* mutants have a reduction of hemato-vascular

progenitor cells, we performed WISH on embryos at the 4-6 somite stage and examined expression of

the earliest known regulators of endothelial and hematopoietic cell differentiation: *etv2* and *tal1,* both of

which are expressed in the posterior lateral plate mesoderm (PLPM) during early somitogenesis stages

(Qian et al., 2005; Reischauer et al., 2016; Stainier et al., 1995a; Thompson et al., 1998). We bred

arnt1^{bcm2/+}; arnt2^{bcm3/+} adults to each other to generate *arnt1/2* mutant, *arnt1* mutant, *arnt2* mutant, and

wildtype embryos. We then performed WISH to examine the expression of *etv2* (**Fig. 3A and B**, n=516

embryos from 5 clutches) or *tal1* (**Fig. 3C and D**, n=665 embryos from 3 clutches). 8% and 6% of

embryos showed reduced expression of *etv2* and *tal1*, respectively, neither of which is statistically

significantly different from the expected Mendelian ratio of 6.25% double homozygous mutant embryos

(binomial test, *etv2* n=516 *p>0.05*; *tal1* n=665 *p>0.05*). To confirm that the observed reduction of *etv2*

and *tal1* in these embryos correlates with the *arnt1/2* double homozygous mutant genotype, following

WISH we genotyped 24 of the embryos with absent or reduced expression of *etv2* and 17 with absent

or reduced expression of *tal1*. 100% of the low/no expressing *etv2* embryos were *arnt1/2* double

homozygous mutants, while 83% of the low/no expressing *tal1* embryos were *arnt1/2* double

homozygous mutants, demonstrating a strong correlation between the reduced expression of *etv2* and

tal1 and the *arnt1/2* double homozygous genotype. Additionally, we genotyped 26 of the embryos with

normal expression of *etv2* and 7 with normal expression of *tal1*. 7.7% of the normal expressing *etv2*

embryos were *arnt1/2* mutants (genotypes of the 26 embryos: 2 *arnt1/2* mutants, 1 wildtype, 3

arnt1$^{bcm2/bcm2}$, 4 *arnt1$^{bcm2/bcm2}$;arnt2$^{bcm3/+}$*, 4 *arnt1$^{bcm2/+}$*, 5 *arnt1$^{bcm2/+}$;arnt2$^{bcm3/+}$*, 4 *arnt1$^{bcm2/+}$;arnt2$^{bcm3/bcm3}$*,

1 *arnt2$^{bcm3/bcm3}$*, and 3 *arnt2$^{bcm3/+}$*), while 0% of the normal expressing *tal1* embryos were *arnt1/2*

mutants (genotypes of the 7 embryos: 1 *arnt1$^{bcm2/+}$;arnt2$^{bcm3/+}$*, 1 *arnt1$^{bcm2/+}$;arnt2^{bcm3bcm3}*, 1 *arnt1$^{bcm2/+}$*, 1

arnt2$^{bcm3/+}$, and 3 *arnt1$^{bcm2/bcm2}$;arnt2$^{bcm3/+}$*). We conclude that *arnt1* and *arnt2* are required for hemato-

vascular progenitor specification. Previous findings demonstrated that *npas4l* is a master regulator of

hemato-vascular progenitor specification (Reischauer et al., 2016). Together, these results demonstrate

that *npas4l*, *arnt1* and *arnt2* are co-master regulators of hemato-vascular specification.

304   Npas4l interacts with both Arnt1 and Arnt2

Npas4l is a Class I bHLH-PAS domain transcription factor (Reischauer et al., 2016), and as

such, must heterodimerize with Class II bHLH-PAS transcription factors, like Arnt1 and Arnt2, to initiate

gene transcription . To determine if Npas4l forms a complex with either Arnt1 or Arnt2, we performed a

coimmunoprecipitation experiment. Due to the absence of commercially available antibodies that

recognize zebrafish Npas4l, Arnt1 or Arnt2 proteins, we co-expressed tagged versions of zebrafish

npas4l (Npas4l-V5), *arnt1* (Arnt1-myc), and *arnt2* (Arnt2-HA) in HEK293T cells. We performed co-

immunoprecipitation studies and found that Arnt1-myc interacts with Npas4l-V5 and that Arnt2-HA

interacts with Npas4l-V5 (**Fig. 4A, B**). These data suggest that Npas4l uses either Arnt1 or Arnt2 or

both as a binding partner. Furthermore, these data indicate that Arnt1 and Arnt2 may be functionally

interchangeable and redundant, at least in the case of Npas4l.

Arnt1 and Arnt2 cannot functionally compensate for one another in the case of Ahr2

Our results demonstrate that Arnt1 and Arnt2 are functionally interchangeable and redundant when it comes to interacting with Npas4l in the context of hemato-vascular development. It is unclear if Arnt1 and Arnt2 are functionally interchangeable with all Class I bHLH-PAS domain transcription factors or just with Npas4l. To determine if *arnt1/arnt2* redundancy is unique to Npas4l, we examined if *arnt1* and *arnt2* are interchangeable in the case of a different class I bHLH-PAS transcription factor, the aryl hydrocarbon receptor 2 (Ahr2). Previous studies demonstrate that Ahr2 can bind DNA in the presence of either Arnt1 or Arnt2 *in vitro* (Andreasen et al., 2002; Hirose et al., 1996; Tanguay et al., 2000), but whether Arnt1 and Arnt2 bind Ahr2 interchangeably *in vivo* is not known. Ahr2 is a ligand-dependent transcription factor. A known ligand of Ahr2 is 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), which we have previously shown induces cardiotoxicity in zebrafish larvae and requires wild-type *ahr2* (Souder and Gorelick, 2019*)*. We asked whether TCDD causes cardiotoxicity in *arnt1* or *arnt2* single mutants, reasoning that if Ahr2 interacts with Arnt1 or Arnt2 interchangeably, then only *arnt1/2* double mutants, but not single mutants, would be resistant to TCDD exposure. We treated wildtype, *arnt1* mutants, and *arnt2* mutants with either vehicle (0.1% DMSO) (**Fig. 5A, B, and C**) or with 10 ng/mL TCDD (**Fig. 5D, E, and F**) from 1 dpf to 3 dpf. Wildtype and *arnt2* mutants treated with TCDD exhibit cardiac edema and the heart fails to loop properly (**Fig. 5D,** and **F**), unlike vehicle treated siblings (**Fig. 5A** and **C**). In contrast, *arnt1* mutants treated with TCDD showed normal heart looping and lack cardiac edema (**Fig. 5E**), like their vehicle treated siblings (**Fig. 5B**). This suggests that *arnt1,* but not *arnt2,* is required for Ahr2-dependent TCDD toxicity. This also shows that *arnt1* and *arnt2* are not always redundant *in vivo* and that the Class I dimerization partner may influence whether the Class II dimerization partners are redundant.

DISCUSSION

Class I bHLH-PAS proteins must dimerize with Class II bHLH-PAS proteins to transcribe their target genes (Ema et al., 1996; Erbel et al., 2003; Jiang et al., 1996; Lees and Whitelaw, 1999; Michaud et al., 2000; Wu and Rastinejad, 2017). It has previously been established that Npas4l, a Class I bHLH-PAS transcription factor, is required for hemato-vascular development in zebrafish (Reischauer et al., 2016). To determine the Class II bHLH-PAS binding partner of Npas4l, we generated *arnt1* mutant and *arnt2* mutant zebrafish and expected one of these mutants to mimic the *npas4l* mutant phenotype. However, both *arnt1* mutant and *arnt2* mutant larvae are viable to 5dpf and are indistinguishable from wildtype embryos by eye. Our results demonstrate that double homozygous *arnt1/2* mutant embryos have a similar phenotype to *npas4l* mutants (Reischauer et al., 2016; Stainier et al., 1995b; Thompson et al., 1998) and that Npas4l can function with either Arnt1 or Arnt2. Thus, Arnt1 and Arnt2 act redundantly to specify hemato-vascular progenitor cells.

Endothelial cell development and hematopoietic cell development are closely related during embryogenesis. Both cell types primarily arise from the lateral plate mesoderm (LPM) and share a common progenitor (Qian et al., 2005; Reischauer et al., 2016; Stainier et al., 1995a; Thompson et al., 1998). Additionally, during definitive hematopoiesis, some HSCs are derived from specialized endothelial cells, called hemogenic endothelium (Boisset et al., 2010; Kissa and Herbomel, 2010). Both *npas4l* and *arnt1/2* mutants lack all blood cells but do have a small population of endothelial cells, which is restricted to the most posterior region of the animal (Figure 2 and (Reischauer et al., 2016)). In zebrafish, there is a population of endothelial cells in the caudal region of the embryo that are not derived from the LPM and differentiate in the absence of *npas4l* expression (Pak et al., 2020). We hypothesize that *arnt1* and *arnt2* are

required for the differentiation of endothelial cells from the LPM but are not required for the

differentiation of endothelial cells, such as the caudal population, from outside the LPM. The

transcriptomic signature of the caudal population of endothelial cells was distinct from

endothelial cells that require *npas4l* for differentiation, with genes for somitogenesis and

neurogenesis highly enriched compared to LPM-derived endothelial cells (Pak et al., 2020).

Based on this transcriptome analysis, we speculate that the caudal population of endothelial

cells fails to express genes required for hemogenic endothelium and that no blood cells are

derived from the caudal population of endothelial cells. This is consistent with the observation

that the caudal population of endothelial cells, but not blood cells, are present in *npas4l* or

arnt1/2 mutants.

Our results show that either Arnt1 or Arnt2, Class II bHLH-PAS proteins, can act as a

required co-factor for Npas4l-dependent mesodermal specification to hematopoietic and

endothelial progenitors. Likewise, Arnt1 is essential for adult hematopoietic stem cells in mice

and is required for survival of hematopoietic progenitor cells in the fetal liver (Krock et al.,

2015). Homozygous Arnt1 mutant mice are not viable beyond E10.5 and show severe defects

in vascularization, suggesting that Arnt1 is essential for normal vascular development (Kozak

et al., 1997; Maltepe et al., 1997). The role of Arnt1 during mammalian embryonic

hematopoiesis remains unclear.

Based on these data we speculate that the mammalian equivalent to zebrafish Npas4l

is likely to be a Class I bHLH-PAS protein that heterodimerizes with ARNT1. Zebrafish Npas4l

shows the most sequence homology to mouse Npas4 (Reischauer et al., 2016). Although

Npas4 is known to dimerize with Arnt1 and Arnt2 *in vivo* (Brigidi et al., 2019), mammalian

Npas4 is unlikely to be the functional equivalent to zebrafish Npas4l because homozygous

Npas4 mutant mice are viable to adulthood and do not show any overt vascular or

hematopoietic defects (Bloodgood et al., 2013).

Similarly, we speculate that Npas1, 2, and 3 are unlikely to be the mammalian

equivalent to Npas4l, as mice mutant for these genes are viable to adulthood and no

endothelial or hematopoietic phenotype has been reported (Dudley et al., 2003; Erbel-Sieler et

al., 2004). Additionally, we think that the Sim (Sim1/2) and Circadian Locomotor Output Cycles

Kaput (CLOCK) genes are unlikely to the be the master regulators of hemato-vascular

specification in mice or humans because mutant mice appear to have normal blood cells and

vasculature (DeBruyne et al., 2007; Epstein et al., 2000; Holder et al., 2004). We cannot

exclude the possibility that Npas1, 2, 3 and 4 proteins act redundantly to specify hemato-

vascular progenitors in mammalian embryos because mice with mutations in multiple Npas

genes have not been reported.

Three other Class I bHLH-PAS proteins, Ahr, Hif1α, and Hif2α, are known to be

involved in mammalian hematopoiesis and vascularization. Lack of Ahr *in vitro* and *in vivo*

promotes adult hematopoietic progenitor expansion and resulted in aberrant ratios and

functionality of some blood cell populations, suggesting that Ahr could be functionally relevant

during definitive hematopoiesis (Singh et al., 2011; Smith et al., 2013). However, there is no

evidence that Ahr is expressed during primitive hematopoiesis in mice and Ahr mutant mice

are viable to adulthood and do not show any vascular defects (Singh et al., 2011).

Hif1α and Hif2α are involved in mouse vascular development and hematopoiesis (Iyer

et al., 1998; Krock et al., 2015; Peng et al., 2000; Scortegagna et al., 2003; Takubo et al.,

2010). However, the role of Hif1α and Hif2α during early mouse embryonic hematopoiesis and

endothelial development is not well characterized. Hif1α is required for cell cycle quiescence of

adult hematopoietic stem cells (Takubo et al., 2010). Hif1α mRNA is detected during mouse

primitive hematopoiesis and Hif1α homozygous mutant mice die *in utero* before E11.5 (Iyer et

al., 1998; Kotch et al., 1999). The Hif1α mutant mice are reported to have defects in blood

vessels, increased death of vascular-supporting mesenchymal cells, and cardiovascular

defects (Iyer et al., 1998; Kotch et al., 1999). In mice, Hif2α mRNA is expressed during primary

hematopoiesis specifically within the yolk sac, where primary hematopoiesis originates

(Flamme et al., 1997). Global knockout of *Hif2α* is reported to be embryonic or postnatal lethal

depending upon the mutation and strain of mouse used (Compernolle et al., 2002; Peng et al.,

2000; Tian et al., 1998). Embryonic lethal Hif2α mutants showed severe vascular

disorganization and increased frequency of hemorrhaging *in utero* (Compernolle et al., 2002;

Peng et al., 2000). Both *Hif1α* and *Hif2α* mutants have endothelial and blood cells, thus neither

mutation mimics the zebrafish *npas4l* phenotype. But both *Hif1α* and *Hif2α* mutant mice do

have a vascular phenotype (Compernolle et al., 2002; Compernolle et al., 2003; Iyer et al.,

1998; Peng et al., 2000; Ryan et al., 1998). Although neither Hif1α nor Hif2α single mutant

mice recapitulate the phenotype observed in zebrafish *cloche* or *arnt1;arnt2* mutants, the HIFs

are strong candidates to be the functional equivalent of Npas4l.

One reason we may not see an Npas4l-like phenotype in Hif1α or Hif2α single mutant

mice could be because of genetic compensation. Genetic compensation occurs in both

zebrafish and mice (Bunton-Stasyshyn et al., 2019; Rossi et al., 2015). There is evidence in

zebrafish and mice of *Hif1α* and *Hif2α* functionally compensating for one another. Zebrafish

with reduced expression of all 4 Hif1 and Hif2 transcripts *(hif1aa, hif1ab, hif2aa, hif2ab)* lack

hemogenic endothelium, suggesting that at least one HIF gene is required for the specification

of hemogenic endothelium (Gerri et al., 2018). These results also indicate that *hif1α* and *hif2α*

are functionally redundant for the generation of hemogenic endothelium (Gerri et al., 2018).

Global knockout of mouse *Hif1α* is embryonic lethal (Compernolle et al., 2003; Iyer et al., 1998;

Ryan et al., 1998) and global knockout of mouse *Hif2α* is embryonic or postnatal lethal

depending upon the strain of mouse used (Compernolle et al., 2002; Peng et al., 2000; Tian et

al., 1998).

To our knowledge, no one has generated a mouse in which both *Hif1α* and *Hif2α* are

globally knocked-out. Future studies could examine if global or mesodermal-specific knockout

of both *Hif1α* and *Hif2α* genes in mouse results in a phenotype similar to knockout of both

arnt1 and *arnt2* in zebrafish. Our results demonstrate that under some conditions, Arnt1 and

Arnt2 act redundantly. In other contexts, Arnt1 is the preferred binding protein for a class I PAS

transcription factor. The extent to which this occurs during different developmental stages *in*

vivo, or in different cell types, warrants further study.

Acknowledgements

We thank Lauren Pandolfo and the staff of the Baylor College of Medicine aquatics facility for taking

care of our zebrafish colony. This work funded by NIH grants R01ES026337 and P30ES030285.

447   References:

Andreasen, E.A., Hahn, M.E., Heideman, W., Peterson, R.E., Tanguay, R.L., 2002. The zebrafish
(Danio rerio) aryl hydrocarbon receptor type 1 is a novel vertebrate receptor. Molecular pharmacology
62, 234-249.

Aplan, P.D., Nakahara, K., Orkin, S.H., Kirsch, I.R., 1992. The SCL gene product: a positive regulator
of erythroid differentiation. The EMBO journal 11, 4073-4081.

Bloodgood, B.L., Sharma, N., Browne, H.A., Trepman, A.Z., Greenberg, M.E., 2013. The activity-
dependent transcription factor NPAS4 regulates domain-specific inhibition. Nature 503, 121-125.

Boisset, J.C., van Cappellen, W., Andrieu-Soler, C., Galjart, N., Dzierzak, E., Robin, C., 2010. In vivo
imaging of haematopoietic cells emerging from the mouse aortic endothelium. Nature 464, 116-120.

Brigidi, G.S., Hayes, M.G.B., Delos Santos, N.P., Hartzell, A.L., Texari, L., Lin, P.A., Bartlett, A., Ecker,
J.R., Benner, C., Heinz, S., Bloodgood, B.L., 2019. Genomic Decoding of Neuronal Depolarization by
Stimulus-Specific NPAS4 Heterodimers. Cell 179, 373-391 e327.

Brownlie, A., Donovan, A., Pratt, S.J., Paw, B.H., Oates, A.C., Brugnara, C., Witkowska, H.E., Sassa,
S., Zon, L.I., 1998. Positional cloning of the zebrafish sauternes gene: a model for congenital
sideroblastic anaemia. Nat Genet 20, 244-250.

Bunton-Stasyshyn, R.K.A., Wells, S., Teboul, L., 2019. When all is not lost: considering genetic
compensation in laboratory animals. Lab Anim (NY) 48, 282-284.

Casie Chetty, S., Rost, M.S., Enriquez, J.R., Schumacher, J.A., Baltrunaite, K., Rossi, A., Stainier,
D.Y., Sumanas, S., 2017. Vegf signaling promotes vascular endothelial differentiation by modulating
etv2 expression. Dev Biol 424, 147-161.

Compernolle, V., Brusselmans, K., Acker, T., Hoet, P., Tjwa, M., Beck, H., Plaisance, S., Dor, Y.,
Keshet, E., Lupu, F., Nemery, B., Dewerchin, M., Van Veldhoven, P., Plate, K., Moons, L., Collen, D.,
Carmeliet, P., 2002. Loss of HIF-2alpha and inhibition of VEGF impair fetal lung maturation, whereas
treatment with VEGF prevents fatal respiratory distress in premature mice. Nat Med 8, 702-710.

Compernolle, V., Brusselmans, K., Franco, D., Moorman, A., Dewerchin, M., Collen, D., Carmeliet, P.,
2003. Cardia bifida, defective heart development and abnormal neural crest migration in embryos
lacking hypoxia-inducible factor-1alpha. Cardiovasc Res 60, 569-579.

Costa, G., Kouskoff, V., Lacaud, G., 2012. Origin of blood cells and HSC production in the embryo.
Trends Immunol 33, 215-223.

DeBruyne, J.P., Weaver, D.R., Reppert, S.M., 2007. CLOCK and NPAS2 have overlapping roles in the
suprachiasmatic circadian clock. Nat Neurosci 10, 543-545.

Dobrzycki, T., Krecsmarik, M., Bonkhofer, F., Patient, R., Monteiro, R., 2018. An optimised pipeline for
parallel image-based quantification of gene expression and genotyping after in situ hybridisation. Biol
Open 7.

Dooley, K.A., Davidson, A.J., Zon, L.I., 2005. Zebrafish scl functions independently in hematopoietic
and endothelial development. Dev Biol 277, 522-536.

Doss, M.X., Gaspar, J.A., Winkler, J., Hescheler, J., Schulz, H., Sachinidis, A., 2012. Specific gene
signatures and pathways in mesodermal cells and their derivatives derived from embryonic stem cells.
Stem Cell Rev Rep 8, 43-54.

Dudley, C.A., Erbel-Sieler, C., Estill, S.J., Reick, M., Franken, P., Pitts, S., McKnight, S.L., 2003.
Altered patterns of sleep and behavioral adaptability in NPAS2-deficient mice. Science 301, 379-383.

Dumont, D.J., Fong, G.H., Puri, M.C., Gradwohl, G., Alitalo, K., Breitman, M.L., 1995. Vascularization of
the mouse embryo: a study of flk-1, tek, tie, and vascular endothelial growth factor expression during
development. Dev Dyn 203, 80-92.

Ema, M., Morita, M., Ikawa, S., Tanaka, M., Matsuda, Y., Gotoh, O., Saijoh, Y., Fujii, H., Hamada, H.,
Kikuchi, Y., Fujii-Kuriyama, Y., 1996. Two new members of the murine Sim gene family are
transcriptional repressors and show different expression patterns during mouse embryogenesis.
Molecular and cellular biology 16, 5865-5875.

Epstein, D.J., Martinu, L., Michaud, J.L., Losos, K.M., Fan, C., Joyner, A.L., 2000. Members of the
bHLH-PAS family regulate Shh transcription in forebrain regions of the mouse CNS. Development 127,
4701-4709.

Erbel, P.J., Card, P.B., Karakuzu, O., Bruick, R.K., Gardner, K.H., 2003. Structural basis for PAS
domain heterodimerization in the basic helix--loop--helix-PAS transcription factor hypoxia-inducible
factor. Proc Natl Acad Sci U S A 100, 15504-15509.

Erbel-Sieler, C., Dudley, C., Zhou, Y., Wu, X., Estill, S.J., Han, T., Diaz-Arrastia, R., Brunskill, E.W.,
Potter, S.S., McKnight, S.L., 2004. Behavioral and regulatory abnormalities in mice deficient in the
NPAS1 and NPAS3 transcription factors. Proc Natl Acad Sci U S A 101, 13648-13653.

Ferrara, N., Carver-Moore, K., Chen, H., Dowd, M., Lu, L., O'Shea, K.S., Powell-Braxton, L., Hillan,
K.J., Moore, M.W., 1996. Heterozygous embryonic lethality induced by targeted inactivation of the
VEGF gene. Nature 380, 439-442.

Flamme, I., Frohlich, T., von Reutern, M., Kappel, A., Damert, A., Risau, W., 1997. HRF, a putative
basic helix-loop-helix-PAS-domain transcription factor is closely related to hypoxia-inducible factor-1
alpha and developmentally expressed in blood vessels. Mech Dev 63, 51-60.

Gerri, C., Marass, M., Rossi, A., Stainier, D.Y.R., 2018. Hif-1alpha and Hif-2alpha regulate hemogenic
endothelium and hematopoietic stem cell formation in zebrafish. Blood 131, 963-973.

Gong, B., Liang, D., Chew, T.G., Ge, R., 2004. Characterization of the zebrafish vascular endothelial
growth factor A gene: comparison with vegf-A genes in mammals and Fugu. Biochim Biophys Acta
1676, 33-40.

Gunaseeli, I., Doss, M.X., Antzelevitch, C., Hescheler, J., Sachinidis, A., 2010. Induced pluripotent
stem cells as a model for accelerated patient- and disease-specific drug discovery. Curr Med Chem 17,
759-766.

Hill, A.J., Heiden, T.C., Heideman, W., Peterson, R.E., 2009. Potential roles of Arnt2 in zebrafish larval
development. Zebrafish 6, 79-91.

Hirose, K., Morita, M., Ema, M., Mimura, J., Hamada, H., Fujii, H., Saijo, Y., Gotoh, O., Sogawa, K., Fujii-Kuriyama, Y., 1996. cDNA cloning and tissue-specific expression of a novel basic helix-loop-helix/PAS factor (Arnt2) with close sequence similarity to the aryl hydrocarbon receptor nuclear translocator (Arnt). Molecular and cellular biology 16, 1706-1713.

Hirschi, K.K., 2012. Hemogenic endothelium during development and beyond. Blood 119, 4823-4827.

Holder, J.L., Jr., Zhang, L., Kublaoui, B.M., DiLeone, R.J., Oz, O.K., Bair, C.H., Lee, Y.H., Zinn, A.R., 2004. Sim1 gene dosage modulates the homeostatic feeding response to increased dietary fat in mice. American journal of physiology. Endocrinology and metabolism 287, E105-113.

Huber, T.L., Kouskoff, V., Fehling, H.J., Palis, J., Keller, G., 2004. Haemangioblast commitment is initiated in the primitive streak of the mouse embryo. Nature 432, 625-630.

Hughes, M.C., Zimmer, A.M., Perry, S.F., 2019. Role of internal convection in respiratory gas transfer and aerobic metabolism in larval zebrafish (Danio rerio). Am J Physiol Regul Integr Comp Physiol 316, R255-R264.

Iyer, N.V., Kotch, L.E., Agani, F., Leung, S.W., Laughner, E., Wenger, R.H., Gassmann, M., Gearhart, J.D., Lawler, A.M., Yu, A.Y., Semenza, G.L., 1998. Cellular and developmental control of O2 homeostasis by hypoxia-inducible factor 1 alpha. Genes & development 12, 149-162.

Jao, L.E., Wente, S.R., Chen, W., 2013. Efficient multiplex biallelic zebrafish genome editing using a CRISPR nuclease system. Proc Natl Acad Sci U S A 110, 13904-13909.

Jiang, B.H., Rue, E., Wang, G.L., Roe, R., Semenza, G.L., 1996. Dimerization, DNA binding, and transactivation properties of hypoxia-inducible factor 1. J Biol Chem 271, 17771-17778.

Kalev-Zylinska, M.L., Horsfield, J.A., Flores, M.V., Postlethwait, J.H., Vitas, M.R., Baas, A.M., Crosier, P.S., Crosier, K.E., 2002a. Runx1 is required for zebrafish blood and vessel development and expression of a human RUNX1-CBF2T1 transgene advances a model for studies of leukemogenesis. Development 129, 2015-2030.

Kalev-Zylinska, M.L., Horsfield, J.A., Flores, M.V.C., Postlethwait, J.H., Vitas, M.R., Baas, A.M., Crosier, P.S., Crosier, K.E., 2002b. Runx1 is required for zebrafish blood and vessel development and expression of a human RUNX1-CBF2T1 transgene advances a model for studies of leukemogenesis. Development 129, 2015-2030.

Kataoka, H., Hayashi, M., Nakagawa, R., Tanaka, Y., Izumi, N., Nishikawa, S., Jakt, M.L., Tarui, H., Nishikawa, S., 2011. Etv2/ER71 induces vascular mesoderm from Flk1+PDGFRalpha+ primitive mesoderm. Blood 118, 6975-6986.

Kissa, K., Herbomel, P., 2010. Blood stem cells emerge from aortic endothelium by a novel type of cell transition. Nature 464, 112-115.

Kobayashi, I., Kobayashi-Sun, J., Kim, A.D., Pouget, C., Fujita, N., Suda, T., Traver, D., 2014. Jam1a-Jam2a interactions regulate haematopoietic stem cell fate through Notch signalling. Nature 512, 319-323.

Kotch, L.E., Iyer, N.V., Laughner, E., Semenza, G.L., 1999. Defective vascularization of HIF-1alpha-null
embryos is not associated with VEGF deficiency but with mesenchymal cell death. Dev Biol 209, 254-
267.

Kozak, K.R., Abbott, B., Hankinson, O., 1997. ARNT-deficient mice and placental differentiation. Dev
Biol 191, 297-305.

Krock, B.L., Eisinger-Mathason, T.S., Giannoukos, D.N., Shay, J.E., Gohil, M., Lee, D.S., Nakazawa,
M.S., Sesen, J., Skuli, N., Simon, M.C., 2015. The aryl hydrocarbon receptor nuclear translocator is an
essential regulator of murine hematopoietic stem cell viability. Blood 125, 3263-3272.

Labun, K., Montague, T.G., Gagnon, J.A., Thyme, S.B., Valen, E., 2016. CHOPCHOP v2: a web tool
for the next generation of CRISPR genome engineering. Nucleic Acids Res 44, W272-276.

Labun, K., Montague, T.G., Krause, M., Torres Cleuren, Y.N., Tjeldnes, H., Valen, E., 2019.
CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. Nucleic Acids Res 47,
W171-W174.

Lam, E.Y., Chau, J.Y., Kalev-Zylinska, M.L., Fountaine, T.M., Mead, R.S., Hall, C.J., Crosier, P.S.,
Crosier, K.E., Flores, M.V., 2009. Zebrafish runx1 promoter-EGFP transgenics mark discrete sites of
definitive blood progenitors. Blood 113, 1241-1249.

Lauter, G., Soll, I., Hauptmann, G., 2011. Multicolor fluorescent in situ hybridization to define abutting
and overlapping gene expression in the embryonic zebrafish brain. Neural Dev 6, 10.

Lee, R.K., Stainier, D.Y., Weinstein, B.M., Fishman, M.C., 1994. Cardiovascular development in the
zebrafish. II. Endocardial progenitors are sequestered within the heart field. Development 120, 3361-
3366.

Lees, M.J., Whitelaw, M.L., 1999. Multiple roles of ligand in transforming the dioxin receptor to an active
basic helix-loop-helix/PAS transcription factor complex with the nuclear protein Arnt. Molecular and
cellular biology 19, 5811-5822.

Liao, E.C., Paw, B.H., Oates, A.C., Pratt, S.J., Postlethwait, J.H., Zon, L.I., 1998. SCL/Tal-1
transcription factor acts downstream of cloche to specify hematopoietic and vascular progenitors in
zebrafish. Genes & development 12, 621-626.

Liao, W., Bisgrove, B.W., Sawyer, H., Hug, B., Bell, B., Peters, K., Grunwald, D.J., Stainier, D.Y., 1997.
The zebrafish gene cloche acts upstream of a flk-1 homologue to regulate endothelial cell
differentiation. Development 124, 381-389.

Liao, W., Ho, C.Y., Yan, Y.L., Postlethwait, J., Stainier, D.Y., 2000. Hhex and scl function in parallel to
regulate early endothelial and blood differentiation in zebrafish. Development 127, 4303-4313.

Lohr, H., Ryu, S., Driever, W., 2009. Zebrafish diencephalic A11-related dopaminergic neurons share a
conserved transcriptional network with neuroendocrine cell lineages. Development 136, 1007-1017.

Maltepe, E., Schmidt, J.V., Baunoch, D., Bradfield, C.A., Simon, M.C., 1997. Abnormal angiogenesis
and responses to glucose and oxygen deprivation in mice lacking the protein ARNT. Nature 386, 403-
407.

Marchi, D., Santhakumar, K., Markham, E., Li, N., Storbeck, K.H., Krone, N., Cunliffe, V.T., van Eeden,
F.J.M., 2020. Bidirectional crosstalk between Hypoxia-Inducible Factor and glucocorticoid signalling in
zebrafish larvae. PLoS Genet 16, e1008757.

Michaud, J.L., DeRossi, C., May, N.R., Holdener, B.C., Fan, C.M., 2000. ARNT2 acts as the
dimerization partner of SIM1 for the development of the hypothalamus. Mech Dev 90, 253-261.

Moore, C., Richens, J.L., Hough, Y., Ucanok, D., Malla, S., Sang, F., Chen, Y., Elworthy, S., Wilkinson,
R.N., Gering, M., 2018. Gfi1aa and Gfi1b set the pace for primitive erythroblast differentiation from
hemangioblasts in the zebrafish embryo. Blood Adv 2, 2589-2606.

Moore, J.C., Sheppard-Tindell, S., Shestopalov, I.A., Yamazoe, S., Chen, J.K., Lawson, N.D., 2013.
Post-transcriptional mechanisms contribute to Etv2 repression during vascular development. Dev Biol
384, 128-140.

Okuda, T., van Deursen, J., Hiebert, S.W., Grosveld, G., Downing, J.R., 1996. AML1, the target of
multiple chromosomal translocations in human leukemia, is essential for normal fetal liver
hematopoiesis. Cell 84, 321-330.

Paik, E.J., Zon, L.I., 2010. Hematopoietic development in the zebrafish. Int J Dev Biol 54, 1127-1137.

Pak, B., Schmitt, C.E., Choi, W., Kim, J.D., Han, O., Alsio, J., Jung, D.W., Williams, D.R., Coppieters,
W., Stainier, D.Y.R., Jin, S.W., 2020. Analyses of Avascular Mutants Reveal Unique Transcriptomic
Signature of Non-conventional Endothelial Cells. Front Cell Dev Biol 8, 589717.

Palis, J., Robertson, S., Kennedy, M., Wall, C., Keller, G., 1999. Development of erythroid and myeloid
progenitors in the yolk sac and embryo proper of the mouse. Development 126, 5073-5084.

Parant, J.M., George, S.A., Pryor, R., Wittwer, C.T., Yost, H.J., 2009. A rapid and efficient method of
genotyping zebrafish mutants. Dev Dyn 238, 3168-3174.

Pelster, B., Burggren, W.W., 1996. Disruption of hemoglobin oxygen transport does not impact oxygen-
dependent physiological processes in developing embryos of zebra fish (Danio rerio). Circulation
Research 79, 358-362.

Peng, J., Zhang, L., Drysdale, L., Fong, G.H., 2000. The transcription factor EPAS-1/hypoxia-inducible
factor 2alpha plays an important role in vascular remodeling. Proc Natl Acad Sci U S A 97, 8386-8391.

Pongratz, I., Antonsson, C., Whitelaw, M.L., Poellinger, L., 1998. Role of the PAS domain in regulation
of dimerization and DNA binding specificity of the dioxin receptor. Mol Cell Biol 18.

Prasch, A.L., Tanguay, R.L., Mehta, V., Heideman, W., Peterson, R.E., 2006. Identification of zebrafish
ARNT1 homologs: 2,3,7,8-tetrachlorodibenzo-p-dioxin toxicity in the developing zebrafish requires
ARNT1. Molecular pharmacology 69, 776-787.

Qian, F., Zhen, F., Ong, C., Jin, S.W., Meng Soo, H., Stainier, D.Y., Lin, S., Peng, J., Wen, Z., 2005.
Microarray analysis of zebrafish cloche mutant using amplified cDNA and identification of potential
downstream target genes. Dev Dyn 233, 1163-1172.

Reischauer, S., Stone, O.A., Villasenor, A., Chi, N., Jin, S.W., Martin, M., Lee, M.T., Fukuda, N.,
Marass, M., Witty, A., Fiddes, I., Kuo, T., Chung, W.S., Salek, S., Lerrigo, R., Alsio, J., Luo, S., Tworus,

D., Augustine, S.M., Mucenieks, S., Nystedt, B., Giraldez, A.J., Schroth, G.P., Andersson, O., Stainier,
D.Y., 2016. Cloche is a bHLH-PAS transcription factor that drives haemato-vascular specification.
Nature 535, 294-298.

Romano, S.N., Edwards, H.E., Souder, J.P., Ryan, K.J., Cui, X., Gorelick, D.A., 2017. G protein-
coupled estrogen receptor regulates embryonic heart rate in zebrafish. PLoS Genet 13, e1007069.

Rossi, A., Kontarakis, Z., Gerri, C., Nolte, H., Holper, S., Kruger, M., Stainier, D.Y., 2015. Genetic
compensation induced by deleterious mutations but not gene knockdowns. Nature 524, 230-233.

Ryan, H.E., Lo, J., Johnson, R.S., 1998. HIF-1 alpha is required for solid tumor formation and
embryonic vascularization. The EMBO journal 17, 3005-3015.

Scortegagna, M., Morris, M.A., Oktay, Y., Bennett, M., Garcia, J.A., 2003. The HIF family member
EPAS1/HIF-2alpha is required for normal hematopoiesis in mice. Blood 102, 1634-1640.

Shalaby, F., Rossant, J., Yamaguchi, T.P., Gertsenstein, M., Wu, X.F., Breitman, M.L., Schuh, A.C.,
1995. Failure of blood-island formation and vasculogenesis in Flk-1-deficient mice. Nature 376, 62-66.

Shu, L.P., Zhou, Z.W., Zhou, T., Deng, M., Dong, M., Chen, Y., Fu, Y.F., Jin, Y., Zhou, S.F., He, Z.X.,
2015. Ectopic expression of Hoxb4a in hemangioblasts promotes hematopoietic development in early
embryogenesis of zebrafish. Clinical and experimental pharmacology & physiology 42, 1275-1286.

Singh, K.P., Garrett, R.W., Casado, F.L., Gasiewicz, T.A., 2011. Aryl Hydrocarbon Receptor-Null Allele
Mice Have Hematopoietic Stem/Progenitor Cells with Abnormal Characteristics and Functions. Stem
Cells and Development 20, 769-784.

Smith, B.W., Rozelle, S.S., Leung, A., Ubellacker, J., Parks, A., Nah, S.K., French, D., Gadue, P.,
Monti, S., Chui, D.H., Steinberg, M.H., Frelinger, A.L., Michelson, A.D., Theberge, R., McComb, M.E.,
Costello, C.E., Kotton, D.N., Mostoslavsky, G., Sherr, D.H., Murphy, G.J., 2013. The aryl hydrocarbon
receptor directs hematopoietic progenitor cell expansion and differentiation. Blood 122, 376-385.

Souder, J.P., Gorelick, D.A., 2019. ahr2, But Not ahr1a or ahr1b, Is Required for Craniofacial and Fin
Development and TCDD-dependent Cardiotoxicity in Zebrafish. Toxicol Sci 170, 25-44.

Stainier, D.Y., Lee, R.K., Fishman, M.C., 1993. Cardiovascular development in the zebrafish. I.
Myocardial fate map and heart tube formation. Development 119, 31-40.

Stainier, D.Y., Weinstein, B.M., Detrich, H.W., 3rd, Zon, L.I., Fishman, M.C., 1995a. Cloche, an early
acting zebrafish gene, is required by both the endothelial and hematopoietic lineages. Development
121, 3141-3150.

Stainier, D.Y., Weinstein, B.M., Detrich, H.W., Zon, L.I., Fishman, M.C., 1995b. Cloche, an early acting
zebrafish gene, is required by both the endothelial and hematopoietic lineages. Development 121,
3141-3150.

Takubo, K., Goda, N., Yamada, W., Iriuchishima, H., Ikeda, E., Kubota, Y., Shima, H., Johnson, R.S.,
Hirao, A., Suematsu, M., Suda, T., 2010. Regulation of the HIF-1alpha level is essential for
hematopoietic stem cells. Cell stem cell 7, 391-402.

Tanguay, R.L., Andreasen, E., Heideman, W., Peterson, R.E., 2000. Identification and expression of
alternatively spliced aryl hydrocarbon nuclear translocator 2 (ARNT2) cDNAs from zebrafish with
distinct functions. Biochim Biophys Acta 1494, 117-128.

Thisse, C., Thisse, B., 2008. High-resolution in situ hybridization to whole-mount zebrafish embryos.
Nat Protoc 3, 59-69.

Thompson, M.A., Ransom, D.G., Pratt, S.J., MacLennan, H., Kieran, M.W., Detrich, H.W., 3rd, Vail, B.,
Huber, T.L., Paw, B., Brownlie, A.J., Oates, A.C., Fritz, A., Gates, M.A., Amores, A., Bahary, N., Talbot,
W.S., Her, H., Beier, D.R., Postlethwait, J.H., Zon, L.I., 1998. The cloche and spadetail genes
differentially affect hematopoiesis and vasculogenesis. Dev Biol 197, 248-269.

Tian, H., Hammer, R.E., Matsumoto, A.M., Russell, D.W., McKnight, S.L., 1998. The hypoxia-
responsive transcription factor EPAS1 is essential for catecholamine homeostasis and protection
against heart failure during embryonic development. Genes & development 12, 3320-3324.

Vogeli, K.M., Jin, S.W., Martin, G.R., Stainier, D.Y., 2006. A common progenitor for haematopoietic and
endothelial lineages in the zebrafish gastrula. Nature 443, 337-339.

Wang, Q., Stacy, T., Binder, M., Marin-Padilla, M., Sharpe, A.H., Speck, N.A., 1996. Disruption of the
Cbfa2 gene causes necrosis and hemorrhaging in the central nervous system and blocks definitive
hematopoiesis. Proceedings of the National Academy of Sciences of the United States of America 93,
3444-3449.

Weng, W., Nagai, H., Hamidi, S., Sheng, G., 2020. NPAS4L is involved in avian hemangioblast
specification. Haematologica.

Westerfield, M., 2000. The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio Rerio).
4th ed. University of Oregon Press, Eugene, OR.

Wu, D., Rastinejad, F., 2017. Structural characterization of mammalian bHLH-PAS transcription factors.
Curr Opin Struct Biol 43, 1-9.

Zhang, C., Featherstone, M., 2020. A zebrafish hox gene acts before gastrulation to specify the
hemangioblast. Genesis, e23363.
694

Gene	Allele	CRISPR target	Oligo 1	Oligo 2
arnt1	*bcm1*	GGATGGTTTATAGGTGCCAT<u>CGG</u>	TAGGATGGTTTATAGGTGCCAT	AAACATGGCACCTATAAACCAT
arnt1	*bcm2*	GGCGACTGTGAGGATGAGTA<u>TGG</u>	TAGGCGACTGTGAGGATGAGTA	AAACTACTCATCCTCACAGTCG
arnt2	*bcm3*	GGTCTCGGTCCGTCTAAGGA<u>GGG</u>	TAGGTCTCGGTCCGTCTAAGGA	AAACTCCTTAGACGGACCGAGA

698   Table 1. CRISPR target sequence and oligos used to generate *arnt1* and *arnt2* mutants. PAM sequence in

699   CRISPR target is underlined

Figure 1

Figure 1. *arnt1/2* double mutant embryos, but not *arnt1* or *arnt2 single* mutants, lack circulating blood, have cardiac edema and atrial enlargement. (A) bHLH-PAS transcription factors act as heterodimers, where a class I and indicated class II protein interact and bind DNA. Asterisks indicate proteins encoded by genes present in zebrafish but absent in humans. (**B,C**) Generation of zebrafish with mutations in arnt1 and arnt2 genes. The amino acid sequences for the zebrafish Arnt1 (**B**), and Arnt2 proteins (**C**) are represented by the black line. Basic helix-loop-helix (bHLH), Per-Arnt-Sim (PAS) domains, and the motif C-terminally located to the PAS (PAC) are represented by the labeled bubbles. The numbers below each domain indicate the starting and ending amino acid (AA) for each motif. The *arnt1^bcm1* mutation causes a premature stop codon to occur at AA139 and loss of the PAS-B, PAC, and majority of the PAS-A. The *arnt1^bcm2* mutation causes a premature stop codon to occur at AA250 and loss of the PAS-B and PAC domains (**B**). The *arnt2^bcm3* mutation causes a premature stop codon to occur at AA287 and loss of the PAS-B and PAC domains (**C**). (**D-I**) Live embryos from *arnt1^bcm2/+;arnt2^bcm3/+* heterozygous parents were imaged at 2 days post fertilization (dpf) prior to genotyping. All double homozygous *arnt1^bcm2/bcm2; arnt2^bcm3/bcm3* (*arnt1/2* mutant) embryos showed cardiac edema and atrial enlargement (**G** and **G'**). The wildtype (**D** and **D'**), *arnt1^bcm2/bcm2* (*arnt1* mutant) (**E** and **E'**), and *arnt2^bcm3/bcm3* (*arnt2* mutant) (**F** and **F'**) siblings of the *arnt1/2* mutants showed normal heart development. Black arrows in panels D'-G' indicate the atrium and ventricle of the heart. Additionally, all siblings of *arnt1/2* mutants had circulating blood (**H** and **M1**) while *arnt1/2* mutants lacked visible circulting blood cells (**I** and **M2**). The white arrows in panel **H** indicate blood cells in the dorsal aorta. Images are representative of 7 different cluches, each cluch contained between 20-300 embryos. For all images of zebrafish embryos (D-I) anterior is towards the right and doral is towards the top of the image. The scale bar in Panel D applies to Panels D-G and is 1000 μm in lenght. The scale bar in Panel D' apples to Panels D'-G' and is 100 μm in length.

705   Figure 2

Figure 2. *arnt1/2* double homozygous mutants show reduced levels of *flk-1*-positive endothelial cells and *runx-1*-positive cells in blood island (A-H). Embryos were probed for *flk-1* (A-D) or *runx-1* (E-H) using whole-mount in situ hybridization at 2 days post fertilization (dpf). Wild-type embryos were derived from an in-cross of laboratory strain AB parents (A and E). *arnt1* mutant embryos were derived from *arnt1;arnt2* heterozygous parents or are maternal zygotic derived from *arnt1* homozygous parents (MZ*arnt1*). *arnt2* mutant embryos were derived from heterozygous parents. *arnt1/2* double homozygous mutant embryos (D and H) were derived from *arnt1;arnt2* double heterozygous parents and identified by cardiovascular phenotype prior to fixation. Panels A'-D' are enlarged images of each embryo's trunk near the end of the yolk extension (area indicated by red box in panels A-D). (A'-D') Black arrows indicate *flk-1* staining in the dorsal longitudinal anastomotic vessels (DLAV), white arrows indicate the dorsal aorta and caudal vein (DA/CV), arrowheads indicate inter-segmental vessels. (E-H) *runx-1* labeling near the end of the yolk extension, arrows indicate blood island region. n=2-5 clutches for each genotype, 10-100 embryos per clutch. All embryos are oriented with anterior to the right and dorsal towards the top. The scale bar in D applies to A-D, the scale bar in D' applies to A'-D', and the scale bar in H applies to E-H. All scale bars are 10 μm.

Figure 3. arnt1/2 mutants show decreased etv2 and tal1 expression in the posterior lateral plate mesoderm (PLPM) at 4-6 somite stages. Whole-mount in situ hybridization for *etv2* and *tal1* expression was performed on 4-6 somite stage embryos which were offspring of *arnt1^bcm2/+; arnt2^bcm3/+* adult zebrafish. (A) Expression of *etv2* was observed in the PLPM in 92% of embryos (n=477 of 516 total embryos). (B) Limited or no expression of *etv2* was observed in 8% of embryos (n=39 of 516), consistent with the expected Mendelian ratios for *arnt1/2* mutants (6.25%). 24 of the 39 presumptive *arnt1/2* mutants were genotyped and 24/24 embryos were confirmed *arnt1/2* mutants. (C) Expression of *tal1* was observed in 94% of embryos (n=624 of 665 total embryos). (D) Limited or no expression of *tal1* was observed in 6% of embryos (n=41 of 665), consistent with the expected Mendelian ratios for *arnt1/2* mutants (6.25%). 17 of the 41 presumptive *arnt1/2* mutants were genotyped and 14/17 embryos were confirmed *arnt1/2* mutants. All embryos are oriented with dorsal towards the top of the image and posterior is closest to the reader. Scale bar is 10 μm and applies to all images.

711   Figure 4

Figure 4. Arnt1 and Arnt2 co-immunoprecipitate with Npas4l. (A) HEK293T cells were transfected with zebrafish Npas4l-V5 and Arnt1-myc. Cells were homogenized (input) and immunoprecipitated (IP) with antibodies aganist V5 or an immunoglobulin G control (IgG). Samples were run on an immunoblot and probed using myc antibodies (anti-myc) to detect the presence of Arnt1-myc (top) or samples were probed using V5 antibodies (anti-V5) to verify sucessful pulldown of Npas4l-V5 (bottom). **(B)** HEK293T cells were transfected with zebrafish Npas4l-V5 and Arnt2-HA. Cells were homogenized and immunoprecipitated with antibodies aganist HA or an IgG control. Samples were run on an immunoblot and probed using V5 antibodies (anti-V5) to detect the presence of Npas4l-V5 (top) or samples were probed using HA antibodies (anti-HA) to verify sucessful pulldown of Arnt2-HA (bottom).

Figure 5. Mutation in *arnt1,* but not *arnt2,* prevents AHR2-dependent TCDD toxicity. Wild type or mutant larvae were exposed to 0.1% DMSO (**A, B, C**), or 2,3,7,8-tetrachlorodibenzodioxin (TCDD) (**D, E, F**) from 1-3 dpf and imaged at 3 dpf. Wildtype embryos exposed to TCDD (**D**) lack an S-shaped cardiac loop and exhibit cardiac edema, unlike their vehicle treated siblings (**A**). *arnt1* homozygous mutant larvae (*arnt1bcm2/bcm2*) treated with vehicle and TCDD (**B,E**) show normal heart looping and do not have cardiac edema. *arnt2* homozyous mutant larvae (*arnt1bcm3/bcm3*) treated with vehicle (**C**) show expected heart looping similar to wild type siblings, but *arnt2* homozyous mutant larvae treated with TCDD (**F**) lack an S-shaped cardiac loop similar to wild type and exhibit cardiac edema similar to TCDD treated wild-type embryos. Black arrows indicate location of the heart. Embryos were genotyped following imaging and representative images are shown. TCDD treated wildtype siblings of *arnt1* mutant embryos, n= 9 embryos across 3 clutches; TCDD treated *arnt1* mutant larvae, n=21 larvae across 3 clutches. TCDD treated wildtype siblings of *arnt2* larvae, n=19 embryos across 3 cluches; TCDD treated *arnt2* mutant larvae, n=22 embryos across 4 cluches. All embryos are oriented with anterior to the right and dorsal towards the top of the image. Scale bar is 100 μm.