Hemato-vascular specification requires *arnt1* and *arnt2* genes in zebrafish embryos

Authors: Hailey E. Edwards, Mary Jane Edgington, Jaclyn P. Souder, Daniel A. Gorelick

Center for Precision Environmental Health, Department of Molecular and Cellular Biology, Baylor

College of Medicine, Houston, Texas, USA

Correspondence to DAG gorelick@bcm.edu

ABSTRACT

During embryonic development, a subset of cells in the mesoderm germ layer are specified as hemato-

vascular progenitor cells, which then differentiate into endothelial cells and hematopoietic stem and

progenitor cells. In zebrafish, the transcription factor *npas4l*, also known as *cloche*, is required for the

specification of hemato-vascular progenitor cells. However, it is unclear if *npas4l* is the sole factor at the

top of the hemato-vascular specification cascade. Here we show that *arnt1* and *arnt2* genes are

required for hemato-vascular specification. We found that *arnt1;arnt2* double homozygous mutant

zebrafish embryos (herein called *arnt1/2* mutants), but not *arnt1* or *arnt2* single mutants, lack blood

cells and most vascular endothelial cells. *arnt1/2* mutants have reduced or absent expression of *etsrp*

and *tal1*, the earliest known endothelial and hematopoietic transcription factor genes. *npas4l* and the

arnt genes are PAS domain-containing bHLH transcription factors that function as dimers. We found

that Npas4l binds both Arnt1 and Arnt2 proteins *in vitro*, consistent with the idea that PAS domain-

containing bHLH transcription factors act in a multimeric complex to regulate gene expression. Our

results demonstrate that *npas4l*, *arnt1* and *arnt2* act together to regulate endothelial and hematopoietic

cell fate, where each gene is necessary, but by itself not sufficient, to drive hemato-vascular

specification. Our results also demonstrate that *arnt1* and *arnt2* act redundantly in a transcriptional

complex containing *npas4l*, but do not act redundantly when interacting with another PAS domain-

containing bHLH transcription factor, the aryl hydrocarbon receptor. Altogether, our data enhance our

understanding of hemato-vascular specification and the function of PAS domain-containing bHLH

transcription factors.

INTRODUCTION

Endothelial and hematopoietic cells are derived from a common mesodermal progenitor (Huber

et al., 2004; Kataoka et al., 2011; Liao et al., 2000). In mice and zebrafish, fully multipotent

hematopoietic stem cells (HSCs) are derived from specialized endothelial cells, called hemogenic

endothelium (Hirschi, 2012; Kobayashi et al., 2014; Ramírez-Bergeron et al., 2004) The first hemato-

vascular progenitors were traced back to the lateral plate mesoderm of zebrafish and in the

extraembryonic yolk sac mesoderm of mice (Lee et al., 1994; Palis et al., 1999; Stainier et al., 1993).

The complete mechanism of the specification of these hemato-vascular progenitors from the

multipotent mesoderm remains elusive in mammals.

Many genes involved in embryonic hematopoiesis and vascular development are evolutionally

conserved between zebrafish and mammals like mice and humans (Begley et al., 1989; Casie Chetty et

al., 2017; Dooley et al., 2005; Ferrara et al., 1996; Herbert and Stainier, 2011; Kalev-Zylinska et al.,

2002; Kataoka et al., 2011; Okuda et al., 1996; Wang et al., 1996). In zebrafish, the *neuronal PAS4-like*

gene (*npas4l*), formerly known as *cloche,* is the earliest known transcription factor to drive the

specification of hemato-vascular progenitors (Reischauer et al., 2016). *npas4l* mutants have a

distinctive phenotype as they lack most endothelial and all hematopoietic cells (Stainier et al., 1995;

Thompson et al., 1998).

Npas4l is a basic helix-loop-helix Per-Arnt-Sim (bHLH-PAS) transcription factor and is thought

to be the master regulator of hemato-vascular specification (Reischauer et al., 2016). bHLH-PAS

transcription factors are classified as Class I or Class II, which must dimerize to regulate gene

transcription or repression (**Figure 1A**) (Card et al., 2005; Edwards and Gorelick, 2022; Pongratz et al.,

1998). Npas4l is a Class I bHLH-PAS protein, but its Class II dimerization partner during hemato-

vascular progenitor specification is not known. We hypothesized that Arnt1 or Arnt2 is the Class II

dimerization partner of Npas4l in zebrafish because *arnt1* and *arnt2* genes are expressed at the same

stages of embryo development as hemato-vascular specification (Baranasic et al., 2022; Prasch et al.,

2006; Reischauer et al., 2016; Tanguay et al., 2000).

To test this hypothesis, we examined hemato-vascular development in zebrafish with predicted

loss-of-function mutations in *arnt1* and *arnt2* genes. We find that Arnt1 and Arnt2 are essential for

hemato-vascular specification. Double homozygous *arnt1;arnt2* zebrafish embryos (herein called

arnt1/2 mutants) phenocopy *npas4l* mutants and show lack of circulating blood, absence of

hematopoietic stem and progenitor cells, and dramatic reduction in the number of vascular endothelial

cells. We find that expression of *npas4l*-target genes requires *npas4l* and *arnt1* or *arnt2*. Additionally,

we show that zebrafish Npas4l protein interacts with both zebrafish Arnt1 and zebrafish Arnt2 proteins

in vitro. Together, our results suggest that Arnt1 and Arnt2 act redundantly and form a transcriptional

complex with Npas4l that sits atop the hemato-vascular specification cascade.

MATERIALS AND METHODS

Zebrafish

Adult zebrafish were raised at 28.5°C on a 14-h light, 10-h dark cycle in the BCM Zebrafish Research

Facility in an Aquaneering recirculating water system (Aquaneering, Inc., San Diego, CA) and a

Tecniplast recirculating water system (Tecniplast USA, Inc., West Chester, PA). Wild-type zebrafish

were AB strain (Westerfield, 2000) and *arnt1* and *arnt2* mutant lines were generated on the AB strain.

Once the mutant lines were established, they were bred onto the *Tg(Fli1:EGFP)^{y1}* transgenic

background (Lawson and Weinstein, 2002). All procedures were performed in accordance with and

approved by the BCM Institutional Animal Care and Use Committee.

Embryo collection

Adult zebrafish were allowed to spawn naturally in pairs or in groups. Embryos were collected in

intervals of 20 minutes to ensure precise developmental timing or staged following collection, placed in

60 cm^2 Petri dishes at a density of no more than 100 per dish in E3B media (60X E3B: 17.2g NaCl,

0.76g KCl, 2.9g CaCl$_2$-2H$_2$O, 2.39g MgSO$_4$ dissolved in 1L Milli-Q water; diluted to 1X in 9L Milli-Q

water plus 100 µL 0.02% methylene blue), and then stored in an incubator at 28.5°C on a 14-h light,

10-h dark cycle.

Genotyping adult zebrafish

Genomic DNA was isolated from tail biopsies from individual adult zebrafish and incubated in 50 µL

ELB (10 mM Tris pH 8.3, 50 mM KCl, 0.3% Tween 20) with 0.5 µL proteinase K (800 U/ml, NEB) in 96

well plates, one sample per well, at 55°C for 8 hours. Proteinase K was inactivated by incubation at

98°C for 10 minutes and DNA was stored at -20°C. Genotyping was performed by PCR and high-

resolution melting curve analysis (HRMA) as described (Parant et al., 2009; Romano et al., 2017). All

melting curves were generated with a Bio-Rad CFX96 or CFX Opus 96 Real-Time System over a 70-

95°C range and analyzed with the Bio-Rad CFX Manager 3.1 or the Bio-Rad CFX Maestro 4.1

software. All mutations were confirmed by TA cloning and sequencing. All primers used for HRMA can

be found in **Table S1**.

CRISPR-Cas9 Mutant Generation

Cas9 mRNA and gRNAs for *arnt1* and *arnt2* mutants were generated as previously described (Romano

et al., 2017; Souder and Gorelick, 2019). Cas9 mRNA was transcribed from a linearized pT3TS-nCas9n

plasmid, Addgene #46757 (Jao et al., 2013). The target sequences were identified using CHOPCHOP

(Labun et al., 2016; Labun et al., 2019). Oligonucleotides were annealed to each other and cloned into

a pT7-gRNA plasmid, Addgene #46759 (Jao et al., 2013). Oligonucleotides utilized are shown in **Table**

S1. gRNAs were synthesized from plasmids using a MEGAshortscript T7 Transcription Kit (Invitrogen

AM1354) and purified.

One-cell-stage embryos were injected using glass needles pulled on a Sutter Instruments

Fleming/Brown Micropipette Puller, model P-97 and a regulated air-pressure micro-injector (Harvard

Apparatus, NY, PL1–90). Each embryo was injected with a 1nL solution containing one gRNA for *arnt1*

and one gRNA for *arnt2* (30 ng/µL per target), Cas9 mRNA (150 ng/µL), and 0.1% phenol red. Mixtures

were injected into the yolk of each embryo. Approximately 100 injected embryos per gRNA pair were

raised to adulthood and crossed to AB zebrafish to generate F1 embryos. F1 offspring with heritable

mutations were sequenced to identify mutations predicted to cause loss of function.

Live Imaging

Embryos were imaged with a Nikon SMZ25 microscope equipped with a Hamamatsu ORCA-Flash4.0

digital CMOS camera, or with a Nikon SMZ18 microscope equipped with a Nikon DS-Fi3 camera.

Images were equally adjusted for brightness and contrast in Adobe Photoshop CC 2020. Embryos and

larvae were anesthetized with 0.04% tricaine and imaged in Petri dishes containing E3B. All embryos

were genotyped after imaging.

Whole-mount *in situ* hybridization (WISH)

We used the following DNA plasmids to generate RNA probes: *egfl7, lmo2* (Sun et al., 2018), *etsrp* (gift

of Dr. Nathan Lawson, Addgene plasmid # 49005) (Moore et al., 2013)*, tal1* (gift of Dr. Arne Lekven)*,*

npas4l (gift of Dr. Leonard Zon) (Reischauer et al., 2016)*, runx1* and *kdrl* (gifts of Dr. Mary Goll) (Li et

al., 2015). For *lmo2* and *egfl7,* RNA was extracted from 10 pooled 1 dpf wild-type embryos using TRIzol

Reagent (ThermoFisher Scientific, 15596026) and purified using RNA clean & concentration kit (Zymo

Research, R1019). cDNA was synthesized using iScript cDNA synthesis (BioRad, 1708890). *lmo2* and

egfl7 were PCR amplified and Topo TA cloned into pCRII vector (Invitrogen, K4600-01). Primers used

to amplify *lmo2* and *egfl7* can be found in **Table S1**. All DNA clones were verified by sequencing.

Digoxigenin-labeled antisense RNA probes were synthesized using T7 or SP6. Colorimetric *in situ*

hybridization (ISH) was performed on fixed zebrafish embryos and larvae as previously described, with

5% dextran in the hybridization buffer (Lauter et al., 2011; Thisse and Thisse, 2008). Following ISH,

embryos were washed several times in 1x PBST (phosphate buffered saline, 0.01% Tween), cleared

with glycerol, or mounted in 3% methylcellulose on a class coverslip, and imaged on either a Nikon

SMZ18 stereoscopic microscope or an inverted Nikon ECLIPSE Ti2-E microscope. Embryos were then

genotyped. If embryos were older than 24 hpf, genomic DNA was extracted in 20 μL ELB (10 mM Tris

pH 8.3, 50 mM KCl, 0.3% Tween 20) with 0.2 μL proteinase K (800 U/ml, NEB) in 96 well plates, one

sample per well, at 55°C for 2 hours. If the embryos were 24 hpf or younger, they were first de-

crosslinked in 300 mM NaCl for 4 hours at 65°C and held at 4-8°C until DNA could be extracted.

Genomic DNA was extracted using the HotSHOT protocol (Dobrzycki et al., 2018). Briefly, embryos

were suspended in 10-20 μL of lysis buffer (50 mM NaOH) and incubated at 95°C for 30 minutes,

cooled to 4°C, then 1/10th the volume of neutralization buffer (1M Tris-HCl, pH 8.0) was added. Embryo

lysate was incubated at 4°C overnight before starting PCR reactions. DNA regions of interest were

amplified using one of two methods. The genes of interest were PCR amplified using JumpStart

REDTaq ReadyMix (Sigma-Milipore, P0982) according to the manufacturer's protocols. Alternatively,

the DNA regions of interest were PCR amplified in two successive reactions using HS Taq (Takara,

R007). The first PCR amplified a larger fragment of *arnt1* (about 700 bp) or *arnt2* (about 970 bp). Then

the product of the larger fragment was used as a template to amplify a smaller fragment of *arnt1* (about

400 bp) or *arnt2* (about 650 bp). After PCR amplification, the PCR products were sequenced to

determine the genotype. All primer sequences can be found in **Table S1.**

Whole-mount *in situ* hybridization chain reaction (WIHCR)

HCR Antisense RNA probe sets for *arnt1, arnt2,* and *npas4l,* HCR amplifiers, and HCR buffers were

obtained from Molecular Instruments, Inc (https://www.molecularinstruments.com/). Each gene had a

probe set size of 20. WIHCR was performed on fixed AB embryos between 4-6 somites, as previously

described (Choi et al., 2018; Ibarra-García-Padilla et al., 2021). Following HCR, embryos were washed

several times in 1x PBST and cleared in glycerol. All embryos were mounted in 75% glycerol and

imaged on a Nikon Ti2-E inverted microscope with Yokogawa CSU-W1 spinning disk confocal and

Photometrics Prime95b sCMOS camera.

WIHCR Image Processing and Analysis

All images used in processing and analysis were captured on the same day, using identical acquisition

controls. We selected only images where a clear, bilateral view of the presumptive lateral plate

mesoderm (LPM) and presumptive notochord was available (N=6 embryos from a single clutch, and

N=4 negative control embryos from the same clutch). Images of anterior and posterior views of the

embryos were collected. To calculate the number of 3D objects in a stack, we used Image J, FIJI

version 2.3.0/1.53q (Schindelin et al., 2012), and the 3D-OC plug-in (Bolte and Cordelières, 2006). A

custom threshold for each channel was applied to each image to filter out background autofluorescence

seen in the negative control embryos. The number of detected objects for probed samples was then

normalized to negative control by subtracting the average number of detected objects from control

embryos from the number of detected objects in probed embryos. We used ImageJ to calculate the

adjusted integrated intensity of probed and control embryos (Schindelin et al., 2012). First, maximum

intensity projections (MIP) of the z-stacks were created, then the region of interest (ROI) was drawn

around the areas with the highest concentration of *npas4l* expression (presumptive LPM) or around the

corresponding region of the presumptive LPM in the negative control embryos. The percent area of

pixels and the mean gray value for each channel were calculated and limited to the threshold. The

adjusted integrated density was then calculated by multiplying the percent area and the mean gray

value.

O-dianisidine labeling of blood cells

Zebrafish embryos were stained at approximately 72 hours post fertilization. Embryos were

dechorionated using 0.2% pronase in E3B media for 5-10 minutes and were gently triturated with a

1000µl pipette. Embryos were transferred to glass vials and submerged in o-dianisidine working

solution, prepared as described (Kafina and Paw, 2018). Vials were kept in the dark for 5 minutes

during staining. Samples were then fixed in 4% formaldehyde for 30 minutes at room temperature.

Embryos were washed in PBS-Tween 0.1% before mounting for imaging in 3% methylcellulose on a

glass dish or coverslip. Embryos were genotyped after imaging.

Plasmid synthesis for microinjection and *in vitro* transfection

Cloning of plasmids for expression in cultured cells was performed as follows. For npas4l-V5, we took

pCS2+npas4lORF (gift of Dr. Leonard Zon) that contains the open reading frame (ORF) of zebrafish

npas4l downstream of the CMV promoter. This plasmid was modified to include a C-terminal V5 tag

using Q5 Site-Directed Mutagenesis Kit (New England Biolabs, #E0554S). For arnt1-myc, we

commercially synthesized a plasmid containing CMV promoter driving a codon-optimized zebrafish

arnt1 ORF (ENSDART00000081852.5) with a C-terminal myc tag (Twist Bioscience, San Francisco,

CA). For arnt2-myc, we commercially synthesized a gene block of the zebrafish *arnt2* ORF

(ENSDART00000158162.2, Integrated DNA Technologies). The gene block was integrated into a

plasmid containing a CMV promoter using pcDNA3.3-TOPO TA cloning kit (Invitrogen, K8300-01) in

accordance with manufacturer's instructions. A C-terminal HA tag was added using Q5 Site-Directed

Mutagenesis Kit (New England Biolabs, #E0554S). For microinjection, an SP6 promoter was inserted

upstream of the transcriptional start using Q5 Site-Directed Mutagenesis Kit (New England Biolabs,

#E0554S). All plasmids were confirmed by sequencing.

Cell Culture and Transient Transfection

HEK293T cells (gift from Dr. Charles Foulds), were thawed and maintained at 37°C in humidified 5%

CO_2 atmosphere incubator. Cells were cultured in DMEM supplemented with 10% fetal bovine serum

(FBS; Gibco, 2062564) and 1% penicillin-streptomycin (Gibco, 15140-122). After 24 hours of passaging

in a 100mm plate at a density of 2.2×10^6 cells per dish and reaching at least 70 confluency, the media

was changed to DMEM supplemented with 1% dialyzed FBS. Cells were transiently transfected with 12

µg of total DNA and Lipofectamine 2000 according to the manufacturer's instructions (Invitrogen,

2400439) with *npas4l-v5* together with either *arnt1-myc* or *arnt2-HA* plasmid DNA.

Co-immunoprecipitation and Western blot

Transiently transfected cells were washed with ice cold PBS and lysed in 500 µL of 1x RIPA buffer

containing Pierce Protease (ThermoScientific, A32955). Cells were sonicated in 6 x 30 second intervals

using a Bioruptor 300 Machine (Diagenode), then centrifuged for 10 minutes at 10,000g at 4°C. The

lysate (supernatant) was removed from the pellet and placed in fresh 1.5mL Eppendorf tube. Cell lysate

was precleared with Protein A beads (Invitrogen #159 18-014) for 30 minutes 4°C, incubated with either

5 µg of V5 antibody (ThermoFisher Scientific R960-25) or myc antibody (SantaCruz, SC-40) diluted

1:200 for 1 hour, then immunoprecipitated with 20 µL of Protein A beads for 1 hour at 4°C. The beads

were washed three times with RIPA buffer and once with PBS and finally suspended in 30µL of PBS.

Equal volume of 2x Laemmli Sample Buffer (BioRad, cat# 1610737) was added to each sample and

boiled for 5 minutes at 95°C then placed on ice. Samples were run on 4-20% Mini-PROTEAN TGX gels

(BioRad, cat# 4561093) and transferred onto PVDF membrane using Trans-Blot Turbo Transfer

System (BioRad) using the pre-defined Mixed MW program. The membranes were washed several

times with 1x Tris-Buffered Saline, 0.1% Tween 20 (TBST) (GBiosciences, R043) and incubated for 1

hour with agitation in 5% milk in 1x TBST. Membranes were probed with 1:1000 V5-HRP

(ThermoFisher Scientific, R960-25), 1:1000 myc-HRP (SantaCruz, SC-40), or 1:200 HA (Cell Signaling,

3724) antibodies rocking overnight at 4°C. Membranes were washed 5 times with 1xTBST on orbital

rocker for 15 minutes at room temperature. Membranes probed with the anti-HA antibody were then

probed with 1:10,000 IRDye 800CW donkey anti-rabbit (Licor, 926-32213) in 2.5%-5% milk in 1x TBST

for 1 hour and then washed multiple times with 1xTBST on orbital rocker for 5 minutes prior to imaging.

Membranes probed with HRP conjugated antibodies were developed using Clarity™ Western ECL

Substrate (BioRad, 170-5061). Membranes were imaged using BioRad ChemiDoc™ MP Imaging

System.

Zebrafish embryo injection

Embryo injections were performed as described (Westerfield, 2000) with the following modifications.

Plasmids containing either *npas4l, arnt1,* or *arnt2* were linearized by restriction enzyme digestion.

mRNA was transcribed using the mMessage mMachine kit (Invitrogen, AM1340) and purified with the

RNA clean & concentration kit (Zymo Research, R1019). Embryos were microinjected at the 1-cell

stage with 1 nL total volume containing either *npas4l* mRNA (50 pg mRNA) or a mixture of *arnt1* and

arnt2 mRNA (50 pg each for 100 pg total mRNA). All solutions were injected into the yolk.

TCDD exposure

At 1-day post fertilization (dpf), embryos were exposed to 10 ng/mL 2,3,7,8-tetrachlorodibenzodioxin

(TCDD, AccuStandard Inc, D404N) or vehicle (0.1% DMSO). All treatments were performed on a single

clutch of at least 20 embryos. Embryos treated with vehicle were from the same clutches as embryos

treated with TCDD. Chemical exposures were performed on at least 2 clutches per genotype. Embryos

were raised at 28.5°C on a 14-h light, 10-h dark cycle until 3 dpf, when they were imaged and then

genotyped.

Statistical Analysis

All Statistical analyses were performed using Prism version 9.4.1 (GraphPad Software, Boston, MA).

Mendelian Frequencies Analysis: For the Chi-squared analysis of embryo and adult genotype ratios we

eliminated all 0 values from the analysis.

WISH Phenotype Frequency Analysis: For WISH images of *arnt1/2* mutants**,** we used a binomial test to

compare the observed distribution of embryos with the reduced expression phenotype to the calculated

expected number of *arnt1/2* mutants. Additionally, we performed a Spearman rank correlation analysis

on genotyped embryos to determine if there was a correlation between the number of mutant alleles

and the observed phenotype. Each genotype was assigned a number 0-4: with 0 indicating no mutant

alleles (wild-type siblings), 1 indicating one mutant allele (*arnt1* heterozygous and *arnt2* heterozygous

siblings), 2 indicating two mutant alleles (*arnt1* homozygous, *arnt2* homozygous, or *arnt1/2*

heterozygous siblings), 3 indicating three mutant alleles (*arnt1* homozygous; *arnt2* heterozygous and

arnt1 heterozygous; *arnt2* homozygous siblings), and 4 indicating 4 mutant alleles (*arnt1/2* mutants).

The overserved phenotypes were assigned a number value: normal expression was coded as a 1 and

reduced expression was coded as 0.

WIHCR Image Analysis: For WIHCR image analysis, the variance was not assumed for *npas4l* and

arnt2 samples. *arnt1* samples did not have a normal distribution, thus a non-parametric Mann-Whitney

test was performed. We were able to identify an outlier in the *arnt1* integrated intensity data using

Grubb's test ($\alpha=0.05$), however, when this outlier was removed from analysis there was no change in

significance or Mann-Whitney U value.

TCDD Treatment Analysis: For TCDD treated embryos, unpaired Welch's *t*-tests were performed to

compare the average number of embryos with cardiotoxicity to the average number of embryos with a

normal phenotype from a given genotype.

    RESULTS

arnt1/2 **homozygous mutants lack circulating blood**

To determine if *arnt1* or *arnt2* is required for hemato-vascular development, we generated

heritable mutant alleles for *arnt1 (arnt1^bcm1^* and *arnt1^bcm2^*) and *arnt2 (arnt2^bcm3^*) using CRISPR/Cas9. For

*arnt1^bcm1^*, we generated zebrafish with a 5 bp insertion in exon 5, which results in a frameshift at amino

acid (AA) 126 and a premature stop codon in the first PAS domain at AA139. For *arnt1^bcm2^*, we

generated zebrafish with an 11 bp deletion in exon 10, which results in a frameshift at AA230 and a

premature stop codon between the two PAS domains at AA250 (**Figure 1B**). We did not observe any

phenotypic differences between homozygous *arnt1^bcm/bcm1^* and *arnt1^bcm2/bcm2^* mutant embryos or larvae,

these fish will herein be called *arnt1* mutants. During embryogenesis and larval development, *arnt1*

mutants appear to have a normal phenotype and are indistinguishable from age-matched wild-type

larvae (**Figure 1D-E**). *arnt1* mutants are viable to adulthood and are fertile, consistent with previously

published *arnt1* mutants (Marchi et al., 2020). Maternal zygotic *arnt1* mutants did not show a

developmental phenotype and were indistinguishable from wild-type or zygotic *arnt1* mutants. For

*arnt2^bcm3^*, we generated zebrafish with a 3 bp deletion and 1 bp insertion in exon 9, which resulted in a

missense mutation at AA286 and a premature stop codon between the two PAS domains at AA287

(**Figure 1C**). During embryogenesis and larval development, *arnt2^bcm3/bcm3^* embryos, herein called *arnt2*

mutants, were indistinguishable from wild-type embryos (**Figure 1D-F**). *arnt2* mutants were viable at 5

days post fertilization (dpf) but did not survive to adulthood, consistent with other independently

generated *arnt2* mutant zebrafish lines (Hill et al., 2009; Löhr et al., 2009).

Because Arnt1 and Arnt2 can interact with multiple Class I bHLH-PAS proteins (**Figure 1A**), we

examined if embryos that lacked both *arnt1* and *arnt2* had a hemato-vascular phenotype. To generate

these embryos, we crossed *arnt1^bcm2/+^;arnt2^bcm3/+^* adult fish to each other. We observed all genotypes

at the expected Mendelian ratios in offspring at 24 to 48 hours post fertilization (**Table 1**). However,

*arnt1^bcm2/bcm2^; arnt2^bcm3/bcm3^* embryos, herein called *arnt1/2* mutants, showed a clear phenotype (**Figure**

1G). These embryos lack all circulating blood (**Figure 1H-L, and Movies 3, 4, 7, 8**). All other

intermediate *arnt1;arnt2* genotypes, e.g., *arnt1$^{bcm2/+}$;arnt2$^{bcm3/+}$*, appeared phenotypically wild-type

(**Figure 1H, J, K, and Movies 1, 2, 5, 6**). These results suggest that *arnt1* and *arnt2* function

redundantly because only the *arnt1/2* double homozygous mutants show a phenotype. Additionally, the

arnt1/2 mutant phenotype appears identical to the previously described *npas4l* mutant phenotype, also

known as *cloche* (Reischauer et al., 2016; Stainier et al., 1995). Like *npas4l* mutants, *arnt1/2* mutants

exhibit pericardial edema and have an enlarged heart atrium, likely a secondary effect of the lack of

blood and endocardial cells (**Figure 1G, Movies 3, 7**). Taken together, these data suggest that *arnt1*

and *arnt2* have a similar function as *npas4l*.

Next, we examined the frequencies of the genotypes in adult fish from *arnt1$^{bcm2/+}$; arnt2$^{bcm3/+}$*

parents. Given that *arnt2* mutants are not viable as adults, these adults can be one of six possible

genotypes. Interestingly, we did not observe the expected Mendelian ratios of alleles in these adult fish

(**Table 2**; X^2(df=5, N=138) = 45.57, p<0.0001, N=138 offspring from two different clutches). There is a

reduction in the number of observed *arnt1$^{bcm2/bcm2}$;arnt2$^{bcm3/+}$* heterozygous adults compared to the

expected number. We hypothesize that the number of *arnt1* and *arnt2* wild-type alleles correlates with

the likelihood of juvenile fish surviving to adulthood.

***arnt1/2* mutants have reduced expression of hematopoietic and endothelial cells**

We hypothesized that the Arnt proteins are dimerization partners with Npas4l and that *arnt1/2*

mutants would exhibit the same phenotypes as *npas4l* mutants. Therefore, we examined *arnt1/2*

mutant embryos for additional phenotypes found in *npas4l* mutants. Previous studies determined that

npas4l mutants lack almost all endothelial cells, except a population of endothelial cells located in the

lower trunk and caudal tail of the larvae (Liao et al., 1997; Reischauer et al., 2016; Stainier et al., 1995).

To determine if our *arnt1/2* mutants showed a similar reduction in endothelial cells, we performed

whole-mount *in situ* hybridization (WISH) on 2 dpf embryos for *kdrl* (*flk-1*). *kdrl*, a widely used marker of

endothelial cells in blood vessels, is essential for the development of blood vessels in mice and

zebrafish (Dumont et al., 1995; Liao et al., 1997; Shalaby et al., 1997). We found that *arnt1* mutants,

arnt2 mutants, and wild-type embryos showed clear expression of *kdrl* in blood vessels at 2 dpf (**Figure**

2A-C, N=2-5 biological replicates per genotype per probe, 10-100 embryos examined per biological

replicate). In contrast, *arnt1/2* mutants had a reduced number of *kdrl*-positive cells at 2 dpf (**Figure 2D**).

The *kdrl*-positive cells present in the *arnt1/2* mutant were in the caudal tail and likely comprised a

portion of the caudal vein (**Figure 2D'**). To determine if this population of cells was present earlier in

development we bred *arnt1^{bcm2/+}; arnt2^{bcm3/+}* onto a Tg(*fli1:egfp^{y1}*) transgenic background, which labels

all blood vessels and the pharyngeal arches of the developing embryos. At 1 dpf, *arnt1* mutants, *arnt2*

mutants, and wild-type embryos had GPF-positive vasculature structures (**Figure 2I-K,** N=2 biological

replicates, with 21-131 embryos per clutch, 232 embryos total). Within the tails of these embryos the

dorsal aorta, caudal vein, and intersegmental vessels are clear and present (**Figure 2I'-K'**). In contrast,

arnt1/2 mutants had reduced GFP-positive vascular structures (**Figure 2L).** Within these embryos, only

the extreme caudal vein was fluorescent (**Figure 2L'**). This population of endothelial cells is also seen

in *npas4l* mutants (Liao et al., 1997). Together these results suggest that *arnt1* or *arnt2* is required for

npas4l-dependent blood vessel development.

In addition to a clear reduction of endothelial cells, *npas4l* mutants show a reduction in the

number of hematopoietic stem cells (HSC) (Stainier et al., 1995; Thompson et al., 1998). To determine

if *arnt1/2* mutants contain HSCs, we examined *runx1* expression by WISH. *runx1* is a marker of HSCs

and is required for definitive hematopoiesis in zebrafish (Kalev-Zylinska et al., 2002). At 48 hpf, *runx1*-

positive HSCs are known to localize to the area between the dorsal aorta and the posterior caudal vein

(Lam et al., 2009; North et al., 2009). At 2 dpf, *arnt1* mutants, *arnt2* mutants, and wild-type embryos all

contain *runx1*-positive cells near the dorsal aorta (**Figure 2E-G**). However, *arnt1/2* mutants have little or

no *runx1* labeling in this region (**Figure 2H**). This suggests that *arnt1* or *arnt2* is necessary for

hematopoietic cell development during embryogenesis, just like *npas4l* (Liao et al., 1997; Liao et al.,

1998; Stainier et al., 1995; Thompson et al., 1998).

Hematopoietic and endothelial progenitor populations are reduced in *arnt1/2* mutants

Hemato-vascular progenitor specification from the multipotent mesoderm begins in mid-gastrula, around shield stage, and peaks during segmentation (Vogeli et al., 2006). Npas4l is thought to directly upregulate target genes that cause cells to differentiate into endothelial cells and hematopoietic stem and progenitor cells. If Npas4l, Arnt1 and Arnt2 act in tandem to regulate target gene expression, then we expect that Npas4l-target genes would be downregulated in *arnt1/2* mutants. Therefore, we hypothesized that Npas4l-target genes would be downregulated in *arnt1/2* mutants. We focused on four putative *npas4l* target genes known to be important for the hemato-vascular lineage: *etsrp (etv2), tal1, lmo2 (scl),* and *egfl7* (Marass et al., 2019; Sumanas and Lin, 2006; Sumanas et al., 2005). We bred *arnt1^{bcm2/+}; arnt2^{bcm3/+}* adults to each other to generate embryos that were *arnt1/2* mutant, *arnt1* mutant, *arnt2* mutant, wild-type, and all other intermediate genotypes (referred to as *arnt1/2* siblings). At the 2-6 somite stages (about 10-14 hpf), *npas4l* mutant embryos show a reduction of expression in *etsrp*, *tal1,* and *lmo2* (Reischauer et al., 2016; Thompson et al., 1998). We examined the expression of these genes in *arnt1/2* mutants and *arnt1/2* siblings between 4-6 somite stage (**Figure 3**). We found that 7% of embryos from *arnt1^{bcm2/+};arnt2^{bcm3/+}* parents showed reduced expression of *etsrp* compared to siblings, (**Figure 3A-B**, *N*=4 biological replicates, 80-250 embryos per clutch, 604 embryos total). This did not differ from the expected Mendelian ratio of *arnt1/2* mutants (6.25%; binomial test, *p*>0.05). To confirm that these embryos were in fact *arnt1/2* mutants, following WISH we genotyped 24 embryos with reduced expression of *etsrp* and found that 100% were *arnt1/2* mutants. We also genotyped 26 embryos with normal expression of *etsrp,* only two of which were *arnt1/2* mutants (**Table S2**). To examine if there is an association between the number of mutant alleles and the observed phenotype, we performed a Spearman Rank correlation test (see Methods for details) and found that there is a statistically significant correlation between the number of mutant alleles and the observed phenotype (*Spearman correlation,* r= -0.9, *p*<0.001). We conclude that the expression of *etsrp* is reduced in *arnt1/2* mutants compared to wild-type.

We found that 7% of embryos from arnt$1^{bcm2/+}$; arnt$2^{bcm3/+}$ parents showed reduced expression of

tal1 compared to arnt1/2 siblings (**Figure 3C-D,** N=6 biological replicates, 60-300 embryos per clutch,

681 embryos total). This did not differ significantly from the expected Mendelian ratio of arnt1/2 mutants

(binomial test, $p>0.05$). To confirm these embryos were in fact arnt1/2 mutants, we genotyped 18

embryos with reduced expression and confirmed that 83% (15 of 18 embryos) of these embryos were

arnt1/2 mutants (**Table S3**). We genotyped 11 embryos with normal expression of tal1, none of which

were arnt1/2 mutants (**Table S3**). To examine if there is a relationship between the number of mutant

alleles and the observed phenotype, we performed a Spearman Rank correlation test and found that

there is a statistically significant correlation between the number of mutant alleles and the observed

phenotype (Spearman correlation, r= -0.6, $p<0.005$). We conclude that the expression of tal1 is reduced

in arnt1/2 mutants compared to wild-type.

We found that 11% of embryos from arnt$1^{bcm2/+}$;arnt$2^{bcm3/+}$ parents showed reduced expression

of lmo2 compared to siblings (**Figure 3D-E,** N=2 biological replicates, 95-175 embryos per clutch, 270

embryos total). This did significantly differ from the expected 6.25% Mendelian ratio of arnt1/2 mutants

(binomial test, $p<0.01$). To test for a phenotype-genotype correlation, we genotyped 185 embryos

following labeling for lmo2: 160 embryos with normal lmo2 expression and 25 embryos with reduced

lmo2 expression. Of the 25 embryos with reduced expression of lmo2, we observed 8 embryos that

were arnt1/2 mutants, 8 embryos with three mutant alleles, 4 embryos with two mutant alleles, 3

embryos with one mutant allele, and 2 wild-type embryos (**Table S4**). Of the 160 embryos with normal

expression of lmo2, 5 embryos were arnt1/2 mutants, 36 embryos had three mutant alleles, 64 embryos

had two mutant alleles, 46 embryos had one mutant allele, and 9 were wild-type embryos (**Table S4**).

To examine if there is a relationship between the number of mutant alleles and the observed

phenotype, we performed a Spearman rank correlation and found that there is a statistically significant

correlation between the number of mutant alleles and the observed phenotype (Spearman correlation, r

= -0.2, $p<0.01$). We conclude that the expression of lmo2 is reduced in arnt1/2 mutants compared to

wild-type.

We examined the expression of *egfl7* at 10-somite stage embryos from *arnt1*^{bcm2/+}*; arnt2*^{bcm3/+}

parents. We observed *egfl7* expression in portions of the ALPM and PLPM in wild-type and in 95% of

the embryos from *arnt1*^{bcm2/+}*; arnt2*^{bcm3/+} parents, but no expression of *egfl7* in the anterior of 5% of

embryos from *arnt1*^{bcm2/+}*; arnt2*^{bcm3/+} parents, N=11 of 214 embryos (**Figure S1,** N=2 biological

replicates, 94-120 embryos per clutch, 214 total embryos, and age-matched wild-type embryos, N=1

biological replicate, 28 embryos). This did not differ from the expected Mendelian ratio of *arnt1/2*

mutants (6.25%; binomial test, *p*>0.05). To examine if there is a relationship between the number of

mutant alleles and the observed phenotype, we performed a Spearman Rank correlation test and found

that there is a statistically significant correlation between the number of mutant alleles and the observed

phenotype (*Spearman correlation,* r= -0.5, *p*<0.001) (**Table S5**). We conclude that the expression of

egfl7 is reduced in *arnt1/2* mutants compared to wild-type.

These results support the idea that *npas4l*, *arnt1* and *arnt2* regulate identical target genes for

hemato-vascular specification. *arnt1* and *arnt2* could interact with *npas4l* at the same level of the

pathway, consistent with bHLH-PAS transcription factors acting as heterodimers. Alternatively, it is

possible that *arnt1* and *arnt2* act upstream of *npas4l*. To distinguish among these ideas, we tested

whether *npas4l* expression was reduced in *arnt1/2* mutants. At the 2-4 somite stage, *arnt1/2* mutants

exhibit normal expression of *npas4l* in the lateral plate mesoderm, indistinguishable from age-matched

wild-type embryos (**Figure S2, Table S6,** N=2 biological replicates, 68-100 embryos per clutch, 168

total embryos). This result indicates that *npas4l* expression occurs independently of *arnt1* and *arnt2*.

We conclude that *npas4l* does not act downstream of *arnt1* or *arnt2*. Taken together, our results

suggest that *arnt1* and *arnt2* act in conjunction with *npas4l* during hemato-vascular specification*.*

arnt1 and arnt2 expression is ubiquitous and colocalizes with npas4l expression

If *arnt1, arnt2* and *npas4l* act together to specify hemato-vascular progenitor cells, then all three

genes should be expressed in the same cells in the lateral plate mesoderm. To determine if *arnt1,*

arnt2, and *npas4l* are expressed in the same cells, we performed whole-mount *in situ* hybridization

chain reaction (WIHCR) on wild-type zebrafish embryos at the 4-somite stage (approximately 10.5 hpf).

At this stage, *npas4l* is robustly expressed in the LPM (Reischauer et al., 2016). We found that *arnt1*

and *arnt2* are expressed ubiquitously throughout the embryo at this stage, although both genes appear

to be expressed at low levels (**Figure 4A and Figure S3A**, N=3 biological replicates, 5-9 embryos per

replicate, 22 embryos total).

Because no previous publications examined the spatial distribution of *arnt1* or *arnt2* mRNA in

embryos younger than 24 hpf (Löhr et al., 2009), we tested whether background or autofluorescence

influenced our results. Samples probed for *arnt1*, *arnt2,* and *npas4l* were compared to negative control

embryos, embryos from the same clutch that underwent the same WIHCR protocol, but in the absence

of anti-sense RNA probe sets (**Figure 4B and Figure S3B**, N=3 biological replicates, 6-12 embryos per

replicate, 28 embryos total)*.* We verified that the observed *npas4l, arnt1,* and *arnt2* expression was

specific using two different image analysis tools. First, we analyzed representative images in 3-

dimensions using the ImageJ plug-in "3D Count Objects" (Bolte and Cordelières, 2006). The number of

objects above a set threshold was measured for each channel and then normalized to background by

subtracting the average number of objects detected in the corresponding channel of the negative

control embryos. *npas4l-, arnt1-*, and *arnt2*-positive objects were all detected in the posterior of the

embryos (**Figure 4C-E**). Second, we analyzed the expression of *arnt1* and *arnt2* in the presumptive

LPM by drawing regions of interest (ROIs) encompassing the observed *npas4l* signal*,* and then

measured the percent area of expression and the mean gray intensity above the threshold for each

channel. The integrated intensity was calculated by multiplying the percent area by the mean grey

intensity, thus taking both brightness and area of signal into consideration (**Figure 4F-H, Figure S3 C-**

E). We found that *npas4l, arnt1,* and *arnt2* all showed statistically significantly more expression in

experimental embryos compared to the negative control, as measured by both number of objects and

integrated intensity of objects in the LPM (**Figure 4F-H,** *npas4l* Welsh's t-test, $p<0.0001$; *arnt2* Welsh's

t-test, $p<0.0001$; *arnt1* Mann-Whitney test, *$p<0.0001$*).

We asked whether the expression of *npas4l*, *arnt1* and *arnt2* was the same in the anterior LPM

(ALPM) and posterior LPM (PLPM). *npas4l* and its associated downstream targets exhibit more

expression in the PLPM compared to the ALPM in embryos throughout segmentation (Liao et al., 1997;

Liao et al., 2000; Marass et al., 2019; Mattonet et al., 2022; Qian et al., 2005; Stainier et al., 1995). At 4-

6 somite stage embryos, we found that expression of *npas4l* and *arnt2* was greater in the PLPM than in

the ALPM, but the expression of *arnt1* was similar in both regions (**Figure S3C, S3E,** *npas4l* Welsh's t-

test, p<0.0001; *arnt2* Welsh's t-test, *p<0.005*). We conclude that *arnt1* and *arnt2* are expressed within

the *npas4l*-positive LPM, consistent with our hypothesis that all 3 genes together regulate the

specification of hemato-vascular progenitor cells.

Expression of *npas4l*-target genes requires *npas4l*, *arnt1* and *arnt2*

npas4l is not expressed until tailbud stage (Reischauer et al., 2016). A previous study

demonstrated that ectopic expression of *npas4l* in shield-stage embryos, prior to the onset of

expression of endogenous *npas4l*, resulted in upregulation of and ectopic expression of early hemato-

vascular genes *etsrp* and *tal1* (Reischauer et al., 2016). We suspect that upregulation of *npas4l* target

genes occurred because *arnt1* and *arnt2* are ubiquitously expressed at shield stage. Single cell RNA

sequencing studies detected *arnt1* and *arnt2* mRNA throughout gastrulation, which encompasses

shield stage, in zebrafish embryos (Baranasic et al., 2022). To test whether expression of *npas4l*-target

genes requires *npas4l*, *arnt1,* and *arnt2*, we injected wild-type embryos at the 1-cell stage with either

npas4l mRNA or *arnt1* and *arnt2* mRNA. The embryos were fixed at shield stage, and assayed for

expression of *tal1* or *lmo2* using WISH. Wild-type embryos injected with *npas4l* mRNA exhibited robust

expression of *tal1* compared to uninjected control embryos (**Figure 5A-B**). In contrast, embryos

injected with *arnt1* and *arnt2* mRNA showed little or no expression of *tal1*, similar to uninjected embryos

(**Figure 5C**). This suggests that neither *arnt1* nor *arnt2* can drive expression of *tal1* in the absence of

npas4l. To directly test whether *arnt1* and *arnt2* can drive expression of *tal1* in the absence of *npas4l*,

we injected *npas4l* mutant embryos with *npas4l* RNA or with *arnt1* and *arnt2* RNA (**Figure 5D-F)**.

Embryos injected with *npas4l* mRNA exhibited robust expression of *tal1* (**Figure 5E**), consistent with

previously published results (Reischauer et al., 2016). However, all the embryos (derived from

npas4l$^{s5/+}$ parents crossed to each other) injected with *arnt1* and *arnt2* mRNA showed little to no

expression of *tal1* (**Figure 5F**). Since 25% of these embryos are *npas4l$^{s5/s5}$*, we conclude that *arnt1* and

arnt2 cannot drive *tal1* expression in the absence of *npas4l*. Finally, to test if *npas4l* upregulates target

genes in the absence of *arnt1* and *arnt2*, we injected *arnt1/2* mutant embryos with *npas4l* mRNA. We

found that *arnt1/2* siblings showed upregulation of *npas4l*-target gene *lmo2*, but *arnt1/2* mutants did not

show upregulation of *lmo2* (**Figure 5G-H**)*. From these results, we conclude that expression of *npas4l*

and either *arnt1* or *arnt2* are required for upregulation of *npas4l*-target genes.

Npas4l interacts with both Arnt1 and Arnt2

Npas4l is a Class I bHLH-PAS domain transcription factor (Reischauer et al., 2016), and as

such, must heterodimerize with a Class II bHLH-PAS transcription factor, like Arnt1 and Arnt2, to

regulate gene transcription. Considering that (1) bHLH-PAS transcription factors function as dimers, (2)

arnt1 and *arnt2* fail to upregulate hemato-vascular genes in the absence of *npas4l*, and (3) *npas4l* fails

to upregulate hemato-vascular genes in *arnt1/2* mutants, therefore we hypothesize that Arnt1 and Arnt2

proteins form a transcription complex with Npas4l protein to regulate gene expression. To determine if

Npas4l interacts with Arnt1 or Arnt2, we performed a coimmunoprecipitation experiment. Due to the

absence of commercially available antibodies that recognize zebrafish Npas4l, Arnt1 or Arnt2 proteins,

we co-expressed tagged versions of zebrafish *npas4l* (Npas4l-V5), *arnt1* (Arnt1-myc), and *arnt2* (Arnt2-

HA) in HEK293T cells. We performed co-immunoprecipitation studies and found that Arnt1-myc

interacts with Npas4l-V5 and that Arnt2-HA interacts with Npas4l-V5 (**Figure 6**, N= 2 biological

replicates per condition). These results suggest that Npas4l uses either Arnt1 or Arnt2 or both as

binding partners. Furthermore, these data indicate that Arnt1 and Arnt2 may be functionally

interchangeable and redundant, at least in the case of Npas4l.

Arnt1 and Arnt2 cannot functionally compensate for one another in the case of Ahr2

Our results demonstrate that Arnt1 and Arnt2 are functionally interchangeable and redundant

when interacting with Npas4l in the context of hemato-vascular development. It is unclear if Arnt1 and

Arnt2 are functionally interchangeable with all Class I bHLH-PAS domain transcription factors or just

with Npas4l. To determine if *arnt1/arnt2* redundancy is unique to Npas4l, we examined if *arnt1* and

arnt2 are interchangeable with a different class I bHLH-PAS transcription factor, the aryl hydrocarbon

receptor 2 (Ahr2). Previous studies demonstrate that Ahr2 can bind DNA in the presence of either Arnt1

or Arnt2 *in vitro* (Andreasen et al., 2002; Hirose et al., 1996; Tanguay et al., 2000), but whether Arnt1

and Arnt2 bind Ahr2 interchangeably *in vivo* is not known. Ahr2 is a ligand-dependent transcription

factor. A known ligand of Ahr2 is 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), which we have previously

shown induces cardiotoxicity in zebrafish larvae and requires *ahr2* (Souder and Gorelick, 2019). We

asked whether TCDD causes cardiotoxicity in *arnt1* or *arnt2* single mutants, reasoning that if Ahr2

interacts with Arnt1 or Arnt2 interchangeably, then only *arnt1/2* double mutants, but not single mutants,

would be resistant to TCDD exposure. We treated clutches of larvae from wild-type parents (N= 3

biological replicates, 10-60 larvae per clutch per drug treatment, 157 larvae total), *arnt1* heterozygous

parents (N= 3-4 biological replicates per allele, 21-36 larvae per clutch per drug treatment, 533 larvae

total, 248 larvae from allele *bcm1* and 285 larvae allele from *bcm2*), or *arnt2* heterozygous parents

(N=3 biological replicates, 40-45 embryos per clutch per drug treatment, 253 larvae total) with either

0.1% DMSO (vehicle) or with 10 ng/mL TCDD from 1 dpf to 3 dpf **(Figure S4)**. Wild-type larvae

exposed to TCDD exhibited pericardial edema and abnormal heart looping, consistent with previous

results (**Figure 7A, D**). *arnt2* mutants treated with TCDD exhibited a phenotype similar to wild-type

(**Figure 7C, F**). In contrast, *arnt1* mutants treated with TCDD showed normal heart looping and no

pericardial edema, similar to vehicle-treated larvae (**Figure 7B, E**). This demonstrates that *arnt1,* but

not *arnt2,* is required for Ahr2-dependent TCDD toxicity. We conclude that *arnt1* and *arnt2* are not

always functionally redundant *in vivo*. Certain Class I bHLH-PAS transcription factors may prefer to

function with a specific Class II bHLH-PAS transcription faction.

DISCUSSION

We find that Arnt1 and Arnt2 are obligate coregulators of Npas4l. Together, *npas4l*, *arnt1* and *arnt2*

genes regulate specification of hemato-vascular cells. *npas4l*, *arnt1* and *arnt2* are each necessary, but

on their own not sufficient, for hemato-vascular specification. Npas4l serves as a master regulator, with

Arnt proteins acting as permissive coregulators. Arnt1 and Arnt2 are functionally redundant in the case

of Npas4l but are not functionally redundant in the case of Ahr2, where there is a preference for Ahr2-

Arnt1 complex.

Class I bHLH-PAS proteins must dimerize with Class II bHLH-PAS proteins to transcribe their target

genes (Card et al., 2005; Ema et al., 1996; Hirose et al., 1996; Jiang et al., 1996; Lees and Whitelaw,

1999; Michaud et al., 2000; Wu and Rastinejad, 2017). It has previously been established that Npas4l, a

Class I bHLH-PAS transcription factor, is required for hemato-vascular development in zebrafish

(Reischauer et al., 2016). To determine the Class II bHLH-PAS binding partner of Npas4l, we

generated *arnt1* mutant and *arnt2* mutant zebrafish and expected one of these mutants to mimic the

npas4l mutant phenotype. However, both *arnt1* mutant and *arnt2* single mutant larvae were viable to 5

dpf and were grossly indistinguishable from wild-type embryos. Only *arnt1/2* double mutant embryos

exhibited a cardiovascular phenotype. Our results demonstrate that double homozygous *arnt1/2* mutant

embryos have a similar phenotype to *npas4l* mutants (Reischauer et al., 2016; Stainier et al., 1995;

Thompson et al., 1998) and that Npas4l can function with either Arnt1 or Arnt2. Thus, Arnt1 and Arnt2

act redundantly to specify hemato-vascular progenitor cells in zebrafish.

Are *arnt1* and *arnt2* functionally redundant with all Class I bHLH-PAS transcription factors? We

argue no. In the case of the class I transcription factor Ahr2, both Arnt1 and Arnt2 bind Ahr2 *in vitro*

(Lanham et al., 2011; Tanguay et al., 2000). However, we find that only Arnt1, and not Arnt2, is required

for Ahr2-dependent cardiotoxicity in zebrafish embryos. This argues that *in vivo*, Ahr2 preferentially

interacts with Arnt1 over Arnt2. Another class I protein, Sim1a, appears to preferentially interact with

Arnt2 *in vivo*. *arnt2* mutant zebrafish embryos have fewer dopaminergic neurons in the brain compared

to wild-type embryos (Löhr et al., 2009). The *arnt2* mutant phenotype is similar to the phenotype when

Sim1a expression is knocked down (Löhr et al., 2009). These results suggest that in the case of the

growth and/or survival of dopaminergic neurons, Sim1a preferentially interacts with Arnt2 versus Arnt1.

To what degree this is due to differences in expression levels, versus an intrinsic structural preference

for a particular protein-protein interaction, requires further study. Such interactions may also depend on

the cell-type and developmental stage where the bHLH-PAS proteins function. Arnt1 and Arnt2 appear

to be expressed ubiquitously during embryonic development, which suggests that such interactions

may be due to factors other than expression levels.

Endothelial cell development and hematopoietic cell development are closely related during

embryogenesis. Both cell types primarily arise from the lateral plate mesoderm (LPM) and share a

common progenitor (Qian et al., 2005; Reischauer et al., 2016; Stainier et al., 1995; Thompson et al.,

1998). Additionally, during definitive hematopoiesis, some HSCs are derived from specialized

endothelial cells, called hemogenic endothelium (Boisset et al., 2010; Kissa and Herbomel, 2010). Both

npas4l and *arnt1/2* mutants lack all blood cells but do have a small population of endothelial cells,

which is restricted to the most posterior region of the animal (Figure 2 and (Reischauer et al., 2016)). In

zebrafish, there is a population of endothelial cells in the caudal region of the embryo that are not

derived from the LPM and differentiate in the absence of *npas4l* expression (Pak et al., 2020). We

hypothesize that *arnt1* and *arnt2* are required for the differentiation of endothelial cells from the LPM

but are not required for the differentiation of endothelial cells, such as the caudal population, from

outside the LPM. The transcriptomic signature of the caudal population of endothelial cells was distinct

from endothelial cells that require *npas4l* for differentiation, with genes for somitogenesis and

neurogenesis highly enriched compared to LPM-derived endothelial cells (Pak et al., 2020). Based on

this transcriptome analysis, we speculate that the caudal population of endothelial cells fails to express

genes required for hemogenic endothelium and that no blood cells are derived from the caudal

population of endothelial cells. This is consistent with the observation that the caudal population of

endothelial cells, but not blood cells, are present in *npas4l* or *arnt1/2* mutants.

Arnt proteins interact with multiple class I bHLH-PAS domain proteins to regulate cardiovascular

development. Our results focused on the role of Arnt proteins in hemato-vascular specification,

however Arnt proteins are likely involved in later steps of endothelial cell development. Differentiation of

hemogenic endothelium in zebrafish and in mice is dependent upon several Class I bHLH-PAS

transcription factors: Hypoxia inducible factor 1 alpha (mouse *Hif1α*, zebrafish genes *hif1aa*, *hif1ab)*

and hypoxia inducible factor 2 alpha (mouse *Hif2α,* zebrafish genes *epas1a*, *epas1b*) (Gerri et al., 2018;

Ramírez-Bergeron et al., 2004). Given that these HIF proteins are Class I bHLH-PAS transcription

factors, it is likely that one or both Arnt proteins are required for HIF-dependent differentiation of

hemogenic endothelium.

The function of Arnt proteins in zebrafish cardiovascular development is likely conserved in mice.

Arnt1 is essential for the viability of adult hematopoietic stem cells in mice and is required for the

survival of hematopoietic progenitor cells in the fetal mouse liver (Krock et al., 2015). Homozygous

Arnt1 mutant mice are not viable beyond E10.5 and show severe defects in vascularization, suggesting

that Arnt1 is essential for normal vascular development (Kozak et al., 1997; Maltepe et al., 1997). The

degree to which Arnt2 contributes to hemato-vascular development in mice is not well-understood.

Arnt2 has primarily been studied within the context of neural development (Hao et al., 2013; Hosoya et

al., 2001; Jain et al., 1998), however, there is some evidence to suggest that *Arnt1* and *Arnt2* may

exhibit functional redundancy during development of non-neural tissues (Keith et al., 2001; Sekine et

al., 2006). We observed lower than expected numbers of $arnt1^{bcm2/bcm2}$; $arnt2^{bcm3/+}$ adult zebrafish

derived from $arnt1^{bcm2/+}$; $arnt2^{bcm3/+}$ parents. This suggests that *arnt1* and *arnt2* have a dose-dependent

effect on the survival or fitness of the animal, where anything less than four wild-type alleles (two *arnt1*

alleles, two *arnt2* alleles) is deleterious. A similar dose-dependent effect on survival can be seen in

mouse embryos from $Arnt1^{+/-}$; $Arnt2^{+/-}$ parents (Keith et al., 2001). The mechanisms behind these dose-

dependent gene effects are not known.

While zebrafish *arnt1* and *arnt2* genes are conserved in mammals, there is no known equivalent of

zebrafish *npas4l* in mammalian genomes. Zebrafish Npas4l shows the highest sequence homology to

mouse Npas4 (Reischauer et al., 2016). Although Npas4 is known to dimerize with Arnt1 and Arnt2 *in*

vivo (Brigidi et al., 2019), mammalian Npas4 is unlikely to be the functional equivalent to zebrafish

Npas4l because homozygous Npas4 mutant mice are viable to adulthood and do not show any overt

vascular or hematopoietic defects (Bloodgood et al., 2013). Given our results that *npas4l* requires *arnt1*

or *arnt2* to act as co-regulators of hemato-vascular specification, and the abundance of conserved

genes that regulate vascular development in zebrafish and humans (Hogan and Schulte-Merker, 2017),

we speculate that the functional mammalian equivalent of *npas4l* is a Class I bHLH-PAS transcription

factor, or multiple such factors, that function together with Arnt1 or Arnt2.

Acknowledgements

We thank Lauren Pandolfo and the staff of the Baylor College of Medicine aquatics facility for taking
care of our zebrafish colony, Rosa Uribe and Adam Howard from Rice University for help with WIHCR,
and Paula Pimienta-Ramirez and Yunping Lei in the CPEH DNA sequencing core at BCM for help with
sequencing embryos following in situ hybridization. This work was supported by NIH grants
R01ES026337 (to DAG), P30ES030285 (pilot award to DAG), and T32DK060445 (to MJE).
625

627
628

Genotype	Expected Percentage	Expected n=356	Actual Percentage	Actual n=356
$arnt1^{+/+}; arnt2^{+/+}$	6.3%	22.25	6.2%	22
$arnt1^{+/+}; arnt2^{bcm3/+}$	12.5%	44.5	14.3%	51
$arnt1^{+/+}; arnt2^{bcm3/bcm3}$	6.3%	22.25	4.5%	16
$arnt1^{bcm2/+}; arnt2^{+/+}$	12.5%	44.5	11.8%	42
$arnt1^{bcm2/+}; arnt2^{bcm3/+}$	25%	89	25.3%	90
$arnt1^{bcm2/+}; arnt2^{bcm3/bcm3}$	12.5%	44.5	11.8%	42
$arnt1^{bcm2/bcm2}; arnt2^{+/+}$	6.3%	22.25	5.3%	19
$arnt1^{bcm2/bcm2}; arnt2^{bcm3/+}$	12.5%	44.5	13.5%	48
$arnt1^{bcm2/bcm2}; arnt2^{bcm3/bcm3}$	6.3%	22.25	7.3%	26

Table 1. Genotypes of embryos from $arnt1^{bcm2/+}; arnt2^{bcm3/+}$ parents follow expected Mendelian ratios. The frequencies of the observed genotypes were not significantly different from the expected genotypes, X^2 (8, N=356) = 4.38, $p>0.05$.

Genotype	Expected Percentage	Expected n=138	Actual Percentage	Actual n=138
arnt1^{+/+}; *arnt2*^{+/+}	7.69%	10.6	7.24%	10
arnt1^{+/+}; *arnt2*^{bcm3/+}	15.4%	21.25	15.2%	21
arnt1^{+/+}; *arnt2*^{bcm3/bcm3}	0%	0	0%	0
arnt1^{bcm2/+}; *arnt2*^{+/+}	13.3%	18.4	30.4%	42
arnt1^{bcm2/+}; *arnt2*^{bcm3/+}	30.8%	42	43.5%	60
arnt1^{bcm2/+}; *arnt2*^{bcm3/bcm3}	0%	0	0%	0
arnt1^{bcm2/bcm2}; *arnt2*^{+/+}	7.69%	10.6	1.45%	2
arnt1^{bcm2/bcm2}; *arnt2*^{bcm3/+}	15.4%	21.25	2.17%	3
arnt1^{bcm2/bcm2}; *arnt2*^{bcm3/bcm3}	0%	0	0%	0

632
633

Table 2. Genotype frequencies of viable adult fish from *arnt1*$^{bcm2/+}$; *arnt2*$^{bcm3/+}$ parents. Offspring were raised to adulthood (≥3 months old) and genotyped. The expected Mendelian ratios were adjusted to reflect the fact that *arnt2*$^{bcm3/bcm3}$ mutants are not viable to adulthood. The frequencies of the observed genotypes were significantly different from the expected genotypes, with fewer *arnt1*$^{bcm2/bcm2}$; *arnt2*$^{bcm3/+}$ observed. X^2 (5, N=138) = 45.57, p<0.0001.

Figure 1. *arnt1/2* double mutant embryos, but not *arnt1* or *arnt2 single* mutants, lack circulating blood, have cardiac edema and atrial enlargement. (A) bHLH-PAS transcription factors act as heterodimers: a Class I protein and Class II protein interact and bind DNA. Asterisks indicate proteins encoded by genes present in zebrafish but absent in humans. (**B,C**) Generation of zebrafish with mutations in *arnt1* and *arnt2* genes. The amino acid sequences for the zebrafish Arnt1 (**B**), and Arnt2 proteins (**C**) are represented by the black line. Basic helix-loop-helix (bHLH), Per-Arnt-Sim (PAS) domains, and the motif C-terminally located to the PAS (PAC) are represented by the labeled bubbles. The numbers below each domain indicate the starting and ending amino acid (AA) for each motif. (**B**) The *arnt1^{bcm1}* mutation causes a premature stop codon to occur at AA139 and loss of the PAS-B, PAC, and majority of the PAS-A. The *arnt1^{bcm2}* mutation causes a premature stop codon to occur at AA250 and loss of the PAS-B and PAC domains. (**C**) The *arnt2^{bcm3}* mutation causes a premature stop codon to occur at AA287 and loss of the PAS-B and PAC domains. (**D-I**) Live images of embryos from the indicated genotype at 2 days post fertilization (dpf). All double homozygous *arnt1^{bcm2/bcm2}; arnt2^{bcm3/bcm3}* embryos showed cardiac edema and atrial enlargement (**G** and **G'**). The wildtype (**D** and **D'**), *arnt1^{bcm2/bcm2}* (**E** and **E'**), and *arnt2^{bcm3/bcm3}* (**F** and **F'**) siblings of the double homozygous mutants (*arnt1^{bcm2/bcm2}; arnt2^{bcm3/bcm3}*) showed normal heart development. Black arrows in panels D'-G' indicate the atrium and ventricle of the heart. Images are representative of 7 different clutches, each clutch contained 20-300 embryos. All siblings of double homozygous mutants (*arnt1^{bcm2/bcm2}; arnt2^{bcm3/bcm3}*) had circulating blood (**H** and **Movie 1, Movie 2, Movie 5, Movie 6**) while *arnt1/2* mutants lacked visible circulating blood cells (**I** and **Movie 3, Movie 4, Movie 7, Movie 8**). The black arrows in panel **H** indicate blood cells. The arrowheads in **H** and **I** indicate non-blood cells found in all embryos. (**J-L**), o-dianisidine staining to label blood cells in embyos 3 days post fertilization shows that *arnt1^{bcm2/bcm2}; arnt2^{bcm3/bcm3}* double homozygous mutants lack blood cells, while all other genotypes have clearly labeled blood cells. The arrow in **J** and **K** points to the duct of Cuvier, and the arrowheads in **K** show circulating blood cells. All embryos were derived from *arnt1^{bcm2/+};arnt2^{bcm3/+}* parents, images are representative of 1 clutch, containing 98 embryos. (**J**) N=96 of 98 embryos showed clear o-dianisidine staining and none of these embryos were *arnt1/2* mutants. (**K**) N=2 of 98 embryos showed a lack of o-dianisidine staining and both embryos were confirmed *arnt1/2* mutants. For all images anterior is towards the left and dorsal is towards the top, except for panel **J** which shows a ventral view of the embryos. Scale bars: D-G, 1 mm; D'-G' and H-K, 100 μm.

Figure 2. *arnt1/2* mutants show reduced levels of endothelial cells and *runx1*-positive hematopoietic cells. (A-L) Embryos were probed for *kdrl* (A-D) or *runx1* (E-H) using whole-mount in situ hybridization at 2 days post fertilization (dpf). (A-D) *kdrl* staining in wild-type, *arnt1* mutant, *arnt2*

mutant, and *arnt1/2* mutant embryos. (A'-D') Images of each embryo's trunk near the end of the yolk

extension (indicated by the box in panels A-D). Black arrowheads indicate *kdrl* staining in the dorsal

longitudinal anastomotic vessels, black arrows indicate the dorsal aorta and caudal vein, and white

arrowheads indicate inter-segmental vessels. (E-H) Black arrows indicate *runx1* labeling in the

presumptive hematopoietic stem cell niche. (I-L) Wild-type and mutant embryos on Tg(*fli1:EGFP*[y1])

transgenic background were imaged at 1 dpf and then genotyped. (I'-L') Enlarged images of each

embryo's trunk near the end of the yolk extension (indicated by the box in panels I-L). White

arrowheads correspond to inter-segmental vessels. e: eye, pa: pharyngeal arch, da: dorsal aorta, cv:

caudal vein. The fraction in the bottom right corner indicates the number of embryos of the given

genotype with the represented phenotype over the total number of embryos of the given genotype from

a single clutch. Scale bars,100 μm.

Figure 3. *arnt1/2* mutants show decreased *etsrp*, *tal1*, and *lmo2* expression in the lateral plate

mesoderm at 4-6 somite stages. Whole-mount in situ hybridization for *etsrp*, *tal1,* and *lmo2*

expression was performed on 4-6 somite stage embryos that were offspring of *arnt1*[bcm2/+]*;arnt2*[bcm3/+]

adult zebrafish. (**A**) Expression of *etsrp* was observed in the anterior and posterior lateral plate

mesoderm (ALPM, PLPM) in 93% of embryos. (**B**) Limited or no expression of *etsrp* was observed in

7% of embryos, consistent with the expected Mendelian ratio of 6.25% for *arnt1*[bcm2/+]*;arnt2*[bcm3/+] (*arnt1/2*

mutants). (**C**) Expression of *tal1* was observed in the ALPM and PLPM in 93% of embryos. (**D**) Limited

or no expression of *tal1* was observed in 7% of embryos, consistent with the expected Mendelian ratio

of 6.25% for *arnt1/2* mutants. (**E**) Expression of *lmo2* was observed in 89% of embryos. (**F**) Limited or

no expression of *lmo2* was observed in 11% of embryos, which is slightly more than the expected

Mendelian ratio of 6.25% for *arnt1/2* mutants. We genotyped a subset of each group of embryos (see

Supplemental Tables 2-4) and observed a correlation between number of mutant alleles and

expression of *etsrp*, *tal1*, and *lmo2* genes. Spearman correlation, *etsrp* r= -0.9, p<0.001; *tal1* r= -0.6,

p<0.005; *lmo2* r= -0.2, p<0.01. All embryos are oriented with dorsal towards the top. Fractions in

bottom right corners refer to the number of embryos with the indicated phenotype over the total number

of embryos examined. Scale bars, 100 μm

Figure 4. *arnt1* and *arnt2* are ubiquitously expressed at the 4-somite stage and co-localized with

***npas4l* expression.** Whole-mount *in situ* hybridization chain reaction was performed on wild-type (AB)

embryos at the 4 somite stage. (**A**) Embryos were probed for *npas4l, arnt1,* and *arnt2* expression and

counter-stained with DAPI. *arnt1* and *arnt2* labeling was detected ubiquitously, while *npas4l* labeling

was localized to the lateral plate mesoderm. (A') Higher magnification views of the boxed areas in (A)

shows co-localization of *arnt1, arnt2,* and *npas4l* in the posterior lateral plate mesoderm, outlined in

cyan. (**B**) To demonstrate that the observed expression of *arnt1* and *arnt2* is not autofluorescence of

the embryo or the yolk, wild-type embryos were processed identically to those in (A), but no anti-sense

RNA probe sets were added. (B') Higher magnification views of the boxed areas in (B) shows what

background/autofluorescence looks like in the presumptive posterior lateral plate mesoderm. All images

in (**A, B**) are maximum intensity projections from z-stacks taken every 2.5 μm. In merged panels,

npas4l is green, *arnt1* is magenta, *arnt2* is cyan. The posterior portion of the embryo is in view, dorsal is

towards the top of the image. Scale bars, 100 μm. (**C-E**) To validate the observed expression of *arnt1*

and *arnt2,* we quantified the number of detectable objects above a set threshold in the whole embryo

across each z-plane for 4 negative control embryos and 6 experimental embryos probed for *arnt1*, *arnt2*

and *npas4l*. We then normalized the number of detected objects by subtracting the average number of

objects detected in the negative controls from the number of detected objects in the experimental

embryos. Each point on these graphs represents the total number of objects above threshold for a

single embryo. (**F-H**) To demonstrate that *arnt1* and *arnt2* expression occurs in the lateral plate

mesoderm (LPM), we calculated the integrated intensity of fluorescence (pixel area, pixel intensity) over

each side of the posterior LPM in maximum intensity projections from control and experimental

embryos. Each point represents the integrated intensity from one side of the posterior LPM from a

single embryo (right or left side), where each embryo has two bilateral *npas4l*-positive regions of the

posterior LPM. Points of the same color are measurements from opposite sides of the posterior LPM

from the same embryo. *npas4l*, *arnt1*, and *arnt2* labeling was statistically significantly increased in the

lateral plate mesoderm compared to negative control embryos. *npas4l* Welch's t-test, t(11)=8.6,

p<0.0001; *arnt1* Mann-Whitney, U=0, p<0.0001; *arnt2* Welch's t-test, t(11)=7.0, p<0.0001

Figure 5. Expression of *npas4l*-target genes requires *npas4l, arnt1* and *arnt2*. Zebrafish embryos,

derived from the indicted parental genotypes, were injected at the one-cell stage with 50 pg of *npas4l*

mRNA or with 50 pg each of *arnt1* and *arnt2* mRNA (100 pg total RNA). Embryos were fixed at shield

stage and probed for *tal1* or *lmo2, npas4l*-target genes, using whole mount in situ hybridization.

Embryos were then genotyped. (A-C) In wild-type embryos, *tal1* is expressed following injection of

npas4l RNA. *tal1* mRNA was not detected in uninjected embryos or following injection of *arnt1* and

arnt2 RNA. (D-F) Heterozygous *npas4l* mutant fish were crossed to each other and the resulting

embryos were injected with the indicated RNA. *npas4l* RNA, but not *arnt1* and *arnt2* RNA, upregulated

expression of *tal1*. Genotype of embryos in (E) n = 11 *npas4l*$^{+/+}$, n = 15 *npas4l*$^{s5/+}$, n = 10 *npas4l*$^{s5/s5}$, (F)

n = 12 *npas4l*$^{+/+}$, n = 20 *npas4l*$^{s5/+}$, n = 8 *npas4l*$^{s5/s5}$. (G-I) *arnt1*$^{bcm2/+}$*;arnt2*$^{bcm3/+}$ adults were crossed to

each other and the resulting embryos were injected with *npas4l* RNA. Double homozygous embryos

failed to upregulate *lmo2* (panel I) whereas sibling embryos upregulated *lmo2* (H). Genotype of

embryos in (H) n = 1 *arnt1$^{+/+}$;arnt2$^{+/+}$*, n = 7 *arnt1$^{bcm2/+}$;arnt2$^{bcm3/+}$*, n = 1 *arnt1$^{+/+}$;arnt2$^{bcm3/+}$*, n = 8

arnt1$^{bcm2/+}$;arnt2$^{+/+}$, n= 4 *arnt1$^{+/+}$;arnt2$^{bcm3/bcm3}$*, n= 7 *arnt1$^{bcm2/bcm2}$;arnt2$^{+/+}$*, n = 7 *arnt1$^{bcm2/+}$;arnt2$^{bcm3/bcm3}$*,

n = 4 *arnt1$^{bcm2/bcm2}$;arnt2$^{bcm3/+}$*, (I) n = 3 *arnt1$^{bcm2/bcm2}$;arnt2$^{bcm3/bcm3}$*. Embryos are oriented in a lateral view

with the animal pole towards the top. Fractions in the bottom right corner indicate the number of

embryos with the represented phenotype over the total number of embryos assayed. Scale bar in (A)

applies to all panels, 100 μm.

Figure 6. Arnt1 and Arnt2 co-immunoprecipitate with Npas4l. (A) HEK293T cells were transfected

with zebrafish Npas4l-V5 and Arnt1-myc. Cells were homogenized (input) and immunoprecipitated (IP)

with antibodies against V5 or an immunoglobulin G control (IgG). Samples were run on an immunoblot

and probed using myc antibodies (anti-myc) to detect the presence of Arnt1-myc (top) or samples were

probed using V5 antibodies (anti-V5) to verify successful pulldown of Npas4l-V5 (bottom). **(B)** HEK293T

cells were transfected with zebrafish Npas4l-V5 and Arnt2-HA. Cells were homogenized and

immunoprecipitated with antibodies against HA or an IgG control. Samples were run on an immunoblot

and probed using V5 antibodies (anti-V5) to detect the presence of Npas4l-V5 (top) or samples were

probed using HA antibodies (anti-HA) to verify successful pulldown of Arnt2-HA (bottom).

Figure 7. Mutation in *arnt1,* but not *arnt2,* prevents AHR2-dependent TCDD toxicity. Wild-type or

mutant larvae were exposed to (**A, B, C**) 0.1% DMSO, or (**D, E, F**) 2,3,7,8-tetrachlorodibenzodioxin

(TCDD) from 1-3 dpf and imaged at 3 dpf. (**A, D**) Wildtype larvae exposed to TCDD lack an S-shaped

cardiac loop and exhibit cardiac edema, unlike vehicle treated wild-type siblings. (**B,E**) *arnt1* mutant

larvae (*arnt1$^{bcm2/bcm2}$*) treated with vehicle and TCDD show normal heart looping and do not have

cardiac edema. (**C,F**) *arnt2* mutant larvae (*arnt2$^{bcm3/bcm3}$*) treated with vehicle show expected heart

looping similar to wild type siblings, but *arnt2* homozygous mutant larvae treated with TCDD lack an S-

shaped cardiac loop similar to wild type and exhibit cardiac edema similar to TCDD treated wild-type

larvae. Black arrows indicate location of the heart. Larvae were genotyped following imaging and

representative images are shown. TCDD treated wild-type siblings of *arnt1* mutant larvae, n= 9 larvae

across 3 clutches; TCDD treated *arnt1* mutant larvae, n=21 larvae across 3 clutches. TCDD treated

wild-type siblings of *arnt2* larvae, n=19 larvae across 3 clutches; TCDD treated *arnt2* mutant larvae,

n=22 larvae across 4 clutches. All larvae are oriented with anterior to the left and dorsal towards the

top. Scale bar, 100 μm.

770
771
772

Andreasen, E. A., Spitsbergen, J. M., Tanguay, R. L., Stegeman, J. J., Heideman, W. and Peterson, R. E. (2002). Tissue-specific expression of AHR2, ARNT2, and CYP1A in zebrafish embryos and larvae: effects of developmental stage and 2,3,7,8-tetrachlorodibenzo-p-dioxin exposure. *Toxicol. Sci.* **68**, 403–419.

Baranasic, D., Hörtenhuber, M., Balwierz, P. J., Zehnder, T., Mukarram, A. K., Nepal, C., Várnai, C., Hadzhiev, Y., Jimenez-Gonzalez, A., Li, N., et al. (2022). Multiomic atlas with functional stratification and developmental dynamics of zebrafish cis-regulatory elements. *Nat. Genet.* **54**, 1037–1050.

Begley, C. G., Aplan, P. D., Denning, S. M., Haynes, B. F., Waldmann, T. A. and Kirsch, I. R. (1989). The gene SCL is expressed during early hematopoiesis and encodes a differentiation-related DNA-binding motif. *Proc Natl Acad Sci USA* **86**, 10128–10132.

Bloodgood, B. L., Sharma, N., Browne, H. A., Trepman, A. Z. and Greenberg, M. E. (2013). The activity-dependent transcription factor NPAS4 regulates domain-specific inhibition. *Nature* **503**, 121–125.

Boisset, J.-C., van Cappellen, W., Andrieu-Soler, C., Galjart, N., Dzierzak, E. and Robin, C. (2010). In vivo imaging of haematopoietic cells emerging from the mouse aortic endothelium. *Nature* **464**, 116–120.

Bolte, S. and Cordelières, F. P. (2006). A guided tour into subcellular colocalization analysis in light microscopy. *J. Microsc.* **224**, 213–232.

Brigidi, G. S., Hayes, M. G. B., Delos Santos, N. P., Hartzell, A. L., Texari, L., Lin, P.-A., Bartlett, A., Ecker, J. R., Benner, C., Heinz, S., et al. (2019). Genomic Decoding of Neuronal Depolarization by Stimulus-Specific NPAS4 Heterodimers. *Cell* **179**, 373-391.e27.

Card, P. B., Erbel, P. J. A. and Gardner, K. H. (2005). Structural basis of ARNT PAS-B dimerization: use of a common beta-sheet interface for hetero- and homodimerization. *J. Mol. Biol.* **353**, 664–677.

Casie Chetty, S., Rost, M. S., Enriquez, J. R., Schumacher, J. A., Baltrunaite, K., Rossi, A., Stainier, D. Y. R. and Sumanas, S. (2017). Vegf signaling promotes vascular endothelial differentiation by modulating etv2 expression. *Dev. Biol.* **424**, 147–161.

Dooley, K. A., Davidson, A. J. and Zon, L. I. (2005). Zebrafish scl functions independently in hematopoietic and endothelial development. *Dev. Biol.* **277**, 522–536.

Dumont, D. J., Fong, G. H., Puri, M. C., Gradwohl, G., Alitalo, K. and Breitman, M. L. (1995). Vascularization of the mouse embryo: a study of flk-1, tek, tie, and vascular endothelial growth factor expression during development. *Dev. Dyn.* **203**, 80–92.

Edwards, H. E. and Gorelick, D. A. (2022). The evolution and structure/function of bHLH-PAS transcription factor family. *Biochem. Soc. Trans.* **50**, 1227–1243.

Ema, M., Morita, M., Ikawa, S., Tanaka, M., Matsuda, Y., Gotoh, O., Saijoh, Y., Fujii, H., Hamada, H., Kikuchi, Y., et al. (1996). Two new members of the murine Sim gene family are transcriptional

repressors and show different expression patterns during mouse embryogenesis. *Mol. Cell. Biol.* **16**,
5865–5875.

Ferrara, N., Carver-Moore, K., Chen, H., Dowd, M., Lu, L., O'Shea, K. S., Powell-Braxton, L.,
Hillan, K. J. and Moore, M. W. (1996). Heterozygous embryonic lethality induced by targeted
inactivation of the VEGF gene. *Nature* **380**, 439–442.

Gerri, C., Marass, M., Rossi, A. and Stainier, D. Y. R. (2018). Hif-1α and Hif-2α regulate hemogenic
endothelium and hematopoietic stem cell formation in zebrafish. *Blood* **131**, 963–973.

Hao, N., Bhakti, V. L. D., Peet, D. J. and Whitelaw, M. L. (2013). Reciprocal regulation of the basic
helix-loop-helix/Per-Arnt-Sim partner proteins, Arnt and Arnt2, during neuronal differentiation. *Nucleic*
Acids Res. **41**, 5626–5638.

Herbert, S. P. and Stainier, D. Y. R. (2011). Molecular control of endothelial cell behaviour during
blood vessel morphogenesis. *Nat. Rev. Mol. Cell Biol.* **12**, 551–564.

Hill, A. J., Heiden, T. C. K., Heideman, W. and Peterson, R. E. (2009). Potential roles of Arnt2 in
zebrafish larval development. *Zebrafish* **6**, 79–91.

Hirose, K., Morita, M., Ema, M., Mimura, J., Hamada, H., Fujii, H., Saijo, Y., Gotoh, O., Sogawa, K.
and Fujii-Kuriyama, Y. (1996). cDNA cloning and tissue-specific expression of a novel basic helix-
loop-helix/PAS factor (Arnt2) with close sequence similarity to the aryl hydrocarbon receptor nuclear
translocator (Arnt). *Mol. Cell. Biol.* **16**, 1706–1713.

Hirschi, K. K. (2012). Hemogenic endothelium during development and beyond. *Blood* **119**, 4823–
4827.

Hogan, B. M. and Schulte-Merker, S. (2017). How to plumb a pisces: understanding vascular
development and disease using zebrafish embryos. *Dev. Cell* **42**, 567–583.

Hosoya, T., Oda, Y., Takahashi, S., Morita, M., Kawauchi, S., Ema, M., Yamamoto, M. and Fujii-
Kuriyama, Y. (2001). Defective development of secretory neurones in the hypothalamus of Arnt2-
knockout mice. *Genes Cells* **6**, 361–374.

Huber, T. L., Kouskoff, V., Fehling, H. J., Palis, J. and Keller, G. (2004). Haemangioblast
commitment is initiated in the primitive streak of the mouse embryo. *Nature* **432**, 625–630.

Jain, S., Maltepe, E., Lu, M. M., Simon, C. and Bradfield, C. A. (1998). Expression of ARNT, ARNT2,
HIF1?, HIF2? and Ah receptor mRNAs in the developing mouse. *Mech. Dev.* **73**, 117–123.

Jiang, B. H., Rue, E., Wang, G. L., Roe, R. and Semenza, G. L. (1996). Dimerization, DNA binding,
and transactivation properties of hypoxia-inducible factor 1. *J. Biol. Chem.* **271**, 17771–17778.

Kafina, M. D. and Paw, B. H. (2018). Using the zebrafish as an approach to examine the mechanisms
of vertebrate erythropoiesis. *Methods Mol. Biol.* **1698**, 11–36.

Kalev-Zylinska, M. L., Horsfield, J. A., Flores, M. V. C., Postlethwait, J. H., Vitas, M. R., Baas, A.
M., Crosier, P. S. and Crosier, K. E. (2002). Runx1 is required for zebrafish blood and vessel
development and expression of a human RUNX1-CBF2T1 transgene advances a model for studies
of leukemogenesis. *Development* **129**, 2015–2030.

Kataoka, H., Hayashi, M., Nakagawa, R., Tanaka, Y., Izumi, N., Nishikawa, S., Jakt, M. L., Tarui, H.
and Nishikawa, S.-I. (2011). Etv2/ER71 induces vascular mesoderm from Flk1+PDGFRα+ primitive
mesoderm. *Blood* **118**, 6975–6986.

Keith, B., Adelman, D. M. and Simon, M. C. (2001). Targeted mutation of the murine arylhydrocarbon
receptor nuclear translocator 2 (Arnt2) gene reveals partial redundancy with Arnt. *Proc Natl Acad Sci*
USA **98**, 6692–6697.

Kissa, K. and Herbomel, P. (2010). Blood stem cells emerge from aortic endothelium by a novel type
of cell transition. *Nature* **464**, 112–115.

Kobayashi, I., Kobayashi-Sun, J., Kim, A. D., Pouget, C., Fujita, N., Suda, T. and Traver, D. (2014).
Jam1a-Jam2a interactions regulate haematopoietic stem cell fate through Notch signalling. *Nature*
512, 319–323.

Kozak, K. R., Abbott, B. and Hankinson, O. (1997). ARNT-deficient mice and placental
differentiation. *Dev. Biol.* **191**, 297–305.

Krock, B. L., Eisinger-Mathason, T. S., Giannoukos, D. N., Shay, J. E., Gohil, M., Lee, D. S.,
Nakazawa, M. S., Sesen, J., Skuli, N. and Simon, M. C. (2015). The aryl hydrocarbon receptor
nuclear translocator is an essential regulator of murine hematopoietic stem cell viability. *Blood* **125**,
3263–3272.

Lam, E. Y. N., Chau, J. Y. M., Kalev-Zylinska, M. L., Fountaine, T. M., Mead, R. S., Hall, C. J.,
Crosier, P. S., Crosier, K. E. and Flores, M. V. (2009). Zebrafish runx1 promoter-EGFP transgenics
mark discrete sites of definitive blood progenitors. *Blood* **113**, 1241–1249.

Lanham, K. A., Prasch, A. L., Weina, K. M., Peterson, R. E. and Heideman, W. (2011). A dominant
negative zebrafish Ahr2 partially protects developing zebrafish from dioxin toxicity. *PLoS ONE* **6**,
e28020.

Lawson, N. D. and Weinstein, B. M. (2002). In vivo imaging of embryonic vascular development using
transgenic zebrafish. *Dev. Biol.* **248**, 307–318.

Lees, M. J. and Whitelaw, M. L. (1999). Multiple roles of ligand in transforming the dioxin receptor to
an active basic helix-loop-helix/PAS transcription factor complex with the nuclear protein Arnt. *Mol.*
Cell. Biol. **19**, 5811–5822.

Lee, R. K., Stainier, D. Y., Weinstein, B. M. and Fishman, M. C. (1994). Cardiovascular development
in the zebrafish. II. Endocardial progenitors are sequestered within the heart field. *Development* **120**,
3361–3366.

Liao, W., Bisgrove, B. W., Sawyer, H., Hug, B., Bell, B., Peters, K., Grunwald, D. J. and Stainier,
D. Y. (1997). The zebrafish gene cloche acts upstream of a flk-1 homologue to regulate endothelial
cell differentiation. *Development* **124**, 381–389.

Liao, E. C., Paw, B. H., Oates, A. C., Pratt, S. J., Postlethwait, J. H. and Zon, L. I. (1998). SCL/Tal-1
transcription factor acts downstream of cloche to specify hematopoietic and vascular progenitors in
zebrafish. *Genes Dev.* **12**, 621–626.

Liao, W., Ho, C. Y., Yan, Y. L., Postlethwait, J. and Stainier, D. Y. (2000). Hhex and scl function in
parallel to regulate early endothelial and blood differentiation in zebrafish. *Development* **127**, 4303–
4313.

Löhr, H., Ryu, S. and Driever, W. (2009). Zebrafish diencephalic A11-related dopaminergic neurons
share a conserved transcriptional network with neuroendocrine cell lineages. *Development* **136**,
1007–1017.

Maltepe, E., Schmidt, J. V., Baunoch, D., Bradfield, C. A. and Simon, M. C. (1997). Abnormal
angiogenesis and responses to glucose and oxygen deprivation in mice lacking the protein ARNT.
Nature **386**, 403–407.

Marass, M., Beisaw, A., Gerri, C., Luzzani, F., Fukuda, N., Günther, S., Kuenne, C., Reischauer, S.
and Stainier, D. Y. R. (2019). Genome-wide strategies reveal target genes of Npas4l associated with
vascular development in zebrafish. *Development* **146**,.

Marchi, D., Santhakumar, K., Markham, E., Li, N., Storbeck, K.-H., Krone, N., Cunliffe, V. T. and
van Eeden, F. J. M. (2020). Bidirectional crosstalk between Hypoxia-Inducible Factor and
glucocorticoid signalling in zebrafish larvae. *PLoS Genet.* **16**, e1008757.

Mattonet, K., Riemslagh, F. W., Guenther, S., Prummel, K. D., Kesavan, G., Hans, S.,
Ebersberger, I., Brand, M., Burger, A., Reischauer, S., et al. (2022). Endothelial versus
pronephron fate decision is modulated by the transcription factors Cloche/Npas4l, Tal1, and Lmo2.
Sci. Adv. **8**, eabn2082.

Michaud, J. L., DeRossi, C., May, N. R., Holdener, B. C. and Fan, C. M. (2000). ARNT2 acts as the
dimerization partner of SIM1 for the development of the hypothalamus. *Mech. Dev.* **90**, 253–261.

North, T. E., Goessling, W., Peeters, M., Li, P., Ceol, C., Lord, A. M., Weber, G. J., Harris, J.,
Cutting, C. C., Huang, P., et al. (2009). Hematopoietic stem cell development is dependent on blood
flow. *Cell* **137**, 736–748.

Okuda, T., van Deursen, J., Hiebert, S. W., Grosveld, G. and Downing, J. R. (1996). AML1, the
target of multiple chromosomal translocations in human leukemia, is essential for normal fetal liver
hematopoiesis. *Cell* **84**, 321–330.

Pak, B., Schmitt, C. E., Choi, W., Kim, J.-D., Han, O., Alsiö, J., Jung, D.-W., Williams, D. R.,
Coppieters, W., Stainier, D. Y. R., et al. (2020). Analyses of Avascular Mutants Reveal Unique
Transcriptomic Signature of Non-conventional Endothelial Cells. *Front. Cell Dev. Biol.* **8**, 589717.

Palis, J., Robertson, S., Kennedy, M., Wall, C. and Keller, G. (1999). Development of erythroid and
myeloid progenitors in the yolk sac and embryo proper of the mouse. *Development* **126**, 5073–5084.

Pongratz, I., Antonsson, C., Whitelaw, M. L. and Poellinger, L. (1998). Role of the PAS domain in
regulation of dimerization and DNA binding specificity of the dioxin receptor. *Mol. Cell. Biol.* **18**,
4079–4088.

Prasch, A. L., Tanguay, R. L., Mehta, V., Heideman, W. and Peterson, R. E. (2006). Identification of
zebrafish ARNT1 homologs: 2,3,7,8-tetrachlorodibenzo-p-dioxin toxicity in the developing zebrafish
requires ARNT1. *Mol. Pharmacol.* **69**, 776–787.

Qian, F., Zhen, F., Ong, C., Jin, S.-W., Meng Soo, H., Stainier, D. Y. R., Lin, S., Peng, J. and Wen,
Z. (2005). Microarray analysis of zebrafish cloche mutant using amplified cDNA and identification of
potential downstream target genes. *Dev. Dyn.* **233**, 1163–1172.

Ramírez-Bergeron, D. L., Runge, A., Dahl, K. D. C., Fehling, H. J., Keller, G. and Simon, M. C.
(2004). Hypoxia affects mesoderm and enhances hemangioblast specification during early
development. *Development* **131**, 4623–4634.

Reischauer, S., Stone, O. A., Villasenor, A., Chi, N., Jin, S.-W., Martin, M., Lee, M. T., Fukuda, N.,
Marass, M., Witty, A., et al. (2016). Cloche is a bHLH-PAS transcription factor that drives haemato-
vascular specification. *Nature* **535**, 294–298.

Sekine, H., Mimura, J., Yamamoto, M. and Fujii-Kuriyama, Y. (2006). Unique and overlapping
transcriptional roles of arylhydrocarbon receptor nuclear translocator (Arnt) and Arnt2 in xenobiotic
and hypoxic responses. *J. Biol. Chem.* **281**, 37507–37516.

Shalaby, F., Ho, J., Stanford, W. L., Fischer, K. D., Schuh, A. C., Schwartz, L., Bernstein, A. and
Rossant, J. (1997). A requirement for Flk1 in primitive and definitive hematopoiesis and
vasculogenesis. *Cell* **89**, 981–990.

Souder, J. P. and Gorelick, D. A. (2019). ahr2, But Not ahr1a or ahr1b, Is Required for Craniofacial
and Fin Development and TCDD-dependent Cardiotoxicity in Zebrafish. *Toxicol. Sci.* **170**, 25–44.

Stainier, D. Y., Lee, R. K. and Fishman, M. C. (1993). Cardiovascular development in the zebrafish. I.
Myocardial fate map and heart tube formation. *Development* **119**, 31–40.

Stainier, D. Y., Weinstein, B. M., Detrich, H. W., Zon, L. I. and Fishman, M. C. (1995). Cloche, an
early acting zebrafish gene, is required by both the endothelial and hematopoietic lineages.
Development **121**, 3141–3150.

Sumanas, S. and Lin, S. (2006). Ets1-related protein is a key regulator of vasculogenesis in zebrafish.
PLoS Biol. **4**, e10.

Sumanas, S., Jorniak, T. and Lin, S. (2005). Identification of novel vascular endothelial-specific genes
by the microarray analysis of the zebrafish cloche mutants. *Blood* **106**, 534–541.

Tanguay, R. L., Andreasen, E., Heideman, W. and Peterson, R. E. (2000). Identification and
expression of alternatively spliced aryl hydrocarbon nuclear translocator 2 (ARNT2) cDNAs from
zebrafish with distinct functions. *Biochim. Biophys. Acta* **1494**, 117–128.

Thompson, M. A., Ransom, D. G., Pratt, S. J., MacLennan, H., Kieran, M. W., Detrich, H. W., Vail,
B., Huber, T. L., Paw, B., Brownlie, A. J., et al. (1998). The cloche and spadetail genes
differentially affect hematopoiesis and vasculogenesis. *Dev. Biol.* **197**, 248–269.

Vogeli, K. M., Jin, S.-W., Martin, G. R. and Stainier, D. Y. R. (2006). A common progenitor for
haematopoietic and endothelial lineages in the zebrafish gastrula. *Nature* **443**, 337–339.

Wang, Q., Stacy, T., Binder, M., Marin-Padilla, M., Sharpe, A. H. and Speck, N. A. (1996).
Disruption of the Cbfa2 gene causes necrosis and hemorrhaging in the central nervous system and
blocks definitive hematopoiesis. *Proc Natl Acad Sci USA* **93**, 3444–3449.

Westerfield, M. (2000). *The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio*
Rerio). 4th ed. Eugene, OR: University of Oregon Press.

Wu, D. and Rastinejad, F. (2017). Structural characterization of mammalian bHLH-PAS transcription
factors. *Curr. Opin. Struct. Biol.* **43**, 1–9.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplemental files

Supplemental Table 1. Oligonucleotide primers and probes used in this study

Supplemental Table 2. Genotype versus phenotype correlation of embryos probed for *etsrp* (related to Figure 3)

Supplemental Table 3. Genotype versus phenotype correlation of embryos probed for *tal1* (related to Figure 3)

Supplemental Table 4. Genotype versus phenotype correlation of embryos probed for *lmo2* (related to Figure 3)

Supplemental Table 5. Genotype versus phenotype correlation of embryos probed for *egfl7* (related to Supplemental Figure 1)

Supplemental Table 6. Genotype versus phenotype correlation of embryos probed for *npas4l* (related to Supplemental Figure 2)

Supplemental Figure 1. *arnt1/2* mutants show reduced expression of *egfl7* in the lateral plate mesoderm (related to Figure 3)

Supplemental Figure 2. *arnt1/2* mutants and *arnt1/2* siblings have normal expression of *npas4l* (related to Figure 3)

Supplemental Figure 3. *npas4l* and *arnt2* mRNA, but not *arnt1*, are enriched in the posterior lateral plate mesoderm (related to Figure 4)

Supplemental Figure 4. *arnt1* mutants, but not *arnt2* mutants, are resistant to TCDD toxicity (related to Figure 7)

Movie 1.mp4 *arnt1$^{bcm2/+}$;arnt2$^{bcm3/+}$* 50 hpf, head and heart
Movie 2.mp4 *arnt1$^{bcm2/+}$;arnt2$^{bcm3/+}$* 50 hpf, blood flow in tail
Movie 3.mp4 *arnt1$^{bcm2/bcm2}$;arnt2$^{bcm3/bcm3}$* 50 hpf, head and heart
Movie 4.mp4 *arnt1$^{bcm2/bcm2}$;arnt2$^{bcm3/bcm3}$* 50 hpf, blood flow in tail
Movie 5.mp4 *arnt1$^{bcm2/+}$;arnt2$^{bcm3/+}$* 2 dpf, head and heart
Movie 6.mp4 *arnt1$^{bcm2/+}$;arnt2$^{bcm3/+}$* 2 dpf, blood flow in tail
Movie 7.mp4 *arnt1$^{bcm2/bcm2}$;arnt2$^{bcm3/bcm3}$* 2 dpf, head and heart
Movie 8.mp4 *arnt1$^{bcm2/bcm2}$;arnt2$^{bcm3/bcm3}$* 2 dpf, blood flow in tail

Movies 1 and 2 from the same embryo, movies 3 and 4 from the same embryo, movies 5 and 6 from the same embryo, movies 7 and 8 from the same embryo.

Primer	Forward 5'-3'	Reverse 5'-3'	Reference
bcm1-gRNA	TAGGATGGTTTATAGGTGCCAT	AAACATGGCACCTATAAACCAT	
bcm1-HRMA	ATGGCCGTATCACACATGAAG	AGACCTGGTCTGTGAGAAAGGAT	
bcm1-seq	AAATGACCGCCTACATCA	TGCCTACAGTGCATTGTT	
bcm2-gRNA	TAGGCGACTGTGAGGATGAGTA	AAACTACTCATCCTCACAGTCG	
bcm2-HRMA	GTGAAGAAGGAGGGTGGTCA	CATTCTGCAGATGAAGGAGCG	
bcm2-ISHseq	GCAAATATATCTTAAGGGGGCCAAT	TCCAGGTCAGTGTCTTCACCA	
bcm2-ISHseq	TCAGACATACTGTGAAACATTTCTTGC	CACCATGAAATAGGAGCAGCTTG	
bcm2-seq	AAACTCACTATCCTGCGCATG	AGACCTGGTCTGTGAGAAAGGAT	
bcm3-gRNA	TAGGTCTCGGTCCGTCTAAGGA	AAACTCCTTAGACGGACCGAGA	
bcm3-HRMA	TGTGAGCCGAATGTTGGCT	ACAACAGAGTACTGCGCTTCA	
bcm3-ISHseq	CCAGGTAAAAACATGACCTGTTCC	CATTGAAACAATATGCAGCAATGA	
bcm3-ISHseq	TGTCTACAAAATCCTCCACACTGCT	TTGAATATTTCAAAATCACATGCAG	
bcm3-seq	ACATGACCTGTTCCATACC	TCAGTGCTTCAGTGTCTT	
egfl7- probe	GCGCGGTGCTCTTCATCAG	AGCCTCCTTACACTCGTCCACATC	
lmo2-probe	GTGCGGCAGATGGTCTTTCGGAACC	CTGCAACCCAACACTGGGAAACCTG	(Sun *et al*. 2018)

Table S1. Oligonucleotide primers and probes used in this study

Genotype		Phenotype (number of embryos)	
Allele designation	Number of mutant alleles	Normal expression	Reduced expression
arnt1 $^{bcm2/bcm2}$; arnt2 $^{bcm3/bcm3}$	4	2	24
arnt1 $^{bcm2/bcm2}$; arnt2 $^{bcm3/+}$	3	4	0
arnt1 $^{bcm2/+}$; arnt2 $^{bcm3/bcm3}$		4	0
arnt1 $^{bcm2/+;}$ arnt2 $^{bcm3/+}$	2	5	0
arnt1 $^{bcm2/bcm2}$		3	0
arnt2 $^{bcm3/bcm3}$		2	0
arnt1 $^{bcm2/+}$	1	4	0
arnt2 $^{bcm3/+}$		3	0
wild-type	0	1	0

Table S2. Genotype versus phenotype correlation of embryos probed for *etsrp* (See Figure 3). There was a significant correlation between the number of mutant alleles and *etsrp* expression phenotype, $r(47)=-0.9, p<0.001$

Genotype		Phenotype (number of embryos)	
Allele designation	Number of mutant alleles	Normal expression	Reduced expression
arnt1 ^{bcm2/bcm2} ; *arnt2* ^{bcm3/bcm3}	4	1	14
arnt1 ^{bcm2/bcm2} ; *arnt2* ^{bcm3/+}	3	3	3
arnt1 ^{bcm2/+} ; *arnt2* ^{bcm3/bcm3}		1	0
arnt1 ^{bcm2/+;} *arnt2* ^{bcm3/+}	2	1	0
arnt1 ^{bcm2/bcm2}		0	0
arnt2 ^{bcm3/bcm3}		0	0
arnt1 ^{bcm2/+}	1	1	0
arnt2 ^{bcm3/+}		2	0
wild-type	0	0	0

Table S3. Genotype versus phenotype correlation of embryos probed for *tal1* (See Figure 3). There was a significant correlation between the number of mutant alleles and *tal1* expression phenotype, *r(25)=-0.6, p<0.001*

Genotype		Phenotype (number of embryos)	
Allele designation	Number of mutant alleles	Normal expression	Reduced expression
arnt1 $^{bcm2/bcm2}$; arnt2 $^{bcm3/bcm3}$	4	5	7
arnt1 $^{bcm2/bcm2}$; arnt2 $^{bcm3/+}$	3	23	5
arnt1 $^{bcm2/+}$; arnt2 $^{bcm3/bcm3}$		13	3
arnt1 $^{bcm2/+;}$ arnt2 $^{bcm3/+}$	2	47	3
arnt1 $^{bcm2/bcm2}$		5	0
arnt2 $^{bcm3/bcm3}$		10	2
arnt1 $^{bcm2/+}$	1	28	1
arnt2 $^{bcm3/+}$		19	2
wild-type	0	7	2

Table S4. Genotype versus phenotype correlation of embryos probed for *lmo2* (See Figure 3). There was a significant correlation between the number of mutant alleles and *lmo2* expression phenotype, *r(184)=-0.2, p<0.005*

Genotype		Phenotype (number of embryos)	
Allele designation	Number of mutant alleles	Normal expression	Reduced expression
arnt1 $^{bcm2/bcm2}$; *arnt2* $^{bcm3/bcm3}$	4	2	6
arnt1 $^{bcm2/bcm2}$; *arnt2* $^{bcm3/+}$	3	4	0
arnt1 $^{bcm2/+}$; *arnt2* $^{bcm3/bcm3}$		6	1
arnt1 $^{bcm2/+;}$ *arnt2* $^{bcm3/+}$	2	6	1
arnt1 $^{bcm2/bcm2}$		13	2
arnt2 $^{bcm3/bcm3}$		2	0
arnt1 $^{bcm2/+}$	1	36	0
arnt2 $^{bcm3/+}$		3	0
wild-type	0	12	0

Table S5. Genotype versus phenotype correlation of embryos probed for *egfl7* (See Supplemental Figure 1). There was a significant correlation between the number of mutant alleles and *egfl7* expression phenotype, *r(94)=-0.5, p<0.001*

Genotype		Phenotype (number of embryos)	
Allele designation	Number of mutant alleles	Normal expression	Reduced expression
arnt1 $^{bcm2/bcm2}$; *arnt2* $^{bcm3/bcm3}$	4	2	0
arnt1 $^{bcm2/bcm2}$; *arnt2* $^{bcm3/+}$	3	4	0
arnt1 $^{bcm2/+}$; *arnt2* $^{bcm3/bcm3}$		11	0
arnt1 $^{bcm2/+;}$ *arnt2* $^{bcm3/+}$	2	19	1
arnt1 $^{bcm2/bcm2}$		3	1
arnt2 $^{bcm3/bcm3}$		5	0
arnt1 $^{bcm2/+}$	1	11	0
arnt2 $^{bcm3/+}$		5	0
wild-type	0	8	0

Table S6. Genotype versus phenotype correlation of embryos probed for *npas4l* (See Supplemental Figure 2). There was no significant correlation between the number of mutant alleles and *npas4l* expression phenotype, *r(69)=-0.05, p>0.05*

Supplemental Figure 1. *arnt1/2* mutants show reduced expression of *egfl7* in the lateral plate mesoderm. Whole-mount in situ hybridization for *egfl7* was performed on 10 somite stage embryos from *arnt1bcm2/+;arnt2bcm3/+* parents (N=2 biological replicates, 94-120 embryos per clutch, 214 total embryos) or from wildtype (AB) parents (N=1 biological replicate, 28 total embryos). (**A** and **D**) expression of *egfl7* in wildtype embryos is restricted to the anterior lateral plate mesoderm (ALPM) (**A**) and the posterior lateral plate mesoderm (PLPM) (**D**). (**B** and **E**) Expression of *egfl7* was observed in the ALPM (**B**) and the PLPM (**E**) in 95% of embryos from *arnt1bcm2/+; arnt2bcm3/+* parents (n=203 out of 214 total embryos). (**C** and **F**) Limited or no expression of *egfl7* was observed in the ALPM and PLPM in 5% of embryos from *arnt1bcm2/+;arnt2bcm3/+* parents (n=11 out of 214 total embryos), which is consistent with the expected Mendelian ratio of 6.25% *arnt1bcm2/bcm2;arnt2bcm3/bcm3* (*arnt1/2* mutants, binomial test, $p<0.05$). All embryos are oriented with dorsal towards the top of the image. Fractions in bottom right corners refer to the number of embryos with the indicated phenotype over the total number of embryos examined. See **Supplemental Table 5** for genotype frequencies. Scale bar, 100μm.

Supplemental Figure 2. *arnt1/2* mutants and *arnt1/2* siblings have normal expression of *npas4l*. Whole-mount in situ hybridization for *npas4l* on 4 somite stage embryos from *arnt1^{bcm2/+}; arnt2^{bcm3/+}* parents (N=2 biolgical replicates, 68-100 embryos per clutch, n= 168 embryos total) or from wild-type (AB) parents (N=1 biological replicate, n=27 embryos total). (**A**) Expression of *npas4l* was observed in the anterior lateral plate mesoderm (ALPM) of wild-type embryos. (**B**) Expression of *npas4l* was observed in the ALPM embryos from *arnt1^{bcm2/+}; arnt2^{bcm3/+}* parents, including the *arnt1/2* mutants. (**C**) Expression of *npas4l* was observed in the PLPM of wild-type embryos. (**D**) Expression of *npas4l* was observed in the posterior lateral plate meosderm (PLPM) embryos from *arnt1^{bcm2/+}; arnt2^{bcm3/+}* parents, including the *arnt1/2* mutants. Embryos oriented with dorsal towards the top of the image. The fractions in the right-hand corners of **C** and **D** refer to the number of embryos with the represented phenotypes over the total number of embryos. Scale bar, 100 μm.

Supplemental Figure 3. Expression of *npas4l*, *arnt1* and *arnt2* mRNA in the anterior lateral plate mesoderm. Companion to Figure 4, which shows expression in the posterior lateral plate mesoderm. Whole-mount *in situ* hybridization chain reaction was performed on wild-type (AB) embryos at the 4 somite stage. **(A)** Embryos were probed for *npas4l, arnt1,* and *arnt2* expression and counter-stained with DAPI. **(A')** Higher magnification views of the boxed areas in (A) shows the anterior lateral plate mesoderm, outlined in cyan. N=3 biological replicates, 5-9 embryos per replicate, 22 embryos total. **(B)** To demonstrate that the observed labeling is not autofluorescence of the embryo or the yolk, wild-type embryos were processed identically to those in (A), but no anti-sense RNA probe sets were added. The views in (B) are identical magnification as those in (A'). N=3 biological replicates, 6-12 embryos per replicate, 28 embryos total. Images are maximum intensity projections from z-stacks taken every 2.5 µm. In merged panels, *npas4l* is green, *arnt1* is magenta, *arnt2* is cyan. The anterior portion of the embryo is in view, dorsal is towards the top of the image. Scale bars = 100 µm. **(C-E)** *npas4l* and *arnt2* mRNA, but not *arnt1*, are enriched in the posterior lateral plate mesoderm. Quantification of the fluorescence integrated intensity of *npas4l, arnt1,* and *arnt2* labeling within the *npas4l*-positive cells of the anterior and posterior lateral plate mesoderm (see panel A and Figure 4). We calculated the integrated intensity of fluorescence (pixel area, pixel intensity) over each side of the anterior or posterior LPM in maximum intensity projections. Each point represents the integrated intensity from one side of the LPM from a single embryo (right or left side), where each embryo has two bilateral *npas4l*-positive regions of the LPM. Points of the same color are measurements from opposite sides of the LPM from the same embryo. 6 representative embryos were used for this analysis, in 2 of these embryos the anterior LPM was damaged during imaging and could not be analyzed. **(C)** *npas4l* expression was statistically significantly increased in the posterior versus anterior LPM, Welch's t-test, *t(12.36)=6.4, p<0.0001.* **(D)** *arnt1* was expressed at similar levels in the anterior and posterior, Mann-Whitney, U=45, p>0.05. **(E)** *arnt2* expression was statistically significantly increased in the posterior versus anterior LPM, Welch's t-test, *t(11)= 5.0, p<0.0005.*

Supplemental Figure 4. *arnt1* mutants, but not *arnt2* mutants, are resistant to TCDD toxicity. Representative images of larvae at 3 days post fertilization (dpf) exposed to either DMSO or TCDD. Mutant larvae were derived from heterozygous parents and genotyped following imaging (denoted as wild-type +/+, heterozygous -/+, or homozygous -/- in panels **C-H**). (**A-H**) overview images of treated larvae, red asterisks denote larvae detailed in panels A'-H' and A''-H''. Black arrows in panels A''-H'' point to the heart within the pericardial cavity. Pericardial edema and incomplete heart looping is present in 100% of wild-type (n=3 biological replicates per genotype, 10-60 larvae per clutch per treatment, 157 larvae total) and *arnt2* mutant larvae (n=3 biological replicates per genotype, 40-50 larvae per clutch per treatment, 253 larvae total). In contrast, 100% of *arnt1* mutants (3-4 clutches per allele, 21-36 larvae per clutch per treatment, 533 larvae total, 248 larvae from allele bcm1 and 285 larvae allele from bcm2) are resistant to TCDD cardiotoxicity. Scale bars, 100 µm. (**I** and **J**) Percent larvae from each clutch demonstrating cardiotoxicity (red dots/red bars), mortality (black dots/gray bars), or normal development (white dots/white bars) at 3 dpf. All clutches were derived from heterozygous parents and embryos were analyzed prior to genotyping. Each clutch contain a mix of wild-type, heterozygous, and homozygous larvae. Each dot represents the percent of larvae from a single clutch and bars represent the mean. (**J**) In clutches derived from *arnt2*bcm3/+ parents, nearly 100% of embryos exhibited cardiotoxicity following TCDD exposure. In clutches derived from *arnt1* heterozygous parents, approximately 75% of embryos exhibited cardiotoxicity or death, while 25% of embryos exhibited a normal phenotype (blue arrows), consistent with the expected Mendelian ratio of *arnt1* homozygotes. Normal larvae were not observed in wild-type or *arnt2* mixed clutches exposed to TCDD. (**K-M**) We genotyped the embryos from (**J**) and graphed the number of embryos per clutch exhibiting cardiotoxicity (red dots/red bars) or normal development (white dots/white bars). Each dot represents the number of larvae from a single clutch. (**K,L**) All *arnt1* homozygous mutants (*bcm1* or *bcm2*) exhibited normal phenotype following TCDD exposure (blue arrows), while *arnt1* heterozygotes and (**M**) *arnt2* mutants exhibited cardiotoxicity. Unpaired Welch's t-tests, * *p*<0.05, ** *p*<0.005, and *** *p*<0.001.