Cost-effectiveness of quadrivalent versus trivalent inactivated influenza vaccines for the Portuguese elderly population

Diana Tavares[1¶], Helena Mouriño[1*¶], Cristina Antón Rodríguez[2¶], Carlos Martín Saborido[3¶]

[1]Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal

[2]Facultad de Medicina, Universidad Francisco de Vitoria, Madrid, España

[3]Hospital La Paz Institute for Health Research, IP Research on Evidence and Decision Making Group, Higher Education Center Hygiea. UDIMAMadrid, Madrid. España

* Corresponding author

E-mail: mhnunes@fc.ul.pt

¶ This authors contributed equally to this work.

Abstract

Background

Quadrivalent Inactivated Vaccine (QIV) is expected to replace Trivalent Inactivated Vaccine (TIV) over time. In Portugal, TIV is free of charge for risk groups, which include older adults. In its turn, QIV – which provides broader protection as it includes an additional lineage B strain – was introduced in Portugal in October 2018, but only from the influenza season 2019/20 was provided free of charge for the risk groups. This study evaluates the cost-effectiveness of switching from TIV to QIV from the National Health Service perspective in the Portuguese elderly mainland population (≥ 65 years old).

Methods

A decision tree model was developed to compare TIV and QIV, based on Portuguese hospitalization data for the 2015/16 influenza season. The primary health economic outcome under consideration was the Incremental Cost-Effectiveness Ratio (ICER). In addition, a one-Way Sensitivity Analysis was performed to evaluate the impact of model parameters on the ICER; Probabilistic Sensitivity Analysis was also carried out to analyze the robustness of the base case results.

Results

The high cost of QIV (approximately three times the cost of TIV) would lead to a total increment of 5,283,047 €, and the resulting ICER would be 26,403,007€/QALY, mainly above the usual willingness-to-pay threshold.

Conclusions

From the National Health Service perspective, our findings reveal that QIV is not cost-effective for the Portuguese elderly population due to the high cost. If the QIV costs were the same as the TIV, then QIV would be cost-effective.

Keywords: Influenza vaccines; Decision tree; Cost-Effectiveness analysis; One-way sensitivity analysis; Probabilistic sensitivity analysis

1   Introduction

Seasonal influenza is an acute respiratory disease caused by infection with influenza viruses(1). There are

four types of seasonal influenza viruses: A, B, C and D(1). Only influenza A and B viruses circulate and

cause seasonal epidemics of disease(1). The infection may cause signs and symptoms like fever, cough,

headache, muscle and joint pain and weakness(1).

Influenza outbreaks are recorded every year. In temperate zones of the northern and southern

hemispheres, epidemics occur during the winter, while in the tropics they occur throughout the year(2,3).

In Portugal, influenza outbreaks are characterized by significant morbidity in the general population and

increased mortality rates. The high-risk groups are: elderly (≥65 years old), patients with chronic or

immunosuppressive conditions aged six months or older, pregnant women, health professionals and other

healthcare givers(4).

An influenza pandemic can appear when a new and different type of influenza A virus emerges, and it

can infect humans who are not immunized(1,5). The most recent pandemic was in 2009, caused by the

A(H1N1)2009 virus, after which it became a seasonal influenza serotype(1). In Portugal, 1,189 people

were hospitalized, around 10% in intensive care units, resulting in 124 deaths(6).

According to the World Health Organization (WHO), vaccination is the most effective way to prevent

seasonal influenza and subsequent severe outcomes(1). Therefore, vaccination should be administered

annually to provide optimal protection. Influenza Surveillance System in Portugal is usually activated on

week 40 (October) of a given year (n) and lasts up to week 20 (April) of the following year (n+1) (7,8).

Thus, it is recommended to administer the vaccine during autumn and winter, preferably until the end of

the year n (4). The development of immunity against influenza viruses takes about two weeks after

vaccination(9). However, vaccine effectiveness depends on the influenza subtypes included in the

vaccines (a mismatch occurs when the viruses in the vaccine are different from the circulating viruses) and

the population vaccination coverage. Unfortunately, several European countries reported a decline in

vaccination coverage among older people from 2008/09 to 2014/15 seasons. Conversely, Portugal

reported an increase in vaccination uptake throughout these seasons(10). In the 2015/16 season, influenza

vaccination coverage was 50.1% in Portugal(7). The coverage rate has still increased in the last few years.

For instance, in the 2017/18 season, it has increased to 60.8% for people aged 65 years or above(11).

A Cochrane study on this topic concluded that the risk of influenza decreases from 6% to 2.5%, and the

risk of Influenza-Like Illness (ILI) reduces from 6% to 3.5%, between unvaccinated and vaccinated

groups (≥ 65 years old) during a single season(12). In the European Union, it is estimated that influenza

vaccination prevents up to 37,000 deaths each year(13).

There are two main types of seasonal influenza vaccines, namely Inactivated Influenza Vaccine (IIV)

and Lived Attenuated Influenza Vaccine (this one is not within the scope of the paper)(5,12). The different

IIV developed are: trivalent, trivalent adjuvanted and quadrivalent. Trivalent Inactivated Vaccine (TIV) is

the most common IIV and protect against three different influenza viruses (two influenza A strains and

one influenza B lineage). In some EU/EEA countries, the trivalent adjuvanted vaccine is available for

older people to empower immune response(5,14). Quadrivalent inactivated Influenza Vaccine (QIV) is

available since the 2014/2015 season in some EU/EEA countries, and it is expected to replace TIV over

time(5). It contains one more influenza B strain in addition to those included in TIV, which provides wider

protection against influenza viruses. TIV is available in all EU/EEA countries, while QIV is only available

in some of them. In Portugal, QIV was launched in October 2018, but only from the influenza season

2019/20 it was provided free of charge for the risk groups.

Influenza is, thus, a major global health threat that affects all countries, not only from a health

perspective but also from economic and social approaches. Therefore, it is crucial to carry out economic

evaluations of the influenza vaccination programmes.

This work aims to estimate the cost-effectiveness of switching from TIV to QIV based on the National

Health Service perspective. The results are presented as incremental cost per unit of effect, described by

the Incremental Cost-Effectiveness Ratio (ICER). As far as we know, no cost-effectiveness analysis of

Portuguese influenza vaccination has been reported in the literature yet.

Methods

Key Features of the Economic Evaluation

The target population is the population of mainland Portugal aged 65 and above because they are one of the most relevant influenza risk groups. In 2015, they represented 21% of the Portuguese population(15). This group is usually retired, so productivity losses are not relevant. The cost-effectiveness evaluation was performed from the National Health Service (NHS) perspective, where only direct costs were considered. Thus, only the medical costs paid by the NHS are considered here.

The Influenza Surveillance System runs from week 40 of a specific year until week 20 of the following year, so the time horizon established for this work was one year. Therefore, data under analysis refers to the 2015/16 influenza season.

As stated above, economic evaluations are performed to identify, measure, value and compare the costs and consequences of different therapeutic alternatives in what respects to influenza vaccination. To perform the cost-effectiveness analysis, health outcomes generated by the two alternative interventions under analysis are assessed through the Quality-Adjusted Life Year (QALY). It measures the length of life-years (i.e., quantity gains) and the quality of life (i.e., quality gains). The quality of life is quantified by the notion of utility: the greater the preference for a particular state, the greater its utility. Utilities incorporate in the analysis the preferences of individuals for different treatment-related outcomes. Another key measure is the concept of quality-of-life loss due to the disease (influenza, in this case), that is, disutility. It corresponds to the reverse of utility, i.e., decrease in utility due to serious health complications. These quantities are obtained by Health-related quality of life (HRQoL) instruments, such as the EQ-5D family of questionnaires. Studies about utilities and disutilities for the Portuguese population are very scarce. In this work, utilities associated with different health states were obtained from a study on *EQ-5D Portuguese Population norms*(16). Data available were stratified by age, but unfortunately, there was no information for people aged 65 or over. Therefore, the utilities for that age group were derived by interpolation (see S1 Appendix). Moreover, no Portuguese studies on disutilities

were found for influenza. Such data were extracted from a Spanish study (17) (see S1 Appendix). All

these results were thus combined in a single index, QALY(18,19).

Incremental Cost-Effectiveness Ratio (ICER) is the primary cost-effectiveness outcome used in this

work(19). To make recommendations to policymakers, ICER is usually compared with an established

threshold, called Willingness-To-Pay (WTP). Some decision-makers have established a Willingness-To-

Pay (WTP) threshold, that refers to the maximum ICER accepted for an intervention to be considered

cost-effective. In Portugal, there is no established cost-effectiveness threshold for health interventions.

Some Portuguese authors refer to a WTP threshold of 30,000€/QALY(20,21), while others consider the

cost-effectiveness threshold twice the Gross Domestic Product (GDP) per capita or Gross National Income

(GNI) per capita(21,22). In 2015, the Portuguese GNI per capita was 16,887.36€, resulting in a ceiling

ratio of approximately 34,000€ (twice the GNI)(23,24). In this paper, the ICER will be used to provide

information on the extra amount that is necessary to pay in order to gain an extra QALY when the most

effective alternative is chosen(19,25).

Decision Tree

The cost-effectiveness analysis was based on a decision tree model, a very powerful tool in this context

because it incorporates probabilities of the outcomes, resource costs, vaccination coverage rate, prices of

the vaccines, efficacy, utilities and disutilities. It consists of a series of pathways representing possible

prognoses for each alternative therapy being evaluated. A static model was used, and no herd effects of

vaccination were evaluated.

One decision tree model was developed for each alternative therapy (TIV and QIV). For convenience,

each decision tree was split into two different diagrams referring to vaccinated and unvaccinated people.

For illustration, Fig 1 displays the branch of the decision tree for vaccinated individuals.

Fig 1. Decision tree for the cost-effectiveness analysis. Branch for the vaccinated people aged ≥ 65 years old.

ILI, influenza-like-illness; H, healthy; D, death; Conf. Influenza, Confirmed Influenza; GP Consultation, General Practitioner Consultation; Pneu. Hosp., Hospitalization due to pneumonia; RD Hosp., Hospitalization due to respiratory disease; HD Hosp., Hospitalization due to heart disease; Inf. Hosp., Hospitalization due to Influenza.

It was assumed that influenza confirmation (through diagnostic tests) was carried out in the context of General Practitioner (GP) consultation, pneumonia hospitalization, respiratory disease hospitalization or heart disease hospitalization. Although it is known that influenza may be diagnosed in many other situations, the diseases described above are usually associated with complications due to influenza in the elderly. It should be noted that influenza event (as a secondary diagnosis) may have occurred before or after hospital admission, in which principal diagnoses was pneumonia, respiratory disease or heart disease. It is important to note that the "Healthy" state is assumed to include all individuals who did not die, regardless of their health conditions.

Input Parameters and Computation of the Probabilities

The decision tree is powered by different types of parameters. A detailed description of each of these measures is provided in the S2 Appendix (which includes S1 Table). A summary of the base case values for the parameters under consideration is displayed in Table 1. The computation of the different parameters involved in the decision tree pathways for the TIV – probabilities, costs and QALY – are given in the S1 Fig and S2 Table. Similar calculations are made for the QIV scenario.

121   **Table 1. Input parameters for the decision tree.** Base case values.

Input Parameters	Base Case	Source
Population	2,067,654	(26)
Vaccination Coverage Rate	0.501	(7)
Trivalent Vaccine Effectiveness	0.580	(12)
Quadrivalent Vaccine Effectiveness	0.599	(27–35)
Probability of Influenza-like-illness	0.007	(7,12,26,31,36)
Probability of Confirmed Influenza	0.278	(31)
Probability of GP Consultation	0.952	(31,36)
Probability of Hospitalization due to Influenza	0.091	(37)
Probability of death when Hosp. Influenza	0.079	(37)
Probability of Hospitalization due to Pneumonia	0.008	(37)
Probability of death due to Hosp. Pneumonia	0.381	(37)
Probability of Hospitalization due to Respiratory Disease	0.014	(37)
Probability of death when Hosp. RD	0.050	(37)
Probability of Hospitalization due to Heart Disease	0.026	(37)
Probability of death when Hosp. HD	0.110	(37)
Probability of death when no confirmed influenza	0.043	(26,38)
Cost of GP consultation	31.000	(39)
Cost of Hospitalization due to Influenza	2644.011	(37,39)
Cost of death when Hospitalization due to Influenza	6364.597	(37,39)
Cost of Hospitalization due to Pneumonia	2906.749	(37,39)
Cost of death when Hospitalization due to Pneumonia	11298.219	(37,39)
Cost of Hospitalization due to Respiratory Disease	2534.841	(37,39)
Cost of death when Hospitalization due to RD	1363.460	(37,39)
Cost of Hospitalization due to Heart Disease	2978.842	(37,39)
Cost of death when Hospitalization due to HD	7785.951	(37,39)
Cost of antiviral treatment	18.300	(40,41)
Cost of trivalent vaccine	2.576	(42)
Cost of quadrivalent vaccine	7.692	(43,44)

Cost of vaccine administration	3.700	(39)
Disutility associated with ILI without influenza confirmation	0.009	(17)
Disutility associated with no hospitalized influenza	0.009	(17)
Disutility associated with hospitalization due to influenza	0.031	(17)
Disutility associated with hospitalization due to pneumonia	0.031	(17)
Disutility associated with hospitalization due to respiratory disease	0.031	(17)
Disutility associated with hospitalization due to heart disease	0.031	(17)
Utility associated with healthy population	0.625	(16,26)

Scenario Analysis

The most common ways to characterize parameter uncertainty are: One-way Sensitivity Analysis (OWSA)

and Probabilistic Sensitivity Analysis (PSA)(19,25).

OWSA consists of varying the point estimates of the input parameters one at a time within a plausible

range of values and assessing the impact on model outcomes(19,25), which corresponds to the ICER.

When the Confidence Interval (CI) for a specific parameter was known from the literature, these limits

were used as the OWSA range. For parameters with no CI available, a range of ±20% was used to assess

model's sensitivity to parameters variation. A tornado diagram was used to report OWSA results.

A scenario analysis of the vaccination coverage rate, TIV effectiveness and QIV effectiveness variation

was also carried out. The cost of the quadrivalent vaccine was also studied in detail, as it was identified as

a critical parameter(45). For vaccination coverage rate and QIV cost, a sequence of parameters values was

generated with an increment of 0.05. For TIV and QIV effectiveness, the increment used was 0.005.

Instead of representing the parameters as single point estimates as in OWSA, in PSA parameters are

represented by random variables following a particular distribution. The S2 Table displays the base case

values used for each input parameter of the decision tree models. It also shows the probability distribution

(with information on location, shape, scale, when appropriate) used for each parameter under

consideration. A total of 1,000 Monte Carlo (MC) simulations were performed.

140   Results

141   *Base Case Analysis*

Table 2 summarises the base case results of shifting from TIV to QIV in the elderly, based on the 2015/16

influenza season data. Results show a difference in confirmed influenza cases of approximately 37. As a

result, about 36 GP consultations could have been averted and, therefore, 1,103€ saved. In addition, five

hospitalizations, including hospitalizations due to influenza, pneumonia, respiratory disease and heart

disease, could have been averted and one life saved.

Table 2. Base Case. Results of the cost-effectiveness evaluation comparing TIV and QIV.

	TIV	QIV	Difference (QIV-TIV)
Events			
GP Consultations	2631.35	2595.77	-35.58
Hospitalizations due to Influenza	239.00	235.77	-3.23
Deaths due to Influenza Hospitalization	19.00	18.74	-0.26
Hospitalizations due to Pneumonia	21.00	20.72	-0.28
Deaths due to Pneumonia Hospitalization	8.00	7.89	-0.11
Hospitalizations due to RD	40.00	39.46	-0.54
Deaths due to RD Hospitalization	2.00	1.97	-0.03
Hospitalizations due to HD	73.00	72.01	-0.99
Deaths due to HD Hospitalization	8.00	7.89	-0.11
Vaccine Doses	1,035,895	1,035,895	0.00
Costs			
GP Consultations	81,572 €	80,469 €	-1,103 €
Hospitalizations due to Influenza	710,464 €	701,189 €	-9,275 €

Deaths due to Influenza Hospitalization	121,552 €	119,934 €	-1,617 €
Hospitalizations due to Pneumonia	128,213 €	126,508 €	-1,705 €
Deaths due to Pneumonia Hospitalization	90,401 €	89,189 €	-1,211 €
Hospitalizations due to RD	99,125 €	97,841 €	-1,285 €
Deaths due to RD Hospitalization	2,731 €	2,697 €	-34 €
Hospitalizations due to HD	256,048 €	252,687 €	-3,361 €
Deaths due to HD Hospitalization	62,303 €	61,471 €	-831 €
Vaccine Doses	2,668,465 €	7,968,557 €	5,300,092 €
Total	8,018,570 €	13,301,618 €	5,283,047 €
QALYs			
Total	1265456.95	1265457.15	0.20
ICER (€/QALY)			26,403,007

Regarding hospitalization and death costs, caution must be taken in the interpretation of the results.

The cost of death, when hospitalized, includes all hospitalization costs, and in the same way, the

hospitalization costs also include the costs of people who died. Thus, 15,625€ could have been saved in

hospitalizations: 9,275 € in hospitalizations due to influenza; 1,705 € in hospitalizations due to

pneumonia; 1,285€ in hospitalizations due to respiratory disease; and 3,361 € in hospitalizations due to

heart disease. From that value, 3,694€ are related to hospitalizations of patients who died (Table 2).

In what concerns the number of vaccinated people (mentioned in Table 2 as vaccine doses), the value

is the same for both strategies. It was an expected result because the vaccination coverage rate applied to

TIV and QIV models was the same (the paper's focus is to evaluate the cost-effectiveness of shifting from

TIV to QIV). However, there is a difference in the expected costs computed from the two strategies

because they depend on the cost of each vaccine. A total of 1,035,895 vaccine doses were expected to

have been administered to individuals aged ≥65 years old in the 2015/16 season. For the TIV strategy, this

represented spending 2,668,465 €, while for QIV corresponds to 7,968,557 €, that is, shifting from TIV to

QIV would result in an additional cost of 5,300,092 € (Table 2).

The expected difference in costs between TIV and QIV is 5,283,047 €, while the difference in effects

(QALYs) is expected to be 0.20. These values result in a base case ICER of 26,403,007€/QALY (Table

2). The plot of the differences in costs and effects on the cost-effectiveness plane shows the value under

consideration is in the first quadrant. Thus, for a cost-effectiveness threshold of 34 000€/QALY (twice the

GNI per capita), the new vaccine would not be cost-effective.

One-way Sensitivity Analysis

Fig 2 summarizes the one-way sensitivity analysis using the tornado diagram.

Fig 2. One-way sensitivity analysis. Tornado diagram.

Fig 2 reveals that the disutility associated with ILI when no confirmed influenza, and the disutility

associated with no hospitalized influenza have the highest impact on the ICER. For example, if the first

parameter mentioned above assumes the value of 0.007 (lower bound), it results in an ICER of 19,219,556

€/QALY; and the value of 0.011 (upper bound) results in an ICER of 42,160,989 €/QALY. This can be

explained by the increase in incremental QALYs, yielded by decreasing this parameter, resulting in a

reduced ICER and vice-versa. In contrast, the variation of the second parameter within the 95% CI (0.007,

0.011) produced ICERs of 39,019,821€/QALY and 19,951,738€/QALY, respectively. Thus, while the

pathway corresponding to vaccinated patients with ILI but not confirmed influenza is more populated in

the QIV model than in TIV, the same is not valid for the pathway referring to vaccinated patients with no

hospitalized influenza. Thus, a reduction in disutility associated with no hospitalized influenza would

decrease incremental QALYs and increase ICER.

The cost of the quadrivalent vaccine is the third parameter appearing in the tornado diagram (Fig 2).

The variation of the cost of the quadrivalent vaccine within a ±20% range results in an ICER of

18,438,140 €/QALY (lower bound) and 34,367,874€/QALY (upper bound), respectively. The high impact

of QIV's cost on ICER was expected if attention is paid to the difference in costs of QIV and TIV vaccine

doses (Table 2). The cost of QIV's vaccines doses is largely above the cost of TIV's vaccine doses.

Further analysis was performed, varying the QIV cost within a broader range (Fig 3). For a QIV cost equal

to TIV cost, an ICER of -85,182€/QALY was obtained, meaning that shifting from TIV to QIV would be

cost-saving, as expected. However, a value higher than 2.576€ would quickly lead to being not cost-

effective.

Fig 3. One-way sensitivity analysis of the cost of QIV. Variation between 2.576€ (the cost of TIV) and

9.756€ with increments of 0.05€.

The probability of ILI and the probability of confirmed influenza also have a high impact on ICER (Fig

2). Varying these parameters 20% downward and upward produced ICERs of 33,025,054€/QALY and

21,988,309€/QALY, respectively. Thus, as the probability of ILI and/or the probability of confirmed

influenza increases, the ICER decreases.

One-way sensitivity analyses of the effectiveness of QIV and TIV are displayed in S2 Fig and S3 Fig,

respectively. If the TIV effectiveness is equal to QIV effectiveness, the ICER is not defined as it results in

a difference in effects equal to zero and, therefore, a division by zero, and a vertical asymptote at 59.91%

is shown (S2 Fig). As TIV effectiveness approaches 59.91% from the left, the ICER tends to positive

infinity. Otherwise, as TIV effectiveness approaches 59.91% from the right, the ICER tends to negative

infinity. Such behavior is explained by the difference in effects close to zero when TIV and QIV

effectiveness are close. A negative ICER is produced for TIV effectiveness values higher than QIV

effectiveness as the incremental QALYs value is negative. Similarly, S3 Fig shows a vertical asymptote

when QIV effectiveness equals 58%; ICER tends to infinity when it approaches the vertical asymptote

from the right. Finally, the higher the effectiveness of quadrivalent vaccines (>58%), the lower the ICER.

Although vaccination coverage rate does not significantly impact ICER as the above parameters do, a

detailed study was also performed for this parameter (S4 Fig). A higher coverage rate would result in

decreased ICER. As recommended by the World Health Assembly, for a coverage rate of 75%,

21,010,680 €/QALY would be obtained, corresponding to a reduction of 20,42%.

Probabilistic Sensitivity Analysis

Probabilistic sensitivity analysis allows one to evaluate the robustness of the base case results. To attain

this goal, a probability distribution was assigned to each parameter of the model to reflect parameter

uncertainty due to sampling errors. Parameters values were then sampled 1,000 times; expected costs and

QALY were recorded from the 1,000 MC simulations. Finally, some empirical statistics such as mean,

standard deviation and confidence intervals were computed.

In general, the results from PSA (S3 Table) are similar to those obtained from the base case analysis

(Table 2). The estimated mean costs difference between QIV and TIV is 5,293,456 € with an interval

estimate of (4,375,217€; 6,088,637€), while the estimated mean effect difference corresponds to 0.20

QALYs with an interval varying from 0.06 to 0.43 QALYs. On its turn, mean ICER is estimated at

34,501,793€/QALY (13,052,665€/QALY; 87,418,632€/QALY). This value is about 31% higher than in

the base case result.

S5 Fig displays the scatterplot of the 1,000 MC simulations to compare TIV and QIV, that is, the

graphical representation of costs against QALYs. The base case results are also shown. QALY of both

interventions varies within a similar range of values (it varies between 1,155,190 and 1,375,507). On the

other hand, the difference between TIV and QIV in terms of costs is very well delimited, being QIV more

costly than TIV (QIV: mean=13,323,148€; SD=987,347€; TIV: mean=8,029,692€; SD=548,739€).

Fig 4 shows the results of the simulations in a cost-effectiveness plane, where the incremental costs are

plotted against the incremental effects. According to Fig 4, the cost-effectiveness results are robustly

located in the first quadrant of the cost-effectiveness plane, and all simulated ICERs are higher than CE

thresholds previously established.

Fig 4. Probabilistic sensitivity analysis for TIV and QIV. Cost-effectiveness plane based on data from

the 2015/16 influenza season.

Discussion

To the best of our knowledge, this is one of the first published studies performing a cost-effectiveness

evaluation of seasonal influenza vaccination in Portugal. Only a few papers studied the burden of disease

and vaccination effectiveness in the Portuguese population, for instance, the international I-MOVE+

project(46–50).

We evaluated the cost-effectiveness of switching from TIV to QIV based on the 2015/16 influenza

season data. Base case results revealed that the universal substitution of TIV with QIV would result in an

ICER of 26,403,007€ per QALY gained. As such, for the cost-effectiveness thresholds of 30,000€/QALY

or 34,000€/QALY, QIV is not cost-effective. Probabilistic sensitivity analysis enhanced the robustness of

the base case results. The ICER is much higher than any possible ceiling ratio established by NHS. Such

results are comparable to those obtained for the first year of QIV administration to Hong Kong elderly

(≥80 years old) when a cost-effectiveness analysis over nine seasons was performed(51). The need for a

longer time horizon study is here emphasized.

The majority of the published international papers(45,52–60)does not report such a high ICER, and

QIV is usually identified as cost-effective compared with TIV. As cited above, one of the reasons might

be using of a long time horizon to calculate QIV effectiveness, which allowed to assume a few seasons

that match the TIV with the circulating strains and therefore low mismatching (see S2 Appendix). Another

reason might be the cost of QIV, which in this study is about three times the TIV. Regarding the

quadrivalent vaccine, its cost is here 200% higher than TIV. However, in the literature, the additional cost

of QIV is rarely higher than 100% of the TIV cost(52,54–56), which might be one of the reasons why QIV

is cost-effective in some studies. To be cost-effective, the cost of QIV would need to be the same as TIV

in our case.

In the literature, the probability of influenza-like-illness and the probability of confirmed influenza was

found to be higher than those used here. For example, Capri et al.(60) applied a probability of influenza of

6.4%, resulting from the product of ILI attack rate (16.8%) by influenza virus isolation rate (32.1%), while

Van Bellinghen et al.(55) used a probability of symptomatic influenza infection of 6.17%. In our study,

the probability of confirmed influenza was derived from a Portuguese report on influenza surveillance,

and the probability of ILI was determined from other input values. It is comparable to the value reported

by INSA for the 2015/16 influenza season(31).

One-way sensitivity analysis demonstrated that our model is more sensitive to disutility associated with

ILI, disutility associated with no hospitalized influenza, cost of quadrivalent vaccine and probabilities of

confirmed influenza and ILI. The most relevant input parameters obtained from the OWSA are in line

with other OWSA results in the literature(52,55,56). Considering the PSA, the results are in line with the

base case conclusions and confirm their robustness.

A limitation of our study has to do with the fact that the model is static in a way that herd effects of

immunization are not considered, and only direct protection is captured. Thus, the impact of vaccination

on the burden of the disease might be underestimated. In the future, a dynamic transmission model might

be a possible approach to account for the impact of vaccination on influenza transmission(45,56). In

addition, side effects of vaccination were not included in this study, but QIV safety is assumed to be

comparable to that of TIV, and side effects are assumed to be mild and temporary. Thus, no impact is

expected on the ICER(56).

Summing up, this study contributes to understanding the impact of annual influenza epidemics on

health economics and public health in Portugal.

279 Acknowledgments

280   H. Mouriño was supported by the National Funding from FCT – Fundação para a Ciência e Tecnologia,

281   under the project: UIDB/04561/2020.

282 References

1.   WHO | Influenza (Seasonal). WHO [Internet]. 2018; Available from:

http://www.who.int/mediacentre/factsheets/fs211/en/

2.   Shi A, Zeng D, Lin S-S, Caldwell L. Harrison's - Principles of Internal medicine. 16th ed.

McGraw-Hill; 2011. 515 p.

3.   Villacampa T. Manual Curso Intensivo Mir Asturias - Enfermedades Infecciosas. 2017.

4.   Direção Geral de Saúde. Orientação Vacinação contra a gripe. Época 2017/2018. 2017;1–6.

5.   Types of Influenza Viruses | Seasonal Influenza (Flu) | CDC [Internet]. [cited 2018 Mar 6].

Available from: https://www.cdc.gov/flu/about/viruses/types.htm

6.   George F. História da gripe. 2014;1–28.

7.   Sousa Uva M, Kislaya I, Roquette R, Rodrigues AP, Machado A. Vacinação antigripal da

população portuguesa na época 2015/2016: estudo na amostra ECOS. 2016 Oct 13 [cited 2018 Sep

1]; Available from: http://repositorio.insa.pt/handle/10400.18/4013

8.   Instituto Nacional de Saúde Dr. Ricardo Jorge. Influenza Vaccine Effectiveness in Portugal.

Season 2015-16 Report. 2017. 44 p.

9.   Key Facts About Influenza (Flu) | Seasonal Influenza (Flu) | CDC [Internet]. [cited 2018 Mar 9].

Available from: https://www.cdc.gov/flu/keyfacts.htm

10.   Jorgensen P, Mereckiene J, Cotter S, Johansen K, Tsolova S, Brown C. How close are countries of

the WHO European Region to achieving the goal of vaccinating 75% of key risk groups against

influenza? Results from national surveys on seasonal influenza vaccination programmes,

2008/2009 to 2014/2015. Vaccine [Internet]. 2017;36(4):442–52. Available from:

https://doi.org/10.1016/j.vaccine.2017.12.019

11.   Machado A, Torres AR, Kislaya I, Neto M. Vacinação antigripal da população portuguesa em

2016/2017 e 2017/2018. Cobertura e características do ato vacinal. [Internet]. Lisboa; 2019.

Available from: http://repositorio.insa.pt/bitstream/10400.18/5700/3/INSA_Relatorio_Vacinacao-

antigripal-epocas-2016-2017_2017-2018.pdf

12.   Demicheli V, Jefferson T, C DP, Ferroni E, Thorning S, Re T, et al. Vaccines for preventing

influenza in the elderly (Review). 2018;(2).

13.   Influenza - European Commission [Internet]. [cited 2018 Mar 15]. Available from:

https://ec.europa.eu/health/vaccination/influenza_en

14.   Types of seasonal influenza vaccine |WHO. 2018 Mar 22 [cited 2018 Mar 22]; Available from:

http://www.euro.who.int/en/health-topics/communicable-diseases/influenza/vaccination/types-of-

seasonal-influenza-vaccine

15.   Krahn M, Kuntz KM, Meltzer DO, Owens DK, Russell LB, Siegel JE, et al. Recommendations for

Conduct, Methodological Practices, and Reporting of Cost-effectiveness Analyses: Second Panel

on Cost-Effectiveness in Health and Medicine. JAMA. 2016;316(10):1093–103.

16.   Ferreira LN, Ferreira PL, Pereira LN, Oppe M. EQ-5D Portuguese population norms. Qual Life

Res. 2014;23(2):425–30.

17. Hollmann M, Garin O, Galante M, Ferrer M, Dominguez A, Alonso J. Impact of Influenza on

Health-Related Quality of Life among Confirmed (H1N1)2009 Patients. PLoS One. 2013;8(3):1–

10.

18. Greenland S. Sensitivity analysis, Monte Carlo risk analysis, and Bayesian uncertainty assessment.

Risk Anal [Internet]. 2001;21(4):579–83. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/11726013

19. Drummond MF, Sculpher MJ, Claxton K, Stoddart GL, Torrance GW. Methods for the Economic

Evaluation of Health Care Programmes. Fourth. Oxford; 2015.

20. Laires PA, Ejzykowicz F, Hsu T-Y, Ambegaonkar B, Davies G. Cost–effectiveness of adding

ezetimibe to atorvastatin vs switching to rosuvastatin therapy in Portugal. J Med Econ [Internet].

2015;18(8):565–72. Available from:

http://www.tandfonline.com/doi/full/10.3111/13696998.2015.1031794

21. Cameron D, Ubels J, Norström F. On what basis are medical cost-effectiveness thresholds set?

Clashing opinions and an absence of data: a systematic review. Glob Health Action [Internet].

2018;11(1). Available from: https://doi.org/10.1080/16549716.2018.1447828

22. Marques A, Lourenço Ó, Ortsäter G, Borgström F, Kanis JA, da Silva JAP. Cost-Effectiveness of

Intervention Thresholds for the Treatment of Osteoporosis Based on FRAX®in Portugal. Calcif

Tissue Int. 2016;99(2):131–41.

23. Areia M, Spaander MCW, Kuipers EJ, Dinis-Ribeiro M. Endoscopic screening for gastric cancer:

A cost-utility analysis for countries with an intermediate gastric cancer risk. United Eur

Gastroenterol J. 2018;6(2):192–202.

24. World Development Indicators | DataBank [Internet]. [cited 2018 Sep 15]. Available from:

http://databank.worldbank.org/data/reports.aspx?source=2&country=PRT

25.   Briggs A, Sculpher M, Claxton K. Decision Modelling for Health Economic Evaluation. Oxford:

344   Oxford University Press; 2006. 445 p.

345   26.   Portal do Instituto Nacional de Estatística [Internet]. [cited 2018 May 4]. Available from:

346   https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&indOcorrCod=0000256&co

347   ntexto=bd&selTab=tab2&xlang=en

348   27.   Guiomar R, Conde P, Cristóvão P, Pechirra P, Nunes B. Programa Nacional de Vigilância da Gripe

349   - Relatório da Época 2012/2013. Inst Nac Saúde Doutor Ricardo Jorge, IP. 2013;

350   28.   Guiomar R, Conde P, Cristóvão P, Pechirra P, Nunes B. Programa Nacional de Vigilância da Gripe

351   - Relatório da Época 2013/2014. Inst Nac Saúde Doutor Ricardo Jorge, IP. 2015;

352   29.   Guiomar R, Costa I, Cristóvão P, Pechirra P, Rodrigues AP, Nunes B. Programa Nacional de

353   Vigilância da Gripe - Relatório da época 2014/2015. Inst Nac Saúde Doutor Ricardo Jorge, IP.

354   2015;

355   30.   Guiomar R, Pechirra P, Cristóvão P, Costa I, Conde P, Rodrigues AP, et al. Programa Nacional de

356   Vigilância da Gripe - Relatório da época 2016/2017. Inst Nac Saúde Doutor Ricardo Jorge, IP.

357   2017;

358   31.   Guiomar R, Pechirra P, Rodrigues AP, Silva SP, Machado A, Nunes B. Programa Nacional de

359   Vigilância da Gripe - Relatório da época 2015/2016. Instituto Nacional de Saúde Doutor Ricardo

360   Jorge, editor. Inst Nac Saúde Doutor Ricardo Jorge, IP [Internet]. 2016; Available from:

361   http://repositorio.insa.pt//handle/10400.18/4044

362   32.   Gonçalves P, Pechirra P, Conde P, Guiomar R, Nunes B. Programa Nacional de Vigilância da

363   Gripe - Relatório da Época 2010/2011. Inst Nac Saúde Doutor Ricardo Jorge, IP. 2013;

364   33.   Gonçalves P, Pechirra P, Conde P, João I, Guiomar R, Nunes B. Programa Nacional de Vigilância

365   da Gripe - Relatório da Época 2011/2012. Inst Nac Saúde Doutor Ricardo Jorge, IP. 2013;

366   34.   Petri E, Ruiz-aragón J. A Comparison of Cost Effectiveness of Seasonal Influenza vaccines in

367   Spain. In: IVW conference, Albufeira, 6th -9th October 2015. 2015. p. 7015.

368   35.   Instituto Nacional de Saúde Doutor Ricardo Jorge. Boletim de Vigilância Epidemiológica da Gripe

369   - semana 20. 2018.

370   36.   Inverno e Saúde - Evolução Semanal [Internet]. [cited 2018 May 9]. Available from:

371   http://saudesazonal.min-saude.pt/#consultaSindromeGripal/149/757

372   37.   Administração Central do Sistema de Saúde I.P. Base de dados Central de Morbilidade Hospitalar.

373   38.   Portal do Instituto Nacional de Estatística [Internet]. [cited 2018 May 4]. Available from:

374   https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&indOcorrCod=0008180&co

375   ntexto=bd&selTab=tab2

376   39.   Coelho PP, Morgado C, Albuquerque D De, Rodrigues M. Diário da República 1.ª série, Nº153 -

377   Portaria 234/2015 de 7 de agosto - Anexo-II. 2015.

378   40.   BASE:contratos públicos online [Internet]. [cited 2018 May 4]. Available from:

379   http://www.base.gov.pt/Base/pt/Homepage

380   41.   Direção Geral de Saúde. Orientação 007/2015 Terapêutica e quimioprofilaxia da gripe sazonal.

381   2014;1–7.

382   42.   Serviços Partilhados do Ministério da Saúde. Contrato Fornecimento de V13 - Vacina Contra a

383   Gripe Inativada [UDOSE] à Administração Regional de Saúde do Alentejo. 2016.

384   43.   BASE:contratos públicos online [Internet]. [cited 2020 Jan 27]. Available from:

385   http://www.base.gov.pt/Base/pt/Pesquisa/Contrato?a=5864096

386   44.   Portal do INE - Índice de preços no consumidor (Taxa de variação média dos últimos 12 meses -

387   Base 2012 - %) por Localização geográfica e Agregados especiais; Mensal [Internet]. [cited 2020

Jan 13]. Available from:

https://www.ine.pt/xportal/xmain?xpid=INE&xpgid=ine_indicadores&indOcorrCod=0002390&co

ntexto=pi&selTab=tab0

45.   Boer PT, Crépey P, Pitman RJ, Macabeo B, Chit A, Postma MJ. Cost-Effectiveness of

Quadrivalent versus Trivalent Influenza Vaccine in the United States. Value Heal [Internet].

2016;19(8):964–75. Available from: http://dx.doi.org/10.1016/j.jval.2016.05.012

46.   Rondy M, Kissling E, Gherasim A, Pebody R, Trebbien R, Pozo F, et al. Interim 2017 / 18

influenza seasonal vaccine effectiveness : combined results from five European studies. Euro

Surveill. 2018;23(9):1–12.

47.   Rondy M, Gherasim A, Casado I, Launay O, Rizzo C, Pitigoi D, et al. Low 2016/17 season vaccine

effectiveness against hospitalised influenza A(H3N2) among elderly: awareness warranted for

2017/18 season. Eurosurveillance. 2017 Oct 12;22(41).

48.   Rondy M, Larrauri A, Casado I, Alfonsi V, Pitigoi D, Launay O, et al. 2015 / 16 seasonal vaccine

effectiveness against hospitalisation with influenza A (H1N1) pdm09 and B among elderly people

in Europe : results from the I-MOVE + project. Euro Surveill. 2015;

49.   Nunes B, Machado A, Pechirra P, Falcão I, Gonçalves P, Conde P, et al. Efetividade da vacina

antigripal na {é}poca 2010-2011 em Portugal: resultados do projeto EuroEVA. Rev Port Med

Geral E Fam. 2012;28(4):271–84.

50.   Kissling E, Valenciano M, Pozo F, Vilcu A-M, Reuss A, Rizzo C, et al. 2015/16 I-MOVE/I-

MOVE+ multicentre case control study in Europe: moderate vaccine effectiveness estimates

against influenza A(H1N1)pdm09 and low estimates against lineage mismatched influenza B

among children. Influenza Other Respi Viruses. 2017 Nov 10;

51.   You JHS, Ming WK, Chan PKS. Cost-effectiveness analysis of quadrivalent influenza vaccine

versus trivalent influenza vaccine for elderly in Hong Kong. BMC Infect Dis. 2014;14(1):1–7.

52.   You JHS, Ming WK, Chan PKS. Cost-effectiveness of quadrivalent influenza vaccine in Hong

Kong - A decision analysis. Hum Vaccines Immunother. 2015;11(3):564–71.

53.   Chit A, Roiz J, Aballea S. An assessment of the expected cost-effectiveness of quadrivalent

influenza vaccines in Ontario, Canada using a static model. PLoS One [Internet]. 2015;10(7):1–15.

Available from: http://dx.doi.org/10.1371/journal.pone.0133606

54.   Chit A, Roiz J, Briquet B, Greenberg DP. Expected cost effectiveness of high-dose trivalent

influenza vaccine in US seniors. Vaccine [Internet]. 2015;33(5):734–41. Available from:

http://dx.doi.org/10.1016/j.vaccine.2014.10.079

55.   Van Bellinghen LA, Meier G, Van Vlaenderen I. The potential cost-effectiveness of quadrivalent

versus trivalent influenza vaccine in elderly people and clinical risk groups in the UK: A lifetime

multi-cohort model. PLoS One. 2014;9(6).

56.   Dolk C, Eichner M, Welte R, Anastassopoulou A, Van Bellinghen LA, Poulsen Nautrup B, et al.

Cost-Utility of Quadrivalent Versus Trivalent Influenza Vaccine in Germany, Using an Individual-

Based Dynamic Transmission Model. Pharmacoeconomics. 2016;34(12):1299–308.

57.   García A, Ortiz de Lejarazu R, Reina J, Callejo D, Cuervo J, Morano Larragueta R. Cost–

effectiveness analysis of quadrivalent influenza vaccine in Spain. Hum Vaccines Immunother.

2016;12(9):2269–77.

58.   Meier G, Gregg M, Poulsen Nautrup B. Cost-effectiveness analysis of quadrivalent influenza

vaccination in at-risk adults and the elderly: an updated analysis in the UK. J Med Econ [Internet].

2015;18(9):746–61. Available from:

http://www.tandfonline.com/doi/full/10.3111/13696998.2015.1044456

59.   Mullikin M, Tan L, Jansen JP, Van Ranst M, Farkas N, Petri E. A Novel Dynamic Model for

434   Health Economic Analysis of Influenza Vaccination in the Elderly. Infect Dis Ther.

435   2015;4(4):459–87.

436 60. Capri S, Barbieri M, de Waure C, Boccalini S, Panatto D. Cost-effectiveness analysis of different

437   seasonal influenza vaccines in the elderly Italian population. Hum Vaccin Immunother.

438   2018;14(6):1331–41.

439 61. MC W, G T, Mcguire A. QALYs : The Basics. Value Heal. 2009;12(1):S5–9.

440 62. Uva MS, Kislaya I, Roquette R, Rodrigues AP, Machado A. Vacinação antigripal da população

441   portuguesa na época 2015/2016. Estudo na amostra ECOS. 2016; Available from:

442   http://www.onsa.pt/conteu/estud_vacina-anti-gripal_onsa.pdf

445 **Supporting Information**

446 **S1 Appendix.**

447 **S2 Appendix. Model input parameters.**

448 **S1 Table. Proportion of B lineage influenza virus not included in the seasonal trivalent**

449 **vaccines from 2010/11 to 2017/18 influenza seasons.**

450 **S2 Table. Description of the variables that take part of S1 Fig.**

451 **S3 Table. Input parameters.** Base case value; SD; parameters of the probability distribution.

452 **S4 Table. Probabilistic sensitivity analysis.** Assessing the 2015/16 influenza season-related

453 uncertainty.

S1 Fig. Probability, cost and QALY calculations by decision tree pathway for individuals

aged ≥65 years old who received TIV.

S2 Fig. One-way sensitivity analysis of TIV effectiveness. Variation between 34% and 74%

with increments of 0.005.

S3 Fig. One-way sensitivity analysis of QIV effectiveness. Variation between 58% (TIV

effectiveness) and 100% with increments of 0.005.

S4 Fig. One-way sensitivity analysis of vaccination coverage rate. Variation between 0% and

100% with increments of 0.05.

S5 Fig. Probabilistic sensitivity analysis for TIV and QIV. Scatter plot based on the 2015/16

influenza season.

Figure 3

Figure 4

Figure 1

Figure 2