1

B1 SINE-binding ZFP266 impedes reprogramming through suppression of chromatin opening by

pioneering factors

Daniel F Kaemena[1], Masahito Yoshihara[2], James Ashmore[1], Meryam Beniazza[1], Suling Zhao[1], Mårten

Bertenstam[3], Victor Olariu[3], Shintaro Katayama[2,4,5], Keisuke Okita[6], Simon Tomlinson[1], Kosuke Yusa[7,8] &

Keisuke Kaji[1]

[1]MRC Centre for Regenerative Medicine, University of Edinburgh, Edinburgh BioQuarter, 5 Little France

Drive, Edinburgh, EH16 4UU, Scotland, UK. [2]Department of Biosciences and Nutrition, Karolinska Institutet,

141 83 Stockholm, Sweden. [3]Computational Biology and Biological Physics, Lund University, Lund, Sweden.

[4]Research Programs Unit, Stem Cells and Metabolism Research Program (STEMM), University of Helsinki,

00014 Helsinki, Finland. [5]Folkhälsan Research Center, 00014 Helsinki, Finland. [6]Center for iPS Cell Research

and Application, Kyoto University, Kyoto, Japan. [7]Stem Cell Genetics, Wellcome Sanger Institute, Hinxton,

Cambridge, CB10 1SA, UK. [8]Stem Cell Genetics, Institute for Frontier Life and Medical Sciences, Kyoto

University, Kyoto, Japan.

20 Correspondence: keisuke.Kaji@ed.ac.uk

Abstract

Successful generation of induced pluripotent stem cells (iPSCs) via the overexpression of *Oct4 (Pou5f1)*, *Sox2*, *Klf4* and *c-Myc* (OSKM) highlights the power of transcription factor (TF)-mediated cellular conversions. Nevertheless, iPSC reprogramming is inherently inefficient and understanding the molecular mechanisms underlying this inefficiency holds the key to control cellular identity successfully. Here, we report 16 novel reprogramming roadblock genes identified by CRISPR/Cas9-mediated genome-wide knockout (KO) screening. Of these, disruption of KRAB zinc finger protein (KRAB-ZFP) *Zfp266* strongly and consistently enhanced iPSC generation in several iPSC reprogramming settings, emerging as the most robust roadblock. Further analyses revealed that ZFP266 bound Short Interspersed Nuclear Elements (SINEs) adjacent to OSK binding sites and impedes chromatin opening. This work serves as a resource for better understanding reprogramming mechanisms and proposes SINEs as a critical genetic element that regulates chromatin accessibility at enhancers for efficient pluripotency induction.

Introduction

The reprogramming of somatic cells into iPSCs via the overexpression of *Oct4 (Pou5f1)*, *Sox2*, *Klf4* and *c-Myc* (OSKM) has provided an important tool for medical research and cell therapies[1]. Equally importantly, the generation of fully functional iPSCs indistinguishable from ESCs from somatic cells has clearly demonstrated that cellular identity can be completely converted from one to another by simple overexpression of master transcription factors. This has provided a model system to understand how to control cellular identity. Inhibition of *Trp53* and *Cdkn1a* (*p21*) revealed OSKM-induced apoptosis and senescence as a major roadblock of iPSC generation[2–8]. Knockdown of *Dot1l* and *Suv39h1* demonstrated H3K79me and H3K9me3 as critical epigenetic modifications that impede this cell conversion[9,10]. Thus, identifying genes that act against successful reprogramming offers the foundation to understand critical molecular mechanisms involved in pluripotency induction.

Transposable elements (TEs), which consist of about 40% of mouse and human genome, take part in gene expression regulation as cis-regulatory elements in the genome or non-coding RNAs[11], and importance of

many types of TEs in pluripotent cells has been described[12–15]. Knockdown of the long interspersed element

1 (LINE1) inhibits mouse ESC self-renewal and induces transition to a 2C state[12]. KLF4 activates transcription

of HERVH during reprogramming, and whose down-regulation is critical for exit from the pluripotent state

of human iPSCs[13]. KLF4 is also responsible for opening thousands of genomic loci in naïve human ESCs via

binding to HERVH, HERVK, and non-LTR retrotransposon SVA subgroups[14]. Similarly, chromatin accessibility

of SINEs, which constitute ~25% of TEs, is particularly high in mouse pre-implantation embryos and ESCs[15],

however the functional importance of this has not yet been addressed. Krüppel-associated box (KRAB) zinc-

finger proteins (ZFPs) are the largest transcription factor family in mouse and human genomes with over

300 members, which have evolved to supress rapidly mutating TEs, and about two thirds of KRAB-ZFPs bind

to TEs[16]. Nevertheless, only a small number of KRAB-ZFPs that bind to SINEs is known[17].

Here, we report an unbiased genome-wide CRISPR KO screen with a library containing 90,230 sgRNAs

targeting 18,424 protein coding genes. This screen identified 16 genes as novel reprogramming roadblocks,

as well as 8 previously reported inhibitory genes. Of those, KO of the KRAB-ZFP gene *Zfp266* has

accelerated the kinetics of reprogramming and improved efficiency of iPSC generation 4- to 10-fold in

various reprogramming contexts. We revealed that ZFP266 binds to SINEs adjacent to OSKM binding sites

during reprogramming and impedes chromatin opening. Furthermore, replacing its KRAB co-suppressor

interacting domain with a co-activator interacting domain converted ZFP266 from a reprogramming

inhibitor to a reprogramming facilitator. This indicated that SINEs next to OSKM binding sites during

reprogramming were critical genetic elements that modulate efficiency of OSKM-mediated iPSC generation.

This work serves as a resource for better understanding of reprogramming mechanisms and highlights

SINEs as novel transposable elements (TEs) involved in pluripotency induction.

A CRISPR/Cas9-mediated genome-wide KO screen identified 16 novel reprogramming roadblock genes

We have previously generated a Cas9 expressing mouse ES cell line, named Cas9 TNG MKOS, with a *Nanog-*

GFP-ires-Puro reporter and a doxycycline-inducible *MKOS-ires-mOrange* polycistronic reprogramming

cassette in the *Sp3* locus (Supplemental Figure S1A and S1B)[18,19]. Efficient KO by lentiviral sgRNA delivery in

both Cas9 TNG MKOS ESCs and mouse embryonic fibroblasts (MEFs), generated through morula

aggregation of these ESCs, was confirmed using sgRNA against ICAM1 and CD44, respectively, resulting in

>80% loss of protein within 72 hours (Supplemental Figure S1C and S1D). Reprogramming of Cas9 TNG

MKOS MEFs following sgRNAs transduction against known roadblock genes, *Trp53* and *Rb1*[3,4,6–8,20], and

essential genes, *Pou5f1* and *Kdm6a*[21], reproduced the expected reprogramming enhancement and

reduction phenotypes, respectively (Supplemental Figure S1E-S1G), confirming that the CRISPR-based KO

system as a powerful tool to investigate gene function in reprogramming. We then performed genome-

wide KO screening using a previously published lentiviral sgRNA library[22], with an optimized reprogramming

condition consisting of 8 days of reprogramming factor expression followed by 8 days of puromycin

selection for *Nanog*-GFP[+] iPSCs (Figure 1A and Supplemental Figure S1H-S1K). This condition resulted in an

average coverage of ~170 MEFs/sgRNA/screening replicate. Genomic DNA from flow-sorted *Nanog*-GFP[+]

iPSCs was then collected in triplicate, and integrated sgRNAs were Illumina-sequenced after PCR

amplification alongside the original sgRNA plasmid library.

The normalized read counts of all sgRNAs and the results of the statistical tests to identify novel roadblock

genes with a software, MAGeCK[23], are available in Supplemental Tables S1, S2 and at https://kaji-crispr-

screen-updated.netlify.app. Using a false discovery rate (FDR) <0.1 as a cut-off, we identified 24 genes as

reprogramming roadblocks (Figure 1B). This included 16 novel candidates as well as 5 previously

characterized genes: *Trp53*, *Cdkn1a*, *Dotl1* and AP-1 transcription factor components *Jun* and *Fosl2*[24,25], and

3 genes previously uncharacterized yet identified in other screens, *Men1, Gtf2i and Cdk13*[26,27], signifying

the robustness of our screen (Figure 1C and 1D). When the top 3 ranked sgRNA for each gene were

individually tested, transduction of *Trp53* and *Cdkn1a* sgRNAs produced the largest increase in *Nanog*-GFP[+]

colony numbers (Figure 1E and 1F), although they also significantly increased the number of partially

reprogrammed colonies (Figure 1F and Supplemental Figure S2A). Transduction of sgRNAs targeting all

other genes, except for the previously reported roadblock *Cdk13,* enhanced *Nanog*-GFP[+] colony formation

between 2- and 6-fold (Figure 1E and Supplemental Figure S2A), verifying the inhibitory effects of the novel

reprogramming roadblock genes. Expression of the validated 23 roadblock genes during reprogramming did

not follow any common particular pattern and many of them exhibited consistently low expression,

compared to the common housekeeping genes or the reprogramming and pluripotency marker genes[28]

(Supplemental Figure S2B). This highlights the advantage of functional screening to identify their inhibitory

effects over expression profiling.

Of the 16 novel roadblock genes, KO of 8 resulted in a >4-fold enhancement similar to or better than

previously reported roadblocks (Figure 1E). We therefore further characterise these 8 novel reprogramming

roadblocks; *Fam122a, Zfp266, Bcorl1, Usp28, Usp34, Zc3h10, Scaf8* and *Spop* (Figure 1F, blue), alongside

the previously reported roadblocks *Trp53, Cdkn1a, Men1, Dot1l* and *Gtf2i* (Figure 1F, green), as 13 top

roadblocks.

***Zfp266* KO consistently enhances and accelerates the attainment of pluripotency**

Reprogramming roadblock function is influenced by multiple elements such as the stoichiometry or

expression levels of OSKM, culture conditions, and starting cell types[18]. Thus, we examined the KO effects

of our 13 top roadblocks in different reprogramming contexts. We first performed *piggyBac* transposon-

based reprogramming with a *MKOS* or STEMCCA (*OKSM*) reprogramming cassette[29,30] (Supplemental Figure

S3A). The STEMCCA cassette expresses lower levels of KLF4 protein due to an N-terminal truncation

followed by a 2A peptide[31], resulting in inefficient mesenchymal-epithelial transition (MET) and a higher

proportion of partially reprogrammed cells[18,32]. *piggyBac* delivery of the MKOS cassette together with each

sgRNA against all the 13 roadblocks enhanced reprogramming of *Cas9 Nanog*-GFP MEFs as seen with Cas9

TNG MKOS MEFs before (Figure 2A and 2B), despite a markedly lower KO efficiency with the *piggyBac*

system compared to lentiviral sgRNA delivery (Supplemental Figure S3B). In STEMCCA-mediated *piggyBac*

reprogramming, KO of all, except *Cdkn1a*, *Fam122a* and *Zc3h10,* increased the number of *Nanog*-GFP$^+$

colonies. Particularly, the KO of *Zfp266* increased numbers of *Nanog*-GFP$^+$ colonies ~10-fold with almost all

colonies expressing *Nanog*-GFP (Figure 2C and 2D), while *Cdkn1a*, *Fam122a* knockout drastically increased

only *Nanog*-GFP$^-$ colony numbers (Figure 2C and 2D). When *piggyBac MKOS*+sgRNA vectors were used to

reprogram *Cas9* expressing neural stem cells (NSCs), sgRNAs against *Men1, Fam122a, Zfp266* and *Usp34*

increased reprogramming efficiency, with KO of *Zfp266* again leading to the greatest enhancement in

NANOG[+] colony formation (~5-fold) (Figure 2E and 2F). When we explored reprogramming kinetics using

Cas9 TNG MKOS MEFs by assessing expression changes of reprogramming markers, CD44, ICAM1 and

Nanog-GFP[33], KO of 5 genes, *Men1*, *Gtf2i, Dot1l, Zfp266* and *Zc3h10,* demonstrated the accelerated

reprogramming (Figure 2G, 2H, and Supplementary Figure 3C). In summary, KO of *Zfp266* exhibited the

most context-independent and robust reprogramming enhancement amongst all roadblock genes we

identified. We therefore investigated further how *Zfp266* impedes the reprogramming process.

ZFP266 impedes activation of pluripotency genes via its KRAB domain

Zfp266 encodes a KRAB-ZF protein with a singular KRAB-A module in the N-terminus and putative DNA

binding domain with 12x C2H2 type zinc finger array in the C-terminus (Figure 3A). KRAB domains are

known to interact with co-suppressor KAP-1/TRIM28, a scaffold protein that can recruit epigenetic

modifiers and promote the formation of heterochromatin and transcriptional repression[34], suggesting that

ZFP266 acts as a suppressor. Consistent with the reprogramming enhancement by *Zfp266* KO,

overexpression of exogenous *Zfp266* completely hindered reprogramming (Figure 3B). Exogenous

overexpression of *Zfp266* mutants either lacking the entire KRAB domain or containing point mutations

which disrupt the interaction with KAP-1/TRIM28[35,36] could not abolish *Zfp266* sgRNA-mediated

reprogramming enhancement even though the *Zfp266* mutants were resistant to the sgRNA (Figure 3C and

3D). This clearly demonstrates that Zfp266 inhibits reprogramming through its KRAB-domain.

To assess further the function of zfp266, we examined the gene expression changes associated with

depletion of *Zfp266* (Supplemental Tables S3). RNA-seq of MEFs 4 days after *Zfp266* sgRNA transduction

revealed only 4 differentially expressed genes (DEGs) (FDR<0.05, log2FC<|1|), demonstrating that loss of

ZFP266 alone was not sufficient to cause drastic gene expression changes in MEFs (Figure 3E, Supplemental

Tables S4). In contrast, the number of DEGs between *Zfp266* KO and wild-type cells rapidly increased during

reprogramming from 24 at day 3 to 222 at day 5, and to 1761 at day 7 (Figure 3E, Supplemental Tables S5-

S7). The majority of DEGs at day 3 and day 5 were upregulated (75% and 67% respectively, Figure 3E),

consistent with the predicted role for *Zfp266* as a transcriptional suppressor. Enhanced up-regulation of

pluripotency-associated genes *Piwil2* and *Dppa5a* at day 3, *Nanog, Esrrb, Dppa2,* and *Tcl1, etc* at day 5 was

already detected in *Zfp266* KO (Figure 3E). Furthermore, over 60% (11/18), 80% (120/149), 69% (575/835)

of up-regulated DEGs in *Zfp266* KO cells at day 3, 5, 7 of reprogramming were genes more highly expressed

in ESCs compared to MEFs (FDR<0.05, log2FC<|1|) (Figure 3E, green, Supplemental Tables S8). Gene

ontology (GO) enrichment analysis identified 'stem cell population maintenance' in day 5 and 'response to

leukaemia inhibitory factor' in day 7 up-regulated DEGs as one of the most enriched terms (Figure 3F),

while downregulated DEGs at day 7 were significantly enriched in developmental and differentiation terms

(Figure 3F). Principal component analysis (PCA) also indicated that gene expression changes that have

occurred in *Zfp266* KO reprogramming at day 5 and 7 reflected an accelerated transition towards a

pluripotent state (Figure 3G). Taken together, these data indicate that *Zfp266* KO enhances and accelerates

reprogramming by permitting a more efficient activation of pluripotency genes by OSKM.

***Zfp266* KO in MEFs results in chromatin opening at the SINE-containing Zfp266 binding sites**

One possible mechanism by which ZFP266 impedes reprogramming is that ZFP266 binds and locks the

pluripotency loci in MEFs and other differentiated cells. To investigate this possibility, we mapped ZFP266

binding sites in MEFs using DamID-seq, which does not required specific antibodies[37,38]. This identified

15,119 unique ZFP266 binding sites (Figure 4A), predominantly situated in introns or intergenic regions

(Supplementary Figure 4A, Supplemental Tables S9). These ZFP266 binding sites have low chromatin

accessibility as measured by ATAC-seq in MEFs, 72 hours after reprogramming as well as in iPSCs[39] (Figure

4A). ZFP266 binding sites were predominantly enriched for somatic AP-1 TF motifs (Figure 4B), and little

OSKM binding was observed at the same loci in 48hr reprogramming or ESC ChIP-Seq datasets[40,41]

(Supplementary Figure 4B and 4C). This disputed the idea that ZFP266 functions as a suppressor at the

pluripotency-related gene loci in differentiated cells. Thus, we investigated whether any TE families were

enriched in ZFP266 binding sites, since many KRAB-ZFPs are known to bind and supress transcription of

TEs[16]. In line with this, we found the 15,119 ZFP266 DamID-seq peaks in MEFs were highly enriched in SINEs

with about two thirds (10,523) overlapping with SINEs (Figure 4C and 4D). Of SINE subfamilies, B1 SINEs in

particular exhibited both the most significant enrichment and the most abundant overlap with ZFP266

binding sites (Figure 4C and 4D). Furthermore, de novo motif analysis of ZFP266 binding sites identified 3

long de novo motifs which all corresponded to parts of the B1 SINE consensus sequence (Figure 4E),

suggesting ZFP266 might bind B1 SINEs.

We next examined how depletion of ZFP266 might affect chromatin accessibility. To this end, we

performed ATAC-Seq of *Zfp266* KO MEFs and identified 479 more open regions (MORs) compared to WT

MEFs, while only one locus was found to be a more closed (Figure 4F and Supplemental Tables S10).

Considering the predicted ZFP266 function as a repressor, we next examined whether ZFP266 binds to the

MORs in wild-type MEFs. Although only about 25% (123/479) of *Zfp266* KO MEF MORs overlapped with

ZFP266 DamID-seq peaks, non-overlapped MORs also had increased DamID-seq signals albeit at a lower

level (Figure 4G), unlike randomly selected control regions with similar chromatin accessibility

(Supplementary Figure 4D). This suggests that more than 25% of MORs are likely bound by ZFP266, while

they were not identified as a 'peak' with our DamID-seq due to the cut-off criteria and/or technical

limitations. In fact, similar to ZFP266 DamID-seq peaks in MEFs, *Zfp266* KO MEF MORs were mainly located

in intergenic regions and introns (Supplementary Figure 4E), and enriched in AP-1 TF motifs, with 87% of all

Zfp266 KO MEF MORs containing at least one AP-1 TF motif (Figure 4H). *De novo* motif discovery analysis

also identified 5 motifs that overlap with the B1 SINE consensus sequence (Figure 4I), consistent with the

fact that SINE was the most enriched repetitive element (Figure 4J) and 92% (1418/1522) of *Zfp266* KO MEF

MORs had at least one SINE (Figure 4K). Overall, these data suggest that ZFP266 binds to SINEs in MEFs to

keep target loci closed, and removal of ZFP266 allows TFs that binds nearby, like AP-1 TFs, to facilitate

chromatin opening (Figure 4L). However, ZFP266 does neither bind to nor regulate pluripotency gene loci in

MEFs.

Zfp266 KO in reprogramming results in chromatin opening at SINE-containing OSK binding sites

In order to address why *Zfp266* KO results in significant reprogramming enhancement, we performed

ATAC-seq 72 hours after reprogramming with and without *Zfp266* KO. Similar to the KO effects in MEF,

Zfp266 KO reprogramming cells exhibited 1522 MORs, majority of which were situated in intergenic regions

and introns (Supplementary Figure 5A), and only 86 more closed regions compared to wild-type cells

(Figure 5A, Supplemental Tables S11). They were also significantly enriched in SINEs, particularly B1 SINE

(Figure 5B), with >90% (1459/1522) of MORs containing at least one SINE (Figure 5C). *De novo* motif

discovery analysis also identified motifs that correspond to the B1 SINE consensus sequence as the most

significant motifs (Figure 5D). However, these loci hardly overlapped with *Zfp266* KO MEF MORs (Figure 5E),

suggesting a context dependency for which loci become more open in the absence of ZFP266. The overlap

with MEF ZFP266 DamID-seq peaks was also minimal, with only ~10% of MORs overlapping (Supplemental

Figure S5B), and non-overlapped MORs did not have increased DamID-seq signals compared to the control

regions with a similar chromatin accessibility (Supplemental Figure S5B). This indicated that upon OSKM

expression ZFP266 changes their binding sites, in which ZFP266 regulates chromatin accessibility. TF motif

enrichment analysis revealed that KLF, SOX and the OCT4::SOX2 motifs were highly enriched in *Zfp266* KO

reprogramming MORs, with particularly KLF family (KLF1, KLF5, KLF4, KLF9, KLF12) motifs identified in >90%

of these MORs, while AP-1 TF motifs were also enriched (Figure 5F). Next, we classified *Zfp266* KO

reprogramming MORs into two groups using *K*-means clustering (Figure 5G). The first cluster (121 regions)

are open in both wild-type and *Zfp266* KO MEFs, and then become more closed upon reprogramming,

while *Zfp266* KO cells are more resistant to this closing (cluster 1, Figure 5G). The second cluster contained

the majority (1401) of *Zfp266* KO reprogramming MORs, which are closed in both wild-type and *Zfp266* KO

MEFs, and become open following reprogramming, an effect which is enhanced when *Zfp266* is knocked

out (cluster 2, Figure 5G). Cluster 2 indicates that removal of ZFP266 facilitates OSKM-mediated chromatin

opening. In fact, we observed OSK binding in cluster 2 reprogramming MORs with particularly strong KLF4

signal using published reprogramming 48-hour ChIP-Seq datasets[41] (Figure 5H). Similar OSK binding was

observed in an ESC ChIP-Seq dataset albeit to a lesser extent[40] (Supplemental Figure S5C), and about one

third of the cluster 2 loci have an open chromatin state in iPSCs (Supplemental Figure S5D), suggesting that

some of the MORs are OSK targets in pluripotent cells. Interestingly, while both OSK binding and motifs

were enriched close to the MOR peak summit (within 70 bp) (Figure 5H and 5I), SINEs were depleted from

summits and were instead enriched immediately upstream or downstream (~70 bp away from the summit),

therefore being located immediately adjacent to OSK motifs and binding sites (Figure 5I and 5J).

Furthermore, B1 SINEs within the MORs were likely orientated such that the 5' head sequence was

positioned inwards facing towards the peak summit (Figure 5I and 5J). This positional and directional bias

within the MOR was exclusive to B1 SINE subfamilies as B2 SINEs exhibited no such bias (Supplemental

Figure S5E). Based on these data together with a report that somatic TFs' binding sites drastically change

upon OSKM expression[40], we speculated that ZFP266 was relocated to B1 SINEs next to OSK binding sites

during reprogramming, where it then impeded chromatin opening.

Facilitating chromatin opening at ZFP266 targeted SINEs enhances reprogramming

In order to validate binding of ZFP266 to SINEs, we generated an activator version of *Zfp266* with the KRAB

domain replaced by a flexible linker and three transactivating domains VP64, p65 and Rta (VPR)[42] (Figure

6A), and performed luciferase reporter assays using HEK293 cells. Clear luciferase expression was observed

when the VPR-*Zfp266*, but not BFP or VPR only controls, was co-transfected with a reporter plasmid

containing the B1 SINE consensus sequence upstream of a SV40 minimal promoter (Figure 6B). Co-

expression of wild-type *Zfp266* alongside VPR-*Zfp266* attenuated this reporter expression (Figure 6C),

confirming ZFP266 specifically binds B1 SINEs. We next examined whether VPR-ZFP266 can bind to *Zfp266*

KO reprogramming MORs using the *Luciferase* assay. We selected SINE containing MORs in three genes,

B3gnt3, *Piwil2* and *Snx20*, whose transient up-regulation during reprogramming was significantly

augmented by *Zfp266* KO (Figure 6D). These loci are closed in MEFs, open up more in *Zfp266* KO cells upon

reprogramming, and are bound by KLF4 at 48 hours of reprogramming (Figure 6E). Each MOR was cloned

upstream of a minimal SV40 promoter in both a forward and reverse orientation in a luciferase reporter

vector. We found that VPR-*Zfp266* could also enhance luciferase expression from these MOR reporter

vectors (Figure 6F), while co-expression of wild-type *Zfp266* would then ablate it (Figure 6G). Deleting B1

SINE sequences from the *B3gnt3*, *Piwil2* and *Snx20* MORs diminished VPR-*Zfp266*'s ability to enhance

luciferase expression, confirming that ZFP266 binds to reprogramming MORs specifically via B1 SINE

sequences (Figure 6H-6J). We also confirmed that OSKM expression enhanced luciferase expression from

the *B3gnt3* and Snx20 MOR containing reporter vectors in MEFs (Figure 6K), indicating the MORs have

OSKM-dependent enhancer activity. In addition, removing B1 SINE sequences from the *Snx20* MOR led to

an increase in *Luciferase* expression following OSKM induction (Figure 6L), demonstrating B1 SINEs function

to repress reprogramming factor-mediated transactivation presumably via endogenous ZFP266 binding.

Finally, overexpression of the VPR-*Zfp266* together with OSKM lead to accelerated and enhanced

reprogramming with a robust appearance of *Nanog*-GFP[+] colonies by day 9 (Figure 6M and 6N). Taken

together, we propose a model where 1) ZFP266 binds to B1 SINEs adjacent to OSK binding sites during

reprogramming and acts to impede chromatin opening, and 2) KO of *Zfp266* (or recruitment of co-

activators to these loci) tips the balance in favour of OSK, allowing them to establish a more open

chromatin state to drive gene activation necessary for successful reprogramming (Figure 6O).

Discussion

Reprogramming towards iPSCs is a conflict between OSKM transcription factors trying to establish a

pluripotent state and somatic factors trying to resist this disruption in cell identity. We have identified 16

novel reprogramming roadblock genes whose depletion facilitates OSKM-mediated pluripotency induction.

One of the most robust roadblock genes, *Zfp266*, is recruited to OSK bound regulatory elements through

the recognition of adjacent B1 SINEs, where it impedes chromatin opening via its KRAB domain. This

probably underlays the accelerated up-regulation of pluripotency gene by OSKM in both *Zfp266* KO and

VPR-*Zfp266* overexpression reprogramming. In fact, many of *Zfp266* KO MORs at 72 hours of

reprogramming have an open chromatin state and bound by OSK in iPSCs/ESCs, and several of them are

associated with pluripotency genes or other genes highly expressed in ESCs, including *Pou5f1, Sall4, Zfp42,*

Klf2, Piwil2, Fbxo15, *Dnmt3l, Tet1/2* (Supplemental Tables S12). These genes were more efficiently up-

regulated in *Zfp266* KO reprogramming compared to the control group (Supplemental Tables S3). Loss of

ZFP266 in MEFs lead to only 4 DEGs, suggesting that ZFP266 may not play a significant role in a static state

but act as a safeguard against drastic changes of cellular states mediated by newly expressed TFs and/or

extracellular ques, such as cytokines. During early embryo development, i.e. 2-8 cell stage, ICM and ESCs,

SINEs, particularly B1 SINEs, are enriched in the open chromatin regions, but not in subsequent

developmental stages[15]. The data from International Mouse Phenotyping Consortium shows that only 6.5%

of pups from *Zfp266* heterozygous intercrosses are homozygous for the KO allele, presenting incomplete

penetrance and possible roles of ZFP266 during development while there might be compensation

mechanisms (https://www.mousephenotype.org/data/genes/MGI:1924769). Further investigation might

reveal roles of ZFP266 in B1 SINE region closing and neighbouring gene regulation during embryo

development. B1 SINEs within *Zfp266* KO reprogramming MORs displayed a distinct positioning bias

enriched at the flanks of MOR summits, and an orientation bias with head towards the summits. A recent

publication identified enrichment of B1 SINEs at the flanks of CD8+ T-cell specific enhancers[43]. Our work

suggests that this positioning bias could be a much more general feature of B1-SINE linked regulatory

elements and provides evidence that SINEs can affect chromatin states via KRAB-ZFPs. The head-to-tail

orientation bias of B1 SINEs is an intriguing novel observation of our work. Stably positioned nucleosomes

are highly enriched in SINE and LINE retrotransposons in human[44]. Thus, SINEs may have a further

architectural or organisational role of chromatin at the regulatory elements, while it could also be

influenced by surrounding DNA sequences and/or other proteins as about 20-25% of B1 SINE had the

reversed tail-to-head orientation towards the MOR summits. KLF4 has been shown to bind primate/human

specific TEs in naive human ESCs and during reprogramming[13,14]. Interestingly, the *de novo* motif identified

in *Zfp266* KO reprogramming MORs, but not Zfp266 KO MEF MORs or ZFP266 DamID-seq peaks, has a full

KLF4 binding motif with a base switch at the 5' end of B1 SINE consensus sequence, in addition to a partial

KLF4 motif identified in nucleosome enriched KLF4 binding sites[45] (Supplementary Figure S5F). While B1

SINEs are restricted to rodents, the enrichment of KLF4 binding in *Zfp266* KO reprogramming MORs with

SINEs indicates conserved function of KLF4 to regulate gene expression via TE containing regions in

reprogramming/pluripotency across species, which is of clear interest for further investigation. It has been

reported that ~2/3 of human KRAB-ZFPs bind to TEs genome-wide, and KRAB-ZFPs supress not only TEs, but

also expression of genes nearby[16]. Our results suggest a possibility that other KRAB-ZFPs could act as

barriers in different TF-mediated cell conversions or differentiation of pluripotent cells to specific cell types,

and therefore elimination of those obstacles might realize more efficient cellular identity changes. Our

CRISPR/Cas9-mediated genome-wide KO screening also identified several other novel genes whose

inhibition of pluripotency induction would have never been predicted from transcriptomic analyses. Further

understanding how these genes hamper OSKM-mediated reprogramming will bring us a better

understanding of how to control cellular identities.

Methods

Cell culture

ESCs and MEFs were cultured in ESC medium and in MEF medium, respectively, as described previously[18].

Reprogramming was performed in reprogramming medium (ESC medium supplemented with 300 ng ml^{-1} of

doxycycline (Sigma) and 10 μg ml^{-1} of L-ascorbic acid or 2-Phospho-L-ascorbic acid trisodium salt (Vitamin C)

(Sigma). NSCs were cultured in NSC complete medium consisting of DMEM/F-12 Media, 1:1 Nutrient

Mixture (Sigma), 1X N2 supplement (Thermo Fisher Scientific), 1X B27 supplement (Thermo Fisher

Scientific), 8 mM glucose (Sigma), 100 U ml^{-1} Pencillin/Streptomycin (ThermoFisher Scientific), 0.001% BSA

(ThermoScientific), 0.05 mM β-mercaptoethanol (Thermo Fisher Scientific), supplemented with 10 ng ml^{-1}

mouse EGF (Peprotech) and 10 ng ml^{-1} human FGF2 (Peprotech).

Plasmids

Plasmids used in this work are summarized in Supplemental Table S13. The plasmids and their sequences

are available upon request.

Generation of Cas9 TNG MKOS ESC line and Cas9 TNG MKOS MEFs

The *Rosa26* targeting vector carrying EF1α-hCas9-ires-neo cassette (Addgene #67987) was electroporated

into TNG MKOS ESC line[18,22]. Correct targeting was confirmed by Southern blotting using KpnI and MscI

digested genome DNA for a 5' and 3' probe, respectively. The 5' and 3' probes were generated from PCR

amplicon using the following primers, 5' forward CAAGTGCTCCATGCTGGAAGGATTG, 5' reverse

TGATTGGGGAGGATCCAGATGGAG, 3' forward GGATTGCACGCAGGTTCTCCG 3' reverse

CGCCGCCAAGCTCTTCAGCAA and genome DNA (for 5' probe) or the targeting vector (for 3' probe) as a

template. Cas9 TNG MKOS MEFs were isolated from E12.5 chimeric embryos generated via morula

aggregation and the proportion of transgenic MEFs from each embryo was assessed measuring % of

mOrange$^+$ cells after exposing 1/10th of the dissociated cells to Dox for 2 days as describe previously[18].

sgRNA screen

The sgRNA library (Addgene #67988) was prepared as describe before[22]. 9×10^6 high contribution (>98%

mOrange$^+$ 2 days after addition of dox) TNG MKOS Cas9 MEF plated across 90 wells of 6-well culture plates

were exposed to lentiviral sgRNA library at MOI=2 for 4 hrs. We used MOI=2 (infection efficiency ~86%) in

order to increase coverage of the sgRNA library, presuming the scarcity of reprogramming relevant genes

and the negligible probability of the same neutral sgRNAs being repeatedly present in combination with

relevant sgRNAs. After viral containing media was removed, the cells were cultured in 3 ml of

reprogramming medium. Medium was replaced once 3 ml a day for the first 3 days, and then twice 4 ml a

day from day 4 of reprogramming. From day 8, the media was switched to ESC medium supplemented with

puromycin (1 µg ml^{-1}) and medium was replenished twice a day with 4 ml / well until day 16. Puromycin

resistant, *Nanog*-GFP$^+$ cells were then sorted using the FACS AriaII (BD Biosciences) and stored at -80°C as

cell pellets before extraction of genomic DNA. Screening was performed in triplicate. Genomic DNA from

3×10^7 sorted GFP$^+$ cells was extracted using the Blood & Cell Culture DNA Maxi Kit (Qiagen). Amplification

of sgRNA regions from the extracted genome and the original sgRNA plasmid library, and Illumina

sequencing was performed as described before[46]. sgRNA read count data was analysed with MAGeCK

(version 0.4.4)[23] and genes with enriched and depleted sgRNAs were detected using the test command

(default parameters).

Cas9 TNG MKOS MEF reprogramming

0.25×10^4 Cas9 TNG MKOS MEFs were mixed with 9.5×10^4 WT MEFs (129 strain) and seeded in gelatine-

coated wells of 6-well plates. Cells were transduced with sgRNAs lentiviruses at an MOI of 3 with 8 µg ml^{-1}

Polybrene (Merck-Millipore) for 4 hours and then reprogramming was initiated by addition of

reprogramming medium. On day 14, whole well colony images were taken using the Celigo S Cell

Cytometer (Nexcelom) and the number of *Nanog*-GFP$^+$ and *Nanog*-GFP$^-$ colonies were counted. The images

shown for illustration were stitched using Celigo S Cell Cytometer and processed using ImageJ.

piggyBac reprogramming of MEFs with sgRNA expression and/or Zfp266 cDNAs

Nanog-GFP MEFs with or without Cas9 expression from the *Rosa* locus isolated from E12.5 embryos, or wild

type MEFs were plated at 1.5x10^5 cells per well in a gelatine-coated 6-well plate. 24 hrs later co-

transfection of a Dox-inducible *piggyBac* transposon vector carrying the *tetO-MKOS-ires-mOrange* or *tetO-*

STEMCCA-ires-mOrange cassette with sgRNA expression cassette, *PB-CA-rtTA* vector with/without carrying

a *P2A*-linked *Zfp266* cDNAs, and *pCMV-hyPBase* was performed using 750 ng each DNA and 9 µl of

FugeneHD (Promega) as per manufacturer's instructions[18,47,48]. 24 hrs later reprogramming was initiated

with reprogramming medium. Medium was changed every 2 days. For colony counting, whole well colony

images were taken on day 14 using the Celigo S Cell Cytometer (Nexcelom) and colonies were counted with

ImageJ.

piggyBac reprogramming of NSCs with sgRNA expression

A GFP sgRNA vector was delivered into Cas9 and GFP expressing NSCs using nucleofection with the SG Cell

Line 4DNucleofector X Kit (Lonza) as per manufacturer's instructions[49]. GFP$^-$ NSCs were sorted using the

FACS AriaII (BD Biosciences) and plated at clonal density. Individual clones were picked and genotyped to

confirm GFP KO. NSCs were reprogrammed by nucleofection of a Dox-inducible *piggyBac* transposon vector

carrying the *tetO-MKOS-ires-mOrange* cassette with/without a sgRNA expression cassette, *PB-CA-rtTA*

vector and *pCMV-hyPBase*. 2x10^5 NSCs for essential gene expression were nucleofected with 750 ng each of

the above-mentioned plasmids using SG Cell Line 4DNucleofector X Kit (Lonza) as per manufacturer's

instructions. Cells were then plated on a layer of wild type MEF feeder cells seeded the day before at a

density of 1 x10^5 cells per well in a gelatin-coated 6-well plate. One day post-nucleofection, reprogramming

was initiated with NSC complete medium supplemented with 100 U ml^{-1} human LIF, 0.3µg ml^{-1} of

doxycycline (Sigma) and 10 µg ml^{-1} of L-ascorbic acid or 2-Phospho-L-ascorbic acid trisodium salt (Sigma)

(sigma). After 6 days, the medium was switched to serum-free N2B27-based medium (containing

DMEM/F12 medium supplemented with N2 combined 1:1 with Neurobasal® medium supplemented with

B27; all from Thermo Fisher Scientific), MEK inhibitor (PD0325901, 0.8 µM, Axon Medchem), GSK3b

inhibitor (CHIR99021, 3.3 µM, Axon Medchem), 1 µg ml^{-1} of doxycycline (Sigma) and 10 µg ml^{-1} of L-ascorbic

acid or 2-Phospho-L-ascorbic acid trisodium salt (sigma). At day 16 of reprogramming, immunofluorescence

for NANOG was performed as follows: cells fixed with 4% paraformaldehyde for 10 minutes on day 14 were

permeabilized in 0.1% Triton-X in PBS for 1 hours, blocked in 5% BSA in PBS with 0.1% Tween20 for 1 hour

at room temperature, and then stained in blocking solution with a primary antibody for NANOG (eBioMLC-

51, Thermofisher Scientific) overnight at 4 °C. The next day, an AlexaFluor488 conjugated secondary

antibody (A-21208, Invitrogen) was applied in blocking solution for 45 minutes at room temperature before

washing and imaging. Whole well images were taken using the Celigo S Cell Cytometer (Nexcelom) and

colonies were counted with ImageJ.

CD44, ICAM1, *Nanog*-GFP expression analysis during reprogramming

Cells harvested at different time points of reprogramming were stained in FACS buffer for 30 min at 4°C and

washed with FACS buffer prior to acquisition with LSR Fortessa (BD Biosciences) cytometer. The following

antibodies from eBioscience were used: ICAM1-biotin (Clone: 13-0541; Dilution: 1/100), CD44-APC (Clone:

17-0441; Dilution 1/300), streptavidin-PE-Cy7 (Clone: 25-4317-82; Dilution: 1/1500). Dead cells were

excluded using LIVE/DEAD™ Fixable Near-IR Dead Cell Stain Kit (ThermoFisher Scientific, Dilution: 1/1500).

Data were analysed using Flowjo v10.

RNA-Seq

Sample Preparation. For Wt and *Zfp266* KO MEF samples, 1 x 10^5 Cas9 TNG MKOS MEFs were transduced

with either a non-targeting control sgRNA or *Zfp266* sgRNA lentivirus at an MOI of 3 with 8 µg ml^{-1}

polybrene (Merck-Millipore) for 4 hours. After additional 96 hours culture in MEF media, the cells were

harvested for RNA extraction. For reprogramming samples, 0.25 x 10^4 Cas9 TNG MKOS MEFs were mixed

with 9.5 x 10^4 WT MEFs (129 strain) and seeded in gelatine-coated wells of 6-well plates. Cells were

transduced with either a non-targeting control sgRNA or *Zfp266* sgRNA lentivirus at an MOI of 3 with 8 µg

ml^{-1} Polybrene (Merck-Millipore) for 4 hours, before being recovered for 24 hours in MEF media. After 24

hours reprogramming was initiated by addition of reprogramming medium. Cells were harvested at day 3,

day 5 and day 7 of reprogramming, respectively, and 1 x 10^5 of mOrange$^+$ OSKM expressing cells were

sorted with the FACS AriaII (BD Biosciences) per sample. *Nanog*-GFP$^+$ iPSCs were harvested at day 15, and

sorted with the FACS AriaII (BD Biosciences) into 96-well plates. Sorted iPSCs were cultured in ESC medium

with puromycin (1 µg ml^{-1}) to select for transgene independent clones and KO of *Zfp266* was confirmed by

genotyping. *Zfp266* KO ESCs were generated by transfecting Clone J ESCs with a *Zfp266* sgRNA plasmid

expressing BFP. Single BFP$^+$ ESCs were then sorted with the FACS AriaII (BD Biosciences) into 96-well plates

48 hours after transfection. Clones which became BFP- negative (i.e. shed the sgRNA plasmid) were

selected and KO of *Zfp266* was confirmed by genotyping. 1 x 10^5 iPSCs or ESCs were used for RNA

preparation. Cells were homogenised with the QIAshredder kit (Qiagen) and total RNA was extracted from

all samples using the RNeasy Plus Micro Kit (Qiagen). Libraries were prepped with the NEB Ultra II stranded

mRNA Library prep kit (NEB). RNA-Seq libraries were sequenced with NextSeq, 75SE.

Read processing. For each sequencing run, a quality control report was generated using FastQC and

Illumina TruSeq adapter sequences were removed using Cutadapt[50]. Sequencing runs from the same

biological sample were then concatenated and mapped to the GRCm38 reference genome using STAR[51].

Differential analysis. For each biological sample, aligned sequencing reads were first assigned to genomic

features (e.g., genes) using Rsubread[52] and a count table was generated. Differential expression analysis

was then performed with DESeq2[53], and statistically significant genes (e.g., FDR < 0.05 and

log2FoldChange > 1) were identified using the standard workflow. Importantly, although the data

represents a control and treatment time-series experiment, we opted to combine the factors of interest

into a single factor for easier comprehension. Gene ontology analysis for differentially expressed genes was

performed using the goseq package[54].

Downstream analysis. For exploratory analysis and visualization, a batch-corrected and regularized log

matrix of expression values was used. The count table was first transformed to stabilize the variance across

the mean using the rlog function from DESeq2 and then unwanted batch effects (e.g., library preparation

date) were removed using the removeBatchEffect function from limma[55].

DamID-seq

Sample Preparation. 1 x 10[5] WT MEFs (129 strain) were nucleofected with either *PGK-mO-Dam* or *PGK-mO-Dam-Zfp266* plasmids using the P4 Primary Cell 4D-Nucleofector X Kit (Lonza). 5 replicates were performed in total. Cells were recovered in MEF media for 48 hours before 3 x 10[4] - 1.6 x 10[5] GFP+ cells per sample were sorted with the FACS AriaII (BD Biosciences). Genomic DNA was isolated with Quick-gDNA™ MicroPrep (ZymoResearch) and 32 ng genomic DNA/sample was used for DamID-seq library preparation as previously described[37]DamID libraries were sequenced with NextSeq, 40PE.

Read processing. For each sequencing run, a quality control report was generated using FastQC and Illumina Nextera adapter sequences were removed using Cutadapt. Sequencing runs from the same biological sample were then concatenated and mapped to the GRCm38 reference genome using BWA[56]. Uninformative and spurious alignments were subsequently filtered using a combination of SAMtools[57] and BEDtools[58] commands. Specifically, reads mapped to the mitochondrial chromosome and reads mapped to blacklisted regions were filtered.

Peak calling. For each biological sample, aligned sequencing reads were assigned to genomic features (e.g., DpnII restriction fragments) using Rsubread and a count table was generated. Statistically significant regions of Dam-fusion protein binding (e.g., FDR < 0.05 and log2FoldChange > 1) were detected using the callPeak command from Daim[37]. For further details, please refer to the original manuscript describing the Daim software[37]. The regions were then annotated and analysed for gene and genome ontology enrichment using the annotatePeaks command from HOMER[59].

Downstream analysis. Heatmaps of read coverage at Dam-fusion binding regions were produced using the computeMatrix and plotHeatmap commands from deepTools[60]. When plotting heatmaps, a total of 5 peaks identified exactly over *Zfp266* exons (chr9:20495068-20521417) were removed from the *Zfp266* DamID peak regions due to the high signal intensity caused by the *PGK-mO-Dam-Zfp266* plasmid. *De novo* motif discovery and was performed using the MEME-ChIP tool from the MEME suite (version 5.1.1)[61]. Motif enrichment analysis was performed using findMotifsGenome command from HOMER[59] as DamID-seq's large perk size was not optimal for the MEME-ChIP tool. Genome browser images of peak regions and read coverage were composed using the Integrative Genomics Viewer[62].

ATAC-seq

Sample Preparation. Cas9 TNG MKOS MEFs were thawed, plated and transduced in the same manner as

samples prepared for RNA-Seq. After 24 hours reprogramming was initiated by addition of reprogramming

medium for reprogramming samples, while MEF samples were maintained in MEF media. Cells were

harvested 96 hours after sgRNA transduction (which was 72 hours after OSKM induction for reprogramming

samples) and sorted with the FACS AriaII (BD Biosciences). Cells were then processed for ATAC-Sequencing

according to the Omni-ATAC protocol[63]. Briefly, 5×10^4 sorted Cas9 TNG MKOS MEFs per sample were

washed with cold 1x PBS then pelleted before the supernatant was discarded. Cell pellets were then gently

resuspended in 50 μl of lysis buffer (48.5 μl resuspension buffer, 0.5 μl 10% NP-40 (Sigma) 0.5 μl 10%

Tween-20 (Sigma), 0.5 μl 1% Digitonin (Promega) (resuspension buffer: 500 μl 1M Tris-HCl, pH7.5

(ThermoFisher), 100 μl 5M NaCl (Sigma), 150 μl 1M $MgCl_2$ (Sigma), 49.25 ml nuclease-free H_2O) and

incubated on ice for 3 minutes. Then, 1 ml of wash buffer (990 μl resuspension buffer, 10 μl Tween-20

(Sigma)) was added to the tubes before they were gently inverted and then centrifuged for 10 minutes at

500 x g, at 4 °C. Supernatants were then carefully aspirated. Nuclei pellets were then resuspended in 50 μl

of transposition mix (2.5 μl Tn5 transposase, 25 μl 2x TD buffer (both Illumina), 0.5 μl 1% Digitonin

(Promega), 0.5 μl 10% Tween-20 (Sigma), 16.5 μl 1x PBS, 5μl nuclease-free H_2O) and incubated in a

thermomixer at 37 °C, 1000 rpm for 30 minutes. Transposed DNA was then purified with the Zymo DNA

Clean and Concentrator-5 Kit (Zymo Research) and eluted in 21 μl nuclease-free H_2O. All purified DNA (~20

μl) was used for PCR amplification with NEBNext High Fidelity 2x MasterMix (NEB) and optimum cycle

number was determined by qPCR, as per the protocol. Amplified DNA was then purified with double-sided

bead purification using AMPure XP magnetic beads (Beckman Coulter). Library concentration was

determined with Qubit (ThermoFisher) and fragment size/quality with TapeStation (Agilent). ATAC libraries

were sequenced with NextSeq, 40PE.

Read processing. For each sequencing run, a quality control report was generated using FastQC and

Illumina Nextera adapter sequences were removed using Cutadapt[50]. Sequencing runs from the same

biological sample were then concatenated and mapped to the GRCm38 reference genome using BWA[56].

Duplicate reads caused by PCR amplification were subsequently identified using the MarkDuplicates

command from Picard (https://broadinstitute.github.io/picard/). Uninformative and spurious alignments

were next filtered using a combination of SAMtools[57] and BEDtools[58] commands. Specifically, reads mapped

to the mitochondrial chromosome, reads mapped to blacklisted regions, reads marked as duplicates, and

reads not properly paired (e.g., reads that aren't FR orientation or with an insert size greater than 2 kb)

were filtered.

Peak calling. For each biological sample, statistically significant regions of chromatin accessibility (e.g., FDR

< 0.1) were detected using the callpeak command from MACS2[64] (https://github.com/macs3-

project/MACS). For downstream analyses, a consensus set of peaks was created by taking the union across

all biological samples with the multiinter command from BEDtools[58].

Differential analysis. For each biological sample, aligned sequencing reads were first assigned to genomic

features (e.g., consensus set of peaks) using Rsubread[52] and a count table was generated. Differential

accessibility analysis was then performed with DESeq2[65] and statistically significant peaks (e.g., FDR < 0.05

and log2FoldChange > 1) were identified using the standard workflow.

Downstream analysis. Heatmaps of read coverage at chromatin accessibility regions were produced using

the computeMatrix and plotHeatmap commands from deepTools[60]. K-means clustering was used to

partition the regions into two distinct categories of reprogramming MORs. Genome browser images of

peak regions and read coverage were composed using the Integrative Genomics Viewer[62]. Peaks were

annotated against mm10 with annotatePeaks.pl from the HOMER suite (version 4.11)[59]. *De novo* motif

discovery and enrichment analysis of MORs were performed using the *Zfp266* KO samples' narrowpeak

summits within MORs with the MEME-ChIP tool from the MEME suite (version 5.1.1)[61]. The number of SINE

elements around peaks were counted using the BEDTools window command in a window of ±500 bp from

the summits of the peaks. ATAC-seq data of iPSCs were retrieved from GSE98124[39].

Luciferase Reporter Assays

The pGL3 reporter plasmid containing the SV40 early promoter (Promega) was used for all luciferase

reporter assays along with an internal control Renilla plasmid (Promega). Luciferase activity was measured

with the GloMax 96-microplate luminometer (Promega) using the Dual-Glo Luciferase Assay System

(Promega). For assays performed in HEK293 cells, $0.5\text{-}1\text{x}10^4$ cells were plated per well of a 96-well plate 24

hours prior to transfection. Transfection mixes were prepared as follows; 100 ng pGL3 reporter plasmid, 0.5

ng Renilla plasmid and 100 ng overexpression plasmid (BFP/VPR/VPR-*Zfp266*/Wt *Zfp266*) were mixed in

Opti-MEM I Reduced Serum Medium (Gibco) up to 100μl. Fugene HD Transfection Reagent (Promega) was

then added at a ratio of 3:1 (reagent:DNA) and 5-10 μl was added to each well of cells. Luciferase activity

was measured 48 hours after transfection. For assays performed in MEF/reprogramming cells, $1\text{x}10^4$ TNG

MKOS MEFs were plated per well of a 96-well plate 24 hours prior to transfection, either in MEF media or

ES media +dox (300 ng/ml) to induce OSKM expression. Transfection mixes were prepared as such; 1 μg

pGL3 reporter plasmid, 10 ng Renilla plasmid were mixed in Opti-MEM I Reduced Serum Medium up to 100

μl. Fugene HD Transfection Reagent was then added at a ratio of 4:1 (reagent:DNA) and 20μl was added to

each well of MEFs/reprogramming cells. Luciferase activity was measured 48 hours after transfection.

END NOTE

Acknowledgements

We thank I. Chambers for providing TNG ESC line, F. Rossi and C. Cryer for assistance with flow cytometry,

Biomed unit staff for mouse husbandry, the Wellcome Sanger Institute sequencing facility for gRNA

sequencing, EMBL GeneCore for RNA-seq, ATAC-seq and DamID-seq, A. Soufi and members of the Kaji and

Soufi labs for comments on the manuscript. Some of the computations for this work were enabled by

resources in project SNIC 2017/7-317 provided by the Swedish National Infrastructure for Computing (SNIC)

at the Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX). This work was

supported by European Research Council (ERC) grants ROADTOIPS (no. 261075) and MRC senior non-

clinical fellowship (MR/N008715/1) funded for K.K. We also thank the generous support from Baillie Gifford

for the collaboration between CiRA and MRC CRM, from Japan Agency for Medical Research and

Development (AMED) for CiRA. K.Y. was supported by the Wellcome Trust (206194). D.F.K., J.A. and M.Y.

was supported by the BBSRC (EASTBIO doctoral training partnership), Principal's Career Development

scholarship from the University of Edinburgh, and Japan Society for the Promotion of Science (JSPS)

Overseas Research Fellowships, respectively. V.O. and M.Bertenstam were supported by the Swedish

Foundation for Strategic Research (A3 04 159p). V.O. was also supported by the Swedish Research Council

(Vr 621-2008-3074).

Author Contribution

D.F.K. designed and performed sgRNA screen, validation and characterization of the roadblock genes. M.A,

J.A., S.K. and S.T. contributed to the analyses of the gRNA sequencing, RNA-seq, ATAC-seq, ChIP-seq and

DamID-seq data sets. M.B. and S.Z. provided technical support. M.Bertenstam and V.O. generated the

screening data website. K.O. advised and supervised D.F.K. K.Y. provided the gRNA library, the Rosa26-Cas9

targeting vector, and advised on the screen. K.K. conceived the study, supervised experiment design and

data interpretation, and wrote the manuscript with D.F.K.

Author Information

The authors declare no competing financial interests. Correspondence and requests for materials should be

address to K.K. (keisuke.kaji@ed.ac.uk).

References

1.   Takahashi, K. & Yamanaka, S. Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult

Fibroblast Cultures by Defined Factors. *Cell* **126**, 663–676 (2006).

2.   Zhao, Y. *et al.* Two Supporting Factors Greatly Improve the Efficiency of Human iPSC Generation. *Cell*

Stem Cell **3**, 475–479 (2008).

3.   Banito, A. *et al.* Senescence impairs successful reprogramming to pluripotent stem cells. *Genes Dev.*

23, 2134–2139 (2009).

4.   Hong, H. *et al.* Suppression of induced pluripotent stem cell generation by the p53-p21 pathway.

Nature **460**, 1132–1135 (2009).

5.   Kawamura, T. *et al.* Linking the p53 tumour suppressor pathway to somatic cell reprogramming.

Nature **460**, 1140–1144 (2009).

6.   Li, H. *et al.* The Ink4/Arf locus is a barrier for iPS cell reprogramming. *Nature* **460**, 1136–1139 (2009).

7.   Marión, R. M. *et al.* A p53-mediated DNA damage response limits reprogramming to ensure iPS cell

genomic integrity. *Nature* **460**, 1149–1153 (2009).

8.   Utikal, J. *et al.* Immortalization eliminates a roadblock during cellular reprogramming into iPS cells.

Nature **460**, 1145–1148 (2009).

9.   Onder, T. T. *et al.* Chromatin-modifying enzymes as modulators of reprogramming. *Nature* **483**, 598–

602 (2012).

10.   Soufi, A., Donahue, G. & Zaret, K. S. Facilitators and Impediments of the Pluripotency

Reprogramming Factors' Initial Engagement with the Genome. *Cell* **151**, 994–1004 (2012).

11.   Bourque, G. *et al.* Ten things you should know about transposable elements. *Genome Biol. 2018 191*

19, 1–12 (2018).

12.   Percharde, M. *et al.* A LINE1-Nucleolin Partnership Regulates Early Development and ESC Identity.

Cell **174**, 391-405.e19 (2018).

13.   Ohnuki, M. *et al.* Dynamic regulation of human endogenous retroviruses mediates factor-induced

reprogramming and differentiation potential. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 12426–12431

(2014).

14.   Pontis, J. *et al.* Hominoid-Specific Transposable Elements and KZFPs Facilitate Human Embryonic

Genome Activation and Control Transcription in Naive Human ESCs. *Cell Stem Cell* **24**, 724-735.e5

(2019).

15.   Lu, J. Y. *et al.* Genomic Repeats Categorize Genes with Distinct Functions for Orchestrated

Regulation. *Cell Rep.* **30**, 3296-3311.e5 (2020).

16.   Yang, P., Wang, Y. & Macfarlan, T. S. The Role of KRAB-ZFPs in Transposable Element Repression and

Mammalian Evolution. *Trends Genet.* **33**, 871–881 (2017).

17.   Wolf, G. *et al.* Krab-zinc finger protein gene expansion in response to active retrotransposons in the

murine lineage. *Elife* **9**, 1–22 (2020).

18.   Chantzoura, E. *et al.* Reprogramming Roadblocks Are System Dependent. *Stem Cell Reports* **5**, 350–

364 (2015).

19.   Ruetz, T. *et al.* Constitutively Active SMAD2/3 Are Broad-Scope Potentiators of Transcription-Factor-

Mediated Cellular Reprogramming. *Cell Stem Cell* **21**, 791-805.e9 (2017).

20.   Kareta, M. S. *et al.* Inhibition of Pluripotency Networks by the Rb Tumor Suppressor Restricts

Reprogramming and Tumorigenesis. *Cell Stem Cell* **16**, 39–50 (2015).

21.   Mansour, A. A. *et al.* The H3K27 demethylase Utx regulates somatic and germ cell epigenetic

reprogramming. *Nature* **488**, 409–413 (2012).

22.   Tzelepis, K. *et al.* A CRISPR Dropout Screen Identifies Genetic Vulnerabilities and Therapeutic Targets

in Acute Myeloid Leukemia. *Cell Rep.* **17**, 1193–1205 (2016).

23.   Li, W. *et al.* MAGeCK enables robust identification of essential genes from genome-scale

CRISPR/Cas9 knockout screens. *Genome Biol.* **15**, 554 (2014).

24.   Liu, J. *et al.* The oncogene c-Jun impedes somatic cell reprogramming. *Nat. Cell Biol.* **17**, 856–67

(2015).

25.   Onder, T. T. *et al.* Chromatin-modifying enzymes as modulators of reprogramming. *Nat. 2012*

4837391 **483**, 598–602 (2012).

26.   Yang, C. S., Chang, K. Y. & Rana, T. M. Genome-wide Functional Analysis Reveals Factors Needed at

the Transition Steps of Induced Reprogramming. *Cell Rep.* **8**, 327–337 (2014).

27.   Michlits, G. *et al.* CRISPR-UMI: Single-cell lineage tracing of pooled CRISPR-Cas9 screens. *Nat.*

Methods **14**, 1191–1197 (2017).

28.   O'Malley, J. *et al.* High-resolution analysis with novel cell-surface markers identifies routes to iPS

617   cells. *Nat. 2013 4997456* **499**, 88–91 (2013).

618   29.   Sommer, C. A. *et al.* Induced Pluripotent Stem Cell Generation Using a Single Lentiviral Stem Cell

619   Cassette. *Stem Cells* **27**, 543–549 (2009).

620   30.   Kaji, K. *et al.* Virus-free induction of pluripotency and subsequent excision of reprogramming factors.

621   *Nature* **458**, 771–775 (2009).

622   31.   Reinhardt, A., Kagawa, H. & Woltjen, K. N-Terminal Amino Acids Determine KLF4 Protein Stability in

623   2A Peptide-Linked Polycistronic Reprogramming Constructs. *Stem Cell Reports* **14**, 520–527 (2020).

624   32.   Kim, S. Il *et al.* KLF4 N-terminal variance modulates induced reprogramming to pluripotency. *Stem

625   Cell Reports* **4**, 727–743 (2015).

626   33.   O'Malley, J. *et al.* High-resolution analysis with novel cell-surface markers identifies routes to iPS

627   cells. *Nature* **499**, 88–91 (2013).

628   34.   Ecco, G., Imbeault, M. & Trono, D. KRAB zinc finger proteins. *Development* **144**, 2719–2729 (2017).

629   35.   Margolin, J. F. *et al.* Krüppel-associated boxes are potent transcriptional repression domains. *Proc.

630   Natl. Acad. Sci.* **91**, 4509–4513 (1994).

631   36.   Peng, H., Ivanov, A. V., Oh, H. J., Lau, Y. F. C. & Rauscher, F. J. Epigenetic Gene Silencing by the SRY

632   Protein Is Mediated by a KRAB-O Protein That Recruits the KAP1 Co-repressor Machinery. *J. Biol.

633   Chem.* **284**, 35670–35680 (2009).

634   37.   Tosti, L. *et al.* Mapping transcription factor occupancy using minimal numbers of cells in vitro and in

635   vivo. *Genome Res.* **28**, 592–605 (2018).

636   38.   Vogel, M. J., Peric-Hupkes, D. & van Steensel, B. Detection of in vivo protein–DNA interactions using

637   DamID in mammalian cells. *Nat. Protoc. 2007 26* **2**, 1467–1478 (2007).

638   39.   Benchetrit, H. *et al.* Direct Induction of the Three Pre-implantation Blastocyst Cell Types from

639   Fibroblasts. *Cell Stem Cell* **24**, 983-994.e7 (2019).

40.   Chronis, C. *et al.* Cooperative Binding of Transcription Factors Orchestrates Reprogramming. *Cell*

168, 442-459.e20 (2017).

41.   Roberts, G. A. *et al.* Dissecting OCT4 defines the role of nucleosome binding in pluripotency. *Nat.*

Cell Biol. 2021 238 **23**, 834–845 (2021).

42.   Chavez, A. *et al.* Highly efficient Cas9-mediated transcriptional programming. *Nat. Methods 2015*

124 **12**, 326–328 (2015).

43.   Ye, M. *et al.* Specific subfamilies of transposable elements contribute to different domains of T

lymphocyte enhancers. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 7905–7916 (2020).

44.   Li, C. & Luscombe, N. M. Nucleosome positioning stability is a modulator of germline mutation rate

variation across the human genome. *Nat. Commun. 2020 111* **11**, 1–13 (2020).

45.   Soufi, A. *et al.* Pioneer Transcription Factors Target Partial DNA Motifs on Nucleosomes to Initiate

Reprogramming. *Cell* **161**, 555–568 (2015).

46.   Koike-Yusa, H., Li, Y., Tan, E. P., Velasco-Herrera, M. D. C. & Yusa, K. Genome-wide recessive genetic

screening in mammalian cells with a lentiviral CRISPR-guide RNA library. *Nat. Biotechnol. 2013 323*

32, 267–273 (2013).

47.   Woltjen, K. *et al.* piggyBac transposition reprograms fibroblasts to induced pluripotent stem cells.

Nat. 2009 4587239 **458**, 766–770 (2009).

48.   Yusa, K., Zhou, L., Li, M. A., Bradley, A. & Craig, N. L. A hyperactive piggyBac transposase for

mammalian applications. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 1531–1536 (2011).

49.   Bressan, R. B. *et al.* Efficient CRISPR/cas9-assisted gene targeting enables rapid and precise genetic

manipulation of mammalian neural stem cells. *Dev.* **144**, 635–648 (2017).

50.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal **17**, 10–12 (2011).

51.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

52.   Liao, Y., Smyth, G. K. & Shi, W. The R package Rsubread is easier, faster, cheaper and better for

alignment and quantification of RNA sequencing reads. *Nucleic Acids Res.* **47**, e47–e47 (2019).

53.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biol.* **15**, 550 (2014).

54.   Young, M. D., Wakefield, M. J., Smyth, G. K. & Oshlack, A. Gene ontology analysis for RNA-seq:

accounting for selection bias. *Genome Biol.* **11**, 1–12 (2010).

55.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and

microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

56.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics **25**, 1754–1760 (2009).

57.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079

(2009).

58.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics **26**, 841–842 (2010).

59.   Heinz, S. *et al.* Simple Combinations of Lineage-Determining Transcription Factors Prime cis-

Regulatory Elements Required for Macrophage and B Cell Identities. *Mol. Cell* **38**, 576–589 (2010).

60.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis.

Nucleic Acids Res. **44**, W160–W165 (2016).

61.   Bailey, T. L., Johnson, J., Grant, C. E. & Noble, W. S. The MEME Suite. *Nucleic Acids Res.* **43**, W39–

W49 (2015).

62.   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol. 2011 291* **29**, 24–26 (2011).

63.   Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and enables interrogation

of frozen tissues. *Nat. Methods 2017 1410* **14**, 959–962 (2017).

64.   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, 1–9 (2008).

65.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq

689   data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

692
Figure. 1. A genome-wide CRISPR screen identifies reprogramming roadblock genes. A. Schematic
diagram of the screening strategy. sgRNA library infected Cas9 TNG MKOS MEFs were cultured in +dox for 8
days then in -dox and +Puro for additional 8 days. Integrated sgRNAs were amplified from *Nanog*-GFP+ cells
for Illumina sequencing. **B.** Enrichment FDR ranking with MAGeCK. 24 genes, including 8 previously
reported (green) and 16 novel (blue) roadblock genes, were identified using a cut-off of FDR<0.1. **C**.
Normalized sgRNA read counts in the initial plasmid library versus mutant iPSC pool. sgRNAs against
previously reported roadblock genes (red/orange) and genes essential for reprogramming (green) exhibited
expected enrichment/depletion respectively (left). sgRNAs against 16 novel roadblock genes identified in
this screen are highlighted in blue (right). **D**. Enrichment rank, FDR, and function of the 24 discovered
roadblock genes. **E**. Validation of the screen result with 3 individual sgRNAs per gene. This graph is a
summary of 5 data sets shown in Supplemental Figure S2A. Error bars indicate SEM, ****p<0.0001,
***p<0.001, **p<0.01,*p<0.05 based on an unpaired two-tailed t-test **F**. Representative whole-well images
of KO reprogramming of 13 top roadblocks from E. Previously reported and novel roadblock genes were
labelled in green and and blue, respectively. Red; mOrange, Green; *Nanog*-GFP.

Figure 2. Characterization of the roadblock gene KO in different reprogramming systems and kinetics. A,

C. *Cas9* expressing *Nanog*-GFP MEF reprogramming with MKOS (A), STEMCCA (C) *piggyBac* transposons

with sgRNA expression at day 15. Red; mOrange, Green; *Nanog*-GFP. **B,D.** *Nanog*-GFP+ and *Nanog*-GFP-

mean colony numbers from A and C. **E.** *Cas9* expressing NSC reprogramming with MKOS *piggyBac*

transposons with sgRNA expression at day 15. Green; immunofluorescence for NANOG. **F.** NANOG+ and

NANOG- mean colony numbers from E. **G.** Accelerated CD44/ICAM/*Nanog*-GFP expression changes by

sgRNA expression against the roadblock genes (n=2). Red; *Nanog*-GFP- cells, Green; *Nanog*-GFP+ cells. **H.**

Quantification of *Nanog*-GFP+ cells from day 6 to 10 of reprogramming. The graph represents an average of

2 independent experiments. For all graphs error bars indicate SEM, ****p<0.0001, ***p<0.001,
**p<0.01,*p<0.05 based on an unpaired two tailed t test.

Figure 3. ZFP266 impedes activation of pluripotency genes via its KRAB domain. A. Diagram of *Zfp266* Wt
and mutants. A red bar indicates a silent mutation that confers sgRNA resistance. KRAB domain deletion
mutants with (ΔKRAB+L) and without a linker (ΔKRAB) do not have the sgRNA target sequence. DV-AA,
DEW-AAA, MLE-AAA mutants have alanine substitutions in the indicated critical amino acids in the KRAB
domain[28,29]. Triple mutant contains all the alanine substitutions. **B.** *Nanog*-GFP MEF reprogramming with
MKOS *piggyBac* transposons and BFP or Wt *Zfp266* cDNA overexpression, imaged at day 15. Red; mOrange,
Green; *Nanog*-GFP. Error bars indicate SEM, *p<0.05 based on an unpaired two-tailed t-test. **C.** *Cas9*
Nanog-GFP MEF reprogramming with MKOS *piggyBac* transposons, *Zfp266* sgRNA expression as well as
BFP, Wt *Zfp266, or* mutant *Zfp266* cDNA overexpression, imaged at day 15. Red; mOrange, Green; *Nanog*-
GFP. **D**. Mean *Nanog*-GFP⁺ colony numbers of C. Error bars indicate SEM, **p<0.01 based on a one-way
ANOVA test. **E.** RNA-Seq volcano plot of *Zfp266* KO vs Wt MEF, day 3, 5 and 7 of reprogramming. Up-
regulated and down-regulated genes in KO cells are shown to the right and left of the plot, respectively
(cut-off FDR<0.05, fold change >2). ESC- and MEF-associated genes (FDR<0.05, fold change >2 in ESCs vs
MEFs) are highlighted in green and red. Graphs below volcano plots show the number of ESC-associated,
MEF-associated and other genes within D3, D5, D7 reprogramming differentially expressed genes (DEGs). **F.**
GO enrichment analysis of upregulated and downregulated genes in *Zfp266* KO reprogramming. **G.** Principal
component analysis of *Zfp266* Wt and KO RNA-Seq samples. Blue dots indicate Wt samples, red dots
indicate KO samples, three samples per timepoint.

Figure 4. *Zfp266* KO in MEFs results in chromatin opening at the SINE-containing ZFP266 binding sites. A.
ZFP266 DamID-seq signals in MEFs, ATAC-seq signals in MEFs, +72hours of reprogramming, and iPSCs at the
ZFP266 DamID-seq peak loci. **B.** Motif enrichment analysis with HOMER on ZFP266 DamID-seq peaks. **C.**
Significance and fold enrichment ratio of transposable element (TE) families overlap with Dam-ZFP266
peaks. Green dots indicate significantly enriched SINEs, blue dots indicate other significantly enriched TEs.
D. Number of ZFP266 DamID-seq peaks that overlap with TEs. **E.** *De novo* motif discovery analysis with
MEME on ZFP266 DamID-seq peaks. The identified motifs correspond to parts of the B1 SINE consensus
sequence, indicated by matching colours. **F.** Volcano plot of *Zfp266* KO vs Wt MEF ATAC-seq. Green and red
dots indicate more open regions (MORs) and more closed regions in *Zfp266* KO MEFs, respectively
(FDR<0.05). **G.** ATAC-seq and ZFP266 DamID-seq signals in the *Zfp266* KO MEF MORs, overlapped (top) and
non-overlapped (bottom) with ZFP266 DamID peaks in MEFs. **H.** Motif enrichment analysis on *Zfp266* KO
MEF MORs. Percentages of MORs containing each motif and AP-1 motif are indicated. **I**. *De novo* motif
discovery analysis on *Zfp266* KO MEF MORs. The top five most significant motifs correspond to parts of the
B1 SINE consensus sequence, indicated by matching colours. **J**. Significance and fold enrichment ratio of
transposable element (TE) families overlap with Zfp266 KO MEF MORs. Green dots indicate significantly
enriched SINEs, blue dots indicate other significantly enriched TEs. **K.** Number of *Zfp266* KO MEF MORs that

754   overlap with TEs. **L.** Examples of *Zfp266* KO MEF MORs (Blue) with SINE (black), overlapping with ZFP266
755   DamID-seq peaks (red).

Figure 5. Reprogramming *Zfp266* KO MEFs results in chromatin opening at OSK bound, B1 SINE containing loci. A. ATAC-seq volcano plot of *Zfp266* KO vs Wt reprogramming (+72hours after OSKM induction). Green and red dots indicate more open regions (MORs) and more closed regions in *Zfp266* KO reprogramming, respectively (FDR<0.05). **B.** Significance and fold enrichment ratio of transposable element (TE) families overlap with Zfp266 KO MEF MORs. Green dots indicate significantly enriched SINEs, blue dots indicate other significantly enriched TEs. **C.** Number of *Zfp266* KO reprogramming MORs that overlap with TEs. **D.** *De novo* motif discovery analysis on *Zfp266* KO reprogramming MORs. The motifs correspond to parts of the B1 SINE consensus sequence are indicated by matching colours. **E.** Overlap between *Zfp266* KO MEF MORs and *Zfp266* KO reprogramming MORs. **F.** Motif enrichment analysis with *Zfp266* KO reprogramming MOR peak summits and percentages of MORs with KLF, SOX, OCT (POU) family and OCT4::SOX2 motifs. **G.** Classification of *Zfp266* KO reprogramming MORs based on ATAC-seq signals in MEF and reprogramming 72 hours. **H.** Reprogramming 48 hours OSKM ChIP-Seq heatmap plots at *Zfp266* KO reprogramming MORs. **I.** KLF, SOX, OCT (POU) family, OCT4::SOX2 and SINE motif distribution within *Zfp266* KO reprogramming MORs. 70 bp from the summit is highlighted in pink. The colours of B1 SINE motifs correlate to those in D. **J.** Orientation-biased distribution of B1 SINE elements within *Zfp266* KO reprogramming MORs. Head of B1 SINE tends to locate on the MOR summit side. 70 bp from the summit is highlighted in pink.

Figure 6. ZFP266 binds to B1 SINEs in Zfp266 KO reprogramming MORs that impede OSKM-mediated gene activation. A. Diagram of Wt ZFP266 and a synthetic activator version of ZFP266, VPR-ZFP266. **B, C.** Luciferase reporter assay with either an empty reporter vector pGL3 or with a reporter vector with the B1 SINE consensus sequence, co-expressed with either BFP, VPR only or VPR-*Zfp266* expression vectors (**B**), with either BFP or Wt *Zfp266* expression vectors in the presence of VPR-ZFP266 (**C**) in HEK293 cells. RLU: Relative Light Units, *p<0.05, ****p<0.0001 based on a two-way ANOVA test. **D.** *B3gnt3*, *Piwil2* and *Snx20* mRNA expression from the *Zfp266* Wt and KO reprogramming RNA-seq data. **E.** ATAC-seq, ChIP-seq signals at the *B3gnt3*, *Piwil2* and *Snx20* MORs, cloned in both forward (F) and reverse (R) directions (relative to gene orientation) for luciferase reporter assays (highlighted in orange). **F-J**. Luciferase reporter assay with an empty reporter vector pGL3 or vectors containing *B3gnt3*, *Piwil2* and *Snx20* MORs co-transfected with

either BFP, VPR only or VPR-*Zfp266* expression vectors (**F**), co-transfected with either BFP or Wt *Zfp266*
expression vectors in the presence of VPR-ZFP266 (**G**), an empty reporter vector pGL3, vectors containing
B3gnt3 (**H**), *Piwil2* (**I**) and *Snx20* (**J**) MORs with (ΔB1) or without (Wt) B1 SINE deletion co-transfected with
either BFP, VPR only or VPR-*Zfp266* expression vectors in HEK293 cells. **K, L**. Luciferase reporter assay with
empty reporter vector pGL3 or vectors containing *B3gnt3*, *Piwil2* and *Snx20* MORs (**K**), *Snx20* MOR reporter
with (ΔB1) or without (Wt) B1 SINE deletion (**L**), using MEF with or without OSKM expression. *p<0.05,
p<0.01, ***p<0.001, ****p<0.0001 based on a two-way ANOVA test. **M. Day 9 and day 14 after OSKM
induction with overexpression of either BFP, *VPR* only or *VPR-Zfp266*. Red; mOrange, Green; *Nanog*-GFP. **N**.
Quantification of *Nanog*-GFP$^+$ colony numbers at day 9 and day 14. ****p<0.0001 based on a one-way
ANOVA test. **O**. Mechanistic model of how *Zfp266* KO enhances reprogramming. ZFP266 recruited by OSKM
to their target loci binds to adjacent B1 SINE and impedes chromatin opening (top), thus *Zfp266* KO results