Opposite-sex associations are linked with annual fitness, but sociality is stable over

lifetime.

*Jamie Dunning[1], Terry Burke[2], Alex Hoi Hang Chan[1,3,4], Heung Ying Janet Chik[5,6], Tim

Evans[7], Julia Schroeder[1].

[1]Department of Life Sciences, Imperial College London, United Kingdom

[2]Ecology and Evolutionary Biology, School of Biosciences, The University of Sheffield,

United Kingdom

[3]Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Germany

[4]Max Plank Institute of Animal Behaviour, Germany

[5]Groningen Institute for Evolutionary Life Sciences, University of Groningen, Netherlands

[6]School of Natural Sciences, Macquarie University, Australia

[7]Center for Complexity Science, Imperial College London, United Kingdom

*Corresponding author – jamiedunning8@googlemail.com

____

Abstract

Animal sociality, an individual's propensity to associate with others, has fitness

consequences through mate choice, for example, directly, by increasing the pool of

prospective partners, and indirectly through increased survival, and individuals benefit

from both. Annually, fitness consequences are realised through increased mating success

and subsequent fecundity. However, it remains unknown whether these consequences

translate to life-time fitness. Here, we quantified social associations and their link to fitness

annually and over lifetime, using a multi-generational, genetic pedigree. We used social

network analysis to calculate variables representing different aspects of an individual's

sociality. Sociality showed high within-individual repeatability. We found that birds with

more opposite-sex associates had higher annual fitness than those with fewer, but this did

not translate to lifetime fitness. Instead, for lifetime fitness, we found evidence for

stabilizing selection on opposite-sex sociality, and sociality in general, suggesting that

reported benefits are only short-lived in a wild population, and that selection favours an

average sociality.

Keywords: sociality, social network analysis, selection, social behaviour, genetic pedigree,
long-term fitness, de-lifing, lifetime reproductive success

Background

Some individuals are consistently more sociable than others, demonstrated by within-individual repeatability of social traits across vertebrate groups (Aplin et al. 2015; Thys et al. 2017; Dimitriadou, Croft and Darden 2019; Plaza et al. 2019; Beck, Valcu, and Kempenaers 2020; Proops et al. 2021; Strickland et al. 2021). This variation in individual sociality is positively linked with fitness in some taxa (Silk 2007; Silk et al. 2009) and is therefore expected to be subject to selection (Krause and Ruxton 2002). Fitness is a relative measure of an individual's genetic contribution to the population in the next generation, and thus, can only be represented comprehensively and precisely by measures of traits spanning an organism's lifetime (Endler 1986; Blankenhorn 2010; Reid et al. 2019; Moiron, Charmantier and Bouwhuis 2022). A comprehensive definition of fitness is fundamental to understand the evolutionary pressures that shape variation in sociality. In practice, however, many studies must rely on fitness correlates instead (e.g., number of broods, or survival, instead of genetic contribution). For example, in mammal societies, both variation in within- and between-sex affiliations (Archie et al. 2014) have been linked to lifetime fitness correlates, survival and longevity (Cameron, Setsaas and Linklater 2009; Silk, Alberts and Altman 2003; Silk et al. 2010; Stanton and Mann 2012, but also see Thompson and Cords 2018). Whereas, in birds, the subject of this study, the use of fitness correlates (eggs laid, chicks fledged, within-year survival etc.) are frequent over more precise fitness measures (Moiron, Charmantier and Bouwhuis 2022), that require intensive field work over a long period of time.

Although some benefits are linked with sociality during the breeding period most of these tend to be short-term and contextual (Bebbington et al. 2017; Riehl and Strong 2018). Instead, benefits associated with reproduction are often linked with non-breeding sociality (Firth and Sheldon 2016; Kohn 2017; Maldonado-Chaparro et al. 2018; Beck, Farine, and Kempenaers 2020), when group cohesion is stronger (Silk et al. 2014; Kurvers 2020). Sociality may influence fitness in different ways, through benefits to reproductive success or increased survival, and so, the mechanism of selection acting on social traits may also

differ. For example, communal foraging between socially associated individuals during the

non-breeding period facilitates resource information transfer (Aplin et al. 2012; Firth,

Sheldon and Farine 2016; Hillemann et al. 2020) and reduces predation risk (Cresswell

1994; Cresswell and Quinn 2011; Sorato et al. 2012), increasing survival. However, these

benefits may also incur costs associated with competition for resources and mate choice

(Birkhead and Biggins 1987; Le Galliard et al. 2005; Forstmeier et al. 2011; Mayer and

Pasinelli 2013; Grant and Grant 2019; but also see Lea et al. 2010). Sociality may also

benefit individuals who hold more central social network positions or have access to

opposite-sex associates, through enhansed mate choice (McDonald 2007; Oh and Badyaev

2010; Firth et al. 2018; Beck, Farine, and Kempenaers 2021). Although the association

between annual fitness correlates and non-breeding sociality has been well described,

testing how selection acts on social traits requires lifetime fitness measures, and remains

unresolved.

With the recent development of tools to construct and analyse social networks (Wey et al.

2008; Farine and Whitehead 2015) the study of sociality has become popular among

behavioural ecologists. Yet, to describe the association between sociality and fitness any

potential study must first overcome two problems: (1) A social association must be clearly

defined relative to the behavior of the study system (Figure 1; Psorakis et al. 2012, 2015),

and (2) to study the evolution of social behaviour, precise measures of individual fitness

must be quantified to test for correlation with a social trait – selection. Although annual

fitness correlates are widely used, lifetime fitness is more precise and without the

stochasticity of annual measures (Dobson, Murie and Viblanc 2020; Alif et al. 2022) and

thus, can better describe selection pressure acting on a trait (Endler 1986; Blankenhorn

2010; Reid et al. 2019). However, lifetime fitness requires wild animals to be monitored

throughout their whole lives, and all breeding attempts, and the fates of their offspring,

must be recorded to determine recruitment. All of these require a multi-generational,

genetic pedigree (Kruuk 2004; Korsten et al. 2013).

Our study system, an island population of house sparrows *Passer domesticus* (hereafter,

sparrow/s) where we monitor all individuals from birth to death, without capture bias

(Simons et al. 2015), overcomes both problems. (1) We have sociality data from birds that
are electronically registered visiting a feeder. Social centrality measures are repeatable
across different timescales and contexts in this and other populations (Plaza et al. 2019).
(2) We have lifetime recruitment data available, from a multi-generational genetic pedigree
that, because our population is closed, meaningly there is no movement of sparrows to or
from the island, and that our study covers all sparrows on the island, we can use to
compute precise annual and lifetime fitness estimates (Schroeder et al. 2015; Alif et al.
2022).

We tested predictions based on arguments presented above to understand the potential for
selection on sociality: (1) We confirmed that the social traits we measured were
meaningful by testing for individual repeatability of sociality; (2) We tested the prediction
that non-breeding sociality has fitness benefits, either driven by reproductive success
through opposite-sex association or through increased survival through network centrality
measures and is subject to selection.

108
109
110 Methods
111

Study system
We used data from the Lundy sparrow system, a long-term study based on the island of
Lundy (51.11N, 4.40W), ≈19 km off North Devon, UK, in the Bristol Channel. The sparrows
on Lundy breed in nest boxes, sited in groups around the only village on the island. The
island is rodent-free and therefore the sparrows have no predators but for the occasional
vagrant raptor. House sparrows are a model organism in behavioral ecology and evolution,
and much is known about their biology, physiology and life-history (Andersson 1994;
Sánchez-Tójar et al. 2018). House sparrow are socially monogamous, but 25% of broods
show they can be genetically promiscuous (Schroeder et al. 2016). On Lundy, they have on
average 2-3 broods of 4-5 eggs per breeding season (Westneat et al. 2014). The sex ratio is
stable, and the mean lifespan of recruits is three years (Alif et al. 2022). Although sparrows
are territorial during the breeding season, during the non-breeding period they form

gregarious groups that forage communally for seed and at supplementary bird feeders

(Summer-Smith 1963), both of which are available year-round on Lundy.

Most sparrows were first captured, and tissue sampled in nest boxes at their natal site

during the breeding season (April to August) or using mist nets during the post-fledging

period (Schroeder et al. 2011; Girndt et al. 2019). Tissue samples were either blood or

mouth swabs and were stored in ethanol and refrigerated at $3°C$ prior to analysis. We

genotyped sparrow DNA at <22 microsatellite loci suitable for parentage assignment in

sparrows (Dawson et al. 2012). Using the genetic data, we assembled a near-complete

genetic pedigree (Schroeder et al. 2015, 2016), which at the time of writing spans twenty

years, 2000–2019, and 8,379 individuals. We fitted all sparrows with a unique

combination of a coded metal ring and three coloured leg rings. We also provided each

sparrow with a subcutaneous Passive Identification Transponder (PIT tag; TROVANID100:

11.5 × 2.1 mm and 0.1 g), under the skin of the breast, which we have previously shown

have no detrimental effect on subsequent fitness (for details see Schroeder et al. 2011).

These tagged individuals were then recorded when they visited a custom made 19.8cm x

19.8cm Radio Frequency Identification antenna (RFID; DorsetID) mounted on a seed

reservoir (for photo see Sánchez-Tójar et al. 2017; Brandl et al. 2019), positioned centrally

within our study site. The feeder was open access, and explicitly not limited to a single bird

feeding at one time, as is the case at hanging bird feeders (Youngblood 2019; Beck, Farine

and Kempenaers 2020). Our feeder recorded visiting birds every day that the island

generators were running (6am – midnight, seven days a week).

145
Social centrality measures

To quantify the sociality of individual sparrows we calculated measures of social centrality

(hereafter centrality measures) using presence data from the RFID antenna, collected

during the non-breeding periods, November–January, of 2015/16 (15 weeks) and 2016/17

(13 weeks, hereafter referred to events). An association between birds observed from

these data can reflect either, individuals that choose to maintain some social cohesion,

which we consider to be a non-random association, or individuals without pre-established

social cohesion who coincide in time and space, which we consider random mixing. These

data record the presence of birds without distinguishing between these two types of

association. Further, dyadic interactions can also be aggressive interactions, and our data

cannot exclude these cases. Although in sparrows', dominance hierarchies are typically

linear, there is no evidence for a correlation with reproductive fitness (Sánchez-Tójar

2018).

The 'gambit of the group' is a common approach used to identify discrete groups among all

associating individuals (Whitehead and Dufault 1999; Figure 1A). However, given the

gregarious nature of sparrows and the high activity at our feeder, at which non-discrete

groups of sparrows accumulated, the gambit of the group approach overestimated

associations between individuals (Figure 1A; also see Ferreira et al. 2020). One solution to

this would be to define a non-random association where two individuals overlapped by a

defined period (Δ_t) at our bird feeder. However, in our system, again due to the near

constant visitations, this resulted in linear network structures, e.g., linking the first bird to

the second, then the second to the third, and so on (Figure 1B). To account for the social

behaviour of sparrows, we derived a method to infer non-random associations that

assumed that non-random social associations are established before they attend the feeder

(suggested by Summer-Smith 1963). We therefore defined an association as two

individuals that arrived to feed within 150 seconds (Δ_t) of each other. Here an arrival is

defined as the (re)appearance of the individual at the feeder after being absent for a period

of minimally >300 seconds (Δ_l). We defined that Δ_t = 150 seconds was sufficient to detect

and link individuals who arrive together in a group (see Figure 1C), and the resulting data

better sampled non-random associations between individuals in our system, from

watching sparrows in the field and on pre-recorded footage, arriving at our feeder (Plaza et

al., 2019).

From the resulting association matrices of the two events, 2015/16 and 2016/17, we built

a series of weighted, non-directional, social networks (hereafter, network/s), where the

vertices represent individual sparrows and interconnecting edges their associations. First,

we built individual networks for each of the 15 weeks in 2015/16, and 13 weeks in

2016/17, to estimate within-individual repeatability in centrality network metrics. This

was to validate these centrality measures against the individual repeatability already

demonstrated in this system using both RFID and video data (Plaza et al. 2019). Then, we

built a network for each event to extract non-breeding sociality. Finally, we also built two

bipartite networks from each event (sub-graphs), which only considered association

strength between opposite-sex individuals.

From the first non-breeding networks we extracted three centrality measures representing

different aspects of sociality for each individual using the 'iGraph' R package (Csardi and

Nepusz 2006): We selected centrality measures to reflect aspects of an individual's social

preference, following similar studies on sociality (McDonald 2007; Farine and Whitehead

2015; Beck, Farine and Kempenaers 2021). Degree represents the number of associates,

and may impact fitness through enhanced mate choice, where individuals position

themselves alongside others of lesser quality (Oh and Badyaev 2010;), or where same-sex

associates benefit reproduction in cooperative breeding species (Bebbington et al. 2017;

Riehl and Strong 2018). Strength represents the quality of those relationships and may

influence the structure and behaviour of reproductive communities (Firth and Sheldon

2016; Culina, Firth and Hinde 2020). We calculated strength following (Farine 2013), using

the sum of dyadic Simple Ratio Indices (the association probability between a dyad, from 0,

never associated, to 1, always associated), which we transformed to give a measure of net

association quality (Eq. 1, where S denotes strength, $d_{(i)}$ and sum $SRI_{(i)}$ the degree and SRI

of a given individual respectively and N(V) = the number of vertices in a network)

$$(Eq.\,1)\quad S\,=\,d_{(i)}\frac{SRI_{(i)}}{N(V)}$$

We used eigenvector centrality (following McDonald 2007, hereafter centrality) to quantify

the influence of an individual to all others within the network (Newman 2004; Oh and

Badyaev 2010); Finally, we extracted opposite-sex degree from the two bipartite sub-

graphs, the number of opposite-sex associations, which we used to represent an

individual's pool of potential reproductive partners, as a fourth measure of sociality

(following Beck, Farine and Kempenaers 2021). Given the high density of sparrows visiting

the feeder, and the frequency at which those birds were detected we did not threshold our

networks (using only a sample of the birds attending an antenna, following Farine and

Whitehead 2015) to maintain network structure, although only birds who arrived in a dyad

(where degree > 0) were included in our networks.

Fitness measures

For each of the sparrows that survived to the following breeding period we used the

genetic pedigree to calculate two fitness measures, and for each of these, we calculated one

annual measure, and one across the lifetime of the sparrow. Both fitness measures are

based on the number of recruits, and we have shown that they correlate well with

reproductive value, thus represent fitness in an evolutionarily meaningful way (Alif et al.

2022). We defined recruits as offspring that survived and produced genetic offspring

themselves.

(1) *Recruits.* For the number of annual recruits, we summed individual recruits within the

breeding year following the social events. We then again summed individual recruits across

a lifetime, or up to 2020 as a measure of lifetime recruits. Note that the latter category only

contained five sparrows that were still alive at the point of census, and as such, our

recruitment data can be considered near complete. We excluded birds which did not

survive to breed, and yearlings which had zero recruits.

(2) *De-lifed fitness.* As a second fitness measure, we used de-lifed fitness (p_{ti}; Eq. 2), which

estimates an individual's genetic contribution to the population (Coulson et al. 2006). De-

lifed fitness is a retrospective measure of realized fitness, relative to the population each

year, calculated by removing (de-lifing) an individual and its offspring from the pedigree

and recalculating the resulting change in population growth.

$$(Eq.\,2) \quad p_{ti} = \frac{\xi_{t(i)} - w_{t(i)}}{N_t - 1}$$

Here, p_{ti} is the contribution of individual ($_i$) to population growth during a specific period t.

Further, $\xi_{t(i)}$ is a measure of individual performance, here the number of surviving

offspring of individual i at the end of the breeding period t. We added a value of one if the

individual i itself survived to the next breeding period $t + 1$. The population size at time t is

N_t at the beginning of each breeding cycle (here April). To estimate the individual's

contribution to population growth, we use w_t, which represents the ratio of the population

size at $t + 1$ to the population size at t. This de-lifed fitness p_{ti} is an annual value per

individual, and we calculated it for all birds which produced at least one recruit. We then

also summed p_{ti}, within individuals as a lifetime de-lifed fitness measure, p_i.

Individual repeatability in sociality

First, we validated our centrality measures by confirming that they were repeatable within-

individuals between weeks. We modelled degree, strength, centrality and opposite-sex

degree respectively, as response variables each in four Gaussian linear mixed models

against the intercept, and with bird identity modelled as a random effect to compensate for

repeat identities between years. We then divided the variance explained by bird identity by

the total phenotypic variance of the trait to quantify repeatability (see Nakagawa and

Schielzeth 2010). We ran repeatability models using package MCMCglmm default

parameters and priors – the models converged robustly and reliably.

Selection on sociality – annual and lifetime fitness

We quantified the association between centrality measures and fitness. As all four

centrality measures are inherently correlated, we modeled each separately to avoid

collinearity (Webster, Schneider and Vander Wal 2020). For each centrality measure, we

ran two models, with annual recruits and annual de-lifed fitness as the response variables.

In the models with annual numbers of recruits as response, we assumed a Poisson error

distribution with a log link function, and in the models explaining de-lifed fitness we

assumed a Gaussian error distribution. We mean-centered all centrality measures within

each year, eliminating between-year differences, and modelled them as fixed covariates.

We also added each sociality variable as a quadratic effect to test for stabilizing or

disruptive selection where averages are favored over the extremes (Wolf et al. 2007). Bird

identity was modelled as a random effect on the intercept to account for pseudo-

replication, and cohort to account for environmental stochasticity. We modelled fixed

effects for sex (male, 1 or female, 0) and age (in calendar years) and age as a quadratic

effect, to account for variation in fitness as explained by demography (Schroeder et al.

2012). We added sex as an interaction term with age to account for the extra-pair

behaviour of older males (Girndt et al. 2018).

We modelled lifetime recruits and lifetime de-lifed fitness in the same way as the annual

ones, but instead of age we used lifespan, or maximum age at year 2020. Because each bird

was only represented once in this dataset, we only modelled cohort as a random effect.

We used Bayesian Markov Monte-Carlo simulations, using MCMCglmm (Hadfield 2010), to

run all models. For all models we used inverse Wishart priors for random effects, and ran

each over 343,000 iterations, with a burn-in of 3,000 and a thinning interval of 200. We

visually checked the posterior trace plots for all model outputs and ensured that

autocorrelation was below 0.1 and effective sample sizes between 1,000 and 2,000. The

fixed effects were considered statistically significant when the 95% credible interval (CI) of

its posterior distribution did not span zero.

282
Null models and dominance interactions

We ran a node-permutation null-model by shuffling the identities of birds visiting the

feeder between existing arrival times in our association matrices, thereby breaking any link

between sociality and fitness, over 1000 randomized permutations (following Farine,

2017). We used these randomized association matrices to construct 1000 new networks

and extracted the mode for our four centrality measures. We used these randomized

centrality measures to re-run all fitness models. Finally, to exclude the possibility that

dominance was interacting with our observed centrality measures, we tested for

correlations between the centrality measures and dominance from videos recorded during

the same period of our social network events. We represented individual dominance by

calculating ELO ratings, based on antagonistic interactions at the bird feeder (for further

details see Sánchez-Tójar 2018). We did not include the randomised centrality measures

from our null models in these correlations.

Results

The data consisted of 150 individual birds making 410,114 visits to the RFID feeder within

our study period (mean = 2,734 visits per bird, SD = 8,116), across both events. Excluding

birds that died prior to the start of our study or those that were ringed after, 160 tagged

birds survived in our system in November 2015, an additional 90 birds were tagged prior

to the 2017 breeding period, although not all survived to sampling. After constructing the

arrival networks, we identified 3,783 associations between 118 PIT tagged individuals

during 2015/2016, and 874 associations between 69 individuals in 2016/2017. These

networks contained 66.3% of 122 and 26.3% of 205 breeding birds in 2016 and 2017

respectively. Combined, we had 130 records for annual and lifetime fitness from 102

individuals, 33 were recorded in both years (for summary statistics see Table 1). Degree

and opposite-sex degree are closely correlated, implying that those with more opposite-sex

associates also tend to have more associates of either sex (supplementary material table 1)

Individual repeatability in sociality

We confirmed individual repeatability by week in all four centrality measures between 15

weeks in 2015/16: Degree, R=0.29 (0.15 – 0.39), Strength, R=0.22 (0.10 – 0.32), Centrality,

015 (0.03 – 0.27) and, Opposite sex degree, 0.27 (0.13 – 0.4); and, 13 weeks in 2016/17:

Degree, R=0.29 (0.15 – 0.39), Strength, R=0.22 (0.10 – 0.32), Centrality, 015 (0.03 – 0.27)

and, Opposite sex degree, 0.27 (0.13 – 0.4) (Table 1).

Selection on sociality – annual and lifetime fitness

Opposite-sex degree had a statistically significant linear relationship with the number of

annual recruits and annual de-lifed fitness. Strength and centrality had a negative quadratic

association with annual recruitment (Figure 2; Table 2; Table 3). Age and sex both also

predicted annual recruits, with younger individuals and females recruiting more offspring.

Age also positively predicted annual de-lifed fitness (Table 2; Table 3).

None of our centrality measures statistically significantly predicted lifetime recruitment

(Supplementary material table 4), however, all four had a statistically significant negative

quadratic relationship with lifetime de-lifed fitness (Figure 2).

328
329

Null models and dominance interactions

We found no link between fitness and sociality, nor any evidence of selection from our null

models (supplementary material figure 3). Likewise, dominance was not strongly

correlated with any centrality measure, implying that our method of assigning associations

based on arrival time, rather than shared space at a bird feeder, is unlikely to be influenced

by dominance (supplementary material table 1).

Table 1. Summary statistics for recruitment and centrality measures for individual house sparrows on Lundy Island during

two non-breeding events (November–January 2015/2016 and 2016/2017). Each measure is given as mean, standard

deviation and sample size, and including repeatability (R) and 95% CI for centrality measures.

Summary statistics of recruitment				
Fitness Variable	Scale	2015/16 Mean, SD (n = 118)		2016/17 Mean, SD (n = 69)
Recruits	Annual	2.65, 1.98		0.78, 1.07
	Lifetime	5.34, 3.76		4.15, 3.86
Summary statistics and repeatability of centrality measures				
Centrality measure	Mean, SD	R, 95CI	Mean, SD	R, 95CI
Degree	64.11, 32.03	0.29, 0.15–0.39	25.33, 17.37	0.35, 0.16–0.49
Strength	0.06, 0.05	0.22, 0.10–0.32	0.09, 0.13	0.33, 0.16–0.48
Centrality	0.35, 0.30	0.15, 0.03–0.27	0.25, 0.31	0.23, 0.10–0.36
Opposite-sex degree	32.00, 15.86	0.27, 0.13–0.4	12.18, 9.19	0.45, 0.24–0.61

342 Table 2. Annual recruitment model outputs from GLMMs for each of our four centrality measures (Degree, Strength, Centrality
343 and Opposite-sex degree), derived of 410,114 visits to an RFID baited feeder by 150 individuals. Centrality measure of house
344 sparrows on Lundy Island, modeled against annual recruits. We inferred significance where the 95% CI did not span zero,
345 positive effects on the response variable are highlighted in red, and negative in blue.

Annual recruits	Posterior mode 95% credible intervals			
Variable	Degree	Strength	Centrality	Opposite-sex degree
Fixed terms				
(Intercept)	3.49 1.46—6.11	3.69 1.58—6.46	4.19 1.60—6.31	3.78 1.58—6.50
Centrality measure	0.17 -0.01—0.34	0.22 -0.03—0.44	0.26 -0.01—0.45	0.25 0.05—0.41
Centrality measure²	-0.1 -0.26—0.06	-0.12 -0.32—0.07	-0.16 -0.34—0.08	-0.04 -0.16—0.10
Sex (male)	-0.67 -1.41—0.05	-0.7 -1.46—-0	-0.74 -1.48—-0	-0.57 -1.47—0
Age	-1.14 -2.03—-0.30	-1.15 -2.04—-0.28	-1.27 -1.99—-0.26	-1.07 -2.00—-0.26
Age²	0.05 -0.07—0.17	0.04 -0.08—0.17	0.06 -0.08—0.16	0.03 -0.07—0.17
Age*Sex (male)	0.21 -0.01—0.48	0.22 -0.02—0.46	0.28 -0.02—0.49	0.21 0—0.49
Random terms				
*Bird ID	0 0—0.13	0 0—0.12	0 0—0.12	0 0—0.13
*Cohort	1.88 0.25—15.69	1.5 0.19—15.57	2.09 0—15.89	2.29 0.11—16.90
Residuals	0 0—0.27	0 0—0.28	0 0—0.29	0 0—0.24

347   Table 3. Lifetime recruitment model outputs from GLMMs for each of our four centrality measures (Degree, Strength,
348   Centrality and Opposite-sex degree), derived of 410,114 visits to an RFID baited feeder by 150 individuals. Centrality measure
349   of house sparrows on Lundy Island, modelled against lifetime recruits. We inferred significance where the 95% CI do not span
350   zero, positive effects on the response variable are highlighted in red, and negative in blue. († Age in lifetime models denotes
351   either lifespan, or age in 2020, whichever is greatest)

Lifetime recruits	Posterior mode, 95% credible intervals (lower – upper)			
Variable	Degree	Strength	Centrality	Opposite-sex degree
Fixed term				
(Intercept)	0.85	0.85	0.99	0.8
	0—1.93	0.02—1.95	-0.06—1.87	-0.03—1.83
Centrality measure	0.1	0.1	0.12	0.14
	-0—0.25	-0.09—0.3	-0.07—0.31	-0.05—0.25
Centrality measure^2	-0.05	-0.01	-0.07	-0.03
	-0.19—0.11	-0.24—0.11	-0.24—0.11	-0.16—0.12
Sex (male)	0.33	0.28	0.28	0.48
	-0.35—0.99	-0.36—0.91	-0.33—0.98	-0.24—1.03
†Age	0.1	0.11	0.09	0.12
	0—0.23	0—0.24	0.01—0.25	0—0.24
†Age*Sex (male)	-0.06	-0.07	-0.07	-0.06
	-0.20—0.07	-0.19—0.07	-0.2—0.06	-0.21—0.04
Random effects				
*Cohort	NA	0.47	0.45	0.39
	0.07—2.78	0.07—2.76	0.06—3	0.10—2.77
Residuals	NA	0.12	0.11	0.14
	0—0.27	0.01—0.29	0.02—0.28	0—0.26

Discussion

We found evidence for annual fitness benefits of sociality, where individuals with more

opposite-sex associates had higher fitness in the breeding period, than those with fewer,

but that this did not translate to lifetime fitness. For lifetime fitness, we found evidence for

stabilising selection on sociality, including opposite-sex degree, suggesting that such

benefits are only short-lived, or contextual, in a wild population.

We constructed our networks by linking dyads of birds that arrived together to a bird

feeder, but ignored the time that they spent there, to eliminate most random associations.

Other studies have adapted similar approaches in high-density and open feeder systems, or

have considered the same implicit problems (Gomes, Boogert and Cardoso, 2021). Ferreira

et al. (2020), for example, identified flocks arriving, but then defined associations by spatial

proximity at a series of feeder boxes. Further research could optimize our approach for

other systems, either by refining the time after which an individual is determined to have

left the feeder ($\triangle_i$), or similarly, the time it takes for all members of a group to interact with

the feeder upon arrival ($\triangle_t$). Further work may also consider defining associations only

where a dyad visit together more often than would be expected by chance but doing so

must also consider some method of retaining peripheral associations. Although RFID

systems sample sociality well at a feeder, we cannot be sure that sociality traits are

maintained in other contexts – future works might consider tracking social behaviour

across time and space (For example see Ripperger and Carter 2021).

Where previous studies on wild birds have suggested links between aspects of sociality and

annual reproductive success (for examples see Firth and Sheldon 2016; Kohn 2017; Beck,

Farine, and Kempenaers 2020), we were also able to use lifetime measures, which better

reflect the genetic contribution of the individual to population growth, and thus, fitness.

This also allowed us to also describe how selection acts upon sociality across the

population. We found that sociality had little influence on fitness at the annual scale, apart

from for opposite-sex association, which was linked to increased recruitment and de-lifed

fitness. Our study corroborates that annual fitness benefits described elsewhere,

particularly regarding mate choice (Oh and Badyaev 2010; Beck, Farine, and Kempenaers

2021, Beck, Farine, and Kempenaers 2020) directly translate into increased annual fitness.

At the lifetime scale, our study also provides some insight into the evolution of social

behaviors, which we found to be maintained at the population average through stabilizing

selection. We are therefore, to the best of our knowledge, the first study to link sociality

with lifetime fitness benefits in a wild bird (but see Formica et al. 2021; Philson and

Blumstein 2022). Our results may also suggest a mechanism for selection on sociality

through enhanced mate choice, but the impact on survival was difficult to determine in this

study. Sociality is predicted to increase survival through reduced predation risk or

information transfer (Sorato et al. 2012; Hillemann et al 2020), but we found no evidence

to suggest that either was selected for, through higher centrality, in our analyses.

Stabilizing selection in this case may be driven by factors such as high mate fidelity or

changing sociality with age (Oh and Badyaev 2010; Albery et al. 2022), removing the need

to constantly maintain opposite-sex associations over lifetime while maintaining individual

fitness. However, those opposite-sex associations may also be beneficial in an extra-pair

context from the male perspective (Beck, Farine, and Kempenaers 2020) and requires

further research.

Our centrality measures were associated with lifetime, but not with annual de-lifed fitness,

and only opposite-sex degree was associated with recruitment at the annual scale. We

found no relationship between social centrality and dominance in our study using arrival

time to define sociality, but aggressive interactions are probably also reduced over the

non-breeding period (Summer-Smith 1963). None of our centrality metrics were linked

with recruitment at the lifetime scales. Overall, de-lifed fitness better represents fitness as

it is a relative measure of the contribution to population growth (Alif et al. 2022). The

number of recruits, while intuitively appealing, is not relative, and in good years, more

birds may have a higher number of recruits, while in poor years, having one recruit may be

an achievement. As such, this measure is not always comparable between years and may

explain our results. Further, recruitment is also dependent on parental effects and

relationships within the breeding season, which were not quantified here, although they

have been suggested elsewhere (Bebbington et al. 2017; Riehl and Strong 2018), whereas

de-lifed fitness also captures long-term survival. We found that older males recruited more

offspring, likely by virtue of older males siring more extra-pair offspring (Girndt et al.

2018). Likewise, younger birds had lower annual de-lifed fitness, because younger birds

had not recruited any offspring in previous years that would contribute to their current de-

lifed fitness.

In conclusion, we suggest a link between opposite-sex association and reproductive success

at the annual scale, suggesting a mechanism for selection to shape social behaviour. At the

lifetime scale we suggest that selection on sociality is stable, suggesting greater fitness for

those at the population mean, in a wild population of passerine birds.

420
421
422
Acknowledgments

We would like to thank the Lundy Landmark trust and the Lundy Field Society for their

ongoing support, particularly Dean Jones, Rosie Ellis and Tom Carr. This research was

supported by the QMEE CDT, funded by NERC grant number NE/P012345/1 (JD), a

fellowship from the Volkswagen Foundation (JS), a grant from the German Research

Foundation: Deutsche Forschungsgemeinschaft (JS), a European Research Council grant,

CIG PCIG12-GA-2012-333096 (JS), and by NERC grant NE/J024597/1 (TB). We also thank

two anonymous reviewers for their helpful feedback.

Data availability

Analyses reported in this article can be reproduced using the data provided by Dunning,

Jamie et al. (2022), Opposite-sex associations are linked with annual fitness, but sociality is

stable over lifetime, Dryad, Dataset, https://doi.org/10.5061/dryad.z08kprrhd

Bibliography

Alif, Ž., J. Dunning, H.Y.J. Chik., T. Burke., and J Schroeder. 2022. What is the best fitness
measure in wild populations? A case study on the power of short-term fitness proxies to
predict reproductive value. *PLoS One*. 17(4):
e0260905. doi.org/10.1371/journal.pone.0260905

Andersson, M. 1994. Sexual Selection. Princeton University Press, New Jersey.

Aplin, L. M., D. R. Farine, J. Morland-Ferron, and B. C. Sheldon. 2012. Social networks predict
patch discovery in a wild population of songbirds. *Proc. R. Soc. B.* 279(1745): 4199–4205.
doi.org/10.1098/rspb.2012.1591

Aplin, L. M., D. R. Farine, J. Morand-Ferron, E. F. Cole, A. Cockburn, and B. C. Sheldon. 2013.
Individual personalities predict social behaviour in wild networks of great tits (*Parus*
major). *Ecol. Lett.* 16 (11): 1365–72. doi.org/10.1111/ele.12181.

Aplin, L. M., J. A. Firth, D. R. Farine, B. Voelkl, R. A. Crates, A. Culina, C. J. Garroway. 2015.
Consistent individual differences in the social phenotypes of wild great tits, *Parus major*.
Anim. Behav. 108: 117-127. doi.org/10.1016/j.anbehav.2015.07.016.

Aplin, L. M., D. R. Farine, R. P. Mann, and B. C. Sheldon. 2014. Individual-level personality
influences social foraging and collective behaviour in wild birds. *Proc. R. Soc. B.. Sci.* 281
(1789): 20141016. doi.org/10.1098/rspb.2014.1016.

Archie, E. A., J. Tung., M. Clark., J. Altmann., and S.C. Alberts. 2014. Social affiliation matters:
Both same-sex and opposite-sex relationships predict survival in wild female baboons.
Proc. R. Soc. B. 281(1793): 20141261. doi.org/10.1098/rspb.2014.1261

Bebbington, K., S. A. Kingma, E. A. Fairfield, H. L. Dugdale, J. Komdeur, L. G. Spurgin, and D. S.
Richardson. 2017. Kinship and familiarity mitigate costs of social conflict between
Seychelles warbler neighbors. *PNAS.* 114(43): E9036 - E9045.
doi.org/10.1073/pnas.1704350114.

Beck, K. B., M. Valcu, and B. Kempenaers. 2020. Analysis of within-individual variation in
extrapair paternity in blue tits (*Cyanistes caeruleus*) shows low repeatability and little
effect of changes in neighborhood. *Behav. Ecol.* 31(6):1303—1315.
doi.org/10.1093/beheco/araa069.

Beck, K. B., D. R. Farine, and B. Kempenaers. 2020. Winter associations predict social and
extra-pair mating patterns in a wild songbird. *Proc. R. Soc. B.* 287(1921): 20192606.
doi.org/10.1098/rspb.2019.2606.

Beck, K. B., D. R. Farine, and B. Kempenaers. 2021. Social network position predicts male
mating success in a small passerine. *Behav. Ecol.* 32(5):856 – 864.
doi.org/10.1093/beheco/arab034.

Birkhead, T. R., and J. D. Biggins. 1987. Reproductive synchrony and Extra-pair copulation
in birds. *Ethology*. 74(4): 320 - 334. doi.org/10.1111/j.1439-0310.1987.tb00942.x.

Blankenhorn, W. 2010. *The quantitative study of sexual and natural selection,* in, *Animal*
Behaviour: Evolution and Mechanisms. P. Kappeler (Ed.). Springer. Germany.

Brandl, H. B., D. R. Farine, C. Funghi, W. Schuett, and S. C. Griffith. 2019. Early-life social
environment predicts social network position in wild zebra finches. *Proc. R. Soc. B.* 286:
20182579—20182579. doi.org/10.1098/rspb.2018.2579.

Cameron, E. Z., T.H. Setsaas, and W.L. Linklater. 2019. Social bonds between unrelated
females increase reproductive success in feral horses. *PNAS*. 106(33):13850 - 13853.
doi10.1073pnas.0900639106

Carter, K. D., R. Brand, J.K. Carter, B. Shorrocks, and A.W. Goldizen. 2013. Social networks,
long-term associations and age-related sociability of wild giraffes. *Anim. Behav*. 86(5):901 -
910. doi.org/10.1016/j.anbehav.2013.08.002.

Coulson, T., T. G. Benton, P. Lundberg, S. R. X. Dall, B. E. Kendall, and J. M. Gaillard. 2006.
Estimating individual contributions to population growth: Evolutionary fitness in
ecological time. *Proc. R. Soc. B.* 273 (1586): 547–55. doi.org/10.1098/rspb.2005.3357.

Croft, D. P., R. James, and J. Krause. 2008. *Exploring Animal Social Networks*. Princeton
University Press, NJ.

Cresswell, W. 1994. Flocking is an effective anti-predation strategy in redshanks, Tringa
totanus. *Anim. Behav.* 47(2): 433 - 442. doi.org/10.1006/anbe.1994.1057.

Cresswell, W., and J. L. Quinn. 2011. Predicting the optimal prey group size from predator
hunting behaviour. *J. Anim. Ecol.* 80(2):310-319. doi.org/10.1111/j.1365-
2656.2010.01775.x.

Csardi, G., and T. Nepusz. 2006. The igraph software package for complex network
research. *Inter J. Complex Syst.* 1695(5): 1 - 9.

Culina, A., J.A. Firth, and C.A. Hinde. 2020. Familiarity Breeds Success: Pairs That Meet
Earlier Experience Increased Breeding Performance in a Wild Bird Population. *Proc. R. Soc.*
B. Proc. R. Soc. B. 287 (1941): 20201554.

Dawson, D. A., G.J. Horsburgh, A.P. Krupa, Stewart, S. Skjelseth, H. Jensen, A.D. Ball, L.G.
Spurgin, M.E. Mannarelli, S. Nakagawa, J. Schroeder, C. Vangestel, G.N. Hinten and T. Burke.
2012. Microsatellite resources for Passeridae species: A predicted microsatellite map of the
house sparrow *Passer domesticus*. *Mol. Ecol. Res*. 12(3): 501 – 523. doi:10.1111/j.1755-
0998.2012.03115.x.

Dimitriadou, S., D.P. Croft, D.P. and S.K. Darden. 2019. Divergence in social traits in
Trinidadian guppies selectively bred for high and low leadership in a cooperative
context. *Sci Rep* 9(17194):. doi.org/10.1038/s41598-019-53748-4

Dobson, F., J. Murie and V. Viblanc. 2020. Fitness Estimations for Ecological Studies: An
Evalualtion in Columbian Ground Squirrels. *Front. Ecol, Evol.* 8.
doi.org/10.3389/fevo.2020.00216

Endler, J.A., 1986. Natural Selection in the Wild. Princeton University Press, New Jersey.

Falconer, D.S. and T.F.C. Mackay. 1996. Introduction to Quantitative Genetics, Edinburgh.

Farine, D. R., and H. Whitehead. 2015. Constructing, conducting and interpreting animal
social network analysis. *J. Anim. Ecol.* 84(5):1144 - 1163. doi.org/10.1111/1365-
2656.12418.

Farine, D. R. 2017. A guide to null models for animal social network analysis. *Methods Ecol.*
Evol. 8(10): 1309 – 20. doi.org/10.1111/2041-210X.12772.

Farine, D. R., and B. C. Sheldon. 2015. Selection for territory acquisition is modulated by
social network structure in a wild songbird. *J. Evol. Biol.* 28(3): 547 - 556.
doi.org/10.1111/jeb.12587.

Farine, D. R. 2013. Animal Social Network Inference and Permutations for Ecologists in R
using asnipe. Meth. *Ecol. Evol.* 4(12): 1187 – 1194. doi/full/10.1111/2041-210X.12121.

Ferreira, A. C., R. Covas, L. R. Silva, S. C. Esteves, I. F. Duarte, R. Fortuna, F. Theron, C.
Doutrelant, and D. R. Farine. 2020. How to make methodological decisions when inferring
social networks. *Ecol. Evol.* 10 (17): 9132 – 9143. doi.org/10.1002/ece3.6568.

Firth, J. A., E. F. Cole, C. C. Ioannou, J. L. Quinn, L. M. Aplin, A. Culina, K. McMahon, and B. C.
Sheldon. 2018. Personality shapes pair bonding in a wild bird social system. *Nat. Ecol. Evol.*
2(11):1696 - 1699. doi.org/10.1038/s41559-018-0670-8.

Firth, J. A., and B. C. Sheldon. 2016. Social carry-over effects underpin trans-seasonally
linked structure in a wild bird population. *Ecol. Let.* 19(11):1324 - 1332.
doi.org/10.1111/ele.12669.

Firth, J. A., B. C. Sheldon, and D. R. Farine. 2016. Pathways of information transmission
among wild songbirds follow experimentally imposed changes in social foraging structure.
Biol. Lett. 12(6). doi.org/10.1098/rsbl.2016.0144.

Formica, V., H. Donald, H. Marti, Z. Iregebay, and E. Brodie III. 2021. Social network position
experiences more variable selection than weaponry in wild subpopulations of forked
fungus beetles. *J Anim Ecol.* 90(1): 168-182. doi.org/10.1111/1365-2656.13322

Forstmeier, W., K. Martin, E. Bolund, H. Schielzeth, and B. Kempenaers. 2011. Female
extrapair mating behavior can evolve via indirect selection on males. *PNAS.* 108(26): 10608
- 10613. doi.org/10.1073/pnas.1103195108.

Franks, D. W., G. D. Ruxton, and R. James. 2010. Sampling animal association networks with
the gambit of the group. *Behav. Ecol. Sociobiol.* 64(3): 493 - 503. doi.org/10.1007/s00265-
009-0865-8.

Girndt, A., C. W. T. Chng, T. Burke, and J. Schroeder. 2018. Male age is associated with extra
pair paternity, but not with extra-pair mating behaviour. *Sci. Rep.* 8:8378.
doi.org/10.1038/s41598-018-26649-1

Girndt, A., G. Cockburn, A. Sánchez-Tójar, M. Hertel, T. Burke, and J. Schroeder. 2019. Male
age and its association with reproductive traits in captive and wild house sparrows. *J. Evol.*
Biol. 32(12):1431 - 1443. doi.org/10.1111/jeb.13542.

Grant, P.R., and B.R. Grant. 2019. Adult sex ratio influences mate choice in Darwin's finches.
PNAS. 116(25): 12372 - 12382. doi.org/10.1073/pnas.1903838116.
562

563   Gomes, A.C.R., N.J. Boogert, G.C. Cardoso. 2021. Network structure and the optimization of
564   proximity-based association criteria. *Meth. Ecol. Evo.* 12(1):88 –
565   100. doi.org/10.1111/2041-210X.13387.
566

Hadfield, J.D. 2010. MCMC methods for multi-response generalized linear mixed models:
The MCMCglmm R package. *J. Stat. Softw.* 33 (2):1 - 22. doi.org/10.18637/jss.v033.i02.

Hillemann, F., E. F. Cole, B. C. Sheldon, and D. R. Farine. 2020. Information use in foraging
flocks of songbirds: no evidence for social transmission of patch quality. *Anim. Behav.* 165:
35–41. doi.org/10.1016/j.anbehav.2020.04.024.

Kohn, G. M. 2017. Friends give benefits: autumn social familiarity preferences predict
reproductive output. *Anim. Behav.* 132: 201 – 208. doi.org/10.1016/j.anbehav.2017.08.013.

Korsten, P., T. Van Overveld, F. Adriaensen, and E. Matthysen. 2013. Genetic integration of
local dispersal and exploratory behaviour in a wild bird. *Nat. Comm.* 4: 2362.
doi.org/10.1038/ncomms3362.

Krause, J. U., and G. D. Ruxton. 2002. Living in groups. Oxford Series in Ecology and
Evolution. Oxford: Oxford University Press.

Kruuk, L. E. B. 2004. Estimating genetic parameters in natural populations using the
"animal model." *Proc. R. Soc. B.* 359(1446):873 - 890. doi.org/10.1098/rstb.2003.1437.

Kurvers, R. H. J. M., L. Prox, D. R. Farine, C. Jongeling, and L. Snijders. 2020. Season-specific
carryover of early life associations in a monogamous bird species. *Anim. Behav.* 164:25–37.
doi.org/10.1016/j.anbehav.2020.03.016.

Lea, A.J., D.T. Blumstein, T.W. Wey, and J.G.A. Martin. 2010. Heritable victimization and the
benefits of agonistic relationships. *PNAS.* 107(50): 21587–21592.
doi.org/10.1073/pnas.1009882107.

Le Galliard, J. F., P. S. Fitze, R. Ferrière, and J. Clobert. 2005. Sex ratio bias, male aggression,
and population collapse in lizards. *PNAS.*102(50): 18231–18236.

Maldonado-Chaparro, A. A., P. O. Montiglio, W. Forstmeier, B. Kempenaers, and D. R. Farine.
2018. Linking the fine-scale social environment to mating decisions: a future direction for
the study of extra-pair paternity. *Biol. Rev.* 93(3): 1558—1577.
doi.org/10.1111/brv.12408.

Mayer, C., and G. Pasinelli. 2013. New support for an old hypothesis: Density affects extra-
pair paternity. *Ecol. Evol.* 3(3):694 - 705. doi.org/10.1002/ece3.489.

McDonald, D. B. 2007. Predicting fate from early connectivity in a social network. *PNAS.*
104(26):10910–10914. doi.org/10.1073/pnas.0701159104.

McDonald, G.C., N. Engel, S. S. Ratão, T. Székely, and A. Kosztolányi. 2020. The impact of
social structure on breeding strategies in an island bird. *Sci. Rep.* 10(1): 13872.
doi.org/10.1038/s41598-020-70595-w.

Moiron, M., Charmantier, A. and Bouwhuis, S. 2022. The quantitative genetics of fitness sin
a wild seabird. *Evolution*. Doi:10.1111/evo.14516 [online ahead of print]

Nakagawa, S., and H. Schielzeth. 2010. Repeatability for Gaussian and non-Gaussian data: a
practical guide for biologists. *Biol. Rev*. 85(4): 935 - 956. 10.1111/J.1469-
185x.2010.00141.X
605
Oh, K. P., and A. V. Badyaev. 2010. Structure of social networks in a passerine bird:
Consequences for sexual selection and the evolution of mating strategies. *Am. Nat.* 176(3):
E80—E89. doi.org/10.1086/655216.

Philson, C.S., Blumstein, D.T. 2022. Group social structure has limited impact on
reproductive success in a wild mammal. *Behav. Ecol.*
arac102, https://doi.org/10.1093/beheco/arac102

Plaza, M., T. Burke, T. Cox, A. F. Carroll, A. Girndt, G. Halford, D. A. Martin, et al. 2019. Social
network node-based metrics can function as proxies for animal personality traits. *J. Evol.*
Biol. 33(11): 1634 - 1642. doi.org/10.32942/OSF.IO/YVQ9D.

Proops, L., C. A. Troisi, T. K. Kleinhappel, and T. Romero. 2021. Non ⍰ random associations
in group housed rats (*Rattus norvegicus*). *Sci. Rep.* 11(1):15349. doi.org/10.1038/s41598-
021-94608-4.

Psorakis, I., S. J. Roberts, I. Rezek, and B. C. Sheldon. 2012. Inferring social network
structure in ecological systems from spatiotemporal data streams. *J. R. Soc. Interface.* 9(76):
3055 – 3066. doi.org/10.1098/rsif.2012.0223.

Psorakis, I., B. Voelkl, C. J. Garroway, R. Radersma, L. M. Aplin, R. A. Crates, A. Culina, D. R.
Farine, J. A. Firth, C. A. Hinde, L. R. Kidd, N. D. Milligan, S. J. Roberts, B. Verhelst and B. C.
Sheldon . 2015. Inferring social structure from temporal data. *Behav. Ecol. Sociobiol.* 69: 857
- 866. doi.org/10.1007/s00265-015-1906-0.

R Core Team. 2022. R: A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical Computing. http//www.R-project.org/.

Reid, J.M., Nietlisbach, P., Wolak, M.E., Keller, L.F. and Arcese, P. 2019. Individuals expected
genetic contributions to future generations, reproductive value, and short-term metrics of
fitness in free-living song sparrows (*Melospiza meolodia*). *Evolution Letters.* 3(3):271 – 285.
doi/pdf/10.1002/evl3.118

Riehl, C., and M. J. Strong. 2018. Stable social relationships between unrelated females
increase individual fitness in a cooperative bird. *Proc. R. Soc. B.* 285(20180130).
doi.org/10.1098/rspb.2018.0130.

Ripperger, S.P and Carter, G.C. 2021. Social Foraging in Vampire Bats Is Predicted by Long-
Term Cooperative Relationships. *PLoS Biology* 19 (9):e3001366.

Sánchez-Tójar, A., I. Winney, A. Girndt, M. J. P. Simons, S. Nakagawa, T. Burke, and J.
Schroeder. 2017. Winter territory prospecting is associated with life-history stage but not
activity in a passerine. *J. Avian Biol.* 48(3):407 - 416. doi.org/10.1111/jav.01055.

Sánchez-Tójar, A., Nakagawa, S., Sánchez-Fortún, M., Martin, D. A., Ramani, S., Girndt, A.,
Bókony, V., Kempenaers, B., Liker, A., Westneat, D. F., Burke, T., and Schroeder, J. 2018.
Meta-analysis challenges a textbook example of status signalling and demonstrates
publication bias. *ELife.* 7: e37385. doi.org/10.7554/eLife.37385
643
644 Sánchez-Tójar, A. 2018. *The evolution of social dominance in house sparrows*. http://nbn-
645 resolving.de/urn:nbn:de:bsz:352-2-9850l4uzuvk43
646
Schroeder, J., I. R. Cleasby, S. Nakagawa, N. Ockendon, and T. Burke. 2011. No evidence for
adverse effects on fitness of fitting passive integrated transponders (PITs) in wild house
sparrows *Passer domesticus. J. Avian Biol.*42(3):271 - 275. doi.org/10.1111/j.1600-
048X.2010.05271.x.

Schroeder, J., S. Nakagawa, I. R. Cleasby, and T. Burke. 2012. Passerine birds breeding under
chronic noise experience reduced fitness. *PLoS One*. 7(7): E39200.
doi.org/10.1371/journal.pone.0039200.

Schroeder, J., T. Burke, M.E. Mannarelli, D.A. Dawson, and S. Nakagawa. 2012A. Maternal
effects and heritability of annual productivity. *J. Evol. Biol.* 24(1):149 - 156. doi:
10.1111/j.1420-9101.2011.02412.x.
657

Schroeder, J., S. Nakagawa, M. Rees, M. E. Mannarelli, and T. Burke. 2015. Reduced fitness in
progeny from old parents in a natural population. *PNAS.* 112(13): 4021 - 4025.
doi.org/10.1073/pnas.1422715112.

Schroeder, J., Y.H. Hsu, I. Winney,M. Simons, S. Nakagawa, and T. Burke. 2016. Predictably
philandering females prompt poor paternal provisioning. *Am. Nat.* 180(2): 219 – 230.
doi.org/10.1086/687243.
664
665   Silk, J. B., S.C. Alberts, and J. Altmann. 2003. Social Bonds of Female Baboons Enhance Infant
666   Survival. *Science, 302*(5648): 1227–1231. doi.org/10.1126/science.1088474
667
Silk, J. B. 2007. The adaptive value of sociality in mammalian groups. *Philos. Trans. R. Soc.*
*Biol. Sci.*362(1480): 539 - 59. doi.org/10.1098/rstb.2006.1994.

Silk, J. B., J. C. Beehner, T. J. Bergman, C. Crockford, A. L. Engh, L. R. Moscovice, R. M. Wittig,
R. M. Seyfarth, and D. L. Cheney. 2009. The benefits of social capital: Close social bonds
among female baboons enhance offspring survival. *Proc. R. Soc. B..* 276 (1670): 3099–3104.
doi.org/10.1098/rspb.2009.0681.

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., Wittig, R.
M., Seyfarth, R. M., and Cheney, D. L. 2010. Strong and consistent social bonds enhance the
longevity of female baboons. *Curr. Biol. 20*(15): 1359–1361.
doi.org/10.1016/j.cub.2010.05.067.
678
Silk, M. J., D. P. Croft, T. Tregenza, and S. Bearhop. 2014. The importance of fission-fusion
social group dynamics in birds. *Ibis.* 156(4):701 - 715. doi.org/10.1111/ibi.12191.

Simons, M. J. P., I. Winney, S. Nakagawa, T. Burke, and J. Schroeder. 2015. Limited catching
bias in a wild population of birds with near-complete census information. *Ecol. Evol.*
5(16):3500 - 3506. doi.org/10.1002/ece3.1623.

Sorato, E., P.R, Gullet, S.C. Griffith, and A.F. Russell. 2012. Effects of Predation Risk on
Foraging Behaviour and Group Size: Adaptations in a Social Cooperative Species. *Animal*
Behaviour 84 (4): 823–34.

Summer-Smith, D. 1963. *The House Sparrow*. Collins. UK.

Stanton, M. A., and J. Mann. 2012. Early Social Networks Predict Survival in Wild Bottlenose
Dolphins. *PLoS One*. 7(10):E47508. doi.org/10.1371/journal.pone.0047508.

Strickland, K., D. J. Mitchell, C. Delmé, and C. H. Frère. 2021. Repeatability and heritability of
social reaction norms in a wild agamid lizard. *Evolution.* 75(8): 1953–65.
doi.org/10.1111/evo.14298.

Thompson, N. A., and Cords, M. 2018. Stronger social bonds do not always predict greater
longevity in a gregarious primate. *Ecol. Evl. 8*(3): 1604–1614. doi.org/10.1002/ece3.3781
695

Thys, B., M. Eens, S. Aerts, A. Delory, A. Iserbyt, and R. Pinxten. 2017. Exploration and
sociability in a highly gregarious bird are repeatable across seasons and in the long-term
but are unrelated. *Anim. Behav.* 123: 339–48: doi.org/10.1016/j.anbehav.2016.11.014.

Voelkl, B., J. A. Firth, and B. C. Sheldon. 2016. Nonlethal predator effects on the turn-over of
wild bird flocks. *Sci. Rep.* 6: 33476. doi.org/10.1038/srep33476.

Webber, Q. M. R., D. C. Schneider and E. Vander Wal. 2020. Is less more? A commentary on
the practice of 'metric hacking' in animal social network analysis. *Anim. Behav.*, *168*: 109–
120. doi.org/10.1016/j.anbehav.2020.08.011.

Westneat, D. F., Bókony, V., Burke, T., Chastel, O., Jensen, H., Kvalnes, T., Lendvai, Á. Z., Liker,
A., Mock, D., Schroeder, J., Schwagmeyer, P. L., Sorci, G., & Stewart, I. R. K. 2014. Multiple
aspects of plasticity in clutch size vary among populations of a globally distributed
songbird. *J. Anim. Ecol. 83*(4): 876–887. doi.org/10.1111/1365-2656.12191
708
Wey, T., D. T. Blumstein, W. Shen, and F. Jordán. 2008. Social network analysis of animal
behaviour: a promising tool for the study of sociality. *Anim. Behav.* 75(2): 333 – 44.
doi.org/10.1016/j.anbehav.2007.06.020.

Whitehead, H., and S. Dufault. 1999. Techniques for Analyzing Vertebrate Social Structure
Using Identified Individuals: Review and Recommendations. *Adv. Study Behav.* 28: 33 - 74.
doi.org/10.1016/S0065-3454(08)60215-6.

Wolf, M., G.S. van Doorn, O. Leimar, F.J. Weissing. 2007. Life-history trade-offs favour the
evolution of animal personalities. *Nature*, 447(7144): 581 - 584.
717
Youngblood, M. 2019. A Raspberry Pi-based, RFID-equipped birdfeeder for the remote
monitoring of wild bird populations. *Ring. Migr. 34*(1): 25 – 32.
doi.org/10.1080/03078698.2019.1759908

Figure 1. Three versions of a simulated event (A, B and C) show the interval over which five individuals (1–5, black/grey bars)

spent at a resource over time (t), and the derived social networks from each approach: A = gambit of the group, which links all

individuals in a discrete group equally; B = time-window overlap (by Δ_t), which links individuals who overlap at a resource;

and C = arrival time (developed for this study), which links individuals who arrive together to a resource. Red boxes denote

the time period during which individuals are considered to be associated, and overlaps represent an association: A, all

individuals within a group; B, where they are physically present at the same time (red box), or shortly after they depart to

account for birds which were present, but not currently being recorded by the antenna, in that case, over-lapping by Δ_t (red

over-hanging box, typically a few seconds); or, C, where they arrive within Δ_t of each other, but the subsequent time spent at a

feeder is irrelevant. However, note that the function of Δ_t differs between B and C; Where in B, Δ_t functions to detect when

birds are in the same place but where one (or more) are not currently being detected by the antenna, in C the function is to

link all individuals which arrive together, while ignoring those already present at the resource, which has the potential to link

two separate groups in A and B. In the case of C, an additional interval (Δ_i) is required to define when birds have left the

resource, after which they can be recorded as arriving again.

Figure 2. De-lifed fitness as response variables against Centrality measures from 8 linear mixed models, at two scales, from the

Lundy Island house sparrows, derived of 410,114 visits to an RFID baited feeder by 150 individuals: Explanatory variables for

Annual de-lifed fitness (A); and Lifetime de-lifed fitness (B), where [2] denotes a quadratic function, also shown in the four

adjacent panels for A and B, and their 95% credible intervals. Credible intervals are given as solid bars for each explanatory

variable, where a solid point denotes the posterior mode. Black bars denote no effect on the response variable; red denote a

positive and blue, a negative, relationship with the response. In adjacent panels, quadratic functions of each response variable

presented in A and B (on the Y axis: A Centrality, Degree, Opp. Degree, Strength, and B Centrality, Degree, Opp. Degree,

Strength). Blue curves represent a negative interaction with fitness measures (on the X axis). Measures with no effect are not

shown in figure. We found no link to sex, and age was also subject to stabilising selection, (given in supplementary material

table 4).