Isolated Grauer's gorilla populations differ in diet and gut

microbiome

Alice Michel[1,2], Riana Minocher[1,7], Peter Niehoff[1], Yuhong Li[1,8], Kevin Nota[3], Maya A. Gadhvi[1],

Jiancheng Su[1], Neetha Iyer[2], Amy Porter[2], Urbain Ngobobo-As-Ibungu[4], Escobar Binyinyi[4], Radar

Nishuli Pekeyake[5], Laura Parducci[3,9], Damien Caillaud[2], Katerina Guschanski[1,6]

[1]Animal Ecology, Department of Ecology and Genetics, Evolutionary Biology Centrum, Uppsala

University, Sweden

[2]Department of Anthropology, University of California, Davis, CA, USA

[3]Plant Ecology, Department of Ecology and Genetics, Evolutionary Biology Centrum, Uppsala

University, Sweden

[4]The Dian Fossey Gorilla Fund International, Kinshasa, DRC

[5]Institut Congolais pour la Conservation de la Nature, Kinshasa, DRC

[6]Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, UK

[7]Department of Human Behaviour, Ecology and Culture, Max-Planck Institute for Evolutionary

Anthropology, Leipzig, Germany

[8]Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences, University of

Groningen, The Netherlands

[9]Department of Environmental Biology, Sapienza University of Rome, 00185 Rome, Italy

Short running title: Diet and gut microbiome of Grauer's gorillas

Correspondence:

Alice Michel: aljmichel@ucdavis.edu; Katerina Guschanski: katerina.guschanski@ed.ac.uk

Keywords: Metabarcoding, 16s rRNA, *trn*L, critically endangered, genetic diversity, faecal DNA

Abstract

Conservation efforts tend to focus on populations that are genetically differentiated without paying attention to their ecological differentiation. However, isolated populations may be ecologically unique, an important aspect for the design of appropriate conservation measures for endangered species. Here we investigate the interplay between diet and gut microbiome in several geographically isolated and genetically differentiated populations of the critically endangered Grauer's gorilla. We find that dietary and gut microbial profiles are population-specific, likely due to geographic isolation and environmental differences. In addition, social groups within each population also differed in diet and, to a lesser extent, in gut microbial composition and diversity. Individuals at low elevation consumed a larger variety of plant taxa than those at high elevation, consistent with the notion that dietary choice is constrained by food availability that changes with elevation. Despite no detectable correlation between the diet and gut microbiome in richness or evenness, dietary and gut microbial composition covaried significantly. As we did not find evidence for an effect of genetic relatedness on the composition of the gut microbiome of Grauer's gorillas, this pattern is likely a result of long-term social, ecological, and geographic factors acting on both diet and microbiome. These results reveal that isolated and genetically distinct populations of Grauer's gorillas are also ecologically distinct, highlighting the need to dedicate separate conservation efforts for each population.

Introduction

Biodiversity world-wide is declining rapidly due to habitat loss, overexploitation, and climate change (Ceballos et al., 2020; Haddad et al., 2015; IUCN, 2021; Ripple et al., 2019). The aim of conservation is to prevent this decline and to preserve unique species and populations (DeWoody et al., 2021; Teixeira & Huber, 2021). Across a species' range, variation in environmental conditions combined with restricted gene flow can lead to population differentiation, due to both neutral (drift) and adaptive (selection) processes. From a conservation perspective, such differentiated populations are important, as they contain unique traits that could be lost from the species' pool of variation if the individual populations were to disappear (Cadotte et al., 2011). Therefore, effective conservation requires investigating the ecology and genetics of populations. This necessitates a holistic approach.

Non-invasive faecal sampling is a powerful tool for genetic studies of elusive animals (Beja-Pereira et al., 2009; Schwartz et al., 2007). In addition to DNA from the individual itself, faecal DNA also contains clues to its ecology, including diet and gut microbiome (Valentini et al., 2009). Food consumed reflects individual feeding behaviour and health, and can be informative about foraging strategies, ranging patterns, and social and cultural behaviour (Kamil et al., 2012). Further, diet can reflect local adaptation, if populations specialise due to local availability of different foods or different foraging behaviours and cultures (Alberdi et al., 2020).

Tightly linked to diet, the gut microbiome impacts long-term health and behaviour of its host, by aiding digestion, modulating dietary choice, and stimulating the immune system (Agranyoni et al., 2021; Colston & Jackson, 2016; Davidson et al., 2020; Ley et al., 2008; Moran et al., 2019). Isolated populations in different habitats have been shown to possess unique gut microbiomes, which is likely a result of ecological and genetic differences (Bueno de Mesquita et al., 2021; Couch et al., 2020; Fietz et al., 2018). The gut microbiome may even facilitate local adaptation and drive genomic evolution (Alberdi et al., 2016; Rudman et al., 2019). It is therefore important to consider gut microbiome when evaluating conservation needs and values (Trevelline et al., 2019; West et al., 2019).

Here, we jointly investigate dietary and gut microbial diversity and composition in isolated and genetically differentiated populations of Grauer's gorillas (*Gorilla beringei graueri*) using DNA metabarcoding. This gorilla subspecies is endemic to the eastern Democratic Republic of Congo (DRC)

and is classified as critically endangered due to recent drastic population decline (IUCN, 2016; Plumptre

et al., 2016, 2021). Our study populations occupy the low and high altitudinal extremes of the species'

range (600-700m, 800m and 2100-2400m), with limited connectivity and gene flow between

populations (Baas et al., 2018; Plumptre et al., 2021). We hypothesized that ecological differences

would cause diet and gut microbiome to differ between these isolated populations, co-varying with host

genetic differentiation. In addition, we investigated whether the taxonomic composition of the gut

microbiome is influenced by the dietary profile and examined microbiome stability in individuals

through time.

Materials and Methods

Ethics Statement

This study was conducted in compliance with legal requirements of the DRC and the animal use policies of UC Davis. Data collection protocols were approved by Institut Congolais pour la Conservation de la Nature. Samples were collected non-invasively, without approaching or disturbing wild animals.

Sample collection

Faecal samples (n=220) were opportunistically collected from night nests of Grauer's gorillas in eastern DRC between 2015 and 2018 at three sites: Kahuzi-Biega National Park (2.32˚S, 28.72˚E; KBNP; 2500 m above sea level [asl]), Nkuba Conservation Area in Walikale territory, North Kivu (1.38˚S, 27.47˚E; NCA; 600 m asl), and Maiko National Park (0.87˚S, 27.35˚E; MNP; 830 m asl; **Figure 1**). In KBNP, gorillas in the Chimanuka group were habituated to human presence and samples were collected from identified individuals immediately upon defecation. All other samples were collected from night nests without knowledge of individual identity following the two-step collection method (Nsubuga et al., 2004). Geographic location and altitude were recorded using handheld GPS for all sampling sites except for the Mankoto group in KBNP, for which this information is missing. We assigned age classes in the field based on dung diameter, as follows: infant < 4cm, sharing a nest with an adult; juvenile/subadult <5cm, own nest; and adults >5cm (McNeilage et al., 2006; Schaller, 1963). For the Chimanuka group, age classes of identified individuals were known from observations. We included biological replicates for several individuals for genotyping and gut microbiome analyses.

DNA extraction

Faecal samples were exported to Uppsala University, Sweden, for molecular analysis. DNA was extracted from 50 mg of dried material using the DNeasy PowerSoil DNA Extraction Kit (Qiagen)

in a dedicated primate faecal extraction laboratory. We implemented the following modifications to the

manufacturer's protocol: faecal samples were incubated under shaking (500 RPM) in the C1 solution

overnight at 23°C. They were then transferred into a heating block and incubated at 65°C for 10 minutes,

followed by bead beating on a vortex at maximum speed for 1 hour. Incubation in C2 and C3 solution

was on ice. We incubated the samples in C6 solution at room temperature for 5 minutes before elution.

Gorilla genotyping, individual identification, relatedness and population differentiation analyses

We genotyped all 220 samples at 12 microsatellite loci (vWF, D1s550, D4s1627, D5s1457,

D5s1470, D6s474, D6s1056, D7s817, D8s1106, D10s1432, D14s306, and D16s2624) following the

two-step multiplex protocol (Arandjelovic et al., 2009) and sexed them with the amelogenin assay

(Bradley et al., 2001). PCR products were run on an agarose gel to confirm amplification success and

absence of contamination in blanks. Up to four loci were pooled, based on fluorophores and product

sizes, and run on the ABI GeneAnalyzer (ThermoFisher Scientific). We scored genotypes manually in

GeneMapper v5.0 (Chatterji & Pachter, 2006) and used Cervus v3.0.7 (Kalinowski et al., 2007) to

identify individuals. Samples were considered from the same individual if their genotypes had no

mismatches and matched at five or more loci, with the probability of identity assuming full-sibling

relationship (PIDsib) less than 0.05. We manually generated consensus individual genotypes from

matching samples, taking into account their time and place of sample collection, and evidence about

the presence of other individuals from the same group.

We tested for deviations from Hardy-Weinberg equilibrium, heterozygote deficiency, and

linkage disequilibrium at each locus in GenePop v4.7.5 (Raymond & Rousset, 1995; Rousset, 2008).

Genetic population structure was assessed using STRUCTURE v2.3.4 (Porras-Hurtado et al., 2013)

with 20 independent runs for K = 1-11 (corresponding to the eleven social groups), an 100,000-iteration

burn-in, and data collection for 1,000,000 runs, assuming population admixture and correlated allele

frequencies (Falush et al., 2003). Results from different runs of K were merged in CLUMPP (Jakobsson

& Rosenberg, 2007; Kopelman et al., 2015), and analysed and visualized in 'pophelper' in R (Francis,

2017; R Core Team, 2021). The most likely value of K was determined using ΔK (Evanno et al., 2005).

We used the 'adegenet' R package for Principle Component Analysis (PCA) based on individual

genotypes (Jombart, 2008). Population differentiation statistics F_{ST} and F'_{ST} (Meirmans & Hedrick,

2011) were calculated in GenoDive v3.04 (Meirmans, 2020), and significance assessed with 9999

permutations. We compared genetic relatedness between populations and social groups using an

AMOVA in the R package 'poppr' (Kamvar et al., 2014) and calculated pairwise relatedness (*r*)

between all individuals within KBNP and NCA separately in ML-Relate (Kalinowski et al., 2006).

Characterization of gorilla diet

We characterized the diet of 92 unique individuals identified by genotyping (see Results), from

nine social groups and two lone silverbacks (**Table S1, S2**). Where possible, we selected a single nest

site per group, to maximize the number of unique individuals. When multiple samples per individual

were available, we chose the sample that performed best in genotyping as indication of quality.

We amplified the P6 loop of the *trn*L chloroplast intron (Taberlet et al., 2007), a locus that has

been successfully used for dietary metabarcoding in primates, and for which a large database of tropical

plants is available (Mallott, Garber, et al., 2018). We used the standard *trn*L g and h primers (**Table

S3**), tagged with 96 eight-base-pair (bp) barcodes. Each barcode differed from all others at a minimum

of three positions. DNA amplifications were carried out in 20 μl reactions containing 2 μl faecal DNA

extract, 1 U Platinum II *Taq* Hot-Start DNA polymerase, 1x Platinum II Buffer, 0.2 mM each dNTP, 2

mM $MgCl_2$, and 1 μM each primer. Each DNA sample was amplified twice. The duplicates were placed

randomly on different PCR plates to avoid potential batch effects and biases due to cross-contamination

of sample and/or barcoded primer (**Table S1**). We included one PCR negative and two to three empty

wells per plate, to check for contamination during PCR (Taberlet et al., 2018). In addition, we included

five DNA extraction blanks. PCR conditions consisted of 2 minutes denaturation at 94˚C followed by

35 cycles of 94˚C for 30 seconds, 51˚C for 30 seconds, and 68˚C for 15 seconds, without final extension.

PCR products were checked on a 2% agarose gel to confirm amplification without contamination.

The barcoded PCR products were pooled column-wise (16 μl for each sample, duplicates in

separate pools), mixed with 640 μl PB Buffer, and purified using MinElute columns (Qiagen, The

Netherlands), eluting in 50 µl EB buffer. Double-indexed next-generation sequencing libraries (Kircher

et al., 2012) were prepared as detailed (Brealey et al., 2020; Rohland et al., 2015) but using not-barcoded

incomplete adapters after blunt-end repair. Two library preparation blanks were carried through all

steps. Each pool was quantified using qPCR with PreHyb primers (**Table S3**; Rohland et al., 2015) and

amplification settings as in Brealey et al. (2020).

Each sample pool and both library blanks received a unique combination of indices (**Table S1**).

For indexing PCR, we used the same reaction mixture and cycling conditions as Brealey et al. (2020).

The number of cycles ranged from 8 to 10, depending on the copy number estimated from qPCR (**Table

S1**). Library preparation blanks were amplified for 10 cycles to maximize capture of potential

contaminants. We performed MinElute purification and quantified indexed pools with qPCR, as above,

using i7 and i5 primers (Rohland et al., 2015, **Table S3**). Indexed sample pools were combined in

equimolar amounts, except for library preparation blanks, of which we added 0.5 µl each into the final

pool, corresponding to the lowest amount added for any sample. The final sequencing pool was cleaned

using AmPure XP beads (Beckman Coulter, USA) with two elutions (0.5x followed by 1.8x), which

remove very large fragments and fragments <100bp, respectively. This size selection is optimized for

the retention of *trn*L amplicons (~10-150bp in length + 148 bp of barcoded and indexed adapters).

Elution was performed in 30 µl of EB buffer. The cleaned library pool was quantified using both a

Qubit High Sensitivity fluorometer and 2200 TapeStation and sequenced at the Uppsala Science for

Life Laboratory on a single MiSeq lane with 150 bp paired-end sequencing with version 2 chemistry.

Sequence processing and analysis was done in OBITools v1.2.13 (Boyer et al., 2016). Paired

reads with quality scores >40 and overlap >10 bp were retained and merged. Sample of origin for each

read was established through its index and barcode, requiring an exact sequence match. Sequences were

clustered into molecular operational taxonomic units (MOTUs), each representing a unique plant taxon

(Valentini et al., 2009). A large number of MOTUs had fewer than 10 sequences across all samples and

were removed as recommended (*e.g.*, Shehzad et al., 2012). We also removed sequences that differed

by exactly one nucleotide from a more abundant sequence and had a total count less than 5% of the

more abundant sequence, following Boyer et al. (2016).

Finally, taxonomic assignment used a custom-made reference database (below). Based on a

frequency plot of identity to the reference database (**Figure S1**) and similar *trn*L-based studies of

tropical primate diet (*e.g.,* Quémére et al., 2013), we removed sequences below an identity threshold

of 0.90. Below this, sequences were regarded as likely chimeric, enriched in sequencing or PCR

errors. No singletons were present after this filtering step.

Compiling plant trn*L reference database*

We built a local DNA barcoding reference library by downloading all 324,502 available

sequences from NCBI GenBank using the search query: "(trnL[All Fields] OR complete genome[All

Fields]) AND (plants[filter] AND (chloroplast[filter] OR plastid[filter]))" (last accessed 2 December

2021). In OBITools v1.2.13, the sequence list was annotated with taxonomy information downloaded

from NCBI (ftp://ftp.ncbi.nih.gov/pub/taxonomy/taxdump.tar.gz, last accessed 3 December 2021). To

complete the database of *trn*L genes, we followed established protocol (Boyer et al., 2016), using the

same *trn*L g-h primers as in the wet laboratory to extract *trn*L variants *in silico* in the program ecoPCR

v2.1 (Ficetola et al., 2010). We kept sequences that were between 10 to 230 base pairs long with at most

three primer mismatches total (Taberlet et al., 2018). The final database contained 21,308 *trn*L *in silico*

amplicons, in 608 families and 5,662 genera.

To evaluate the resolution of our reference database with respect to local plant diversity, we

compared plant taxa present in our database to a list of plants known to occur in the Kahuzi and Itebero

regions of KBNP (Yumoto et al., 1994). To enable this comparison, we updated the taxonomic

classification of the KBNP plant list (Yumoto et al., 1994) by searching for species names in the Global

Biodiversity Information Facility (GBIF). The updated list contained 328 taxa, in 81 unique families

and 234 genera. Of these, all families and 77.4% of genera were present in our *trn*L database.

Characterization of gorilla gut microbiome

We determined gut microbial composition of 86 samples belonging to 68 individuals from

KBNP and NCA populations (**Table S2**). (Samples from MNP were not yet available at the time of

microbiome characterization.) The dataset included multiple samples from 15 individuals (2-3 each),

collected 0-257 days apart (mean = 27 days, median = 0), which we used to evaluate intra- versus inter-

individual variation in microbiome composition. We also carried nine random extraction blanks through

the entire data generation process, on par with samples.

The V4 region of the *16S rRNA* gene was amplified with primers 515F/806R (**Table S3**) for

each sample in duplicate. The PCR reaction contained 2 µl of extracted DNA, 5 µM each of the forward

and reverse primer, 1x Phusion High-Fidelity Buffer, 0.02 units Phusion HF DNA polymerase (2U/µl),

0.012 mg DMSO and 0.05 µM (each) dNTPs, with the volume made up to 20 µl with Ultrapure H2O.

Thermal cycling conditions were as follows: 30 seconds at 98°C, 25 cycles of 98°C for 10 seconds,

52°C for 20 seconds and 72°C for 20 seconds, and 10 minutes at 72°C. PCR cycles were limited to 25

to minimize the risk of unspecified products and chimeras. Duplicate reactions were pooled and cleaned

with AmPure beads (Qiagen).

Next-generation sequencing libraries were prepared from PCR products following the double-

barcoding, double-indexing strategy (Kircher et al., 2012; Meyer & Kircher, 2010; Rohland et al., 2015;

van der Valk et al., 2017). As a result, each sample had a unique combination of two barcodes and two

indices, which enabled bioinformatic filtering of potential chimeric molecules and misassigned reads

resulting from index hopping (van der Valk et al., 2017, 2020). For indexing, we determined the suitable

number of PCR cycles (8-11) based on qPCR of barcoded libraries, as above. Indexed libraries were

quantified by qPCR and pooled in equimolar amounts for sequencing on a single MiSeq lane, using

version 2 chemistry and 250 bp paired end sequencing at the Uppsala Science for Life Laboratory

sequencing facility.

Sequencing reads were demultiplexed and adapters removed using a Python script (Brealey et

al., 2021). We followed established protocol to estimate microbial amplicon sequence variants (ASVs)

using DADA2 (Callahan et al., 2016), rather than clustering sequences, which avoids biases due to

arbitrary similarity thresholds (Edgar, 2018). Forward and reverse reads were truncated to 200 and 150

bp, respectively, at which point read quality scores dropped below 35. We merged paired-end reads,

requiring an overlap of at least 12 bp, and removed sequences outside the 250-256 bp range and those

with any barcode mismatch, as recommended (Callahan et al., 2016).

Taxonomy was assigned using the SILVA 132 reference database, released in December 2017

(Quast et al., 2012). Species-level assignment required a strict 100% match (Edgar, 2018). We removed

singletons and ASVs labelled 'Unassigned', 'Eukaryota', "mitochondria", or "chloroplast". We

retained Archaea, although archaeal amplification from the V4 region of the *16S rRNA* is limited

(Raymann et al., 2017), because within-dataset comparisons are nonetheless informative. We built a

bacterial phylogenetic tree by aligning sequences to the Greengenes 13_5 mega-phylogeny (203,452

99% OTUs; DeSantis et al., 2006) in SEPP using default parameters (Mirarab et al., 2012).

Statistical analyses of trn*L and* 16S *datasets*

To examine dietary and microbiome diversity, we analysed the *trn*L and *16S rRNA*

metabarcoding datasets, after first filtering out rare sequence variants below 0.5% relative abundance

in at least one sample, as suggested (Deagle et al., 2019). We evaluated sampling effort and sequencing

depth accumulation curves in the R packages 'vegan' (Oksanen et al., 2020) and 'ranacapa' (Kandlikar

et al., 2018), respectively. We checked whether the predicted number of taxa (asymptote of the

sequencing accumulation curve) minus actual number of taxa (richness) related to any of the considered

biological variables or sequencing depth (read count) using a generalised linear model (GLM) with

quasi-Poisson error distribution in the R package 'lme4' (Bates et al., 2015).

We calculated two alpha diversity metrics for each dataset: richness, or the number of taxa, and

Shannon's diversity index, or evenness (Chao et al., 2014). As recommended by McMurdie & Holmes

(2013), we did not rarefy to minimum sequencing depth. To test the effects of population, social group,

altitude, sex, and age class on diversity metrics, we fitted a GLM with quasi-Poisson (for richness) or

gamma (for evenness) error distribution with log link, followed by Tukey honestly significant

difference (HSD) *post-hoc* comparisons between levels of categorical variables that were overall

significant (χ^2 test with Bonferroni correction) (Lenth et al., 2021) .

To assess trends in diet and microbiome beta diversity, or composition, we followed a strategy

designed for the compositional nature of metabarcoding data (Gloor et al., 2017; Weiss et al., 2017).

We used Bayesian multiplicative zero replacement and then centred and log-ratio (CLR) transformed

each dataset using the R packages 'zcompositions' (Palarea-Albaladejo & Martín-Fernández, 2015) and

'compositions' (van den Boogaart & Tolosana-Delgado, 2008). For the microbiome dataset, we

secondarily used Phylogenetic Isometric Log-Ratio Transform (phILR) to compute compositional

abundance at phylogenetic balances (Silverman et al., 2017). To evaluate variation in composition of

diet and microbiome, we computed Aitchison's dissimilarity (Euclidean distance between CLR values)

(Aitchison et al., 2000). To quantitatively estimate which factors best predict variation in diet and gut

microbiome, we modelled the composition in CLR (or phILR) transformed space as a function of

ecological and biological variables using PERMANOVA, via function *adonis2* in 'vegan' (Anderson

& Walsh, 2013). The predictor variables were population, social group, sex, age class, and altitude.

Sequencing read count was kept as the first predictor, even if $p > 0.05$. *Post-hoc* comparisons between

levels of overall significant variables were done with Bonferroni correction using 'pairwiseAdonis'

(Arbizu, 2020). We assessed the temporal stability of individual microbiomes using a pairwise

Wilcoxon test between Aitchison dissimilarities of replicate and unique individuals. The influence of

genetic distance (*1 - genetic relatedness*) was modelled separately within each population using Mantel

and partial Mantel tests (controlling for social group identity).

We estimated the covariance between diet and microbiome using a co-inertia analysis between

the two matrices in the package 'omicade4' and calculated the RV coefficient (Escoufier, 1973; Robert

& Escoufier, 1976) and its significance using a Monte Carlo test with 999 permutations (Meng et al.,

2014). To compare the effects of diet and other variables on the gut microbiome, we fit a Multiple

Regression on Matrices (MRM) model (Lichstein, 2007), an extension of the partial Mantel test, in

'ecodist' (Goslee & Urban, 2007). The explanatory variables were straight-line geographic distance,

altitude difference, diet composition (Aitchison distance), and social group and population as binary

(same, 0, or different, 1). Significance was assessed using 999 permutations of the response variable,

the Aitchison distance matrix of gut microbiome composition.

301   Differences in beta diversity can be due to differential abundance of a few key organisms, or

302 subtle differences across an entire community. To identify dietary and microbial taxa that may have

303 driven compositional differences, we used the R package 'ALDEx2' and focused on significant

304 differences (Wilcoxon rank sum test with correction for false discovery rate (FDR) $p < 0.05$) with effect

305 sizes >1, as recommended (Gloor et al., 2017).

Results

Study populations of Grauer's gorillas are genetically differentiated

We identified 92 unique individuals in the three study populations by microsatellite genotyping: 59 in KBNP, 28 in NCA, and 5 in MNP (**Figure 1A, Table S2**). Individuals belonged to six different social groups in KBNP, two in NCA, and one in MNP. We sampled four to 17 individuals from each social group. In addition, there were two solitary adult male gorillas (lone silverbacks) in KBNP.

Social group		N (f,m)	Year	Month	Altitude (m)	$N_{genotypes}$
🟡 Bonane		1 (0,1)	2016	June	2328	1
🟢 Chimanuka		14 (4,9)	2016	April-June	2300	44
🟢 Mankoto	Kahuzi-Biega NP	12 (1,11)	2016	July	NA	13
🔵 Mpungwe		8 (6,2)	2016	June	2200	21
🟣 Mufanzala 2		4 (3,1)	2016	July	2443	5
🟣 Mugaruka		1 (0,1)	2016	June	2123	2
🟣 Namadiriri		13 (8,5)	2016	June	2174	23
🟤 Nouvelle Famille		6 (5,1)	2016	July	2282	16
🟥 Bansamba		11 (9,2)	2015	June	619	
🟥 Bansamba	Nkuba-Biruwe	3 (1,2)	2016	April	630	46
🟧 Bansamba		3 (0,3)	2018	July	675	
🟩 Membe 1		11 (5,6)	2017	August	650	33
🔺 Maiko	Maiko	5 (3,2)	2016	November	835	6

Figure 1. Geographic location and genetic differentiation of Grauer's gorilla populations. (**A**) Broad geographic location of Grauer's gorilla populations in Maiko National Park (MNP; designated with cyan on the left-hand side of the map), Nkuba Conservation Area (NCA; green) and Kahuzi-Biega National Park (KBNP; brown). Multiple social groups were collected in NCA and KBNP. Each social group is given its own symbol colour, whereas repeated sampling of the same social group is shown by multiple identical or, for the group sampled in multiple years, similarly coloured symbols. The inset (**B**) zooms in on KBNP. Circle colours designate

different social groups, as shown in the table (C). Note that multiple circles are present for the Chimanuka group, consistent with repeated, opportunistic sampling of identified individuals. Geographic coordinates were not available for Mankoto group. (**C**) Overview of sample size for dietary characterization (N=number of unique individuals), sex composition (f=number of females, m=number of males), collection year and month, altitude and the total number of successfully genotyped samples from each group. Gut microbiome data is not available for groups shaded in grey. Group Bansamba (NCA) was sampled repeatedly, but only few samples were included in dietary analyses in later years (3 in 2016 and 3 in 2018). (D) PCA of genetic distances among individuals, coloured by population. KBNP is clearly separated from NCA and MNP.

None of the 12 microsatellite loci deviated from Hardy-Weinberg equilibrium after Bonferroni correction for multiple testing ($p > 0.1$). On average, there were 6.1 alleles per locus. The average observed and expected heterozygosities were 0.66 and 0.68, respectively. The test for global heterozygote deficiency was not significant overall ($p = 0.6$) or in any population ($p > 0.4$). The test for genotypic linkage disequilibrium using log likelihood ratio statistic with 66 pairwise comparisons between the 12 loci was not significant for any pair ($p > 0.1$). Thus, we assumed linkage equilibrium and considered all loci in further analyses.

Analysis of the three populations using STRUCTURE (Porras-Hurtado et al., 2013) revealed two distinct genetic groups (optimal K=2 according to ΔK; Evanno et al., 2005; **Figure S2**). The clusters differentiated gorillas in high-altitude KBNP from those in low-altitude NCA and MNP (**Figure S3**). Only three KBNP individuals (from two different social groups) showed a higher genetic similarity to the low altitude populations. This population structure was consistent with the PCA, which divides gorilla genotypes in KBNP from NCA and MNP (**Figure 1B**). All three populations were significantly differentiated from one another ($p < 0.001$), with largest differences between MNP and KBNP (F'_{ST} = 0.45; **Table S4A**), which are furthest apart geographically (215km). Individuals within social groups were more closely related than individuals in different groups in the same population (AMOVA ϕ = 0.12, $p < 0.001$; **Table S4B**), consistent with gorilla social structure (Harcourt & Stewart, 2013).

Negative controls in trn*L and* 16S rRNA *metabarcoding*

Metagenomic studies rarely report treatment of negative control samples, although stringent quantification of contamination is of utmost importance for any molecular study. To quantify contamination in the diet (*trn*L) dataset, we included and analysed five DNA extraction blanks, two

PCR negative controls, five unused barcode combinations, all sequenced in duplicates, and two library preparation negative controls (**Table S2**; **Table S5**). There were 286 *trn*L reads belonging to 19 different taxa in the negative controls in total. With two exceptions, the detected taxa showed very low abundance in the blanks (1-3 reads) but were highly abundant in the samples (average > 200,000 reads). This pattern is consistent with low-level cross-contamination from high quantity into low quantity samples typical for large-scale sequencing studies (Eisenhofer et al., 2019). The two exceptions with high abundance in blanks (77% and 15% of all blank reads, respectively) had low abundance in the samples (<0.01%) and were in the families Fagaceae (chestnut) and Rutaceae (citrus). We thus concluded that they may derive from reagent contamination. Both taxa were automatically removed from the dataset during abundance-based filtering, as they were below the specified 0.5% cut-off. Zero reads were identified with unused barcode combinations, suggesting that cross-contamination of barcodes during PCR and library preparation was negligible (Taberlet et al., 2018).

In the microbiome (*16S rRNA*) dataset, five out of nine DNA extraction blanks (**Table S2**) contained zero reads. The remaining four blanks had six to 72 reads each, in ten different *16S* ASVs (**Table S6**). These ten taxa were among the most abundant in the samples and thus, as above, likely represent cases of low-level cross-contamination. We found no indication of ASVs exclusive to the blanks, and hence all ASVs were retained in the dataset.

Diet of Grauer's gorillas

We characterised the diet of 92 Grauer's gorilla individuals using the chloroplast *trn*L *P6 loop* locus. After data filtering, we retained 536,160 *trn*L sequencing reads, 45% of raw reads, belonging to 120 unique taxa (**Table S7A**). PCR duplicates were more similar to each other than to other samples in alpha and beta diversity ($p < 0.001$, **Figure S4**), and hence their sequencing data were pooled. Sample accumulation curves suggested that additional samples would uncover only few novel *trn*L variants (**Figure S5A**), except in MNP, where only five individuals were sampled. Although deeper sequencing could have potentially uncovered novel taxa (**Figure S5B**), sequencing depth did not have a strong effect on *trn*L richness (three additional taxa per 47,000 reads, $p = 0.03$). Furthermore, the discrepancy

per sample in observed minus predicted richness did not corelate with sequencing depth ($p = 0.3$, nor

with any of the tested variables (population $p = 0.9$; social group $p = 0.06$; sex $p = 0.6$; age class $p =$

0.8; altitude $p = 0.4$). Taken together, this suggests that our dataset is suitable to carry out comparisons

across study populations.

Of the 120 detected taxa, 115 could be identified to at least the order level (in 29 different

orders), 110 to family (in 49 families), and 44 to genus (in 35 genera) (**Table S8**). Based on previous

studies of Grauer's gorilla diet in KBNP, NCA, and Mt. Tshiaberimu (Kambale, 2018; van der Hoek,

Pazo, et al., 2021; Yamagiwa et al., 1994, 2005; Yumoto et al., 1994) and plant distribution in the region

(Spira et al., 2018), we manually confirmed that all taxa assignments were possible. This revealed 21

taxa that have not been previously reported to be consumed by gorillas but occur in the study region

(**Table S8**).

Each Grauer's gorilla faecal sample contained 36 – 80 *trn*L taxa (mean 58.52 ± 10.83) (**Figure

2A**). The most abundant taxon across the dataset was *Urera* sp. (MOTU1, 25.6% of all reads), followed

by Apocynaceae sp. (MOTU2, 14.8%), Apocynoideae sp. (MOTU5, 6.7%), Urticaceae sp. (MOTU6,

4.6%), and *Maesa lanceolata* (MOTU9, 4.3%) (**Table S8**). The five most abundant plant families were

Urticaceae (31.9%), Apocynaceae (23.0%), Moraceae (7.1%), Myristicaceae (5.0%), and Primulaceae

(4.4%).

400
Figure 2. (**A**) Plants consumed by Grauer's gorillas in KBNP, NCA, and MNP, with the 15 most abundant taxa
across all samples indicated in the legend on the right. Populations are designated with coloured bars below

(Maiko NP cyan, Nkuba Conservation Area green, Kahuzi-Biega NP brown). (**B**) Hierarchical cluster dendrogram
of Ward's sum of squares based on minimum variance of squared dissimilarities (Murtagh & Legendre, 2014) of
centred-log-ratio (CLR) transformed taxon abundance. Relative taxon abundance clustered by population (aligned
to the bars below (A)) and by social group, shown as coloured tips in the dendrogram (see legend to right of (C)).
(**C**) Non-metric multidimensional scaling (NMDS) ordination of *trn*L CLR-transformed Aitchison distances,
representing dietary composition dissimilarity across KBNP, NCA, and MNP. Groupings by population (shape)
and social group (colours) were significant (PERMANOVA p < 0.001, R^2 = 0.28, 0.22, respectively).

Geography, altitude and social group identity influence dietary diversity and composition in Grauer's

gorillas

Dietary richness and evenness differed significantly by population and social group identity (p

< 0.001, **Table S9**). Both richness and evenness were higher in low altitude populations (NCA and

MNP) than in high-altitude KBNP (mean richness: 66.8 taxa in MNP, 65.6 in NCA vs. 54.5 in KBNP,

evenness: 10.0 in MNP, 8.4 in NCA, vs. 5.4 in KBNP, p < 0.001; **Figure S6**). Within the KBNP

population, the diets of social groups Namadiriri and Nouvelle Famille were richer than other groups

(>62 vs. <53 plant taxa), and the diets of groups Namadiriri and Mankoto were more even (p < 0.01;

Table S9; **Figure S6**). One of these groups, Namadiriri, was sampled at lower elevation than any other

social group in KBNP. In this population, one standard deviation of altitudinal increase (84 m) from the

mean (2250m asl) corresponded in a 6% decrease in richness and 25% decrease in dietary evenness (p

< 0.001; **Figure S7**). In contrast, neither individual's sex nor age class (age class available for N=70)

had an effect on dietary richness or evenness (p > 0.3).

Hierarchical clustering of dietary composition first separated high altitude (KBNP) from low

altitude (NCA and MNP) locations (**Figure 2B**). Within populations, individuals clustered together by

social group. NMDS ordination showed a similar pattern (**Figure 2C**). After accounting for slight

differences in sequencing depth, dietary composition was significantly influenced by population (p <

0.001), which explained 27.9% of the variance, and social group (p < 0.001), which explained an

additional 21.6%, but not sex (p = 0.7) nor age class (p = 0.2; **Table 1**). All between-group *post-hoc*

comparisons were significant (p < 0.05), suggesting that the group Chimanuka was not an outlier,

though it was sampled from April-May 2016 rather than June-July 2016 like other KBNP groups.

Table 1. PERMANOVA model of factors influencing dietary composition

Variable	Df	R^2	F	p	Post-hoc tests	
					Significant pairwise comparisons	$p_{Bonferroni}$
Read Count	1	0.01	1.73	0.05	–	
Population	2	0.279	17.80	<0.001	NCA – KBNP	< 0.001
					KBNP – MNP	< 0.001
					MNP – NCA	< 0.001
Social Group	6	0.216	4.56	<0.001	Chimanuka – Nouvelle Famille	0.02
					Chimanuka – Mankoto	0.005
					Chimanuka – Mpungwe	0.005
					Chimanuka - Mufanzala2	0.02
					Chimanuka – Namadiriri	0.005
					Nouvelle Famille – Mankoto	0.02
					Nouvelle Famille – Mpungwe	0.02
					Nouvelle Famille – Namadiriri	0.005
				KBNP	Mankoto – Mpungwe	0.005
					Mankoto - Mufanzala2	0.03
					Mankoto – Namadiriri	0.005
					Mpungwe – Namadiriri	0.005
					Mufanzala2 – Namadiriri	0.02
				NCA	Membe1 – Bansamba	0.005
Sex	1	0.005	0.82	0.6	–	
Age class†	2	0.022	1.28	0.1	–	

†Age class (Infants (N=7), Juveniles/subadults (N=21), Adults (N=42)) was modelled separately using a reduced dataset, since only 70/92 samples had age estimates. In this model the other predictor variables had similar estimates, so for all other variables the model for the complete dataset (n=92) is presented.

We identified differentially abundant taxa by population and social group using ALDEx2 (**Figure 3**, **Table S10**). Eight taxa were enriched in KBNP relative to MNP and NCA (Wilcoxon test with FDR < 0.05) and were considered important contributors to population dietary uniqueness (effect size > 1). These were dominated by the family Urticaceae, especially the genus *Urera*, and also included *Piper sp.* (Piperaceae), *Maesa lanceolata* (Primulaceae), and taxa in Apocynaceae and Solanaceae. In NCA, seven taxa including *Megaphrynium, Marantochloa, Aframomum, Palisota,* and Myristicaceae spp., were enriched relative to the other populations. We found 14 enriched taxa in MNP, notably Apocynaceae spp. Out of the 21 previously undescribed food items, 13 were more common in low-altitude populations in comparison to the high-altitude population. These taxa included *Gnetum* sp., more abundant in NCA relative to KBNP and MNP (MOTU34, effect = 0.93) and Humiriaceae sp., enriched in both NCA and MNP relative to KBNP (MOTU48, effect = 0.79) (**Table S8, S9**). Within

451   populations, each social group consumed twelve differentiating taxa on average (range = 0 – 30, median

452   = 8).

453
Figure 3. Grauer's gorilla diet across three study populations. Differentially abundant taxa are coloured according
to the population in which they are significantly more abundant (ALDEx2 Wilcoxon test $p < 0.05$). For taxa that
differ between two or more population pairs, the colour corresponds to the population where the effect size is
greatest. Gray taxa do not differ significantly in abundance between populations. Branch lengths do not reflect
phylogenetic distance. This diagram was generated with the 'metacoder' package in R (Foster et al., 2017).

Gut microbiome of Grauer's gorillas in Kahuzi-Biega National Park and Nkuba Conservation Area

We sequenced the *V4* region of *16S rRNA* to characterize the microbiome of 88 samples from

a subset of individuals for which we had dietary data (n=70, **Figure 1A**; **Table S2**). Two samples had

low read count (5 and 348, compared to other samples: mean=43,611 ± 11,357) and were thus excluded.

Our dataset contained multiple individuals (n=15) that were sampled repeatedly (2-3 times each),

allowing us to assess individual gut microbiome stability over time. Microbiome composition was more

similar in samples from the same individual collected zero to 257 days apart (mean compositional

dissimilarity = 35.0) than between different individuals from the same social group sampled zero to 231

days apart (Wilcoxon Test *p* < 0.001). Therefore, for further analysis we focused on the 68 unique

individuals and picked the same faecal samples used for dietary analyses. The final dataset contained

2,965,516 reads in 421 unique microbial taxa (**Table S11**).

The sample accumulation analyses suggested that additional sampling of faeces from more

individuals could uncover novel gut commensals at the population level (**Figure S8A**). However, per

sample sequencing depth was sufficient to obtain a good representation of host microbiome diversity

(**Figure S8B**). We detected 16 different phyla and 48 different families of microorganisms in the gut

microbiome of Grauer's gorillas. All taxa were identified at least to the family level; 309 taxa were

refined further to genus and 17 to species (**Table S11**). None were closely related to dominant soil

microorganisms (Delgado-Baquerizo et al., 2018). There were seven Archaea in our dataset, from the

Methanomethylophilaceae and Methanobacteriaceae families. Each faecal sample contained on average

200.29 ± 19.6 unique taxa (min = 160, max = 237), each with average abundance 0.2% ± 0.4%. The

most abundant taxa were *Christensenellaceae* R-7 group sp. (3.07% per sample), *Flexilinea* sp.

(2.97%), and *Faecalibacterium* sp. (2.46%). Eleven taxa were present in every gorilla faecal sample,

constituting a "core microbiome" of Grauer's gorillas (**Table 2**).

At a broader taxonomic scale, the average Grauer's gorilla gut microbiome was dominated by

the phyla Firmicutes (64.7%), Bacteroidetes (21.0%), Spirochaetes (3.8%), Chloroflexi (3.0%),

Proteobacteria (2.9%), and Actinobacteria (1.9%) (**Figure S9**). The most abundant families were

Ruminococcaceae (25.1%), Lachnospiraceae (14.8%), and Rikenellaceae (13.1%) (**Figure 4A**).

Table 2. Grauer's gorilla core microbiome, taxa present in 100% of samples.

Core Microbiome	Abundance†	Taxonomy to genus
ASV1	2.97%	Bacteria; Chloroflexi; Anaerolineae; Anaerolineales; Anaerolineaceae; *Flexilinea*
ASV3	2.41%	Bacteria; Firmicutes; Clostridia; Clostridiales; Family XIII; *AD3011 group*
ASV6	2.13%	Bacteria; Firmicutes; Erysipelotrichia; Erysipelotrichales; Erysipelotrichaceae; *UCG-004*
ASV22	0.94%	Bacteria; Bacteroidetes; Bacteroidia; Bacteroidales; Prevotellaceae; *Prevotella 7*
ASV21	0.87%	Bacteria; Firmicutes; Clostridia; Clostridiales; Ruminococcaceae; *UCG-005*
ASV31	0.85%	Bacteria; Firmicutes; Clostridia; Clostridiales; Lachnospiraceae; *Oribacterium*
ASV30	0.82%	Bacteria; Proteobacteria; Gammaproteobacteria; Betaproteobacteriales; Burkholderiaceae; *Sutterella*
ASV33	0.74%	Bacteria; Firmicutes; Clostridia; Clostridiales; Ruminococcaceae; *Ruminiclostridium 9*
ASV59	0.41%	Bacteria; Firmicutes; Clostridia; Clostridiales; Ruminococcaceae; *UCG-002*
ASV70	0.35%	Bacteria; Actinobacteria; Coriobacteriia; Coriobacteriales; Eggerthellaceae; *Senegalimassilia*
ASV152	0.15%	Bacteria; Bacteroidetes; Bacteroidia; Bacteroidales; Prevotellaceae; NA

†Abundance per sample, averaged over the entire dataset (N=68).

Diversity and composition of the gut microbiome in Grauer's gorillas correlates with population and

social group identity

Opposite to observations for diet, gut microbiome richness and evenness were higher in the

high-elevation population (richness: mean KBNP = 202.2 taxa vs. NCA = 190.2; p = 0.05; evenness:

83.7 vs. 74.2, p = 0.01), though neither diversity metric was related to altitude within KBNP (p = 0.07,

0.9; **Table S12**; **Figure S10**). Microbial richness did not differ by sex (p = 0.2), but females had higher

microbial evenness than males (85.7 vs. 78.5, p = 0.002). There were no differences by age class (p =

0.3, 0.1). In KBNP, as with diet, the gorilla group Namadiriri had the highest microbial richness and

evenness (**Table S12**). Neither gut microbial richness nor evenness were directly related to dietary

richness or evenness (p > 0.2), but dietary composition, scaled to one-dimension by NMDS, did

correlate with richness (p = 0.002; **Table S12**), a result we investigated further in the following section.

The composition of individual gut microbiome differed between the two populations (KBNP

and NCA) (**Figure 4B-D**). Population explained 10.5% of the total variance, and social group in KBNP

explained an additional 17.8% (p < 0.001; **Table 3**). Intergroup differences were not only due to

Chimanuka group (**Table 3**). Overall, gut microbiome dissimilarity was largest between individuals of

different populations, followed by individuals from different social groups, and individuals sharing a

social group were least dissimilar (**Figure S11**). We modelled altitude separately, as it was distinct for each population and social group (**Figure 1**). It explained 12.7% of the variance across populations (N=56, $p < 0.001$) but only 3.8% in KBNP (N=45, $p = 0.01$). Host genetics did not explain variance in gut microbiome composition in KBNP when social group was considered as a factor ($\rho = 0.015$, p = 0.3). Despite low sample size, for the one social group in NCA (N=11), we also found no correlation between gut microbiome and genetic distance ($\rho = -0.080$, p = 0.7). Microbiome composition did not differ by sex or age class ($p > 0.05$, **Table 3**). Results using phylogeny-informed (phILR) distances were qualitatively similar (**Table S13**).

We identified 111 taxa that differed significantly in abundance between NCA and KBNP ($p < 0.05$), of which 42 had effect size greater than one (**Table S14**). At the family-level, gorilla gut microbiomes in KBNP had a higher abundance of Muribaculaceae and Erysipelotrichaceae, while gut microbiomes in NCA had more Spirochaetaceae and Christensenellaceae. At a finer phylogenetic level, populations differed in abundance of specific ASVs belonging to common, shared families like Rikenellaceae, Lachnospiraceae, and Ruminococcaceae.

Table 3. PERMANOVA model of factors influencing microbiome composition

Variable	Df	R^2	F	p	Post-hoc tests Significant pairwise comparisons	$p_{Bonferroni}$
Read Count	1	0.019	1.68	0.05	–	
Population	1	0.105	7.97	<0.001	KBNP – NCA	< 0.001
Social Group	5	0.178	2.18	<0.001	Chimanuka – Nouvelle Famille	0.004
					Chimanuka – Mankoto	0.01
					Chimanuka – Mpungwe	0.007
					Chimanuka – Namadiriri	0.004
					Nouvelle Famille – Mankoto	0.02
					Nouvelle Famille – Namadiriri	0.01
					Mankoto – Mpungwe	0.007
					Mankoto – Namadiriri	0.004
Sex	1	0.014	1.18	0.2	–	
Age class†	2	0.031	1.22	0.2	–	

†Age class (Infants (N=3), Juveniles/subadults (N=21), Adults (N=38)) was modelled separately using a reduced dataset, since only 62/68 samples had age estimates. In this model the other predictor variables had similar estimates, so for all other variables the model for the complete dataset (n=68) is presented.

Figure 4. Gut microbiome composition at the family level (**A**) and differentiation based on compositional
abundances of microbial ASVs (**B-D**). Gut microbiome clusters by population in CLR Aitchison distances
dendrogram based on Ward's clustering criterion (Murtagh and Legendre 2014) (B), CLR Aitchison distances
using NMDS (PERMANOVA $R^2 = 0.14$, p = 0.001) (C), and phILR Euclidean distances using NMDS ($R^2 = 0.15$, p = 0.001) (D).

Dietary and microbial profiles are correlated with each other

Although neither dietary richness nor evenness directly correlated with gut microbial richness

or evenness ($\rho < 0.2$, $p = 0.2$; **Figure 5A**), compositional differences in dietary and gut microbial

profiles showed significant co-inertia (RV = 0.557, $p < 0.001$; **Figure 5B**) and were correlated ($\rho =$

0.32, $p < 0.001$; **Figure 5C**). Microbiome composition is known to change with diet in individuals and

also differs between populations with different dietary preferences (*e.g.,* Muegge et al., 2011;

Youngblut et al., 2019). To investigate whether dietary differences across our study populations could

explain differences in gut microbial composition or, alternatively, if external factors influenced diet and

microbiome independently, we compared the effects of diet composition and geographical and social

variables on microbiome composition using an MRM model (**Table 4**). Population and social group

were the only predictor variables significantly correlated with microbiome composition, while neither

dietary composition nor geography had an effect after accounting for social group and population of

origin.

Table 4. MRM model comparing the effects of geography, diet, and sociodemographic factors on Grauer's gorilla
gut microbiome composition†

N	Explanatory variables	Spearman's ρ	p	R²	F-statistic
	Geographic distance	0.07	0.4		
	Altitudinal difference	-0.08	0.6		
56	Diet composition†	-0.19	0.07	0.278	98.35
	Population	**0.64**	**< 0.001**		
	Social group	**0.45**	**< 0.001**		

550 † Microbiome and diet composition in Aitchison distances.

551
552 **Figure 5.** Relationship between diet and gut microbiome. (**A**) Microbiome and dietary richness, assessed as per-
553 sample total sequence count, are not correlated (p = 0.2). (**B**) High multiple co-inertia (MCIA) between
554 microbiome and diet composition in CLR-transformed space with Aitchison distance (RV = 55.7%, MC p < 0.001
555 based on 999 permutations). (**C**) Compositional differences (Aitchison distances) in diet and microbiome between
556 samples (i.e., individual gorillas) are correlated in matrix regression.
557

Discussion

We jointly characterized genetic and dietary profiles of three isolated populations of Grauer's gorillas, one of which has never been studied before (Maiko NP). For two of these populations (Nkuba Conservation Area and Kahuzi-Biega NP), we also characterized the gut microbiome. These complementary datasets enabled direct evaluation of the role of social factors, host relatedness, and diet on gut microbial diversity and composition, thus allowing us to speculate on the importance of ecological differentiation for geographically isolated populations. Whether such a relationship exists and how it is determined is a central topic in ecology (Amato et al., 2019; Bolnick et al., 2014; Bueno de Mesquita et al., 2021; Gomez et al., 2016; Hicks et al., 2018; Muegge et al., 2011). We find that diet and microbiome profiles are correlated and differ between the study populations, likely as a result of habitat differences and social factors. Habitat- or behavioural-driven differentiation in dietary and gut microbial composition, concurrent with genetic differentiation, might be an indicator of processes relevant to local adaptation (Alberdi et al., 2016, 2020; Fietz et al., 2018; Henry et al., 2019; Kolodny et al., 2020; Moeller & Sanders, 2020), which is of particular relevance to the design of effective conservation policy for these critically endangered apes.

Molecular analyses complement field observations of Grauer's gorilla diet

In addition to traditional methods based on direct observations and feeding traces, molecular approaches provide further opportunities to detect rare dietary items, revealing a more complete picture of cryptic and rare species' nutritional ecology (Ait Baamrane et al., 2012; Bradley et al., 2007; Goldberg et al., 2020; Hibert et al., 2013; Mallott, Garber, et al., 2018; Nawaz et al., 2019; Osman et al., 2020; Quéméré et al., 2013; Valentini et al., 2009). The approach is scalable to many individuals, is not biased towards habituated or approachable individuals, and provides data which can be compared across species (Kartzinel et al., 2019). Importantly, analysis of faecal samples minimizes disturbance and avoids influencing animal behaviour, aligning data collection with conservation goals (Jewell, 2013).

Nevertheless, as every method, molecular dietary analyses are not perfect. For instance, while

Grauer's gorillas predominantly feed on vegetative plants, they occasionally consume insects and fungi

(van der Hoek, Pazo, et al., 2021; Yamagiwa et al., 1991). Yet by choosing a chloroplast gene, *trn*L,

we restricted dietary characterization in this study to plants only. There are also additional technical

limitations. By relying on a reference database for identification of plant taxa, metabarcoding

approaches are necessarily limited by the content of these databases (Taberlet et al., 2018). Similarly,

chosen thresholds for sequence identity and relative abundance could impact results. We used a

sequence identity threshold of 90%, similar to other studies of tropical mammal diets via *trn*L, which

use thresholds between 90-95% (Hibert et al., 2013; Quéméré et al., 2013; Srivathsan et al., 2016). We

also set a relative abundance threshold to greater than 0.05% in at least one sample, to filter out potential

sequencing and PCR errors, chimeric sequences, etc. It is possible, particularly due to the small size

and high variability of the *trn*L locus, that our cut-offs removed some genuine dietary items.

Additionally, estimated abundances of the different plant taxa may not accurately reflect dietary

abundances (Deagle et al., 2019). DNA copy number can be biased by plant tissue type (*i.e.,* fruit, pith,

leaves, the latter of which contain more chloroplasts; Egea et al., 2010), relative digestibility (*i.e.,*

amount of fibre), and PCR amplification success (reviewed by Deagle et al., 2019). Some of these issues

also arise with other approaches. For example, accuracy of macroscopic faecal analysis depends on the

types of tissues consumed and the extent of digestion (King & Schoenecker, 2019), while observational

studies can overestimate the dietary importance of foods with longer handling times (Yamagiwa et al.,

2013).

Due to these limitations, it is important to establish how well the molecular results correspond

to traditional dietary analyses. The most abundant plant families detected in our study – Urticaceae,

Apocynaceae, Moraceae, Myristicaceae, and Primulaceae – were reported to be commonly consumed

by Grauer's gorillas in KBNP, based on feeding observations and faeces dissections (Yamagiwa et al.,

2005), and in NCA, based on feeding traces (van der Hoek, Pazo, et al., 2021). Vice versa, we detected

all top plant families reported in these observation-based studies, suggesting good correspondence

between our metabarcoding results and the literature on gorilla diet (van der Hoek, Pazo, et al., 2021;

Yamagiwa et al., 2005).

It is noteworthy that we have detected plants from which gorillas are known to primarily

consume either vegetative parts (*e.g.,* Basellaceae, Oleaceae, Poaceae in KBNP and Myristicaceae,

Marantaceae, Arecaceae in NCA) or fruits (*e.g.,* Moraceae, Euphorbiaceae, Sapindaceae, Piperaceae,

Primulaceae in KBNP and Vitaceae, Moraceae, Apocynaceae in NCA). This suggests that the *trn*L

approach can capture both vegetation and fruits in the diet. In concordance with previous investigations

of the diet of gorillas in KBNP (Yamagiwa et al., 2005), *Urera* sp. was the most common taxon in our

study (average abundance 21%). It seems likely that this taxon corresponds to *Urera hypselodendron*,

which gorillas in KBNP ate on 95% of days (Yamagiwa et al., 2005). However, *Urera* bark is very

fibrous, and an alternative explanation is that it preserves well in faeces and is thus easily detected.

On average, we detected 59 plant taxa per sample, more than the maximum number of plant

species Grauer's gorillas were reported to eat in one day in KBNP (17; Yamagiwa *et al.* 2005). It is

possible that gorillas have more complex diets than focal follows, direct observations (usually limited

to a few hours per day), or macroscopic faecal inspections suggest. Gorilla retention time can range

from 24 to 60 hours in zoos (Remis, 2000), and therefore varied feeding or movement through varied

habitats over several days could partially explain this discrepancy. Overall, the number of molecularly

identified plant taxa (120) is similar to previous studies, which reported 116 food plants in KBNP

(Yamagiwa et al., 2005) and over 100 in NCA (van der Hoek et al. 2021).

Divergent dietary ecology between isolated populations

We found that the dietary profiles of gorillas differed significantly by population and by social

group, with a unique suite of plants characterizing each population. Moreover, gorillas in NCA and

MNP tended to have more species-rich diets compared to KBNP. These two sites are 1500 m lower in

elevation than KBNP. Even within the KBNP population, we observed the highest dietary diversity in

the social group sampled at the lowest elevation (Namadiriri, 100 m below other social groups).

Globally, plant diversity tends to decrease with elevation (Rahbek, 1995), and this trend occurs within

KBNP too (Imani et al., 2016). Such an altitudinal diversity gradient is relevant for mountain gorillas,

who consume fewer species and spend more time feeding on herbaceous vegetation at higher elevation

(Ganas et al., 2004). Taken together, our data thus support the theory that feeding in eastern gorillas is

at least partially determined by plant availability, with larger dietary repertoire at lower elevations.

Another plausible interpretation is that gorillas in low altitude NCA and MNP have greater daily ranges.

This could be driven by the search for fruit, suggested in NCA (van der Hoek, Binyinyi, et al., 2021).

Or, gorillas may travel further to acquire sufficient nutrition due to potential degradation of habitat in

the low-elevation sites, relative to protected KBNP (Maldonado et al., 2012). Such increased mobility

could place individuals at even greater risk to poaching.

We identified 21 taxa that have not previously been reported in the Grauer's gorilla diet. Some

of these occur in KBNP (*e.g.*, various Apocynaceae tribes, non-native Cactaceae, and Laurales like

Ocotea usambarensis; Spira et al., 2018; Yumoto et al., 1994), and some are eaten by mountain gorillas

(*e.g., Solanum,* Solanoideae; Rothman et al., 2014; Watts, 1984) or western lowland gorillas (*e.g.,*

Beilschmiedia, Laruales; Remis et al., 2001). Others, such as *Gnetum sp.* and Humiriaceae sp., have not

been recorded in KBNP, but are western lowland gorilla foods (Rogers et al., 2004; Takenoshita &

Yamagiwa, 2008). These taxa were significantly more common in the understudied NCA and MNP

sites, suggesting that gorillas incorporate locally available plants into their diet. Another possibility is

that these taxa are environmental "contaminants", such as building material of night nests from which

samples were collected.

Other studies have documented the importance of season in gorilla diet, as gorillas seasonally

increase fruit consumption (van der Hoek, Pazo, et al., 2021; Yamagiwa & Basabose, 2006). Our ability

to evaluate the effect of season was limited by sampling. For example, gorillas in MNP had a

compositionally distinct diet, and these samples were collected later in the year. Still, the two

populations sampled at similar timepoints, NCA and most of KBNP, also diverged in diet and

microbiome. It is thus possible that the divergences that we observed are partially due to seasonal

factors, but further sampling is necessary to elucidate the effect of season.

The gut microbiome of Grauer's gorillas resembles that of other gorillas

In general, the microbial families and phyla dominating the Grauer's gorilla gut microbiome

were similar to those reported in other great apes (top phyla: Firmicutes, Bacteroidetes, Spirochaetes,

Chloroflexi, Actinobacteria, Proteobacteria; Campbell et al., 2020; Gomez et al., 2016; Hicks et al.,

2018; Nishida & Ochman, 2019). Like other gorilla subspecies, the abundance of Chloroflexi and

Spirochaetes in the wild Grauer's gorilla gut microbiome is strikingly high compared to Old World

monkeys and humans (Gomez et al. 2016, Hicks et al. 2018, Campbell et al. 2020). However, there

were differences in the relative abundance of each of the major phyla compared to wild western lowland

gorillas, with the Grauer's gorilla gut microbiomes showing proportionally fewer Actinobacteria and

Chloroflexi, but more Firmicutes, Bacteroidetes, and Spirochaetes (Campbell et al., 2020; Fontsere et

al., 2021; Hicks et al., 2018). Specifically, the relative abundance of Firmicutes in Grauer's gorilla

microbiomes was higher (64.7% average) compared to western lowland (42-53%; Campbell et al.,

2020; Fontsere et al., 2021; Hicks et al., 2018) and mountain gorillas (34%, Gomez et al. 2016). The

abundance of Bacteroidetes (21.0%) was intermediate between western lowland (10-19%; Campbell et

al., 2020; Fontsere et al., 2021; Hicks et al., 2018) and mountain gorillas (30%, Gomez et al. 2016).

Higher ratios of Firmicutes-to-Bacteroidetes are thought to indicate greater energy extraction potential

(Mariat *et al.* 2009) and decline in captivity (Clayton et al. 2018). Thus, the relatively high F:B ratio in

Grauer's gorillas here might facilitate efficient metabolism of their fibrous food resources [log(F:B) =

1.12 in this study compared to log(F:B) < 1 in wild western lowland gorillas, Hicks et al. 2018]. Further,

we detected higher gut microbiome alpha diversity in Grauer's gorillas (13 phyla/sample) than reported

for western lowland gorillas (9), chimpanzees (9), and humans (<5, Hicks et al*., 2018), suggesting either

higher diversity of Grauer's gorilla microbiomes or better characterisation of rare phyla in our study

(*e.g.,* Patescibacteria, Epsilonbacteraeota).

We identified eleven microbial taxa present in every individual in this dataset. As these are

shared by individuals living up to 175 km apart, they may be functionally important. Many of the these

are carbohydrate- and fibre-degraders related to common members of western lowland gorilla

microbiomes that were much more common in wild vs. captive individuals, e.g. *Flexilinea* (Chloroflexi;

Anaerolineaceae)*, Prevotella 7* (Bacteroidetes; Prevotellaceae)*,* and *Ruminococcus* (Firmicutes;

Ruminococcaceae) (Campbell et al., 2020; Hicks et al., 2018; Narat et al., 2020). However, some of

these core taxa were not particularly common in previous studies of gorilla microbiomes, *e.g.,*

Senegalimassilia (Actinobacteria; Eggerthellaceae) and *Sutterella* (Gammaproteobacteria;

Burkholderiaceae), though they are common in the gut of baboons (Grieneisen et al., 2021), geladas

(Baniel et al., 2021), white-faced capuchins, (Mallott, Amato, et al., 2018), and woolly monkeys

(Quiroga-González et al., 2021). The gut microbiome of Grauer's gorillas appeared to be relatively

stable through time, as evidenced by greater similarity amongst samples from the same individuals

collected almost one year apart compared to other individuals from the same social group, a finding

generally in agreement with studies on the human gut microbiome (Björk et al., 2019; Faith et al., 2013).

Gut microbiome differs between populations and social groups

The Grauer's gorilla gut microbiome differed between populations separated by large

geographic and genetic distances and also across social groups. As we found no effect of genetic

relatedness, it is likely that contact and social association with groupmates explains why individuals

from the same social group show greater similarity in their gut microbiome than to individuals from

other social groups within the same population. Time spent in close proximity offers opportunities for

transfer of gut microbiota, such as through grooming and coprophagy (Graczyk & Cranfield, 2003). An

effect of social association on the gut microbiome has previously been identified for a range of primates,

including chimpanzees (Degnan et al., 2012; Moeller et al., 2016), baboons (Tung et al., 2015), colobus

monkeys (Wikberg et al., 2020), black howler monkeys (Amato et al., 2017), sifakas (Perofsky et al.,

2017, 2021), and humans (Dill-McFarland et al., 2019), as well as other group-living animals, *e.g.*

bighorn sheep (Couch et al., 2020). However, this does not preclude an effect of heritability.

Longitudinal studies in Amboseli baboons have shown that there is a high heritability of the primate

gut microbiome, which cannot easily be detected in shorter-term studies (Grieneisen et al., 2021).

Alternatively, it may not be social interaction *per se* but rather shared ecological factors within

groups that drive the observed patterns. While diet was less important than population and social group

identity in explaining variation in the gut microbiome, it is possible that dietary similarity among

members of the same group drives convergence in the microbiome over longer time periods. This would

imply limited between-group homogenization in diet, which could occur if groups have limited range

overlap or restrict their foraging to particular areas. Since plant diversity in KBNP varies with altitude

(Imani et al., 2016), this is possible, particularly if gorilla groups forage only within their preferred

altitudinal range, similar to the proposed patterns for female dispersal in the Bwindi mountain gorillas

(Guschanski et al., 2008).

The gorilla gut microbiome is intertwined with diet

Experimental work in captive settings has identified diet as one of the most important factors

determining microbiome diversity and composition (Clayton et al., 2018). Studies of wild primate

populations frequently use proxies for diet, such as seasonal changes or environmental variables (e.g.

forest type, habitat degradation) (Amato et al. 2019; Baniel et al. 2021; Bueno de Mesquita et al. 2021;

Clayton et al. 2018; Correa et al. 2021; Gomez et al. 2015, 2016; but see Hicks et al. 2018 for joint diet

and gut microbiome study). We add to this work by simultaneously characterizing diet and gut

microbiome in two populations of wild Grauer's gorillas. In accordance with expectations from

experimental work, we indeed detect high co-inertia and correlation between diet and microbiome

composition.

Despite this global trend, population and social group membership were the only significant

predictors of microbiome composition. This could indicate that microbial changes reflect long-term

dietary patterns more consistent with social and ranging patterns than with diet at the precise time of

sampling. Other local variables, such as abiotic and biotic environmental factors and the social

environment may have additional effects on the gut microbiome. If diet and microbiome are linked, a

more varied diet might be expected to select for a more complex gut microbiome. However, in contrast

to findings in healthy humans (Heiman & Greenway, 2016), the low-altitude population that had higher

dietary richness did not show higher gut microbial richness than the high-altitude population. In fact,

the population with greater dietary evenness, NCA, had significantly lower gut microbial evenness. To

understand if this relative lack of microbial diversity indicates nutritional redundancy in the diet of

NCA gorillas requires further investigation into the nutritional diversity of the plant species consumed.

Divergence by population and social group in Grauer's gorilla gut microbiomes tended to

derive from differences in specific taxa within universally common bacterial families, especially

Lachnospiraceae and Rikellenaceae. This observation emphasizes the trend towards consistency of

digestive function across populations of the same species (Amato et al., 2019). There were, however,

several family-level differences. For instance, *Treponema* (Spirochaetaceae) were more abundant in

NCA, whereas *Prevotella* (Prevotellaceae) were more abundant in KBNP. Hicks et al. (2018) found

Treponema in western gorillas associated with high-fibre fallback foods, like Marantaceae, Moraceae,

and Zingeraceae. The genus *Prevotella*, meanwhile, was enriched during periods of high fruit

abundance (*e.g.,* Annonaceae, and Urticaceae) (Hicks et al., 2018). *Prevotella* is common in

carbohydrate-rich diets in other studies as well, and is sometimes associated with captivity (Campbell

et al., 2020; Clayton et al., 2016; Hicks et al., 2018; Precup & Vodnar, 2019). Consistent with these

reports, we found greater abundance of Marantaceae, Moraceae, Commelinaceae, and Zingeraceae in

diets of NCA Grauer's gorillas, which had more *Treponema*, and greater abundance of Urticaceae,

Primulaceae, Sapindaceae, and Piperaceae in KBNP, where *Prevotella* was more abundant. While

speculation about functional correlations is possible, alternate approaches, such as shotgun

metagenomics coupled with deeper characterization of global bacterial diversity and metabolic function

would further enhance our understanding of the relationship between diet and microbial communities.

Conclusions

We provide evidence that fragmented and genetically differentiated populations of critically

endangered Grauer's gorillas differ in dietary and gut microbiome composition, suggesting that they

might be locally adapted. We emphasize the utility of faecal sampling for minimally-invasive

population monitoring of different aspects of endangered species biology, from genetics to ecology to

foraging behaviour. Despite its limitations, a molecular approach can reveal otherwise clandestine

insights into the ecology of elusive animals and is particularly powerful when combined with traditional

field observational methods. Our results stress the importance of protecting geographically-isolated

wildlife populations, as safeguarding ecological and genetic biodiversity is the objective of species

conservation.

Acknowledgements

We thank L'Institut Congolais pour la Conservation de la Nature (ICCN) and local landowners and community members, for permitting us to work in the Kahuzi-Biega National Park and the Nkuba Conservation Area. We are indebted to numerous rangers and field assistants in Kahuzi-Biega National Park as well as field assistants, gorilla trackers, and local community members in Nkuba Conservation Area who, through their effort and commitment, support this project and ensure the survival of the Grauer's gorillas. This research was supported by the Royal Physiographic Society of Lund Jan Löfqvist and Nilsson-Ehle Endowments to KG and the Erasmus Mundus Master Programme in Evolutionary Biology Consortium Scholarship to AM. The Dian Fossey Gorilla Fund's work in DRC was supported by individual donations and grants from the Great Ape Conservation Fund of the US Fish and Wildlife Service, the Arcus Foundation, the Daniel L. Thorne Foundation, and the Turner Foundation. Sequencing was performed by the SNP&SEQ Technology Platform in Uppsala. The SNP&SEQ Technology Platform is part of the National Genomics Infrastructure Sweden and Science of Life Laboratory. The SNP&SEQ Platform is also supported by the Swedish Research Council and the Knut and Alice Wallenberg Foundation. We also acknowledge the National Bioinformatics Infrastructure for providing computational resources to this project.

Author contribution

AM, RM and KG planned the study design. AM generated dietary data and performed all analyses. RM generated gut microbial data. PN, YL, MAG, and JS generated gorilla genotyping data. KN and LP provided reagents and expertise for dietary analyses. NI, AP, UN, EB, RNP, DC, and KG collected faecal samples and provided support in the field. DC, LP and KG provided project supervision. KG supervised the experiments and data analyses. AM and KG wrote the manuscript, with contribution from all authors. All authors reviewed and approved of the final manuscript.

Data availability

Sequences for the gut microbiome and diet generated in this project have been uploaded to the European Nucleotide Archive (ENA) under Accession no.: PRJEB49814.

References

Agranyoni, O., Meninger-Mordechay, S., Uzan, A., Ziv, O., Salmon-Divon, M., Rodin, D., Raz, O., Koman, I., Koren, O., Pinhasov, A., & Navon-Venezia, S. (2021). Gut microbiota determines the social behavior of mice and induces metabolic and inflammatory changes in their adipose tissue. *Npj Biofilms and Microbiomes*, *7*(1), 1–14. https://doi.org/10.1038/s41522-021-00193-9

Ait Baamrane, M. A., Shehzad, W., Ouhammou, A., Abbad, A., Naimi, M., Coissac, E., Taberlet, P., & Znari, M. (2012). Assessment of the food habits of the Moroccan dorcas gazelle in M'Sabih Talaa, west central Morocco, using the trnL approach. *PLoS ONE*, *7*(4). https://doi.org/10.1371/journal.pone.0035643

Aitchison, J., Barceló-Vidal, C., Martín-Fernández, J. A., & Pawlowsky-Glahn, V. (2000). Logratio analysis and compositional distance. *Mathematical Geology*, *32*(3), 271–275.

Alberdi, A., Aizpurua, O., Bohmann, K., Zepeda-Mendoza, M. L., & Gilbert, M. T. P. (2016). Do vertebrate gut metagenomes confer rapid ecological adaptation? *Trends in Ecology and Evolution*, *31*(9), 689–699. https://doi.org/10.1016/j.tree.2016.06.008

Alberdi, A., Razgour, O., Aizpurua, O., Novella-Fernandez, R., Aihartza, J., Budinski, I., Garin, I., Ibáñez, C., Izagirre, E., Rebelo, H., Russo, D., Vlaschenko, A., Zhelyazkova, V., Zrnčić, V., & Gilbert, M. T. P. (2020). DNA metabarcoding and spatial modelling link diet diversification with distribution homogeneity in European bats. *Nature Communications*, *11*(1), 1–8. https://doi.org/10.1038/s41467-020-14961-2

Amato, K. R., G. Sanders, J., Song, S. J., Nute, M., Metcalf, J. L., Thompson, L. R., Morton, J. T., Amir, A., J. McKenzie, V., Humphrey, G., Gogul, G., Gaffney, J., L. Baden, A., A.O. Britton, G., P. Cuozzo, F., Di Fiore, A., J. Dominy, N., L. Goldberg, T., Gomez, A., … R. Leigh, S. (2019). Evolutionary trends in host physiology outweigh dietary niche in structuring primate gut microbiomes. *ISME Journal*, *13*(3), 576–587. https://doi.org/10.1038/s41396-018-0175-0

Amato, K. R., Van Belle, S., Di Fiore, A., Estrada, A., Stumpf, R., White, B., Nelson, K. E., Knight, R., & Leigh, S. R. (2017). Patterns in gut microbiota similarity associated with degree of sociality among sex classes of a neotropical primate. *Microbial Ecology*, *74*(1), 250–258. https://doi.org/10.1007/s00248-017-0938-6

Anderson, M. J., & Walsh, D. C. I. (2013). PERMANOVA, ANOSIM, and the Mantel test in the face of heterogeneous dispersions: what null hypothesis are you testing? *Ecological Monographs*, *83*(4), 557–574.

Arandjelovic, M., Guschanski, K., Schubert, G., Harris, T. R., Thalmann, O., Siedel, H., & Vigilant, L. (2009). Two-step multiplex polymerase chain reaction improves the speed and accuracy of genotyping using DNA from noninvasive and museum samples. *Molecular Ecology Resources*, *9*(1), 28–36. https://doi.org/10.1111/j.1755-0998.2008.02387.x

Arbizu, P. M. (2020). *pairwiseAdonis: pairwise multilevel comparison using adonis*. https://github.com/pmartinezarbizu/pairwiseAdonis.

Baas, P., van der Valk, T., Vigilant, L., Ngobobo, U., Binyinyi, E., Nishuli, R., Caillaud, D., & Guschanski, K. (2018). Population-level assessment of genetic diversity and habitat fragmentation in critically endangered Grauer's gorillas. *American Journal of Physical Anthropology*, *165*(3), 565–575. https://doi.org/10.1002/ajpa.23393

Baniel, A., Amato, K. R., Beehner, J. C., Bergman, T. J., Mercer, A., Perlman, R. F., Petrullo, L., Reitsema, L., Sams, S., Lu, A., & Snyder-Mackler, N. (2021). Seasonal shifts in the gut microbiome indicate plastic responses to diet in wild geladas. *Microbiome*, *9*(1), 1–20. https://doi.org/10.1186/s40168-020-00977-9

Bates, D., Maechler, M., Bolker, B. M., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

Beja-Pereira, A., Oliveira, R., Alves, P. C., Schwartz, M. K., & Luikart, G. (2009). Advancing ecological understandings through technological transformations in noninvasive genetics. *Molecular Ecology Resources*, *9*(5), 1279–1301. https://doi.org/10.1111/j.1755-0998.2009.02699.x

Björk, J. R., Dasari, M., Grieneisen, L., & Archie, E. A. (2019). Primate microbiomes over time: Longitudinal answers to standing questions in microbiome research. *American Journal of Primatology*, *81*(10–11), 1–23. https://doi.org/10.1002/ajp.22970

Bolnick, D. I., Snowberg, L. K., Hirsch, P. E., Lauber, C. L., Knight, R., Caporaso, J. G., & Svanbäck, R. (2014). Individuals' diet diversity influences gut microbial diversity in two freshwater fish (threespine stickleback and Eurasian perch). *Ecology Letters*, *17*(8), 979–987. https://doi.org/10.1111/ele.12301

Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., & Coissac, E. (2016). obitools: A unix-inspired software package for DNA metabarcoding. *Molecular Ecology Resources*, *16*(1), 176–182.

Bradley, B. J., Chambers, K. E., & Vigilant, L. (2001). Accurate DNA-based sex identification of apes using non-invasive samples. *Conservation Genetics*, *2*(2), 179–181. https://doi.org/10.1023/A:1011847528045

Bradley, B. J., Stiller, M., Doran-Sheehy, D. M., Harris, T., Chapman, C. A., Vigilant, L., & Poinar, H. (2007). Plant DNA sequences from feces: Potential means for assessing diets of wild primates. *American Journal of Primatology*, *69*(6), 699–705. https://doi.org/10.1002/ajp.20384

Brealey, J. C., Leitão, H. G., Hofstede, T., Kalthoff, D. C., & Guschanski, K. (2021). The oral microbiota of
wild bears in Sweden reflects the history of antibiotic use by humans. *Current Biology*, *31*(20), 4650–
4658.e6. https://doi.org/10.1016/j.cub.2021.08.010
Brealey, J. C., Leitão, H. G., Van-Der-Valk, T., Xu, W., Bougiouri, K., Dalén, L., & Guschanski, K. (2020).
Dental calculus as a tool to study the evolution of the mammalian oral microbiome. *Molecular Biology*
and Evolution, *37*(10), 3003–3022. https://doi.org/10.1093/molbev/msaa135
Bueno de Mesquita, C. P., Nichols, L. M., Gebert, M. J., Vanderburgh, C., Bocksberger, G., Lester, J. D., Kalan,
A. K., Dieguez, P., McCarthy, M. S., Agbor, A., Álvarez Varona, P., Ayimisin, A. E., Bessone, M.,
Chancellor, R., Cohen, H., Coupland, C., Deschner, T., Egbe, V. E., Goedmakers, A., … Dunn, R. R.
(2021). Structure of chimpanzee gut microbiomes across tropical Africa. *MSystems*, *6*(3).
https://doi.org/10.1128/msystems.01269-20
Cadotte, M. W., Carscadden, K., & Mirotchnick, N. (2011). Beyond species: Functional diversity and the
maintenance of ecological processes and services. *Journal of Applied Ecology*, *48*(5), 1079–1087.
https://doi.org/10.1111/j.1365-2664.2011.02048.x
Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2:
High-resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*(7), 581–583.
https://doi.org/10.1038/nmeth.3869
Campbell, T. P., Sun, X., Patel, V. H., Sanz, C., Morgan, D., & Dantas, G. (2020). The microbiome and
resistome of chimpanzees, gorillas, and humans across host lifestyle and geography. *ISME Journal*, *14*(6),
1584–1599. https://doi.org/10.1038/s41396-020-0634-2
Ceballos, G., Ehrlich, P. R., & Raven, P. H. (2020). Vertebrates on the brink as indicators of biological
annihilation and the sixth mass extinction. *Proceedings of the National Academy of Sciences of the United*
States of America, *117*(24), 13596–13602. https://doi.org/10.1073/pnas.1922686117
Chao, A., Gotelli, N. J., Hsieh, T. C., Sander, E. L., Ma, K. H., Colwell, R. K., & Ellison, A. M. (2014).
Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species
diversity studies. *Ecological Monographs*, *84*(1), 45–67.
Chatterji, S., & Pachter, L. (2006). Reference based annotation with GeneMapper. *Genome Biology*, *7*(4), 1–10.
Clayton, J. B., Al-Ghalith, G. A., Long, H. T., Tuan, B. Van, Cabana, F., Huang, H., Vangay, P., Ward, T.,
Minh, V. Van, Tam, N. A., Dat, N. T., Travis, D. A., Murtaugh, M. P., Covert, H., Glander, K. E., Nadler,
T., Toddes, B., Sha, J. C. M., Singer, R., … Johnson, T. J. (2018). Associations between nutrition, gut
microbiome, and health in a novel nonhuman primate model. *Scientific Reports*, *8*(1), 1–16.
https://doi.org/10.1038/s41598-018-29277-x
Clayton, J. B., Vangay, P., Huang, H., Ward, T., Hillmann, B. M., Al-Ghalith, G. A., Travis, D. A., Long, H. T.,
Van Tuan, B., Van Minh, V., Cabana, F., Nadler, T., Toddes, B., Murphy, T., Glander, K. E., Johnson, T.
J., & Knights, D. (2016). Captivity humanizes the primate microbiome. *Proceedings of the National*
Academy of Sciences of the United States of America, *113*(37), 10376–10381.
https://doi.org/10.1073/pnas.1521835113
Colston, T. J., & Jackson, C. R. (2016). Microbiome evolution along divergent branches of the vertebrate tree of
life: what is known and unknown. *Molecular Ecology*, *25*(16), 3776–3800.
https://doi.org/10.1111/mec.13730
Correa, F., Torti, V., Spiezio, C., Checcucci, A., Modesto, M., Borruso, L., Cavani, L., Mimmo, T., Cesco, S.,
Luise, D., Randrianarison, R. M., Gamba, M., Rarojoson, N. J., Sanguinetti, M., Di Vito, M., Bugli, F.,
Mattarelli, P., Trevisi, P., Giacoma, C., & Sandri, C. (2021). Disentangling the possible drivers of Indri
indri microbiome: A threatened lemur species of Madagascar. *Frontiers in Microbiology*, *12*(August), 1–
12. https://doi.org/10.3389/fmicb.2021.668274
Couch, C. E., Arnold, H. K., Crowhurst, R. S., Jolles, A. E., Sharpton, T. J., Witczak, M. F., Epps, C. W., &
Beechler, B. R. (2020). Bighorn sheep gut microbiomes associate with genetic and spatial structure across
a metapopulation. *Scientific Reports*, *10*(1), 1–10. https://doi.org/10.1038/s41598-020-63401-0
Davidson, G. L., Raulo, A., & Knowles, S. C. L. (2020). Identifying microbiome-mediated behaviour in wild
vertebrates. *Trends in Ecology and Evolution*, *35*(11), 972–980. https://doi.org/10.1016/j.tree.2020.06.014
Deagle, B. E., Thomas, A. C., McInnes, J. C., Clarke, L. J., Vesterinen, E. J., Clare, E. L., Kartzinel, T. R., &
Eveson, J. P. (2019). Counting with DNA in metabarcoding studies: How should we convert sequence
reads to dietary data? *Molecular Ecology*, *28*(2), 391–406. https://doi.org/10.1111/mec.14734
Degnan, P. H., Pusey, A. E., Lonsdorf, E. V., Goodall, J., Wroblewski, E. E., Wilson, M. L., Rudicell, R. S.,
Hahn, B. H., & Ochman, H. (2012). Factors associated with the diversification of the gut microbial
communities within chimpanzees from Gombe National Park. *Proceedings of the National Academy of*
Sciences of the United States of America, *109*(32), 13034–13039.
https://doi.org/10.1073/pnas.1110994109
Delgado-Baquerizo, M., Oliverio, A. M., Brewer, T. E., Benavent-González, A., Eldridge, D. J., Bardgett, R. D.,
Maestre, F. T., Singh, B. K., & Fierer, N. (2018). A global atlas of the dominant bacteria found in soil.
Science, *359*(6373), 320–325. https://doi.org/10.1126/science.aap9516

DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., Huber, T., Dalevi, D., Hu, P., & Andersen, G. L. (2006). Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Applied and Environmental Microbiology*, *72*(7), 5069–5072.

DeWoody, J. A., Harder, A. M., Mathur, S., & Willoughby, J. R. (2021). The long-standing significance of genetic diversity in conservation. *Molecular Ecology*, *30*(17), 4147–4154. https://doi.org/10.1111/mec.16051

Dill-McFarland, K. A., Tang, Z. Z., Kemis, J. H., Kerby, R. L., Chen, G., Palloni, A., Sorenson, T., Rey, F. E., & Herd, P. (2019). Close social relationships correlate with human gut microbiota composition. *Scientific Reports*, *9*(1), 1–10. https://doi.org/10.1038/s41598-018-37298-9

Edgar, R. C. (2018). Updating the 97% identity threshold for 16S ribosomal RNA OTUs. *Bioinformatics*, *34*(14), 2371–2375. https://doi.org/10.1093/bioinformatics/bty113

Egea, I., Barsan, C., Bian, W., Purgatto, E., Latché, A., Chervin, C., Bouzayen, M., & Pech, J. C. (2010). Chromoplast differentiation: Current status and perspectives. *Plant and Cell Physiology*, *51*(10), 1601–1611. https://doi.org/10.1093/pcp/pcq136

Eisenhofer, R., Minich, J. J., Marotz, C., Cooper, A., Knight, R., & Weyrich, L. S. (2019). Contamination in low microbial biomass microbiome studies: issues and recommendations. *Trends in Microbiology*, *27*(2), 105–117.

Escoufier, Y. (1973). Le traitement des variables vectorielles. *Biometrics*, *29*(4), 751. https://doi.org/10.2307/2529140

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology*, *14*(8), 2611–2620.

Faith, J. J., Guruge, J. L., Charbonneau, M., Subramanian, S., Seedorf, H., Goodman, A. L., Clemente, J. C., Knight, R., Heath, A. C., Leibel, R. L., Rosenbaum, M., & Gordon, J. I. (2013). The long-term stability of the human gut microbiota. *Science*, *341*(6141). https://doi.org/10.1126/science.1237439

Falush, D., Stephens, M., & Pritchard, J. K. (2003). Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics*, *164*(4), 1567–1587.

Ficetola, G. F., Coissac, E., Zundel, S., Riaz, T., Shehzad, W., Bessière, J., Taberlet, P., & Pompanon, F. (2010). An in silico approach for the evaluation of DNA barcodes. *BMC Genomics*, *11*(1), 1–10.

Fietz, K., Rye Hintze, C. O., Skovrind, M., Kjærgaard Nielsen, T., Limborg, M. T., Krag, M. A., Palsbøll, P. J., Hestbjerg Hansen, L., Rask Møller, P., & Gilbert, M. T. P. (2018). Mind the gut: Genomic insights to population divergence and gut microbial composition of two marine keystone species. *Microbiome*, *6*(1), 1–16. https://doi.org/10.1186/s40168-018-0467-7

Fontsere, C., Frandsen, P., Hernandez-Rodriguez, J., Niemann, J., Scharff-Olsen, C. H., Vallet, D., Le Gouar, P., Ménard, N., Navarro, A., Siegismund, H. R., Hvilsom, C., Gilbert, M. T. P., Kuhlwilm, M., Hughes, D., & Marques-Bonet, T. (2021). The genetic impact of an Ebola outbreak on a wild gorilla population. *BMC Genomics*, *22*(1). https://doi.org/10.1186/s12864-021-08025-y

Foster, Z. S. L., Sharpton, T. J., & Grünwald, N. J. (2017). Metacoder: An R package for visualization and manipulation of community taxonomic diversity data. *PLoS Computational Biology*, *13*(2), e1005404.

Francis, R. M. (2017). pophelper: an R package and web app to analyse and visualize population structure. *Molecular Ecology Resources*, *17*(1), 27–32.

Ganas, J., Robbins, M. M., Nkurunungi, J. B., Kaplin, B. A., & McNeilage, A. (2004). Dietary variability of mountain gorillas in Bwindi Impenetrable National Park, Uganda. *International Journal of Primatology*, *25*(5), 1043–1072. https://doi.org/10.1023/B:IJOP.0000043351.20129.44

Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V., & Egozcue, J. J. (2017). Microbiome datasets are compositional: and this is not optional. *Frontiers in Microbiology*, *8*, 2224.

Goldberg, A. R., Conway, C. J., Tank, D. C., Andrews, K. R., Gour, D. S., & Waits, L. P. (2020). Diet of a rare herbivore based on DNA metabarcoding of feces: Selection, seasonality, and survival. *Ecology and Evolution*, *10*(14), 7627–7643. https://doi.org/10.1002/ece3.6488

Gomez, A., Petrzelkova, K., Yeoman, C. J., Vlckova, K., Mrázek, J., Koppova, I., Carbonero, F., Ulanov, A., Modry, D., Todd, A., Torralba, M., Nelson, K. E., Gaskins, H. R., Wilson, B., Stumpf, R. M., White, B. A., & Leigh, S. R. (2015). Gut microbiome composition and metabolomic profiles of wild western lowland gorillas (Gorilla gorilla gorilla) reflect host ecology. *Molecular Ecology*, *24*(10), 2551–2565. https://doi.org/10.1111/mec.13181

Gomez, A., Rothman, J. M., Petrzelkova, K., Yeoman, C. J., Vlckova, K., Umaña, J. D., Carr, M., Modry, D., Todd, A., Torralba, M., Nelson, K. E., Stumpf, R. M., Wilson, B. A., Blekhman, R., White, B. A., & Leigh, S. R. (2016). Temporal variation selects for diet-microbe co-metabolic traits in the gut of Gorilla spp. *ISME Journal*, *10*(2), 514–526. https://doi.org/10.1038/ismej.2015.146

Goslee, S. C., & Urban, D. L. (2007). The ecodist package for dissimilarity-based analysis of ecological data. *Journal of Statistical Software*, *22*(1), 1–19.

Graczyk, T. K., & Cranfield, M. R. (2003). Coprophagy and intestinal parasites: implications to human-habituated mountain gorillas (Gorilla gorilla beringei) of the Virunga Mountains and Bwindi Impenetrable

Forest. *Primate Conservation*, *9*, 58–64.

Grieneisen, L., Dasari, M., Gould, T. J., Björk, J. R., Grenier, J. C., Yotova, V., Jansen, D., Gottel, N., Gordon,
J. B., Learn, N. H., Gesquiere, L. R., Wango, T. L., Mututua, R. S., Warutere, J. K., Siodi, L., Gilbert, J.
A., Barreiro, L. B., Alberts, S. C., Tung, J., … Blekhman, R. (2021). Gut microbiome heritability is nearly
universal but environmentally contingent. *Science*, *373*(6551), 181–186.
https://doi.org/10.1126/science.aba5483

Guschanski, K., Caillaud, D., Robbins, M. M., & Vigilant, L. (2008). Females shape the genetic structure of a
gorilla population. *Current Biology*, *18*(22), 1809–1814. https://doi.org/10.1016/j.cub.2008.10.031

Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., Lovejoy, T. E., Sexton, J.
O., Austin, M. P., Collins, C. D., Cook, W. M., Damschen, E. I., Ewers, R. M., Foster, B. L., Jenkins, C.
N., King, A. J., Laurance, W. F., Levey, D. J., Margules, C. R., … Townshend, J. R. (2015). Habitat
fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*, *1*(2), 1–10.
https://doi.org/10.1126/sciadv.1500052

Harcourt, A. H., & Stewart, K. J. (2013). *Gorilla Society: Conflict, Compromise, and Cooperation Between the*
Sexes. The University of Chicago Press. https://doi.org/10.7208/chicago/9780226316048.001.0001

Heiman, M. L., & Greenway, F. L. (2016). A healthy gastrointestinal microbiome is dependent on dietary
diversity. *Molecular Metabolism*, *5*(5), 317–320. https://doi.org/10.1016/j.molmet.2016.02.005

Henry, L. P., Bruijning, M., Forsberg, S. K. G., & Ayroles, J. F. (2019). Can the microbiome influence host
evolutionary trajectories? *BioRxiv*, 700237. https://doi.org/10.1101/700237

Hibert, F., Taberlet, P., Chave, J., Scotti-Saintagne, C., Sabatier, D., & Richard-Hansen, C. (2013). Unveiling
the diet of elusive rainforest herbivores in next generation sequencing era? The tapir as a case study. *PLoS*
ONE, *8*(4). https://doi.org/10.1371/journal.pone.0060799

Hicks, A. L., Lee, K. J., Couto-Rodriguez, M., Patel, J., Sinha, R., Guo, C., Olson, S. H., Seimon, A., Seimon,
T. A., Ondzie, A. U., Karesh, W. B., Reed, P., Cameron, K. N., Lipkin, W. I., & Williams, B. L. (2018).
Gut microbiomes of wild great apes fluctuate seasonally in response to diet. *Nature Communications*,
9(1). https://doi.org/10.1038/s41467-018-04204-w

Imani, G., Zapfack, L., Kalume, J., Riera, B., Cirimwami, L., & Boyemba, F. (2016). Woody vegetation groups
and diversity along the altitudinal gradient in mountain forest: case study of Kahuzi-Biega National Park
and its surroundings, RD Congo. *Journal of Biodiversity and Environmental Sciences*, *8*(6), 134–150.
http://www.innspub.net

IUCN. (2016). *Gorilla beringei ssp. graueri*. The IUCN Red List of Threatened Species.
https://doi.org/https://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T39995A17989838.en

IUCN. (2021). *The IUCN Red List of Threatened Species*. https://www.iucn.org/.

Jakobsson, M., & Rosenberg, N. A. (2007). CLUMPP: a cluster matching and permutation program for dealing
with label switching and multimodality in analysis of population structure. *Bioinformatics*, *23*(14), 1801–
1806.

Jewell, Z. (2013). Effect of monitoring technique on quality of conservation science. *Conservation Biology*,
27(3), 501–508. https://doi.org/10.1111/cobi.12066

Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics*,
24(11), 1403–1405.

Kalinowski, S. T., Taper, M. L., & Marshall, T. C. (2007). Revising how the computer program CERVUS
accommodates genotyping error increases success in paternity assignment. *Molecular Ecology*, *16*(5),
1099–1106. https://doi.org/10.1111/j.1365-294X.2007.03089.x

Kalinowski, S. T., Wagner, A. P., & Taper, M. L. (2006). ML-RELATE: A computer program for maximum
likelihood estimation of relatedness and relationship. *Molecular Ecology Notes*, *6*(2), 576–579.
https://doi.org/10.1111/j.1471-8286.2006.01256.x

Kambale, E. S. (2018). *Diet selection strategies of Grauer's gorillas (Gorilla beringei graueri) in relation to*
nutritional benefits and exposure to heptotocix phytochemicals in Mount Tshiabirimu Forest, Virunga
National Park, DRC. Makerere University.

Kamil, A. C., Krebs, J. R., & Pulliam, H. R. (2012). *Foraging behavior*. Springer Science & Business Media.

Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: an R package for genetic analysis of
populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, *2*, e281.

Kandlikar, G. S., Gold, Z. J., Cowen, M. C., Meyer, R. S., Freise, A. C., Kraft, N. J. B., Moberg-Parker, J.,
Sprague, J., Kushner, D. J., & Curd, E. E. (2018). ranacapa: An R package and Shiny web app to explore
environmental DNA data with exploratory statistics and interactive visualizations. *F1000Research*, *7*.

Kartzinel, T. R., Hsing, J. C., Musili, P. M., Brown, B. R. P., & Pringle, R. M. (2019). Covariation of diet and
gut microbiome in African megafauna. *Proceedings of the National Academy of Sciences of the United*
States of America, *116*(47), 23588–23593. https://doi.org/10.1073/pnas.1905666116

King, S. R. B., & Schoenecker, K. A. (2019). Comparison of methods to examine diet of feral horses from
noninvasively collected fecal samples. *Rangeland Ecology and Management*, *72*(4), 661–666.
https://doi.org/10.1016/j.rama.2019.02.005

Kircher, M., Sawyer, S., & Meyer, M. (2012). Double indexing overcomes inaccuracies in multiplex sequencing on the Illumina platform. *Nucleic Acids Research*, *40*(1), e3–e3.

Kolodny, O., Callahan, B. J., & Douglas, A. E. (2020). The role of the microbiome in host evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *375*(1808), 20190588. https://doi.org/10.1098/rstb.2019.0588

Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). CLUMPAK: a program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources*, *15*(5), 1179–1191.

Lenth, R., Singmann, H., Love, J., Buerkner, P., & Herve, M. (2021). Emmeans: Estimated marginal means, aka least-squares means. *R Package Version 1.7.0*, *1*(1), 3.

Ley, R. E., Lozupone, C. A., Hamady, M., Knight, R., & Gordon, J. I. (2008). Worlds within worlds: Evolution of the vertebrate gut microbiota. *Nature Reviews Microbiology*, *6*(10), 776–788. https://doi.org/10.1038/nrmicro1978

Lichstein, J. W. (2007). Multiple regression on distance matrices: a multivariate spatial analysis tool. *Plant Ecology*, *188*(2), 117–131.

Maldonado, O., Aveling, C., Cox, D., Nixon, S., Merlo, D., Pintea, L., & Williamson, E. A. (2012). *Grauer's Gorillas and Chimpanzees in Eastern Democratic Republic of Congo (Kahuzi-Biega, Maiko, Tayna and Itombwe Landscape): Conservation Action Plan 2012–2022*. IUCN.

Mallott, E. K., Amato, K. R., Garber, P. A., & Malhi, R. S. (2018). Influence of fruit and invertebrate consumption on the gut microbiota of wild white-faced capuchins (Cebus capucinus). *American Journal of Physical Anthropology*, *165*(3), 576–588. https://doi.org/10.1002/ajpa.23395

Mallott, E. K., Garber, P. A., & Malhi, R. S. (2018). Trnl outperforms rbcl as a DNA metabarcoding marker when compared with the observed plant component of the diet of wild white-faced capuchins (Cebus capucinus, Primates). *PLoS ONE*, *13*(6), 1–16. https://doi.org/10.1371/journal.pone.0199556

McMurdie, P. J., & Holmes, S. (2013). phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PloS One*, *8*(4), e61217.

McNeilage, A., Robbins, M. M., Gray, M., Olupot, W., Babaasa, D., Bitariho, R., Kasangaki, A., Rainer, H., Asuma, S., Mugiri, G., & Baker, J. (2006). Census of the mountain gorilla Gorilla beringei beringei population in Bwindi Impenetrable National Park, Uganda. *ORYX*, *40*(4), 419–427. https://doi.org/10.1017/S0030605306001311

Meirmans, P. G. (2020). genodive version 3.0: Easy-to-use software for the analysis of genetic data of diploids and polyploids. *Molecular Ecology Resources*, *20*(4), 1126–1131.

Meirmans, P. G., & Hedrick, P. W. (2011). Assessing population structure: FST and related measures. *Molecular Ecology Resources*, *11*(1), 5–18.

Meng, C., Kuster, B., Culhane, A. C., & Gholami, A. M. (2014). A multivariate approach to the integration of multi-omics datasets. *BMC Bioinformatics*, *15*(1), 1–13.

Meyer, M., & Kircher, M. (2010). Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harbor Protocols*, *5*(6), 1–11. https://doi.org/10.1101/pdb.prot5448

Mirarab, S., Nguyen, N., & Warnow, T. (2012). SEPP: SATé-enabled phylogenetic placement. *Biocomputing*, 247–258.

Moeller, A. H., Foerster, S., Wilson, M. L., Pusey, A. E., Hahn, B. H., & Ochman, H. (2016). Social behavior shapes the chimpanzee pan-microbiome. *Science Advances*, *2*(1). https://doi.org/10.1126/sciadv.1500997

Moeller, A. H., & Sanders, J. G. (2020). Roles of the gut microbiota in the adaptive evolution of mammalian species: Microbiotas alter mammalian adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *375*(1808), 1–10. https://doi.org/10.1098/rstb.2019.0597rstb20190597

Moran, N. A., Ochman, H., & Hammer, T. J. (2019). Evolutionary and ecological consequences of gut microbial communities. *Annual Review of Ecology, Evolution, and Systematics*, *50*, 451–475. https://doi.org/10.1146/annurev-ecolsys-110617-062453

Muegge, B. D., Kuczynski, J., Knights, D., Clemente, J. C., González, A., Fontana, L., Henrissat, B., Knight, R., & Gordon, J. I. (2011). Diet drives convergence in gut microbiome functions across mammalian phylogeny and within humans. *Science*, *332*(6032), 970–974. https://doi.org/10.1126/science.1198719

Murtagh, F., & Legendre, P. (2014). Ward's hierarchical agglomerative clustering method: which algorithms implement Ward's criterion? *Journal of Classification*, *31*(3), 274–295.

Narat, V., Amato, K. R., Ranger, N., Salmona, M., Mercier-Delarue, S., Rupp, S., Ambata, P., Njouom, R., Simon, F., Giles-Vernick, T., & LeGoff, J. (2020). A multi-disciplinary comparison of great ape gut microbiota in a central African forest and European zoo. *Scientific Reports*, *10*(1), 1–15. https://doi.org/10.1038/s41598-020-75847-3

Nawaz, M. A., Valentini, A., Khan, N. K., Miquel, C., Taberlet, P., & Swenson, J. E. (2019). Diet of the brown bear in Himalaya: Combining classical and molecular genetic techniques. *PLoS ONE*, *14*(12), 1–21. https://doi.org/10.1371/journal.pone.0225698

Nishida, A. H., & Ochman, H. (2019). A great-ape view of the gut microbiome. *Nature Reviews Genetics*,

20(4), 195–206. https://doi.org/10.1038/s41576-018-0085-z

Nsubuga, A. M., Robbins, M. M., Roeder, A. D., Morin, P. A., Boesch, C., & Vigilant, L. (2004). Factors
affecting the amount of genomic DNA extracted from ape faeces and the identification of an improved
sample storage method. *Molecular Ecology*, *13*(7), 2089–2094. https://doi.org/10.1111/j.1365-
294X.2004.02207.x

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos,
P., Henry, M., & Stevens, M. H. H. (2020). Vegan community ecology package: ordination methods,
diversity analysis and other functions for community and vegetation ecologists. *R Package Version 2.5-7*.

Osman, N. A., Abdul-Latiff, M. A. B., Mohd-Ridwan, A. R., Yaakop, S., Nor, S. M., & Md-Zain, B. M. (2020).
Diet composition of the wild stump-tailed macaque metabarcoding approach: A preliminary study.
Animals, *10*, 1–14.

Palarea-Albaladejo, J., & Martín-Fernández, J. A. (2015). zCompositions—R package for multivariate
imputation of left-censored data under a compositional approach. *Chemometrics and Intelligent*
Laboratory Systems, *143*, 85–96.

Perofsky, A. C., Ancel Meyers, L., Abondano, L. A., Di Fiore, A., & Lewis, R. J. (2021). Social groups
constrain the spatiotemporal dynamics of wild sifaka gut microbiomes. *Molecular Ecology*, *September*, 1–
17. https://doi.org/10.1111/mec.16193

Perofsky, A. C., Lewis, R. J., Abondano, L. A., Difiore, A., & Meyers, L. A. (2017). Hierarchical social
networks shape gut microbial composition in wild verreaux's sifaka. *Proceedings of the Royal Society B:*
Biological Sciences, *284*(1868). https://doi.org/10.1098/rspb.2017.2274

Plumptre, A. J., Kirkby, A., Spira, C., Kivono, J., Mitamba, G., Ngoy, E., Nishuli, R., Strindberg, S., Maisels,
F., Buckland, S., Ormsby, L., & Kujirakwinja, D. (2021). Changes in Grauer's gorilla (Gorilla beringei
graueri) and other primate populations in the Kahuzi-Biega National Park and Oku Community Reserve,
the heart of Grauer's gorilla global range. *American Journal of Primatology*, *83*(7).
https://doi.org/10.1002/ajp.23288

Plumptre, A. J., Nixon, S., Kujirakwinja, D. K., Vieilledent, G., Critchlow, R., Williamson, E. A., Nishuli, R.,
Kirkby, A. E., & Hall, J. S. (2016). Catastrophic decline of world's largest primate: 80% loss of grauer's
gorilla (Gorilla beringei graueri) population justifies critically endangered status. *PLoS ONE*, *11*(10).
https://doi.org/10.1371/journal.pone.0162697

Porras-Hurtado, L., Ruiz, Y., Santos, C., Phillips, C., Carracedo, Á., & Lareu, M. (2013). An overview of
STRUCTURE: applications, parameter settings, and supporting software. *Frontiers in Genetics*, *4*, 98.

Precup, G., & Vodnar, D. C. (2019). Gut Prevotella as a possible biomarker of diet and its eubiotic versus
dysbiotic roles: A comprehensive literature review. *British Journal of Nutrition*, *122*(2), 131–140.
https://doi.org/10.1017/S0007114519000680

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2012). The
SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic*
Acids Research, *41*(D1), D590–D596.

Quéméré, E., Hibert, F., Miquel, C., Lhuillier, E., Rasolondraibe, E., Champeau, J., Rabarivola, C., Nusbaumer,
L., Chatelain, C., Gautier, L., Ranirison, P., Crouau-Roy, B., Taberlet, P., & Chikhi, L. (2013). A DNA
metabarcoding study of a primate dietary diversity and plasticity across its entire fragmented range. *PLoS*
ONE, *8*(3). https://doi.org/10.1371/journal.pone.0058971

Quiroga-González, C., Cardenas, L. A. C., Ramírez, M., Reyes, A., González, C., & Stevenson, P. R. (2021).
Monitoring the variation in the gut microbiota of captive woolly monkeys related to changes in diet during
a reintroduction process. *Scientific Reports*, *11*(1), 1–11. https://doi.org/10.1038/s41598-021-85990-0

R Core Team. (2021). R: A language and environment for statistical computing. *R Foundation for Statistical*
Computing. https://www.r-project.org/

Rahbek, C. (1995). The elevational gradient of species richness: a uniform pattern? *Ecography*, *18*(2), 200–205.
https://doi.org/10.1111/j.1600-0587.1995.tb00341.x

Raymann, K., Moeller, A. H., Goodman, A. L., & Ochman, H. (2017). Unexplored archaeal diversity in the
great ape gut microbiome. *MSphere*, *2*(1). https://doi.org/10.1128/msphere.00026-17

Raymond, M., & Rousset, F. (1995). GENEPOP (Version 1.2): Population Genetics Software for Exact Tests
and Ecumenicism. *Journal of Heredity*, *86*(3), 248–249.
https://doi.org/10.1093/oxfordjournals.jhered.a111573

Remis, M. J. (2000). Initial studies on the contributions of body size and gastrointestinal passage rates to dietary
flexibility among gorillas. *American Journal of Physical Anthropology*, *112*(2), 171–180.
https://doi.org/10.1002/(SICI)1096-8644(2000)112:2<171::AID-AJPA4>3.0.CO;2-F

Remis, M. J., Dierenfeld, E. S., Mowry, C. B., & Carroll, R. W. (2001). Nutritional aspects of western lowland
gorilla (Gorilla gorilla gorilla) diet during seasons of fruit scarcity at Bai Hokou, Central African
Republic. *International Journal of Primatology*, *22*(5), 807–836.
https://doi.org/10.1023/A:1012021617737

Ripple, W. J., Wolf, C., Newsome, T. M., Betts, M. G., Ceballos, G., Courchamp, F., Hayward, M. W., Van

Valkenburgh, B., Wallach, A. D., & Worm, B. (2019). Are we eating the world's megafauna to extinction? *Conservation Letters*, *12*(3), 1–10. https://doi.org/10.1111/conl.12627

Robert, P., & Escoufier, Y. (1976). A unifying tool for linear multivariate statistical methods: The RV-coefficient. *Applied Statistics*, *25*(3), 257. https://doi.org/10.2307/2347233

Rogers, M. E., Abernethy, K., Bermejo, M., Cipolletta, C., Doran, D., Mcfarland, K., Nishihara, T., Remis, M., & Tutin, C. E. G. (2004). Western gorilla diet: A synthesis from six sites. *American Journal of Primatology*, *64*(2), 173–192. https://doi.org/10.1002/ajp.20071

Rohland, N., Harney, E., Mallick, S., Nordenfelt, S., & Reich, D. (2015). Partial uracil – DNA – glycosylase treatment for screening of ancient DNA. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *370*(1660), 1–11. https://doi.org/10.1098/rstb.2013.0624

Rothman, J. M., Nkurunungi, J. B., Shannon, B. F., & Bryer, M. A. H. (2014). High altitude diets: Implications for the feeding and nutritional ecology of mountain gorillas. In *High Altitude Primates* (pp. 247–264). Springer New York. https://doi.org/10.1007/978-1-4614-8175-1_14

Rousset, F. (2008). GENEPOP'007: A complete re-implementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources*, *8*(1), 103–106. https://doi.org/10.1111/j.1471-8286.2007.01931.x

Rudman, S. M., Greenblum, S., Hughes, R. C., Rajpurohit, S., Kiratli, O., Lowder, D. B., Lemmon, S. G., Petrov, D. A., Chaston, J. M., & Schmidt, P. (2019). Microbiome composition shapes rapid genomic adaptation of Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(40), 20025–20032. https://doi.org/10.1073/pnas.1907787116

Schaller, G. E. (1963). *The mountain gorilla: Ecology and behavior.* University of Chicago Press.

Schwartz, M. K., Luikart, G., & Waples, R. S. (2007). Genetic monitoring as a promising tool for conservation and management. *Trends in Ecology and Evolution*, *22*(1), 25–33. https://doi.org/10.1016/j.tree.2006.08.009

Shehzad, W., Riaz, T., Nawaz, M. A., Miquel, C., Poillot, C., Shah, S. A., Pompanon, F., Coissac, E., & Taberlet, P. (2012). Carnivore diet analysis based on next-generation sequencing: Application to the leopard cat (Prionailurus bengalensis) in Pakistan. *Molecular Ecology*, *21*(8), 1951–1965.

Silverman, J. D., Washburne, A. D., Mukherjee, S., & David, L. A. (2017). A phylogenetic transform enhances analysis of compositional microbiota data. *Elife*, *6*, e21887.

Spira, C., Mitamba, G., Kirkby, A., Katembo, J., Kambale, C. K., Musikami, P., Dumbo, P., Byaombe, D.-D., Plumptre, A. J., & Maisels, F. (2018). *Inventaire de la biodiversite dans le Parc National de Kahuzi-Biega, Republique Democratique du Congo.*

Srivathsan, A., Ang, A., Vogler, A. P., & Meier, R. (2016). Fecal metagenomics for the simultaneous assessment of diet, parasites, and population genetics of an understudied primate. *Frontiers in Zoology*, *13*(1), 1–13. https://doi.org/10.1186/s12983-016-0150-4

Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). *Environmental DNA: For biodiversity research and monitoring*. Oxford University Press.

Taberlet, P., Coissac, E., Pompanon, F., Gielly, L., Miquel, C., Valentini, A., Vermat, T., Corthier, G., Brochmann, C., & Willerslev, E. (2007). Power and limitations of the chloroplast trnL (UAA) intron for plant DNA barcoding. *Nucleic Acids Research*, *35*(3), e14–e14. https://doi.org/10.1093/nar/gkl938

Takenoshita, Y., & Yamagiwa, J. (2008). Estimating gorilla abundance by dung count in the northern part of Moukalaba-Doudou National Park, Gabon. *African Study Monographs*, *Suppl. 39*(April), 41–54. http://repository.kulib.kyoto-u.ac.jp/dspace/handle/2433/66239

Teixeira, J. C., & Huber, C. D. (2021). The inflated significance of neutral genetic diversity in conservation genetics. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(10), 1–10. https://doi.org/10.1073/pnas.2015096118

Trevelline, B. K., Fontaine, S. S., Hartup, B. K., & Kohl, K. D. (2019). Conservation biology needs a microbial renaissance: A call for the consideration of host-associated microbiota in wildlife management practices. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1895). https://doi.org/10.1098/rspb.2018.2448

Tung, J., Barreiro, L. B., Burns, M. B., Grenier, J. C., Lynch, J., Grieneisen, L. E., Altmann, J., Alberts, S. C., Blekhman, R., & Archie, E. A. (2015). Social networks predict gut microbiome composition in wild baboons. *ELife*, *2015*(4), 1–18. https://doi.org/10.7554/eLife.05224

Valentini, A., Pompanon, F., & Taberlet, P. (2009). DNA barcoding for ecologists. *Trends in Ecology and Evolution*, *24*(2), 110–117. https://doi.org/10.1016/j.tree.2008.09.011

van den Boogaart, K. G., & Tolosana-Delgado, R. (2008). "Compositions": a unified R package to analyze compositional data. *Computers & Geosciences*, *34*(4), 320–338.

van der Hoek, Y., Binyinyi, E., Ngobobo, U., Stoinski, T. S., & Caillaud, D. (2021). Daily travel distances of unhabituated Grauer's gorillas (Gorilla beringei graueri) in a low elevation forest. *Folia Primatologica*, *92*(2), 112–125. https://doi.org/10.1159/000514626

van der Hoek, Y., Pazo, W. D., Binyinyi, E., Ngobobo, U., Stoinski, T. S., & Caillaud, D. (2021). Diet of

Grauer's gorillas (Gorilla beringei graueri) in a low-elevation forest. *Folia Primatologica*, *92*(2), 126–138. https://doi.org/10.1159/000515377

van der Valk, T., Lona Durazo, F., Dalén, L., & Guschanski, K. (2017). Whole mitochondrial genome capture from faecal samples and museum-preserved specimens. *Molecular Ecology Resources*, *17*(6), e111–e121. https://doi.org/10.1111/1755-0998.12699

van der Valk, T., Vezzi, F., Ormestad, M., Dalén, L., & Guschanski, K. (2020). Index hopping on the Illumina HiseqX platform and its consequences for ancient DNA studies. *Molecular Ecology Resources*, *20*(5), 1171–1181. https://doi.org/10.1111/1755-0998.13009

Watts, D. P. (1984). Composition and variability of mountain gorilla diets in the Central Virungas. *American Journal of Primatology*, *7*(4), 323–356. https://doi.org/10.1002/ajp.1350070403

Weiss, S., Xu, Z. Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., Lozupone, C., Zaneveld, J. R., Vázquez-Baeza, Y., & Birmingham, A. (2017). Normalization and microbial differential abundance strategies depend upon data characteristics. *Microbiome*, *5*(1), 1–18.

West, A. G., Waite, D. W., Deines, P., Bourne, D. G., Digby, A., McKenzie, V. J., & Taylor, M. W. (2019). The microbiome in threatened species conservation. *Biological Conservation*, *229*(November 2018), 85–98. https://doi.org/10.1016/j.biocon.2018.11.016

Wikberg, E. C., Christie, D., Sicotte, P., & Ting, N. (2020). Interactions between social groups of colobus monkeys (Colobus vellerosus) explain similarities in their gut microbiomes. *Animal Behaviour*, *163*, 17–31. https://doi.org/10.1016/j.anbehav.2020.02.011

Yamagiwa, J., & Basabose, A. K. (2006). Diet and seasonal changes in sympatric gorillas and chimpanzees at Kahuzi-Biega National Park. *Primates*, *47*(1), 74–90. https://doi.org/10.1007/s10329-005-0147-7

Yamagiwa, J., Basabose, A. K., Kahekwa, J., Bikaba, D., Ando, C., Matsubara, M., Iwasaki, N., & Sprague, D. S. (2013). Long-Term Research on Grauer's Gorillas in Kahuzi-Biega National Park, DRC: Life History, Foraging Strategies, and Ecological Differentiation from Sympatric Chimpanzees. In P. M. Kappeler & D. P. Watts (Eds.), *Long-Term Field Studies of Primates* (pp. 1–460). Springer. https://doi.org/10.1007/978-3-642-22514-7

Yamagiwa, J., Basabose, A. K., Kaleme, K., & Yumoto, T. (2005). Diet of Grauer's gorillas in the montane forest of Kahuzi, Democratic Republic of Congo. *International Journal of Primatology*, *26*(6), 1345–1373. https://doi.org/10.1007/s10764-005-8856-8

Yamagiwa, J., Mwanza, N., Yumoto, T., & Maruhashi, T. (1991). Ant eating by eastern lowland gorillas. *Primates*, *32*(2), 247–253. https://doi.org/10.1007/BF02381183

Yamagiwa, J., Mwanza, N., Yumoto, T., & Maruhashi, T. (1994). Seasonal change in the composition of the diet of eastern lowland gorillas. *Primates*, *35*(1), 1–14. https://doi.org/10.1007/BF02381481

Youngblut, N. D., Reischer, G. H., Walters, W., Schuster, N., Walzer, C., Stalder, G., Ley, R. E., & Farnleitner, A. H. (2019). Host diet and evolutionary history explain different aspects of gut microbiome diversity among vertebrate clades. *Nature Communications*, *10*(1), 1–15. https://doi.org/10.1038/s41467-019-10191-3

Yumoto, T., Yamagiwa, J., Mwanza, N., & Maruhashi, T. (1994). List of Plant Species Identified in Kahuzi-Biega National Park, Zaïre. *Tropics*, *3*(3/4), 295–308.