Short title: Light regulates nicotine biosynthesis

Mutation in shoot-to-root mobile transcription factor, ELONGATED HYPOCOTYL 5, leads to low
nicotine levels in tobacco

Deeksha Singh[1,2], Shambhavi Dwivedi[3], Hiteshwari Sinha[1,2], Nivedita Singh[1,$], & Prabodh Kumar
Trivedi[1,2,3,*]

[1]CSIR-National Botanical Research Institute, Council of Scientific and Industrial Research (CSIR-NBRI),

Rana Pratap Marg, Lucknow-226001, INDIA

[2]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad- 201002, India

[3]Central Institute of Medicinal and Aromatic Plants (CSIR-CIMAP), Lucknow, India.

[$]Present address:

Department of Botany, University of Lucknow, Lucknow-226007, India.

Corresponding author:

Prabodh Kumar Trivedi,
CSIR-National Botanical Research Institute (CSIR-NBRI), Rana Pratap Marg, Lucknow-226001, INDIA;
Telephone number: +91 522 2297958; e-mail address: prabodht@hotmail.com; prabodht@nbri.res.in
16

ABSTRACT

Tobacco remains one of the most commercially important crops due to the parasympathomimetic alkaloid nicotine used in cigarettes. Most genes involved in nicotine biosynthesis are expressed in root tissues; however, their light-dependent regulation has not been studied. We identified the ELONGATED HYPOCOTYL 5 homolog, NtHY5, from *Nicotiana tabacum* and demonstrated its role in nicotine biosynthesis. We report the development of CRISPR/Cas9-based mutant plants, *NtHY5^{CR}*, and show down-regulation of the nicotine biosynthetic pathway, whereas NtHY5 overexpression (NtHY5OX) plants show the opposite effect. Grafting experiments using wild-type, *NtHY5^{CR},* and NtHY5OX indicated that NtHY5 moves from shoot-to-root to regulate nicotine biosynthesis in the root tissue. We conclude that shoot HY5, directly or through enhancing expression of the HY5 in the root, promotes nicotine biosynthesis. The CRISPR/Cas9-based mutants developed, in this study; with low nicotine accumulation in leaves could help people to overcome their nicotine addiction and the risk of death from tobacco use.

Keywords: *Arabidopsis thaliana,* genome-editing, light-dependent regulation, low nicotine, *NtHY5,* tobacco

INTRODUCTION

Nicotine is the most abundant alkaloid in *Nicotiana tabacum* (Dewey and Xie, 2013), contributing to 90% of the total alkaloids. The development of tobacco plants with significantly less nicotine using genetic engineering techniques is urgently needed to reduce the death rate and diseases caused by tobacco consumption, as the tobacco epidemic is one of the biggest threats to public health, killing millions of people every year around the world (WHO, 2015, 2021). Nicotine is structurally composed of pyridine and pyrrolidine rings (Hibi et al., 1994). The biosynthesis of nicotine occurs in the cortical cells of roots through various enzymatic steps (**Figure 1A**) and is cloistered in vacuoles (Shoji et al., 2009). It is further transported to leaves via xylem (Dewey et al., 2013; Shoji et al., 2021; Hayashi et al., 2020), where it is used for defence against herbivory by the plant (Steppuhn et al., 2004). Quinolinate phosphoribosyl transferase (QPT) (Sinclaie et al., 2000) acts on nicotinic acid and catalyze the formation of the pyridine ring (Hashimotoet et al., 1994). The pyrrolidine ring, on the other hand, is formed from putrescine derived directly from ornithine by ornithine decarboxylase (ODC) (Imanshi et al., 1998). Putrescine N-methyltransferase (PMT) converts putrescine to N-methyl-putrescine (Riechers et al., 1999), which is further oxidized by N-methyl-putrescine oxidase (MPO) and cyclized to N-methyl pyrrolinium cation (Naconsie et al., 2014). The cation finally condenses with nicotinic acid catalysed by NADPH-dependent PIP reductase A622 (DeBoer et al., 2009) and the berberine bridging enzyme-like (BBL) proteins to form nicotine (Kajikawa et al., 2011; Dewey et al., 2013; Hibi et al., 1994; Shoji et al., 2021; Shoji et al., 2011).

Previous research has suggested that a number of regulatory factors are involved in the production of nicotine. According to one study, there are two AP2/ERF genes, NtORC1/ERF221 and NtJAP1/ERF10, both of them positively control the PMT gene. In contrast, NtORC1/ERF221 overexpression only causes high nicotine alkaloid levels (De Boer et al., 20011). Microarray analysis using *Nicotiana tabacum* NIC2 mutant identified several AP2/ERF genes which can regulate nicotine biosynthesis. Seven clustered AP2/ERF genes at the NIC2 locus belong to the group IX AP2/ERF subfamily. One of them, NtERF189, encodes for subclade 2-1 of group IX AP2/ERF protein, is MeJA sensitive, and was expressed in roots, where nicotine is primarily synthesized. The master regulatory role of NtERF189 in nicotine biosynthesis in tobacco is suggested by the fact that knockout mutant of NtERF189 and NtERF199 synthesize reduced

nicotine (Hayashi et al., 2020). The NtERF189 protein recognizes GCC-box-like elements,

(A/C)GC(A/C)(A/C)NCC, in the promoter of nicotine biosynthetic enzyme genes (Shoji et al.,

2012). NtERF32 (EREBP2), another AP2/ERF gene, has also been discovered, which is not a part

of the NIC2 locus. Analysis suggested that NtERF32 may act as a transcriptional activator to

regulate the expression of multiple genes involved in nicotine synthesis by interacting with the

GCC-box-like element (Sears et al., 2014). According to one study, MYC2-type bHLH

transcription factors play a role in regulating the biosynthesis of nicotine. While their

overexpression marginally boosted PMT expression and nicotine content, RNA silencing of

NbbHLH1 and NbbHL2 significantly lowered the expression levels of the genes encoding the

nicotine biosynthesis enzymes. The target GCC-box of NtORC1/ERF221, where NbnHLH1 can

bind, is located proximal to the G-box in the PMT promoter. By interactions with both

components, NtORC1/ERF221 and NbbHLH1 work together to control PMT expression.

Moreover, *N. tabacum* has been found to contain homologous genes of NbbHLH1 and NbbHLH2.

The transcript levels of NIC2-locus AP2/ERF genes, transporter genes, and nicotine biosynthesis

enzymes are favorably regulated by NtMYC2a and NtMYC2b (Shoji et al., 2011). The direct

interaction between NtMYC2 and NtJAZ1 has also been confirmed. The basic helix-loop-helix

protein MYC2 and an orthologous pair of ethylene response factor (ERF) family proteins, ERF189

and ERF199, form a cascade to regulate the nicotine production pathway by specifically

recognizing the GC-rich sequences in the promoter regions of their downstream genes (Shoji et

al., 2010; Shoji et al., 2013; Shoji and Yuan et al., 2021). NtMYB305a has recently been revealed

to bind to the JA-responsive GAG region in the NtPMT1a promoter to positively regulate nicotine

biosynthesis (Bian et al., 2022). Despite the fact that these studies have been done on

transcriptional regulation of nicotine biosynthesis, the light-dependent regulation of the nicotine

pathway in tobacco is not explored yet, although studies suggested the presence of two functionally

important *cis*-elements, a G-box element and a GCC-box element, (Shoji et al., 2000; Oki and

Hashimoto et al., 2004; Xu and Timko, 2004) in the promoter region of PMT.

Light plays a crucial role in plant developmental processes such as flowering, seed development,

organ development, seed germination and seedling de-etiolation (Chory et al., 2010; Vanhaelewyn

et al., 2016; Gangappa et al., 2016; Bhatnagar et al., 2020). Many transcription factors of various

families, including bHLH, bZIP, MYB, GATA and Zinc-finger are activated after light perception

by photoreceptors (Chory et al., 2010). Constitutive nuclear-localized ELONGATED

HYPOCOTYL 5 (HY5), a bZIP family transcription factor, acts downstream to the photoreceptors

and plays a central role in seedling development via integrating light and phytohormone signaling

pathways (Koornneef et al., 1980; Osterlund et al., 2000; Ulm et al., 2012; Li et al., 2013). In

Arabidopsis, HY5 acts as a master regulator by binding to *cis*-regulatory elements of more than

3000 genes directly (Zhang et al., 2011) and regulates flavonoid and terpenoid biosynthesis (Bhatia

et al., 2018; Michael et al., 2021). Though extensive characterization of HY5 and flavonoid

biosynthesis has been done in several plant species, but, there are no reports of characterization of

N. tabacum HY5 and its role in light-regulated photomorphogenesis as well as nicotine and

flavonoid biosynthesis in the plant.

In this study, to decipher the function of HY5 of tobacco in nicotine and flavonoid biosynthesis,

we identified and functionally characterized NtHY5 through complementing *Arabidopsis* HY5

mutant (*athy5*), developing NtHY5-overexpressing lines and CRISPR/Cas9-based mutants. In our

study, through different combinations of grafting experiments, we showed that in response to light,

NtHY5 moves from shoot-to-root and may activate the expression of *NtHY5* in roots as well as

bind directly to the promoter of genes involved in nicotine biosynthesis. CRISPR/Cas9-based HY5

mutants accumulated significantly lower amounts of nicotine compared to wild-type tobacco

plants, suggesting the involvement of light-associated signaling in the regulation of nicotine

biosynthesis in roots.

RESULTS

RESULTS

Identification of light-associated signaling component involved in nicotine biosynthesis

Based on the presence of G and GCC box in the *NtPMT* promoter described in the previous studies

(Xu et al., 2017; Shoji et al., 2000; Oki and Hashimoto et al., 2004; Xu and Timko et al., 2004),

we hypothesized that light could be the factor affecting the nicotine biosynthesis in tobacco.

Previous studies have also established that HY5 binds to G-box (CACGTG) in the promoters of

light-responsive genes (Zhang et al., 2011). To validate the hypothesis, tobacco seedlings were

grown in the light, keeping roots in the dark to maintain natural conditions, and continuously dark

for 10 days. Phenotypically, light- and dark-grown seedlings showed substantial differences in

hypocotyl length, cotyledon opening and chlorophyll content (**Figure 1B**). The gene expression

analysis in roots for nicotine pathway genes, and seedlings for flavonoid pathway genes, suggested

a higher expression of genes involved in the nicotine biosynthesis pathway (*NtODC, NtPMT,*

NtMPO, NtQPT and NtA622) (**Figure 1C**) and flavonoid biosynthesis (*CHS, CHI, FLS and DFR*)

(**Supplemental Figure S1A**) in light-grown compared to dark-grown seedlings. Expression of the

regulatory gene of the flavonoid pathway, *NtMYB12*, was also higher in light-grown than dark-

grown seedlings (**Supplemental Figure S1A**). We measured the nicotine content in the hypocotyl

region of seedlings, and analysis suggested significantly higher nicotine content in light-grown

compared to dark-grown seedlings (**Figure 1D**). We also evaluated the content of flavonols

(kaempferol, quercitin, and rutin), another significant secondary metabolite in tobacco. Analysis

suggested a significantly higher accumulation of flavonols (**Supplemental Figure S1B**) in light-

grown seedlings compared to dark-grown seedlings. Additionally, 10-day-dark grown seedlings

were exposed to light for 0.5, 2, 4, and 24 hours to observe the light-responsiveness of nicotine

pathway genes. The analysis suggested enhanced expression of most of the genes until 4 hours of

light exposure. *NtMPO* and *NtODC* showed the maximum induction (**Figure 1E**).

As HY5 is a key light signaling component in several plants (Nawkar et al., 2017; Burman et al.,

2018), we identified the tobacco HY5, NtHY5, and investigated its function in nicotine and

flavonol accumulation. NtHY5 showed more than 90% sequence similarity with *Arabidopsis* HY5,

AtHY5. Sequence analysis of NtHY5 suggested that it contains 4 exons and 3 introns similar to

AtHY5 (**Supplemental Figure S2A**). Amino acid sequence analysis suggested that NtHY5

contains leucine-rich zinc finger motif and the VPE motif (COP1-interaction motif) (Holm et al.,

2002) and has a high degree of similarity with SlHY5 and AtHY5 (**Supplemental Figure S2B**).

The conserved domain database analysis revealed that NtHY5 is a member of the bZIP

superfamily, similar to HY5 proteins in other plant species (**Supplemental Figure S2C**).

Phylogenetic and pairwise distance analysis suggested that NtHY5 is closer to CaHY5 and SlHY5

than AtHY5 (**Supplemental Figure S3**).

The changes in phenotype, expression of genes and metabolite content in the dark- and light-grown

seedlings led us to quantify the expression of identified NtHY5 gene in the dark- and light-grown

seedlings. The expression of NtHY5 was significantly low in dark-grown seedlings compared to

light-grown seedlings (**Figure 1F**). As the content of flavonols changes in response to light

(Pandey et al., 2014; Bhatia et al., 2018), we studied the responsiveness of flavonols (kaempferol

and quercitin) towards light after exposing 5-day-old dark-grown WT seedlings to light conditions

for 3 h. The analysis suggested a higher accumulation in dark-to-light exposed seedlings compared

to only dark-grown seedlings (**Supplemental Figure S4**).

To investigate the spatio-temporal expression of *NtHY5* and nicotine pathway genes, a comparative

analysis was carried out at different stages of tobacco development. The analysis suggested 4- to

5-fold higher expression of NtHY5 (**Supplemental Figure S5A**) in the leaf, compared with the

seedling, root, stem, and flower of the plant. Expression of *NtODC, NtQPT, NtPMT, NtMPO* and

NtA622 was several folds higher in root in comparison to leaf, flower, seedling, and stem

(**Supplemental Figure S5B**).

NtHY5 interacts with the promoters of genes involved in nicotine and flavonoid biosynthesis

According to the aforementioned data, structural genes of the nicotine biosynthesis (*NtPMT*,

NtQPT, and *NtODC*) and flavonol regulatory genes (*NtMYB12*) have substantially higher

expression in the light compared to dark. It was, therefore, intriguing to analyze whether or not the

light signaling component, NtHY5, directly regulates these genes. *in-silico* analysis suggested that

promoters of *NtPMT*, *NtQPT*, *NtODC*, and *NtMYB12* contain potential LREs (GATA, BOX-4,

ACE, and LTR) upstream of the transcription start site (**Supplemental Table S1-S4**). Analysis

suggested the presence of G-boxes with core motif ACGT at -168 bp (G-box I) and -1040 bp (G-

box II) in the *NtPMT* promoter, -178 bp (G-box I) upstream of the *NtODC* transcription start site

and -36 bp (G-box I) upstream of the *NtQPT* promoter from the transcription start site (**Figure

2A**). NtMYB12 contains G-boxes at -1249 bp and -207 bp upstream of the transcription start site

(**Supplemental Figure S6A**).

An electrophoretic mobility shift assay (EMSA) was employed to analyze the *in vitro* interaction

between ACGT (core motif of G-boxes) and recombinant 6XHis-tagged NtHY5 (NtHY5His)

expressed in *E. coli*. In the EMSA with DIG-labeled G-box probe (G-BOX I and G-BOX II for

PMT and G-BOX I for ODC and QPT) (**Figure 2B**) and NtMYB12 probe (G-BOX I and G-BOX

II) (**Supplemental Figure S6B**), a band shift was observed upon addition of the NtHY5-His

protein. At the same time, no band shift was observed with the mutated probe (mPMT, mODC and

mQPT). In the competition assay, as the concentration of unlabeled probe or cold probe (unPMT,

unODC, unQPT) (**Figure 2C-F**) and (unNtMYB12) (**Supplemental Figure S6C and D**) increases

from 10X to 50X, the intensity of binding got diminished. These findings suggested that NtHY5

interacts with promoters of *NtPMT*, *NtODC*, and *NtQPT* to regulate the expression of these genes

leading to the regulation of nicotine content, whereas to regulate flavonoid content, it binds to

NtMYB12.

NtHY5 is a functional ortholog of AtHY5

To investigate the biological function of NtHY5, the homozygous transgenic lines in the

Arabidopsis HY5 mutant (*athy5* background) were generated and used for detailed analysis.

Phenotypic complementation by NtHY5 in terms of hypocotyls length, root length and rosette

diameter was analysed (**Figure 3A**). The average hypocotyl lengths of 7-d-old wild-type (WT),

hy5 mutant, and NtHY5;*hy5* seedlings were approximately 0.2, 0.7, and 0.4 mm, respectively,

demonstrating that complemented lines have recovered their lost hypocotyl length (**Figure 3B**).

As AtHY5 plays a significant role in light-dependent root growth in *Arabidopsis* (Zhang et al.,

2017), the root length of 7-day-old light-grown WT, *hy5* mutant, and NtHY5;*hy5* complemented

Arabidopsis seedlings were measured. Analysis suggested that the root length of complemented

seedlings is recovered to WT seedlings (**Figure 3C**). The average rosette diameter of the *hy5*

mutant was about 6 cm, whereas the diameters of the WT and complemented lines were 4 and 3

cm, respectively, demonstrating the recovery of rosette size as well (**Figure 3D, Supplemental**

Figure S7A). Similarly, seeds of the *hy5* mutant also regained the size similar to WT after

complementation (**Figure 3E and 3F**). These results suggest that NtHY5 could complement the

phenotypic function of the AtHY5 in the *hy5* mutant.

NtHY5 can compensate for the loss at transcript and metabolite levels in *Arabidopsis hy5*

mutant

To understand the function of NtHY5 at transcript and metabolite levels, the expression of flavonol

biosynthetic pathway genes was studied in WT, *hy5* and NtHY5;*hy5* complemented seedlings. The

analysis suggested that the expression of the NtHY5 in the *hy5* mutant background restored the

expression of genes associated with the phenylpropanoid pathway (*AtCHS*, *AtCHI*, *AtFLS* and

AtDFR) (**Figure 3G**). The levels of flavonols (kaempferol and quercetin) were also restored to WT

in NtHY5;*hy5* lines compared with the *hy5* mutant **(Figure 3H)**. To determine the responsiveness

of complemented lines towards light, expression of genes associated with flavonoid biosynthesis

was study in 5-day-old dark-grown seedlings subsequently exposed to light for 3h. The transcript

level of structural genes increased on exposure to light for 3h in complemented lines. By contrast,

hy5 mutant showed inhibited light-dependent induction of gene expression **(Figure 3I)**. In addition, expression of MYB family transcription factors also enhanced in NtHY5 complemented lines after 3h of light induction compared with *hy5* mutant seedlings **(Supplemental Figure S7B)**.

The phenylpropanoid pathway synthesizes various secondary metabolites, including flavonols, flavones, flavonols, isoflavones, flavan-3-ol, proanthocyanidins (PAs), anthocyanin and lignin. Phenylalanine is reported to be a common precursor for all of them (Vogt et al., 2010). Thus, we studied the effect of NtHY5 on lignin and anthocyanin production in *Arabidopsis* complemented lines. For lignin staining, the stem cross-sections of the WT, *hy5*, and NtHY5;*hy5* were used. The results showed a decreased lignification in the interfascicular and vascular tissues (**Supplemental Figure S7C**). Additionally, compared to WT and *hy5* mutant plants, NtHY5;*hy5* had a shorter vascular bundle in length (**Figure 3J**) this suggested the vascular damage due to lesser lignin. As AtHY5 is known to be involved in anthocyanin and chlorophyll biosynthesis (Oyama et al., 1997; Chattopadhyay et al., 1998), we estimated anthocyanin and chlorophyll contents of WT, *hy5*, and NtHY5;*hy5* seedlings. In 7-day-old light-grown seedlings, the total chlorophyll content of NtHY5;*hy5* was higher than WT and *hy5* (**Supplemental Figure S7D**)*.* The anthocyanin content in NtHY5;*hy5* was higher than *hy5* seedlings but found to be comparatively lesser than WT (**Supplemental Figure S7E**). These findings imply that NtHY5 can compensate for the loss in the *athy5* mutant at the transcript and metabolite levels.

Nicotine and flavonoid production in tobacco is regulated by NtHY5

To gain a deeper insight into NtHY5 function and its involvement in nicotine and flavonoid, we developed NtHY5 mutant plants using the CRISPR–Cas9 approach and NtHY5 overexpression (NtHY5OX) plants. To mutate NtHY5, we used guide RNAs (gRNAs) targeting the coding region of NtHY5 (**Supplemental Figure S8A**). The nucleotide sequence analysis of the mutants revealed deletions and insertions in the coding regions of NtHY5 (**Supplemental Figure S8B**). The mutants (*NtHY5^{CR}* plants) showed frame-shift mutations leading to truncated peptides (**Supplemental Figure S8C**). In 20-day-old light-grown seedlings where roots were covered to inhibit light illumination to maintain natural condition (**Figure 4A**), qRT-PCR expression analysis revealed the downregulation of NtHY5 in mutants, while overexpression seedlings showed a higher expression (**Figure 4B**). We also analyzed the expression of nicotine pathway genes (*NtODC, NtPMT, NtMPO, NtQPT* and *NtA622*). The analysis suggested a significant downregulation of

genes involved in nicotine biosynthesis in *NtHY5^{CR}* seedlings, whereas overexpression seedlings

showed an increase in expression in comparison to WT (**Figure 4C**). Additionally, structural genes

of the phenylpropanoid pathway, such as *NtCHS, NtCHI, NtFLS*, and *NtDFR,* and regulatory gene,

NtNtMYB12, were downregulated in *NtHY5^{CR}* seedlings and the opposite was observed for

overexpression lines (**Supplemental Figure S9A and B**).

Phytochemical analysis suggested up to 50% reduction and 75% increase in nicotine level in 20-

day-old seedlings in *NtHY5^{CR}* and NtHY5OX lines, respectively, compared to WT (**Figure 4D**).

The levels of flavonols (kaempferol, quercetin, rutin and CGA) and anthocyanin in mutant and

overexpression seedlings also got modulated (**Supplemental Figure S10A-D**; **Supplemental**

Figure S11A and B) in *NtHY5^{CR}* and NtHY5OX seedlings compared to WT. Thus, the decline in

nicotine and flavonoid content in mutant lines and the increase in overexpression lines suggest an

involvement of the NtHY5 in the biosynthesis of these molecules in tobacco seedlings.

NtHY5 regulates light-dependent expression of genes involved in nicotine biosynthesis

In order to determine the involvement of light on the expression of genes involved in nicotine and

flavonoid biosynthesis, *NtHY5^{CR}* and NtHY5OX seedlings were grown in the continuous dark for

10 days, followed by exposure to light for 3h. The expression analysis suggested a significantly

increased gene expression after 3h exposure to light to WT and NtHY5OX seedlings (**Figure 4E**

and F). However, such an increase in the expression was not observed in *NtHY5^{CR}* seedlings. We

also analysed the expression of NtHY5 and flavonoid pathway genes and results suggested that

exposure to 3h of light can induce the expression significantly in NtHY5OX but not in *NtHY5^{CR}*

plants (**Supplemental Figure S12A and B**). This induction led to enhanced accumulation of

nicotine in the plants **(Figure 4G)**. These observations suggested NtHY5-dependent light-

responsiveness of genes involved in nicotine and flavonoid biosynthesis in tobacco.

NtHY5 of shoot regulates nicotine biosynthesis in root

To determine light- and NtHY5-dependent regulation of nicotine biosynthesis in roots (which are

not usually exposed to light), grafting experiment using shoot (scion) and root (rootstocks) of 7-

day-old WT, NtHY5OX and *NtHY5^{CR}* seedlings in different combinations were carried out. The

grafting experiments included NtHY5OX/*NTHY5^{CR}* (NtHY5OX shoot grafted over *NtHY5^{CR}* root),

NtHY5^{CR}/HY5OX (*NtHY5^{CR}* shoot grafted over NtHY5OX root), NtHY5OX/NtHY5OX

(NtHY5OX shoot grafted over NtHY5OX root), *NtHY5^{CR}/NtHY5^{CR}* (*NtHY5^{CR}* shoot grafted over

NtHY5^{CR} root), NtHY5OX/WT (NtHY5OX shoot grafted over WT root), *NtHY5^{CR}*/WT (*NtHY5^{CR}*

shoot grafted over WT root) and WT/WT (WT shoot grafted over WT root). After three days of

grafting, the roots of seedlings were covered to grow them in the dark, followed by sampling of

roots after two weeks (**Supplemental Figure S13A**). The expression analysis of genes and

nicotine content were measured in two ways; one in which HY5OX/HY5OX were taken as control

and another with WT/WT as control.

Expression analysis of NtHY5 (**Supplemental Figure S13B**) and nicotine pathway genes for

different grafting combinations suggested a significantly higher expression when NtHY5OX was

used as a scion (NtHY5OX/*NtHY5^{CR}*). The highest expression was observed when NtHY5OX was

used for both scion and rootstock (NtHY5OX/NtHY5OX) and the lowest when *NtHY5^{CR}* were

used as both scion and rootstock (*NtHY5^{CR}/NtHY5^{CR}*) (**Figure 5A**). Similar to the expression

pattern, the accumulation of nicotine followed the pattern with the highest accumulation for

NtHY5OX/NtHY5OX grafted union followed by HY5OX/*HY5^{CR}*, whereas the lowest for

NtHY5^{CR}/NtHY5^{CR} grafted union followed by *NtHY5^{CR}*/NtHY5OX (**Figure 5B**). Additionally,

when WT/WT was taken as control, *NtHY5^{CR}*/WT showed the lesser expression of NtHY5 (**Figure**

5C) and nicotine pathway genes (**Figure 5D**) compared to NtHY5OX/WT. Similar to the

expression pattern, nicotine accumulation was also less in *NtHY5^{CR}*/WT compared to

NtHY5OX/WT (**Figure 5E**). These observations suggest that shoot NtHY5 may activate the root

local HY5 which in turn binds to promoter of nicotine pathway genes. Alternatively, shoot HY5

may directly regulate the expression of nicotine pathway genes and nicotine biosynthesis. Similar

to our study, previous studies have also demonstrated the movement of AtHY5 from shoot-to-root

to regulate itself and other targets (Chen et al., 2016; Sharma et al., 2022).

NtHY5 regulates nicotine biosynthesis in mature tobacco plants

Our abovementioned results using NtHY5OX and *NtHY5^{CR}* seedlings suggested that NtHY5

regulates nicotine biosynthesis in tobacco roots. We explored the impact of NtHY5 on nicotine

synthesis in mature plants. The expression analysis suggested increased expression of genes

involved in nicotine biosynthesis in NtHY5OX compared to *NtHY5^{CR}* and WT roots (**Figure 6A**

and B). At the metabolite level, leaves of NtHY5OX accumulated significantly higher nicotine

levels compared to WT and *NtHY5^CR* leaves. There was 92% and 70% reduction in nicotine content

in *NtHY5^CR* leaves compared to NtHY5OX and WT leaves, respectively (**Figure 6C**).

We also analyzed the effect of light-dark transition on nicotine biosynthesis using NtHY5OX and

NtHY5^CR plants. Light-grown mature tobacco plants were completely covered with black paper for

24 h before sampling for the analysis **(Figure 6D)**. Roots of both NtHY5OX and *NtHY5^CR* plants

showed reduced expression of nicotine pathway genes during the light-to-dark transition.

However, the reduction was substantially higher in *NtHY5^CR* plants than in NtHY5OX when

compared to WT (**Figure 6E**). The modulation in the expression of genes led us to determine

nicotine concentration in mature leaves. The analysis suggested a higher accumulation of nicotine

in NtHY5OX leaves than in *NtHY5^CR* compared to WT after the light-to-dark transition **(Figure**

6F). This suggested that dark negatively regulated nicotine synthesis in mature tobacco plants.

NtHY5 regulates growth and development of tobacco

Previous study suggested that flavonols can also function as an endogenous polar auxin

transport inhibitor to regulate shoot and root growth under various conditions (**Peer et al.,**

2004, Santelia et al., 2008) although the relation between hormone signaling and NtHY5 could

be the subject of further study and reduction in nicotine content leads to positive regulation of root

growth (**Hildreth et al., 2011**). We subsequently studied the effect of modulated nicotine and

flavonoids on the growth and development of tobacco. A comparative analysis between NtHY5OX

and *NtHY5^CR* transgenic lines suggested differences in the growth of NtHY5OX and *NtHY5^CR*

light-grown seedlings (**Supplemental Figure S14A**). Several previous studies have demonstrated

that AtHY5 regulates the hypocotyl length by repressing the expression of hypocotyl cell

elongation-related genes (Liu et al., 2013). To validate the role of NtHY5 in hypocotyl length

elongation in tobacco, we studied the difference in hypocotyl length in NtHY5OX and *NtHY5^CR*

plants. The analysis suggested that the hypocotyl length of *NtHY5^CR* mutant seedlings was

significantly higher compared to overexpression lines (**Supplemental Figure S14B**). At the same

time, the hypocotyl length of the dark-grown overexpression lines was significantly higher

compared to *NtHY5^CR* mutant plants (**Supplemental Figure S14C and D**). We also measured the

root length and bending angle of 10-d-old light-grown seedlings. Analysis suggested the higher

root length of *NtHY5^CR* seedlings compared to WT and NtHY5OX, whereas the root bending angle

of NtHY5OX seedlings was higher compared to WT and *NtHY5^{CR}* plants (**Supplemental Figure**
S14E and **F**).

We also observed the modulation in seed size; *NtHY5^{CR}* plants produced smaller seeds
(**Supplemental Figure S15 A and B**) compared to NtHY5OX and WT (**Supplemental Figure**
S15C). In addition to these factors, we observed the difference in height and chlorophyll content
of the young plants. The plant height of 40-days-old *NtHY5^{CR}* plants was significantly higher than
NtHY5 overexpression and WT plants (**Supplemental Figure S15D and E**), whereas chlorophyll
was found to be lesser in *NtHY5^{CR}* plants compared to NtHY5OX and WT (**Supplemental Figure**
S15F).

NtHY5 enhances the salt stress tolerance in tobacco
As HY5 is known to play a role in abiotic stress tolerance, the response of NtHY5 under salt stress
was analyzed through leaf disc assay using NaCl (200 mM), and NtHY5OX, *NtHY5^{CR}* and WT
plants. After 3 days of salt stress, higher chlorosis was observed in *NtHY5^{CR}* leaves compared to
leaves of NtHY5OX and WT plants (**Supplemental Figure S16A and B**). Chlorosis affects the
survival of leaves by decreasing the chlorophyll content (Chen et al., 2019). Thus, the survival of
leaves was measured by analyzing the total chlorophyll content as well as chlorophyll a and b. A
higher chlorophyll content was observed in NtHY5OX leaves compared with *NtHY5^{CR}* and WT
leaves after 3 days of salt stress (**Supplemental Figure S16C**). It is well known that salt stress
leads to excessive accumulation of ROS, leading to cell death (Chen et al., 2019). Higher NBT and
DAB staining were observed in *NtHY5^{CR}* compared with NtHY5OX and WT (**Supplemental**
Figure S16D and E), respectively, suggesting the higher ROS formation in *NtHY5^{CR}* leaves, thus
above study concluded that NtHY5 might play a crucial role in salt stress tolerance in tobacco by
inhibiting ROS accumulation. A higher ROS accumulation results in lower antioxidant activity in
NtHY5^{CR} lines, whereas lower ROS accumulation leads to the higher antioxidant activity in
NtHY5OX lines compared to WT (**Supplemental Figure S17**).

Discussion

The nicotine biosynthesis pathway in terms of intermediate steps and enzymes is well explored
(Dewey and Xie, 2013). Additionally, several previous studies suggested the role of various

transcription factors belonging to different families, for example, helix–loop–helix (bHLH)
transcription factors (e.g. MYC2a/b), MYC2-type bHLH TFs NbbHLH1 and NbbHLH2,
AP2/ERF genes, NtORC1/ERF221 and NtJAP1/ERF10, and MYB (NtMYB305a), in regulating
nicotine biosynthesis (Song et al., 2011; Shoji et al., 2011; Shear et al., 2014; Liu et al., 2019; Bian
et al., 2021). In the past, other factors, including insect attack, wounding, and jasmonate (JA) have
been shown to be involved in the biosynthesis of nicotine (Imanishi et al., 1998; Shi et al., 2006;
Bozorov et al., 2017; Li et al., 2018; Liu et al., 2020). Previous study suggested the presence of
several G and GCC boxes in the promoter regions of nicotine biosynthesis genes (Zhang et al.,
2011). It has also been established that HY5 binds to G-box (CACGTG) in the promoters of light-
responsive genes. Therefore, we hypothesized that NtHY5 could regulate the nicotine production
in tobacco.

Previous studies in *Arabidopsis* and other plant species suggested HY5 as a master regulator of
photomorphogenesis and a key signaling component of light. It promotes photomorphogenesis
under a broad spectrum of wavelengths (Ulm et al., 2004; Lee et al., 2007; Nawkar et al., 2017).
However, the functional characterization of tobacco HY5 is not done yet. In this study, we have
identified the NtHY5 and demonstrated that it regulates nicotine biosynthesis in tobacco, extending
our understanding of NtHY5-mediated regulation of secondary plant metabolism. The present
work shows the involvement of light signaling component in nicotine and flavonoid regulation of
tobacco as accumulation of nicotine as well as flavonols also found to be higher in light-grown
seedlings compared to dark-grown seedlings (**Figure 1D and Supplemental Figure S1B**). WT
dark-grown seedlings showed the induction in the expression of nicotine pathway genes on
exposure to light and continuously increased with increase in time interval till saturation (**Figure
1E**). Thus, above results led us to hypothesize the involvement of NtHY5 as one of the potential
factors responsible for light-mediated nicotine and flavonoid regulation in tobacco. A higher
expression of identified NtHY5 in light-grown seedlings as well as our further analysis in which
we have exposed 10-d-old dark-grown seedlings to light of different time interval suggested the
significant induction in expression of nicotine pathway genes (**Figure 1E and F**) as well as
indicated, NtHY5 as light-responsive key regulatory factor in nicotine and flavonoid biosynthesis
(**Supplemental Figure S1**) in tobacco.

Prior studies have established that HY5 acts as a transcription factor that predominantly binds to
the LREs like *ACGT*-containing *cis*-element (e.g., *G-box* and *T/G-box*) and controls the
expression of numerous target genes in response to light signals (Song et al., 2008). To investigate
the binding of NtHY5 to the promoter of nicotine pathway genes, *in silico* analysis was carried out
and the result revealed the presence of cis-acting LREs upstream of the transcription start site on
promoters of PMT, QPT, ODC and NtMYB12. Similar to our result, there are previous studies,
suggesting the presence of GCC and G-box elements in the promotor regions of nicotine pathway
genes (Todd et al., 2010; Shoji and Hashimoto, 2011; Zhang et al., 2012). Our EMSA analysis
validated NtHY5 intraction with the promoters of genes involved in nicotine pathway and
flavonoid regulatory transcription factor, suggesting the involvement of NtHY5 in nicotine and
flavonoid biosynthesis (**Figure 2 and Supplemental Figure S6**).

Previously, several HY5 orthologs from different plant species have been identified and
characterized (Nishimura et al., 2002; Burman et al., 2018). OsZIP48 is reported as a functional
ortholog of AtHY5 in rice as it was able to rescue the phenotype of *Arabidopsis hy5* mutant plants
after complementing with OsZIP48. Similarly, overexpression of STF1, the HY5 ortholog in
soybean, can fully complement the *hy5* phenotype (Song et al., 2008). As NtHY5 was identified
for the first time in the current study thus complementation study has been done to identify whether
NtHY5 can fully complement the *Arabidopsis hy5* or not? The current study demonstrated that
NtHY5 partially complemented *Arabidopsis hy5* mutant for phenotypes such as hypocotyl length,
root length, seed size and rosette diameter. This study also indicated the fully complementation at
the level of expression of structural and regulatory genes of flavonoid biosynthesis and
accumulation of metabolites, such as flavonoid, anthocyanin, lignin and chlorophyll in *NtHY5;*
hy5 compared to *hy5* (**Figure 3 and Supplemental Figure S7).** These findings suggest that NtHY5
had a conserved function across different species, and it regulates flavonoid biosynthesis in
complemented lines of *Arabidopsis*.

To decipher the molecular mechanism through which light regulate nicotine and flavonoid
biosynthesis in tobacco, CRISPR/Cas9-mediated knockout and overexpression tobacco plants for
NtHY5 were developed. As per previous studies, nicotine synthesis takes place in the root, which
is subsequently transported to the leaf via the xylem (Hayashi et al., 2020). Thus, in the current
study, analysis of nicotine content and pathway gene expression was carried out at seedling (roots

were kept in the dark) as well as mature plant stage in mutant and overexpressed plants of NtHY5

compared to WT. The expression analysis of seedlings as well as roots of mature plants indicated

a higher expression of nicotine pathway genes in seedlings and roots of NtHY5OX mature plants,

whereas expression was lower in seedlings and roots of *NtHY5^{CR}* plants in comparison to WT

(**Figure 4C, 6A and 6B**). Quantitative estimation of nicotine content in seedlings suggested up to

50% reduction and four-fold increase in nicotine level in *NtHY5^{CR}* and NtHY5OX lines,

respectively. At the mature plant stage, leaves of NtHY5OX accumulated 92% higher nicotine

levels compared to WT and *NtHY5^{CR}* leaves showed 70% reduction in nicotine content in

comparison to WT. Thus, current study reveals that NtHY5 regulates nicotine synthesis not only

at seedling but also at the mature plant stage. The CRISPR/cas9 mutated NtHY5 could not bind to

promoters of nicotine pathway genes, resulting into a lower accumulation of nicotine. The current

study also indicated that NtHY5 is induced by light supported by dark-to-light induction analysis.

In OE lines, NtHY5 regulates its own expression in light, while in dark it undergoes degradation

and is unable to regulate its own expression. This leads to a lower expression in the dark compared

to seedlings exposed to 3h light after 10-day growth in dark **(Supplemental Figure S12 A)**. Less

nicotine-containing edited tobacco plants will be beneficial for mankind by helping adults to

switch instead of continuing smoking cigarettes because nicotine is part of what makes cigarettes

addictive.

As tobacco roots are known to synthesize nicotine (Hayashi et al., 2020) and are typically in the

dark, we studied how shoot NtHY5 regulates nicotine production in roots. We carried out grafting

experiments in which several hypocotyl graft chimeras with roots covered in darkness were

examined. We observed that a NtHY5OX scion facilitated an enhanced nicotine accumulation (as

opposed to a *NtHY5^{CR}* scion) (**Figure 5, Supplemental Figure S13**), indicating that a shoot-

derived and mobile NtHY5 signal controls nicotine biosynthesis in roots, controlling both by

directly modulating nicotine pathway genes and automatically activating local HY5 in the roots.

Similar to our findings, a previous study suggested that *Arabidopsis* HY5 regulates growth in

response to light, is a shoot-to-root mobile signal that mediates light promotion of root growth and

nitrate uptake (Chen et al., 2016).

Our tissue-specific expression analysis also suggested the lower expression of nicotine pathway

genes in leaves compared to roots. In the past, several studies have also suggested the biosynthesis

of nicotine in the cortical cells of roots through various enzymatic steps, which is further

transported to leaves vacuoles via xylem (Shoji et al., 2009; Dewey et al., 2013; Hayashi et al.,

2020; Shoji et al., 2021). This was the reason for using roots for all the expression analysis whereas

young and mature leaves were used for the nicotine quantitative analysis. However, it is possible

that NtHY5 might regulates nicotine biosynthesis in the leaves but to a lesser extent.

In case of $NtHY5^{CR}$/NtHY5OX grafted seedling, no functional HY5 is present in shoot which can

move to root to activate root local HY5; thus, expression of nicotine pathway genes is lower. In

addition, $NtHY5^{CR}$/NtHY5OX grafted seedling, though HY5 in root is expressing under control of

CaMV35S promoter, the possibility of degradation of this by root-specific COP1 cannot be ruled

out. This might be one of the reasons for the low nicotine accumulation in $NtHY5^{CR}$/NtHY5OX

grafted seedling. NtHY5 transcript levels are very high in roots of NtHY5OX/$NtHY5^{CR}$ grafted

seedlings. There could be a possibility that transcript levels observed are from the mutated gene or

shoot NtHY5 transcripts which move to the roots. There are a few previous studies that suggest

that transcripts could also move between scion and rootstock. For example, CmGAI transcripts in

tomato, BEL5 transcripts in tomato and transcripts in grapes have been demonstrated to be present

from scion to rootstock (Yin et al., 2022; Haywood et al., 2005; Banerjee et al., 2006; Yang et al.,

2015).

We also demonstrated the regulation of flavonols and anthocyanin biosynthesis by NtHY5 as they

are found to be reduced in NtHY5 knockout seedlings in comparison to WT (**Supplemental**

Figure S10 and S11). We found a significant increase in flavonol content, but interestingly

decrease in anthocyanin content in NtHY5OX lines, suggesting the diversion of flux towards

flavonol biosynthesis. Thus, these results provided a deeper insight of light-mediated regulatory

mechanism involved in nicotine and flavonoid biosynthesis in tobacco.

HY5 and specific MYBs are known to be involved in plant growth and development (Zhong et al.,

2008; McCarthy et al., 2009; Gangappa et al., 2016; Misra et al., 2010). OsbZIP48 ortholog is

reported to cause a reduction in cell size, thus reducing the plant height of OsbZIP48

overexpression transgenic plants (Burman et al., 2018). HY5 negatively regulates hypocotyl

elongation by suppressing auxin signaling (Cluis et al., 2004; Sibout et al., 2006). Our results

suggested that NtHY5 regulates flavonol content in tobacco. Similarily, previous study suggested

that flavonols can function as an endogenous polar auxin transport inhibitor to regulate shoot and

root growth under various conditions (Peer et al., 2004; Santelia et al., 2008). Thus, we next
determined the impact of NtHY5 on plant growth parameters since, no data is available on the
involvement of NtHY5 in the growth and development of tobacco via regulating secondary
metabolites. We conducted a comparison study between the transgenic NtHY5OX and $NtHY5^{CR}$
lines. The results suggested that modulation in the NtHY5 expression causes changes in nicotine
and flavonoid concentration leading to variation in various morphological characteristics, with
contrasting phenotypes between NtHY5OX and $NtHY5^{CR}$ plants. Hypocotyls length of mutants
was significantly longer than overexpressed and WT seedlings grown in light for 10 days.
Interestingly, shorter hypocotyl length was observed in mutants in comparison to overexpress and
WT in dark-grown seedlings (**Supplemental Figure S14 A-D**). This suggested that mutation in
NtHY5 might have eradicated the light-dependent repression of cell elongation and does not
promote cell elongation independent of the light stimulus.

In tobacco, one previous study suggested that RNAi silencing of NUP1, a plasma membrane-
localized nicotine transporter, causes a reduction in nicotine content which in turn regulates the
root growth of seedlings positively (Hildreth et al., 2011). In our study, we demonstrated that low
nicotine content, accompanied by low flavonols, causes an increase in root length in $NtHY5^{CR}$
seedlings compared to overexpression lines and WT. It is earlier reported that the asymmetric
accumulation of flavonoids in the root tips of *Arabidopsis* affects the direction of root growth (Wan
et al., 2018). In agreement, root bending angle of 10-day-old light-grown NtHY5 overexpression
and $NtHY5^{CR}$ seedlings of the tobacco was measured. The result suggested the higher root bending
angle in NtHY5OX lines in comparison to $NtHY5^{CR}$ and WT (**Supplemental Figure S14F**).
Interestingly, analysis also showed the modulation in tobacco seed size and height of the 40-d-old
plants (**Supplemental Figure S15B and FF**). Thus, results indicated that NtHY5 plays a pivotal
role in tobacco growth and development through auxin signaling, although the relation between
hormone signaling and NtHY5 could be the subject of further study.

Our study suggested the regulation of NtMYB12 by the NtHY5 through interaction with its
promoter. One recent study demonstrated that overexpression of NtMYB12 causes more flavonol
accumulation and a higher ROS scavenging property, enhancing tobacco tolerance to low Pi stress
(Song et al., 2020). There are some studies which suggest that tobacco responds to salt stress by
increased activity of antioxidant enzymes (Sun et al., 2020. In our study, antioxidant activity was

measured, and the analysis suggested higher activity in overexpression lines compared to WT and

NtHY5^{CR} plants. The high antioxidant activity led to lesser chlorosis in salt-treated young leaves

of NtHY5OX lines compared with *NtHY5^{CR}* **(Supplemental Figure S16A-C)**, suggesting that

higher NtHY5 expression leads to higher ROS scavenging ability in overexpression lines. As

tobacco *NtHY5^{CR}* mutants were salt-sensitive, thus more ROS accumulated in salt-treated mutant

leaves **(Supplemental Figure S16D and E)**. Thus, our study suggests that NtHY5 promotes salt

stress tolerance at least partially through the antioxidant activity of flavonoids **(Supplemental**

Figure S17). To understand whether nicotine also has the antioxidant activity, a detailed regulatory

mechanism needs to be studied.

We propose the model (**Figure 7**), which depicts that HY5 is a shoot-root mobile signal that

mediates the light-regulated synthesis of nicotine and flavonoids. This coupling is achieved via the

regulation of nicotine and flavonoid pathway genes through interaction of the NtHY5 with the G-

Boxes present within the promoters of different genes. NtHY5 is also found to modulate growth

and development, as evident through modulated seed size, plant height and root length.

Additionally, NtHY5 provides salt tolerance via diminishing ROS accumulation evident through

DAB and NBT staining. Thus, this study provides an essential insight into a molecular strategy to

develop tobacco having significantly low nicotine through manipulating the NtHY5 gene of

tobacco.

Materials and Methods

Isolation and sequence analysis of NtHY5. To identify the ortholog of AtHY5 in tobacco,

BLAST searches were performed on National Center for Biotechnology Information

(http://www.ncbi.nlm.nih.gov/BLAST). The sequence was retrieved and amino acid sequence was

analyzed for the leucine rich bZIP Zinc finger domain and VPE domain using multallignment tool.

Conserved domain database (CCD) was employed to ensure the bZIP family of the identified

NtHY5 protein. Promoter sequences (~1.5 kb upstream of translation start site) of NtMYB12 gene

was retrieved from the respective databases AGRIS (Palaniswamy et al., 2006) and PlantCare

(Lescot et al., 2002), which was utilized for the scanning of light-responsive cis-elements (LREs).

Homologs of AtHY5 that belongs to different families were identified by using the National Center

for Biotechnology Information BLASTn tool. The genes retrieved were checked for the presence

of the requisite domains using InterProScan (Quevillon et al., 2005) (http://

www.ebi.ac.uk/Tools/pfa/iprscan/). A phylogenetic tree was constructed using the Neighbor-

Joining (NJ) algorithm in MEGA6.0 software (Kumar et al., 2008).

Plant material, growth conditions and treatments. Leaves of *Nicotiana tabacum* cv. *Petit*

Havana (NtPH) have been used for raising the transgenic plants. Tobacco plants were grown in

glass house at 25°C±2°C and 16 h/8 h light-dark photoperiods for harvesting of tissues belonging

to different developmental stage. Seeds were surface sterilized using 70% EtOH for 1 min and

then dipped in 50% bleach for 10 min, washed repeatedly with autoclaved Milli-Q water and grown

with roots covered in dark for 20 days. For dark treatment, seeds were surface sterilized and grown

for 10 days in absence of light, 24°C temperature, and 60% relative humidity on one-half-strength

Murashige and Skoog (MS) medium (Sigma) plates after stratification for 2 to 3 d at 4°C in the

dark and subsequently transferred to culture room with 16 h/8 h light-dark photoperiods and 150–

180 mm m^{-2}s $^{-1}$ light intensity. For light-to-dark treatment glass house grown 5-month old tobacco

plants were completely covered with black paper for 24h before sampling. In mature plants, three

biological replicates of each line were grown in glasshouse till maturity. Plants were then uprooted

and the roots were cut from the root-shoot junction and sampled for further analysis. In seedlings,

seeds were surface sterilized and grown with roots covered in dark for 20 days following which

the roots were cut from root-shoot junction in dark room and sampled for further analysis.

Arabidopsis (Col-0) was used as the wild-type plant in complementation study. Seeds were surface

sterilized and after stratification transferred to a growth chamber under controlled conditions of

16-h-light/8-h-dark photoperiod cycle, 22°C temperature, 150 to 180 μmol m^{-2}s^{-1} light intensity,

and 60% relative humidity for 10 days unless mentioned otherwise.

Plasmid construction and generation of *Arabidopsis* and tobacco transgenic and mutant

plants. To overexpress NtHY5, the NtHY5 was amplified from the single-stranded

complementary DNA (cDNA) library prepared from 10-day-old tobacco seedlings through PCR

using specific primers. The full-length open reading frame of the NtHY5 cDNA, under the control

of cauliflower mosaic virus 35S promoter in the binary vector pBI121 (Clontech, USA) was

transferred into *Agrobacterium tumefaciens* strain GV3101 and used to transform *Arabidopsis* by

floral dip method and tobacco plants by leaf disk method (Horsch et al., 1985). Several transgenic

tobacco lines constitutively expressing NtHY5 cDNA were selected on the basis of RT-PCR.

Seeds were harvested, sterilized and plated on solid half strength MS medium supplemented with

100 mg/l kanamycin. Antibiotic resistant plants were shifted to the glass house and grown until

maturity.

For generation of mutant plants of NtHY5, CRISPR/Cas9 technology was used. A 20-bp gRNA

was selected from the identified coding region. NtHY5 gRNAs was cloned into the binary vector

pHSE401 using the BsaI restriction site (Xing et al., 2014). This vector contains Cas9

endonuclease-encoding gene under dual CaMV35S promoter as well as genes encoding neomycin

phosphotransferase and hygromycin phosphotransferase as selection markers. All the constructs

were sequenced from both the orientations using plasmid-specific forward and vector reverse

primers after that transferred to GV3101 and then transformed to tobacco.

Gene expression analysis. Total RNA was isolated using a Spectrum Plant Total RNA kit (Sigma

Aldrich) following the manufacturer's instruction. For qRT-PCR analysis, 2 µg of DNA free RNA

was reverse transcribed using a RevertAid H minus first-strand cDNA synthesis Kit (Fermentas)

according to the manufacturer's instructions. qRT-PCR was performed using Fast Syber Green

mix (Applied Biosystems) in a Fast 7500 Thermal Cycler instrument (Applied Biosystems).

Expression was normalized using Tubulin (NtTUB, XM_009622503; AtTUB, NM_125664) and

analyzed through the comparative CT method (Livak and Schmittgen 2001). For each experiment,

three independent RNA preparations and total nine technical replicates (three for each RNA

preprations) were used. The oligonucleotide primers used to study expression of different genes

were designed using the Primer Express 3.0.1 tool (Applied Biosystems) and information is

provided in Supplementary Table S5.

Tobacco grafting. Grafting between wild type, NtHY5OX, *NtHY5^{CR}* lines was performed

according to method described by Rus et al. (2006). Seedlings grown on one-half-strength MS

media for 7 days were used for grafting. Cuts between the scion and stock were made at an angle

of 90° and were then placed in desired combinations. The grafted seedlings remained on the media

plate for a period of 3 days to allow the formation of the graft union. Roots of the grafted seedlings

were further covered in darkness for 2 more weeks prior to sampling. For further experiments,

only the grafted seedlings were used. Seedlings which did not produce proper graft (scion and

rootstock did not bind with each other and rootstock ultimately died) were discarded.

Extraction and quantitative estimation of nicotine. Nicotine extraction was carried out with slight modifications (Khan et al., 2017). About 300 mg freeze-dried tissue samples for mature leaves and 300 mg fresh weight tissue for seedlings/ grafted seedlings were grinded to powder and extracted in 2.0 mL 75 % methanol. The extract was ultrasonicated for 60 min, and then centrifuged (14,000 rpm) for 10 min. The supernatant was concentrated to dry powder and then dissolved in 1mL HPLC grade methanol. Extracts were filtered through 0.2 mm filter (Millipore, USA) and subjected to HPLC analysis. Lab Solutions software (Shimadzu) was used for the quantification of nicotine through HPLC. Column used was Phenomenex Luna C18(2) column (250 mm x 4.6 mm x 5μ). Mobile phase consisted of water: methanol: 0.1 M buffer acetate (pH4.5): acetonitrile: acetic acid in the ratio 74: 3: 20: 2:1 (Solvent A) with pH adjusted to 4.2 with triethylamine. Solvent A was maintained at 100% throughout the run with a slight modification in the gradient of flow rate of 1ml/min in 0-15min, 1.0-1.5ml/min in 15-15.5min, 1.5ml/min in 15.5-24.5min, 1.5-1.0ml/min in 24.5-25min and 1ml/min in 25-30min. The samples were analysed by HPLC–PDA and chromatograms were recorded at 254nm (Pereira et al., 2001).

Extraction and quantitative estimation of flavonols. Extraction of flavonols was carried out by grinding the plant material (300mg) into the fine powder in liquid N_2 and then extracted with 80% methanol overnight at room temperature with brief agitation. The extract was hydrolyzed with an equal amount of 6 N HCl at 70°C for 40 min followed by addition of an equal amount of methanol to prevent the precipitation of the aglycones (Jiang et al. 2015). Extracts were filtered through 0.2 mm filter (Millipore, USA) before HPLC. For non-hydrolyzed extracts, samples were extracted as described previously (Pandey et al., 2014). The mobile phase was a gradient prepared from 0.05% (v/v) ortho-phosphoric acid in HPLC-grade water (component A) and methanol (component B). Before use, the components were filtered through 0.45-mm nylon filters and de-aerated in an ultrasonic bath. The gradient from 25 to 50% B in 0–3 min, 50 to 80% B in 3–18 min, 80 to 25% B in 25 min and 25% B in 30 min was used for conditioning of the column with a flow rate of 1 ml/min. All the samples were analysed by HPLC–PDA with a Waters 1525 Binary HPLC Pump system comprising PDA detector (Niranjan et al., 2011).

Extraction of total anthocyanin, chlorophyll and lignin staining. For anthocyanin, 300 mg tissue were boiled for 3 min in extraction solution (propanol: HCl: H2O, 18:1:81), and incubated at room temperature in the dark for at least 2 h. Samples were centrifuged, and the absorbance of the supernatants was measured at 535 and 650 nm. Anthocyanin content was calculated as

$$\text{Anthocyanin content} = (A_{535} - 2.2 \times A_{650}) / \text{g FW.}$$

Chlorophyll content was measured by using the 300mg tissue of 10-d-old WT and transgenic

seedlings. Sample was crushed in 2 ml (80 %, v/v) chilled acetone. Homogenized tissues were

centrifuged at 7,800g for 10 min. Supernatant was collected and absorbance was recorded at 663,

645, 510 and 480 nm. Total chlorophyll content was calculated in mg g^{-1} fresh wt using formula

(Machlachalan and Zalik, 1963).

To visualize lignified cells in stems, hand-cut sections were stained using phloroglucinol (Sigma-

Aldrich) for 1 min and visualized on a Leica DM2500 microscope (Sharma et al., 2020).To

measure the length of vascular bundle, automated length measuring scale of Leica DM2500 was

used.

Determination of total antioxidant activity. A method with slight modification was used to

determine the antioxidant activity. DPPH (1, 1-diphenyl-2-picrylhydrazyl free radical) assay was

used in determining the free radical scavenging activity (Wong et al., 2006). In brief, 0.1 mM

solution of DPPH was prepared in methanol. The initial absorbance of DPPH in methanol was

measured at 515 nm and did not change throughout the period of assay. An aliquot of the samples

(50–100 ml) was used to make up volume to 1.5 ml with DPPH solution. The sample was incubated

at room temperature for 30 min. The absorbance of the samples was recorded at 515 nm using

Ultraspec 3000 UV/Vis spectrophotometer (Pharmacia Biotech Ltd., Cambridge, CB4, 4FJ, UK).

Trolox was used to make the calibration curve using different dilutions. The antioxidant capacity

based on the DPPH free radical scavenging ability of the extract was expressed as mM Trolox

equivalents per gm of plant material on fresh weight basis.

Expression and purification of NtHY5. The complete open reading frame of NtHY5 was

amplified and cloned into pET-28b(+) vector (Novagen, Germany). The construct was transformed

into *E. coli* BL21 (DE3) pLysS (Invitrogen, USA) for prokaryotic expression, induced by IPTG.

Recombinant 6X-Histagged HY5 was purified using a Ni-NTA column (Nucleopore, India) and

the eluted and was quantified through Bradford assay.

Electrophoretic mobility shift assay. The 2nd generation DIG Gel Shift EMSA kit (Roche, USA)

was used to label the probes with digoxigenin as per manufacturer's instructions. Labelled probes

were incubated at 21°C for 30 min in binding buffer [100 mM HEPES (pH 7.6), 5 mM EDTA, 50

mM $(NH4)_2SO_4$, 5 mM DTT, Tween 20, 1% (w/v), 150 mM KCl] with or without recombinant

protein (500 ng). Unlabeled probes were added to the reaction solution in increasing concentrations

to test specific binding. The binding reaction was resolved on a 6% polyacrylamide gel in 0.5X

TBE (pH 8.0) buffer and was semi-dry blotted (Transblot, BIO-RAD, USA) onto a positively

charged nylon membrane (BrightStar, Invitrogen, USA) followed by UV cross-linking. The

membrane was finally incubated with CSPD chemiluminescent solution and exposed to X-ray blue

film (Retina, India).

Salt stress treatment. Salt stress analysis was performed by using the young leaves from 2^{nd} and

3^{rd} nodes from 1-month-old *NtHY5OX*, *NtHy5CR* and WT plants to determine the comparative

response during the leaf discs' salt assay. These leaves were transferred into salt solution of 200

mM. The treatment was carried out at 25 °C for 3 days. In order to determine the effect of salinity

stress, survival of leaves and chlorophyll content of NaCl treated leaves were analysed. All the

data were recorded from 0 to 3 day of NaCl treatment. The experiment was repeated three times.

Nitro blue tetrazolium (NBT) and 3,3′-diaminobenzidine (DAB) staining. Superoxide (O_2-)

staining was performed by infiltration by the nitro blue tetrazolium (NBT) method (Liu et al.,

2018; Song et al., 2019) with some modification. WT as well as transgenic seedlings were grown

for 10 days on half strength MS media then transferred to 200 mM salt solution for 3h. After 3

hours of salt stress seedlings were dipped in 0.5 mg/ml NBT solution for 2h followed by removal

of chlorophyll by heating at $70^{o}C$ temperature until chlorophyll is removed. Leaves were directly

visualized for NBT staining.

Hydrogen peroxide (H_2O_2) staining of 10-day light grown tobacco seedlings was performed by

staining with 3,3′-diaminobenzidine (DAB) using an adaptation of previous methods (Zhao et al

2016). WT as well as transgenic seedlings were grown for 10 days on half strength MS media then

transferred to 200 mM salt solution for 3h. After 3 h of salt stress, seedlings were dipped in 1mg/ml

DAB solution for 5 hours. Following the incubation, DAB staining solution was replaced with

bleaching solution (ethanol: acetic acid: glycerol = 3:1:1). After that chlorophyll was removed by

heating at $90-95^{o}C$ temperature. Leaves were directly visualized for DAB staining.

Statistical analysis. Data are plotted as means±SE (n=3) in the figures. Data of flavonol and

nicotine accumulation were evaluated for statistical significance using the Two-tailed Student's t-

test using Graphpad prism 5.01 software. Asterisks indicate significance levels with (* $P < 0.1$; **

$P < 0.01$; *** $P < 0.001$) as confidence intervals.

Author contributions

P.K.T. designed and supervised this study. D.S. designed and performed most of the experiments; HS, NS and SD provided help in conducting various experiments; P.K.T. analyzed the data; D.S., P.K.T. wrote the article; All authors read, contributed, and approved the article.

Acknowledgment

P.K.T. acknowledges CSIR and DBT, New Delhi, for financial support in the form of projects on pathway engineering and genome editing. P.K.T. also acknowledges Department of Science and Technology, New Delhi for JC Bose National Fellowship. DS acknowledge University Grant Commission, New Delhi for Senior and Junior Research Fellowship respectively.

Competing interests: Authors declare no competing interest.

Supplemental Data

Supplemental Figure S1. Estimation of flavonol content in WT tobacco.

Supplemental Figure S2. Sequence analysis of NtHY5 gene.

Supplemental Figure S3. Phylogenetic tree of NtHY5.

Supplemental Figure S4. Estimation of flavonol content in WT tobacco.

Supplemental Figure S5. Tissue-specific expression of NtHY5 and nicotine pathway genes.

Supplemental Figure S6. In vitro interaction between NtHY5 and the ACGT motif of G-BOX present at the promoters of *NtMYB12*.

Supplemental Figure S7. Complementation of *Arabidopsis hy5* mutant by NtHY5.

Supplemental Figure S8. Position of gRNA in NtHY5 sequences and amino acid sequence of truncated protein in mutants.

Supplemental Figure S9. NtHY5 modulates the gene expression of flavonoid biosynthesis genes.

Supplemental Figure S10. NtHY5 modulates flavonoid content in tobacco.

Supplemental Figure S11. NtHY5 leads to alteration in accumulation of anthocyanin content.

Supplemental Figure S12. Light responsiveness of NtHY5 and flavonol pathway genes.

Supplemental Figure S13. NtHY5 moves from shoot to root to regulate nicotine biosynthesis.

Supplemental Figure S14. NtHY5 regulates various morphological characterstics of tobacco seedlings.

Supplemental Figure S15. Comparative analysis of HY5OX vs. *NtHY5^{CR}*.

Supplemental Figure S16. Effect of salt stress on NtHY5 transgenic lines of tobacco.

Supplemental Figure S17. Measurement of antioxidant activity in NtHY5OX and *NtHY5^{CR}* lines
in comparison to WT.

Supplemental Table S1. Putative cis-acting light responsive elements in NtPMT promoter.

Supplemental Table S2. Putative cis-acting light responsive elements in NtQPT promoter.

Supplemental Table S3. Putative cis-acting light responsive elements in NtODC promoter.

Supplemental Table S4. Putative cis-acting light responsive elements in NtMYB12 promoter.

Supplemental Table S5. Oligonucleotides used for development of constructs and expression
analysis.

Gene accession numbers
NtHY5 (XM_009590053), NtMYB12 (XM_016624824), NtODC (NW_017671182), NtQPT
(NW_017672768), NtPMT (NW_017670241), NtTUB (XM_009622503), AtTUB (NM_125664).

Data availability
All data generated or analyzed during this study are included in this published article (and its
supplementary information files).

Conflict of interest
The authors declare that they have no conflict of interest.
751

 Figure Legends

Figure 1. Light-mediated modulation of nicotine and flavonol content in WT tobacco. A,
Diagrammatic representation of nicotine pathway of tobacco. B, Phenotype of 10-d-dark and light
grown WT seedlings. C, Expression analysis of nicotine pathway genes (*ODC, PMT, MPO, QPT*
and A622) in WT seedlings grown in light (root were covered to inhibit light illumination) and
dark for 10-days through qRT-PCR. D, Quantitative estimation of nicotine in 10-day-dark and
light grown seedlings (using only hypocotyl region) through HPLC. E, Time dependent qRT-PCR
expression analysis of nicotine pathway genes (ODC, MPO, PMT, QPT and A622) in 10-day dark
grown WT tobacco seedlings subsequently transferred to light for 0.5, 2, 4 and 24 hours. F,
Relative expression of NtHY5 in 10-d-dark and light grown seedlings. Statistical analysis was
performed using two-tailed Student's t-test. Error bars represent SE of means (n=3) For hypocotyl
and root length (n=15). Tubulin was used as endogenous control to normalize the relative
expression levels. Error bars represent standard deviation. Asterisks indicate a significant
difference, *P < 0.1, **P < 0.01, ***P < 0.001.

Figure 2. In vitro interaction between NtHY5 and the ACGT motif of G-BOX present at the
promoters of *NtPMT, NtODC and NtQPT* genes. A, Representation of the probes (G-BOXI, G-
mut I, G-BOXII and G-mut II) designed upstream of the transcription start site of NtPMT, NtODC
and NtQPT. G-BOX is altered by multiple base substitutions. B, EMSA (Electrophoretic mobility
shift assay) for the binding of 6X-His-HY5 (NtHY5) with core ACE-DIG element (HY5 binding
site that is G-BOXI and G-BOXII) present in the LRE motifs of the NtPMT, NtODC and NtQPT
respectively. Upper and lower arrows indicate shift and free probe respectively. C-F, EMSA with
competition between digoxigen in labelled G-BOXI, unlabeled G-BOX I (cold), G-BOXII and
unlabeled G-BOX II (cold) probe to bind to 6X-His-HY5 respectively. Superscript values 10X,
20X, 30X and 50X represent increasing amount of cold probe NtPMT-P1, NtPMT-P2, NtQPT and
NtODC respectively. Upper and lower arrows indicate shift and free probe respectively.

Figure 3. Complementation of *Arabidopsis hy5* mutant by NtHY5. A, Phenotype of 7-d-old
light-grown WT (wild type; Col-0), *hy5* and NtHY5OX; *hy5* seedlings (L1 and L2 are two
independent transgenic lines overexpressing NtHY5 in *hy5* mutant background). B, Hypocotyl
length of 7-d-old white light grown WT, *hy5*, L1 and L2 seedlings. C, Root length of 7-d-old white
light grown WT, hy5, L1 and L2. D, Rosette diameter of 30-day-old WT, *hy5*, L1 and L2 plants.
E, Representative photographs of the seed size of WT, *hy5* and transgenic lines (L1 and L2). Bars
= 1 mm. F, Measurement of seed size of 30-day-old WT, *hy5*, L1 and L2 plants. G, Relative
expression of pheylpropanoid pathway genes (*AtCHS, AtCHI, AtFLS and AtDFR*) in 7-d-light
grown WT, *hy5*, L1 and L2 seedlings. H, Quantitative estimation of flavonol (Kaempferol and
Quercitin) content in 7-d-old light grown WT, *hy5*, L1 and L2 seedlings through HPLC. I,
Expression analysis of *AtCHS, AtCHI, AtFLS* and *AtDFR* in WT, *hy5*, L1 and L2 seedlings grown
for 5 days in dark followed by an exposure to light for 3hr. J, Length of vascular bundle of 30-day-
oldstem of WT, *hy5*, L1 and L2 plants. Statistical analysis was performed using two-tailed
Student's t-test. Error bars represent SE of means (n=3). For hypocotyl and root length (n=10-12).

Tubulin was used as endogenous control to normalize the relative expression levels. Error bars
represent standard deviation. Asterisks indicate a significant difference, *P < 0.1, **P < 0.01, ***P
< 0.001.

Figure 4. Overexpression of NtHY5 and CRISPR-Cas9-mediated knockout of NtHY5
modulates gene expression and accumulation of metabolite in tobacco seedlings. A, Visual
representation of the 20-day-old seedlings with root covered with black paper after germination
for 5-days to inhibit light exposure. Roots were covered with black paper for 14 days. B, Relative
expression of NtHY5 in 20-day-old light grown WT, NtHY5OX and *NtHY5^{CR}* lines of tobacco. B,
C, Relative transcript abundance of nicotine pathway genes (*NtPMT, NtODC, NtQPT, NtMPO,*
NtBBL and NtA622) in 20-day-old light grown WT, NtHY5OX and *NtHY5^{CR}* lines. D,
Quantification of nicotine content in 20-day light-grown WT, NtHY5OX and *NtHY5^{CR}* leaves
through HPLC. E and F, Relative transcript abundance of nicotine pathway genes (*NtODC,*
NtPMT, NtMPO, NtQPT, NtBBL and NtA622) in 10-day-old dark-grown seedlings and seedlings
transferred light for 3h in NtHY5OX and *NtHY5^{CR}* seedlings through qRT-PCR. G, Quantification
of nicotine content in 10-day-dark grown seedlings and seedlings transferred light for 3h in
NtHY5OX and *NtHY5^{CR}* seedlings through HPLC. Roots and hypocotyls were used for all the
expression analysis and nicotine quantification, respectively. Tubulin was used as the endogenous
control to normalize the relative expression levels. The statistical analysis was performed using
two-tailed Student's t-tests. The data are plotted as means ± SD (n=3). The error bars represent
standard deviations. The asterisks indicate significant difference, *P< 0.1; **P< 0.01;
***P< 0.001.

Figure 5. NtHY5 moves from shoot to root to modulate nicotine levels in tobacco. A, Relative
expression of nicotine pathway genes (*NtODC, NtPMT, NtQPT, NtMPO, NtBBL and NtA622*) in
roots of grafted NtHY5OX and *NtHY5^{CR}* seedlings in several combinations keeping
HY5OX/HY5OX as control. B, Quantitative estimation of nicotine content in grafted seedlings of
WT, NtHY5OX and *NtHY5^{CR}* lines. Epicotyl tissue (300 mg fresh weight) was used for the
analysis. C, Relative expression of NtHY5 in roots of grafted WT, NtHY5OX and *NtHY5^{CR}*
seedlings in several combinations keeping *WT/WT* as control. D, Relative expression of nicotine
pathway genes (*NtODC, NtPMT, NtQPT, NtMPO, NtBBL and NtA622*) in roots of grafted WT,
NtHY5OX and *NtHY5^{CR}* seedlings in several combinations keeping *WT/WT* as control. E,
Quantitative estimation nicotine content in grafted seedlings of WT, NtHY5OX and *NtHY5^{CR}* lines
keeping *WT/WT* as control using HPLC. Roots and hypocotyls were used for all the expression
analysis and nicotine quantification, respectively. TUBULIN was used as the endogenous control
to normalize the relative expression levels. The statistical analysis was performed using two-tailed
Student's t-tests. The data are plotted as means ± SD (n= 3). The error bars represent standard
deviations. The asterisks indicate significant difference, *P< 0.1; **P< 0.01; ***P< 0.001.

Figure 6. NtHY5 modulates nicotine levels in mature tobacco transgenic lines. A and B
Relative transcript abundance of nicotine pathway genes (*NtODC, NtPMT, NtMPO, NtQPT and*
NtA622) in roots of NtHY5OX and *NtHY5^{CR}* 3-month-old tobacco plants. C, Quantitative

estimation nicotine content in 3-month-old leaves of plant using HPLC. D, Representative
photograph of 5-months-old mature tobacco plant being kept in dark for 24hours. E, Relative
expression analysis of nicotine pathway genes (*NtODC, NtMPO, NtPMT, NtA622, NtQPT and*
NtBBL) in roots of 5-months-old mature NtHY5OX and *NtHY5^{CR}* plants grown in light and
transferred to dark for 24h through qRT-PCR. F, Quantification of nicotine content in in 5-months-
old mature tobacco leaves transferred to dark for 24h in NtHY5OX and *NtHY5^{CR}* lines through
HPLC. Leaf tissue (300 mg dry weight) was used for the analysis. Roots and hypocotyls were
used for all the expression analysis and nicotine quantification, respectively. Tubulin was used as
the endogenous control to normalize the relative expression levels. The statistical analysis was
performed using two tailed Student's t-tests. The data are plotted as means $\pm$ SD (n= 3). For
measurement of root bending angle, root length, hypocotyls length and seed size (n=15). The error
bar represents standard deviations. The asterisks indicate significant difference, *P< 0.1;
**P< 0.01; ***P< 0.001.

Figure 7. Proposed working model showing the role of NtHY5 in nicotine and flavonol
accumulation in tobacco. We propose that NtHY5 moves from shoot to root to upregulate the
expression of nicotine pathway genes in light which results into increased accumulation of nicotine
as well as flavonoid biosynthesis. NtHY5 regulates nicotine and flavonol biosynthesis via binding
to the G-boxes in the promoters of nicotine and flavonol regulatory genes. Higher accumulation
of secondary metabolites leads to less ROS accumulation thus increased salt stress tolerance
ability. Higher nicotine and flavonoids also regulates growth and development of tobacco.
Arrowheads and tees indicate positive and negative regulation, respectively.

References

Agati G, Azzarello E, Pollastri S, Tattini M (2012) Flavonoids as antioxidants in plants: location
and functional significance. Plant Sci **196**: 67–76.

Badola PK, Sharma A, Gautam H, Trivedi PK (2022) MicroRNA858a, its encoded peptide,
and phytosulfokine regulate *Arabidopsis* growth and development. Plant Physiol **189**, 1397-
1415.

Banerjee AK, Chatterjee M, Yu Y, Suh S, Miller WA, Hannapel DJ (2006) Dynamics of a
mobile RNA of potato involved in a long-distance signaling pathway. Plant Cell **18**, 3443–
3457.

Bhatnagar A, Singh S, Khurana JP (2020) HY5-COP1: the central module of light signaling
pathway. J Plant Biochem Biotechnol **29**: 590–610.

Bhatia C, Pandey A, Gaddam SR, Hoecker U, Trivedi PK (2018) Low temperature-enhanced
flavonol synthesis requires light-associated regulatory components in *Arabidopsis thaliana*.
Plant Cell Physiol **59**: 2099–2112.

Bian S, Sui X, Wang J, Tian T, Wang C, Zhao X, Liu X, Fang N, Zhang Y, Liu Y, Du Y,
Wang B, Timko MP, Zhang Z, Zhang H (2022) NtMYB305a binds to the jasmonate-
responsive GAG region of *NtPMT1a* promoter to regulate nicotine biosynthesis. Plant Physiol
188: 151–166.

Blilou I, Xu J, Wildwater M, Willemsen V, Paponov I, Friml J, Heidstra R, Aida M, Palme
K, Scheres B (2005) The PIN auxin efflux facilitator network controls growth and patterning
in *Arabidopsis* roots. Nature **433**: 39–44

Bozorov TA, Dinh ST, Baldwin IT (2017) JA but not JA-Ile is the cell-nonautonomous signal
activating JA mediated systemic defenses to herbivory in *Nicotiana attenuata*. J Int Plant Biol
59: 552–57.

Burman N, Bhatnagar A, Khurana JP (2018) OsbZIP48, a HY5 transcription factor ortholog,
exerts pleiotropic effects in light-regulated development. Plant Physiol **176**: 1262-1285.

Chen X, Yao Q, Gao X, Jiang C, Harberd NP, Fu X (2016) Shoot-to-root mobile transcription
factor hy5 coordinates plant carbon and nitrogen acquisition. *Curr Biol.* **26**: 640–646.

Chattopadhyay S, Ang LH, Puente P, Deng, XW, Wei N (1998) *Arabidopsis* bZIP protein HY5
directly interacts with light-responsive promoters in mediating light control of gene expression.
Plant Cell **10**: 673–683.

X. Chen, Q. Yao, X. Gao, C. Jiang, N.P. Harberd, X. Fu (2016) Shoot-to-root mobile

transcription factor HY5 coordinates plant carbon and nitrogen acquisition, Curr Biol **26:**

640e646.

Chen S, Wu F, Li Y, Qian Y, Pan X, Li F, Wang Y, Wu Z, Fu C, Lin H, Yang A (2019)

NtMYB4 and NtCHS1 are critical factors in the regulation of flavonoid biosynthesis and are

involved in salinity responsiveness. Front Plant Sci **10**: 178.

Chory J (2010) Light signal transduction: an infinite spectrum of possibilities. Plant J **61**: 982-

991.

Cluis CP, Mouchel CF, Hardtke CS (2004) The *Arabidopsis* transcription factor HY5 integrates

light and hormone signaling pathways. Plant J **38**: 332-347.

De Boer KD, Lye JC, Aitken CD, Su AK, Hamill JD (2009) The A622 gene in Nicotiana glauca

(tree tobacco): evidence for a functional role in pyridine alkaloid synthesis. Plant Mol Biol **69**:

299–312.

De Boer K, Lye JC, Aitken CD, Su AK, Hamill JD (2009) The A622 gene in Nicotiana glauca

(tree tobacco): evidence for a functional role in pyridine alkaloid synthesis. Plant Mol Biol **69**:

299–312.

De Boer K, Tilleman S, Pauwels L, Vanden Bossche **R, Sutter De V, Vanderhaeghen R,**

Hilson P, Hamill JD, Goossens A (2011) APETALA2/ETHYLENE RESPONSE FACTOR

and Basic Helix-Loop-Helix Tobacco Transcription Factors Cooperatively Mediate Jasmonate-

Elicited Nicotine Biosynthesis. Plant J. **66**: 1053–1065.

Dewey RE, Xie J (2013) Molecular genetics of alkaloid biosynthesis in Nicotiana tabacum.

Phytochemistry **94**: 10-27.

Doughty J, Aljabri M, Scott RJ (2014) Flavonoids and the regulation of seed size in *Arabidopsis*.

Biochem Soc Trans **42**: 364–369.

Falcone Ferreyra, ML, Rius, SP, Casati P (2012) Flavonoids: biosynthesis, biological functions,

and biotechnological applications. Front Plant Sci **3**: 222.

Friml J, Vieten A, Sauer M, Weijers D, Schwarz H, Hamann T, Offringa R, Jürgens G (2003)

Efflux-dependent auxin gradients establish the apical-basal axis of *Arabidopsis*. Nature **426**:

147–153

Fu B, Ji X, Zhao M, He F, Wang X, Wang Y, Liu P, Niu, L (2016) The influence of light quality
on the accumulation of flavonoids in tobacco (*Nicotiana tabacum* L.) leaves. J Photochem
Photobiol **162**: 544-549.

Gangappa SN, Botto JF (2016) The multifaceted roles of HY5 in plant growth and development.
Mol Plant **9**: 1353–1365.

Geisler M, Blakeslee JJ, Bouchard R, Lee OR, Vincenzetti V, Bandyopadhyay A,
Titapiwatanakun B, Peer WA, Bailly A, Richards EL, et al (2005) Cellular efflux of auxin
catalyzed by the *Arabidopsis* MDR/PGP transporter AtPGP1. Plant J **44**: 179–194

Hashimoto T, Yamada Y (1994) Alkaloid biogenesis: Molecular aspects. Annu Rev Plant Physiol
Plant Mol Biol **45**: 257–28

Hayashi S, Watanabe M, Kobayashi M, Tohge T, Hashimoto T, Shoji T (2020) Genetic
Manipulation of Transcriptional Regulators Alters Nicotine Biosynthesis in Tobacco, *Plant*
Cell Physiol **61**: 1041–1053.

Haywood V, Yu T, Huang N, Lucas WJ (2005) Phloem long-distance trafficking of gibberellic
acid-insensitive RNA regulates leaf development. Plant J. **42**, 49–68.

Hibi N, Higashiguchi S, Hashimoto T, Yamada Y (1994) Gene expression in tobacco low-
nicotine mutants. Plant Cell **6**: 723–735.

Hildreth SB, Gehman EA, Yang H, Lu RH, Ritesh KC, Harich KC, Yu S, Lin J, Sandoe JL,
Okumoto S, Murphy AS, Jelesko JG (2011) Tobacco nicotine uptake permease (NUP1)
affects alkaloid metabolism. Proc Natl Acad Sci USA **108**: 18179–18184.

Holm M, Ma LG, Qu LJ, Deng XW (2002) Two interacting bZIP proteins are direct targets of
COP1-mediated control of light-dependent gene expression in *Arabidopsis*. Genes Dev **16**(10):
1247–1259.

Horsch RB (1985) A simple method for transferring genes into plants. Science **227**: 1229–123.

Imanishi S, Hashizume K, Nakakita M, Kojima H, Matsubayashi Y, Hashimoto T (1998)
Differential induction by methyl jasmonate of genes encoding ornithine decarboxylase and
other enzymes involved in nicotine biosynthesis in tobacco cell cultures. Plant Mol Biol **38**:
1101–1111.

Jiang W, Yin Q, Wu R, Zheng G, Liu J, Dixon RA (2015) Role of a chalcone isomerase-like
protein in flavonoid biosynthesis in *Arabidopsis* thaliana. J Exp Bot **66**: 7165–7179.

Jing Y, Zhang D, Wang X, Tang W, Wang W, Huai, J, Xu G, Chen D, Li, Y, Lin R (2013)
Arabidopsis chromatin remodeling factor PICKLE interacts with transcription factor HY5 to
regulate hypocotyl cell elongation. Plant Cell **25**: 242–256.

Kajikawa M, Shoji T, Kato A, Hashimoto T (2011) Vacuole-localized berberine bridge enzyme-
like proteins are required for a late step of nicotine biosynthesis in tobacco. Plant Physiol **155**:
2010–2022.

Katoh A, Uenohara K, Akita M, Hashimoto T (2006) Early steps in the biosynthesis of NAD
in *Arabidopsis* start with aspartate and occur in the plastid. Plant Physiol **141**: 851–857.

Khan S, Pandey SS, Jyotshna, Shanker K, Khan F, Rahman LU (2017) Cloning and functional
characterization of quinolinic acid phosphoribosyl transferase (QPT) gene of Nicotiana
tabacum. Physiol Plant **160**: 253-265.

Koornneef, M, Rolff, E, Spruit CJP (1980) Genetic control of light inhibited hypocotyl
elongation in *Arabidopsis thaliana* (L.) *Heynh. Zeitschrift für Pflanzenphysiologie* **100**: 147–
160.

Kumar S, Nei M, Dudley J, Tamura K (2008) MEGA: a biologist-centric software for
evolutionary analysis of DNA and protein sequences. Brief Bioinform **9**: 299–306.

Lee J, He K, Stolc V, Lee H, Figueroa P, Gao Y, Zhao H, Lee I, Denga XW (2007) Analysis
of transcription factor HY5 genomic binding sites revealed its hierarchical role in light
regulation of development. Plant Cell **19**: 731–749.

Lescot, M, Dehais G, Thisj K, Marchal Y, Peer Y, Rouze S, Rombauts, S (2020)
RombautsPlantCARE, a database of plant cis-acting regulatory elements and a portal to tools
for in silico analysis of promoter sequences. Nucleic Acids Res **30**: 325-327.

Li C, Teng W, Shi Q, Zhang F (2007). Multiple signals regulate nicotine synthesis in tobacco
plant. Plant Signal Behav **2:**280-1.

Li J, Yang L, Jin D, Nezames CD, Terzaghi W, Deng XW (2013) UV-B-induced
photomorphogenesis in *Arabidopsis*. Protein Cell **4**: 485–492.

Li R, Llorca LC, Schuman MC, Wang Y, Wang L, Joo Y, Wang M, Vass~ao DG, Baldwin
IT (2018) ZEITLUPE in the roots of wild tobacco regulates jasmonate-mediated nicotine
biosynthesis and resistance to a generalist herbivore. Plant Physiol **177**: 833–8.

Liu H, Kotova TI, Timko MP (2019) Increased leaf nicotine content by targeting transcription
factor gene expression in commercial flue-cured tobacco (*Nicotiana tabacum* L.). Genes
(Basel) **10**: 930

Liu X, Qin T, Ma Q (2013) Light-regulated hypocotyl elongation involves proteasome-dependent
degradation of the microtubule regulatory protein WDL3 in *Arabidopsis*. Plant Cell **25**: 1740-
1755.

Liu X, Singh SK, Patra B, Liu Y, Wang B, Wang J, Pattanaik S, Yuan L (2020) Protein-
phosphatase NtPP2C2b and MAP-kinase NtMPK4 act in concert to modulate nicotine
biosynthesis. J Exp Bot **72**: 1661–1676.

Liu ZY, Wang PL, Zhang TQ, Li YB, Wang YY, Gao CQ (2018) Comprehensive analysis of
BpHSP genes and their expression under heat stresses in *Betula platyphylla*. Environ Exp Bot
152: 167–176.

Liva KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
quantitative PCR and the 2∆∆CT I method. Methods **25**: 402–408.

Maclachlan S, Zalik S (1963) Plastid structure, chlorophyll concentration, and free amino acid
composition of a chlorophyll mutant of barley. Can J Bot **41**(7), 1053-1062.

McCarthy RL, Zhong, R, Ye ZH (2009) MYB83 is a direct target of SND1 and acts redundantly
with MYB46 in the regulation of secondary cell wall biosynthesis in *Arabidopsis*. Plant Cell
Physiol **50**: 1950–1964.

Misra P, Pandey A, Tiwari M, Chandrashekar K, Sidhu OP, Asif MH, Chakrabarty D, Singh
PK, Trivedi PK (2010) Modulation of transcriptome and metabolome of tobacco by
Arabidopsis transcription factor, AtMYB12, leads to insect resistance. Plant Physiol **152**: 2258–
2268.

Michael R, Ranjan A, Kumar RS, Parthak PK, Trivedi PK (2020) Light-regulated expression
of terpene synthase gene, AtTPS03, is controlled by the Bzip transcription factor, HY5, in
Arabidopsis thaliana. Biochem Biophys Res Commun **529**: 437–443.

Naconsie M, Kato K, Shoji T, Hashimoto T (2014) Molecular evolution of N-methylputrescine
oxidase in tobacco. Plant Cell Physiol **55**: 436–444.

Nawkar GM, Kang CH, Maibam P, Park JH, Jung YJ, Chae HB, Chi YH, Jung IJ, Kim
WY, Yun DJ (2017) HY5, a positive regulator of light signaling, negatively controls the
unfolded protein response in *Arabidopsis*. Proc Natl Acad Sci USA **114**: 2084–2089.

Nguyen NH, Jeong CY, Kang GH, Yoo SD, Hong SW, Lee H (2015) MYBD employed by HY5

increases anthocyanin accumulation via repression of MYBL2 in *Arabidopsis*. Plant J **84**: 1192-

1205.

Niranjan A (2011) Development and optimization of HPLC-PDA-MS-MS method for

simultaneous quantifcation of three classes of favonoids in legume seeds, vegetables, fruits and

medicinal plants. J Liq Chromatogr Relat Technol **34**: 1729–1742.

Nishimura R, Ohmori M, Fujita H, Kawaguchi M (2002) A Lotus basic leucine zipper protein

with a RING-finger motif negatively regulates the developmental program of nodulation. Proc

Natl Acad Sci USA **99**: 15206–15210.

Noh B, Murphy AS, Spalding EP (2001) Multidrug resistance-like genes of *Arabidopsis* required

for auxin transport and auxin-mediated development. Plant Cell **13**: 2441–2454

Oki H., Hashimoto T (2004) Jasmonate-responsive regions in a *Nicotiana sylvestris PMT* gene

involved in nicotine biosynthesis. Plant Biotechnol **21**: 269–274

Osterlund MT, Hardtke CS, Wei N, Deng XW (2000) Targeted destabilization of HY5 during

light-regulated development of *Arabidopsis*. Nature **405**: 462–466.

Oyama T, Shimura Y, Okada K (1997) The *Arabidopsis* HY5 gene encodes a bZIP protein that

regulates stimulus-induced development of root and hypocotyl. Genes Dev **11**, 2983–2995.

Palaniswamy SK, James S, Sun H, Lamb RS, Davuluri RV, Grotewold E (2006) AGRIS and

AtRegNet: a platform to link cis-regulatory elements and transcription factors into regulatory

networks. Plant Physiol **140**: 818-829.

Pandey A, Misra P, Trivedi PK (2015) Constitutive expression of *Arabidopsis* MYB

transcription factor, *AtMYB11*, in tobacco modulates flavonoid biosynthesis in favor of flavonol

accumulation. Plant Cell Rep **34**: 1515–1528.

Pandey A, Misra, P, Bhambhani S, Bhatia C, Trivedi, PK (2014) Expression of *Arabidopsis*

MYB transcription factor, AtMYB111, in tobacco requires light to modulate flavonol content.

Sci Rep **4**: 5018.

Pereira GR, Marchetti JM, Bentley MVLB (2001) A simple and rapid method for nicotine assay

by HPLC from cutaneous microdialysis samples. Anal Lett **34**: 1669-1676.

Peer WA , Bandyopadhyay A , Blakeslee JJ (2004). Variation in expression and protein

localization of the PIN family of auxin efflux facilitator proteins in flavonoid mutants with

altered auxin transport in *Arabidopsis thaliana*. Plant Cell, **16**: 1898-1911.

Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R (2005) InterProScan: protein domains identifier. Nucleic Acids Res **33**(Web Server issue): W116–W120.

Riechers DE, Timko MP (1999) Structure and expression of the gene family encoding putrescine N-methyltransferase in *Nicotiana tabacum*: New clues to the evolutionary origin of cultivated tobacco. Plant Mol Biol **41**: 387–401.

Rus A, Baxter I, Muthukumar B, Gustin J, Lahner B, Yakubova E, Salt DE (2006) Natural variants of AtHKT1 enhance Na+ accumulation in two wild populations of *Arabidopsis*. PLoS Genet **2**: e210.

Santelia D , Henrichs S, Vincenzetti V, Sauer M, Bigler L (2008) Flavonoids redirect PIN-mediated polar auxin fluxes during root gravitropic responses. J. Biol. Chem. **283**: 31218-31226.

Schulz E, Tohge T, Zuther E, Fernie AR, Hincha DK (2016) Flavonoids are determinants of freezing tolerance and cold acclimation in *Arabidopsis* thaliana. Sci Rep **6**: 34027.

Sears MT, Zhang HB, Rushton PJ, Wu M, Han S, Spano AJ, Timko MP (2014) NtERF32: a non-NIC2 locus AP2/ERF transcription factor required in jasmonate-inducible nicotine biosynthesis in tobacco. Plant Mol Biol **84**: 49–66.

Sharma A, Badola, PK, Bhatia C, Sharma D, Trivedi PK (2020) Primary transcript of miR858 encodes regulatory peptide and controls flavonoid biosynthesis and development in *Arabidopsis*. Nat Plants **6**: 1262–1274.

Sharma A, Badola PK, Gautam H, Gaddam SR, Trivedi PK (2022) HY5 regulates light-dependent expression and accumulation of miR858a-encoded peptide, miPEP858a. Biochem Biophys Res Commun **589**: 204–208.

Shin DH, Choi M, Kim K, Bang G, Cho M, Choi SB, Choi G, Park YI (2013) HY5 regulates anthocyanin biosynthesis by inducing the transcriptional activation of the MYB75/PAP transcription factor in *Arabidopsis*. FEBS Lett **587**: 1543-1547.

Shoji T, Yamada Y, Hashimoto T (2000) Jasmonate induction of putrescine N-methyltransferase genes in the root of *Nicotiana sylvestris*. Plant Cell Physiol. **41**: 831–839

Shoji T, Hashimoto T (2011) Tobacco MYC2 regulates jasmonate-inducible nicotine biosynthesis genes directly and by way of the NIC2-locus ERF genes. Plant Cell Physiol **52**: 1117–1130.

Shoji T, Inai K, Yazaki K, Sato Y, Takase H, Shitan N (2009) Multidrug and toxic compound extrusion-type transporters implicated in vacuolar sequestration of nicotine in tobacco roots. Plant Physiol **149**: 708–718.

Shoji T, Kajikawa M, Hashimoto T (2010) Clustered transcription factor genes regulate nicotine biosynthesis in tobacco. Plant Cell **22**: 3390–3409.

Shoji T, Ogawa T, Hashimoto T (2008) Jasmonate-induced nicotine formation in tobacco is mediated by tobacco COI1 and JAZ genes. Plant Cell Physiol **49**: 1003–1012.

Shoji T, Mishima M, Hashimoto, T (2013) Divergent DNA-binding specificities of a group of ETHYLENE RESPONSE FACTOR transcription factors involved in plant defense. Plant Physiol **162**: 977–990.

Shoji, T Yuan, L (2021) ERF gene clusters: working together to regulate metabolism. Trends Plant Sci **26**: 23–32.

Shoji T, Hashimoto T (2011) Recruitment of a duplicated primary metabolism gene into the nicotine biosynthesis regulon in tobacco. Plant J **67:** 949–959.

Shoji T, Hashimoto T (2012) DNA-Binding and Transcriptional Activation Properties of Tobacco NIC2-Locus ERF189 and Related Transcription Factors. Plant Biotechnol. **29**: 35–42.

Sibout R, Su P, Hettiarachchi C, Holm M, Muday GK, Hardtke CS (2006) Opposite root growth phenotypes of *hy5* versus *hy5 hyh* mutants correlate with increased constitutive auxin signaling. PLOS Genet **2**: e202.

Sinclaie SJ, Murphy KJ, Birch CD, Hamill JD (2000) Molecular characterization of quinolinate phosphoribosyltransferase (QPRTase) in Nicotiana. Plant Mol Biol **44**: 603–617.

Song S, Qi T, Huang H, Ren Q, Wu D, Chang C, Peng W, Liu Y, Peng J, Xie D (2011) The jasmonate-ZIM domain proteins interact with the R2R3–MYB transcription factors MYB21 and MYB24 to affect jasmonate-regulated stamen development in *Arabidopsis*. Plant Cell **23**: 1000–1013.

Song YH, Yoo CM, Hong AP, Kim SH, Jeong HJ, Shin SY, Kim HJ, Yun DJ, Lim CO, Bahk JD (2008) DNA-binding study identifies C-box and hybrid C/G-box or C/A-box motifs as high-affinity binding sites for STF1 and LONG HYPOCOTYL5 proteins. Plant Physiol **146**: 1862-1877.

Song Z, Luo Y, Wang W, Fan N, Wang D, Yang C, Jia H (2020) NtMYB12 positively regulates
flavonol biosynthesis and enhances tolerance to low Pi stress in *Nicotiana tabacum*. Front Plant
Sci **10**: 1683.

Song ZP, Fan NB, Jiao GZ, Liu MH, Wang XY, Jia HF (2019) Overexpression of OsPT8
increases auxin content and enhances tolerance to high-temperature stress in *Nicotiana*
tabacum. Genes **10**: 809.

Steppuhn A, Gase K, Krock B, Halitschke R, Baldwin IT (2004) Nicotine's defensive function
in nature. PLoS Biol **2**: E217.

Stracke R, Favory JJ, Gruber H, Bartelniewoehner L, Bartels S, Binkert M, Funk M,
Weisshaar B, Ulm R (2010) The *Arabidopsis* bZIP transcription factor HY5 regulates
expression of the PFG1/MYB12 gene in response to light and ultraviolet-Bradiation. Plant Cell
Environ **33**: 88–103.

Sun H, Sun X, Wang H (2020) Advances in salt tolerance molecular mechanism in tobacco
plants. Hereditas **157**, 5.

Todd AT, Liu E, Polvi SL, Pammett RT, Page JE (2010) A functional genomics screen
identifies diverse transcription factors that regulate alkaloid biosynthesis in Nicotiana
benthamiana. Plant J **62**: 589–600

Truong HA, Lee S, Trịnh CS, Lee WJ, Chung EH, Hong SW, Lee H (2021) Overexpression
of the *HDA15* Gene Confers Resistance to Salt Stress by the Induction of NCED3, an ABA
Biosynthesis Enzyme. Front Plant Sci **12**: 640443.

Ulm R, Baumann A, Oravecz A, Máté Z, Adám E, Oakeley EJ, Schäfer E, Nagy F (2004)
Genome-wide analysis of gene expression reveals function of the bZIP transcription factor HY5
in the UV-B response of *Arabidopsis*. Proc Natl Acad Sci USA **101**: 1397–1402.

Vanhaelewyn L, Schumacher P, Poelman D, Fankhauser C, Van Der Straeten D,
VandenbusscheF (2016) REPRESSOR OF ULTRAVIOLET-B PHOTOMORPHOGENESIS
function allows efficient phototropin mediated ultraviolet-B phototropism in etiolated
seedlings. Plant Sci **252**: 215-221.

Vogt T (2010) Phenylpropanoid biosynthesis. Mol Plant **3**: 2–20.

Wan J, Zhang P, Wang R, Sun L, Wang W, Zhou H, Xu J (2018) UV-B Radiation Induces
Root Bending Through the Flavonoid-Mediated Auxin Pathway in *Arabidopsis*. Front Plant Sci
9: 618.

WHO (2015) Advisory note: global nicotine reduction strategy. WHO study group on tobacco
product regulation**. Geneva: WHO Press.**

Winkel-shirley B (2002) Biosynthesis of flavonoids and effects of stress. Curr Opin Plant Biol
5: 218-223.

Wong SP, Leong LP, Koh JHW (2006) Antioxidant activities of aqueous extracts of selected
plants. Food Chem **99**: 775–783.

Xing HL, Dong L, Wang ZP, Zhang HY, Han CY, Liu B, Wang XC, Chen QJ (2014)
CRISPR/Cas9 toolkit for multiplex genome editing in plants. BMC Plant Biol **14**: 327.

Xu B, Timko M (2004) Methyl jasmonate induced expression of the tobacco putrescine N-
methyltransferase genes requires both G-box and GCC-motif elements. Plant Mol Biol **55**: 743–
761.

Xu S, Brockmöller T, Navarro-Quezada A, Kuhl H, Gase K, Ling Z, Zhou W, Kreitzer C,
Stanke M, Tang H, Lyons E, Pandey P, Pandey SP, Timmermann B, Gaquerel E, Baldwin
IT (2017) Wild tobacco genomes reveal the evolution of nicotine biosynthesis. Proc Natl Acad
Sci USA **114**: 6133–6138.

Yamawaki S, Yamashino T, Nakanishi H, Mizuno T (2011) Functional characterization of HY5
homolog genes involved in early light-signaling in *Physcomitrella patens*. Biosci Biotechnol
Biochem **75**: 1533–15.

Yang Y, Mao L, Jittayasothorn Y, Kang Y, Jiao C, Fei Z, Zhong G (2015) Messenger RNA
exchange between scions and rootstocks in grafted grapevines. BMC Plant Biol. **15**, 251.

Yin P, Lan B, Liu X, Cui Y, Wang Y, Xu J (2022) Mobile messenger RNAs in grafts of *Salix*
matsudana are associated with plant rooting. Forests. **13**:354.

Zhang H, He H, Wang X, Wang X, Yang X, Li L, Deng XW (2011) Genome-wide mapping of
the HY5-mediated gene networks in *Arabidopsis* that involve both transcriptional and post-
transcriptional regulation. Plant J **65**: 346–358.

Zhang HB, Bokowiec MT, Rushton PJ, Han SC, Timko MP (2012) Tobacco Transcription
factors NtMYC2a and NtMYC2b form nuclear complexes with the NtJAZ1 repressor and
regulate multiple jasmonate-inducible steps in nicotine biosynthesis. Mol Plant **5**: 73–84.

Zhang Y, Li C, Zhang J, Wang J, Yang J, Lv Y (2017) Dissection of HY5/HYH expression in
Arabidopsis reveals a root autonomous HY5-mediated photomorphogenic pathway. PLoS ONE
12: e0180449.

Zhao Q, Ren YR, Wang QJ, Wang XF, You CX, Hao, YJ (2016) Ubiquitination-related MdBT

scaffold proteins target a bHLH transcription factor for iron homeostasis. Plant Physiol **172**:

1973–1988.

Zhong R, Lee C, Zhou J, McCarthy RL, Ye ZH (2008) A battery of transcription factors

involved in the regulation of secondary cell wall biosynthesis in *Arabidopsis*. Plant Cell **20**,

2763–2782.

Zhou Z, Schenke D, Miao Y, Cai D (2017) Investigation of the crosstalk between the flg22 and

the UV-B-induced flavonol pathway in *Arabidopsis thaliana* seedlings. Plant Cell Environ **40**:

453–458.

1167

1168

Figure 1. Light-mediated modulation of nicotine and flavonol content in WT tobacco. A, Diagrammatic representation of nicotine pathway of tobacco. B, Phenotype of 10-d-dark and light grown WT seedlings. C, Expression analysis of nicotine pathway genes (*ODC, PMT, MPO, QPT and A622*) in WT seedlings grown in light (root were covered to inhibit light illumination) and dark for 10-days through qRT-PCR. D, Quantitative

estimation of nicotine in 10-day-dark and light grown seedlings (using only hypocotyl region) through HPLC. E, Time dependent qRT-PCR expression analysis of nicotine pathway genes (ODC, MPO, PMT, QPT and A622) in 10-day dark grown WT tobacco seedlings subsequently transferred to light for 0.5, 2, 4 and 24 hours. F, Relative expression of NtHY5 in 10-d-dark and light grown seedlings. Statistical analysis was performed using two-tailed Student's t-test. Error bars represent SE of means (n=3) For hypocotyl and root length (n=15). Tubulin was used as endogenous control to normalize the relative expression levels. Error bars represent standard deviation. Asterisks indicate a significant difference, *$P < 0.1$, **$P < 0.01$, ***$P < 0.001$.

Figure 2. In vitro interaction between NtHY5 and the ACGT motif of G-BOX present at the promoters of *NtPMT, NtODC and NtQPT* genes. A, Representation of the probes (G-BOXI, G-mut I, G-BOXII and G-mut II) designed upstream of the transcription start site of NtPMT, NtODC and NtQPT. G-BOX is altered by multiple base substitutions. B, EMSA (Electrophoretic mobility shift assay) for the binding of 6X-His-HY5 (NtHY5) with core ACE-DIG element (HY5 binding site that is G-BOXI and G-BOXII) present in the LRE motifs of the NtPMT, NtODC and NtQPT respectively. Upper and lower arrows indicate shift and free probe respectively. C-F, EMSA with competition between digoxigen in labelled G-BOXI, unlabeled G-BOX I (cold), G-BOXII and unlabeled G-BOX II (cold) probe to bind to 6X-His-HY5 respectively. Superscript values 10X, 20X, 30X and 50X represent increasing amount of cold probe NtPMT-P1, NtPMT-P2, NtQPT and NtODC respectively. Upper and lower arrows indicate shift and free probe, respectively.

Figure 3. Complementation of *Arabidopsis hy5* mutant by NtHY5. A, Phenotype of 7-d-old light-grown WT (wild type; Col-0), *hy5* and NtHY5OX; *hy5* seedlings (L1 and L2 are two independent transgenic lines overexpressing NtHY5 in *hy5* mutant background). B, Hypocotyl length of 7-d-old white light grown

WT, *hy5*, L1 and L2 seedlings. C, Root length of 7-d-old white light grown WT, hy5, L1 and L2. D, Rosette diameter of 30-day-old WT, *hy5*, L1 and L2 plants. E, Representative photographs of the seed size of WT, *hy5* and transgenic lines (L1 and L2). Bars = 1 mm. F, Measurement of seed size of 30-day-old WT, *hy5*, L1 and L2 plants. G, Relative expression of pheylpropanoid pathway genes (*AtCHS, AtCHI, AtFLS and AtDFR*) in 7-d-light grown WT, *hy5*, L1 and L2 seedlings. H, Quantitative estimation of flavonol (Kaempferol and Quercitin) content in 7-d-old light grown WT, *hy5*, L1 and L2 seedlings through HPLC. I, Expression analysis of *AtCHS, AtCHI, AtFLS* and *AtDFR* in WT, *hy5*, L1 and L2 seedlings grown for 5 days in dark followed by an exposure to light for 3hr. J, Length of vascular bundle of 30-day-oldstem of WT, *hy5*, L1 and L2 plants. Statistical analysis was performed using two-tailed Student's t-test. Error bars represent SE of means (n=3). For hypocotyl and root length (n=10-12). Tubulin was used as endogenous control to normalize the relative expression levels. Error bars represent standard deviation. Asterisks indicate a significant difference, $*P < 0.1$, $**P < 0.01$, $***P < 0.001$.

Figure 4. Overexpression of NtHY5 and CRISPR-Cas9-mediated knockout of NtHY5 modulates gene expression and accumulation of metabolite in tobacco seedlings. A, Visual representation of the 20-day-old seedlings with root covered with black paper after germination for 5-days to inhibit light exposure. Roots were covered with black paper for 14 days. B, Relative expression of NtHY5 in 20-day-old light grown WT,

NtHY5OX and *NtHY5^{CR}* lines of tobacco. B, C, Relative transcript abundance of nicotine pathway genes (*NtPMT, NtODC, NtQPT, NtMPO, NtBBL and NtA622*) in 20-day-old light grown WT, NtHY5OX and *NtHY5^{CR}* lines. D, Quantification of nicotine content in 20-day light-grown WT, NtHY5OX and *NtHY5^{CR}* leaves through HPLC. E and F, Relative transcript abundance of nicotine pathway genes (*NtODC, NtPMT, NtMPO, NtQPT, NtBBL* and NtA622) in 10-day-old dark-grown seedlings and seedlings transferred light for 3h in NtHY5OX and *NtHY5^{CR}* seedlings through qRT-PCR. G, Quantification of nicotine content in 10-day-dark grown seedlings and seedlings transferred light for 3h in NtHY5OX and *NtHY5^{CR}* seedlings through HPLC. Roots and hypocotyls were used for all the expression analysis and nicotine quantification, respectively. Tubulin was used as the endogenous control to normalize the relative expression levels. The statistical analysis was performed using two-tailed Student's t-tests. The data are plotted as means ± SD (n=3). The error bars represent standard deviations. The asterisks indicate significant difference, *P< 0.1; **P< 0.01; ***P< 0.001.

Figure 5. NtHY5 moves from shoot to root to modulate nicotine levels in tobacco. A, Relative expression of nicotine pathway genes (*NtODC, NtPMT, NtQPT, NtMPO, NtBBL and NtA622*) in roots of grafted NtHY5OX and *NtHY5*^CR seedlings in several combinations keeping *HY5OX/HY5OX* as control. B, Quantitative estimation of nicotine content in grafted seedlings of WT, NtHY5OX and *NtHY5*^CR lines. Epicotyl tissue (300 mg fresh weight) was used for the analysis. C, Relative expression of NtHY5 in roots of grafted WT, NtHY5OX and *NtHY5*^CR seedlings

in several combinations keeping *WT/WT* as control. D, Relative expression of nicotine pathway genes (*NtODC, NtPMT, NtQPT, NtMPO, NtBBL and NtA622*) in roots of grafted WT, NtHY5OX and *NtHY5^{CR}* seedlings in several combinations keeping *WT/WT* as control. E, Quantitative estimation nicotine content in grafted seedlings of WT, NtHY5OX and *NtHY5^{CR}* lines keeping *WT/WT* as control using HPLC. Roots and hypocotyls were used for all the expression analysis and nicotine quantification, respectively. TUBULIN was used as the endogenous control to normalize the relative expression levels. The statistical analysis was performed using two-tailed Student's t-tests. The data are plotted as means ± SD (n= 3). The error bars represent standard deviations. The asterisks indicate significant difference, *P< 0.1; **P< 0.01; ***P< 0.001.

Figure 6. NtHY5 modulates nicotine levels in mature tobacco transgenic lines. A and B Relative transcript abundance of nicotine pathway genes (*NtODC, NtPMT, NtMPO, NtQPT and NtA622*) in roots of NtHY5OX and *NtHY5^{CR}* 3-month-old tobacco plants. C, Quantitative estimation nicotine content in 3-month-old leaves of plant using HPLC. D, Representative photograph of 5-months-old mature tobacco plant being kept in dark for 24hours. E, Relative expression analysis of nicotine pathway genes (*NtODC, NtMPO, NtPMT, NtA622, NtQPT and NtBBL*) in roots of 5-months-old mature NtHY5OX and *NtHY5^{CR}* plants grown in light and transferred to dark for 24h through qRT-PCR. F, Quantification of nicotine content in in 5-months-old mature tobacco leaves transferred to dark for 24h in NtHY5OX and *NtHY5^{CR}* lines through HPLC. Leaf tissue (300 mg dry weight) was used for the analysis. Roots and hypocotyls were used for all the expression analysis and nicotine quantification, respectively. Tubulin was used as the endogenous control to normalize the relative expression levels. The statistical analysis was performed using two tailed Student's t-tests. The data are plotted as means ± SD (n= 3). For measurement of root bending angle, root length, hypocotyls length and seed size (n=15). The error bar represents standard deviations. The asterisks indicate significant difference, *P< 0.1; **P< 0.01; ***P< 0.001.

Figure 7. Proposed working model showing the role of NtHY5 in nicotine and flavonol accumulation in tobacco. We propose that NtHY5 moves from shoot to root to upregulate the expression of nicotine pathway genes in light which results into increased accumulation of nicotine as well as flavonoid biosynthesis. NtHY5 regulates nicotine and flavonol biosynthesis via binding to the G-boxes in the promoters of nicotine and flavonol regulatory genes. Higher accumulation of secondary metabolites leads to less ROS accumulation thus increased salt stress tolerance ability. Higher nicotine and flavonoids also regulates growth and development of tobacco. Arrowheads and tees indicate positive and negative regulation, respectively.