THEWEIGHT: A simple and flexible algorithm for simulating non-ideal, age-structured populations

Robin S. Waples

[1]Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA, 98112 USA

Running head: Simulating non-ideal populations

Key words: Wright-Fisher model, population genetics, computer simulations, reproductive skew, effective population size

Corresponding author:
Robin Waples (robin.waples@noaa.gov)
Northwest Fisheries Science Center
2725 Montlake Blvd. East
Seattle, WA, 98112 USA
(206) 860-3254 (voice); (206) 860-3335 (FAX)

January 2022

Abstract

1. The Wright-Fisher model, which directs how matings occur and how genes are transmitted across generations, has long been a lynchpin of evolutionary biology. This model is elegantly simple, analytically tractable, and easy to implement, but it has one serious limitation: essentially no real species satisfies its many assumptions. With growing awareness of the importance of jointly considering both ecology and evolution in eco-evolutionary models, this limitation has become more apparent, causing many researchers to search for more realistic simulation models.

2. A recently described variation retains most of the Wright-Fisher simplicity but provides greater flexibility to accommodate departures from model assumptions. This generalized Wright-Fisher model relaxes the assumption that all individuals have identical expected reproductive success by introducing a vector of parental weights w that specifies relative probabilities different individuals have of producing offspring. With parental weights specified this way, expectations of key demographic parameters are simple functions of w. This allows researchers to quantitatively predict the consequences of non-Wright-Fisher features incorporated into their models.

3. An important limitation of the Wright-Fisher model is that it assumes discrete generations, whereas most real species are age-structured. Here I show how an algorithm (*THEWEIGHT*) that implements the generalized Wright-Fisher model can be used to model evolution in age-structured populations with overlapping generations. Worked examples illustrate simulation of seasonal and lifetime reproductive success and show how the user can pick vectors of weights expected to produce a desired level of reproductive skew or a desired N_e/N ratio. Alternatively, weights can be associated with heritable traits to provide a simple, quantitative way to model natural selection. Using *THEWEIGHT*, it is easy to generate positive or negative correlations of individual reproductive success over time, thus allowing explicit modeling of common biological processes like skip breeding and persistent individual differences.

4. R code is provided to implement basic features of *THEWEIGHT* and applications described here. However, required coding changes to the Wright-Fisher model are modest, so the real value of the new algorithm is to encourage users to adopt its features into their own or others' models.

1 | INTRODUCTION

The Wright-Fisher (WF) model of reproduction is a lynchpin of evolutionary biology, as it describes how matings occur and how genes are transmitted across generations. This model also provides a direct link between the ecological consequences of abundance, which depend on census size (N), and the evolutionary consequences of abundance, which depend on effective population size (N_e). When populations dynamics follow the standard WF model, the number of offspring per parent is approximately Poisson distributed (with variance ≈ mean), in which case the census and effective sizes are the same. Since the time of the Modern Synthesis (Huxley 1942), theoretical population genetics has played a seminal role in shaping the development of evolutionary biology. Virtually every equation in population genetics includes a term for effective size, and it is routine to adopt the assumptions of the WF model to take advantage of this equivalence between N and N_e. Under WF assumptions (a single closed population of constant N, discrete generations, random mating and random reproductive success), rates of random genetic drift, inbreeding, and loss of neutral genetic variability all depend only on the number of reproductive individuals, so both demographic and evolutionary processes can be modeled with a single parameter for population size. A population that satisfies all of these assumptions is said to be "ideal." These features make the WF model easy to simulate in a computer, and as a consequence it is incorporated into a wide variety of software for forward-in-time and coalescent simulations (Peng et al. 2015; NCI 2018).

Despite its advantages and nearly universal appeal, the standard WF model has an important limitation: its many assumptions are compatible with essentially zero species in nature. Almost all real populations deviate from WF assumptions in ways that cause the effective size to be less than the number of adults, primarily because of greater-than-Poisson variance in offspring number, and these deviations lead to a wide range of N_e/N ratios in nature (Frankham 1995; Palstra and Fraser 2012). From the evolutionary perspective this might not be a problem, as a researcher who wants to model genetic processes in a real population with census and effective sizes of, say, $N = 1000$ and $N_e = 200$, could just model a WF population with $N = N_e = 200$. However, populations with $N = 200$ and $N = 1000$ generally are not equivalent for ecological processes such as competition, predation, and individual behavior, so exclusive reliance on WF assumptions constrains the ability of a model to adequately capture realistic eco-evolutionary dynamics. In recent decades it has become apparent that most problems in biology

are not strictly ecological or evolutionary but rather are eco-evolutionary in nature (Pelletier et

al. 2009), and researchers have increasingly expressed the desire for more realistic models for

population genetic simulations (Haller and Messer 2019).

Because the WF model requires numerous simplifying assumptions, there are many ways

to model non-ideal populations (e.g., Eldon and Wakeley 2006; Der et al. 2011; Chotibut and

Nelson 2017). A common approach to incorporate non-random variation in reproductive success

is to draw offspring numbers from a negative binomial or gamma distribution, which can be

scaled to produce a desired level of reproductive skew. This approach, however, is better suited

to modeling population-level processes than characteristics of individuals. Here I describe in

detail an algorithm, *THEWEIGHT*, which makes a simple change to the WF model that provides

considerable added flexibility to simulate realistic, individual-based demographics. In the

standard WF model, every individual has an equal chance to produce each offspring; in

THEWEIGHT, the user defines a vector of parental weights that specifies the relative probabilities

that each individual will be chosen as a parent for a given offspring. Individual weighting

schemes have been incorporated into some genetic simulators, such as *Nemo* (Guillaume and

Rougemont 2006), *simuPOP* (Peng and Amos 2008), and *SFS_CODE* (Hernandez 2008).

THEWEIGHT takes advantage of recent theoretical developments that show that many key

demographic parameters can be predicted based on simple functions of the vector of parental

weights. However, these results, and the original description of the generalized Wright-Fisher

model (Waples 2020), followed the traditional WF model in assuming that generations are

discrete.

Most real populations are age structured, so the discrete-generation assumption is an

important limitation. Here I illustrate how *THEWEIGHT* algorithm can be used to model eco-

evolutionary processes in iteroparous species with overlapping generations. The treatment here

focuses on the large fraction of species that exhibit strongly seasonal, birth-pulse reproduction

(Caswell 2001). For these species, reproductive success is commonly assessed from two

different perspectives: seasonal (in which offspring number is the total produced by an

individual across a reproductive season—here assumed to be years) and lifetime (in which

lifetime reproductive success (*LRS*) is the cumulative number of offspring produced across an

individual's lifespan). Variance in *LRS* has received considerable attention (Hill 1972; Brown

1988; Kruuk et al. 1999; Tuljapurkar et al. 2020) because it integrates information across

generational time scales and is the most important factor determining effective population size

(N_e) per generation. However, analysis of data on seasonal reproduction provides important

insights into mating systems. Furthermore, seasonal reproduction is much easier to study,

especially in long-lived species, and many monitoring programs routinely estimate the effective

number of breeders per year (N_b) and other population genetic parameters using seasonal data

(Pudovkin et al. 1996; Ruzzante et al. 2016; Ackerman et al. 2017).

Worked examples illustrate how *THEWEIGHT* can be used to model (1) changes in

fecundity with age; (2) variation in expected reproductive success among individuals of the same

age and sex; and (3) positive or negative correlations in individual reproductive success over

time. This latter feature is useful for modeling common biological features such as intermittent

breeding and persistent individual differences (aka "individual quality"; Wilson and Nussey

2010). Using *THEWEIGHT*, it is also easy to simulate null models for age-structured species that

incorporate random survival and/or random variation in reproductive success; these null models

in turn provide important context for evaluating empirical data. Computer code is provided to

illustrate how these features can be implemented, but the required coding changes are simple

enough that users can easily incorporate them into their own or others' models.

2 | METHODS

2.1 | Wright-Fisher reproduction

2.1.1 | The standard Wright-Fisher model

The standard Wright-Fisher model has the following features: population size is

constant; generations are discrete; selection is absent; mating is random; and every individual is

equally likely to be the parent of a given offspring, regardless how many or few offspring that

individual has already produced. Conceptually, reproduction occurs as follows: each individual

contributes an equal and very large number of gametes to a huge (essentially infinite) pool of

gametes, which unite at random to produce zygotes. Random survival of zygotes then produces

the few individuals that form the next generation (Wright 1931). A population meeting all of

these criteria is said to be "ideal" and has the property that the effective population size (N_e)

equals (on average) the adult census size (N).

This latter feature can be illustrated using standard definitions of inbreeding N_e based on

the mean (μ_k) and variance (σ_k^2) among parents in number of offspring (k) (Crow and Denniston

1988):

Inbreeding $N_e = \dfrac{\mu_k N - 1}{\mu_k - 1 + \sigma_k^2/\mu_k} = \dfrac{\mu_k N - 1}{\mu_k - 1 + \phi}$,   (1)

where $\phi = \sigma_k^2/\mu_k$ is the ratio of variance to mean offspring number. Focus is on inbreeding N_e

because it relates to the number of parents and hence is more relevant than variance N_e to the

topics considered here. Furthermore, inbreeding N_e is much less sensitive to effects of sampling

and hence more flexible than variance N_e, especially for applications involving age-structured

populations (Waples 2002, 2020). Equation 1 applies to monoecious/hermaphroditic diploids

with random selfing; slight variations are available for other mating systems. Random

reproductive success leads to multinomial sampling of parents, so $E(\sigma_k^2) \approx \mu_k$. Inserting $\phi = 1$

into Equation 1 yields the result that $N_e \approx (\mu_k N - 1)/\mu_k \approx N$. [The result is exact if one uses the

exact (binomial) variance $E(\sigma_k^2) = \mu_k(N - 1)/N$ rather than the Poisson approximation.]

The WF model is easy to implement in a computer, the only required input parameters

being the number of ideal individuals (N) and, if genotypes are modeled, the number of gene loci

being tracked. Then, only three more steps are required to complete one generation of

reproduction (Figure 1A). In Step 2, each offspring "chooses" its parents randomly and with

replacement from the N potential parents. In Step 3, at each gene locus one of the two alleles

from each parent is randomly chosen to pass on to the offspring. Step 4 completes the cycle, as

offspring become potential parents of the next generation.

2.1.2 | The generalized Wright-Fisher model

Conceptually, the generalized WF model differs in one important way from the standard

version: the parents are allowed to make unequal contributions to the initial gene pool, with the

contributions being proportional to the weight (w_i) for each parent. The first use of a weighted

WF model like this was by Robertson (1961) for the special case of full-sibling families, and the

model was further developed by Waples (2020). Let μ_W and σ_W^2 be the mean and variance,

respectively, of the vector of parental weights, and let $CV_W = \sigma_W/\mu_W$ be the coefficient of

variation of w. [CV_W is the same when calculated using the raw parental weights w or weights

that are standardized so that they sum to 1, so here the raw weights are used]. Then, several key

population genetic parameters can be expressed as simple functions of the squared CV of the

parental weights, $CV_w^2 = \sigma_w^2/(\mu_w)^2$ (Waples 2020):

$$E(\sigma_k^2) = \mu_k[1 + \mu_k CV_w^2] \; ; \qquad\qquad (2)$$

$$E(\phi) = E(\sigma_k^2)/\mu_k = 1 + \mu_k CV_w^2 \; ; \qquad\qquad (3)$$

$$E(\text{Inbreeding } N_e) = \frac{N-1}{1+CV_w^2} \; . \qquad\qquad (4)$$

Equations 2 and 3 are conditional expectations that depend on mean offspring number; they are

expressed in terms of population parameters (μ_k, σ_k^2) but also are unbiased as conditional

expectations for samples (Waples 2020). Equation 4 is a general result for inbreeding N_e that

does not depend on mean offspring number (Waples 2020); for variance N_e, this equation holds

only if $\mu_k = 2$ (Felsenstein 2019). Equation 4 can be converted into an expression that predicts

the N_e/N ratio. If one ignores the "-1" in the numerator of Equation 4, which is of little

importance unless N is very small, and divides both sides by N, the result is:

$$E(N_e/N) \approx \frac{1}{1+CV_w^2} \; . \qquad\qquad (5)$$

When all parental weights are equal, $CV_w = 0$ and Equation 5 predicts $N_e/N = 1$, as expected for

the standard Wright Fisher model.

In computer modeling, the only difference between the traditional and generalized WF

models occurs at Step 2: instead of choosing adults with equal probability to be parents of a

given offspring, the probability a parent will be chosen is proportional to its weight (Figure 1B).

To deal effectively with age structure, it is also important to allow the overall number of

offspring to vary from the constraint in the WF model that $\mu_k = 2$. Modeling separate sexes

requires separate routines for choosing male and female parents, together with separate vectors

of parental weights. Effective size can then be calculated separately for males and females (N_{em},

N_{ef}), and overall N_e can be obtained using Wright's (1938) sex ratio adjustment: $N_e = 4N_{em}N_{ef}$

$/(N_{em} + N_{ef})$.

2.2 | Modeling age-structured populations with *TheWeight* algorithm

2.2.1 | Seasonal reproduction

2.2.1.1 | Age-specific vital rates

For seasonal reproduction in a discrete-time, birth pulse model, age structure can be modeled using age-specific vital rates from a life table: v_x = probability of surviving from age x to $x+1$, b_x = mean number of offspring produced by an individual of age x, and N_x = number of individuals in age class x. In an age-structured model based on *TheWeight*, a simple option is to assign parental weights based on the b_x value for the appropriate age. If all individuals of the same age receive the same weight, then each age class behaves like a mini WF population, with the expected age-specific variance in offspring number being approximately equal to the mean: $E(\sigma^2_{k(x)}) \approx b_x$. In that case, the expected value of the variance-to-mean ratio (ϕ_x) is $E(\phi_x) = E(\sigma^2_{k(x)})/b_x \approx 1$.

Assuming that $\phi_x = 1$ for each age and sex might be reasonable for some populations, but that is not universally the case. Empirical data for age-specific ϕ_x are seldom published, but (for example) for black bears in Michigan, USA, ϕ_x was significantly >1 for both females and males and estimated to be 10 or higher for males aged 6-10 years (Waples et al. 2018a). A more comprehensive life table would then include another age-specific vector of vital rates: ϕ_x.

To model the desired level of overdispersion at each age (indicated by ϕ_x), the following steps can be used to generate N_x parental weights. This process can be done separately for male and female parents of each offspring.

Step 1. Determine the target value for the squared *CV* of parental weights. By rearranging Equation 3,

$$TargetCV^2_{w,x} = (\phi_x-1)/b_x. \tag{6}$$

Step 2. For any given $TargetCV^2_{w,x}$, it is possible to identify a Poisson distribution with mean = variance = λ_x that will produce a vector of parental weights such that $CV^2_{w,x} = var(w_x)/[mean(w_x)]^2 = \lambda_x/(\lambda_x)^2 = 1/\lambda_x$. The required mean of the Poisson distribution is thus $\lambda_x = 1/TargetCV^2_{w,x}$.

Step 3. Simulating N_x random Poisson values using parameter λ_x will thus, on average, produce a vector of parental weights with the desired $CV^2_{w,x}$. This process can be iterated until the

realized $CV_{w,x}^2$ is within a specified tolerance of the target (results presented here used

tolerance = 1%).

With the parental weights specified for each age, seasonal reproduction in the generalized WF

model can proceed using either a two-step or one-step process.

Step 4A. In the two-step process, for each offspring one first determines the age of the parent(s).

This can be done randomly (e.g., by sampling the parent's age from the ages of the adult

lifespan, with relative probabilities being the product $N_x b_x$). Next, the actual parent can be

chosen randomly using the appropriate age-specific vector of parental weights generated in

Step 3.

Step 4B. In the single-step process, adults of all ages (and their weights) are combined into a

single matrix, and the parent(s) for each offspring are chosen based on the combined vector

of parental weights. This requires that the mean weight for each age ($\overline{w}_x$) be proportional to

the age-specific mean fecundity (b_x). If b_x is constant, the weighting scheme described

above already accomplishes this. More generally, if b_x varies with age, then for each age

every parental weight has to be rescaled by multiplying by the factor $F = b_x/\lambda_x$. After

rescaling, $E(\overline{w}_x) = b_x$ for each age, and, although both the mean and variance of the weights

have changed, the squared CV remains the same. Alternatively, before generating the

weights, the initial Poisson factor for each age can be multiplied by the same factor F.

2.2.1.2 | Natural selection

Stabilizing selection is commonly modeled using a Gaussian fitness function applied to a

continuous phenotypic trait z, such as body size (Lande 1976). A general way to implement this

using *THEWEIGHT* is as follows:

Step 1: Define the phenotypic distribution, $p(z)$. z is commonly assumed to have a normal

distribution with mean = μ_Z and variance = σ_Z^2. In that case,

237
$$p(z) = \frac{1}{\sigma_z\sqrt{2\pi}}\, e^{-\frac{[z-\mu_z]^2}{2\sigma_Z^2}} \quad .$$

238   Step 2: Map the phenotypes to the fitness function to produce the distribution of relative

239   individual fitness $p(w)$, which is identical to the distribution of parental weights. Under

240   stabilizing selection, the maximum fitness (w_{max}) occurs at an intermediate phenotype,

241   denoted by θ.

242
$$w = w_{max}\ e^{-\frac{(z-\theta)^2}{2\omega^2}}\quad,$$

where $\omega = \sigma_w$ is the "width" of the fitness surface.

Steps 1 and 2 are commonly used to model selection. The advantage of using *THEWEIGHT* is that

the expected consequences of selection for key demographic and population genetic parameters

can be quantified.

Equation 3 above shows that the expected value of the variance to mean ratio for

offspring number ($E(\sigma_k^2)/\mu_k$) is a simple function of the squared *CV* of parental weights. A

related quantity defined by Crow (1958) is $I = \sigma_k^2/(\mu_k)^2 = \phi/\mu_k$. Commonly referred to as the

Opportunity for Selection, I is the variance in relative fitness, which places an upper limit on the

evolutionary response to selection (Walsh and Lynch 2018). Dividing each side of Equation 3

by the mean offspring number produces this result:

$E(\sigma_k^2)/{\mu_k}^2 = E(I) = 1/\mu_k + CV_W^2.$ (7)

If the phenotypic distribution is continuous, the distributions of fitnesses and parental weights are

also continuous, in which case the means and variances can be calculated using the following

formulas (lower bounds of the integrals are set to 0 because fitnesses and parental weights cannot

be negative):

$E(w) = \int_0^\infty wp(w)dw$;

$E(w^2) = \int_0^\infty w^2 p(w)dw$;

$\sigma_w^2 = E(w^2) - [E(w)]^2$;

$CV_w^2 = \sigma_w^2/[E(w)]^2$.

The above example assumed a Gaussian distribution of selection coefficients, but this approach

can also be used more generally with any selection function (linear, quadratic, etc.). Equations

2-5 and 7 do not depend on the distribution of parental weights—only the *CV*.

2.2.1.3 | Generic weighting schemes

Generic weighting schemes that assign weights randomly to individuals are easy to

implement. One way to do this is to randomly assign potential parents ordered cardinal numbers

$i = 1, 2, \dots N$ and then give each a relative weight that is a function of their assigned number.

Simple examples include power functions ($w_i = i^x = 1^x, 2^x, \dots N^x$) and harmonic functions ($w_i =$

$1/i^x = 1/1^x, 1/2^x, \dots 1/N^x$). Equations 2-5 can be used to predict the consequences of these

weighting schemes for parameters of interest.

2.2.2 | Lifetime reproductive success

Analysis of *LRS* focuses on a birth cohort of individuals that are enumerated at a

specified age (here assumed to be age at maturity). *THEWEIGHT* has three general options for

modeling lifetime reproductive success. The simplest approach is to generate new weights for

each age to match the species' vital rates (as described in Section 2.2.1.1) and then assign those

randomly to the survivors each year. This approach is suitable if it is reasonable to assumes that

an individual's reproductive success is uncorrelated over time (e.g., Felsenstein 1971; Hill 1972).

THEWEIGHT can also be used to model two common scenarios in which this independence

assumption does not hold.

2.2.2.1 | Intermittent breeding

In many species, females (and sometimes males) do not necessarily breed every year,

leading to a strategy that has been called "skip breeding" (Goutte et al. 2011) or "intermittent

breeding" (Shaw and Levin 2011). This type of life history can be found in vertebrate

ectotherms, mammals, birds, and some invertebrates; it can lead to positive or negative

correlations in the number of offspring an individual produces over time, depending on the time

lag and pattern of skip breeding. This life history can be described by a vector θ, the t^{th} element

of which (θ_t) is the probability that an individual will reproduce in the current year, given that it

last reproduced t years before (Shaw and Levin 2011). For example, Waples and Antao (2014)

estimated that $\theta_t = [0.025, 0.443, 0.634, 0.743, 1]$ for $t=1,5$ for female loggerhead turtles,

reflecting substantial costs of migration and reproduction. In contrast, for female black bears in

Michigan $\theta_t = [0.035, 0.94, 1]$ for $t=1,3$ (Waples et al. 2018b), indicating a faster recovery of

reproductive potential. This life-history can easily be modeled with *THEWEIGHT*.

Step 1: For each individual, determine the value of t = the number of years since the individual

last reproduced.

Step 2: With probability θ_t, replace that individual's weight with 0.

2.2.2.2 | Persistent individual differences

Persistent individual differences in reproductive success occur when some individuals are

consistently above or below average (for their age and sex) at producing offspring. In

iteroparous species, persistent individual differences increase lifetime variance in offspring

number and reduce N_e (Lee et al. 2011, 2020). Variation in size at age can easily lead to this

type of pattern: an individual that is larger (and hence more fecund) than others in its cohort at

age x is also likely to have above-average size and fecundity at older ages. Persistent differences

could have a genetic basis or might simply reflect early-life-history environmental "silver spoon"

effects with long-term consequences.

This life history feature can be modeled in *THEWEIGHT* by introducing positive

correlations between parental weights over time, which can be accomplished by retaining

individual weights across lifetimes. But because the target value of desired $CV_{w,x}^2$ is contingent

on b_x, when fecundity changes with age the squared *CV* of the parental weights must also

change. Doing this without distorting the relative rankings of the individual weights is tricky but

can be accomplished as follows:

Step 1. Using the approach outlined in the previous section, generate a vector of parental

weights w_1 for the N_1 age-1 individuals in a cohort, for which expected fecundity is b_1 and

$TargetCV_{w,1}^2 = (\phi_1\text{-}1)/b_1$. The Poisson parameter to produce the desired parental weights at

age 1 is $\lambda_1 = 1/TargetCV_{w,1}^2$. If age-1 individuals are not mature, start this process with the

age at sexual maturity.

Step 2. Now we need a vector of weights w_2 that retain the same framework of individual

differences but produce $TargetCV_{w,2}^2$ at age 2 = $(\phi_2\text{-}1)/b_2$. This can be achieved by adding

a constant C_2 to each individual weight generated at age 1. This has no effect on the

variance of the weights but changes the mean, so it also changes the squared *CV*. Define the

factor $Q_2 = TargetCV_{w,2}^2/TargetCV_{w,1}^2$. Then, $TargetCV_{w,2}^2 = \sigma_{w,2}^2/(\overline{w}_2)^2 =$

$Q_2 * \sigma_{w,1}^2 / (\bar{w}_1)^2$. Now σ_w^2 is the same at ages 1 and 2, so this last equation implies that

$1/(\bar{w}_2)^2 = Q_2/(\bar{w}_1)^2$ and hence $\bar{w}_2 = \bar{w}_1/\sqrt{Q_2}$.

Step 4. If the increase in the mean weight is accomplished by adding a constant, it is also true

that $\bar{w}_2 = \bar{w}_1 + C_2$. Solving these two equation yields the desired result: $C_2 = \bar{w}_1 * [1/\sqrt{Q_2}) -$

1].

Step 5. Repeat this process for each additional year of the maximum adult lifespan (*AL* years).

The result is a N_1 x *AL* matrix of parental weights. Each year, weights for the survivors in a

cohort are taken from the appropriate column of the matrix.

Adding a constant to all individual weights changes their relative values. However, it maintains

the same rank order of weights, and the pairwise correlations of the age-specific vectors of

individual weights are all $r = 1.0$. Weaker positive correlations could be introduced by adding a

random normal error term to the individual weights each year without shuffling them.

2.3 | Simulations

To illustrate calculations described above and evaluate their accuracy, seasonal and

lifetime reproductive success was simulated using *TheWeight* algorithm. Simulations were

conducted in R (R Core Team 2021) using code available in Supporting Information.

3 | RESULTS

3.1 | Seasonal reproduction

3.1.1 | Modeling age-specific vital rates

Table 1 illustrates how to compute the vectors of parental weights when modeling

seasonal reproduction according to age-specific vital rates from an expanded life table, which

also includes age-specific values for ϕ for a single sex, nominally female. The hypothetical

species is first enumerated at age 1, matures at age 3, and has a maximum lifespan of 10 years.

With annual survival = 0.7 and a fixed cohort size of 1000 age-1 females, expected numbers of

females alive at subsequent age are given by the vector N_x. Expected fecundity (b_x) is

proportional to age and is scaled to production of a stable population (which produces 2000

yearling offspring per year, 1000 of each sex). In this species, the overdispersion index ϕ was

also chosen to increase with age, but more slowly than fecundity. The expected age-specific

variance in offspring number is $E(\sigma^2_{k,x}) = b_x * \phi_x$, adjusted by the factor $N_x/(N_x-1)$ to account for

finite sampling. For each age, Table 1 shows how to calculate the $TargetCV^2_{w,x}$ that will

produce the desired level of overdispersion, and the respective Poisson parameter that can be

used to simulate random weights with the desired properties. When using the two-step process

to simulate reproduction (Step 4A in Section 2.2.1.3), the age(s) of the parent(s) of each

offspring are first chosen, and then the actual parent is chosen from that age group, with

sampling probabilities determined by the parental weights. With this two-step approach, the

initial Poisson parameters (λ_x) can be used for each age. In this example, however, reproduction

is modeled in a single step (as in Step 4B in Section 2.2.1.3), and the weights for each age have

to be rescaled so that age-specific $\overline{w}_x = b_x$. This is accomplished by multiplying the initial λ_x by

an age-specific scaling factor. Across 1000 replicate simulated populations, mean realized ϕ_x for

each age closely agreed with the parametric expectations (Table 1). Table 1 also shows how to

calculate expected reproductive-success parameters across adults of all ages ('Exp' column at

bottom of the table), and average values observed in the simulations also agreed closely with

those expectations.

Most life tables do not contain data for the age-specific parameter ϕ_x, so it is common to

assume that all $\phi_x = 1$. The consequences of violation of this assumption can easily be modeled

using *THEWEIGHT* by replacing empirical ϕ_x estimates with 1.0. This was done for the life table

in Table 1, with results shown in Table S1. Across parents of all ages, assuming all $\phi_x = 1$ has no

effect on the overall mean offspring number ($\mu_k = 1.30$), but it is expected to reduce overall σ^2_k

from 5.04 to 1.57 and overall ϕ from 3.88 to 1.21—both reductions of almost 70%. Simulations

using *THEWEIGHT* with ϕ_x set to 1 agreed very closely with these expectations (Table S1).

These examples used data for the entire population. Subsampling offspring is easy to

accomplish using *THEWEIGHT*, as illustrated in the R code provided.

3.1.2 | Modeling selection

In the example for seasonal reproduction shown in Table 1, $\mu_k =1.30$ and the expected

value of CV^2_W is 2.21, so the expected Opportunity for Selection is (from Equation 7) $E(I) =$

$1/1.30 + 2.21 = 2.98$. Interpretation of raw values of I is complicated by its dependence on mean

offspring number, which can reflect experimental design and sampling effort more than

underlying biology (Downhower et al. 1987; Fairbairn and Wilby 2001). A simple way to

resolve this problem is to subtract the expected contribution from random demographic

stochasticity ($1/\mu_k$), which produces an adjusted Opportunity for Selection that is independent of

mean fitness (Waples 2020):

$E(\Delta_I) = E(I) - 1/\mu_k = CV_W^2 \,.$ (8)

The adjusted Opportunity for Selection index, which is simply the squared *CV* of parental

weights and can be calculated directly from the distribution of relative fitness, represents the

component of variance in relative fitness that exceeds that expected under a null model of

random reproductive success.

For the example shown in Table 1, $E(\Delta_I) = 2.21$, and the empirical mean across 1000

replicate simulations was 2.24. This value does not depend on mean offspring number and hence

can be directly compared to similar values obtained in other studies. Knowing CV_W^2 also allows

one to predict the expected reduction in the N_e/N ratio that can be attributed to natural selection.

For this example, $E(N_e/N)$ based on Equation 5 = 1/(1+2.21) = 1/(1+ $E(\Delta_I)$) = 0.311, indicating

that unequal expectations for reproductive success attributed to selection would be expected to

drop the effective population size to less than one third of the number of adults.

3.1.3 | Generic weighting schemes

The consequences for the N_e/N ratio of applying some simple rules for specifying

parental weights are shown in Figure 2. In the first three scenarios, the weights were power

functions that produced increasing amounts of reproductive skew: $w_i = i^1$, i^2, and i^3. The next

three scenarios used the analogous harmonic series: $w_i = 1/i^1$, $1/i^2$, and $1/i^3$; and the final scenario

used an exponential series: $w_i = e^{\,i}$. It is apparent from Figure 2 that the power functions produce

N_e/N ratios that are largely insensitive to *N*, whereas for the other series log(N_e/N) declines

approximately linearly with log(*N*). The number of possible weighting schemes is effectively

unlimited, but these simple examples show that it is easy for users to pick a weighting scheme

that will produce essentially any desired N_e/N ratio or any desired Opportunity for Selection.

3.2 | Lifetime reproductive success

Table 2 uses the vital rates in Table 1 to illustrate how to generate parental weights that

produce persistent individual differences in reproductive success over time. Since ages 1 and 2

are juveniles, this table begins with age at maturity = 3 years. First, the $TargetCV^2_{w,x}$ expected

to produce weights with the desired level of overdispersion at each age is computed as $(\phi_x\text{-}1)/b_x$,

and the Poisson parameter to generate weights for age-3 adults is then $\lambda_3 = 1/TargetCV^2_{w,3}$—all

as in Table 1. Next, age-specific constants are calculated as $C_x = \lambda_3 * (1/\sqrt{Q_x} - 1)$, where $Q_x =$

$TargetCV^2_{w,x}/TargetCV^2_{w,3}$, and adjusted parental weights for subsequent ages are calculated as

$Adj(\overline{w}_x) = \lambda_3 + C_x$. Results of the simulations show that although these adjustments changed

both the means and variances of the age-specific parental weights, they did not change $CV^2_{w,x}$,

and the actual mean ϕ_x values were very close to the parametric expectations for each age (Table

2), just as they were for the example in Table 1. The adjusted weights, however maintain the

relative weights for each individual over time.

Once the matrix of lifetime parental weights is established, it requires only a single line

of code to switch between modeling reproductive success that is independent and perfectly

correlated over time: for independence the parental weights are randomly scrambled every year,

and to model persistent individual differences they are not. This can make a large difference in

terms of reproductive skew: with parental weights randomized each year, the median variance in

LRS across 1000 simulations for the 490 females in each cohort that reached age 3 was 30.7;

with parental weights retained across lifetimes, median $\sigma^2_{k_\bullet}$ more than doubled, to 76.1.

Persistent individual differences also greatly skewed the distribution of *LRS* compared to a null

model of independence (Figure 3). In simulations for which parental weights were retained,

almost half of the females that reached age at maturity never produced an offspring, while some

others produced 100 or more. When weights were shuffled each year, maximum *LRS* was <60

and null parents comprised a bit over a third of each cohort.

The examples in Tables 1 and 2 calculated means and variances of offspring number only

for adults. If juveniles (all of which produce 0 offspring) had also been included, the overall

means would have been lower and the variance-to-mean ratios would have been higher. Hence,

to avoid confusion resulting from apples-and-oranges comparisons, it is important to clarify the

groups of individuals that are included in analysis of reproductive success.

4 | DISCUSSION

4.1 | Model comparison

The Wright-Fisher model—the most widely-used null model in evolutionary biology—has two distinctive features: equiprobability and independence. Under WF, every individual is equally likely to be the parent of any given offspring, so offspring can be thought of as assigned to parents randomly and with equiprobability. Furthermore, each assignment is independent, in the sense that the outcome does not depend on results of any previous assignments. In R (for example), the WF process can be modeled using the 'sample' function, with *size* = the number of offspring and sampling done from a vector of parental names or IDs. Under the defaults, sampling is equiprobable and done with replacement and hence independent.

In the generalized WF model (Waples 2020), the assignments are still independent (within time steps at least) but parental assignment probabilities are allowed to vary according to the vector of parental weights, w. This only requires adding a 'prob = w' argument to the 'sample' function. Unequal parental weights lead to overdispersed variance in offspring number compared to the Poisson expectation, with the degree of overdispersion captured by the age-specific vector $\phi = \sigma_k^2/\mu_k$.

Another common approach to modeling overdispersed variance is to randomly assign numbers of offspring to parents based on a negative binomial or gamma distribution that can be parameterized to generate a target μ_k and σ_k^2 (e.g., Anderson and Dunham 2005). This approach is fast and simple but has some limitations compared to *THEWEIGHT*. First, when offspring numbers are chosen randomly from a distribution, it is not easy (as it is in *THEWEIGHT*) to relate expected reproductive success to traits or attributes of individuals. Second, this same limitation makes it difficult to control the degree of correlation in expected individual reproductive success over time, which can easily be accomplished using *THEWEIGHT*.

4.2 | Age structure

Age structure is one of the most common ways that most real populations depart from WF assumptions. The generalized WF model is well suited for dealing with overlapping generations, and that was the motivation for developing the features of *THEWEIGHT* algorithm described here. In Methods, procedures are outlined that allow users to generate parental weights designed to produce the desired degree of reproductive skew for each age and sex. For

seasonal reproduction, this also allows the user to control the overall degree of reproductive

skew across all individuals reproducing in a single time period. For lifetime reproductive

success, the user has various options for adjusting individual weights (or not) between time steps,

and this allows realistic modeling of *LRS* for common life-history traits like intermittent breeding

and persistent individual differences.

4.3 | Modeling selection

The standard WF model assumes a very large gamete pool equally produced by *N* parents

and random survival of zygotes until the next generation of *N* offspring is produced, leading to

$\mu_k = 2$. The generalized WF model also assumes random survival of zygotes but allows for

unequal parental contributions to the initial gene pool and allows mean offspring number to vary;

this latter feature is important for modeling seasonal reproduction in iteroparous species. As a

consequence, *THEWEIGHT* algorithm is well suited for modeling fertility selection but does not

explicitly model viability selection. However, the latter can also easily be incorporated by

inserting an episode of selective mortality that relates probability of survival to a trait of interest.

4.4 | Null models

Null models are increasingly important in evolutionary biology, including for the analysis

of reproductive success (van Daalen and Caswell 2017; Tuljapurkar et al. 2020). A variety of

null models are easy to implement using *THEWEIGHT*. For seasonal reproduction, one obvious

null model is to treat all adults as a single WF population, which can be accomplished by setting

fecundity to be constant with age and all $\phi = 1$. For *LRS*, randomly scrambling parental weights

each year replicates a null model of independence of reproductive success over time.

Comparison with results for this null model can dramatically illustrate the consequences of a lack

of independence (see Figure 3).

4.5 | Sweepstakes reproductive success

'Sweepstakes' reproductive success begins with ~random production of eggs by a highly

fecund species, followed by an episode of extreme, family-correlated mortality during which all

offspring from most families perish and only a lucky few families survive to produce the new

cohort (Hedgecock 1994). This type of phenomenon has been suggested as a way to explain

'tiny' estimates of the effective size : census size ratio ($N_e/N < 10^{-3}$) in a number of marine

species (Hauser and Carvalho 2008; Hedgecock and Pudovkin 2011). Conceptually,

sweepstakes reproductive success differs from either the standard or generalized WF models in

that assignments of offspring to parents are not independent; instead, they are highly correlated.

Sweepstakes reproduction could potentially be modeled using *THEWEIGHT*; however, because

tiny N_e/N ratios are only possible if all ϕ_x values are very large (Waples 2016), the parental

weights would have to be so highly skewed that only a few parents have any chance of

producing offspring. An alternative would be to explicitly model this lack of independence:

after two parents are randomly selected, they are allowed to produce not a single offspring (as in

the standard and generalized WF models) but some large number (K) of offspring. Waples and

Anderson (2017) used a model like that to evaluate the consequences of purging putative siblings

from population-genetic datasets. If K were large enough, this approach could lead to very high

σ_k^2 and mimic sweepstakes reproductive success.

Acknowledgments

527   I thank Eric Anderson, Tom Reed, and Ryan Waples for useful comments and suggestions.

528   The author declares no conflict of interest.

Data availability statement

531   This study did not generate any new empirical data, except by simulation. Code to

532   conduct the simulations and analyses described here is available in Supporting Information.

REFERENCES

Ackerman, MW, BK Hand, RK Waples, G Luikart, RS Waples, C Steele, BA Garner, J McCane, & M Campbell. (2017). Effective number of breeders from sibship reconstruction: empirical evaluations using hatchery steelhead. Evolutionary Applications 10:146-160.

Anderson, E. C., & Dunham, K. K. (2005). spip 1.0: a program for simulating pedigrees and genetic data in age-structured populations. *Molecular Ecology Notes*, *5*, 459–461.

Caswell, H. (2001). Matrix Population Models, Ed. 2. Sinauer, Sunderland, MA.

Chotibut, T. & Nelson, D.R. (2017). Population genetics with fluctuating population sizes. *Journal of Statistical Physics*, *167*(3-4), 777-791.

Crow, J. F. (1958). Some possibilities for measuring selection intensities in man. Human Biology 30, 1:13.

Crow, J. F., & N. E. Morton. (1955). Measurement of gene frequency drift in small populations. Evolution 9, 202-214.

Crow, J.F., & C. Denniston. (1988). Inbreeding and variance effective population numbers. *Evolution* 42, 482-495.

Der, R., Epstein, C.L. & Plotkin, J.B. (2011). Generalized population models and the nature of genetic drift. *Theoretical population biology*, *80*(2), 80-99.

Downhower, J.F., Blumer, L.S. & Brown, L. (1987). Opportunity for selection: an appropriate measure for evaluating variation in the potential for selection? *Evolution*, **41**, 1395–1400.

Eldon, B. & Wakeley, J. (2006). Coalescent processes when the distribution of offspring number among individuals is highly skewed. *Genetics*, *172*(4), 2621-2633.

Fairbairn, D.J. & Wilby, A.E. (2001). Inequality of opportunity: measuring the potential for sexual selection. *Evolutionary Ecology Research*, *3*(6), 667-686.

Felsenstein, J. (1971) Inbreeding and variance effective numbers in populations with overlapping generations. Genetics 68, 581–597.

Felsenstein, J. (2019). Theoretical evolutionary genetics. Lecture notes, University of Washington, Seattle. Available at: http://evolution.gs.washington.edu/pgbook/pgbook.pdf.

Frankham, R. (1995). Effective population size/adult population size ratios in wildlife: a review. *Genetics Research*, *66*(2), 95-107.

Goutte, A., M. Kriloff, H.Weimerskirch, & O. Chastel. (2011). Why do some adult birds skip breeding? A hormonal investigation in a long-lived bird. Biol. Lett. 7, 790–792.

Guillaume, F., & J. Rougemont. (2006). Nemo: an evolutionary and population genetics programming framework. *Bioinformatics* 22, 2556-2557.

Haller, B.C. & Messer, P.W. (2019. SLiM 3: Forward genetic simulations beyond the Wright–Fisher model. *Molecular biology and evolution*, *36*(3), 632-637.

Hauser, L. & Carvalho, G.R. (2008). Paradigm shifts in marine fisheries genetics: ugly hypotheses slain by beautiful facts. *Fish and Fisheries*, *9*(4), pp.333-362.

Hedgecock, D. (1994). Does variance in reproductive success limit effective population size of marine organisms? in Genetics and Evolution of Aquatic Organisms, A. R. Beaumont, Ed. (Chapman & Hall, 1994), pp. 122–134.

Hedgecock, D, & A. I. Pudovkin. (2011). Sweepstakes reproductive success in highly fecund marine fish and shellfish: A review and commentary. Bull. Mar. Sci. 87:971–1002.

Hernandez RD. (2008). *A flexible forward simulator for populations subject to selection and demography.*, Bioinformatics, 24, 2786–2787.

Hill, W.G., (1972) Effective size of population with overlapping generations. *Theoretical Population Biology* 3, 278–289.

Huxley, J. (1942). Evolution. The modern synthesis. London: George Alien & Unwin Ltd. 645 p.

Kruuk, L.E.B., Clutton-Brock, T.H., Rose, K.E. & Guinness, F.E. (1999). Early determinants of lifetime reproductive success differ between the sexes in red deer. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *266*(1429), pp.1655-1661.

Lande, R. (1976). Natural selection and random genetic drift in phenotypic evolution. *Evolution*, 314-334.

Lee, A.M., S. Engen, & B.-E. Sæther, (2011) The influence of persistent individual differences and age at maturity on effective population size. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 278, 3303-3312.

Lee, A.M., Myhre, A.M., Markussen, S.S., Engen, S., Solberg, E.J., Haanes, H., Røed, K., Herfindal, I., Heim, M. & Sæther, B.E. (2020). Decomposing demographic contributions to the effective population size with moose as a case study. *Molecular ecology*, *29*, 56-70.

NCI (National Cancer Institute). (2018). Genetic simulation resources (GSR). Available from: https://popmodels.cancercontrol.cancer.gov/gsr/, accessed December 2020.

Palstra, F.P., & D.J. Fraser, (2012) Effective/census population size ratio estimation: a compendium and appraisal. *Ecology and Evolution* 2, 2357–2365.

Pelletier F., Garant D., & Hendry A.P. (2009). Eco-evolutionary dynamics. Phil. Trans. R. Soc. B 364, 1483–1489

Peng, B. & Amos, C.I., (2008). Forward-time simulations of non-random mating populations using simuPOP. *Bioinformatics*, *24*(11), 1408-1409.

Peng, B., Chen, H.S., Mechanic, L.E., Racine, B., Clarke, J., Gillanders, E. & Feuer, E.J., (2015). Genetic data simulators and their applications: an overview. *Genetic epidemiology*, *39*(1), 2-10.

Pudovkin, A.I., Zaykin, D.V. & Hedgecock, D. (1996). On the potential for estimating the effective number of breeders from heterozygote-excess in progeny. *Genetics*, *144*(1), pp.383-387.

R Core Team (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Robertson, A. (1961). Inbreeding in artificial selection programmes. *Genetics Research*, *2*(2), 189-194.

Ruzzante, D.E., McCracken, G.R., Parmelee, S., Hill, K., Corrigan, A., MacMillan, J. & Walde, S.J. (2016). Effective number of breeders, effective population size and their relationship with census size in an iteroparous species, Salvelinus fontinalis. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1823), p.20152601.

Shaw, A. K., & S. A. Levin. (2011). To breed or not to breed: a model of partial migration. Oikos 120, 1871–1879.

Tuljapurkar, S., Zuo, W., Coulson, T., Horvitz, C. & Gaillard, J.M., 2020. Skewed distributions of lifetime reproductive success: beyond mean and variance. *Ecology letters*, *23*(4), pp.748-756.

van Daalen, S.F. & Caswell, H. (2017). Lifetime reproductive output: individual stochasticity, variance, and sensitivity analysis. Theor. Ecol., 10, 355–374.

Walsh, B. & Lynch, M., (2018). *Evolution and selection of quantitative traits*. Oxford University Press.

Waples, R.S. (2002). Evaluating the effect of stage-specific survivorship on the N_e/N ratio. Molecular Ecology 11, 1029-1037.

Waples, R.S. (2016). Tiny estimates of the N_e/N ratio in marine fishes: Are they real? J Fish Biology 89:2479-2504.

Waples, R.S. (2020). An estimator of the Opportunity for Selection that is independent of mean fitness. Evolution 74, 1942-1953.

Waples, R.S., & E.C. Anderson. (2017). Purging putative siblings from population genetic datasets: A cautionary view. Molecular Ecology 26:1211-1224

Waples, R.S., & T. Antao. (2014). Intermittent breeding and constraints on litter size: consequences for effective population size per generation (N_e) and per reproductive cycle (N_b). Evolution 68, 1722-1734.

Waples RS, Scribner K, Moore J, Draheim H, Etter D, & Boersen, M. (2018a). Accounting for age structure and spatial structure in eco-evolutionary analyses of a large, mobile vertebrate. Journal of Heredity 109, 709-723.

Waples RS, Grewe PG, Bravington MV, Hillary R, & Feutry P. (2018b). Robust estimates of a high N_e/N ratio in a top marine predator, southern bluefin tuna. Science Advances 4:eaar7759.

Wilson, A.J. & Nussey, D.H. (2010). What is individual quality? An evolutionary perspective. *Trends in ecology & evolution*, *25*(4), pp.207-214.

Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, *16*:97-159.

Wright, S., (1938). Size of population and breeding structure in relation to evolution. *Science*, *87*, 430-431.

Table 1. A female life table for a hypothetical species, illustrating how to generate the desired level of reproductive skew in seasonal reproduction, using parental weights. Each cohort begins with 1000 age-1 females; annual survival of 0.7 and maturity at age 3 produces a total of N_A = 1539 adult (age 3+) females. For each age x: N_x = number of females alive$;$ b_x = expected fecundity$;$ $E(\sigma^2_{k,x}) = \phi_x b_x (N_x-1)/N_x$; $TargetCV^2_{w,x} = (\phi_x-1)/b_x$; initial $\lambda_x = 1/TargetCV^2_{w,x}$; scaling factor = b_x/initial λ_x; rescaled λ_x = initial λ_x*scaling factor = b_x; $\Sigma(k_{i,x})^2$ = sum of squared offspring numbers; $\bar{w}_x$, $\sigma^2_{w,x}$ = mean and variance in parental weights; $\Sigma w_{i,x}$, $\Sigma(w_{i,x})^2$ = sums of parental weights and squared parental weights. Across all N_A adults, expected (Exp) population parameters are calculated as: $\mu_k = 2*N_1/N_A$; $\sigma^2_k = \Sigma(k_i)^2/N_A$; $\phi = \sigma^2_k/\mu_k$; $\bar{w} = \Sigma w_i/N_A$; $\sigma^2_w = \Sigma(w_i)^2/N_A - \bar{w}^2$; $CV^2_w = \sigma^2_w/\bar{w}^2$. Actual ϕ_x for each age and observed (Obs) results across all ages are averages across 1000 replicate simulations.

Age	N_x	b_x	ϕ_x	$E(\sigma^2_{k,x})$	Target $CV^2_{w,x}$	initial λ_x	scaling factor	rescaled λ_x	$\Sigma(k_{i,x})^2$	Actual ϕ_x	$\bar{w}_x$	$\sigma^2_{w,x}$	$\Sigma w_{i,x}$	$\Sigma(w_{i,x})^2$
1	1000	0												
2	700	0												
3	490	0.805	3.00	2.420	2.484	0.403	2.000	0.805	1503.3	2.97	0.805	1.613	394.5	1108.0
4	343	1.073	3.25	3.497	2.097	0.477	2.250	1.073	1594.5	3.23	1.073	2.421	368.0	1225.4
5	240	1.342	3.50	4.717	1.863	0.537	2.500	1.342	1564.2	3.46	1.342	3.369	322.1	1240.8
6	168	1.610	3.75	6.074	1.708	0.585	2.750	1.610	1455.8	3.73	1.610	4.454	270.5	1183.7
7	118	1.878	4.00	7.576	1.597	0.626	3.000	1.878	1310.2	3.93	1.878	5.682	221.6	1086.7
8	82	2.146	4.25	9.233	1.514	0.660	3.250	2.146	1134.8	4.43	2.146	7.061	176.0	956.6
9	58	2.415	4.50	11.058	1.449	0.690	3.500	2.415	979.6	4.71	2.415	8.601	140.1	837.1
10	40	2.683	4.75	13.071	1.398	0.715	3.750	2.683	810.8	5.02	2.683	10.319	107.3	700.7

N_T	3239				**All adults**				10353				2000	8339
N_A	1539				Exp	Obs								
				μ_k	1.30	1.30								
				σ^2_k	5.04	5.10								
				ϕ	3.88	3.92								
				$\bar{w}$	1.30	1.31								
				σ^2_w	3.73	3.87								
				CV^2_w	2.21	2.24								

Table 2. Example illustrating how to generate parental weights to model persistent individual differences in lifetime reproductive success, for a cohort of 490 females that survive to age at maturity (3). The first six columns repeat age-specific data from Table 1. The next three columns show calculation of adjustments to individual age-3 weights to generate weights for subsequent ages. These adjusted weights produce the same $CV^2_{w,x}$ for each age but maintain the age-3 pattern of relative individual fitness. λ_3 is the Poisson parameter that will generate weights with the desired properties for age-3 adults; $Q_x = TargetCV^2_{w,x}/TargetCV^2_{w,3}$; $C_x = \lambda_3 * (1/\sqrt{Q_x} - 1)$; $Adj(\overline{w}_x) = \lambda_3 + C_x$; Actual ϕ_x = means across 1000 replicate simulations.

Age	N_x	b_x	ϕ_x	Target $CV^2_{w,x}$	λ_3	Q_x	C_x	$Adj(\overline{w}_x)$	Actual ϕ_x
3	490	0.805	3.00	2.484	0.403	1.000	0.000	0.403	3.01
4	343	1.073	3.25	2.097	-	0.844	0.036	0.438	3.24
5	240	1.342	3.50	1.863	-	0.750	0.062	0.465	3.52
6	168	1.610	3.75	1.708	-	0.688	0.083	0.485	3.75
7	118	1.878	4.00	1.597	-	0.643	0.099	0.502	4.00
8	82	2.146	4.25	1.514	-	0.610	0.113	0.516	4.22
9	58	2.415	4.50	1.449	-	0.583	0.124	0.527	4.50
10	40	2.683	4.75	1.398	-	0.563	0.134	0.537	4.74

Figure 1. Flowcharts showing algorithms for the WF model (left) and *THE WEIGHT* (right), with differences highlighted in **bold red**.

Figure 2. Predicted effects of seven different parental weighting schemes on the ratio N_e/N based on Equation 5. For each individual i, $i=1,N$, the legend shows the formula for computing the relative parental weight w_i. Overflow issues in R prevent showing results for e^i when $i > 250$.

Figure 3. Distribution of lifetime reproductive success (*LRS*) for populations simulated using the vital rates in Table 2. Results are combined across 1000 replicates. In each replicate, *LRS* was computed for each of 490 age-3 females in a cohort. Open circles show results for simulations in which parental weights were randomly shuffled each year; red Xs show results for simulations in which individuals retained their weights throughout their reproductive lifespan. Numbers with arrows (49.1%; 36.4%) indicate the percentages of individuals in a cohort that produce zero offspring across their lifetime for persistent and independent simulations, respectively.