Unravelling microalgal-bacterial interactions in aquatic ecosystems through 16S rRNA

gene-based co-occurrence networks

B.L.D. Uthpala Pushpakumara[*1], Kshitij Tandon[1], Anusuya Willis[2], Heroen Verbruggen[*1]

[1]School of Biosciences, University of Melbourne, Victoria, 3010, Australia

[2]Australian National Algae Culture Collection, CSIRO, Tasmania 7000, Australia

*Corresponding   authors: e-mail: bpushpakumar@student.unimelb.edu.au,

heroen@unimelb.edu.au

Abstract

Interactions between microalgae and bacteria can directly influence the global biogeochemical

cycles but the majority of such interactions remain unknown. 16S rRNA gene-based co-

occurrence networks have potential to help identify microalgal-bacterial interactions. Here, we

used data from 10 Earth microbiome projects to identify potential microalgal-bacterial

associations in aquatic ecosystems. A high degree of clustering was observed in microalgal-

bacterial modules, indicating densely connected neighbourhoods. *Proteobacteria* and

Bacteroidetes predominantly co-occurred with microalgae and represented hubs of most

modules. Our results also indicated that species-specificity may be a global characteristic of

microalgal associated microbiomes. Several previously known associations were recovered

from our network modules, validating that biologically meaningful results can be inferred using

this approach. A range of previously unknown associations were recognised such as co-

occurrences of *Bacillariophyta* with uncultured *Planctomycetes OM190* and

Deltaproteobacteria order *NB1-j*. *Planctomycetes* and *Verrucomicrobia* were identified as key

associates of microalgae due to their frequent co-occurrences with several microalgal taxa.

Despite no clear taxonomic pattern, bacterial associates appeared functionally similar across

different environments. To summarise, we demonstrated the potential of 16S rRNA gene-based

co-occurrence networks as a hypothesis-generating framework to guide more focused research

on microalgal-bacterial associations.

Keywords: microalgae, bacteria, microalgal-bacterial associations, co-occurrence networks,

16S rRNA gene

Introduction

Microalgae (photosynthetic micro-eukaryotes) and bacteria are the most widespread and

dominant planktonic organisms in aquatic ecosystems. As primary producers, microalgae form

the foundations of the food web and directly influence global carbon and nutrient cycling, as

well as the energy flow in aquatic ecosystems. A wide spectrum of associations between

microalgae and bacteria have been reported, predominantly related to nutrient exchange, with

important bottom-up effects on primary production. Associations include increased

bioavailability of vitamins (e.g.B12[1–3]), metals (e.g. iron[4,5]) and growth promoting hormones[6]

by bacteria in exchange for organic carbon from the algae. Apart from many beneficial

interactions, some bacteria can have negative influences on algae by being algicidal and

opportunistic pathogens[7–9]. A recent study on the bloom-forming *Phaeocystis* algae revealed

that its microbiomes can take both symbiotic and opportunistic modes[10]. Thus, algal-bacterial

interactions can range from mutualistic to parasitic, and a complex array of interactions can be

hypothesised. For instance, previous work suggests that strong associations may exist between

communities of microalgae and bacteria over a large geographical range in the open ocean[11].

However, the knowledge we have so far about specific microalgal-bacterial interactions

represents only a small fraction of what potentially occurs in nature. Microalgae are highly

diverse and innumerable, therefore gathering knowledge on their associations with equally

ubiquitous and diverse bacteria is challenging.

High-throughput DNA sequencing such as 16S and 18S rRNA gene metabarcoding has been used to deliver insights into bacterial and eukaryotic community composition of diverse ecosystems. Microbial community abundance data can be used to identify associations between community members via microbial association networks[12]. Furthermore, combining community data with environmental data can reveal novel connections between microbial communities and their environment[13]. A network consists of nodes representing taxa and edges representing the associations between taxa. Microbial association networks targeting positive correlations are termed co-occurrence networks. A positive relationship is presumed when taxonomically relevant units co-occur or exhibit similarity in their compositions over multiple samples[14] and biologically meaningful groups or communities of a network can be identified with network clustering[15]. Also, network metrics such as node degree, closeness centrality for example, can be used for quantitative description of communities and identify ecologically important taxa[16,17]. Various network analyses have reported co-occurring microbial taxa in different environments, identified clusters of microbes representing metabolic consortia, defined keystone species, documented recurrent microbial modules and helped elucidate microbial dark matter in microbial communities[18–21]. Ecological processes governing community structure, such as niche filtering and habitat preference, are thought to be reflected in co-occurrence network modules[22]. Overall, networks provide insights into the structure of a community and taxon co-occurrences, offering perspectives to use them to generate hypotheses about interactions that can be investigated with more focused studies. Thus, environmental sequencing data combined with network analysis can be a convenient addition to the toolkit for deriving potential interactions between organisms across myriads of environments.

Studies targeting 16S rRNA marker genes are generally focused on the bacterial communities

and often disregard the chloroplast sequences that are also amplified. The shared origin of the

chloroplasts of all oxygenic photosynthetic micro-eukaryotes and cyanobacteria[23] enable the

use of 16S rRNA marker gene to concurrently estimate relative abundances of microalgae and

bacteria across samples, facilitating the construction of correlation networks to identify co-

occurring taxa and speculate on their potential interactions. Robust co-occurrences can be

detected through analysis of a sufficient number of samples, ideally covering temporal and

spatial gradients, as these provide adequate variability in taxon abundances[12]. This can be

coupled with stringent network building steps such as filtration of low prevalence organisms

to reveal significant co-occurrences[24]. The ever-expanding sequence databases can be a great

starting point to investigate such correlations. Analysis of the Tara Ocean dataset based on

interaction networks has shown that abiotic factors are incomplete predictors of community

structure[22]. Similar observations have been shown in phytoplankton blooms where

environmental parameters were insignificant in influencing the community structure of

plankton[25]. These results indicate that biological interactions are more influential in

determining the community structure than environmental factors. Thus, taxon-taxon co-

occurrence networks built on taxon compositions alone could capture potential interactions in

nature. Our work is based on the premise that previously unknown associations between

microalgae and bacteria may be discovered at large scales from existing 16S rRNA gene-based

metabarcoding datasets using co-occurrence networks. In this study, we generated taxon-taxon

co-occurrence networks using ten 16S rRNA gene metabarcoding datasets from the Earth

Microbiome Project (EMP) that have sampled aquatic environments (both marine and

freshwaters). These datasets were individually analysed to create local networks and network

modules were used to identify significant co-occurrences of microalgae and bacteria.

Methods

Data Acquisition

Publicly available EMP datasets[26] were screened using the Qiita portal[27] and 10 studies (4

marine and 6 Freshwater environments) targeting aquatic environments were chosen (Figure

S1). Since the goal of this study was to individually analyse samples representing a particular

environment to generate local networks and reveal co-occurrences, studies targeting any

aquatic samples (water and sediment samples) were selected. To our knowledge, based on

study and sample information provided on qiita portal for the selected studies, no size

fractionation was carried out on the samples (water samples) provided to EMP. The data

analysed in this study have amplified the same 16S rRNA gene V4 hypervariable region using

515F- 'GTGCCAGCMGCCGCGGTAA' and 806R- 'GGACTACHVGGGTWTCTAAT'

primer pair[28,29] and sequenced on Illumina Hiseq2000. Detailed information on the selected

EMP projects can be found in Supplementary material (Table S1) and can also be further

explored using Qiita ids provided using the Qiita portal.

Sequencing Data Workflow

Raw demultiplexed reads were downloaded using the EBI accessions (provided in Qiita portal)

from the European Nucleotide Archive and analysed individually employing a single

bioinformatic pipeline built on Qiime2 version 2019.10[30]. Primer sequences attached to all

reads were trimmed using cutadapt[31]. Sequence denoising, chimera checking and dereplication

was performed in DADA2 [32] to correct sequencing errors and remove low quality bases. Reads

were truncated based on a median quality score of 30. The final outputs of DADA2, an

abundance table of Amplicon Sequence Variants (ASVs) and fasta sequences of the ASVs were

further processed as described hereafter. Taxonomy was assigned against Silva v132 [30] 16S

rRNA gene sequences trained with a Naive Bayes classifier[33]. Chloroplast sequences were

filtered using the Silva taxonomy file by identifying sequences assigned as "chloroplast".

These chloroplast sequences were then classified again using a Qiime2-compatible version of

PhytoRef[34] database accessed at[35]. All the algal taxonomic identities with PhytoRef had a

confidence score of 0.7 or higher (Table S2 and S3). It is important to note that, each algal

node in the networks (see below for details) represents a single ASV, likely representing one

algal species or strain in the environment.

The DADA2 abundance table was filtered to create a bacteria-only abundance table (by

removing mitochondrial, Archaeal and chloroplast ASVs) and a chloroplast-only abundance

table (by retaining only the ASV ids assigned as chloroplast). The bacteria-only table was then

collapsed at the genus level to reduce weak links in downstream network building[36] and was

merged with the chloroplast-only table to create the final abundance table. ASVs identified as

microalgae were not collapsed since the 16S rRNA gene marker is highly conserved in algae

and ASVs likely correspond to species or higher-level taxa[34]. This final abundance table was

filtered to remove low prevalence organisms present in less than 10% of samples to prevent

them from introducing artefacts in network inference.

Correlation analysis and network construction

We used a compositional data analysis tool, FastSpar[37], which is a rapid and parallelizable

implementation of the SparCC algorithm[38] to compute correlations. FastSpar quantifies the

correlation between all ASVs and assesses the statistical significance (p-values) of the inferred

correlations using a bootstrap procedure. The correlation and p-value matrices created by

FastSpar were used to create a network of significant correlations for each study separately.

Co-occurrence networks were created using the igraph package[39] in R studio version 1.4.1106.

Undirected weighted networks were created using statistically ($p < 0.05$) significant

correlations > 0.5 or higher (0.5 for freshwater and 0.6 for marine datasets). Coefficient cut-

off of 0.5 was selected for freshwater datasets as a value above that resulted in fewer algal

nodes in networks compared to 0.5. The idea behind the coefficient cut-off selection was to

preserve as many links as possible with microalgae in resulting networks while choosing a

considerably higher coefficient cut off to report stronger links. Networks were visualised in

Cytoscape version 3.8[40].

Network Analysis and module detection

The Network Analyzer plugin[41] in Cytoscape was used to compute global network properties

of each network to get an overview of node specific and edge-specific attributes. Networks

were checked for the scale free nature by identifying the presence of highly connected nodes

coexisting with nodes with fewer links[42] using the node attributes. In order to identify

communities, networks were clustered using the Cytoscape plugin clusterMaker[43] using the

MCL clustering algorithm[44] with an inflation value of 2.0. Co-occurring microalgae and

bacteria were identified using resulting modules (modules with nodes > or = 4). These co-

occurring taxa (identified beyond the taxonomic rank Phylum) were recorded for each

environment and summarised in heatmaps using Pheatmap v1.0.12[45] in R.

Results and Discussion

16S rRNA gene-based co-occurrence networks can recover microalgal-bacterial

associations

We generated 10 co-occurrences networks representing aquatic environments using publicly

available 16S rRNA gene datasets from the EMP. We reduced noise and false positives by

including ASVs present in at least 10% of samples, filtering statistically insignificant and

weaker correlations. We also used network modules that are indicative of ecological

communities to identify potential interactions. In each co-occurrence network generated, we

observed highly connected nodes coexisting with nodes with fewer links. This indicates the

scale-free nature of the networks, a characteristic in real world networks[46,47]. The clustered

networks (modules) comprised nodes representing microalgae or bacteria and the edges

between them were instances of significant co-occurrences.

Analysis of the microalgal-bacterial modules identified 40 algal nodes co-occurring with at

least one of 76 bacterial nodes in marine environments and 112 microalgal nodes with at least

one of 311 bacterial nodes in freshwater environments (refer Table S2 and S3 for recovered

co-occurrences with the correlation values). These significant co-occurrences inferred in

marine and freshwater environments are summarised in Figures 1 and 2, respectively. We

identified algal nodes at different taxonomic levels although many could not be classified at

lower taxonomic levels such as genus or species. To have consistency in algal node

annotations, we have only used class-level classification in figures. Tables S2 and S3 provide

full taxonomic affiliations of the nodes where possible. Even though taxonomic assignments

were made at higher rank, our ASVs represent taxa at lower taxonomic levels. As the breadth

of species included in reference databases such as PhytoRef increases, it is likely that more

algal ASVs can be assigned to the species-level.

Most modules exhibited high clustering coefficients (> 0.5) indicating that the neighbourhood

of microalgal-bacterial communities are generally densely connected. High clustering

coefficient values in networks were previously suggested as indicative of cross-feeding

relationships and enriched degradation pathways[48]. We identified potential interactions of

bacteria with diverse phyla such as *Cryptophyta*, *Ochrophyta*, *Haptophyta*, *Chlorophyta,*

Streptophyta and *Euglenophta* representing 17 different taxonomic classes. Bacterial nodes in

the marine modules were predominantly represented by *Proteobacteria* and *Bacteroidetes*

followed by *Planctomycetes and Verrucomicrobia.* Similar to marine environments

Proteobacteria and Bacteroidetes dominated the freshwater modules while *Actinobacteria*

and *Verrucomicrobia* were the 2nd and 3rd most common bacterial taxa associated with

microalgae (Figure S2). We also identified hub nodes with the highest node degree (number

of edges connected to a node). Hub nodes represent highly connected nodes and are usually

considered as keystone species[49]. Hub nodes (considering the top ten hubs) in both marine

and freshwater modules were mostly represented by *Alphaproteobacteria*,

Gammaproteobacteria (p_*Proteobacteria*) and *Bacteroidia* (p_*Bacteroidetes*). Other than

these, members of *Planctomycetes* (mostly belonging to c_*Planctomycetacia*) were

commonly found among the top hub nodes in freshwater modules. High prevalence and their

characteristic associations with microalgae may explain the presence of *Proteobacteria* and

Bacteroidetes as hub nodes in most modules[50,51]. The role of *Planctomycetes* in global

nitrogen, carbon and sulphur cycles is gaining attention[52,53]. Thus, microalgal-*Planctomycetes*

associations may be playing a crucial role in the global environmental cycles.

Based on the summarised significant co-occurrences, benthic diatoms exhibited different, and

fewer, associations compared to planktonic diatoms (Figures 1 and 2). For example, bacterial

taxa such as unclassified *Gammaproteobacteria*, *Caldilineaceae* (*Chloroflexi*), *Trichococcus*

(*Firmicutes*), *Hoeflea* (*Proteobacteria*), *Pseudorhodobacter* (*Proteobacteria*) and

Cocleimonas (*Proteobacteria)* exhibited associations only with marine benthic diatoms and

not with marine planktonic diatoms.

We investigated whether this network approach has recovered any known associations

identified and confirmed earlier. We recognized known bacterial associates of microalgae

such as *Flavobacteriales*[54,55], *Rhizobiales*[56,57], *Cytophagales*[58,59] and *Rhodobacterales*[60].

Previously reported associations of microalgal genera with specific bacterial groups were also

recovered in the analysis, including that between *Bacteroidetes* and the cosmopolitan

Prymnesiophyceae genus *Phaeocystis*[61–63] (Table S2) and that of diatoms (*Bacillariophyta*)

with the genus *Flavobacterium* (Table S3)[64–66]. The ability to identify previously known

associations suggests that the 16S rRNA gene-based network approach used in this study can

yield biologically meaningful results.

We also compared co-occurrence networks built in our study with previously published

networks of these data. Previous analysis of lake Mendota samples (EBI accession:

ERP016591) by Kara et al[67] described characteristics of bacterioplankton co-occurrence

networks across three seasons. The network properties of these bacterioplankton networks such

as clustering coefficient (0.167 - 0.256) and the characteristic path length (3.27 - 3.88) were

comparatively lower than those of our microalgal-bacterial network (clustering coefficient >

0.5, characteristic path length = 3.965). Similar to the observation by Gilbert et al[68], the co-

occurrence network built on Western English Channel samples (EBI accession: ERP016541)

featured many *Alphaproteobacteria and Rhodobacterales* nodes.

Both Intrinsic and Extrinsic factors may influence microalgal-bacterial associations

No clear taxonomic pattern was observed among the co-occurring microalgae and bacteria. In

accordance with previous work on phytoplankton-bacteria co-occurrences[67], taxonomically

diverse bacteria co-occurred with different microalgal groups. Moreover, except for a few,

there was no recurrence of bacterial genera (the lowest taxonomic identity of bacterial nodes

in a module) across modules generated in each project. Since each project represents a

specific environment, these observations indicate that each environment harbours its specific

co-occurrence relationships. Project specificity in inferred co-occurrences can clearly be seen

in the generated heatmaps (Figures 1 and 2). Interestingly, a recent study[68] has provided

evidence for biogeographic differentiation of algal microbiomes, showing ecological

boundaries driven by differences in environmental conditions altering the spatial scaling of

the algal microbiomes. Another plausible explanation for such observations is the species

specificity of algal microbiomes. Although some microalgal nodes could not be

taxonomically identified at species level, each microalgal node in a module is an ASV which

likely represents a single microalgal species or a strain. Species specificity of algal

microbiomes has been predominantly shown using algal cultures[50]. In addition to supporting

previous observations, our results show that species-specificity may be a global characteristic

of microalgal microbiomes.

To understand if there is any factor driving general patterns of interactions, some insights may

be gleaned from known functions of the co-occurring bacteria. Our observation was that despite

their taxonomic differences, bacterial associates in modules often shared functional

similarities. For instance, microalgae co-occurred with bacterial groups equipped with specific

metabolic functional potentials such as algal polysaccharide degradation and provision of

vitamins. A few examples of these are, co-occurrences with taxonomically diverse members

of *Bacteroidetes*[63] and *Verrucomicrobia*[69] with polysaccharide degradation ability and

Rhizobiales, *Rhodobacterales* [70] and *SAR116* [71–73] with Vitamin B12 synthesis. Therefore, our

results stemming from observations across multiple datasets suggest that, irrespective of the

environment, microalgae are associated with key functional types of bacteria. As most

microalgal-bacterial functional interactions remain unknown, it is difficult to identify global

trends in key functional types of bacteria associated with microalgal-bacterial communities.

This urges the need for more functional studies to improve our understanding of microalgal-

bacterial communities across the globe.

Emerging microalgal-bacterial associations that can guide functional studies

An advantage of using a network approach is the ability to analyse large scale datasets to

unravel previously unknown associations. We believe that some of the inferred co-occurrences

in this study may help guide focused research to shed light on the functional nature of

interactions. Therefore, we further explored these co-occurrences which were prominent due

to their frequent observations and importance in modules based on topological properties.

For instance, uncultured *Deltaproteobacteria* order *NB1-j* was identified as consistently co-

occurring with *Bacillariophyta* in marine environments. In one environment where the *NB1-j-*

Bacillariophyta link was observed (Surface water samples from Western English Channel, EBI

accession: ERP016541), *Bacillariophyta* was the only microalgal node directly interacting with

NB1-j. In another environment (Seawater metagenome samples from Catlin Arctic survey

(2010 expedition), EBI accession: ERP020022), *NB1-j* was identified as the hub node with the

highest node degree (43), in a module consisting of 77 nodes and 817 edges. This *NB1-j* node

also had the highest closeness centrality (0.69). Closeness centrality measures how close a node

is to another node and helps to identify centrally positioned taxa in the network[24]. Other than

this *NB1-j* node, there were 3 more *NB1-j* nodes in this community. All together these 4 *NB1-*

j nodes were directly interacting (edges) with 57 nodes (out of 73 other nodes) of which 25

were representing microalgal nodes (Figure 3). Out of the 25 microalgal nodes, 13 represented

the taxonomic class *Bacillariophyta*. Some of these *Bacillariophyta* were taxonomically

identified at genus level as *Chaetoceros* and *Fragilariopsis*. Interestingly, except for one, all

these Bacillariophyta nodes represented top hub nodes of the community with node degrees >

26. Most connected taxa are believed to have ecological relevance to the community as their

removal causes the highest impact on many associations[74].

Functional roles of *NB1-j* in marine ecosystems are largely unknown. It is believed to be

involved in hydrocarbon degradation[75] and has been reported mostly from marine sediments

[70,76], mud volcanoes[75], sponges[77] and cyanobacterial mats[78]. To the best of our knowledge, this

Deltaproteobacteria group is not commonly reported in algal microbiomes. However, a recent

study found *NB1-j* ASVs associated with the coral skeleton algal symbiont *Ostreobium*[79]. A

predictive metagenomic approach based on sponge samples originating from reef sites in West

Java (Indonesia) suggest that *NB1-j* may be involved in nitrogen metabolism[77]. It was found

that *NB1-j* was responsible for the elevated predictive gene count corresponding to N-cycling

genes such as those encoding nitric oxide reductase (*norB)*, nitrogenase (*nifD*) and

hydroxylamine reductase. The available information indicates that the nitrogen cycling

capacity of *NB1-j* may underly its association with algae, potentially facilitating the nitrogen

needs of the algae while benefiting from algal organic carbon. However, a repertoire of

interactions, as often expected from a keystone species, can be hypothesised between *NB1-j*

and microalgae in a community.

Bacillariophyta (*Fragilariopsis, Chaetoceros* and many other diatoms unidentified at lower

taxonomic levels) also showed frequent co-occurrences with an uncultured clade of

Planctomycetes, *OM190* (SILVA taxonomy) in both freshwater and marine environments. In

the Catlin Arctic survey samples (ERP02022), *OM190* demonstrated frequent associations with

Bacillariophyta. Three *OM190* nodes were observed with high node degrees (29, 30 and 39)

and these were directly connected to 59 nodes out of a total of 74 other nodes in the community.

From the total of 35 microalgal nodes in this community, *OM190* were directly connected to

27 nodes out of which 15 nodes were represented by *Bacillariophyta* (Figure S3A). Similarly

in a freshwater environment (Lake Superior, Michigan, EBI accession: ERP016492), *OM190*

was identified as the 6th most connected node with a node degree of 89 and was directly

connected to 32 microalgal nodes including *Bacillariophyta* (Figure S3B). As mentioned

previously, highly connected nodes act as hub nodes in the community. Multiple *OM190* acting

as top hub nodes within their habitats indicate that they have high co-occurrences with both

microalgae and bacteria.

OM190 shows deep branching within the *Planctomycetes* group and is usually found in

different environments such as soil and seawater. Members of this clade are usually considered

to be associated with macroalgae[80], such as red[81] and brown algae[82]. Since *OM190* is yet

uncultured, information on its metabolism is scarce. A metagenome-assembled-genome

(MAG) for *OM190* (likely *OM190*) with a rich diversity of secondary metabolite potential has

been reported[83]. The production of secondary metabolites by *OM190*, including antimicrobial

compounds may be one of the underlying reasons for their association with algae as they could

protect the alga from undesired microbes. In a recent study[84], it was shown that diatoms

produce fucose-containing sulfated polysaccharides (FCSP) which can be hypothesised as an

energy source for *OM190*. Many *Planctomycetes* have abundant sulfatases, but these were not

confirmed in the existing *OM190* MAG. In addition to their association with algae, *OM190*

and *NB1-j* seem to have a close relationship with one another. They were recently reported as

abundant co-occurring taxa in Beaufort Sea surface sediments[85]. The reasons for their direct

associations are not known and, along with their respective relationships with algae, this direct

interaction is an interesting topic for future investigation.

More generally speaking, *Planctomycetes* often associate with macroalgae and favour a biofilm

lifestyle[82]. Protein clusters that may be involved in *Planctomycetes* symbiosis or biofilm

maintenance have been reported[53], and production of sulfated polysaccharides by the alga that

serve as the substrate for the abundant sulfatases produced by the *Planctomycetes* contributes to the reasons for successful associations between them[53,80]. Although successful associations are known between macroalgae and *Planctomycetes*, interactions with microalgae are largely unknown. Apart from their associations with diatoms, our network analysis identified taxa representing *Planctomycetes* (c_*Phycisphaerae*, c_*Planctomycetacia*) co-occurring with a range of microalgal groups. In a previous study, *Planctomycetes* closely related to *Pirellula* were identified as one of the dominant lineages associated with diatom blooms[86]. Results of our network analysis demonstrate that associations between *Planctomycetes* and microalgae may be as common as those they maintain with macroalgae.

Verrucomicrobia were also consistently associated with an array of microalgae in fresh and marine environments indicating that they may be common associates of microalgae. Here in our study, all the Verrucomicrobial members co-occurring with microalgae were represented by the taxonomic class *Verrucomicrobiae*. Among the genus-specific associations revealed in our network analysis is that of the Verrucomicrobial genus *Lentimonas* and microalgal genera *Pyramimonas* (c_*Prasinophyceae*), *Phaeocystis*, *Tisochrysis*, *Haptolina*, and *Chrysochromulina* (c_*Prymnesiophyceae*). A recent study showed hundreds of *Lentimonas* enzymes able to digest brown algal fucoidan[87]. Although fucose-containing sulphated polysaccharides were regarded as a macroalgal polysaccharide, microalgae such as diatoms have the potential to produce them[84]. We speculate that other microalgae might also produce this sulfated polysaccharide, or that members of *Lentimonas* may have a broader palate than just FCSP. Although widely distributed, the functional roles of *Verrucomicrobia* in aquatic ecosystems are not well understood due to the lack of cultured strains[88]. Some members of *Verrucomicrobia* have been shown to consume algal extracellular polymeric substances[69,89,90]. However, other factors contributing to their associations are not well understood.

Conclusion

In summary, our study illustrates the promise of using 16S rRNA gene-based co-occurrence networks as a hypothesis-generating framework to guide focused research and speculate on the functional nature of potential interactions. By studying multiple environments based on public datasets, we provided an overview of microalgal-bacterial communities in aquatic ecosystems from a network perspective. This identified a range of associations including previously unknown links that can set the stage for more focused research in the future.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

No data was generated in this study and all the data analysed are publicly available. The EMP datasets analysed are available in European Nucleotide Archive and EMP Qiita portal under the Accession numbers and Qiita ids, respectively, provided in Supplementary material TableS1. FastSpar script used for the correlation analysis, R script used in generating networks and the networks files generated in Cytoscape can be found at

https://melbourne.figshare.com/projects/Unravelling_microalgal-

bacterial_interactions_in_aquatic_ecosystems_through_16S_rRNA_gene-based_co-

occurrence_networks/140071

Competing interests

The authors declare that they have no competing interests.

Funding

We acknowledge the funding from the Melbourne Research Scholarship and the

ResearchPlus Postgraduate Top-Up Scholarship Grants Program and National Research

Collections Australia, CSIRO.

Authors' contribution

UP, HV and AW formulated the project. UP analysed the data and wrote the manuscript with

inputs from all the authors. HV, AW and KT contributed ideas and helped with analysis. All

authors read and approved the final version of the manuscript.

Acknowledgements

Not applicable

References

1. Croft, M. T., Lawrence, A. D., Raux-Deery, E., Warren, M. J. & Smith, A. G. Algae acquire vitamin B12 through a symbiotic relationship with bacteria. *Nature* (2005) doi:10.1038/nature04056.

2. Kazamia, E. *et al.* Mutualistic interactions between vitamin B12-dependent algae and heterotrophic bacteria exhibit regulation. *Environ. Microbiol.* (2012) doi:10.1111/j.1462-2920.2012.02733.x.

3. Bunbury, F. *et al.* Exploring the onset of B12-based mutualisms using a recently evolved Chlamydomonas auxotroph and B12-producing bacteria. *Environ. Microbiol.* **24**, 3134–3147 (2022).

4. Butler, A. Acquisition and utilization of transition metal ions by marine organisms. *Science* (1998) doi:10.1126/science.281.5374.207.

5.   Amin, S. A. *et al.* Photolysis of iron-siderophore chelates promotes bacterial-algal

mutualism. *Proc. Natl. Acad. Sci. U. S. A.* (2009) doi:10.1073/pnas.0905512106.

6.   Amin, S. A. *et al.* Interaction and signalling between a cosmopolitan phytoplankton

and associated bacteria. *Nature* (2015) doi:10.1038/nature14488.

7.   Mayali, X. & Azam, F. Algicidal bacteria in the sea and their impact on algal blooms.

in *Journal of Eukaryotic Microbiology* (2004). doi:10.1111/j.1550-

7408.2004.tb00538.x.

8.   Bagwell, C. E. *et al.* Discovery of bioactive metabolites in biofuel microalgae that

offer protection against predatory bacteria. *Front. Microbiol.* (2016)

doi:10.3389/fmicb.2016.00516.

9.   Hoeger, A. L., Jehmlich, N., Kipping, L., Griehl, C. & Noll, M. Associated bacterial

microbiome responds opportunistic once algal host Scenedesmus vacuolatus is

attacked by endoparasite Amoeboaphelidium protococcarum. *Sci. Rep.* 1–11 (2022)

doi:10.1038/s41598-022-17114-1.

10.   Mars Brisbin, M., Mitarai, S., Saito, M. A. & Alexander, H. Microbiomes of bloom-

forming Phaeocystis algae are stable and consistently recruited, with both symbiotic

and opportunistic modes. *ISME J. 2022* 1–10 (2022) doi:10.1038/s41396-022-01263-

2.

11.   Milici, M. *et al.* Co-occurrence analysis of microbial taxa in the Atlantic ocean reveals

high connectivity in the free-living bacterioplankton. *Front. Microbiol.* **7**, 1–20 (2016).

12.   Barberán, A., Bates, S. T., Casamayor, E. O. & Fierer, N. Using network analysis to

explore co-occurrence patterns in soil microbial communities. *ISME J.* **6**, 343–351

(2012).

13.   Tucker, A. E. & Brown, S. P. Sampling a gradient of red snow algae bloom density

445   reveals novel connections between microbial communities and environmental features.

446   *Sci. Rep.* **12**, 1–15 (2022).

447   14.   Faust, K. & Raes, J. Microbial interactions: From networks to models. *Nat. Rev.*

448   *Microbiol.* **10**, 538–550 (2012).

449   15.   Chun, S. J. *et al.* Network analysis reveals succession of Microcystis genotypes

450   accompanying distinctive microbial modules with recurrent patterns. *Water Res.*

451   (2020) doi:10.1016/j.watres.2019.115326.

452   16.   Huang, S. Back to the biology in systems biology: What can we learn from

453   biomolecular networks? *Briefings Funct. Genomics Proteomics* (2004)

454   doi:10.1093/bfgp/2.4.279.

455   17.   Ma'ayan, A. Introduction to network analysis in systems biology. *Science Signaling*

456   (2011) doi:10.1126/scisignal.2001965.

457   18.   Williams, R. J., Howe, A. & Hofmockel, K. S. Demonstrating microbial co-occurrence

458   pattern analyses within and between ecosystems. *Front. Microbiol.* **5**, 1–10 (2014).

459   19.   Fuhrman, J. A., Cram, J. A. & Needham, D. M. Marine microbial community

460   dynamics and their ecological interpretation. *Nature Reviews Microbiology* (2015)

461   doi:10.1038/nrmicro3417.

462   20.   Chafee, M. *et al.* Recurrent patterns of microdiversity in a temperate coastal marine

463   environment. *ISME J.* **12**, 237–252 (2018).

464   21.   Zamkovaya, T., Foster, J. S., de Crécy-Lagard, V. & Conesa, A. A network approach

465   to elucidate and prioritize microbial dark matter in microbial communities. *ISME J.* **15**,

466   228–244 (2021).

22.   Lima-Mendez, G. *et al.* Determinants of community structure in the global plankton

interactome. *Science (80-.).* (2015) doi:10.1126/science.1262073.

23.   Bennke, C. M. *et al.* The distribution of phytoplankton in the Baltic Sea assessed by a

prokaryotic 16S rRNA gene primer system. *J. Plankton Res.* **40**, 244–254 (2018).

24.   Berry, D. & Widder, S. Deciphering microbial interactions and detecting keystone

species with co-occurrence networks. *Front. Microbiol.* (2014)

doi:10.3389/fmicb.2014.00219.

25.   Needham, D. M. & Fuhrman, J. A. Pronounced daily succession of phytoplankton,

archaea and bacteria following a spring bloom. *Nat. Microbiol.* (2016)

doi:10.1038/nmicrobiol.2016.5.

26.   Thompson, L. R. *et al.* A communal catalogue reveals Earth's multiscale microbial

diversity. *Nature* (2017) doi:10.1038/nature24621.

27.   Gonzalez, A. *et al.* Qiita: rapid, web-enabled microbiome meta-analysis. *Nat. Methods*

(2018) doi:10.1038/s41592-018-0141-9.

28.   Caporaso, J. G. *et al.* Global patterns of 16S rRNA diversity at a depth of millions of

sequences per sample. *Proc. Natl. Acad. Sci. U. S. A.* (2011)

doi:10.1073/pnas.1000080107.

29.   Caporaso, J. G. *et al.* Ultra-high-throughput microbial community analysis on the

Illumina HiSeq and MiSeq platforms. *ISME J.* (2012) doi:10.1038/ismej.2012.8.

30.   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data

science using QIIME 2. *Nature Biotechnology* (2019) doi:10.1038/s41587-019-0209-

9.

31.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal* (2011) doi:10.14806/ej.17.1.200.

32. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina

amplicon data. *Nat. Methods* (2016) doi:10.1038/nmeth.3869.

33. Bokulich, N. A. *et al.* Optimizing taxonomic classification of marker-gene amplicon

sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* (2018)

doi:10.1186/s40168-018-0470-z.

34. Decelle, J. *et al.* PhytoREF: A reference database of the plastidial 16S rRNA gene of

photosynthetic eukaryotes with curated taxonomy. *Mol. Ecol. Resour.* **15**, 1435–1445

(2015).

35. Major oceanic 16S/18S databases in qiime2 format. https://github.com/ndu-

invitae/Oceanic_database/tree/master/PhytoRef.

36. Hemprich-Bennett, D. R., Oliveira, H. F. M., Le Comber, S. C., Rossiter, S. J. &

Clare, E. L. Assessing the impact of taxon resolution on network structure. *Ecology*

(2021) doi:10.1002/ecy.3256.

37. Watts, S. C., Ritchie, S. C., Inouye, M. & Holt, K. E. FastSpar: Rapid and scalable

correlation estimation for compositional data. *Bioinformatics* (2019)

doi:10.1093/bioinformatics/bty734.

38. Friedman, J. & Alm, E. J. Inferring Correlation Networks from Genomic Survey Data.

PLoS Comput. Biol. (2012) doi:10.1371/journal.pcbi.1002687.

39. Csardi, G., & Nepusz, T. The igraph software package for complex network research.

InterJournal, Complex Systems. *igraph Softw. Packag.* (2006).

40. Shannon, P. *et al.* Cytoscape: A software Environment for integrated models of

biomolecular interaction networks. *Genome Res.* (2003) doi:10.1101/gr.1239303.

41.   Assenov, Y., Ramírez, F., Schelhorn, S. E. S. E., Lengauer, T. & Albrecht, M.

514   Computing topological parameters of biological networks. *Bioinformatics* (2008)

515   doi:10.1093/bioinformatics/btm554.

516   42.   Barabási, A. L. Scale-free networks: A decade and beyond. *Science* (2009)

517   doi:10.1126/science.1173299.

518   43.   Morris, J. H. *et al.* ClusterMaker: A multi-algorithm clustering plugin for Cytoscape.

519   *BMC Bioinformatics* (2011) doi:10.1186/1471-2105-12-436.

520   44.   Van Dongen, S. & Abreu-Goodger, C. Using MCL to extract clusters from networks.

521   *Methods Mol. Biol.* (2012) doi:10.1007/978-1-61779-361-5_15.

522   45.   Raivo, K. Pheatmap: pretty heatmaps. *R Pacakage Version* (2012).

523   46.   Albert, R., Jeong, H. & Barabási, A. L. Diameter of the world-wide web. *Nature*

524   (1999) doi:10.1038/43601.

525   47.   Eisenberg, E. & Levanon, E. Y. Preferential attachment in the protein network

526   evolution. *Phys. Rev. Lett.* (2003) doi:10.1103/PhysRevLett.91.138701.

527   48.   Ma, B. *et al.* Earth microbial co-occurrence network reveals interconnection pattern

528   across microbiomes. *Microbiome* **8**, 82 (2020).

529   49.   Wang, H. *et al.* Combined use of network inference tools identifies ecologically

530   meaningful bacterial associations in a paddy soil. *Soil Biol. Biochem.* **105**, 227–235

531   (2017).

532   50.   Goecke, F., Thiel, V., Wiese, J., Labes, A. & Imhoff, J. F. Algae as an important

533   environment for bacteria - Phylogenetic relationships among new bacterial species

534   isolated from algae. *Phycologia* (2013) doi:10.2216/12-24.1.

51.   Krohn-Molt, I. *et al.* Metagenome survey of a multispecies and alga-associated biofilm

revealed key elements of bacterial-algal interactions in photobioreactors. *Appl.*

Environ. Microbiol. (2013) doi:10.1128/AEM.01641-13.

52.   Woebken, D. *et al.* Fosmids of novel marine Planctomycetes from the Namibian and

Oregon coast upwelling systems and their cross-comparison with planctomycete

genomes. *ISME J.* (2007) doi:10.1038/ismej.2007.63.

53.   Faria, M. *et al.* Planctomycetes attached to algal surfaces: Insight into their genomes.

Genomics (2018) doi:10.1016/j.ygeno.2017.10.007.

54.   Barbeyron, T., L'Haridon, S., Corre, E., Kloareg, B. & Potin, P. Zobellia

galactanovorans gen. nov., sp. nov., a marine species of Flavobacteriaceae isolated

from a red alga, and classification of [Cytophaga] uliginosa (ZoBell and Upham 1944)

Reichenbach 1989 as Zobellia uliginosa gen. nov., comb. nov. *Int. J. Syst. Evol.*

Microbiol. (2001) doi:10.1099/00207713-51-3-985.

55.   Nedashkovskaya, O. I. *et al.* Mesonia algae gen. nov., sp. nov., a novel marine

bacterium of the family Flavobacteriaceae isolated from the green alga Acrosiphonia

sonderi (Kütz) Kornm. *Int. J. Syst. Evol. Microbiol.* (2003) doi:10.1099/ijs.0.02626-0.

56.   Kim, B. H., Ramanan, R., Cho, D. H., Oh, H. M. & Kim, H. S. Role of Rhizobium, a

plant growth promoting bacterium, in enhancing algal biomass through mutualistic

interaction. *Biomass and Bioenergy* (2014) doi:10.1016/j.biombioe.2014.07.015.

57.   Rivas, M. O., Vargas, P. & Riquelme, C. E. Interactions of Botryococcus braunii

Cultures with Bacterial Biofilms. *Microb. Ecol.* (2010) doi:10.1007/s00248-010-9686-

6.

58.   Cole, J. J. Interactions between bacteria and algae in aquatic ecosystems. *Annu. Rev.*

Ecol. Syst. Vol. 13 (1982) doi:10.1146/annurev.es.13.110182.001451.

59.   Fulbright, S. P. *et al.* Bacterial community changes in an industrial algae production

560   system. *Algal Res.* (2018) doi:10.1016/j.algal.2017.09.010.

561   60.   Zhou, J. *et al.* Microbial community structure and associations during a marine

562   dinoflagellate bloom. *Front. Microbiol.* (2018) doi:10.3389/fmicb.2018.01201.

563   61.   Simon, M., Glöckner, F. O. & Amann, R. Different community structure and

564   temperature optima of heterotrophic picoplankton in various regions of the Southern

565   Ocean. *Aquat. Microb. Ecol.* (1999) doi:10.3354/ame018275.

566   62.   Brussaard, C. P. D., Mari, X., Van Bleijswijk, J. D. L. & Veldhuis, M. J. W. A

567   mesocosm study of Phaeocystis globosa (Prymnesiophyceae) population dynamics: II.

568   Significance for the microbial community. *Harmful Algae* (2005)

569   doi:10.1016/j.hal.2004.12.012.

570   63.   Janse, I., Zwart, G., Van der Maarel, M. J. E. C. & Gottschal, J. C. Composition of the

571   bacterial community degrading Phaeocystis mucopolysaccharides in enrichment

572   cultures. *Aquat. Microb. Ecol.* (2000) doi:10.3354/ame022119.

573   64.   Grossart, H. P., Levold, F., Allgaier, M., Simon, M. & Brinkhoff, T. Marine diatom

574   species harbour distinct bacterial communities. *Environ. Microbiol.* (2005)

575   doi:10.1111/j.1462-2920.2005.00759.x.

576   65.   Sapp, M. *et al.* Species-specific bacterial communities in the phycosphere of

577   microalgae? *Microb. Ecol.* (2007) doi:10.1007/s00248-006-9162-5.

578   66.   Sapp, M., Wichels, A. & Gerdts, G. Impacts of cultivation of marine diatoms on the

579   associated bacterial community. *Appl. Environ. Microbiol.* (2007)

580   doi:10.1128/AEM.02274-06.

581   67.   Milici, M. *et al.* Bacterioplankton biogeography of the Atlantic ocean: A case study of

582   the distance-decay relationship. *Front. Microbiol.* **7**, 1–15 (2016).

583   68.   Martin, K. *et al.* The biogeographic differentiation of algal microbiomes in the upper

584   ocean from pole to pole. *Nat. Commun.* (2021) doi:10.1038/s41467-021-25646-9.

585   69.   Martinez-Garcia, M. *et al.* Capturing single cell genomes of active polysaccharide

586   degraders: An unexpected contribution of verrucomicrobia. *PLoS One* (2012)

587   doi:10.1371/journal.pone.0035314.

588   70.   Dell'Anno, F. *et al.* Highly Contaminated Marine Sediments Can Host Rare Bacterial

589   Taxa Potentially Useful for Bioremediation. *Front. Microbiol.* (2021)

590   doi:10.3389/fmicb.2021.584850.

591   71.   Newton, R. J. *et al.* Genome characteristics of a generalist marine bacterial lineage.

592   *ISME J.* (2010) doi:10.1038/ismej.2009.150.

593   72.   Sañudo-Wilhelmy, S. A., Gómez-Consarnau, L., Suffridge, C. & Webb, E. A. The role

594   of B vitamins in marine biogeochemistry. *Ann. Rev. Mar. Sci.* (2014)

595   doi:10.1146/annurev-marine-120710-100912.

596   73.   Durham, B. P. *et al.* Cryptic carbon and sulfur cycling between surface ocean

597   plankton. *Proc. Natl. Acad. Sci. U. S. A.* (2015) doi:10.1073/pnas.1413137112.

598   74.   Proulx, S. R., Promislow, D. E. L. & Phillips, P. C. Network thinking in ecology and

599   evolution. *Trends in Ecology and Evolution* (2005) doi:10.1016/j.tree.2005.04.004.

600   75.   Coelho, F. J. R. C. *et al.* Integrated analysis of bacterial and microeukaryotic

601   communities from differentially active mud volcanoes in the Gulf of Cadiz. *Sci. Rep.*

602   **6**, 1–10 (2016).

76.   Queiroz, L. L. *et al.* Bacterial diversity in deep-sea sediments under influence of

asphalt seep at the São Paulo Plateau. *Antonie van Leeuwenhoek, Int. J. Gen. Mol.*

Microbiol. (2020) doi:10.1007/s10482-020-01384-8.

77. de Voogd, N. J., Cleary, D. F. R., Polónia, A. R. M. & Gomes, N. C. M. Bacterial

community composition and predicted functional ecology of sponges, sediment and

seawater from the thousand islands reef complex, West Java, Indonesia. *FEMS*

Microbiol. Ecol. (2015) doi:10.1093/femsec/fiv019.

78. Vigneron, A. *et al.* Multiple strategies for light-harvesting, photoprotection, and

carbon flow in high latitude microbial mats. *Front. Microbiol.* (2018)

doi:10.3389/fmicb.2018.02881.

79. Pushpakumara, B. L. D. U., Tandon, K., Willis, A., Verbruggen, H. The bacterial

microbiome of the coral skeleton algal symbiont Ostreobium shows preferential

associations and signatures of phylosymbiosis. *bioRxiv* (2022)

10.1101/2022.12.13.520198.

80. Lage, O. M. & Bondoso, J. Planctomycetes diversity associated with macroalgae.

FEMS Microbiol. Ecol. (2011) doi:10.1111/j.1574-6941.2011.01168.x.

81. Longford, S. R. *et al.* Comparisons of diversity of bacterial communities associated

with three sessile marine eukaryotes. *Aquat. Microb. Ecol.* (2007)

doi:10.3354/ame048217.

82. Bengtsson, M. M. & Øvreås, L. Planctomycetes dominate biofilms on surfaces of the

kelp Laminaria hyperborea. *BMC Microbiol.* (2010) doi:10.1186/1471-2180-10-261.

83. Ludington, W. B. *et al.* Assessing biosynthetic potential of agricultural groundwater

through metagenomic sequencing: A diverse anammox community dominates nitrate-

rich groundwater. *PLoS One* (2017) doi:10.1371/journal.pone.0174930.

84.   Vidal-Melgosa, S. *et al.* Diatom fucan polysaccharide precipitates carbon during algal

628   blooms. *Nat. Commun.* **12**, 1–13 (2021).

629   85.   Walker, A. M., Leigh, M. B. & Mincks, S. L. Patterns in Benthic Microbial

630   Community Structure Across Environmental Gradients in the Beaufort Sea Shelf and

631   Slope. *Front. Microbiol.* (2021) doi:10.3389/fmicb.2021.581124.

632   86.   Morris, R. M., Longnecker, K. & Giovannoni, S. J. Pirellula and OM43 are among the

633   dominant lineages identified in an Oregon coast diatom bloom. *Environ. Microbiol.*

634   (2006) doi:10.1111/j.1462-2920.2006.01029.x.

635   87.   Sichert, A. *et al.* Verrucomicrobia use hundreds of enzymes to digest the algal

636   polysaccharide fucoidan. *Nat. Microbiol.* (2020) doi:10.1038/s41564-020-0720-2.

637   88.   Bohórquez, J. *et al.* Different types of diatom-derived extracellular polymeric

638   substances drive changes in heterotrophic bacterial communities from intertidal

639   sediments. *Front. Microbiol.* (2017) doi:10.3389/fmicb.2017.00245.

640   89.   Landa, M. *et al.* Phylogenetic and structural response of heterotrophic bacteria to

641   dissolved organic matter of different chemical composition in a continuous culture

642   study. *Environ. Microbiol.* (2014) doi:10.1111/1462-2920.12242.

643   90.   Orsi, W. D. *et al.* Diverse, uncultivated bacteria and archaea underlying the cycling of

644   dissolved protein in the ocean. *ISME J.* (2016) doi:10.1038/ismej.2016.20.

Figure legends

Figure 1: Summarised interactions in marine environments

Correlation heatmap of co-occurring microalgae (columns) and bacteria (rows) in marine

environments. "Taxonomy" represents the taxonomic affiliations of microalgal nodes at class

level. "Taxonomy2" represents the taxonomic affiliations of bacteria (p_, c_, o_, f_ and g_

represent Phylum, Class, Order, Family and Genus, respectively). "Project" represents the

accession numbers of EMP projects and indicate which project the microalgal and bacterial

nodes were recovered from. "Sample_origin" indicates if each sample is planktonic or

benthic. Heatmap colour gradient indicates correlation coefficients. Refer Table S2 for raw

data used to generate the heatmap. Heatmap was generated using Pheatmap v1.0.12

(https://rdrr.io/cran/pheatmap/).

Figure 2: Summarised interactions in Freshwater environments

Correlation heatmap of co-occurring microalgae (columns) and bacteria (rows) in freshwater

environments**.** "Taxonomy" represents the taxonomic affiliations of microalgal nodes at class

level. "Taxonomy2" represents the taxonomic affiliations of bacteria (p_, c_, o_, f_ and g_

represent Phylum, Class, Order, Family and Genus, respectively). "Project" represents the

accessions of EMP projects and indicate which project the microalgal and bacterial nodes

were recovered from. "Sample_origin" indicates if each sample is planktonic or benthic.

Heatmap colour gradient indicates correlation coefficients. Refer Table S3 for raw data used

to generate the heatmap. Heatmap was generated using Pheatmap v1.0.12

(https://rdrr.io/cran/pheatmap/).

Figure 3: Interactions of the Deltaproteobacterial order *NB1-j* with microalgae in a

marine environment.

Figure shows the first neighbours (directly interacting taxa) including microalgal taxa co-

occurring with Deltaproteobacterial order *NB1-j*. Nodes are labelled at the order level (If

unclassified at the order level, higher taxonomic affiliations are provided). The edge width

and node size are continuously mapped to edge weight (correlation strength) and node

degree, respectively. Network image was generated using Cytoscape v3.8

(https://cytoscape.org/)

Supplementary Data

Figure S1: Map indicating the sample collection sites for each EMP project. Sample collection locations for each EMP project are shown on the world map. Geographic coordinates were drawn in R studio (v1.4.1106) using packages rworldmap v1.3-6 (https://rdrr.io/cran/rworldmap/).

Figure S2: Major bacterial Phyla associated with microalgae in freshwater and marine environments

Bars represent the number of bacterial nodes affiliated with each phylum. The total number of bacterial nodes identified to be co-occurring with microalgae in marine environments were 76 and freshwater environments were 311.

Figure S3: *Planctomycetes* class *OM190* interacting with microalgal taxa in a marine (A) and freshwater (B) environment.

Networks (A, B) created from a microalgal-bacterial module by selecting the first neighbours of the *OM190* to showcase the complex interactions with the neighbouring microalgal taxa. The edge width and node size are continuously mapped to edge weight and node degree. Marine and Freshwater environments represent EMP project ERP020022 and ERP016492, respectively. Nodes labelled at order level (If unclassified at the order level, higher taxonomic affiliations are provided). Network image was generated using Cytoscape v3.8 (https://cytoscape.org/).

Table S1: Information on selected EMP studies. Details on Qiita study title, study ids and ENA project accession number.

Table S2: Correlations recovered in marine environments between microalgae and bacteria. Summary of significant co-occurrences recovered from the marine environments. Full taxonomic affiliations of the microalgal and bacterial nodes are provided.

699   **Table S3: Correlations recovered in freshwater environments between microalgae and**

700   **bacteria.** Summary of significant co-occurrences recovered from the freshwater

701   environments. Full taxonomic affiliations of the microalgal and bacterial nodes are provided.