1

Multifunctional role of GPCR signaling in epithelial tube formation

Vishakha Vishwakarma, Thao Phuong Le, and SeYeon Chung*

Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA

*Author for correspondence: seyeonchung@lsu.edu

Key words: *Drosophila*, Salivary gland, Fog signaling, GPCR, Actomyosin

Running title: GPCR signaling in tube formation

ABSTRACT

Epithelial tube formation requires Rho1-dependent actomyosin contractility to generate the cellular forces that drive cell shape changes and rearrangement. Rho1 signaling is activated by G protein-coupled receptor (GPCR) signaling at the cell surface. During *Drosophila* embryonic salivary gland (SG) invagination, the GPCR ligand Folded gastrulation (Fog) activates Rho1 signaling to drive apical constriction. The SG receptor that transduces the Fog signal into Rho1-dependent myosin activation has not been identified. Here, we reveal that the Smog GPCR transduces Fog signal to regulate Rho kinase accumulation and myosin activation in the medioapical region of cells to control apical constriction during SG invagination. We also report on unexpected Fog-independent roles for Smog in maintaining epithelial integrity and organizing cortical actin. Our data supports a model wherein Smog regulates distinct myosin pools and actin cytoskeleton in a ligand-dependent manner during epithelial tube formation.

INTRODUCTION

The formation of three-dimensional tubes by invagination from flat epithelial sheets is a fundamental process in forming organs, such as lungs and kidneys (Andrew and Ewald, 2010). The *Drosophila* embryonic salivary gland (SG) is a premier model system to study the mechanisms underlying epithelial tube morphogenesis (Chung et al., 2014; Girdler and Röper, 2014). The SG begins as a two-dimensional plate of cells on the embryo surface. Neither cell division nor cell death occurs once the SG cells are specified; all the morphogenetic changes arise by changes in cell shape and rearrangement.

A major cell shape change during SG invagination is apical constriction, wherein the apical side of epithelial cells shrinks while keeping the nearly constant volume (Lubarsky and Krasnow, 2003; Martin and Goldstein, 2014). During stage 11, SG cells begin to invaginate at the dorsal/posterior region of the placode through apical constriction (Fig. 1A). This leads to forming a narrow invagination pit through which all SG cells eventually internalize. Apical constriction is observed in both invertebrates and vertebrates, driving morphogenetic processes ranging from gastrulation in *Drosophila* and *Xenopus* to lens formation in the mouse eye to formation of tubular organs such as the *Drosophila* SG and chicken lungs (Kim et al., 2013; Myat and Andrew, 2000; Plageman et al., 2011; Sweeton et al., 1991). Defects in apical constriction result in defects in overall tissue shape, suggesting that coordinated apical constriction is required for final tissue architecture (Chung et al., 2017; Guglielmi et al., 2015; Izquierdo et al., 2018).

Contractile actomyosin networks generate the cellular forces driving coordinated cell behaviors during epithelial morphogenesis (Gorfinkiel and Blanchard, 2011; Levayer and Lecuit, 2012; Munjal and Lecuit, 2014). Non-muscle myosin II (hereafter referred to as myosin) uses its motor activity to pull on actin filaments anchored to adherens junctions (AJs) to apply tension on the junction and neighboring cells. Three distinct pools of myosin generate the contractile forces required for SG invagination (Booth et al., 2014; Chung et al., 2017; Röper, 2012). Medioapical myosin forms in the center of the apical side of the SG cell and drives apical constriction (Booth et al., 2014; Chung et

al., 2017; Röper, 2012). Junctional myosin, associated with AJs of SG cells, is involved

in cell intercalation (Sanchez-Corrales et al., 2018). A supracellular myosin cable

surrounding the entire SG placode generates the tissue-level compressing force (Chung

et al., 2017; Röper, 2012). From studies in the past several years by us and others, we

are beginning to understand how each myosin pool is created and regulated during SG

invagination (Booth et al., 2014; Chung et al., 2017; Le and Chung, 2021; Röper, 2012;

Sanchez-Corrales et al., 2018; Sidor et al., 2020).

Localized activation of the RhoA (Rho1 in *Drosophila*) GTPase is key to creating and/or

polarizing contractile myosin structures in epithelial cells (Blanchard et al., 2010;

Dawes-Hoang et al., 2005; Martin et al., 2009; Mason et al., 2013). Studies in

Drosophila have identified the Folded gastrulation (Fog) pathway as a key signaling

pathway regulating Rho1 signaling during epithelial morphogenesis (Manning and

Rogers, 2014). During *Drosophila* gastrulation, two G protein-coupled receptors

(GPCRs), Smog (ubiquitously expressed) and Mist (mesoderm-specific), respond to

Fog to activate heterotrimeric G proteins, Rho1, and Rho kinase (Rok) to regulate

myosin contractility (Costa et al., 1994; Dawes-Hoang et al., 2005; Kerridge et al., 2016;

Kölsch et al., 2007; Manning et al., 2013; Mason et al., 2013; Parks and Wieschaus,

1991). In our previous study in the SG, we also revealed a tissue-specific regulation of

Fog signaling to activate myosin contractility: the Fork head (Fkh) transcription factor

regulates SG upregulation of *fog*, which promotes accumulation of Rok and myosin

specifically in the medioapical region of SG cells to drive clustered apical constriction

(Chung et al., 2017; Fig. 1A). Yet, it is unknown how the Fog signal is sensed and

transduced into downstream Rho1 signaling during epithelial tube formation.

Studies in early *Drosophila* embryos have suggested a role of GPCR signaling as a

common module for Rho1 activation in different subcellular regions of the cell. In the

mesoderm, Smog and Mist function together to activate myosin in the medioapical

region of cells to drive apical constriction (Kerridge et al., 2016). In the ectoderm, where

no apical constriction occurs, Smog is required for junctional myosin activation to drive

cell intercalation (Kerridge et al., 2016). Recent work further has shown that distinct

RhoGEFs activate myosin contractility in the apical and junctional domain of epithelial

cells under the control of specific G proteins in early *Drosophila* embryos (Garcia De

Las Bayonas et al., 2019). However, whether all three myosin pools in the SG are

regulated by independent mechanisms or by a common module during epithelial tube

formation remains unclear.

The contractile force of the actomyosin networks causes an increase of hydrostatic

pressure in the cytoplasm (Charras et al., 2005). During epithelial morphogenesis, cells

maintain the surface integrity by proper organization of actin at the cell cortex. Loss of

organized cortical actin networks leads to the formation of blebs, spherical protrusions

of the plasma membrane (Charras, 2008). During *Drosophila* gastrulation, blebs

occasionally arise during apical constriction of mesodermal cells; these blebs correlate

with apical F-actin holes (Costa et al., 1994; Jodoin et al., 2015). Disruption of cortical

actin by latrunculin B, which sequesters globular actin (G-actin) and prevents

filamentous actin (F-actin) assembly, enhances blebbing in epithelial cells in early

Drosophila embryos (Kanesaki et al., 2013). Mutations in heterotrimeric G proteins also

enhance blebbing (Kanesaki et al., 2013), suggesting a role of G proteins in stabilizing

cortical actin. The mechanisms of cortical actin regulation during epithelial tube

formation are not well understood.

Here, we reveal multiple roles of Smog GPCR in regulating actomyosin networks during

Drosophila SG tube formation. We show that the GPCR Smog transduces Fog signal to

regulate myosin contractility in the medioapical region of SG cells to drive apical

constriction during SG invagination. Our study supports a model wherein Smog

regulates medioapical and junctional/supracellular myosin pools in a Fog-dependent

and -independent manner, respectively. We further reveal new roles of Smog in

regulating the microtubule networks, key apical and junction proteins, and cortical actin

organization during SG invagination, suggesting multifunctional roles of GPCR signaling

during epithelial tube formation.

RESULTS

Smog transduces Fog signal in the SG

To test the role of Smog in transducing Fog signal during SG invagination, we first examined *smog* transcripts and Smog protein levels in the developing SG. Consistent with the ubiquitous expression of *smog* (Kerridge et al., 2016), fluorescent in situ hybridization revealed low levels of *smog* mRNA expression in the entire embryo, including the SG. Strong *smog* mRNA signals were observed in SG cells near the invagination pit (Fig. 1B, B'), potentially due to the shrinking apical domain in those cells. We also examined Smog protein localization in the SG using a functional Smog-GFP fusion protein expressed under the control of the *sqh* promoter (Kerridge et al., 2016). Smog-GFP signals were detected as small punctate structures enriched near AJs in SG cells (Fig. 1C-C''; AJs were marked with the key AJ protein E-Cadherin (E-Cad). Importantly, Smog-GFP signals were dramatically increased in the entire apical domain of Fog-overexpressing SGs (Fig. 1D-D''; quantification in Fig. 1E), suggesting that Smog is recruited to the apical domain of SG cells by overproduced Fog.

To test whether Smog transduces Fog signal in the SG to regulate myosin contractility, we performed a genetic suppression test. We hypothesized that if Smog is a SG receptor for Fog, knocking down *smog* in the SG should suppress the gain-of-function effect of Fog. We used sqh-GFP, a GFP-tagged version of myosin regulatory light chain (Royou et al., 2004), and Rok-GFP, a GFP-tagged Rok fusion protein (Abreu-Blanco et al., 2014), as readouts of Fog signaling. sqh-GFP signals were significantly increased in the medioapical region of Fog-overexpressing SG cells (compare Fig. 1G', G''' to Fig. 1F', F'''; quantification in Fig. 1H) but not at AJs (Fig. 1I), suggesting increased medioapical myosin by Fog overexpression. Fog-overexpressing SG cells showed highly distorted AJs (Fig. 1G'') compared to control SG cells (Fig 1F''), suggesting increased pulling forces due to increased medioapical myosin in Fog-overexpressing SG cells. Consistent with our finding that Fog signal promotes accumulation of Rok in the medioapical region of SG cells (Chung et al., 2017), quantification of areas occupied by Rok-GFP puncta revealed overaccumulated Rok-GFP signals in the medioapical region of Fog-overexpressing SG cells (compare Fig. 1L', L''' to Fig. 1K', K''';is quantification in Fig. 1N). Knockdown of *smog* in the SG using RNA interference (RNAi)

and the SG-specific *fkh-Gal4* driver (Henderson and Andrew, 2000) both suppressed

the medioapical accumulation of Rok-GFP signals and reduced the magnitude of AJ

distortion (Fig. 1M-M'''; quantification in 1N, O). These data suggest that Smog

transduces Fog signal in the SG to facilitate Rok and myosin accumulation in the

medioapical region of SG cells.

Smog-transfected *Drosophila* S2 cells contract upon Fog signal

To further test the role of Smog in transducing Fog signal to regulate myosin

contractility, we performed an in vitro cell contraction assay using *Drosophila* S2 cells.

S2 cells do not express Fog receptors, and therefore, do not respond to Fog in the

normal culture condition (Manning et al., 2013; Fig. 2A-B'). As shown in a previous

study (Manning et al., 2013), S2 cells transfected with Mist, the mesoderm-specific

receptor for Fog, contracted robustly when cultured in Fog-containing media (Fig. 2C-

D'). Since Smog was not detectable in *Drosophila* S2 cells (modENCODE Cell Line

Expression Data; Flybase), we tested if S2 cells transfected with Smog also contract

upon Fog treatment. Indeed, Smog-transfected cells showed a significantly higher

percentage of contraction upon Fog treatment compared to non-transfected S2 cells

(Fig. 2A-B', E-F'; quantification in Fig. 2G). Thus, Smog can transduce Fog signal to

regulate cellular contractility both in the SG and in vitro.

Smog is required for apical constriction in the SG and epithelial morphogenesis

during embryogenesis

To test the role of Smog in apical constriction of SG cells as a Fog receptor, we

knocked down *smog* in the SG (using *fkh-Gal4*) using RNAi and compared apical areas

of SG cells to those in *fog* mutants. Two independent short hairpin RNAi lines,

TRiP.HMC03192 (a stronger line; Fig. 3B, B'; also used for genetic suppression assay

in Fig. 1M-M''') and *TRiP.GL01473* (a weaker line; Fig. S1G, G'), were used. *smog*

knockdown using either line reduced Smog-GFP signals in the SG, with a stronger

effect with *TRiP.HMC03192*, confirming the effect of the RNAi lines (Fig. S1A-D). Cell

segmentation analysis revealed that SG cells with smaller apical areas showed a less

coordinated spatial distribution in *smog* RNAi SGs compared to control (Fig. 3A-B'; Fig.

S1F-G'), similar to the previously reported *fog* mutant phenotype (Chung et al., 2017;

Fig. S1E, E'). However, quantification of percentage and cumulative percentage of cells

of different apical areas revealed different behaviors between *fog* and *smog*. In *fog*

mutants, the number of constricting cells is not significantly different from wild type

(WT), but cells with small apical areas are dispersed across the entire SG placode

(Chung et al., 2017). On the other hand, SGs with *smog* knockdown with either RNA

line showed larger apical areas compared to control (Fig. 3G; Fig. S1H). The difference

in apical constriction defects between *fog* mutants and *smog* knockdown suggests a

potential Fog-independent role of Smog.

Since *smog* has a strong maternal contribution (Flybase Developmental RNA-Seq), we

also knocked down *smog* both maternally (using *matα-Gal4*; Häcker and Perrimon,

1998) and zygotically in the SG (using *fkh-Gal4*) and observed slightly more severe

apical constriction defects in SG cells (hereafter referred to as *smog* M/Z knockdown;

Fig. 3C, C'; quantification in 3G). Maternal knockdown of *smog* using the stronger RNAi

line (*TRiP.HMC03192*) resulted in severe defects in egg-laying. Therefore, all M/Z

knockdown experiments were performed using the weaker line (*TRiP.GL01473*).

Consistent with the ubiquitous expression of *smog* and its role in epithelial

morphogenesis in the early *Drosophila* embryo (Kerridge et al., 2016), *smog* M/Z

knockdown resulted in a slight disruption of the overall embryonic morphology, including

defects in the head region, a wavy embryo surface, and an irregular ventral midline (Fig.

3M, N). Unlike the circular SG placode of WT, *smog* M/Z knockdown also resulted in a

SG placode elongated along the dorsal/ventral axis (compare Fig. 3A, A' to 3C, C'). 3D

reconstruction of confocal images revealed disrupted epithelial morphology in *smog* M/Z

knockdown, with wide and wavy grooves, enlarged epithelial cells, and an elongated SG

placode (Fig. 3I, J).

smog zygotic null mutants (Kerridge et al., 2016) had a range of defects similar to those

observed with *smog* knockdown. 73.8% of embryos (59/80) showed relatively normal

embryonic morphology with mild apical constriction defects in SG cells, like *smog*

zygotic knockdown (classified as 'smog null [mild]'; Fig. 3D-E', K; quantification in 3H);

26.2% of embryos (21/80) showed a disrupted embryonic morphology with SGs

elongated along the dorsal/ventral axis (Fig. 3F, F', L) and more severe apical

constriction defects (Fig. 3H), like smog M/Z knockdown (classified as 'smog null

[severe]'). All embryos at late stages (stages 15-17) smog null mutants showed

relatively normal embryonic morphology (Fig. S2A, B), suggesting that severely

defective embryos with smog loss do not survive/develop past stage 14. Notably, these

embryos have relatively normal internalized SGs (Fig. S2E-G), except for rare cases of

crooked SG morphology (Fig. S2C, D). Altogether, these data suggest that Smog is

required for coordinated apical constriction during SG invagination and proper epithelial

morphogenesis during embryogenesis.

Smog regulates medioapical and junctional myosin in a Fog-dependent and -

independent manner in the SG

We next examined myosin levels and distribution in the SGs in smog knockdown or

smog mutants using sqh-GFP. Medioapical myosin is the only myosin pool defective in

the SGs of fog mutants (Chung et al., 2017; Fig. S3A-E). Consistent with the idea that

Smog transduces Fog signal in the SG, smog knockdown (either zygotic or M/Z

knockdown) and smog loss resulted in a significant reduction in the intensity of

medioapical myosin in SG cells near the invagination pit, compared to control (Fig. 4A-

D''; quantification in 4I). Moreover, compared to clear web-like structures of medioapical

myosin in control SG cells (Fig. 4A-A''), SG cells in smog knockdown or smog null

mutants showed reduced areas of sqh-GFP puncta (Fig. 4L), with dispersed myosin

along the entire apical surface in smog M/Z knockdown SG cells (Fig. 4C-C'').

Interestingly, the intensity of junctional myosin was also significantly reduced in SG cells

with smog knockdown or smog loss, compared to control (Fig. 4A-D'', quantification in

4J). This phenotype is distinct from what was observed in fog mutants, where junctional

myosin was unaffected in SG cells (Chung et al., 2017; Fig. S3D). The ratio of

medioapical to junctional myosin intensity was significantly reduced in smog M/Z

knockdown but not in smog zygotic knockdown or smog null [mild] SGs (Fig. 4K). We

conclude that Smog is required for maintaining normal levels of both medioapical and

junctional myosin in the SG in a Fog-dependent and -independent manner, respectively.

We also tested Rok distribution in SG cells in *smog* RNAi and *smog* null mutants using

Rok-GFP. Whereas Rok-GFP formed large punctate structures in the medioapical

region in control SG cells (Fig. 4E-E''), Rok-GFP failed to accumulate and was

dispersed along the entire apical domain in SG cells in *smog* knockdown and *smog* null

mutants (Fig. 4F-H''; quantification in 4M), similar to the dispersed Rok-GFP distribution

observed in *fog* mutant SG cells (Chung et al., 2017). These data suggest that Smog is

required for Rok accumulation in the medioapical domain of SG cells.

smog loss leads to defects in the tissue-level myosin cable

Reduced myosin levels and the elongated SG placode in *smog* null [severe] embryos

led us to test the third pool of myosin in the SG, the supracellular myosin cable

surrounding the SG placode. Consistent with the idea that this myosin cable is under

tension (Chung et al., 2017; Röper, 2012), WT SGs showed a relatively smooth tissue

boundary (Fig. 4N-N''). *smog* null [mild] embryos also maintained a smooth SG

boundary comparable to WT (Fig. 4O-O''). However, SGs in *smog* null [severe] embryos

showed an irregular boundary and a less clear myosin cable (Fig. 4P-P'').

We calculated the circularity of the SG placode as a measure of smoothness and

tension of the SG boundary. The circularity of the WT placode was significantly higher

than the circularity of one cell row inside (inner boundary) or outside of the SG placode

(outer boundary), suggesting a higher tension of the myosin cable in WT (Fig. 4Q, R).

The circularity of the placode of *smog* null [mild] embryos was comparable to WT (Fig.

4R). However, the circularity of the SG placode in *smog* null [severe] embryos was

significantly lower than that of WT and was not statistically significant from the inner or

outer boundaries, suggesting a lack of tension at SG boundary in *smog* null [severe]

mutants (Fig. 4R). Consistent with our previous study (Chung et al., 2017), *fog* mutants

did not show defects in the myosin cable (Fig. S3F-H), even in the severely twisted

embryos with extra grooves and folds. Overall, our data suggest that, unlike *fog*, the

loss of *smog* affects all myosin pools in the SG, including the myosin at the SG

boundary, leading to reduced tension at the SG boundary and defects in generating the

compressing force.

Smog is required for proper localization of apical and junctional components and

microtubule networks in epithelial cells

Since we observed a range of defects in whole embryo morphology in *smog* mutants,

we explored the potential cellular basis. In stage 11 *smog* null [severe] embryos, we

occasionally observed large areas in the epidermis where E-Cad (Fig. 5B, B'; compare

to WT in Fig. 5A, A') and the apical determinant protein Crumbs (Crb) (Fig. 5D, D';

compare to WT in Fig. 5C, C') signals were absent or significantly reduced. Relatively

strong signals of the cell membrane marker Gap43-mCherry (Martin et al., 2010) were

observed in these regions (Fig. 5E-E''), suggesting that the cell membrane is still intact.

Cells in these regions had enlarged apical areas compared to neighboring cells with

strong E-Cad signals (Fig. 5E-E''), suggesting reduced epithelial tension and/or integrity

in cells with reduced E-Cad levels. Similarly, in late-stage (stage 14) embryos, Crb

levels were significantly reduced in large areas; in the most severe cases, Crb was

dispersed in the entire epidermis (Fig. S4B, B'). These data suggest that Smog is

required for maintaining epithelial polarity and junctional integrity by regulating E-Cad

and Crb levels or distribution.

To better understand the basis of the reduced E-Cad and Crb in *smog* mutants, we

examined *smog* null mutant SG cells at higher magnification. Even in SGs that do not

contain large patches of E-Cad/Crb loss, E-Cad/Crb signals were discontinuous, with

small gaps in signal along the cell boundary (Fig. 5F-O). Such gaps were observed in

both *smog* null [mild] and *smog* null [severe] embryos. Although the number and the

length of gaps in E-Cad signals in *smog* null [mild] were comparable to those of WT,

they were significantly higher in *smog* null [severe] compared to WT (Fig. 5G, H;

quantification in 5L, M). The number and the length of gaps in Crb signals in SG cells

were significantly higher in both mild and severe *smog* null mutant embryos compared

to WT (Fig. 5J, K; quantification in 5N, O). These data suggest a role of Smog in

regulating Crb and E-Cad levels and continuity during epithelial morphogenesis,

including the SG.

Anisotropic localization of Crb at the SG boundary has been suggested to drive the

formation of supracellular myosin cable in SG (Röper, 2012). Consistent with this, in

control SGs, Crb levels were higher in inside junctions that do not contribute to the

myosin cable, whereas myosin was highly enriched at the junction forming the cable at

the SG boundary (Fig. S4C-E'''). However, in *smog* null [severe] SGs, the overall Crb

intensity was quite low, and Crb did not show an anisotropic localization at the SG

boundary (Fig. S4F-H'''). This data suggests that *smog* loss results in the reduction of

both myosin and Crb, failing to form the supracellular myosin cable at the SG boundary.

Microtubules are required for forming and maintaining apical myosin structures (Booth

et al., 2014; Ko et al., 2019) and for transporting several key apical and junctional

proteins in *Drosophila* tubular organs, including E-Cad and Crb (Le and Chung, 2021; le

Droguen et al., 2015). We recently showed that microtubule-dependent intracellular

trafficking has an important role in regulating apical constriction and medioapical myosin

during SG invagination (Le and Chung, 2021). To examine the microtubule networks in

the SG of WT and *smog* mutant embryos, we stained with antibodies against acetylated

α-tubulin (Ace-tub), a marker of stable, long-lived microtubules, and tyrosinated α-

tubulin (Tyr-tub), a marker of dynamic, short-lived microtubules (Westermann and

Weber, 2003). Compared to WT, both Ace-tub and Tyr-tub signals were reduced in

smog mutants (compare Fig. 5Q, R, U, V to Fig. 5P, T), with a dramatic reduction in

Ace-tub levels in *smog* null [severe] embryos (Fig. 5R), suggesting disrupted

microtubule networks in *smog* mutants. The reduced Ace-tub and Tyr-tub signals were

more prominent in the SG compared to the surrounding epithelial cells; the ratio of Ace-

tub and Tyr-tub intensity in SG cells to the intensity in cells outside the SG was

significantly decreased in *smog* mutants compared to WT (Fig. 5X, Y). A slight but less

prominent reduction in Ace-tub and Tyr-tub signals was also observed in the SG in *fog*

mutants (Fig. 5S, W; quantification in 5X, Y). Overall, this finding suggests that loss of

smog could affect apical pools of myosin, Crb and E-Cad through effects on microtubule
abundance, distribution and/or polarity.

Smog is required for cortical actin organization in the SG
We observed numerous blebs in the apical membrane of SG cells in *smog* null [severe]
mutants, which were most prominent near the invagination pit (compare Fig. 6A-A''' to
Fig. 6B-B'''). Such blebs were not observed in SGs in WT, *smog* null [mild], or *fog*
mutant embryos (Fig. S5A-C''). Rok-GFP was enriched in many of these blebs (Fig. 6B-
B'''), consistent with Rok's recruitment to retracting blebs (Aoki et al., 2016). In *smog*
null [severe] embryos, blebs were also occasionally observed in cells outside the SG
placode (Fig. S5D-E''). During *Drosophila* gastrulation, bleb formation correlates with
cortical F-actin holes (Jodoin et al., 2015), and disruption of cortical actin enhances
blebbing (Kanesaki et al., 2013). We therefore tested if Smog affects the cortical actin
network during apical constriction in the SG. Indeed, phalloidin staining revealed that
whereas strong F-actin signals were observed in the apical domain of SG cells near the
invagination pit in WT, F-actin was reduced in SGs in *smog* mutants (Fig. 6C-E).

Apical F-actin formation in *Drosophila* tubular organs requires the formin family actin
nucleator Diaphanous (Dia) (Massarwa et al., 2009; Rousso et al., 2013). Since the
apical localization and activity of Dia are critical for restricting F-actin formation to the
correct membrane domain, we tested the localization and levels of Dia in the SG in
smog mutants. Similar to reduced F-actin (Fig. 6C-E), we observed reduced Dia signals
in the apical domain of SG cells in *smog* mutants, compared to WT (Fig. 6G-I).
Quantification of Dia levels revealed that the overall Dia intensity is reduced in the SG
placode in *smog* mutants, with a stronger effect in the *smog* null [severe] than *smog* null
[mild] embryos (Fig. 6K). SGs in *fog* mutants appeared to show a slight reduction of F-
actin (Fig. 6F), but Dia levels in *fog* mutants were comparable to WT (Fig. 6J, K). Our
data suggest that Smog is required for the cortical actin networks during SG
invagination, at least in part via modulating Dia levels.

We next tested if the reduction in actin levels in *smog* mutants is due to defects in actin

turnover. To test this, we modulated the level of actin regulators in the SG in *smog*

mutants and assayed the blebbing phenotype. Using RNAi, we knocked down

chickadee (*chic*), which encodes *Drosophila* profilin, a protein that increases F-actin by

promoting actin polymerization (Cooley et al., 1992), in the *smog* mutant background. If

there is a shift from F-actin to monomeric G-actin in *smog* mutants, we expect

enhancement of the disorganized actin phenotype and bleb formation with a reduced

chic level. Blebs were not observed in *smog* null [mild] embryos or in SGs knocked

down for *chic* in the otherwise wild-type background (Fig. 6L-M''). However, when *chic*

was knocked down in the SG in the *smog* mutant background, we observed enhanced

blebbing even in SG cells in *smog* null [mild] embryos (Fig. 6N-N''). Since profilin

promotes formin-mediated actin filament assembly (Romero et al., 2004; Zweifel and

Courtemanche, 2020), our data suggests that reduced profilin by *chic* knockdown

aggravated defects in cortical actin organization with reduced Dia in *smog* mutants.

Overall, our data suggest that Smog is required for cortical actin organization as well as

myosin activation during epithelial morphogenesis.

DISCUSSION

Smog regulates different pools of myosin in a Fog-dependent and -independent

manner during SG invagination

Fog signaling triggers epithelial cell shape changes driving tissue folding and

invagination during development of *Drosophila* and other insects (Benton et al., 2019;

Manning and Rogers, 2014). During *Drosophila* embryogenesis, *fog* is upregulated in

multiple tissues undergoing apical constriction and tissue invagination, such as ventral

furrow (mesoderm), the posterior midgut (endoderm), and the SG (ectodermal

derivative) (Nikolaidou and Barrett, 2004). During *Drosophila* gastrulation, the

ubiquitously expressed GPCR Smog and the mesoderm-specific GPCR Mist respond to

Fog signal to regulate myosin contractility during mesoderm invagination (Kerridge et

al., 2016; Manning et al., 2013). Here, we provide evidence that Smog functions as a

SG receptor for Fog to activate myosin during SG invagination. Knockdown of *smog* by

RNAi suppresses the gain-of-function effect of Fog in the SG (Fig. 1). Also, Smog-

transfected S2 cells contract upon Fog signal (Fig. 2), suggesting that Smog responds

to Fog to regulate myosin contractility both in vivo and in vitro. Our data show that Fog

overexpression leads to Smog recruitment to the medioapical region of SG cells (Fig.

1). Consistent with our data, Fog overexpression induces oligomerization of Smog in

early *Drosophila* embryos (Jha et al., 2018).

Our study also dissects the roles of Smog in myosin activation during SG

morphogenesis and reveals Fog-independent roles of Smog. Whereas *fog* mutants

exhibit a decrease in only the medioapical myosin pool in SG cells (Chung et al., 2017;

Fig. S3), *smog* loss and SG-specific knockdown of *smog* result in a significant reduction

of both medioapical and junctional myosin (Fig. 4). We propose that Smog regulates

junctional myosin during epithelial morphogenesis in response to an unknown,

ubiquitously expressed ligand and regulates medioapical myosin in response to Fog in

tissues with high Fog signals, such as the mesoderm (Kerridge et al., 2016; Manning et

al., 2013) and the SG (this study). Consistent with this model, during *Drosophila*

gastrulation, Smog activates different myosin pools in tissues with different Fog levels.

In the mesoderm, where Fog levels are high, Smog and Mist transduce Fog signal to

activate myosin in the medioapical region and drive apical constriction. In the ectoderm,

where Fog levels are very low, Smog is required for junctional myosin activation to drive

cell intercalation. SG cells undergo both apical constriction and cell intercalation, and

our data shows that Smog regulates both pools of myosin in the SG in a Fog-dependent

and -independent manner. It is possible that in tissues with high Fog signals, an

additional tissue-specific GPCR(s) function with Smog in response to Fog to fine-tune

the Fog signaling for proper apical constriction, like Mist does in the mesoderm. It will be

interesting to determine whether additional SG-specific GPCRs function with Smog

during SG invagination.

The downstream components that transduce signals from the Smog GPCR to regulate

myosin in distinct regions of SG cells await discovery. Good candidates include two Rho

guanine nucleotide exchange factors (RhoGEFs), RhoGEF2 and Dp114RhoGEF, which

have been shown to activate Rho1 signaling at the medioapical and junctional domain

of epithelial cells, respectively (Garcia De Las Bayonas et al., 2019). This process

requires upstream activation by heterotrimeric G proteins $G\alpha_{12/13}$/Cta and $G\beta13F$/$G\gamma1$ at

the medioapical and junctional domains, respectively (Garcia De Las Bayonas et al.,

2019). Thus, distinct ligand-receptor binding with Smog may activate distinct G proteins

and/or RhoGEFs in the medioapical and junctional domains.

Smog regulates cortical actin organization during SG invagination

We also discover that Smog regulates cortical F-actin organization during SG

invagination (Fig. 6). *smog* loss significantly reduces levels of F-actin and the formin

protein Dia (Fig. 6). Rho1 regulates actin polymerization via Dia (Goode and Eck, 2007),

and apical F-actin formation in the *Drosophila* SG requires apical localization and

activity of Dia (Massarwa et al., 2009; Rousso et al., 2013). Our data suggest that Smog

activates Rho1 signaling to regulate both cortical F-actin (via Dia) and myosin (via Rok)

(Fig. 7). *fog* mutants also show slightly reduced F-actin and Dia levels, to a similar level

to *smog* null [mild] embryos (Fig. 7), and these embryos do not show membrane

blebbing (Fig. 7; Fig. S5). It is possible that Smog regulates cortical actin in both Fog-

dependent and independent manners and perhaps subtle changes in Dia and F-actin

levels in *fog* mutants are not enough to cause membrane blebbing. Alternatively,

Smog's role in regulating cortical actin is independent of Fog; reduced F-actin levels in

fog mutants may be a secondary effect due to a defective medioapical myosin pool.

During *Drosophila* gastrulation, *fog* mutants show normal GFP-Moesin signals in the

ventral furrow, suggesting normal cortical actin organization upon *fog* loss (Kanesaki et

al., 2013). $G\alpha_{12/13}$/Cta is a key component downstream of Fog that recruits RhoGEF2 to

the apical membrane to regulate myosin contractility (Barrett et al., 1997; Dawes-Hoang

et al., 2005). Interestingly, mutations in Gαi (G-iα65A) and Gβ13F/Gγ1, but not

$G\alpha_{12/13}$/Cta, disrupt cortical actin organization in early *Drosophila* embryos (Fox and

Peifer, 2007; Kanesaki et al., 2013). Therefore, Smog may signal through Gαi and/or

Gβ13F/Gγ1 to organize cortical actin, independent of Fog and $G\alpha_{12/13}$/Cta.

The link between heteromeric G proteins and Rho1 signaling occurs through activation

of specific RhoGEFs (Vázquez-Victorio et al., 2016). Actin polymerization resulting from

Fog-independent Smog activation may imply the recruitment of distinct proteins

modulating the activation of Rho1-mediated pathways. For example, in the absence of

Fog, a specific RhoGEF(s) downstream of Gαi or Gβ13F/Gγ1 may link Smog and Rho1

signaling to regulate cortical actin organization; upon high Fog signals, a different Rho1

modulator(s) downstream of $G\alpha_{12/13}$/Cta may be recruited to facilitate medioapical

myosin formation. Further work is needed to identify the link between Smog and Rho1

activation in the presence or absence of Fog during SG invagination. It will be also

interesting to determine whether Fog-independent roles of Smog–regulating cortical

actin and junctional myosin–involve common or distinct Rho1 modulators.

Smog regulates epithelial integrity during development

smog null mutant embryos show a range of defects, allowing us to dissect the roles of

Smog in epithelial morphogenesis. Our data reveal that Smog is required for epithelial

integrity by affecting levels and localization of Crb and E-Cad (Fig. 5). Since the

decrease in the level of Crb and E-Cad in *smog* mutants correlates with significantly

reduced myosin levels, disrupted epithelial integrity may be due to disrupted actomyosin

networks. Myosin II is critical for cells to concentrate E-cad at cell-cell contacts (Shewan

et al., 2005). In early *Drosophila* embryos, the absence of contractile myosin leads to

the disassembly of junctions (Weng and Wieschaus, 2016). A possible scenario is that

Smog is required to recruit key Rho1 regulators, such as RhoGEF2, Dp114RhoGEF, or

other SG-upregulated RhoGEFs, to AJs during SG invagination to promote junctional

myosin assembly. Since Dia helps coordinate adhesion and contractility of actomyosin

(Homem and Peifer, 2008) and formin-mediated actin polymerization at AJs stabilizes

E-Cad and maintains epithelial integrity (Rao and Zaidel-Bar, 2016), reduced Dia levels in

smog mutants may also contribute to loss of epithelial integrity.

Reduced and discontinuous signals of apical and junctional components in *smog*

mutant embryos can also be due, in part, to compromised intracellular trafficking.

Disorganized microtubule networks in SGs in *smog* mutants (Fig. 5) support this idea.

Our recent study showed that microtubule- and Rab11-dependent intracellular

trafficking regulates apical myosin pools and apical constriction during SG invagination,

via apical enrichment of Fog and the continuous distribution of Crb and E-Cad along

junctions (Le and Chung, 2021). The mechanism of how Smog affects microtubule

organization remains to be revealed. *fog* mutants also show slightly disrupted

microtubule networks, albeit less prominently than *smog* null [severe] (Fig. 5). Whether

this is a direct effect of loss of *fog* or an indirect result of the disrupted medioapical pool

of myosin remains to be answered. Overall, our findings suggest the multifaceted roles

of Smog during epithelial tube formation in regulating distinct myosin pools and cortical

actin in different subcellular domains in a ligand-dependent manner.

MATERIALS AND METHODS

Fly stocks and genetics

The fly stocks used in this study are listed in Table S1. All the crosses were performed at 25°C.

Immunofluorescence and confocal microscopy

Drosophila embryos were collected on grape juice agar plates supplemented with yeast paste at 25℃. Embryos were dechorionated with 50% bleach. For most samples, embryos were fixed in 1:1 heptane: formaldehyde for 40 minutes and devitellinized with 80% ethanol. For Rok-GFP, sqh-GFP, and phalloidin staining, embryos were hand-devitellinized. Embryos were then stained with primary and secondary antibodies in PBSTB (1X PBS with 0.1% Triton X-100 and 0.2% bovine serum albumin) and PBTr (1X PBS with 0.1% Triton X-100) for ethanol devitalization and hand-devitalization, respectively. Antibodies used in our experiments are listed in Table S2. Embryos were mounted in Aqua-Poly/Mount (Polysciences, Inc) and imaged with a Leica TCS SP8 confocal microscope using 63x, NA 1.4 and 40x, NA 1.3 objectives. Images were acquired as z-stacks (each 0.3 µm apart) that span the apical and junctional domains of cells in the SG placode.

Fluorescent in situ hybridization

Embryos were fixed using formaldehyde/heptane for 30 min followed by devitellinization with methanol. SP6 polymerase-synthesized digoxigenin (DIG)-labeled antisense probe was prepared using the following primers: *smog*-F, 5'-ACAGAGCCCACCTGTGTAGG-3'; *smog*-R, 5'-TCGCTGATCGAAAATGATCTC-3'. Fluorescent in situ hybridization was performed using standard methods as described in (Knirr et al., 1999). Briefly, the embryos were pre-hybridized for 1 hour at 56°C post-fixation. Following this step, the embryos were hybridized with the probe overnight at 56°C. The next day, embryos were stained for DIG using the anti-DIG primary antibody and biotin-conjugated secondary antibody. The embryos were then incubated with AB solution (PK-6100, VECTASTAIN Elite ABC kit). Tyramide signal amplification reaction was performed using Cy3 fluorescent dye (diluted 1:50 using amplification diluent) (Perkin Elmer Life Sciences,

Inc., NEL753001kt). The embryos were co-stained with CrebA and Crb antibodies for

visualizing SG nuclei and apical cell boundaries, respectively.

Total Smog-GFP and Dia intensity

For Smog-GFP quantification, maximum intensity projections of two z-sections that

span the apical and the junctional region of SG cells were used. Intensity means of

Smog-GFP signals of SG cells in the entire SG placode were measured using Imaris

software (Bitplane). The mean intensity of Smog-GFP was normalized by the median

deviation. The integrated density of Smog-GFP was calculated by multiplying the apical

area of the cell with the mean intensity of each cell. Total Smog-GFP signals within the

whole placode were calculated as the sum of the integrated density of all cells. For

background correction, mean gray values of Smog-GFP in ten cells outside of the SG

placode were measured using ImageJ (NIH). The average value of mean gray values of

Smog-GFP in these ten cells was used to subtract the background of the cells inside the

placode from the same embryo. Five SGs were used for quantification for each

genotype except for control in Fig. 1E, where four SGs were used. Statistical

significance was determined using Mann-Whitney U test in the GraphPad Prism

software.

For total Dia intensity quantification, maximum intensity projection of two z-sections of

the most apical region of SG cells was used. The images were then exported to ImageJ

and converted to the RGB stack image type. Mean gray values of Dia signals in the

whole placode were measured. For background subtraction, the average mean gray

value of Dia signals for ten cells outside the SG placode was subtracted from the mean

gray value of Dia intensity of the whole SG placode. These values were scaled by the

mean normalization. The total Dia intensity in the whole placode was calculated by

multiplying the normalized intensity with the entire area of the placode. Statistical

significance was determined using Mann-Whitney U test.

Cell segmentation and apical area quantification

SGs that were invaginated within the range of 4.8-9.9 µm depth were used for

quantification. Two or three z-sections of the apical domain were used to generate a

maximum intensity projection (Leica LasX software). SG cells were marked using CrebA

and segmented along E-Cad signals using the Imaris program. Apical areas of

segmented cells were calculated using Imaris, and cells were color-coded based on

their apical domain size. Frequency distribution was performed using GraphPad Prism.

Apical areas from eight SGs were quantified in control, *smog* knockdown (Z), and *smog*

null [mild] embryos. Six SGs were used for quantification for *smog* knockdown (M/Z)

and *smog* null [severe] embryos. Statistical significance was determined using Mann-

Whitney U test (percentage of cells) and Kolmogorov-Smirnov test (cumulative

percentage of cells).

Quantification of myosin intensity in the medioapical and junctional domains

Maximum intensity projection of two z-sections spanning apical and junctional regions of

SG cells with sqh-GFP and E-Cad signals was used. The images were then exported to

ImageJ and converted to RGB stack image type. Twenty cells near the invagination pit

in each of five SGs were used for quantification. To calculate the medioapical and

junctional myosin intensity, regions were drawn manually along the inner and outer

boundary of the E-Cad signal of each cell, and mean gray values were measured for

medioapical and junctional myosin. For background subtraction, the average mean gray

value of sqh-GFP signals for ten cells outside the SG placode was subtracted from the

mean gray value of medioapical and junctional myosin for each SG cell. SuperPlots

(Lord et al., 2020) were used to address the variability of datasets in each SG. Each

data point in the graph represents one SG consisting of twenty cells used for

quantification. Statistical significance was determined using Welch's t-test. A normal

distribution was tested using Jarque-Bera test (Microsoft Excel) or a bell-shape of data

distribution by making a histogram.

Quantification of areas of Rok-GFP and medioapical myosin particles

A single z-section of the confocal image with the strongest Rok-GFP signals in the

medioapical region was used. Rok-GFP signals were converted to black and white

using the threshold tool in Adobe Photoshop. Junctional Rok-GFP signals were

removed manually based on E-Cad signals. Areas of Rok-GFP puncta were determined

using the Analyze Particles tool in ImageJ. Rok-GFP puncta with area ≥ 0.02 μm^2 was

measured. Fifteen cells in the dorsal posterior region near the invagination pit of the SG

placode were used for quantification. The same strategy was used for sqh-GFP signals

to quantify the area of medioapical myosin. SuperPlots show mean values of data

points of five SGs. Statistical significance was determined using Mann-Whitney U test.

Quantification of the waviness of junctions

Using a single z-section with the highest E-Cad signals and ImageJ software, the

shortest (L_0) and the actual distance (L) between vertices were measured in SG cells.

The ratio of these distances (L/L_0) was used as the waviness of junctions. 10-15 cells

near the invagination pit in each of five SGs were used for quantification. Statistical

significance was determined using Mann-Whitney U test.

Circularity of the SG boundary

Using confocal images and ImageJ, cell boundaries were manually drawn along E-Cad

signals at the SG placode boundary and one cell row outside and inside the placode.

For most samples, strong myosin cable signals at the dorsal, anterior, and posterior

boundaries of the placode and CrebA signals in SG cells were used to determine the

SG boundary. In *smog* null [severe], where myosin signals were significantly reduced in

all cells, CrebA signals were used to determine the boundary. The ventral midline was

used as the ventral boundary of the SG in all cases. The perimeter and the area of the

SG corresponding to these boundaries were measured using ImageJ, and circularity

was calculated using the formula, $C = 4\pi$ area/perimeter^2. For a perfect circle, the value

of circularity should be 1. Five SGs were used for quantification, and statistical

significance was determined using Welch's t-test.

Quantification of number and length of gaps of Crb and E-Cad

The number and the length of gaps for Crb and E-Cad signals were measured using

confocal images and ImageJ. A single z-section with the highest Crb or E-Cad signals

was chosen for quantification. Gaps with a length ≥0.2 μm were used for quantification.

All junctions from ten cells in five SGs were quantified. If there were multiple gaps on a

junction, we added the length of all the gaps on that junction to calculate the ratio of the

length of gaps to junctional length. Statistical significance was determined using Mann-

Whitney U test.

Acetylated and tyrosinated α-tubulin intensity quantification

Three z-sections spanning the most apical region of SG cells were used for

quantification. The SG boundary was drawn manually based on the E-Cad and CrebA

signals. The mean gray value of Ace-tub signals in the whole SG placode and ten cells

outside the SG of the same embryo was measured using ImageJ. The ratio of the mean

gray value of Tyr-tub signals inside the placode to the average mean gray value of cells

outside the placode was calculated and plotted. The same method was used for Tyr-

tub. Five SGs for each genotype were used for quantification. Mann-Whitney U test was

used to calculate the p values.

Cell contractility assay

cDNA clones for *smog* (RE70685) and *mist* (RE13854) were obtained from *Drosophila*

Genomics Resource Center (DGRC). Isolated cDNAs were cloned in-frame with the V5

tag into the pMT-V5-HisB vector (Addgene) using the conventional restriction digestion

and ligation method. Primers with restriction sites used to isolate the open reading

frame are stated below (restriction sites underlined).

smog CDS-EcoRI-5', 5'-CCG<u>GAATTC</u>ATGGAACTGTGCATAGCAAC-3'

smog CDS-NotI-3', 5'-ATTT<u>GCGGCCGC</u>ATTGGTCGTGATTGTATCTTTGG-3'

mist CDS-EcoRI-5', 5'-CCG<u>GAATTC</u>ATGGACAGGAGTCGGAGTAGC-3'

mist CDS-NotI-3', 5'-ATTT<u>GCGGCCGC</u>AGCAAATGGTCTCCATTTTG-3'

S2 and S2-Mt-Fog-myc cells (Manning et al., 2013) were obtained from DGRC. Cells

were tested for contamination regularly. Shields and Sang M3 insect media (HiMedia)

was used with 10% Fetal Bovine Serum (Corning) and 1:1000 Antibiotic-Antimycotic

(Gibco). Cells were cultivated at 25°C either in 25-cm^2 flask in 5 ml media or 75-cm^2

flask in 15 ml media. To prepare for the Fog media, S2-Mt-Fog-myc cells were grown

undisturbed for 4-8 days to attain approximately 100% confluency. 50 µl of 100 mM

CuSO4 was added to nearly 100% confluent cells in 75-cm^2 flask to induce

metallothionein promoter. The Fog-containing media was collected by centrifugation

and further concentrated using 3,000 MW concentrators. The presence of Fog was

confirmed by Western blot using the Myc-antibody (Invitrogen; RRID:AB_2533008).

Non-transfected S2 cells were used as a control.

Cell contractility assay was performed as described in (Manning et al., 2013). Smog- or

Mist-transfected S2 cells were induced with $CuSO_4$. Cells were then transferred to

Concanavalin A-coated coverslips and fixed using 10% paraformaldehyde in PBTr for

15 minutes at room temperature. After being blocked in 5% NGS in PBTr for 20 minutes

at RT, cells were stained with the V5 antibody (Invitrogen; RRID:AB_2556564) to

differentiate between transfected and non-transfected cells. Cell contractility of

transfected cells was monitored using a phase-contrast filter in a Leica DM2500

microscope using the 40X, 0.8 NA objective. Three independent experiments were

performed. Two-way ANOVA was performed to calculate statistical significance.

Acknowledgments

We thank the members of the Chung laboratory for their comments and suggestions. We

thank A. Martin, T. Lecuit and the Bloomington stock center for fly stocks, and D. J.

Andrew and the Developmental Studies Hybridoma Bank for antibodies. We thank

Flybase for the gene information. We are grateful to D. J. Andrew, C. D. Hanlon, J. Kim,

and J. Matthew for their helpful comments on the manuscript. This work is supported by

a start-up fund from Louisiana State University and grants from the Louisiana Board of

Regents Research Competitiveness Subprogram and the National Science Foundation

(MCB 2141387) to S.C.

Figure Legends

Figure 1. Smog transduces Fog signal to activate Rok and myosin in the SG. (A)

Cartoon diagram showing SG placodes (magenta) in the ventral side of a stage 11

Drosophila embryo. Cells that undergo apical constriction near the invagination pit

(asterisk) are shown in purple. In those cells, the Fog ligand (yellow) binds to its GPCR

(blue) to activate medioapical myosin. In all stage 11 SG images, anterior is to the left,

and dorsal is up. (B, B') Fluorescent in situ hybridization of *smog* (magenta) in the WT

SG. Arrowsheads, strong *smog* mRNA signals in apically constricting SG cells. Arrows,

smog signals near the grooves and the ventral midline. Green, CrebA (SG nuclei).

White lines, SG boundary. (C-D'') Smog-GFP (green) signals in control (C-C'') and Fog-

overexpressing SGs (D-D''). Magenta, E-Cad. Yellow arrowheads, puncta of Smog-GFP

signals in the apical region near AJs in control SG cells. Rred arrowheads, considerably

higher Smog-GFP signals in the entire apical domain of Fog-overexpressing SG cells.

Insets, magnified images of yellow boxed regions. (C'', D'') z-sections across the yellow

lines in C and D. (E) Quantification of total Smog-GFP intensity. n=4 SGs (control); 5

SGs (Fog overexpression). *p≤0.05 (Welch's t-test). (F-G''') sqh-GFP (green) and E-

Cad (magenta) signals in control (F-F''') and Fog-overexpressing (G-G''') SGs. (F'-G''')

Magnified images of yellow boxed regions in F and G. Yellow arrowheads, AJs. Cyan

arrowheads, sqh-GFP signals. (H-J) Quantification of medioapical myosin intensity (H),

junctional myosin intensity (I), and the ratio of medioapical to junctional myosin intensity

(J). ns, non-significant; *p≤0.05; **p≤0.01 (Welch's t-test). (K-M''') Rok-GFP (green) and

E-Cad (magenta) signals in SGs. K'-M''', magnified images of yellow boxed regions in

K-M. Yellow arrowheads, AJs. Cyan arrowheads, Rok-GFP puncta. (N, O)

Quantification of the area of Rok-GFP particles (N) and waviness of AJs (O). Asterisks,

invagination pit. n=5 SGs, 75 cells for each genotype in H-J, N, and O. **p≤0.01 (Mann-

Whitney U test).

Figure 2. Smog-transfected S2 cells contract in response to Fog. (A-F') S2 cells

cultured in the absence (A, C, E) or presence (B, D, F) of Fog. (A-B') Non-transfected.

(C-D') Mist-transfected. (E-F') Smog-transfected. A-F, phase contrast; A'-F', merged

images of phase contrast and V5 signals for transfected cells (green). White arrowhead

in E', a Smog-transfected S2 cell not contracting in the absence of Fog. Magenta

arrowheads, contracting cells. (G) Quantification of percentage of contracting cells

among the transfected cells. n=~100 cells per condition from three independent

experiments. ****p≤0.0001 (Two-way ANOVA with Tukey's multiple comparison post

hoc analysis).

Figure 3. *smog* knockdown and loss result in apical constriction defects in the SG

and disrupted epithelial morphology. (A-F') Confocal images of stage 11 SGs

stained for E-Cad (A-F) and corresponding heat maps of apical areas (A'-F').

TRiP.HMC03192 and *TRiP.GL01473* are used for zygotic and M/Z knockdown of *smog*,

respectively. Red and white lines, SG boundary. (G, H) Quantification of percentage

(left) and cumulative percentage (right) of cells with different apical areas. Mann-

Whitney U test (for percentage of cells) and Kolmogorov-Smirnov test (for cumulative

percentage of cells). n=8 SGs (control, 902 cells; *smog* RNAi (Z), 823 cells; *smog* RNAi

(M/Z), 881 cells; WT, 892 cells; *smog* null [mild], 859 cells); 6 SGs (*smog* null [severe],

672 cells). ***p<0.001; ****p<0.0001. (I-L) 3D reconstruction of confocal z-sections of

SGs (Imaris) for WT (I), M/Z knockdown of *smog* (J), *smog* null [mild] (K), and *smog* null

[severe] (L). Green, E-Cad. Magenta, CrebA. Asterisks, invagination pit. Arrows in J and

L, exaggerated grooves. Arrowheads in J and L, enlarged SG cells. (M, N) Tile scan

confocal images of stage 11 embryos in WT (M) and M/Z knockdown of *smog* (N).

Green, E-Cad. Magenta, CrebA. Arrows, the ventral midline. Arrowheads, SGs.

Figure 4. *smog* regulates all three pools of myosin in the SG in a Fog-dependent

and -independent manner. (A-D'') Confocal images of stage 11 SGs with sqh-GFP

(green) and E-Cad (magenta) signals. (A'-D'') Higher magnification of the yellow boxed

region in A-D. (E-H'') Confocal images of SGs with Rok-GFP (green) and E-Cad

(magenta) signals. (E'-H") Magnified images of the yellow boxed region in E-H. (I-M)

Quantification of medioapical myosin intensity (I), junctional myosin intensity (J), ratio of

medial and junctional myosin intensity (K), area of medioapical myosin particles (L), and

area of Rok-GFP particles (M). ns, non-significant; *p≤0.05; **p≤0.01 (Welch's t-test for

I-K; Mann-Whitney U test for L and M). n=5 SGs, 100 cells (I-L) and 5 SGs, 75 cells (M)

729 for each genotype. (N-P'') Confocal images of SGs with sqh-GFP (green) and E-Cad

730 (magenta) signals. Insets, magnified views of the SG boundary. Red arrowheads in N,

731 N'', O and O'', high myosin intensity at the SG boudary in control and *smog* null [mild]

732 embryos. Cyan arrowheads in P and P'', low myosin signals at the SG boundary in the

733 *smog* null [severe] embryo. (Q) Circularity of the SG placode boundary. An example of a

734 WT SG stained for E-Cad (magenta) and sqh-GFP (green) with the SG boundary (red

735 line) and the boundaries one cell row outside (gray line) and inside (blue line) the SG

736 boundary. (R) Quantification of circularity of the SG, outer, and inner boundaries.

737 Asterisks, invagination pit. n=5 SGs for each genotype. ns, non-significant; *p≤0.05;

738 **p≤0.01; ***p≤0.001 (Welch's t-test).

740 **Figure 5. Smog is required for maintaining epithelial integrity and the microtubule**

741 **networks.** (A-D') Confocal images of stage 11 embryos stained for E-Cad (green in A,

742 B), Crb (green in C, D) and CrebA (magenta). (A'-D') Magnified view of the white boxed

743 region in A-D. Arrowheads in B' and D', E-Cad (B') and Crb (D') signals are absent in

744 regions of the epidermis in *smog* mutants. (E) E-Cad (magenta) and Gap43-mCh

745 (green) signals in the epidermis in the *smog* null [severe] embryo. The cell membrane is

746 intact (cyan arrowheads in E'') in regions where E-Cad signals are absent (yellow

747 arrowheads in E'). (F-K) E-Cad (F-H) and Crb (I-K) signals in SGs in WT (F, I), *smog*

748 null [mild] (G, J), and *smog* null [severe] (H, K) embryos. Insets, magnified view of

749 yellow boxed region in the SG. Arrowheads, gaps in E-Cad or Crb signals. (L-O)

750 Quantification of the number of gaps in E-Cad (L) and Crb (N) signals and the ratio of

751 gap length to total junctional length for E-Cad (M) and Crb (O). n=5 SGs, 50 cells per

752 genotype. ns, non-significant; *p≤0.05; **p≤0.01 (Mann-Whitney U test). (P-W) Ace-tub

753 (P-S) and Tyr-tub (T-W) signals in the SG and the surrounding epithelia. Red dotted

754 lines, SG boundary. (X, Y) Quantification of the ratio of Ace-tub (X) and Tyr-tub (Y)

755 intensity in SG cells to the intensity in cells outside the SG placode. n=5 SGs for each

756 genotype. *p≤0.05; **p≤0.01 (Mann-Whitney U test). Asterisks, invagination pit; Pound

757 sign, dividing cells.

Figure 6. Smog is required for maintaining cortical actin networks during SG invagination. (A-B''') Rok-GFP (green) and E-Cad (magenta) in stage 11 SGs in control (A-A''') and *smog* null [severe] (B-B''') embryos. (A'-B'') Higher magnification of the white boxed regions in A and B. (A''', B''') x-z sections across the yellow lines in A and B. Red arrowheads, normal Rok-GFP signals accumulated in the medioapical region of control SG cells. White arrowheads, blebs in SG cells in *smog* mutants enriched with Rok-GFP signals. (C-J) SGs stained for phalloidin (C-F) and Dia (G-J). Compared to strong phalloidin and Dia signals in WT (red arrowheads in C, G), phalloidin and Dia levels are reduced in *smog* and *fog* mutants (cyan arrowheads in D-F, yellow arrowheads in H-J). (K) Quantification of total Dia intensity in the SG placode. n=5 SGs for control and *smog* null [mild], n=4 SGs for *smog* null [severe], n=6 SGs for *fog* null. *p≤0.05; **p≤0.01 (Mann-Whitney U test). (L-N'') Rok-GFP (green) and E-Cad (magenta) in *chic* RNAi in the wild-type (L-L''), *smog* null [mild] (M-M''), and *chic* RNAi in *smog* mutant background (N-N''). (L'-N'') Higher magnification of the white boxed regions in L-N. Red arrowheads in L', L'', M', and M'', normal Rok-GFP signals in *chic* RNAi and smog null [mild] SGs. White arrowheads in N' and N'', membrane blebs with enriched Rok-GFP signals in *smog* null [mild] SG cells with *chic* RNAi. Asterisks, invagination pit.

Figure 7. A proposed model for roles of Smog during epithelial tube formation. During SG invagination, Smog responds to Fog signal to promote Rok accumulation and medioapical myosin formation to control apical constriction (red arrows). Independent of Fog (and responding to an as-yet-unknown ligand X), Smog regulates epithelial integrity by regulating junctional and the supracellular myosin pools, the microtubule networks, and key apical/junctional components (gray arrows). Smog is also required for cortical actin organization during SG invagination, through regulating Dia levels (gray arrows). Different ligands may recruit specific G proteins and RhoGEFs to activate distinct downstream effectors.

References

Abreu-Blanco, M. T., Verboon, J. M. and Parkhurst, S. M. (2014). Coordination of Rho Family GTPase Activities to Orchestrate Cytoskeleton Responses during Cell Wound Repair. *Current Biology* **24**, 144–155.

Afshar, K., Stuart, B. and Wasserman, S. A. (2000). Functional analysis of the Drosophila diaphanous FH protein in early embryonic development. *Development (Cambridge, England)* **127**, 1887–97.

Andrew, D. J. and Ewald, A. J. (2010). Morphogenesis of epithelial tubes: Insights into tube formation, elongation, and elaboration. *Developmental Biology* **341**, 34–55.

Aoki, K., Maeda, F., Nagasako, T., Mochizuki, Y., Uchida, S. and Ikenouchi, J. (2016). A RhoA and Rnd3 cycle regulates actin reassembly during membrane blebbing. *Proceedings of the National Academy of Sciences* **113**, E1863–E1871.

Barrett, K., Leptin, M. and Settleman, J. (1997). The Rho GTPase and a Putative RhoGEF Mediate a Signaling Pathway for the Cell Shape Changes in Drosophila Gastrulation. *Cell* **91**, 905–915.

Benton, M. A., Frey, N., Nunes da Fonseca, R., von Levetzow, C., Stappert, D., Hakeemi, M. S., Conrads, K. H., Pechmann, M., Panfilio, K. A., Lynch, J. A., et al. (2019). Fog signaling has diverse roles in epithelial morphogenesis in insects. *eLife* **8**, e47346.

Blanchard, G. B., Murugesu, S., Adams, R. J., Martinez-Arias, A. and Gorfinkiel, N. (2010). Cytoskeletal dynamics and supracellular organisation of cell shape fluctuations during dorsal closure. *Development* **137**, 2743–2752.

Booth, A. J. R., Blanchard, G. B., Adams, R. J. and Röper, K. (2014). A Dynamic Microtubule Cytoskeleton Directs Medial Actomyosin Function during Tube Formation. *Developmental Cell* **29**, 562–576.

Charras, G. T. (2008). A short history of blebbing. *Journal of Microscopy* **231**, 466–478.

Charras, G. T., Yarrow, J. C., Horton, M. A., Mahadevan, L. and Mitchison, T. J. (2005). Non-equilibration of hydrostatic pressure in blebbing cells. *Nature* **435**, 365–369.

Chung, S., Hanlon, C. D. and Andrew, D. J. (2014). Building and specializing epithelial tubular organs: the Drosophila salivary gland as a model system for

revealing how epithelial organs are specified, form and specialize. *WIREs Dev Biol*

3, 281–300.

Chung, S., Kim, S. and Andrew, D. J. (2017). Uncoupling apical constriction from

tissue invagination. *eLife* **6**, e22235.

Cooley, L., Verheyen, E. and Ayers, K. (1992). chickadee encodes a profilin required

for intercellular cytoplasm transport during Drosophila oogenesis. *Cell* **69**, 173–184.

Costa, M., Wilson, E. T. and Wieschaus, E. (1994). A putative cell signal encoded by

the folded gastrulation gene coordinates cell shape changes during Drosophila

gastrulation. *Cell* **76**, 1075–1089.

Dawes-Hoang, R. E., Parmar, K. M., Christiansen, A. E., Phelps, C. B., Brand, A. H.

and Wieschaus, E. F. (2005). *folded gastrulation* , cell shape change and the

control of myosin localization. *Development* **132**, 4165–4178.

Fox, D. T. and Peifer, M. (2007). Abelson kinase (Abl) and RhoGEF2 regulate actin

organization during cell constriction in *Drosophila*. *Development* **134**, 567–578.

Garcia De Las Bayonas, A., Philippe, J. M., Lellouch, A. C. and Lecuit, T. (2019).

Distinct RhoGEFs Activate Apical and Junctional Contractility under Control of G

Proteins during Epithelial Morphogenesis. *Current Biology* **29**, 3370-3385.e7.

Girdler, G. C. and Röper, K. (2014). Controlling cell shape changes during salivary

gland tube formation in Drosophila. *Seminars in Cell & Developmental Biology* **31**,

74–81.

Goode, B. L. and Eck, M. J. (2007). Mechanism and Function of Formins in the Control

of Actin Assembly. *Annual Review of Biochemistry* **76**, 593–627.

Gorfinkiel, N. and Blanchard, G. B. (2011). Dynamics of actomyosin contractile

activity during epithelial morphogenesis. *Current Opinion in Cell Biology* **23**, 531–

539.

Guglielmi, G., Barry, J. D., Huber, W. and de Renzis, S. (2015). An Optogenetic

Method to Modulate Cell Contractility during Tissue Morphogenesis. *Developmental*

cell **35**, 646–660.

Häcker, U. and Perrimon, N. (1998). *DRhoGEF2* encodes a member of the Dbl family

of oncogenes and controls cell shape changes during gastrulation in *Drosophila*.

Genes & Development **12**, 274–284.

Henderson, K. D. and Andrew, D. J. (2000). Regulation and Function of Scr, exd, and

hth in the Drosophila Salivary Gland. *Developmental Biology* **217**, 362–374.

Homem, C. C. F. and Peifer, M. (2008). Diaphanous regulates myosin and adherens

junctions to control cell contractility and protrusive behavior during morphogenesis.

Development **135**, 1005–1018.

Izquierdo, E., Quinkler, T. and de Renzis, S. (2018). Guided morphogenesis through

optogenetic activation of Rho signalling during early Drosophila embryogenesis.

Nature Communications **9**, 2366.

Jha, A., van Zanten, T. S., Philippe, J.-M., Mayor, S. and Lecuit, T. (2018).

Quantitative Control of GPCR Organization and Signaling by Endocytosis in

Epithelial Morphogenesis. *Current Biology* **28**, 1570-1584.e6.

Jodoin, J. N., Coravos, J. S., Chanet, S., Vasquez, C. G., Tworoger, M., Kingston,

E. R., Perkins, L. A., Perrimon, N. and Martin, A. C. (2015). Stable Force Balance

between Epithelial Cells Arises from F-Actin Turnover. *Developmental Cell* **35**, 685–

697.

Kanesaki, T., Hirose, S., Grosshans, J. and Fuse, N. (2013). Heterotrimeric G protein

signaling governs the cortical stability during apical constriction in Drosophila

gastrulation. *Mechanisms of Development* **130**, 132–142.

Kerridge, S., Munjal, A., Philippe, J.-M., Jha, A., de las Bayonas, A. G., Saurin, A.

J. and Lecuit, T. (2016). Modular activation of Rho1 by GPCR signalling imparts

polarized myosin II activation during morphogenesis. *Nature Cell Biology* **18**, 261–

270.

Kim, H. Y., Varner, V. D. and Nelson, C. M. (2013). Apical constriction initiates new

bud formation during monopodial branching of the embryonic chicken lung.

Development **140**, 3146–3155.

Knirr, S., Azpiazu, N. and Frasch, M. (1999). The role of the NK-homeobox gene

slouch (S59) in somatic muscle patterning. *Development (Cambridge, England)* **126**,

4525–35.

Ko, C. S., Tserunyan, V. and Martin, A. C. (2019). Microtubules promote intercellular

contractile force transmission during tissue folding. *Journal of Cell Biology* **218**,

2726–2742.

Kölsch, V., Seher, T., Fernandez-Ballester, G. J., Serrano, L. and Leptin, M. (2007).

Control of *Drosophila* Gastrulation by Apical Localization of Adherens Junctions and

RhoGEF2. *Science* **315**, 384–386.

Le, T. P. and Chung, S. (2021). Regulation of apical constriction via microtubule- and

Rab11-dependent apical transport during tissue invagination. *Molecular Biology of*

the Cell **32**, 1033–1047.

le Droguen, P.-M., Claret, S., Guichet, A. and Brodu, V. (2015). Microtubule-

dependent apical restriction of recycling endosomes sustains adherens junctions

during morphogenesis of the *Drosophila* tracheal system. *Development* **142**, 363–

374.

Levayer, R. and Lecuit, T. (2012). Biomechanical regulation of contractility: spatial

control and dynamics. *Trends in cell biology* **22**, 61–81.

Lord, S. J., Velle, K. B., Mullins, R. D. and Fritz-Laylin, L. K. (2020). SuperPlots:

Communicating reproducibility and variability in cell biology. *Journal of Cell Biology*

219, e202001064.

Lubarsky, B. and Krasnow, M. A. (2003). Tube morphogenesis: making and shaping

biological tubes. *Cell* **112**, 19–28.

Manning, A. J. and Rogers, S. L. (2014). The Fog signaling pathway: Insights into

signaling in morphogenesis. *Developmental Biology* **394**, 6–14.

Manning, A. J., Peters, K. A., Peifer, M. and Rogers, S. L. (2013). Regulation of

Epithelial Morphogenesis by the G Protein–Coupled Receptor Mist and Its Ligand

Fog. *Science Signaling* **6**, ra98.

Martin, A. C. and Goldstein, B. (2014). Apical constriction: themes and variations on a

cellular mechanism driving morphogenesis. *Development* **141**, 1987–1998.

Martin, A. C., Kaschube, M. and Wieschaus, E. F. (2009). Pulsed contractions of an

actin–myosin network drive apical constriction. *Nature* **457**, 495–499.

Martin, A. C., Gelbart, M., Fernandez-Gonzalez, R., Kaschube, M. and Wieschaus,

E. F. (2010). Integration of contractile forces during tissue invagination. *Journal of*

Cell Biology **188**, 735–749.

Mason, F. M., Tworoger, M. and Martin, A. C. (2013). Apical domain polarization localizes actin–myosin activity to drive ratchet-like apical constriction. *Nature Cell Biology* **15**, 926–936.

Massarwa, R., Schejter, E. D. and Shilo, B.-Z. (2009). Apical Secretion in Epithelial Tubes of the Drosophila Embryo Is Directed by the Formin-Family Protein Diaphanous. *Developmental Cell* **16**, 877–888.

Munjal, A. and Lecuit, T. (2014). Actomyosin networks and tissue morphogenesis. *Development (Cambridge, England)* **141**, 1789–93.

Myat, M. M. and Andrew, D. J. (2000). Organ shape in the Drosophila salivary gland is controlled by regulated, sequential internalization of the primordia. *Development (Cambridge, England)* **127**, 679–91.

Nikolaidou, K. K. and Barrett, K. (2004). A Rho GTPase Signaling Pathway Is Used Reiteratively in Epithelial Folding and Potentially Selects the Outcome of Rho Activation. *Current Biology* **14**, 1822–1826.

Parks, S. and Wieschaus, E. (1991). The drosophila gastrulation gene concertina encodes a Gα-like protein. *Cell* **64**, 447–458.

Plageman, T. F., Chauhan, B. K., Yang, C., Jaudon, F., Shang, X., Zheng, Y., Lou, M., Debant, A., Hildebrand, J. D. and Lang, R. A. (2011). A Trio-RhoA-Shroom3 pathway is required for apical constriction and epithelial invagination. *Development* **138**, 5177–5188.

Rao, M. V. and Zaidel-Bar, R. (2016). Formin-mediated actin polymerization at cell–cell junctions stabilizes E-cadherin and maintains monolayer integrity during wound repair. *Molecular Biology of the Cell* **27**, 2844–2856.

Romero, S., le Clainche, C., Didry, D., Egile, C., Pantaloni, D. and Carlier, M.-F. (2004). Formin Is a Processive Motor that Requires Profilin to Accelerate Actin Assembly and Associated ATP Hydrolysis. *Cell* **119**, 419–429.

Röper, K. (2012). Anisotropy of Crumbs and aPKC Drives Myosin Cable Assembly during Tube Formation. *Developmental Cell* **23**, 939–953.

Rousso, T., Shewan, A. M., Mostov, K. E., Schejter, E. D. and Shilo, B.-Z. (2013). Apical targeting of the formin Diaphanous in Drosophila tubular epithelia. *eLife* **2**, 1–19.

Royou, A., Field, C., Sisson, J. C., Sullivan, W. and Karess, R. (2004). Reassessing
the Role and Dynamics of Nonmuscle Myosin II during Furrow Formation in Early
Drosophila Embryos. *Molecular Biology of the Cell* **15**, 838–850.

Sanchez-Corrales, Y. E., Blanchard, G. B. and Röper, K. (2018). Radially patterned
cell behaviours during tube budding from an epithelium. *eLife* **7**, e35717.

Shewan, A. M., Maddugoda, M., Kraemer, A., Stehbens, S. J., Verma, S., Kovacs,
E. M. and Yap, A. S. (2005). Myosin 2 Is a Key Rho Kinase Target Necessary for
the Local Concentration of E-Cadherin at Cell–Cell Contacts. *Molecular Biology of*
the Cell **16**, 4531–4542.

Sidor, C., Stevens, T. J., Jin, L., Boulanger, J. and Röper, K. (2020). Rho-Kinase
Planar Polarization at Tissue Boundaries Depends on Phospho-regulation of
Membrane Residence Time. *Developmental Cell* **52**, 364-378.e7.

Sweeton, D., Parks, S., Costa, M. and Wieschaus, E. (1991). Gastrulation in
Drosophila: the formation of the ventral furrow and posterior midgut invaginations.
Development (Cambridge, England) **112**, 775–89.

Vázquez-Victorio, G., González-Espinosa, C., Espinosa-Riquer, Z. P. and Macías-
Silva, M. (2016). GPCRs and actin–cytoskeleton dynamics. *Methods in Cell Biology*
132, 165–188.

Weng, M. and Wieschaus, E. (2016). Myosin-dependent remodeling of adherens
junctions protects junctions from Snail-dependent disassembly. *Journal of Cell*
Biology **212**, 219–229.

Westermann, S. and Weber, K. (2003). Post-translational modifications regulate
microtubule function. *Nature Reviews Molecular Cell Biology* **4**, 938–948.

Zweifel, M. E. and Courtemanche, N. (2020). Profilin's Affinity for Formin Regulates
the Availability of Filament Ends for Actin Monomer Binding. *Journal of Molecular*
Biology **432**, 166688.

Figure 1.

Figure 2.

Phase contrast

S2 cells (n.t)
-Fog   +Fog

S2-Mist-V5
-Fog   +Fog

S2-Smog-V5
-Fog   +Fog

10 µm

A   B   C   D   E   F

V5

A'   B'   C'   D'   E'   F'

G

% of contracted cells

ns   ****   ****   ****   ****

n.t.   S2-Mist   S2-Smog

■ -Fog   □ +Fog

Figure 3.

A Control **B** *smog RNAi (Z)* **C** *smog RNAi (M/Z)* **D** WT **E** *smog null [mild, 73.8%]* **F** *smog null [severe, 26.2%]*

E-Cad

5 µm

Apical area (µm²)
30
0

G Percentage of cells — Control, *smog RNAi (Z)*, *smog RNAi (M/Z)* — Apical area (µm²)
Cumulative percentage of cells — Control, *smog RNAi (Z)*, *smog RNAi (M/Z)* — Apical area (µm²)

H Percentage of cells — WT, *smog null [mild]*, *smog null [severe]* — Apical area (µm²)
Cumulative percentage of cells — WT, *smog null [mild]*, *smog null [severe]* — Apical area (µm²)

I WT **J** *smog RNAi (M/Z)* **K** *smog null [mild]* **L** *smog null [severe]* **M** WT **N** *smog RNAi (M/Z)*

E-Cad CrebA

20 µm

50 µm

Figure 4.

Figure 5.

Figure 6.

Figure 7.