1   **Title: Receptor interacting protein kinase-3 mediates both myopathy and cardiomyopathy in**

2   **preclinical animal models of Duchenne muscular dystrophy**

4   Maximilien Bencze[1] ✉ , Baptiste Periou[1] , Isabel Punzón[1], Inès Barthélémy[1], Valentina Taglietti[1],

5   Cyrielle Hou[1], Louai Zaidan[1], Kaouthar Kefi[1], Stéphane Blot[1], Onnik Agbulut[2], Marianne Gervais[1],

6   Geneviève Derumeaux[3], Laurent Tiret[1], François-Jérôme Authier[1], Fréderic Relaix[1]

[1] U955-IMRB, Inserm, UPEC, ENVA, EFS, Team 10, Biology of the Neuromuscular System, Créteil

94000, France.

[2] Sorbonne Université, Institut de Biologie Paris-Seine (IBPS), CNRS UMR 8256, Inserm ERL U1164,

Biological Adaptation and Ageing, 75005, Paris-France.

[3] U955-IMRB, Université Paris-Est Créteil (UPEC), Team 8, Department of Physiology, Henri Mondor

Hospital, FHU-SENEC, AP-HP, Créteil, France.

✉ **Correspondence:** Maximilien Bencze: <u>maximilien.bencze@inserm.fr, m.bencze@ucl.ac.uk</u>

Keywords: Duchenne muscular dystrophy, myonecrosis, programmed cell death, necroptosis,

fibrosis, myogenesis, cardiac failure, animal model.

Abstract

Background

Duchenne muscular dystrophy (DMD) is a progressive muscle degenerative disorder, culminating in a

complete loss of ambulation, hypertrophic cardiomyopathy and a fatal cardiorespiratory failure.

Necroptosis is the form of necrosis that is dependent upon the receptor-interacting protein kinase (RIPK)

3; it is involved in several inflammatory and neurodegenerative conditions. We previously identified

RIPK3 as a key player in the acute myonecrosis affecting the hindlimb muscles of the dystrophic mouse

model, mdx. Whether necroptosis also mediates respiratory and heart disorders in DMD is currently

unknown.

Methods

Evidence of activation of the necroptotic axis was examined in dystrophic tissues from Golden retriever

muscular dystrophy (GRMD) dogs and R-DMDdel52 rats. A functional assessment of the involvement

of necroptosis in dystrophic animals was performed on mdx mice that were genetically depleted for

RIPK3. Dystrophic mice aged from 12 to 18 months were analyzed by histology and molecular biology

to compare the phenotype of muscles from mdx$Ripk3^{+/+}$and mdx$Ripk3^{-/-}$ mice. Heart function was also

examined by echocardiography in 40-week-old mice.

Results

Quantification of *RIPK3* transcripts in sartorius and biceps femoris muscles from GRMD dogs positively

correlated to myonecrosis levels (r=0.81; p=0.0076). *RIPK3* was also found elevated in the diaphragm

(p=0<0.05). In the slow progressing heart phenotype of GRMD dogs, the phosphorylated form of RIPK1

at the Serine 161 site was dramatically increased in cardiomyocytes. A similar p-RIPK1 upregulation

characterized the cardiomyocytes of R-DMDdel52 rats, associated with a marked overexpression *of*

Ripk1 (p=0.007) and *Ripk3* (p=0.008), indicating primed activation of the necroptotic pathway in the

dystrophic heart. Mdx$Ripk3^{-/-}$ mice displayed decreased compensatory hypertrophy of the heart

(p=0.014), and echocardiography showed a 19% increase in the relative wall thickness (p<0.05) and

29% reduction in the left ventricle mass (p=0.0144). Besides, mdx$Ripk3^{-/-}$ mice presented no evidence

of a regenerative default or sarcopenia in skeletal muscles, moreover around 50% less affected by

fibrosis ($p < 0.05$).

51   **Conclusions**

52   Our data provide evidence of the activation of the necroptotic pathway in degenerative tissues from

53   dystrophic animal models, including the diaphragm and the heart. The genetic inhibition of necroptosis

54   in dystrophic mice improves both cardiac function and histological features of muscles, suggesting that

55   prevention of necroptosis is susceptible to providing multiorgan beneficial effects for DMD.

Introduction

Duchenne muscular dystrophy (DMD) is an X-linked genetic disorder, affecting 1/5000 male birth worldwide. Loss-of-function mutations in the dystrophin gene primarily lead to chronic myofibre necrosis (myonecrosis) and severe locomotor muscle disability starting in early childhood. Notably, the absence of dystrophin at the sarcolemma renders myofibre more susceptible to mechanical stress, which favours contraction-induced myonecrosis. Following injury, muscle repair requires myogenesis thanks to the contribution of PAX7-expressing resident muscle stem cells (MuSC)[1]. However, in disabling neuromuscular disorders involving chronic myofiber degeneration such as DMD, regeneration does not fully compensate for chronic myofibre demise. The loss of a sustained homeostatic balance between the two mechanisms leads to the increasing replacement of contractile myofibres by fibrotic or adipose tissue. Biopsies with interstitial fibrosis positively correlate with a poor outcome in DMD boys [2]. Thus, identifying the molecular pathways involved in fibrosis formation is relevant for improving therapeutic perspectives. Dystrophin deficiency also challenges the survival of respiratory and cardiac muscle cells, leading to the premature death of DMD patients from cardiorespiratory weakness or failure [3,4].

Recently, we reported that necroptosis, a receptor-interacting protein kinase (RIPK)1- RIPK3- and mixed lineage domain kinase-like (MLKL)-dependent cell death pathway, is activated in the locomotor musculature of murine dystrophin-deficient muscles. Necroptosis inhibition dampens the peak of myonecrosis in the locomotor muscles of three-week-old mdx mice and improves muscle function in young adults [5]. However, two recent studies reported a link between necroptosis inhibition and a deficit in early myogenesis and myofibre atrophy[6,7]. One of them reported that RIPK3 depletion in mdx MuSCs dramatically impairs muscle function. Repressing necroptosis might therefore lead to two putative conflictual outcomes: a decrease in myofibre degeneration, together with a default in muscle regeneration leading to muscle atrophy and sarcopenia. Consequently, the long-term effects of necroptosis inhibition in dystrophic skeletal and heart muscle remodelling need to be determined, before further envisaging the necroptotic pathway as a putative therapeutic target for DMD.

To this end, we examined the diaphragm and heart for evidence of necroptosis activation from Golden retriever muscular dystrophy (GRMD) dogs. The functional assessment of necroptosis in the

cardiorespiratory system was achieved using mdx*Ripk3*$^{-/-}$ mice. Compared to mdx*Ripk3*$^{+/+}$, locomotor

muscles from mdx*Ripk3*$^{-/-}$ mice displayed a significant reduction of fibrosis. Despite its involvement in

early myogenesis, RIPK3 deficiency did not generate sarcopenia. In the diaphragm, fibrosis reduction

indicates that necroptosis not only participates in the pathology of limbs, but also of respiratory muscles.

Importantly, we found that RIPK3 depletion improves cardiomyopathy by reducing heart hypertrophy

and fibrosis deposition. Together, our data demonstrate that in dystrophic mdx mice, RIPK3 is

dispensable for post-lesional myogenesis and suggests that necroptosis inhibition is beneficial on the

hindlimb, respiratory, and heart muscles in dystrophinopathies. This work opens new avenues for further

research exploring the improvement of patients' health status by reducing necroptosis.

Material and methods

95   **Animals**

Mice and rats were bred in the Institut Mondor de Recherche biomédicale, Créteil, France in a pathogen-

free facility with 12-h light and 12-h dark cycles in accordance with European Directive 2010/63/EU.

Only male mice and rats were used for the experiments. Both wild-type (WT) and DMD rats and mdx

mice were born from the same litter. The GRMD dogs included in this study were housed in the facilities

of the neurobiology laboratory of the Veterinary School of Alfort. Experiments were approved by the

Anses/EnvA/Upec Ethics Committee. All care and manipulations were performed by national and

European legislation on animal experimentation. *Ripk3*$^{-/+}$ mice were kindly provided by Genentech (San

Francisco-CA). mdx*Ripk3*$^{-/-}$ mice were generated by crossing *Ripk3*$^{-/-}$ with mdx mice. The resulting mice

were backcrossed for at least four generations to obtain final mdx*Ripk3*$^{+/-}$ breeders.

106   **Treadmill tests**

107   The running capacity was evaluated on a treadmill (Tecmachine, Medical Development) at a +5% slope.

108   Males of 70- to 76-week-old mice were initially acclimatized to the treadmill for 5 days (10 min/day

109   and 12 m/min). Mice were challenged to an exercise of progressively increased intensity. Speed was

110   increased every 90- to 120 seconds/min and the speed of the last completed running step before

111   exhaustion was considered as the individual maximal aerobic speed.

112
113   **RNA extraction, PCR, and real-time PCR**

Total skeletal muscle RNA was extracted from muscle and heart samples using TRIzol (Thermo Fisher

Scientific) following the manufacturer's instructions. SuperScript III Reverse Transcriptase from the

Invitrogen kit converted RNA into cDNA using the Veriti 96- Well Fast Thermal Cycler (Applied

Biosystems). Gene expression was quantified by real-time qPCR with the StepOnePlus real-time PCR

system (Applied Biosystems) using SYBR Green detection tools (Applied Biosystems). In skeletal

muscle tissue, the expression of each gene was normalized to *Tbp* gene expression. Results are reported

as relative gene expression (2-DDCT). In the heart, from 500 ng of extracted RNA, the first-

strand cDNA was then synthesized using a RevertAid First Strand cDNA Synthesis Kit (Thermo

Fisher Scientific) with random hexamers according to the manufacturer's instructions. Using the Light

Cycler® 480 system (Roche Diagnostics), the reaction was carried out in duplicate for each sample in a

6-µl reaction volume containing 3 µl of SYBR Green Master Mix, 500 nM of the forward and reverse

primers each and 3 µl of diluted (1:25) cDNA. The thermal profile for the SYBR Green qPCR was 95°C

for 8 min, followed by 40 cycles at 95°C for 15 s, 60°C for 15 s and 72°C for 30 s. The mean gene

expression was normalized to *Hprt1* and *Rpl32*. In dogs, RNA was extracted from frozen muscle slides

using the NucleoSpin RNA extraction kit (Macherey Nagel). RNA was quantified by Nanodrop and the

Maxima first strand cDNA master kit (Thermo Fisher) was used to convert 300ng of RNA into

cDNA. Gene expression was quantified by real-time qPCR with the QuantStudio™ 3 Real-Time PCR

System (Applied Biosystems) using SYBR Green detection kit (Thermo Fisher). The reaction was

carried out in triplicates for each sample of diluted (1:10) cDNA. The expression of genes was

normalized to *b-actin* gene expression. Results are reported as relative gene expression (2^{-DDCT}).

Primers used: Mouse: forward *Ripk3* 5′-CGGGCACACCACAGAACAT-3′, *Ripk3* and reverse 5′-

GTAGCACATCCCCAGCACCAC-3′,   forward *Ripk1* 5′-AGAAGAAGGGAACTATTCGC-3′ and

reverse *Ripk1* 5′-TTCTATGGCCTCCACGAT-3', forward *Bnp* 5'-CAGCTCTTGAAGGACCAAGG-

3',   reverse   5'-AGACCCAGGCAGAGTCAGAA-3', forward   *Myh7*   5'-

AGGTGTGCTCTCCAGAATGG-3', reverse 5'-CAGCGGCTTGATCTTGAAGT-3', *Rpl32* - forward

5'-TGGTGAAGCCCAAGATCGTC-3', reverse 5'-GGATCTGGCCCTTGAACCTT-3', *Hprt1* -

forward 5'-CAGGCCAGACTTTGTTGGAT-3', reverse 5'-TTGCGCTCATCTTAGGCTTT-3'. Rat:

forward *Ripk3* 5'-CGTACACGTAGTCCCACTCG-3', reverse 5'- AGGGAGGTGAAGGCTATGGT-

3', forward *Ripk1* 5'- GGTCTCCCATGACCCCTTTG-3', reverse 5'-

GGTAGGTTGGTCTCAGGCAC-3'. Dog: Forward *Ripk1* 5'-GAATGAGTTCAGCCCTGCTC-3',

reverse 5'-CTCGCTCATAGTCGTGGTCA-3', forward *Ripk3* 5'-CAGAGAGGCTCAAGGTCAGG-

3', reverse 5'-CGATGTCTGGGCCACTATCT-3' *b-actin* forward 5'-

CCATCTACGAGGGGTACGCCC-3' reverse TGCTCGAAGTCC AGGGCGACGTA

148   **Histology**

Fibrosis was labelled using Epredia Varistain Gemini ES Automated Slide Stainer. For

immunofluorescence: rat antibody to CD68 (clone FA-11, 137001, 1/50), rabbit antibody to mouse pan-

Laminin (Sigma, L9393, 1/1000), Phospho-RIPK1 (Ser161) antibody (1/100 Invitrogen 105640).

Biopsies from GRMD dogs were processed and analysed as previously described[8]. The dried sections

were stained for 10 minutes in Hematein and 5 minutes in 1% Eosin, dehydrated in four consecutive

baths of ethanol, one bath of xylene and mounted in Canada balsam. One entire section per biopsy was

photographed using an AxioObserver Z1 linked to its ICC1 camera (Zeiss), and the MosaiX application

of the software AxioVision (Zeiss). The image was then analysed using the software Visilog 6.4

(Noesis). A grid of 10 000 mm2 squares was superimposed onto the image of the entire section. At each

intercept of the grid (i.e. every 100 mm) the histological aspect of the underlying tissue was manually

captioned using predefined annotations. The percentage of each type of histological event was

calculated, defined as the percentage of events not corresponding to normal shape fibres. Inflammatory

infiltration in dog biopsies was quantified using CD4, CD8, and CD11b immunostainings. 7μm sections

from each biopsy were fixed in cold acetone-methanol. After having blocked the endogenous peroxidase

activity, the primary antibody (either a rat anti-canine CD4, Serotec H, 1/50, or a rat anti-canine CD8,

Serotec H, 1/50, or a mouse anti-canine CD11b, Serotec H, 1/50) was used. The amount of inflammatory

cells was normalized to the area of the section[8].

Myonecrosis levels in mdx biopsies were determined by measuring the area corresponding to mouse

IgG uptake in myofibres (and not in interstitial areas) and expressed as the percentage of the cross-

sectional area. Muscle fibre minimal Feret diameter, corresponding to the minimum distance between

the two parallel tangents of myofibres, was determined. Images analyses were performed with ImageJ

by a specific self-developed macro that recognizes muscle fibres as previously described [9].

171
172   **Cardiac Phenotyping**

Mice were trained to be grasped since transthoracic echocardiography (TTE) was performed in

conscious, non-sedated mice to avoid any cardiac depressor effect of sedative or anaesthetic agents. Data

acquisition was performed every month by a single operator (JT) as previously described [10]. Images

were acquired from a parasternal position at the level of the papillary muscles using a 13-MHz linear-

array transducer with a digital ultrasound system (Vivid 7, GE Medical System, Horton, Norway). Left

ventricular diameters and ejection fraction, anterior and posterior wall thicknesses were serially obtained

from M-mode acquisition. Relative LV wall thickness (RWT) was defined as the sum of septal and

posterior wall thickness over LV end-diastolic diameter, and LV mass was determined using the

uncorrected cube assumption formula (LV mass=$(IVSd+LVIDd+PWTd)^3-LVIDd^3$). Peak systolic

values of radial SR in the anterior and posterior wall were obtained using Tissue Doppler Imaging (TDI)

as previously described [10]. TDI loops were acquired from the same parasternal view at a mean frame

rate of 514 fps and a depth of 1 cm. The Nyquist velocity limit was set at 12cm/s. Radial SR analysis

was performed offline using the EchoPac Software (GE Medical Systems) by an observer (GD) blinded

to the diet of the animals. The Peak systolic of radial SR was computed from a region of interest

positioned in the mid anterior wall and was measured over an axial distance of 0.6 mm. The temporal

smoothing filters were turned off for all measurements. Because slight respiratory variations exist, we

averaged peak systolic of radial SR on 8 consecutive cardiac cycles.

Results

Firstly, we addressed the relevance of necroptosis in dystrophin deficiency as a potential cell demise

involved in locomotor and respiratory muscles and the heart. Because heart or diaphragm muscle

biopsies cannot be performed in patients, we aimed at examining tissues in different animal models for

DMD. We first investigated degenerating tissues from golden retriever muscular dystrophy (GRMD)

195 dogs that represent a gold standard relevant animal model for DMD[11]. Compared to age-matched wild-

196 type (WT) dogs, 6-month-old-GRMD dogs were characterized by typical high and variable blood

197 creatine kinase levels (p=0.0167) (Figure 1a), assessing a significant ongoing, global myofibre demise.

198 Histological features of myonecrosis in the sartorius and biceps femoris muscles confirmed the presence

199 of recently injured area (Figure 1b). Because myofibre necroptosis comes with an increase of *Ripk3*

200 transcripts in mdx TA muscles, and that RIPK3 upregulation sensitizes muscle cells to TNF-elicited

201 necroptosis[12], we examined whether there was a correlation between muscles presenting high

202 histological necrotic extent and *RIPK3* expression in muscles of GRMD dogs. Sartorius muscles from

203 6- and 12-month-old GRMD dogs were together quantified by qPCR for *RIPK3* transcripts and

204 quantified for evidence of necrotic events and inflammation in hematoxylin & eosin staining.

205 Interestingly, *RIPK3* expression was highly correlated with the extent of myonecrosis (Pearson test,

206 p=0.0076, r=0.8138, n=9 GRMD muscles) (Figure 1b), but not with the inflammatory area (p=0.1409)

207 (Figure 1c), suggesting that *RIPK3* elevation is specific to necrotic demise, rather than post-necrotic

208 events such as myeloid infiltrate. Similarly, *RIPK3* transcripts increased in the GRMD diaphragm

209 compared to age-matched WT dogs, and *RIPK3* expression was also positively correlated with *RIPK3*

210 expression in these muscles (p=0.0307) (Figure 1d-f). Thus, RIPK3-dependant cell death does not only

211 involve ambulatory, but also the main respiratory musculature in the DMD pathogenic course. In the

212 heart, whilst we observed no significant increase in the necrosome transcripts in the left ventricle of 6-

213 month-old GRMD dogs (Figure 1g,h), we found a dramatic increase of immunoreactivity of GRMD

214 dog's cardiomyocytes to antibodies directed against phospho-RIPK1 (Ser161), indicating that RIPK1 is

215 primed for either extrinsic apoptosis or canonical necroptosis (Figure 1i)[13,14].

216 Thus, to determine whether necroptosis has a functional role in progressive cardiomyopathy or in

217 myonecrosis affecting the diaphragm of dystrophin-deficient mice, we generated mdx*Ripk3*[-/-] mice by

218 crossing mdx with *Ripk*3-KO mice, which compromise dystrophic mice for RIPK3-dependent

219 necroptosis[15]. Mdx*Ripk3*[+/+] (hereafter named mdx*Ripk3*[+/+] or mdx) and mdx*Ripk3*[-/-] littermates were

220 kept for ageing to reach the maximum mdx pathogenic state in striated muscles. We first examined

221 whether the long-term genetic inhibition of necroptosis would promote sarcopenia by repressing MuSCs

222 density and the regenerative capacity of skeletal muscle. Indeed, a recent article indicated that

necroptosis inhibition generates a MuSCs depletion by inducing the death of PAX7-positive cells, with

a subsequent deficit in myogenesis associated with a dramatically reduced muscle strength in adult mdx

mice [6]. In biceps from 18-month-old mdx*Ripk3*$^{-/-}$ mice, *Ripk1* transcripts were downregulated compared

to *Ripk3*$^{+/+}$ littermates (Figure 2a), indicating that canonical necroptosis is repressed in mdx*Ripk3*$^{-/-}$

mice. To determine whether the defect of necroptosis signalling compromises muscle function, mdx and

mdx*Ripk3*$^{-/-}$ littermates underwent forced treadmill exercise with progressive increase of speed, until

exhaustion. Control mdx mice were characterized by an important variability in running performance,

expressed in the running distance (Figure 2b) or the maximum aerobic speed before exhaustion

(Figure 2c). mdx*Ripk3*$^{-/-}$ mice showed more homogenous capacities (coefficient of variation:

mdx: 20,99% *versus* mdx*Ripk3*$^{-/-}$ 12.66%). On average, mdx*Ripk3*$^{-/-}$ ran ~ 10% longer than mdx

littermates, despite no statistical difference likely due to the variability of the control mdx phenotype

(p=0.5530). The putative atrophy of *Ripk3*-/- dystrophic muscles was examined using muscle mass

normalized to the individual size of the animals, assessed by the tibial length (TL). TA, EDL and biceps

muscle mass remained unchanged in aged mdx*Ripk3*$^{-/-}$ compared to mdx*Ripk3*$^{+/+}$ mice (Figure 2d).

Although young mdx muscles show extensive necrotic events leading to the accumulation of fibrotic

tissue, muscles from old mice are typically spared. In the rare post-necrotic areas in TAs from old mdx

and mdx*Ripk3*$^{-/-}$ mice, we found an equal density of Ki67-positive cells (Figure 2e,f), suggesting that

necroptosis inhibition does not affect MuSC proliferation in mdx muscle at this age. The mean of pooled

minimum Feret of TA myofibres (Figure 2g), as well as the relative distribution of myofibre size (Figure

2h), were strictly unchanged. Interestingly, we found that RIPK3 depletion reduced fibrosis in the TA

(p=0.0252) (Figure 2i), suggesting a cumulative cytoprotective effect regarding ancient degenerative

events. To further examine any possible evidence of regenerative default in mdx*Ripk3*$^{-/-}$ muscles, we

analysed biceps muscles for *Pax7* (Figure 2j) and *myogenin* (Figure 2k) expression but observed no

difference. Of note, we found a significant downregulation of *myostatin* expression (p=0.0082), a

biomarker of neuromuscular disorders while *follistatin* was not significantly repressed (p=0.4452)

(Figure 2l,m).

We then examined whether the pathogenic features of the mdx diaphragm are improved by RIPK3

depletion (Figure 3). In histology, the localization of blood proteins such as albumin or immunoglobulins

within sarcoplasm is a reliable marker for the necrotic fate of myofibre[16,17]. Muscles from eighteen-

month-old mice showed IgG immunoreactivity almost only localized at the extracellular compartment

(Figure 3a), and rarely within myofibres, indicating no/little ongoing myonecrosis. No difference was

found in both genotypes (unpaired t-test, p=0.1914). The density of myofibres with leaky sarcolemma

was extremely low in both genotypes (2.3 fibres/mm^2 in mdx versus 2.6 fibres/mm^2 in mdx$Ripk3^{-/-}$)

(Figure 3b,c). This was neither associated with high-density cell pockets (Figure 3d) nor with CD68-

positive cell infiltrates (Figure 3e). Furthermore, only a low percentage of centrally nucleated fibres was

found in the mdx diaphragm (5-10% of myofibres) (Figure 3f). These observations suggested that the

diaphragm from 18-month-old mdx mice stand at relative homeostasis with little/no ongoing

myonecrosis, regardless of RIPK3 expression. Examining the profile of myofibre size, we observed a

moderate decrease of myofibre size in mdx$Ripk3^{-/-}$ diaphragm compared to mdx littermates (16.6 in mdx

versus 16.1 in mdx$Ripk3^{-/-}$, p<0.0001, Mann Whitney test, n>16,000 myofibres in each group, included

in six mdx muscles, and eight mdx$Ripk3^{-/-}$ muscles). Dystrophin-deficient muscles are characterized by

an increased heterogeneity in myofibre size, illustrated by an increase of the variance coefficient[18].

RIPK3-deficiency decreased the mean-variance coefficient of myofibres in mdx mice (Figure 3i,

unpaired t-test p=0.0431). Fibrosis deposition in respiratory muscles, such as the diaphragm, is a key

pathogenic event in DMD pathogenesis and correlates with a bad prognosis in patients. To determine

whether RIPK3 also mediates fibrosis deposition in the diaphragm, muscle sections were stained with

Sirius red (Figure 3k), and we observed less collagen deposition in mdx$Ripk3^{-/-}$diaphragms compared to

mdx littermates (Figure 3k, unpaired t-test, p=0.0414). Together, our data demonstrate that RIPK3

participates in diaphragm pathogenesis in mdx mice by promoting interstitial fibrosis.

We next addressed the role of necroptosis in the development of cardiomyopathy upon dystrophin

deficiency. Importantly, end-stage cardiomyopathy is a major concern in the survival of boys and is

associated with hallmarks of cardiomyoblast necrosis and fibrosis deposition. In GRMD dogs, RIPK1

was found phosphorylated at the Ser161 site in the heart (Figure 1i), indicating an early step of activation

of RIPK1-dependant apoptosis or necroptosis[14]. Similar p-RIPK1 immunoreactivity to GRMD dogs was

observed in the hearts of R-DMDdel52 rats (Figure 4a) that develop an early cardiomyopathy[19].

Dystrophin-deficient rats also had a significant increase of *Ripk1* (p=0.0070, t-test) and *Ripk3*

(p=0.0080, t-test) transcripts in heart tissue compared to WT rats (Figure 4b,c), suggesting advanced

necroptosis activation in the heart from R-DMDdel52 rats. This contrasted with dogs that displayed no

significant elevation of *RIPK1* and *RIPK3* transcription (Figure 1g,h), in agreement with rare fresh

necrotic events observed in dystrophic hearts of GRMD dogs. To determine the role of necroptosis in

the progressive cardiomyopathy of dystrophin-deficient hearts, we analysed the heart phenotype of 18-

month-old mdx*Ripk3*[-/-] mice (Figure 4 d-k). RIPK3 depletion decreased cardiac compensatory

hypertrophy in mdx mice (Figure 4d,e). At the investigated age, we found no significant levels of

ongoing degeneration in the heart, displaying little CD68-positive cell infiltration whatever the genotype

(Figure 3f), which suggested an accumulation of past degenerative events with little evolution over time

at this age. Interstitial collagen deposition was reduced by almost 50% in the right ventricle of mdx*Ripk3*[-

[/-] mice compared to their mdx control littermates (Figure 4h,g). Furthermore, the hearts of mdx*Ripk3*[-/-]

mice showed significantly reduced levels of *Bnp* and *Myf7* transcripts, two sensitive markers of cardiac

hypertrophy (Figure 4i,j).

We thus examined whether the improvement of the deleterious phenotype of cardiomyopathy conferred

by RIPK3 depletion was validated at the 2D and functional levels and quantified routine

echocardiographic parameters using the M(otion)-mode (Figure 5). Compared to wild-type mice, aged

mdx mice are generally characterized by eccentric hypertrophy characterized by a transiently increased

LV mass with increased left ventricle (LV) end-diastolic and end-systolic diameter. This is followed by

a decompensated dilated cardiomyopathy associated with a reduction in LV fractional shortening and a

decreased LV mass[20]. We confirmed that our 18-month-old mdx mice suffered from eccentric

hypertrophy with an increased LV mass, associated with a reduced LV internal diameter in diastole

(LVIDd) and systole (LVIDs) (Figure 5a-4c and Supplementary Table 1). Of note, compared to mdx

mice, mdx*Ripk3*[-/-] littermates displayed a 30% reduction of the LV mass linked to a decrease in LVIDd

that contributed to the restoration of the LV relative wall thickness value (Figure 5d), and unchanged

fractional shortening of the LV, strain rate of cardiomyocytes from the anterior wall (Figures 5e, f) and

heart rate (Supplementary Table 1). Altogether, these data suggested that in the mdx model, RIPK3-

dependent necroptosis participates in the long-term development of cardiac hypertrophy and dysfunction

and that in these conditions, RIPK3 deficiency has overall cytoprotective effects against dystrophic-

associated cardiac deleterious phenotype in the mouse.

309   **Discussion**

Necroptosis is now considered a new therapeutic target for several degenerating disorders, including

cardiovascular neurodegenerative, and autoimmune diseases[21]. Dystrophin absence and DUX4

overexpression can lead to myofibre necroptosis *in vivo*[12]. Translating these discoveries into the clinic

for neuromuscular disorders requires an understanding of 1-the long-term consequences of necroptosis

inhibition in dystrophic conditions and 2-the role of necroptosis in the DMD cardiorespiratory

phenotype.

While locomotor tissues can be harvested in DMD patients for diagnostic purposes and could be used

for research purposes, respiratory or heart biopsies were not available. We then used dystrophic dogs

and rats which are highly relevant animal models for DMD and reproduce the main aspects of DMD

pathogenesis[11]. RIPK3 overexpression is a biomarker for myofibre necroptosis in mdx mice, and

sensitizes myoblasts to TNF-elicited necroptosis[5]. Using sartorius and femoral biceps, diaphragm and

cardiac tissues from 6-month-old GRMD dogs, we first assessed different levels of activation of the

early steps of the RIPK1-RIPK3 canonical necroptotic pathway. RIPK3 upregulation in limb muscles,

and diaphragm, and its correlation with the myonecrosis extent (Figure 1b), suggest a role for RIPK3 in

the myonecrosis affecting GRMD dogs, especially in cells expressing p-RIPK1, which prime cells to

programmed necrosis[14]. RIPK3 typically recruits MLKL during final necroptosis induction, and p-

MLKL causes membrane permeability and necrotic morphology. Interestingly, dogs and other species

belonging to the *Carnivora* order express crucial proteins required for necroptosis activation such as

RIPK1, RIPK3, DNA-dependant activator of interferon regulatory factors (DAI) and TIR-domain-

containing adaptor-inducing IFNβ (TRIF), however, they do not express MLK[23]. The molecular

mechanisms of cell death execution in RIPK3-dependant but MLKL-independent necrosis, involvement

remains an open question.

Notably, necrotic demise affecting the diaphragm and heart are key events leading to the short life

expectancy of DMD boys. Herein, we provide evidence that beyond its role in the degeneration of

locomotor muscles, necroptosis also takes part in the respiratory and cardiac phenotypes in the mdx

model of DMD, notably by promoting tissue fibrosis. Accumulation of extracellular matrix deposition

following myonecrosis is a central pathogenic process responsible for the loss of function of dystrophic

muscles, by progressively replacing functional muscle tissue in both the myocardium and skeletal

muscles[2]. Therefore, understanding the molecular pathways directly or indirectly leading to myofibre or

cardiomyocyte death or fibrosis is highly relevant.

In dystrophin-deficient mice, investigating the phenotypes of the diaphragm and heart requires the

generation of dystrophic and necroptosis-incompetent mice that would reach sufficient age to

progressively develop a significant phenotype[24,25]. Compared to young dystrophic TA, the diaphragm

from 18-month-old mdx mice showed IgG immunoreactivity mainly trapped in the large fibrotic area

(Figure 3a) and barely detected within the myofibre compartment (Figure 3c), suggesting little/no

ongoing myonecrosis. The observed low percentage of central nucleation together with low cell

infiltration and high fibrosis indicates that this phenotype is the outcome of accumulating necrotic and

regenerative events over time. Fibrosis decrease was observed in the diaphragm and was associated with

a decrease in the mean myofibres size and size variability (Figure 3g-k). These results are consistent

with a previous report assessing significant myofibre cytoprotection in the diaphragm[26]. We also found

a significant reduction of fibrosis in the TA of old mdx*Ripk3*[-/-] (Figure 2m) compared to mdx littermates,

suggesting that the effects of necroptosis inhibition on muscle histology benefit both locomotor and

respiratory muscles.

Two recent studies reported that necroptosis is involved in post-lesional myogenesis[6,27]. However, the

underlying mechanisms of action of necroptosis on regeneration seem unclear: Sreenivasan and

colleagues observed exclusive expression of RIPK3 in mdx MuSCs and not in myofibres. MuSCs

depleted for RIPK3 led to a dramatic loss of PAX7-expressing cells and severely impaired muscle

function[6]. On another hand, Zhou and colleagues found that myofibres express RIPK3 and MLKL and

are susceptible to necroptotic demise. They demonstrated that myofibre necroptosis indirectly modulates

muscle regeneration by releasing Tenascine-C, which is required for MuSCs proliferation. An effect of

myofibre necroptosis on *Pax7* expression and myofibre size was observed *in vivo* at 7 and 15 days post-

injury[27]. As the full effects on regeneration are generally observed at a later stage, after 4 weeks post-

injury, it is unclear in this study whether the depletion of necroptosis irreversibly stops myogenesis or

only delays it.

Herein, we found no impairment of running performance in dystrophic *Ripk3*[-/-] mice, and no/little effect

on muscle weight, while confirming the downregulation of *Ripk1* in muscle extracts. The weight of limb

muscles was unchanged, as myofibre size (Figure 2a-h). *Pax7* and *myogenin* expression were not

reduced in mdxRipk3-/-mice, suggesting there is no depletion of MuSCs. Together, our results

demonstrate that in mdx mice, a long-term necroptosis inhibition does not lead to sarcopenia of

dystrophic muscle, but rather has beneficial effects since it dampens fibrosis. The lack of observed

sarcopenia in our mdx*Ripk3*[-/-] dystrophic mice could for instance be explained by a delay of myogenesis

when necroptosis is prevented. Indeed, slowing down the kinetics of differentiation does not necessarily

impair regeneration[28]. It is also possible that any effect, positive or negative of RIPK3-deficiency on

regeneration could be mitigated by the cytoprotective effects of necroptosis inhibition. Not to be

forgotten that the regenerative process remains a consequence of previous degenerative events. Hence,

the less degeneration, the less resulting regeneration. Therefore, any protection against myonecrosis is

expected to hide any putative default on regeneration.

In mdx mice, myofibre necrosis does not lead to a dramatic lack of function of the limb, respiratory or

cardiac muscles[29]. On another hand, pathology is dramatically exacerbated in R-DMDdel52 rats with a

reduced life expectancy and cardiomyopathy functionally assessed in young adults[19]. Phospho-Ser161

RIPK1 is strongly upregulated in nuclei from cardiomyocytes, and the RIPK1/3 axis is largely

overexpressed, suggesting ongoing necroptosis activation. In mdx mice, the heart pathogenic phenotype

becomes detectable using M-mode echocardiography only after 10-12 months of age compared to wild-

type mice. Heart weight, histological features, and analysis can discriminate healthy from dystrophic

aged hearts[25,30,31]. Necroptosis depletion reduced heart weight, and biomarkers of hypertrophic

cardiomyopathy such as *Bnp* and *Myh7* transcripts[32,33] and fibrosis (Figures 4). BNP is upregulated in

mdx compared to C57BL/10 control mice and restored to baseline in mdx mice treated by rAAV9-

mediated microdystrophin[34]. The RIPK1/3 axis may therefore represent an interesting therapeutic target

to counteract the development of cardiomyopathy in DMD. The role of canonical necroptosis has been

found in cardiovascular diseases, such as atherosclerosis, ischemia-reperfusion injuries, myocardial

infarction, myocarditis and chemically-induced cardiomyopathy[35]. We failed to observe a significant

decrease in cardiomyocyte demise and necroinflammation. This is likely due to rare cell death events

affecting the cardiomyocytes of mdx mice at this age. Indeed, only a minority of mdx hearts showed

ongoing active necroinflammation area (Figure 4f). However, a reduction of cardiac hypertrophy linked

with a reduction of fibrosis deposition suggests, over time, less accumulation of myocardial cell death

susceptible to be replaced by collagen and overall improved tissue homeostasis.

396
In conclusion, using dogs, rats and mice models of DMD pathogenesis, we found that RIPK3 participates

in the pathogenesis of mdx mice at different levels, mediating necrosis and fibrosis accumulation in

locomotor, respiratory, and cardiac muscles without evidence of a detrimental effect on skeletal muscle

regeneration. This study, therefore, suggests that necroptosis prevention can be beneficial both to

skeletal muscle and heart phenotype in DMD conditions and paves the way for considering necroptosis

as a promising therapeutic target for DMD.

Acknowledgements

405   We thank Dr V. M. Dixit for *Ripk3* knockout mice. We also thank J. Ternacle, B. Drayton Libotte, L.

406   Guillaud and X. Decrouy for technical assistance and C. Laisne and D. Gelperowic for taking good care

407   of animals, and the VOUSH group for fruitful discussions. This work was supported by the Association

408   Française contre les Myopathies (AFM-Téléthon) through the Translamuscle I (#19507) and

409   Translamuscle II (#22946) programs.

Conflict of Interest

412   The authors declare no conflict of interest

Bibliography

1.   Relaix, F. *et al.* Perspectives on skeletal muscle stem cells. *Nat Commun* **12**, 692 (2021).

2.   Desguerre, I. *et al.* Endomysial fibrosis in Duchenne muscular dystrophy: a marker of poor
outcome associated with macrophage alternative activation. *J Neuropathol Exp Neurol* **68**, 762–773
(2009).

3.   Khirani, S. *et al.* Respiratory muscle decline in Duchenne muscular dystrophy. *Pediatr*
Pulmonol **49**, 473–481 (2014).

4.   Fayssoil, A., Abasse, S. & Silverston, K. Cardiac Involvement Classification and Therapeutic
Management in Patients with Duchenne Muscular Dystrophy. *J Neuromuscul Dis* **4**, 17–23 (2017).

5.   Morgan, J. E. *et al.* Necroptosis mediates myofibre death in dystrophin-deficient mice. *Nat*
Commun **9**, 3655 (2018).

6.   Sreenivasan, K. *et al.* Attenuated Epigenetic Suppression of Muscle Stem Cell Necroptosis Is
Required for Efficient Regeneration of Dystrophic Muscles. *Cell Rep* **31**, 107652 (2020).

7.   Zhou, S. *et al.* Myofiber necroptosis promotes muscle stem cell proliferation via releasing
Tenascin-C during regeneration. *Cell Res* **30**, 1063–1077 (2020).

8.   Barthélémy, I. *et al.* Effects of an Immunosuppressive Treatment in the GRMD Dog Model of
Duchenne Muscular Dystrophy. *PLoS ONE* **7**, e48478 (2012).

9.   Reyes-Fernandez, P. C., Periou, B., Decrouy, X., Relaix, F. & Authier, F. J. Automated
image-analysis method for the quantification of fiber morphometry and fiber type population in human
skeletal muscle. *Skelet Muscle* **9**, 15 (2019).

10.   Czibik, G. *et al.* Dysregulated Phenylalanine Catabolism Plays a Key Role in the Trajectory of
Cardiac Aging. *Circulation* **144**, 559–574 (2021).

11.   Wells, D. J. Tracking progress: an update on animal models for Duchenne muscular
dystrophy. *Disease Models & Mechanisms* **11**, dmm035774 (2018).

12.   Morgan, J. E. *et al.* Necroptosis mediates myofibre death in dystrophin-deficient mice. *Nat*
Commun **9**, 3655 (2018).

13.   McQuade, T., Cho, Y. & Chan, F. K.-M. Positive and Negative Phosphorylation Regulates
RIP1 and RIP3-Induced Programmed Necrosis. *Biochem J* **456**, 409–415 (2013).
14.   Zhang, Y. *et al.* RIP1 autophosphorylation is promoted by mitochondrial ROS and is essential
for RIP3 recruitment into necrosome. *Nat Commun* **8**, 14329 (2017).
15.   Newton, K., Sun, X. & Dixit, V. M. Kinase RIP3 is dispensable for normal NF-kappa Bs,
signaling by the B-cell and T-cell receptors, tumor necrosis factor receptor 1, and Toll-like receptors 2
and 4. *Mol Cell Biol* **24**, 1464–1469 (2004).
16.   Straub, V., Rafael, J. A., Chamberlain, J. S. & Campbell, K. P. Animal Models for Muscular
Dystrophy Show Different Patterns of Sarcolemmal Disruption. *J Cell Biol* **139**, 375–385 (1997).
17.   Bencze, M., Periou, B., Baba-Amer, Y. & Authier, F. J. Immunolabelling Myofiber
Degeneration in Muscle Biopsies. *JoVE (Journal of Visualized Experiments)* e59754 (2019)
doi:10.3791/59754.
18.   Briguet, A., Courdier-Fruh, I., Foster, M., Meier, T. & Magyar, J. P. Histological parameters
for the quantitative assessment of muscular dystrophy in the mdx-mouse. *Neuromuscul Disord* **14**,
675–682 (2004).
19.   Taglietti, V. *et al.* Duchenne muscular dystrophy trajectory in R-DMDdel52 preclinical rat
model identifies COMP as biomarker of fibrosis. *Acta Neuropathologica Communications* **10**, 60
(2022).
20.   Quinlan, J. G. *et al.* Evolution of the mdx mouse cardiomyopathy: physiological and
morphological findings. *Neuromuscul Disord* **14**, 491–496 (2004).
21.   Choi, M. E., Price, D. R., Ryter, S. W. & Choi, A. M. K. Necroptosis: a crucial pathogenic
mediator of human disease. *JCI Insight* **4**, e128834 (2021).
22.   Mariot, V. *et al.* RIPK3-mediated cell death is involved in DUX4-mediated toxicity in
facioscapulohumeral dystrophy. *J Cachexia Sarcopenia Muscle* (2021) doi:10.1002/jcsm.12813.
23.   Dondelinger, Y., Hulpiau, P., Saeys, Y., Bertrand, M. J. M. & Vandenabeele, P. An
evolutionary perspective on the necroptotic pathway. *Trends in Cell Biology* **26**, 721–732 (2016).
24.   Stedman, H. H. *et al.* The mdx mouse diaphragm reproduces the degenerative changes of
Duchenne muscular dystrophy. *Nature* **352**, 536–539 (1991).
25.   Quinlan, J. G. *et al.* Evolution of the mdx mouse cardiomyopathy: physiological and
morphological findings. *Neuromuscul Disord* **14**, 491–496 (2004).
26.   Stupka, N., Schertzer, J. D., Bassel-Duby, R., Olson, E. N. & Lynch, G. S. Stimulation of
calcineurin Aα activity attenuates muscle pathophysiology in mdx dystrophic mice. *American Journal
of Physiology-Regulatory, Integrative and Comparative Physiology* **294**, R983–R992 (2008).
27.   Zhou, S. *et al.* Myofiber necroptosis promotes muscle stem cell proliferation via releasing
Tenascin-C during regeneration. *Cell Res* **30**, 1063–1077 (2020).
28.   Bencze, M. *et al.* Proinflammatory Macrophages Enhance the Regenerative Capacity of
Human Myoblasts by Modifying Their Kinetics of Proliferation and Differentiation. *Molecular
Therapy* **20**, 2168–2179 (2012).
29.   Partridge, T. A. The mdx mouse model as a surrogate for Duchenne muscular dystrophy. *The
FEBS Journal* **280**, 4177–4186 (2013).
30.   Zhang, W. *et al.* Abnormal cardiac morphology, function and energy metabolism in the
dystrophic mdx mouse: An MRI and MRS study. *Journal of Molecular and Cellular Cardiology* **45**,
754–760 (2008).
31.   Li, W., Liu, W., Zhong, J. & Yu, X. Early manifestation of alteration in cardiac function in
dystrophin deficient mdx mouse using 3D CMR tagging. *J Cardiovasc Magn Reson* **11**, 40 (2009).
32.   Hasegawa, K. *et al.* Ventricular expression of brain natriuretic peptide in hypertrophic
cardiomyopathy. *Circulation* **88**, 372–380 (1993).
33.   Fananapazir, L., Dalakas, M. C., Cyran, F., Cohn, G. & Epstein, N. D. Missense mutations in
the beta-myosin heavy-chain gene cause central core disease in hypertrophic cardiomyopathy. *PNAS*
90, 3993–3997 (1993).
34.   Shin, J.-H. *et al.* Improvement of cardiac fibrosis in dystrophic mice by rAAV9-mediated
microdystrophin transduction. *Gene Ther* **18**, 910–919 (2011).
35.   Zhe-Wei, S., Li-Sha, G. & Yue-Chun, L. The Role of Necroptosis in Cardiovascular Disease.
Front Pharmacol **9**, 721 (2018).

Figures Bencze et al. 2022

Figure 1: The RIPK1-RIPK3 axis is dysregulated in the skeletal muscles and the heart of

dystrophic dogs.

(a) Quantification of serum CK in six-month-old WT and GRMD dogs. (b) Pearson correlation

coefficient (r) and p-value (p) between necrotic fibre percentage and *RIPK3* transcripts expression

normalized to *ACTIN*. Representative H&E staining of the sartorius muscle corresponding to related

biopsied muscle. Scale bar: 20 µm. (d) Pearson correlation coefficient and p-value between the

percentage of inflammation and *RIPK3* transcripts expression. (d) Quantification of *RIPK3* expression

in the diaphragms of 6-month-old WT and GRMD dogs. (e) Pearson correlation between *RIPK1* and

RIPK3 expression in the diaphragm. (f) H&E staining of the diaphragm muscle. (g) In the heart left

ventricle of GRMD and WT dogs, quantification of *RIPK1* and (h) *RIPK3* transcripts normalized to

ACTIN. (i) Representative image of a left ventricle heart from WT and GRMD dogs labelled with an

antibody directed against the phosphorylated form of RIPK1 (S161). Data are given as means ± s.e.m.

Ns: Not significant, *: p-value < 0.05.

Figure 2. The genetic ablation of RIPK3 does not promote sarcopenia in 18-month-old mdx mice.

Quantification of *Ripk1* transcripts in the eighteen-month-old biceps muscles (a) Seventeen-month-old mdx*Ripk3*+/+ (i.e. mdx) and mdx*Ripk3*-/- male mice were submitted to forced treadmill running with a progressive increase of speed. (b) Endurance curve of running mice against covered distance expressed in cm. (c) Maximum aerobic speed before mice exhaustion. Data are given as means ± s.e.m. mdx*Ripk3*+/+, n=7, mdx*Ripk3*-/-, n=8. Shapiro-Wilk test followed by a Mann-Whitney test. (d) Tibialis anterior (TA), EDL and biceps muscles were harvested, and muscle weight (MW) was determined (and normalized to tibial length (MW/TL). (e) TA immunolabeling using antibodies directed against Ki67 (red). (f) Quantification of the density of Ki67-positive cells. (g) Violin plot of the pooled minimum Feret of TA myofibres (mdx: n=33,142 myofibres out of 7 distinct muscles; mdx*Ripk3*-/-, n=26,469 myofibres out of 9 distinct muscles, Mann-Whitney test p=0.1526). (h) Relative frequencies of minimum Feret from TA myofibres are expressed in percentage. (i) Quantification of collagen deposition using Sirius red dye. The normal distribution of data was tested using the Kolmogorov-

Smirnov test (passing normality test with α=0.05), leading to either t-tests or Mann-Whitney tests. (j)

Quantification of the transcript levels of *Pax7*, (k), *myogenin*, (l) *follistatin* and (m) *myostatin* in the

biceps, by quantitative PCR. Data are given as means ± s.e.m. Ns: Not significant, *: p-value < 0.05,

**: p-value < 0.01. Scale bar:100 μm

Figure 3: Necroptosis participates in the diaphragm fibrosis in mdx mice.

(a) Representative image of IgG uptake in diaphragms of 18-month-old mdx (mdx*Ripk3*[+/+]) and

mdx*Ripk3*[−/−] mice. (b) Quantification of IgG uptake labelling expressed in percentage of cross-

sectional area (%CSA). (c) Quantification of myonecrosis extent. Myonecrosis is defined as the area

occupied by leaky myofibres (i.e immunoreactive for mouse IgG uptake). Quantification of nuclear

density expressed as a percentage of CSA (d), CD68-positive macrophage infiltrates expressed in

percentage of CSA (e), and centrally nucleated myofibres expressed as a percentage of myofibres (f).

(g) Violin plot of the minimum Ferets of diaphragm myofibres. (mdx, n=16873 including 7 distinct

muscles,   mean:18,61 ± 0.06555;   mdx*Ripk3*-/-,   n=31666   including   8   distinct   muscles,

mean:18,21 ± 0.04554, two-tailed Mann-Whitney test p < 0,0001). (h) Relative frequency of minimum

Ferets from diaphragm myofibres. Data are expressed as a percentage. (i) Quantification of the variance

coefficient of the minimum Feret of myofibres, (j) representative picture of Sirius red staining (k)

Quantification of diaphragm fibrosis. The normal distribution of data was tested using the Kolmogorov-

Smirnov test (passing normality test with α=0.05), leading to either t-tests or Mann-Whitney tests. Data

are given as means ± s.e.m. Ns: Not significant, *: p-value < 0.05, ****: p-value < 0.0001.

Figure 4: RIPK3 mediates the compensatory hypertrophy of the dystrophin-deficient heart

(a) Representative pictures of heart labelling using an antibody directed against the phosphorylated form

of RIPK1 at S161 from wild-type (WT) and dystrophin-deficient R-DMDdel52 rats. Quantification of

the rat *Ripk1* (d) and *Ripk3* (c) transcripts in hearts from WT and R-DMDdel52 rats. (d) Heart weights

of 18-month-old mdx mice. (e) Heart weights normalized to the tibial length (TL). (f) Quantification of

CD68-positive macrophage infiltrates. (g) Quantification of the density of CD31-positive microvessels,

(h) representative picture of Sirius red staining of mdx and mdx*Ripk3*⁻/⁻. Scale bar: 1700 μm. (i)

Quantification of RV fibrosis. Hearts from 18-month-old mdx*Ripk3*[+/+] and [-/-] were analyzed for *Bnp* (j),

and *Myh7* (k) mRNA levels by quantitative PCR. The normal distribution of data was tested using the

Kolmogorov-Smirnov test (passing normality test with α=0.05), leading to either t-tests or Mann-

Whitney tests. Data are given as means ± s.e.m. Ns: Not significant, *: p-value < 0.05, **: p-

value < 0.01.

561
Figure 5: The genetic ablation of RIPK3 reduces cardiomyopathy in mdx mice

(a) Representative M-mode echocardiogram traces in mdx and mdx*Ripk3*[-/-] 40-week-old mice showing

a reduced left ventricle (LV) dilation in mdx*Ripk3*[-/-] mice. (b) Quantification of LVIDd, (c) LV mass,

(d) LV relative wall thickness, (e), LV fractional shortening, and (f) strain rate (SR) of the anterior LV

wall. The normal distribution of data was tested using the Kolmogorov-Smirnov test (passing normality

test with α=0.05), leading to t-tests. Data are given as means ± s.e.m. *: p-value < 0.05, ***: p-

value < 0.001.

569
570

571 **Supplementary data**

	Mouse#	Tibial length (mm)	Normalizing factor	HR, beats/min	IVSd, mm	IVSs, mm	LVIDd, mm	LVIDs, mm	PWd, mm	PWs, mm	SRant, s-1	LV RWT, mm	LV mass, mg	LVFS, %	LVEF, %
						Measured values							Calculated values		
mdx	1	21.1	0.912	523	0.78	1.45	3.81	1.85	0.74	1.68	26	0.40	95.75	51	88
	2	18.2	1.056	605	0.94	1.64	3.92	1.70	0.76	1.49	29	0.43	117.26	57	91
	3	19.2	1.006	513	0.85	1.60	3.60	1.56	0.98	1.48	27	0.51	113.44	57	91
	4	19.1	1.011	535	0.86	1.88	3.52	1.63	1.13	1.88	25	0.57	123.85	54	89
	5	18.8	1.027	605	0.85	1.13	3.72	2.38	0.85	1.31	29	0.46	108.05	36	72
	Mean	19.3	1.0	556.2	0.9	1.5	3.7	1.8	0.9	1.6	27.2	0.5	111.7	50.9	86.2
	SD	1.10	0.05	45.22	0.06	0.28	0.16	0.33	0.16	0.22	1.79	0.07	10.60	8.65	8.04
mdxRipk3 -:-	1	18.1	0.932	545	0.74	1.43	3.28	1.48	0.75	1.26	24	0.45	101.18	55	91
	2	17.6	0.961	503	0.89	1.75	3.16	1.11	0.78	1.53	26	0.53	112.28	65	95
	3	17.8	0.950	609	0.79	1.56	2.76	1.03	0.85	1.30	24	0.59	87.89	63	94
	4	20.7	0.816	615	0.59	1.17	2.41	1.34	0.69	1.08	27	0.53	50.19	45	82
	5	19.7	0.855	650	0.74	1.17	2.21	1.19	0.76	1.06	27	0.68	55.91	46	84
	6	19.2	0.879	640	0.76	1.22	2.60	1.46	0.78	1.17	23	0.59	74.12	44	82
	7	18.4	0.915	634	0.76	1.35	2.56	1.46	0.76	1.38	25	0.59	70.62	43	80
	Mean	18.8	0.9	599.4	0.8	1.4	2.7	1.3	0.8	1.3	25.1	0.6	78.9	51.5	86.9
	SD	1.15	0.05	54.76	0.09	0.22	0.39	0.18	0.05	0.17	1.57	0.07	22.86	9.25	6.28

575 **Supplementary table 1: M-Mode echocardiographic values.**