Concurrent fMRI demonstrates propagation of TMS effects across task-related networks

Lifu Deng[1,8], Olga Lucia Gamboa[1,2], Moritz Dannhauer[3], Anshu Jonnalagadda[1], Rena Hamdan[3], Courtney Crowell[1,2,4], Tory Worth[1,8], Angel V. Peterchev[2,3,5,6], Marc A. Sommer[1,3,7,8], Roberto Cabeza[1,8], Lawrence G. Appelbaum[9], Simon W. Davis[1,4,8]

[1] Department of Psychology & Neuroscience, Duke University, Durham, NC 27708, United States of America.

[2] Department of Psychiatry and Behavioral Science, Duke University School of Medicine, Durham, NC 27708, United States of America.

[3] Department of Biomedical Engineering, Duke University, Durham, NC 27708, United States of America.

[4] Department of Neurology, Duke University School of Medicine, Durham, NC 27708, United States of America.

[5] Department of Electrical & Computer Engineering, Duke University, NC 27708, United States of America.

[6] Department of Neurosurgery, Duke University School of Medicine, NC 27708, United States of America.

[7] Department of Neurobiology, Duke University School of Medicine, NC 27708, United States of America.

[8] Center for Cognitive Neuroscience, Duke University, NC 27708, United States of America.

[9] Department of Psychiatry, University of California, San Diego, CA 94143, United States of America

Correspondence should be addressed to Simon W. Davis (simon.davis@duke.edu)

ACKNOWLEDGMENTS

We thank Lamont Conyers and Jennifer Graves for extensive MRI support, and all subjects for their participation. This study was supported by the National Institute of Health, Brain Initiative 1RF1-MH114253, R21-AG058161, and K01-AG053539.

CONFLICT OF INTEREST

A. V. Peterchev has received research funding, travel support, patent royalties, consulting fees, equipment loans, hardware donations, and/or patent application support related to magnetic stimulation from Rogue Research, Tal Medical/Neurex, Magstim, MagVenture, Neuronetics, BTL Industries, and Advise Connect Inspire.

1 ABSTRACT

Transcranial magnetic stimulation (TMS) has become an important technique in both
scientific and clinical practices, and yet our understanding of how the brain responds to TMS is
still limited. Concurrent neuroimaging during TMS may bridge this gap, and emerging evidence
suggests widespread that modulatory effects of TMS may be best captured through changes in
functional connectivity between distributed networks, rather than local changes in cortical
activity. However, the relationship between TMS stimulation parameters and evoked changes in
functional connectivity is unknown. In this study, 24 healthy volunteers received concurrent
TMS-fMRI while performing a dot-motion direction discrimination task. An MR-compatible coil
was used to apply trains of three pulses at 10 Hz rTMS over the primary visual cortex (V1) at
the onset of the dot stimuli with four levels of stimulation intensity (20%, 40%, 80%, and 120%
of resting motor threshold, RMT). Behavioral results demonstrated impairment of motion
discrimination at 80% RMT. FMRI results yielded three findings. First, functional connectivity
between visual and non-visual areas increased as a function of rTMS intensity. Second,
connectivity *within* the visual network was positively associated with motion accuracy, while the
connectivity *between* visual and non-visual regions was negatively associated with motion
accuracy. Lastly, we found that reductions in the similarity between functional and structural
connectivity associated with increasing TMS intensity were constrained to the visual network.
These findings demonstrate spatially dependent nonlinear effects of TMS intensity on brain
functional connectivity that proceed beyond the site of stimulation and influence associated
behavior.

22 1. INTRODUCTION

Transcranial magnetic stimulation (TMS) has been a valuable tool in both cognitive and clinical
neuroscience because it allows for the examination of specific neurocognitive mechanisms and
the development of effective therapies based on these mechanisms. Ideally, the understanding
of mechanisms should precede clinical application, but even the simple dose-response
relationships of TMS are largely unknown outside of the motor cortex. Multiple lines of evidence
in animal models and human studies suggest that TMS induces complex changes in activity
across broad networks of cortical regions, demonstrating clear evidence of polysynaptic effects
(Wang and Voss, 2015; Cocchi et al., 2016; Alekseichuk et al., 2019; Beynel et al., 2020b). In

line with these empirical findings, a recent systematic review of TMS on resting state networks (Beynel et al., 2020a) found that while network-level effects were common in the literature, these effects did not follow the typical frequency-dependent heuristic that helps to predict the local effect of TMS modulation (Luber and Deng, 2016). As such, it can be inferred that network-level modulatory effects depend on not only the parameters of stimulation but also the functional interactions between the stimulated and connected regions.

TMS during concurrent neural recordings has become an exciting technological development that may expand the repertoire of addressable questions by leveraging the rich spatial and temporal information afforded by different imaging methods. Previous studies utilizing concurrent TMS and fMRI are often focused on resting-state brain activity without the presence of cognitive tasks. In such studies, stimulation of primary sensory or motor cortices frequently is reported to exert feed-forward influences on functionally relevant regions, such as the frontal eye fields (Cocchi et al., 2016; Castrillon et al., 2020). Similarly, concurrent TMS during electroencephalographic (EEG) recordings has shown that single-pulse TMS to the dorsal attention network results in widespread activation of that same network (Ozdemir et al., 2020). There is also evidence that these intra-network effects are facilitated by structural connectivity (Momi et al., 2021a), suggesting that stimulation propagates between brain regions that are structurally connected to the stimulation site.

A lingering question is the relevance between these TMS-driven dynamics and cognition, as the effects of TMS has been shown to depend on the cognitive state (Silvanto et al., 2018). For example, when rTMS is applied to motion selective cortex (MT+) during a motion discrimination task, performance is hindered when attention is focused on motion but enhanced when attention is focused on other visual attributes (Brascamp et al., 2010; Zinchenko et al., 2021). Relatedly, connectivity can depend on brain state during motion perception, based on evidence that frontal and occipital connectivity is only present when participants are aware of apparent motion (Sanders et al., 2014). In addition, the timing of stimulation relative to the onset of visual motion strongly determines the influence on performance and EEG responses (Gamboa Arana et al., 2020), implying the different cognitive components affected. Because attention-driven activity during motion-perception tasks is influenced by gating between primary visual cortex and MT+ (Stephan et al., 2008), a network approach may better explain the effects of TMS than focusing on a single region.

Building on past studies on concurrent TMS-neuroimaging, we sought characterize the effects of TMS on network-level functional connectivity (FC) due to different intensities during visual motion perception. Graph theoretical analyses were used to depict the organization of

functional sub-systems and how TMS influenced communication between these systems. Three specific goals were pursued. First, this study investigated whether TMS affected global or local FC, and whether the effects were constrained to the network related to the task. Secondly, this study examined the effects of FC on motion discrimination task performance to better understand how changes in FC mediate behavior. Lastly, structural connectivity (SC) was used to explore how TMS alters the functional network relative to the underlying architecture. This analysis combines structural, functional, and behavioral information to further the understanding of the impacts of TMS on global brain network and behavior.

2. METHODS

2.1 Subjects

A total of 27 healthy volunteers (16 females, mean age = 23.7 years, SD= 2.78) participated in this study. At the beginning of the first visit, all participants were informed about the experimental procedures and were screened for contraindications to MRI and TMS. Subsequently, written informed consent was obtained from each participant. All participants were right-handed according to the Edinburgh Handedness Inventory (Oldfield 1971), had normal or corrected-to-normal vision, and did not have a history of neurological or psychiatric diseases. Participants received $20 per hour for taking part in this study. Procedures in this study conformed to the Declaration of Helsinki and were approved by the Institutional Review Board of Duke University. Data from three participants were excluded from the study as they declared falling asleep during the MRI session. As such, the final dataset analyzed for this study consisted of 24 healthy volunteers (13 females, mean age = 23.6 years, SD= 2.55).

2.2 Experimental design

Participants in this study took part in two experimental sessions. During the first session, participants were provided with screening questionnaires and were administered drug and pregnancy tests to check for eligibility. Resting motor threshold (RMT) was then acquired and subjects were briefly familiarized with the motion discrimination task by performing 1-2 minutes of practice trials. During the second session, participants first practiced the motion

discrimination task in a staircase fashion inside a mock MRI scanner that mimics the
environment of real MRI scanner (projection system, response box, recorded scanner sound,
etc.). Before starting the fMRI-TMS data collection, participants were positioned in the bore of
the MRI scanner and performed one additional staircase of the motion discrimination task to
determine the threshold coherence level used in the subsequent fMRI-TMS task, as described
in *Section 2.3.1*. After fMRI-TMS data acquisition, additional structural and diffusion weighted
images (DWI) were acquired.
100

101
Figure 1. *Experimental design. **(A)** Illustration showing still frames of the visual stimulus along with*
*three TMS pulses (orange markers) presented at 10 Hz at the start of stimulus presentation. **(B)** Coil*
*orientation and stimulation site over Oz as defined in the 10-20 EEG System. **(C)** 3D-printed headrest*
used in the MRI scanner to position the TMS coil over the stimulation site on Oz. Four different intensities
were used: Oz stimulation at 20, 40, 80 and 120% RMT as the main tests and vertex stimulation at 120%
*RMT as control. **(D)** Simulation of the TMS electric field (E-field) strength induced within each cortical*
*node, averaged over all participants. **(E)** Simulated E-field strength (mean ± standard deviation) at the*
target cortical node for each Intensity level across participants.

2.3 Procedures

2.3.1 Motion perception task

Stimuli were generated using MATLAB (Mathworks, Natick, MA, USA) and the Psychtoolbox extension (http://www.psychtoolbox.org). Each trial started with a white fixation-cross shown in the middle of the screen on a black background (see **Fig. 1A**). After 530 ms, white dots (5 pixels each, dot density: 1.41 dots/deg^2, dot- speed: 7.4 °/s) were presented for 150 ms, moving at a pre-determined coherence level (see next paragraph) either upwards or to the left within a circular window on the upper right quadrant of the screen. The center of the dot field was located at 8° from the fixation cross and extended 12° in diameter. Stimuli were projected onto a mirror placed above the scanner head coil using a rear projection system. The projector refresh rate was 60 Hz and viewing distance between the mirror and the projector screen was 70 cm. When needed, vision was corrected using MRI-compatible lenses matching a participant's prescription. Responses were recorded via a fiber-optic response box (Resonance Technology, Inc). Participants had up to 2 s to indicate if dot motion was to the left using the right index finger or upwards using the right middle finger. After each response, a cross colored either green (correct response) or red (incorrect response) was shown during 1 s to provide feedback. Participants were instructed to perform the task as accurately and as quickly as possible, with an emphasis on accuracy.

Before starting the TMS-fMRI data collection during the second session, subjects performed a final practice version of the motion perception task within the bore of the magnet comprising 120 trials in which dot coherence varied from 0% coherence to 100% coherence according to a staircase procedure. Data from this practice session was assessed online to ascertain an individualized coherence value to use in the main task. As in our previous work with this paradigm (Gamboa Arana et al., 2020; Gamboa et al., 2020), the calculation was done by means of a generalized linear model (MATLAB's glmfit function with 'binomial' option) that fit a sigmoid function to the signed (+1/−1 due to dot motion direction) trial coherences and individual correct/incorrect responses to find a particular coherence where 70% of the training trials were correct.

2.3.2 Concurrent TMS–MRI

After the practice task in the mock MRI scanner, subjects were transferred to the MRI scanner room. Participants wore a tightly fitting swimsuit cap where the stimulation points were marked. The main stimulation target was centered over Oz as defined in the 10-20 EEG system, which is

located 10% of the distance between the inion and nasion (approximately 3 cm above the inion

in most adults, see **Fig. 1B**) (Ishikawa et al., 2011). The coil was placed inside a custom

headrest that was 3D printed from semi-flexible material (nGen_FLEX) to be comfortable for the

subject and maintain the TMS coil position on the head during pulses (**Fig. 1C**). The headrest

was specially designed for each stimulation target so that the center of the coil was tangential to

the stimulation point and the induced electric field was perpendicular to the midline, while the

cable-side of the coil was on the right hemisphere. Positioning was adjusted both before the

scan using markers on the fitted swim cap and was verified after the experiment with the use of

four MR-sensitive fiducial markers placed over the borders of the major and minor semi-axes of

the coil. Cushions were used to reduce head motion inside the scanner and participants were

informed about the importance of minimizing head motion during task performance. Participants

wore earplugs to attenuate scanner noise and the TMS clicking sound.

A burst of three biphasic TMS pulses at 10 Hz delivered at motion onset was applied over the

stimulation site using a MagPro R30 magnetic stimulator connected to an MRI-compatible TMS

figure-of-eight coil (MRi-B91 Air-cooled, MagVenture, Farum, Denmark). Stimulation over the

visual area was delivered at 20%, 40%, 80% and 120% of the participant's RMT, presented in

pseudo-randomized order in each of the runs. The task was presented in the form of a slow

event-related design, such that the duration of each trial was 15 s. The TMS device, the

presentation computer, and the MR acquisition were synchronized via an Arduino board (**Fig.

2A**), such that the TMS pulses were delivered reliably in between slice acquisitions.

Furthermore, timing of the MR acquisition rate (30 slices every 2 seconds, or 15 Hz) and TMS

frequency (10 Hz) allowed us to place the TMS pulses consistently inferior (slice 4/30, roughly

at the level of the lower cerebellum) and superior (slice 28/30) locations, ensuring that artifacts

associated with the TMS pulse were spaced at least 4–6 slices away from the principle regions

of interest (i.e., visual cortex, **Fig. 2B**); Arduino code used to time pulses can be found online in

our public repository (https://github.com/ElectricDinoLab). The task was divided into five runs,

with each run containing 32 trials (8 trials for each intensity), giving a total 40 trials per Intensity

over the course of an fMRI session. As each trial comprised three 10 Hz pulses, a total of 120

pulses were administered at each intensity.

Figure 2*. Concurrent TMS-fMRI. **(A)** A generalized wiring diagram. **(B)** Schematic of the timing of*
TMS pulses to avoid concurrent acquisition of data and TMS pulses.

2.3.3 Resting motor threshold (RMT) measurement

To determine the TMS intensities to use during the experiment, the resting motor threshold

(RMT) of the right first dorsal interosseous muscle (FDI) representation in the left primary motor

cortex was obtained in the first visit. During this procedure, biphasic single pulses were sent

with a MagPro R30 stimulator connected to a figure-of-eight coil (MCF-B65 cool coil,

MagVenture, Farum, Denmark) while being guided by a stereotaxic neuronavigation system

(Brainsight, Rogue Research, Canada). Surface electromyogram (EMG) was recorded from the

right FDI muscle by using disposable electrodes (Covidien / Kendall, 133 Foam ECG

Electrodes, Mansfield, USA) in a belly tendon montage. RMT was defined as the minimum

intensity at which a motor evoked potential (MEP) of 50 µV average peak-to-peak amplitude

was elicited when the muscle was at rest (Conforto et al., 2004). The Motor Threshold

Assessment Tool software (MTAT 2.0) (http://www.clinicalresearcher.org) was used to

determine the RMT once a reliable stimulation site for the FDI motor 'hot spot' was found. The

percentage of the maximum stimulator output and the realized peak coil current rate of change

(dI/dt) values were registered to be used in the following session.

2.3.4 fMRI data acquisition & analysis

Functional MR images were obtained with a 3T GE MRI system scanner with a 4-channel

transmit/receive head coil at the Brain Imaging and Analysis Center (BIAC) at Duke University

Medical Center. During the visual task an echoplanar imaging (EPI) sequence was used (320

volumes, 30 axial slices, repetition time/echo time =1500/27 ms; flip angle = 70; FOV = 256

mm^2; matrix size = 64×64; voxel size = 4×4×4 mm^3; slice thickness = 4mm). The first 9 s (6

volumes) of the fMRI record were discarded to allow for T1 equilibrium magnetization.

Anatomical images for co-registration were collected using a T1-weighted MP-RAGE sequence

(repetition time/echo time/inversion time = 6.972/3.01/400 ms, flip angle =11º).

Functional image processing proceeded first by removing slices affected by TMS pulses. This

was necessary because, even though TMS pulses were timed to be delivered in between MR

slice acquisitions, small errors in timing may have resulted in overlap between the TMS pulse

and MR acquisition. We identified slices with a signal magnitude of > 2 SD from the run mean

and visually inspected them for the presence of the TMS artifact. These slices were replaced by

temporal interpolation of the signal values of the same slice from the preceding and succeeding

volumes. Functional images were then preprocessed using image processing tools, including

FLIRT (FMRIB's. Linear Image Registration Tool) and FEAT (FMRIB Expert Analysis Tool) from

FMRIB's Software Library (FSL, https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/). Images were corrected for

slice acquisition timing, motion, and linear trend; motion correction was performed using FSL's

MCFLIRT, and six motion parameters estimated from the step were then regressed out of each

functional voxel using standard linear regression. Images were then temporally smoothed with a

high-pass filter using a 190 s cutoff and normalized to the MNI stereotaxic space. To remove

any additional artifacts associated with TMS pulses, spatiotemporal independent components

analysis (ICA, FSL's MELODIC) was used to identify and remove noise components occurring

in slices collected concurrent with the TMS pulse. White matter and CSF signals were also

removed from the data and regressed from the functional data using the same method as the

motion parameters. BOLD activations during the initial dot motion presentation were entered in

a standard GLM, using HRF-convolved trial regressors and their temporal derivatives.

2.3.5 Electric field modeling

The direct effects of TMS are driven by the strength and spatial distribution of the electric field (E-field) induced in the brain. Thus, simulation of the E-field induced by TMS in individual participants is increasingly recognized as an important step in spatial targeting of specific brain regions and forms a critical link between the externally applied TMS parameters and the neurophysiological response supporting cognitive operations (Peterchev et al., 2012; Gomez et al., 2020). To determine the E-field induced by TMS in the brain of each participant, simulations using T1 images and the finite element method in the SimNIBS software package (version 2.0.1, Thielscher et al., 2015) were conducted (Beynel et al., 2019; Beynel et al., 2020c; Beynel et al., 2021). Twenty-three subjects were included in the E-field analysis and head models were created using the SimNIBS mri2mesh pipeline, featuring five distinct tissue types: skin, skull, cerebrospinal fluid, gray matter, and white matter. The scalp, skull, and cerebrospinal fluid tissue compartments were modeled with isotropic conductivity values of 0.465, 0.01, and 1.654 S/m, respectively. For seventeen subjects, available DWI information was used to estimate anisotropic conductivities for the brain matter using the volume-normalized approach (Gullmar et al., 2010) which kept the geometric mean of the conductivity tensor eigenvalues for the gray and white matter equal to literature-based isotropic values of 0.275 and 0.126 S/m, respectively. DWI data could not be processed for six subjects, for which the latter isotropic conductivity values for gray and white matter were assigned to their corresponding computational tetrahedral elements. For nine subjects, 4 fiducial markers implemented with gel-filled stickers were attached on the active side of the TMS coil to image the location of the coil relative to the head using additional T1 scans. Based on the location of the fiducial markers, the mean (+/- standard deviation) shortest distance between the TMS coil center and the scalp was determined to be 3.8 mm (+/- 3.4 mm) across the nine subjects, and the coil plane deviated from the local scalp tangential plane by only 7.2° (+/- 3.2°), confirming good coil placement. For fifteen subjects it was not possible to consistently image the fiducial markers. Instead, the target Oz scalp site was identified from the MRI data and head model's scalp surface, the coil center was extruded by 4 mm to account for the scalp–coil spacing, and the coil plane was oriented tangentially to the closest scalp surface location. The individual E-field was simulated for TMS coil current rate of change (dI/dt) of 1 A/μs. Leveraging the direct proportionality between the coil current and the E-field strength, the latter was obtained by scaling the E-field solution for the various TMS intensities based on the corresponding dI/dt value computed from the Intensity setting according to the formula dI/dt = 1.9308 × %MSO − 2.4018. The formula was derived by linear regression

250 of the dI/dt values reported by the TMS device for Intensity settings as percentage of maximum

251 stimulator output (%MSO) for 5% MSO steps averaged over three TMS pulses. **Figures 1D** and

252 **1E** present, respectively, a map of the cortical-node E-field strength at 120% RMT averaged

253 across the subjects, and the E-field strength at the target cortical nodes across the four TMS

254 intensity levels.

255 2.3.6 Network Analysis

Functional connectivity (FC) network construction. A consistent parcellation scheme across

all subjects and all modalities (DWI, fMRI) was used. Subjects' T1-weighted images were

segmented using SPM12 (fil.ion.ucl.ac.uk/spm/software/spm12/), yielding a GM and white

matter mask in the T1 native space for each subject. The entire GM was then parcellated into

397 ROIs, each representing a network node by using a sub-parcellated version of the Harvard-

Oxford Atlas (Fornito et al., 2010). Functional connection matrices representing task-related

connection strengths were estimated using a correlational psychophysical interaction (cPPI)

analysis. Briefly, the model relies on the calculation of a PPI regressor for each region, based

on the product of that region's physiological time series and a task regressor convolved with the

HDR, to generate a term reflecting the psychophysical interaction between the seed region's

activity and the specified experimental manipulation. The psychological regressor was multiplied

with two physiological time courses for regions i and j, producing two psychophysiological

interaction regressors, denoted as PPI_i and PPI_j. The partial correlation ($\rho_{XY.Z}$, where X= PPI_i,

Y=PPI_j), was then computed as the partial correlation between two psychophysiological

interaction regressors for a given pair of regions *i* and *j*, controlling for the covariates (denoted

as *z*, which includes the HDR-convolved task regressor and the BOLD time courses for regions *i*

and *j*); the six motion parameters, white matter signal, and CSF signal were also included as

confounds.

Modularity. Functional sub-network community assignment was identified using modularity

analysis (Reichardt and Bornholdt, 2006). A common whole-brain network was obtained by

averaging all functional networks across participants and intensity levels. This common network

was then thresholded to preserve 15% strongest connections and submitted to the modularity

algorithm (gamma = 1). Based on module assignments, functional connections were classified

into between-module connections (connecting regions in different modules) and within-module

connections (connecting regions in the same module).

Structural connectivity (SC) network construction. Information on the structural connections

based on diffusion tractography between each pair of regions in the data was assessed with a

standard DWI processing pipeline used previously in our group (Davis et al., 2019). DWI data
were analyzed using FSL (fsl.fmrib.ox.ac.uk) and MRtrix (mrtrix.org) software packages. Data
were denoised, corrected with eddy current correction, and bias-field corrected. Constrained
spherical deconvolution was used in calculating the fiber orientation distribution; after tracts
were generated, they were filtered using spherical-deconvolution informed filtering of
tractograms (SIFT; Smith et al., 2013). To generate the SC network, the value of SC was
defined as the number of streamlines connecting each pair of ROIs.

2.3.7 Statistical Analysis

Statistical analyses were performed using MATLAB and RStudio. The effect of TMS intensity
on behavior was assessed using repeated-measures ANOVA. The FC network of each
participant under each rTMS intensity was first normalized into z-scores to further control for
motion-induced confound. The between-and within-module FCs were then derived from the z-
scored network. The effect of TMS on FC within and between network was assessed using two-
way repeated-measures ANOVA, in which the Type of FC (between/within) and the Intensity of
TMS (20/40/80/120% RMT) were defined as the two main effects. The relationship between
connectivity and task performance was examined in mix effect regression model using the lme4
package in R, in which the dependent variable was the correct response rate. The random
effect was specified as participant-specific intercept. Variables of interests, estimated as fixed
effects, were TMS intensity, between-module and within-module FC of the visual module, as
well as participant-level covariates such as sex, coherence of the visual stimuli, and resting
motor threshold. The neural effects on performance were assessed using F-contrasts and T-
contrasts. Details about the design matrix and the contrasts can be found in **Figure S1a** and
Figure S1b, respectively, in Supplementary Materials.

3. RESULTS

3.1 Behavioral Results

The accuracy of motion discrimination task (leftwards vs. upwards motion) during concurrent
TMS-fMRI was analyzed to assess the behavioral effect of TMS targeting visual cortex. A one-
way repeated-measures ANOVA revealed significant differences in performance across TMS
intensity levels ($SS_{subjects} = 0.69$, $SS_{condition} = 0.04$, $F_{3,23} = 2.93$, $p = 0.039$). Post hoc, uncorrected
comparisons between TMS Intensity conditions revealed that motion accuracy in the 80% RMT
condition (mean: 58% ± 9%) was significantly lower than 20% RMT (mean: 63% ± 13%, $p =$
0.011), 40% RMT (mean: 62% ± 10%, $p = 0.024$), and 120% RMT (mean: 61% ± 9%, $p =$
0.047) conditions (**Fig. 3A**). No differences in discrimination accuracy were seen between the
highest (120% RMT) and lowest (20% RMT) stimulation intensities ($p > 0.05$), suggesting that
the disruptive effects were attributable to a neural response in visual processing regions rather
than distraction-related confounds (sound, somatic sensations) associated with increasing TMS
intensity. Reaction times (**Fig. 3B**) were not significantly affected by the different TMS
intensities, though correct trials were significantly faster than incorrect trials (paired $t_{23} = 5.33$, p
= 0.0002).

 *Figure 3. Behavioral results. Accuracy **(A)** and reaction times **(B)** for all participants in the dot-motion*
perception task during TMS-fMRI. TMS selectively disrupted motion performance at 80% RMT, relative to
all other stimulation intensities. In contrast, no effects of Intensity were seen on reaction time.

3.2 Delineating task network nodes affected by TMS

The present analysis focused on evaluating network-level, fMRI-based functional connectivity to characterize the effects of stimulation intensity. The network-level effects of TMS were examined on two scales: brain modules, and directly stimulated nodes in visual cortex. Stimulated nodes were defined using E-fields estimated within each individual, localized using fiducial markers placed on the TMS coil; E-fields were then scaled by individual RMT, and then voxelwise E-field values were averaged within a parcellated region of interest to identify the network node with the highest E-field (**Fig. 1D**). Whole brain network data were delineated into meaningful networks to establish reliable boundaries between local and global communities of regions, termed "modules" (also known as "subnetworks"), by using modularity analysis on the average FC matrix (averaging across subject, TMS intensity, and task success conditions). This data-driven modularity analysis parsed 397 ROIs into four highly recognizable modules (**Fig. S2**), including (i) default mode network (DMN), (ii) somatosensory-motor network (SM), (iii) a visual-temporal network complex, and (iv) frontal-parietal control network (FPCN). To obtain a more accurate delineation of the visual system in the Visual-Temporal complex, an additional modularity analysis was performed on it, which in turn produced four subdivisions (**Fig. S3**); among these subdivisions, the two posterior subdivisions were combined into the Visual network (Vis.), and the two anterior, the Temporal network (Tmp.). The final 5-module assignment is shown in **Figure 4A**.

3.3 TMS effects on functional connectivity

3.3.1 TMS effects on visual system functional connectivity

Network-level TMS effects were first evaluated on the task-relevant regions. The 5-module community assignment (**Fig. 4A**) was used to define within-module FC (the average FC within the same module) and between-module FC (the average connectivity between the given module and other modules). For each of the five modules, an Intensity (% RMT) by Type (within/between module connectivity) repeated-measures ANOVA was performed. For all five modules, the within-module FC was significantly stronger than the between-module FC (DMN: $F_{1,21}$ = 11.8, p = 0.0023; SM: $F_{1,21}$ = 40.52, p=1.7e-6; Visual: $F_{1,21}$ = 33.04, p = 7.4e-6; Temporal: $F_{1,21}$ =17.45, p = 3.6e-4; FPCN: $F_{1,21}$ = 17.23, p = 3.9e-4), suggesting that the data-driven

modularity analysis successfully captured meaningful modules of the whole-brain functional

network. For completeness, results for the larger Visual-Temporal complex are shown in **Fig. S2**

in the Supplementary Materials, which were similar to the findings for the Visual network.

*Figure 4. Within and between-module connectivity in response to TMS. (**A**) The 5 modules obtained*
from modularity analysis (DMN = default mode network, SM = somatosensory-motor network, Vis. =
*visual network, FPCN = frontal-parietal control network, Tmp. = temporal network). (**B**) and (**C**) The*
effects of TMS intensity on visual network FC (z-scored); dashed horizontal brace indicates the Intensity x
*Type interaction; horizontal solid lines indicate the main effects of Intensity. (**D**) The effects of TMS*
intensity (expressed as F-values) on the FC between different modules. Error bars indicate standard error
of the mean (SEM). Asterisks (∗) indicate p<0.05.

Crucially, a significant Intensity x Type interaction was found uniquely in the Visual network

($F_{3,69}$ = 3.99, p = 0.011, **Fig. 4B, 4C**) but not in the other modules (all p > .05), suggesting that

the TMS may have a type-specific effect on the functional communication of the targeted visual

system. The effects of TMS intensity on these two types of FC were further examined in two

separate one-way repeated measure ANOVAs, where the intensity effect was significant for

between-module connectivity (i.e., 'Visual to Non-Visual', $F_{3,69}$ = 2.88, p = 0.039, **Fig. 4B**) but

insignificant for within-module connectivity (i.e., 'Visual to Visual', $F_{3,69}$ = 1.69, p = 0.178, **Fig.**

4C). An exploratory analysis suggested that such significant modulation of between-module FC

seemed to be driven by the connections with SM and Temporal modules; both findings are

consistent with the idea that the effects of increasing TMS elicited greater network activity due

to increasing motor (SM) or auditory (Temporal) sensation. No significant effect of TMS intensity

existed between other modules (**Fig. 4D**). In sum, these results showed that TMS modulated FC
of the targeted functional system.

The above results support the idea that intensity effects can be seen at the level of the
stimulated network, but a more rudimentary concern might be that these same effects can also
be seen at the level of the directly stimulated region. To examine this possibility, two non-
overlapping groups of visual ROIs were further investigated: 1) three ROIs in the left occipital
pole that were closest to the TMS coil (Group 1**, "directly-stimulated visual cortex"**), which
also received visual inputs from the right visual field and, and 2) the ROIs in visual network not
directly stimulated (Group 2**, "Indirectly-stimulated visual cortex"**). For each of these two ROI
groups, the average FC to other Visual module ROIs (i.e., 'Within Visual FC') and the average
FC to non-Visual ROIs (i.e., 'Group 1/2 to Non-visual FC') were measured. Results largely
followed similar, but weaker patterns as found in the above paragraph. For *directly-stimulated*
visual cortex (Group 1)**,** the Intensity by Type ANOVA did not produce a significant interaction
($F_{3,69}$ = 1.85, p = 0.15), and two separate one-way ANOVA showed no effect of Intensity in
either "Group 1 to Visual" FC ($F_{3,69}$ = 1.18, p = 0.33) or "Group 1 to Non-Visual" FC ($F_{3,69}$ = 0.26,
p = 0.85). In comparison, the results for *indirectly-stimulated* visual cortex (Group 2) were
largely consistent with the results based on the whole Visual network (**Figs. 4B**, **4C**): the
Intensity by Type ANOVA revealed a significant Intensity x Type interaction ($F_{3,69}$ = 3.12, p =
0.032), and two separate one-way ANOVA showed no effect of Intensity in "Group 2 to Visual"
FC ($F_{3,69}$ = 1.65, p = 0.17) but significant "Group 2 to Non-Visual" FC ($F_{3,69}$ = 3.38, p = 0.023). In
summary, these ancillary findings demonstrated the distributed nature of TMS effects across the
visual system, with changes in FC largely observed in brain regions that were not directly
stimulated by the TMS E-field.

3.3.2 Behavioral correlates of visual system FC

A second goal of this study was to examine the link between FC and performance. Because
TMS led to changes in both performance and visual network FC, links between behavior and FC
should be expected and may shed light on the neural mechanisms of the TMS effect on
performance. In addition, it is possible that within-module FC and between-module FC may
relate to performance in different ways, as the effect of TMS intensity has been shown in the
previous section to interact with FC types (i.e., within- or between-module connectivity).
Therefore, the links between visual system FC and task performance were compared in a mixed
effect regression model in which the correct response rate was predicted by the between-
module and within-module FC of the visual module. The model also accounted for other factors

that may affect performance, including TMS intensity, baseline performance, RMT, and gender

(see **Fig. S1**). The effects of these factors on performance were assessed using F-contrasts

and T-contrasts and are summarized in **Table 1**.

Table 1. *Effects of TMS Intensity (%RMT) and FC type on functional connectivity.*

	Z-stat	F-stat
TMS intensity	-	0.48 (n.s.)
Between-module FC	**-3.02** ∗∗	-
Within-module FC	**2.26** ∗	-
Between vs Within-module	**3.23** ∗∗	-
Between-module FC x Intensity	-	0.045 (n.s.)
Within-module FC x Intensity	-	1.82 (n.s.)

Note: ∗: p<0.05, ∗∗: p<0.01

In contrast to the behavioral findings in *Section 3.1*, once FC measures were considered in

the model, the TMS intensity showed no reliable effect on performance ($F_{3,64.12}$ = 0.48, p = 0.70).

In fact, the between-module FC, previously shown to be significantly modulated by TMS,

negatively predicted response accuracy (beta = -0.27, z = -3.02, p = 0.0025, **Fig. 5A**). The

within-module FC, in contrary, positively predicted response accuracy (beta = 0.12, z = 2.26, p =

0.024, **Fig. 5B**). Such differences in behavioral association were further indicated by a Type by

FC interaction (i.e., between-module minus within-module; beta = 0.39, z = 3.23, p = 0.0013),

suggesting that the effect of TMS on FC during task performance was dependent on the type of

FC. There was no significant Intensity x FC interaction for either within-module FC ($F_{3, 63.54}$ =

1.82, p = 0.15) or between-module FC ($F_{3, 63.84}$ = 0.045, p = 0.99) of the Visual module. In sum,

these results point to the network-level functional communication mediating the TMS-behavior

relationships.

Figure 5. *Relationships between task performance and FC either (**A**) between the Visual system*
*and all non-Visual modules, or (**B**) within the Visual network.*
439

3.3.3 Structural constraints in TMS-induced functional connectivity

The third goal of this study was to determine if FC patterns, constrained by the structural

connectivity (SC) network, can be altered by TMS. As the pattern of functional communication

has been shown to rely on a backbone of white-matter connections (Hermundstad et al., 2013;

Goni et al., 2014), it was predicted that the impact of TMS on FC can be reflected by the level of

consistency between the functional network and its underlying structural network. At the most

basic level, FC-SC similarity can be characterized with Spearman's correlation between

structural and functional networks; implicit in this relatively straightforward approach is the

premise that FC should reflect the underlying SC, and that lower FC-SC similarity indicates a

greater deviation of the FC pattern from a more original configuration constrained by the

structural backbone, hence reflecting a greater disruption of the FC pattern.

For the Visual network, an Intensity by Type repeated measures ANOVA on FC-SC similarity

was employed to assess if and how TMS contributes to the change in FC. The analysis

revealed a significant effect of Intensity ($F_{3,63}$ = 6.17, p = 9.6e-4), as well as a significant

Intensity by Type interaction ($F_{3,63}$ = 4.42, p = 6.9e-3, see **Fig. 6A, 6B**). Separate one-way

repeated measure ANOVAs were performed to further examine how between-module and

within-module connections were affected. As shown in **Figure 6B**, the TMS intensity effect

turned out to be significant for within-module FC-SC similarity ($F_{3,63}$ = 11.18, p = 5.65e-6) in the

Visual network, leading to more distinct patterns of FC at higher intensity levels. In comparison,

TMS intensity showed no effect for between-module FC-SC similarity ($F_{3,63}$ = 0.65, p = 0.59).

Lastly, an exploratory analysis was performed to test whether the FC pattern between other

modules could be affected by TMS (**Fig. 6C**), which further confirmed that the TMS-induced

disruption of FC pattern was specific to the FC within visual system.

Figure 6. *The effects of TMS intensity on visual network FC-SC similarity. Dashed horizontal brace indicates the Intensity x Type interaction; horizontal solid lines indicate main effect of Intensity in the Within Visual (**B**) connections, but not from Visual to Non-Visual (**A**) connections. **C**) The effects of TMS intensity (expressed as F-values) on the patterns of FC (measured as FC-SC similarity) between different modules. (**D**) and (**E**) The relationships between task performance and FC-SC similarity. Error bars indicate standard error of the mean (SEM). Asterisks (∗) indicate p<0.05.*

Next, to determine if FC-SC coupling influenced motion perception accuracy a mixed effect model was performed in which the correct response rate for each participant, under each TMS intensity level, was assigned as the outcome variable. The TMS intensity levels and its interactions with the between-module FC-SC similarity and the within-module FC-SC similarity of the visual module were modeled as fixed effect. Potential covariates, including sex, coherence of dots, and RMT of each subject, were included in the model and treated as fixed effect as well. The effects of these factors on performance were assessed using F-contrasts and T-contrasts, and they are summarized in **Table 2**. Similar to the previous finding in *Section 3.3.2*, with the brain measure (FC-SC similarity) taken into account, TMS intensity no longer showed a significant effect on the correct response rate ($F_{3,56.71} = 0.19$, p = 0.90). The within-module FC-SC similarity of the visual network, previously shown to be modulated by TMS, was positively predicting the correct response rate (beta = 0.31, z = 2.71, p = 0.0068, **Fig. 6E**). The between-module FC-SC similarity of the visual module was not significantly related to correct response rate (beta = -0.04, z = -0.32, p = 0.75, **Fig. 6D**). The Intensity by FC Type interaction

486 effect was found to be trending significant (i.e., between-module minus within-module; beta =
487 0.36, z = 1.77, p = 0.077).

489   ***Table 2.*** *Effects of TMS Intensity (%RMT) and FC-SC similarity on Behavior.*

	Z-stat	F-stat
TMS intensity	-	0.19 (n.s.)
Between-module FC-SC sim.	-0.32 (n.s.)	-
Within-module FC-SC sim.	**2.71** ∗∗	-
Between vs Within-module	1.77 ·	-
Between-module FC-SC sim. x Intensity	-	1.66 (n.s.)
Within-module FC-SC sim. x Intensity	-	0.24 (n.s.)

490   **Note:** ·: p<0.1; ∗: p<0.05, ∗∗: p<0.01
491

4. DISCUSSION

Current views of the neural effects of TMS are changing. Earlier basic science approaches to
TMS in humans focused on the effects of localized stimulation to a specific cortical target. As
both imaging and TMS techniques evolved, a more recent focus on bivariate connections from
the stimulation site to a subcortical target have revealed promising findings, such as the
possibility of targeting subcortical structures with cortical TMS (Wang et al., 2014; Luber et al.,
in press). However, there is a growing appreciation for how stimulation at a single site can
propagate throughout a cortical network (Cocchi et al., 2016) and a growing need to identify
which principles govern that flow. Though there are only a limited number of combined TMS-
EEG or TMS-fMRI studies considering this problem from a network-level perspective, one
general principle that emerges across studies is that local stimulation spreads according to the
functional networks associated with the stimulated node (Momi et al., 2021b). This study
advances the field by considering this phenomenon during the engagement of an active task,
and finds that the effects of TMS on ongoing brain activity are most evident in regions actively
engaged with the task at hand. This is a critical extension of expectations of local cortical
excitation or inhibition with specific stimulation frequencies (Aydin-Abidin et al., 2008), and to
the variability of modulation observed during different brain states (Gollo et al., 2017). By
stimulating with bursts of 10 Hz rTMS at different intensities over the primary visual cortex
during a motion perception task, and concurrent with fMRI imaging, the present study revealed
four novel findings. First, motion processing was selectively impaired at 80% RMT. Second, the
connectivity between the visual system and other cortical communities increased with

stimulation intensity. Third, intra-community connectivity was beneficial, while the extra-
community connectivity induced by TMS was detrimental to performance. Lastly, structural
connectivity constrained this effect, such that FC-SC was strongest in low-intensity conditions,
and was disrupted with increasing intensity, resulting in poorer performance. These findings are
discussed in greater detail below.

Before turning to the connectivity-related findings, it is important to consider the
observation that TMS-related impairment of motion discrimination was selective to the 80%
RMT condition. No differences in discrimination accuracy were seen between the highest (120%
RMT) and lowest (20% RMT) stimulation intensities, suggesting that the disruptive effects were
attributable to a neural response in visual cortex, and not distraction-related confounds
associated with increasing TMS intensity (e.g., noise, cutaneous sensation). Past studies have
reported a wide variety of behavioral effects that result from single or paired pulse TMS over a
range of factors, such as timing, intensity, and the ongoing activation state of the stimulated
region. For example, in primary motor cortex the threshold for inhibitory effects of TMS is
approximately 80% of RMT (Kallioniemi et al., 2014). Thus, TMS intensity can determine the
local balance between inhibitory and excitatory circuit recruitment. In another example, single
pulses of 80% RMT TMS timed to the onset of the N2 latency of MT+ facilitate motion
perception (Gamboa Arana et al., 2020). In addition to the possibility of a specific balance of
inhibitory and excitatory effects, such findings may be explained by a stochastic resonance
account, such that the strength of a stimulus can be improved by externally enhancing the
ongoing neural activity. Despite a similar task paradigm and dosing structure, the current study
instead revealed a disruption of performance at 80% RMT, perhaps due to the differences in
cortical target (V1 vs. MT+), or the use of three pulses (spaced at 10Hz) instead of one pulse of
TMS. Furthermore, this single pulse was timed to the onset of a specific, motion sensitive ERP.
Nonetheless, the current results may abide by the principle that a selective modulation of
performance is based on the engagement with ongoing brain activity, such that that 10 Hz
pulses timed to the onset of motion disrupted the downstream flow of information from primary
visual cortex to motion-sensitive cortex (MT+). This pattern of results also lends support to a
more global model of the effects of TMS on behavior (Silvanto and Cattaneo, 2021) that posits
distinct facilitatory and suppressive ranges as a function of TMS intensity, and which are shifted
by state-dependent changes in neural excitability (e.g., during stimulus processing vs. rest
states).

Turning to the general connectivity changes in response to TMS, the current study first
found that local connectivity within the visual system was more resilient to stimulation intensity

than global connectivity from the visual system to the rest of the brain. Such a result is
consistent with the notion that the network responsible for visual motion is an optimized system
that operates best when allowed to segregate from other cortical networks. In contrast, when
stimulation intensity induced an increased spread of connectivity between modules, this more
global pattern had negative consequences for behavior. In fact, when we examined the source
of these intensity-related increases in between-module FC, we found greater FC with the
somatosensory-motor and temporal networks, consistent with the idea that increasing intensity
of TMS elicited greater network activity due to increasing motor (SM) or auditory (Temporal)
sensation. Thus, an important consideration for studies examining the network-level effects of
TMS is to consider how some of these distant effects can be attributed to the sensory properties
of TMS stimulation, as revealed by conditions that mimic the sound and somatosensory
attributes of TMS while minimizing intracranial E-fields ("sham" conditions). In some cases,
these confounds may be unavoidable (see Conde et al., 2019 for innovative post-processing
techniques) but they are not necessarily a barrier to interpreting effects in non-sensorial regions.

An interesting, if overlooked, observation in many concurrent TMS-fMRI studies is that
the effects of localized TMS are not strongest at the site of stimulation and are often seen in
distant regions of the cortex. While some of these distant effects can be attributed to the
sensory properties of TMS stimulation, discussed above, a recent meta-analysis revealed that
this surprising absence of a local effect generalizes across many different stimulation paradigms
(Rafiei and Rahnev, 2021). More broadly, it is important to consider how such networks may be
influenced or modulated by the level of attention associated with increasing TMS intensity.
Selective attention can modulate the ongoing processing in both visual (Ballesteros et al., 2008)
and motor (Johansen-Berg and Matthews, 2002) regions, and thus it is possible that heightened
responses in sensory regions reflect an increase in the alerting or arousal effect associated with
increasing TMS intensity, in addition to the input from more basic sensory afferents. While not
an explicit focus of this study, such an interpretation is not inconsistent with the finding that
attention-driven MT+ activity during motion-perception tasks is influenced by the gating of V1
and V5 connectivity by the posterior parietal cortex (Silvanto et al., 2005). Taken together, this
perspective links how a network-level interpretation of TMS effects can better explain the
function of single region in a global perspective. While there are many studies on the impact of
TMS on local function of the visual regions during motion paradigms (Campana et al., 2002;
Cowey et al., 2006; Amemiya et al., 2017), there is minimal research on the network-based
influence on this region.

Another important finding was that connectivity *within* the visual network was positively
associated with motion accuracy, whereas the connectivity between visual and other networks
was negatively associated motion accuracy. Furthermore, follow-up tests demonstrated that the
positive correlation between within-module FC and response accuracy was strongest at the
lowest TMS intensity, when our exogenous stimulation was unlikely to have much of an effect
on cortical excitability. In contrast, increasingly worse performance was associated with more
global connectivity at higher levels of TMS intensity (**Table 1**). Such a benefit to more
segregated cortical networks is well known in developmental (Stevens et al., 2012) and aging
(Betzel et al., 2014; Crowell et al., 2020) literatures, but this is the first finding to suggest an
active, parametric perturbation of cortical modularity and its consequences for behavior. An
alternative explanation for these effects is that greater stimulation intensity also activated
auditory and somatosensory-motor regions, and that this influence of non-visual systems
resulted in more interference to the efficient processing of visual stimuli. However, this view is
not supported by the fact that motion discrimination performance was greater at 120% RMT
than 80% RMT, suggesting a non-linear relationship between delivered E-field and its effect on
behavior in this system.

Lastly, an analysis of SC data revealed that the propagation of these effects was largely
constrained by multiple indices of white matter anatomy. Two related findings support the notion
that the FC patterns associated with localized TMS stimulation are constrained by structural
architecture. First, we examined how FC-SC correlations change as a function of TMS intensity.
Collectively, significant effects of both Intensity and a significant Intensity × FC Type interaction,
as well as the decline in within-module connectivity with increasing intensity all suggest that
TMS disrupted "normal" visual network functioning and introducing exogenous stimulation
disrupted successful motion processing by decoupling FC and SC patterns. Second, and
supporting this inference, an Intensity × FC Type Repeated Measures ANOVA revealed a
pattern that mirrors the FC findings above, such that the coupling between structural and FC
predicted task performance for the local within-module regions, but more global between-
module couplings had no influence on motion perception (**Table 2**). In our study the FC-SC
coupling ranged between r = [0.0–0.3], consistent with other many other investigations between
functional and structural network data relying on one-to-one correspondence between FC and
SC connections (Honey et al., 2007; van den Heuvel et al., 2009; Horn et al., 2014). This
suggests that the upper bound of most FC-SC correlations is not very high (Suarez et al., 2020).
Future approaches to characterizing FS-SC relationships may consider the value of derived
properties of individual nodes that describe their global influence, including modal

controllability—which purports to describe the energy associated with transitioning between brain states (Gu et al., 2017; Beynel et al., 2020b).

While the current study represents a novel application of concurrent TMS-fMRI to answer unexplained phenomena describing the spread of focal neurostimulation across the cortex, several limitations were encountered in the pursuit of this goal. First, the study design did not incorporate sham conditions that closely replicate the sound and scalp sensation of TMS without the delivery of significant E-field to the brain. Nevertheless, the results of the parametric investigation of TMS intensity at the target support a cortical origin of the stimulation effects. Second, because the goal of the current study was to investigate multiple stimulation intensities, it was not feasible to complete within a typical MRI scanning session to assess a complementary stimulation site to examine the potential dissociations between task-relevant and task-irrelevant cortical sites. Lastly, the current design relied on a trial-related timing of 10 Hz pulses concurrent with the task; these design choices represent only one of a number of parameters for frequency, number of pulses, and timing relative to the motion perception task. While the prolonged effects of rTMS are generally assumed to relate to synaptic plasticity, the detailed effects of stimulation on a region's excitability via long-term depression or potentiation remains a matter of investigation.

The observed pattern of findings suggests that inducing a more global pattern of connectivity with higher intensity TMS contributes to disruptions in visual motion perception. Our results show that this spread is largely limited to the network of cortical sites involved in the active task (i.e., visual motion), and that TMS-related disruptions of this network are detrimental to cognitive performance. Lastly, we find that the functional connectivity spread with increasing TMS is largely constrained by structural connectivity, suggesting that the use of structural network information can help to predict at least some of the influence of local perturbations on a more global cortical network. Thus, while it is widely known that TMS may disrupt cognition, these results help us to explain *how* TMS disrupts local and global networks to induce perceptual change.

References

Alekseichuk I, Mantell K, Shirinpour S, Opitz A (2019) Comparative modeling of transcranial magnetic and electric stimulation in mouse, monkey, and human. Neuroimage 194:136-148.

Amemiya T, Beck B, Walsh V, Gomi H, Haggard P (2017) Visual area V5/hMT+ contributes to perception of tactile motion direction: a TMS study. Sci Rep 7:40937.

648   Aydin-Abidin S, Trippe J, Funke K, Eysel UT, Benali A (2008) High- and low-frequency repetitive
649   transcranial magnetic stimulation differentially activates c-Fos and zif268 protein
650   expression in the rat brain. Experimental brain research 188:249-261.
651   Ballesteros S, Reales JM, Mayas J, Heller MA (2008) Selective attention modulates visual and
652   haptic repetition priming: effects in aging and Alzheimer's disease. Experimental brain
653   research 189:473-483.
654   Betzel RF, Byrge L, He Y, Goni J, Zuo XN, Sporns O (2014) Changes in structural and
655   functional connectivity among resting-state networks across the human lifespan.
656   Neuroimage 102 Pt 2:345-357.
657   Beynel L, Powers JP, Appelbaum LG (2020a) Effects of repetitive transcranial magnetic
658   stimulation on resting-state connectivity: A systematic review. Neuroimage:116596.
659   Beynel L, Davis S, Crowell C, Hilbig S, Lim W, Nguyen D, Palmer H, Brito A, Peterchev A,
660   Luber B (2019) Online repetitive transcranial magnetic stimulation during working
661   memory in younger and older adults: A randomized within-subject comparison. PloS one
662   14:e0213707.
663   Beynel L, Deng L, Crowell CA, Dannhauer M, Palmer H, Hilbig S, Peterchev AV, Luber B,
664   Lisanby SH, Cabeza R, Appelbaum LG, Davis SW (2020b) Structural Controllability
665   Predicts Functional Patterns and Brain Stimulation Benefits Associated with Working
666   Memory. The Journal of neuroscience : the official journal of the Society for
667   Neuroscience 40:6770-6778.
668   Beynel L, Davis SW, Crowell CA, Dannhauer M, Lim W, Palmer H, Hilbig SA, Brito A, Hile C,
669   Luber B, Lisanby SH, Peterchev AV, Cabeza R, Appelbaum LG (2020c) Site-Specific
670   Effects of Online rTMS during a Working Memory Task in Healthy Older Adults. Brain
671   Sci 10.
672   Beynel L, Dannhauer M, Palmer H, Hilbig SA, Crowell CA, Wang JE, Michael AM, Wood EA,
673   Luber B, Lisanby SH, Peterchev AV, Cabeza R, Davis SW, Appelbaum LG (2021)
674   Network-based rTMS to modulate working memory: The difficult choice of effective
675   parameters for online interventions. Brain Behav 11:e2361.
676   Brascamp JW, Kanai R, Walsh V, van Ee R (2010) Human middle temporal cortex, perceptual
677   bias, and perceptual memory for ambiguous three-dimensional motion. J Neurosci
678   30:760-766.
679   Campana G, Cowey A, Walsh V (2002) Priming of motion direction and area V5/MT: a test of
680   perceptual memory. Cereb Cortex 12:663-669.
681   Castrillon G, Sollmann N, Kurcyus K, Razi A, Krieg SM, Riedl V (2020) The physiological effects
682   of noninvasive brain stimulation fundamentally differ across the human cortex. Sci Adv
683   6:eaay2739.
684   Cocchi L, Sale MV, L LG, Bell PT, Nguyen VT, Zalesky A, Breakspear M, Mattingley JB (2016)
685   A hierarchy of timescales explains distinct effects of local inhibition of primary visual
686   cortex and frontal eye fields. Elife 5.
687   Conde V, Tomasevic L, Akopian I, Stanek K, Saturnino GB, Thielscher A, Bergmann TO,
688   Siebner HR (2019) The non-transcranial TMS-evoked potential is an inherent source of
689   ambiguity in TMS-EEG studies. Neuroimage 185:300-312.
690   Conforto AB, Z'Graggen WJ, Kohl AS, Rosler KM, Kaelin-Lang A (2004) Impact of coil position
691   and electrophysiological monitoring on determination of motor thresholds to transcranial
692   magnetic stimulation. Clinical neurophysiology : official journal of the International
693   Federation of Clinical Neurophysiology 115:812-819.
694   Cowey A, Campana G, Walsh V, Vaina LM (2006) The role of human extra-striate visual areas
695   V5/MT and V2/V3 in the perception of the direction of global motion: a transcranial
696   magnetic stimulation study. Experimental brain research 171:558-562.
697   Crowell CA, Davis SW, Beynel L, Deng L, Lakhlani D, Hilbig SA, Palmer H, Brito A, Peterchev
698   AV, Luber B, Lisanby SH, Appelbaum LG, Cabeza R (2020) Older adults benefit from

more widespread brain network integration during working memory. Neuroimage
218:116959.
Davis SW, Szymanski A, Boms H, Fink T, Cabeza R (2019) Cooperative contributions of
structural and functional connectivity to successful memory in aging. Netw Neurosci
3:173-194.
Fornito A, Zalesky A, Bullmore ET (2010) Network scaling effects in graph analytic studies of
human resting-state FMRI data. Front Syst Neurosci 4:22.
Gamboa Arana OL, Palmer H, Dannhauer M, Hile C, Liu S, Hamdan R, Brito A, Cabeza R,
Davis SW, Peterchev AV, Sommer MA, Appelbaum LG (2020) Intensity- and timing-
dependent modulation of motion perception with transcranial magnetic stimulation of
visual cortex. Neuropsychologia 147:107581.
Gamboa OL, Brito A, Abzug Z, D'Arbeloff T, Beynel L, Wing EA, Dannhauer M, Palmer H, Hilbig
SA, Crowell CA, Liu S, Donaldson R, Cabeza R, Davis SW, Peterchev AV, Sommer MA,
Appelbaum LG (2020) Application of long-interval paired-pulse transcranial magnetic
stimulation to motion-sensitive visual cortex does not lead to changes in motion
discrimination. Neuroscience letters 730:135022.
Gollo LL, Roberts JA, Cocchi L (2017) Mapping how local perturbations influence systems-level
brain dynamics. Neuroimage.
Gomez LJ, Dannhauer M, Koponen LM, Peterchev AV (2020) Conditions for numerically
accurate TMS electric field simulation. Brain stimulation 13:157-166.
Goni J, van den Heuvel MP, Avena-Koenigsberger A, Velez de Mendizabal N, Betzel RF, Griffa
A, Hagmann P, Corominas-Murtra B, Thiran JP, Sporns O (2014) Resting-brain
functional connectivity predicted by analytic measures of network communication. Proc
Natl Acad Sci U S A 111:833-838.
Gu S, Betzel RF, Mattar MG, Cieslak M, Delio PR, Grafton ST, Pasqualetti F, Bassett DS (2017)
Optimal trajectories of brain state transitions. Neuroimage 148:305-317.
Gullmar D, Haueisen J, Reichenbach JR (2010) Influence of anisotropic electrical conductivity in
white matter tissue on the EEG/MEG forward and inverse solution. A high-resolution
whole head simulation study. Neuroimage 51:145-163.
Hermundstad AM, Bassett DS, Brown KS, Aminoff EM, Clewett D, Freeman S, Frithsen A,
Johnson A, Tipper CM, Miller MB, Grafton ST, Carlson JM (2013) Structural foundations
of resting-state and task-based functional connectivity in the human brain. Proc Natl
Acad Sci U S A 110:6169-6174.
Honey CJ, Kotter R, Breakspear M, Sporns O (2007) Network structure of cerebral cortex
shapes functional connectivity on multiple time scales. Proceedings of the National
Academy of Sciences of the United States of America 104:10240-10245.
Horn A, Ostwald D, Reisert M, Blankenburg F (2014) The structural-functional connectome and
the default mode network of the human brain. Neuroimage 102 Pt 1:142-151.
Ishikawa K, Nagai T, Yamada Y, Negi A, Nakamura M (2011) Optimal conditions for multifocal
VEP recording for normal Japanese population established by receiver operating
characteristic analysis. Doc Ophthalmol 122:29-37.
Johansen-Berg H, Matthews PM (2002) Attention to movement modulates activity in sensori-
motor areas, including primary motor cortex. Experimental brain research 142:13-24.
Kallioniemi E, Saisanen L, Kononen M, Awiszus F, Julkunen P (2014) On the estimation of
silent period thresholds in transcranial magnetic stimulation. Clinical neurophysiology :
official journal of the International Federation of Clinical Neurophysiology 125:2247-
2252.
Luber B, Deng Z-D (2016) Application of non-invasive brain stimulation in psychophysiology. In:
Handbook of Psychophysiology (Berntson GG, Cacioppo JT, Tassinary LG, eds), pp
116-150. Cambridge: Cambridge University Press.

Luber B, Davis SW, Deng Z-D, Murphy D, Peterchev AV, Lisanby SH (in press) Using diffusion
tensor imaging to effectively target TMS to deep brain structures. NeuroImage.
Momi D, Ozdemir RA, Tadayon E, Boucher P, Shafi MM, Pascual-Leone A, Santarnecchi E
(2021a) Network-level macroscale structural connectivity predicts propagation of
transcranial magnetic stimulation. Neuroimage 229:117698.
Momi D, Ozdemir RA, Tadayon E, Boucher P, Di Domenico A, Fasolo M, Shafi MM, Pascual-
Leone A, Santarnecchi E (2021b) Perturbation of resting-state network nodes
preferentially propagates to structurally rather than functionally connected regions. Sci
Rep 11:12458.
Ozdemir RA, Tadayon E, Boucher P, Momi D, Karakhanyan KA, Fox MD, Halko MA, Pascual-
Leone A, Shafi MM, Santarnecchi E (2020) Individualized perturbation of the human
connectome reveals reproducible biomarkers of network dynamics relevant to cognition.
Proc Natl Acad Sci U S A 117:8115-8125.
Peterchev AV, Wagner TA, Miranda PC, Nitsche MA, Paulus W, Lisanby SH, Pascual-Leone A,
Bikson M (2012) Fundamentals of transcranial electric and magnetic stimulation dose:
definition, selection, and reporting practices. Brain stimulation 5:435-453.
Rafiei F, Rahnev D (2021) Does TMS increase BOLD activity at the site of stimulation?
[Preprint].
Reichardt J, Bornholdt S (2006) Statistical mechanics of community detection. Phys Rev E Stat
Nonlin Soft Matter Phys 74:016110.
Sanders LL, Auksztulewicz R, Hohlefeld FU, Busch NA, Sterzer P (2014) The influence of
spontaneous brain oscillations on apparent motion perception. Neuroimage 102 Pt
2:241-248.
Silvanto J, Cattaneo Z (2021) Nonlinear interaction between stimulation intensity and initial brain
state: Evidence for the facilitatory/suppressive range model of online TMS effects.
Neuroscience letters 742:135538.
Silvanto J, Lavie N, Walsh V (2005) Double dissociation of V1 and V5/MT activity in visual
awareness. Cerebral cortex 15:1736-1741.
Silvanto J, Bona S, Marelli M, Cattaneo Z (2018) On the Mechanisms of Transcranial Magnetic
Stimulation (TMS): How Brain State and Baseline Performance Level Determine
Behavioral Effects of TMS. Frontiers in psychology 9:741.
Smith RE, Tournier JD, Calamante F, Connelly A (2013) SIFT: Spherical-deconvolution
informed filtering of tractograms. Neuroimage 67:298-312.
Stephan KE, Kasper L, Harrison LM, Daunizeau J, den Ouden HE, Breakspear M, Friston KJ
(2008) Nonlinear dynamic causal models for fMRI. Neuroimage 42:649-662.
Stevens AA, Tappon SC, Garg A, Fair DA (2012) Functional brain network modularity captures
inter- and intra-individual variation in working memory capacity. PloS one 7:e30468.
Suarez LE, Markello RD, Betzel RF, Misic B (2020) Linking Structure and Function in
Macroscale Brain Networks. Trends Cogn Sci 24:302-315.
Thielscher A, Antunes A, Saturnino GB (2015) Field modeling for transcranial magnetic
stimulation: a useful tool to understand the physiological effects of TMS? Ieee Eng Med
Bio:222-225.
van den Heuvel MP, Mandl RC, Kahn RS, Hulshoff Pol HE (2009) Functionally linked resting-
state networks reflect the underlying structural connectivity architecture of the human
brain. Hum Brain Mapp 30:3127-3141.
Wang JX, Voss JL (2015) Long-lasting enhancements of memory and hippocampal-cortical
functional connectivity following multiple-day targeted noninvasive stimulation.
Hippocampus 25:877-883.
Wang JX, Rogers LM, Gross EZ, Ryals AJ, Dokucu ME, Brandstatt KL, Hermiller MS, Voss JL
(2014) Targeted enhancement of cortical-hippocampal brain networks and associative
memory. Science 345:1054-1057.

Zinchenko A, Brunner S, Chen L, Shi Z, Taylor PCJ, Muller HJ (2021) V5/MT+ modulates
spatio-temporal integration differently across and within hemifields: Causal evidence
from TMS. Neuropsychologia 161:107995.