1 **Generalized Michaelis–Menten rate law**
2 **with time-varying molecular concentrations**
3

4 Roktaek Lim[1,7], Thomas L. P. Martin[1,7], Junghun Chae[2,7], WooJoong Kim[2],

5 Cheol-Min Ghim[2,3,*], and Pan-Jun Kim[1,4,5,6,*]

6
[1]Department of Biology, Hong Kong Baptist University, Kowloon, Hong Kong

[2]Department of Physics, Ulsan National Institute of Science and Technology, Ulsan 44919,

Republic of Korea

[3]Department of Biomedical Engineering, Ulsan National Institute of Science and Technology,

Ulsan 44919, Republic of Korea

[4]Center for Quantitative Systems Biology & Institute of Computational and Theoretical

Studies, Hong Kong Baptist University, Kowloon, Hong Kong

[5]State Key Laboratory of Environmental and Biological Analysis, Hong Kong Baptist

University, Kowloon, Hong Kong

[6]Abdus Salam International Centre for Theoretical Physics, 34151 Trieste, Italy

[7]R.L., T.L.P.M., and J.C. contributed equally to this work.

[*]Correspondence:
cmghim@unist.ac.kr (C.-M.G.); panjunkim@hkbu.edu.hk or extutor@gmail.com (P.-J.K.)

20
21 **Abstract**
22

The Michaelis–Menten (MM) rate law has been the dominant paradigm of modeling

biochemical rate processes for over a century with applications in biochemistry, biophysics,

cell biology, systems biology, and chemical engineering. The MM rate law and its remedied

form stand on the assumption that the concentration of the complex of interacting

molecules, at each moment, approaches an equilibrium (quasi-steady state) much faster

than the molecular concentrations change. Yet, this assumption is not always justified. Here,

we relax this quasi-steady state requirement and propose the generalized MM rate law for

the interactions of molecules with active concentration changes over time. Our approach for

time-varying molecular concentrations, termed the effective time-delay scheme (ETS), is

based on rigorously estimated time-delay effects in molecular complex formation. With

particularly marked improvements in protein–protein and protein–DNA interaction

modeling, the ETS provides an analytical framework to interpret and predict rich transient or

rhythmic dynamics (such as autogenously-regulated cellular adaptation and circadian protein

turnover), which goes beyond the quasi-steady state assumption.

Introduction

Since proposed by Henri [1] and Michaelis and Menten [2], the Michaelis–Menten (MM) rate law has been the dominant framework for modeling the rates of enzyme-catalyzed reactions for over a century [1–4]. The MM rate law has also been widely adopted for describing other bimolecular interactions, such as reversible binding between proteins [5–7], between a gene and a transcription factor [8,9], and between a receptor and a ligand [10,11]. The MM rate law hence serves as a common mathematical tool in both basic and applied fields, including biochemistry, biophysics, pharmacology, systems biology, and many subfields of chemical engineering [12]. The derivation of the MM rate law from the underlying biochemical mechanism is based on the steady-state approximation by Briggs and Haldane [3], referred to as the *standard quasi-steady state approximation* (sQSSA) [12–17]. The sQSSA, however, is only valid when the enzyme concentration is low enough and thus the concentration of enzyme–substrate complex is negligible compared to substrate concentration [14]. This condition may be acceptable for many metabolic reactions with substrate concentrations that are typically far higher than the enzyme concentrations.

Nevertheless, in the case of protein–protein interactions in various cellular activities, the interacting proteins as the "enzymes" and "substrates" often show the concentrations comparable with each other [18–20]. Therefore, the use of the MM rate law for describing protein–protein interactions has been challenged in its rationale, with the modified alternative formula from the *total quasi-steady state approximation* (tQSSA) [12,13,21–27]. The tQSSA-based form is generally more accurate than the MM rate law from the sQSSA, for a broad range of combined molecular concentrations and thus for protein–protein interactions as well [12,13,21–27]. The superiority of the tQSSA has not only been proven in the quantitative, but also in the qualitative outcomes of systems, which the sQSSA sometimes fails to predict [12,21]. Later, we will provide the overview of the tQSSA and its relationship with the conventional MM rate law from the sQSSA.

Despite the correction of the MM rate law by the tQSSA, both the tQSSA and sQSSA still rely on the assumption that the concentration of the complex of interacting molecules, at each moment, approaches an equilibrium (quasi-steady state) much faster than the molecular concentrations change [12,14,24]. Although this quasi-steady state assumption may work for a range of biochemical systems, the exact extent of such systems to follow that assumption is not clear. Numerous cellular processes do exhibit active molecular concentration changes over time, such as in signal responses, circadian oscillations, and cell

cycles [6,7,21,28–31], calling for a better approach to even cover the time-varying molecular

concentrations that may not strictly adhere to the quasi-steady state assumption.

In this study, we report the generalization of the MM rate law, whereby the interaction of

time-varying molecular components is more properly described than by the tQSSA and

sQSSA. This generalization is the correction of the tQSSA with rigorously estimated, time-

delay effects affected by free molecule availability. Our formulation, termed the effective

time-delay scheme (ETS), well accounts for the transient or oscillatory dynamics and

experimental data patterns of biochemical systems with the relevant analytical insights,

which are not captured by the previous methods. Surprisingly, we reveal that the existing

quasi-steady state assumption can even fail for extremely slow changes in protein

concentrations under autogenous regulation, whereas the ETS does not. In addition, the ETS

allows the natural explanation of rhythmic degradation of circadian proteins without

requiring explicitly-rhythmic post-translational mechanisms; this is not straightforward

within the quasi-steady state assumption. As an added feat, the ETS improves kinetic

parameter estimation. As demonstrated in a number of contexts such as autogenously-

regulated cellular adaptation and circadian oscillations, our approach offers a useful

theoretical framework to interpret and predict rich transient or rhythmic dynamics of

biochemical systems with a wide range of applicability.

89
90 **Results**
91
92 **Theory development**
93
First, we present the outline of the tQSSA, sQSSA, and our generalized MM rate law.

Consider two different molecules A and B that bind to each other and form complex AB, as

illustrated in Fig. 1(a). For example, A and B may represent two participant proteins in

heterodimer formation, a chemical substrate and an enzyme in a metabolic reaction, and a

solute and a transporter in membrane transport. The concentration of the complex AB at

time t, denoted by $C(t)$, changes over time as in the following equation from the mass-

action law:

$$\frac{\mathrm{d}C(t)}{\mathrm{d}t} = k_a[A(t) - C(t)][B(t) - C(t)] - k_\delta C(t). \tag{1}$$

Here, $A(t)$ and $B(t)$ denote the total concentrations of A and B, respectively, and hence

$A(t) - C(t)$ and $B(t) - C(t)$ are the concentrations of free A and B. The temporal profiles

of $A(t)$ and $B(t)$ are allowed to be very generic, e.g., even with their own feedback effects

as addressed later. k_a denotes the association rate of free A and B. k_δ is the effective

"decay" rate of AB with $k_\delta \equiv k_\mathrm{d} + r_\mathrm{c} + k_\mathrm{loc} + k_\mathrm{dlt}$ where k_d, k_loc, and k_dlt stand for the

dissociation, translocation, and dilution rates of AB, respectively, and r_c for the chemical

conversion or translocation rate of A or B upon the formation of AB. In other words, for the

sake of generality, k_δ is not limited to a dissociation event but encompasses all rate events

to lower the level of AB [Fig. 1(a)].

In the tQSSA, the assumption is that $C(t)$ approaches the quasi-steady state fast enough

each time, given the values of $A(t)$ and $B(t)$ [12,24]. This assumption and the notation $K \equiv$

k_δ/k_a lead Eq. (1) to an estimate $C(t) \approx C_\mathrm{tQ}(t)$ with the following form (Text S1):

$$C_\mathrm{tQ}(t) \equiv \tfrac{1}{2}\{K + A(t) + B(t) - K\Delta_\mathrm{tQ}(t)\}, \tag{2}$$

$$\Delta_\mathrm{tQ}(t) \equiv \sqrt{\left[1 + \frac{A(t)+B(t)}{K}\right]^2 - \frac{4}{K^2}A(t)B(t)}$$

$$= \sqrt{1 + 2\left[\frac{A(t)+B(t)}{K}\right] + \left[\frac{A(t)-B(t)}{K}\right]^2}. \tag{3}$$

Although the tQSSA looks a little complex, it only involves a single parameter K and is easy to

implement in a computer program. As mentioned earlier, the tQSSA is generally more

accurate than the conventional MM rate law [12,13,21–27]. To obtain the MM rate law,

consider a rather specific condition,

$$B(t) \ll K + A(t) \text{ or } A(t) \ll K + B(t). \tag{4}$$

In this condition, the Padé approximant for $C_\mathrm{tQ}(t)$ takes the following form:

$$C_\mathrm{tQ}(t) \approx \frac{A(t)B(t)}{K+A(t)+B(t)}. \tag{5}$$

Considering Eq. (5), Eq. (4) is similar to the condition $C_\mathrm{tQ}(t)/A(t) \ll 1$ or $C_\mathrm{tQ}(t)/B(t) \ll 1$.

In other words, Eq. (5) would be valid when the concentration of AB complex is negligible

compared to either A or B's concentration. This condition is essentially identical to the

assumption in the sQSSA resulting in the MM rate law [14]. In the example of a typical

metabolic reaction with $B(t) \ll A(t)$ for substrate A and enzyme B, Eq. (4) is automatically

satisfied and Eq. (5) further reduces to the familiar MM rate law $C_\mathrm{tQ}(t) \approx A(t)B(t)/[K +$

$A(t)]$, i.e., the outcome of the sQSSA [1–4,12–14]. To be precise, the sQSSA uses the

concentration of free A instead of $A(t)$, but we refer to this formula with $A(t)$ as the sQSSA

because the complex is assumed to be negligible in that scheme. Clearly, K here is the

Michaelis constant, commonly known as K_M.

The application of the MM rate law beyond the condition in Eq. (4) invites a risk of

erroneous modeling results, whereas the tQSSA is relatively free of such errors and has wider

applicability [12,13,21–27]. Still, both the tQSSA and sQSSA stand on the assumption that

$C(t)$ approaches the quasi-steady state fast enough each time before the marked temporal

change of $A(t)$ or $B(t)$. We now relax this quasi-steady state assumption and generalize the

approximation of $C(t)$ to the case of time-varying $A(t)$ and $B(t)$, as the main objective of

this study.

150
151

152
Fig. 1. Generalization of the MM rate law for time-varying molecular concentrations, referred to as

the ETS. (a) Two different molecules A and B bind to each other and form their complex. (b) A TF binds

to a DNA molecule to regulate mRNA expression (RNA polymerase and other molecules are omitted

here). In (a) and (b), the graphs show the comparison among the exact time-course profile of the

complex concentration, the tQSSA-based (a) or QSSA-based (b) profile, and the ETS-based profile. The

relationship between the tQSSA (or QSSA) and the ETS is illustrated through the effective time delay in

the ETS. Notations k_a, k_d, k_δ, t, $\Delta_{tQ}(t)$, K, and $A_{TF}(t)$ are defined in the description of Eqs. (1)–(3)

and (6)–(8). Simulations in (a) and (b) are based on periodic oscillation models in Texts S7 and S8,

respectively, with their parameters in Table S7.

Suppose that $C(t)$ may not necessarily approach the quasi-steady state each time but stays

within some distance from it. As detailed in Text S1, we linearize the right-hand side of Eq.

(1) around $C(t) - C_{tQ}(t)$ and estimate $C(t)$'s solution as the time integral of $C_{tQ}(t')$ (where

$t' \leq t$) with an exponential kernel-like function. The Taylor expansion of $C_{tQ}(t')$ by $t - t'$ is

incorporated into this integral and then its form offers the following approximant for $C(t)$:

$$C_\gamma(t) \equiv \min\left\{C_{tQ}\left\{t - [k_\delta \Delta_{tQ}(t)]^{-1}\right\}, A(t), B(t)\right\}. \tag{6}$$

Although the above $C_\gamma(t)$ looks rather complex, this form is essentially a simple conversion

$t \to t - [k_\delta \Delta_{tQ}(t)]^{-1}$ in the tQSSA. $\min\{\cdot\}$ is just taken for a minor role to ensure that the

complex concentration cannot exceed $A(t)$ or $B(t)$. Hence, the distinct feature of $C_\gamma(t)$ is

the inclusion of an effective time delay $[k_\delta \Delta_{tQ}(t)]^{-1}$ in complex formation. This delay is the

rigorous estimate of the molecular relaxation time during which the effect of instantaneous

$A(t)$ and $B(t)$ is notably sustained in the complex formation, as shown in Text S1. We will

refer to this formulation as the effective time-delay scheme (ETS), and its relationship with

the tQSSA is depicted in Fig. 1(a).

We propose the ETS as the generalization of the MM rate law for time-varying molecular

concentrations that may not strictly adhere to the quasi-steady state assumption. If the

relaxation time in complex formation is so short that the effective time delay in Eq. (6) can

be ignored, the ETS returns to the tQSSA in its form. Surprisingly, we proved that, unlike the

ETS, any simpler new rate law without a time-delay term would not properly work for active

concentration changes over time (Text S3). Nevertheless, one may question the analytical

utility of the ETS, regarding the apparent complexity of its time-delay term. In the examples

of autogenously-regulated cellular adaptation and rhythmic protein turnover below, we will

use the ETS to deliver valuable analytical insights into the systems whose dynamics is

otherwise ill-explained by the conventional approaches.

About the physical interpretation of the ETS, we notice that the effective time delay is

inversely linked to free molecule availability, as $[k_\delta \Delta_{tQ}(t)]^{-1} = k_\delta^{-1}\{1 + K^{-1}[A(t) +$

$B(t) - 2C_{tQ}(t)]\}^{-1}$ from Eq. (2). Here, $A(t) + B(t) - 2C_{tQ}(t) = [A(t) - C_{tQ}(t)] +$

$[B(t) - C_{tQ}(t)]$, which is the total free molecule concentration at the quasi-steady state

each time. In other words, the less the free molecules, the more the time delay, which is at

most k_δ^{-1}. One can understand this observation as follows: $-k_\delta C(t)$ in Eq. (1) gives the

expectation that the decay time-scale (k_δ^{-1}) of the complex may approximate the relaxation

time. Yet, the relaxation time is shorter than k_δ^{-1}, because free A and B are getting depleted

over time as a result of their complex formation and therefore the complex formation rate

$k_a[A(t) - C(t)][B(t) - C(t)]$ in Eq. (1) continues to decline towards quicker relaxation of

the complex level. This free-molecule depletion effect to shorten the relaxation time is

roughly proportional to the free molecule concentration itself (Text S1). Hence, the

relaxation time takes a decreasing function of the free molecule concentration, consistent

with the above observation. Clearly, the free molecule concentration would be low for

relatively few A and B molecules with comparable concentrations—i.e., small $A(t) + B(t)$

and $[A(t) - B(t)]^2$ in Eq. (3). In this case, the relaxation time would be relatively long and

the ETS shall be deployed instead of the tQSSA or sQSSA. We thus expect that protein–

protein interactions would often be the cases in need of the ETS compared to metabolic

reactions with much excess substrates not binding to enzymes, as will be shown later.

Thus far, we have implicitly assumed the continuous nature of molecular concentrations as

in Eq. (1). However, there exist biomolecular events that fundamentally deviate from this

assumption. For example, a transcription factor (TF) binds to a DNA molecule in the nucleus

to regulate mRNA expression and the number of such a TF–DNA assembly would be either 1

or 0 for a DNA site that can afford at most one copy of the TF [Fig. 1(b)]. This inherently

discrete and stochastic nature of the TF–DNA assembly is seemingly contrasted with the

continuous and deterministic nature of the molecular complex in Eq. (1). To rigorously

describe this TF–DNA binding dynamics, we harness the chemical master equation [32] and

introduce quantities $A_{TF}(t)$ and $C_{TF}(t)$, which are the total TF concentration and the TF–

DNA assembly concentration averaged over the cell population, respectively (Text S4).

According to our calculation, the quasi-steady state assumption leads to the following

approximant for $C_{TF}(t)$:

$$C_{TFQ}(t) \equiv \frac{A_{TF}(t)}{V[K + A_{TF}(t)]},$$ (7)

where $K \equiv k_\delta/k_a$ with k_a and k_δ as the TF–DNA binding and unbinding rates, respectively,

and V is the nuclear volume (Text S4). In fact, $C_{TFQ}(t)$ in Eq. (7) corresponds to a special case

of the previously-studied, *stochastic quasi-steady state approximation* (stochastic QSSA)

[33,34] for arbitrary molecular copy numbers such as for multiple DNA binding sites. Of note,

the stochastic QSSA becomes close to the tQSSA as its deterministic version, if $VK\Delta_{tQ}(t) \gg 1$ [34].

$C_{TFQ}(t)$ in Eq. (7) looks very similar to the MM rate law, considering the "concentration" of

the DNA site (V^{-1}). Nevertheless, $C_{TFQ}(t)$ is not a mere continuum of Eq. (5), because the

denominator in $C_{TFQ}(t)$ includes $K + A_{TF}(t)$, but not $K + A_{TF}(t) + V^{-1}$. In fact, the

discrepancy between $C_{TFQ}(t)$ and Eq. (5) comes from the inherent stochasticity in the TF–

DNA assembly (Text S4). In this regard, directly relevant to $C_{\text{TFQ}}(t)$ is the stochastic version

of the MM rate law with denominator $K + A(t) + B(t) - V^{-1}$ proposed by Levine and Hwa

[35], because the DNA concentration $B(t)$ is V^{-1} in our case. $C_{\text{TFQ}}(t)$ is a fundamentally

more correct approximant for the DNA-binding TF level than both the tQSSA and sQSSA in

Eqs. (2) and (5). Therefore, we will just refer to $C_{\text{TFQ}}(t)$ as the QSSA for the TF–DNA

interactions.

Still, the use of $C_{\text{TFQ}}(t)$ stands on the quasi-steady state assumption. We relax this

assumption and generalize the approximation of $C_{\text{TF}}(t)$ to the case of time-varying TF

concentration. In a similar way to obtain $C_{\gamma}(t)$ in Eq. (6), we propose the following

approximant for $C_{\text{TF}}(t)$ (Text S4):

$$C_{\text{TF}\gamma}(t) \equiv C_{\text{TFQ}}\left[t - \frac{k_\delta^{-1}K}{K + A_{\text{TF}}(t)}\right]. \qquad (8)$$

This formula represents the TF–DNA version of the ETS, and its relationship with the QSSA is

illustrated in Fig. 1(b). The time-delay term in Eq. (8) has a similar physical interpretation to

that in Eq. (6). Besides, this term is directly proportional to the probability of the DNA

unoccupancy at the quasi-steady state, according to Eq. (7).

Through numerical simulations of various systems with empirical data analyses, we found

that the ETS provides the reasonably accurate description of the deviations of time-course

molecular profiles from the quasi-steady states (Texts S6–S8). This result was particularly

evident for the cases of protein–protein and TF–DNA interactions with time-varying protein

concentrations. In these cases, the ETS unveils the importance of the relaxation time

(effective time delay) in complex formation to the shaping of molecular profiles, otherwise

difficult to clarify. Yet, the use of the sQSSA or tQSSA is practically enough for typical

metabolic reaction and transport systems, without the need for the ETS. The strict

mathematical conditions for the validity of the ETS as well as those of the quasi-steady state

assumption are derived in Text S5.

259
260 **Autogenous control**

261
Adaptation to changing environments is a process of biological control. The ETS offers an

analytical tool for understanding transient dynamics of such adaptation processes,

exemplified by autogenously regulated systems where TFs regulate their own transcription.

This autogenous control underlies cellular responses to various internal and external stimuli

[36,37]. We here explore the case of positive autoregulation and show that the quasi-steady

state assumption does not even work for extremely-slow protein changes near a tipping

point. The case of negative autoregulation is covered in Text S10.

In the case of positive autoregulation, consider a scenario in Fig. 2(a) that proteins enhance

their own transcription after homodimer formation and this dimer–promoter interaction is

facilitated by inducer molecules. The inherent cooperativity from the dimerization is known

to give a sigmoid TF–DNA binding curve, resulting in abrupt and history-dependent transition

events [36,38]. We here built the full kinetic model of the system without the ETS, tQSSA, or

other approximations of the dimerization and dimer–promoter interaction (Text S9). As the

simulated inducer level increases, Fig. 2(b) demonstrates that an initially low, steady-state

protein level undergoes an abrupt leap at some point η_c, known as a transition or tipping

point. This discontinuous transition with only a slight inducer increase signifies a qualitative

change in the protein expression state. Reducing the inducer level just back to the transition

point η_c does not reverse the protein state, which is sustained until more reduction in the

inducer level [Fig. 2(b)]. This history-dependent behavior, hysteresis, indicates the

coexistence of two different stable states of the protein level (bistability) between the

forward and backward transitions [36,38].

Other than steady states, we examine how fast the system responds to signals. Upon acute

induction from a zero to certain inducer level ($> \eta_c$), the protein level grows over time

towards its new steady state and this response becomes rapider at stronger induction away

from the transition point [Fig. 2(c)]. Conversely, as the inducer level decreases towards the

transition point, the response time continues to increase and eventually becomes diverging

(in this study, response time is defined as the time taken for a protein level to reach 90% of

its steady state). This phenomenon has been called "critical slowing down" [39–41].

Regarding this near-transition much slow protein growth, one may expect that the quasi-

steady state assumption would work properly near that transition point. To test this

possibility, we modified the full model by the tQSSA and QSSA of the dimerization and dimer-

promoter interaction, respectively, and call this modified model the QSSA-based model. For

comparison, we created another version of the model by the ETS of the dimerization and

dimer-promoter interaction and call this version the ETS-based model (Text S9). Across

physiologically-relevant parameter conditions, we compared the QSSA- and ETS-based model

simulation results to the full model's (Text S12 and Table S5). Surprisingly, the QSSA-based

model often severely underestimated the response time, particularly near a transition point,

while the ETS-based response time was relatively close to that from the full model [$P <$

10^{-4} and Text S12; e.g., 8.5-hour shorter and 0.5-hour longer response times in Fig. 2(c)

(left) in the QSSA and ETS cases, respectively].

This unexpected mismatch between the QSSA and full model results comes from the

following factors: because the QSSA model discards the effective time delay in dimerization

and dimer-promoter interaction, this model accelerates positive feedback, transcription, and

protein production, and thus shortens the response time. Near the transition point, although

the protein level grows very slowly, a little higher transcription activity in the QSSA model

substantially advances the protein growth with near-transition ultrasensitivity that we

indicated above. Therefore, the QSSA model shortens the response time even near the

transition point.

(a)

(b)

(c)

(d)

Fig. 2. Positive autoregulation and induction kinetics. (a) Protein production mechanism with positive

autoregulation in the presence of inducers. (b) Bifurcation diagram of the simulated protein level as a

function of η (proxy for an inducer level). The steady state is plotted as η increases (solid line) or

decreases (dashed line). Acute induction can be simulated by a sudden change of $\eta = 0$ to $\eta > \eta_c$ in

the shaded area. (c) Time-series of protein levels from the full, ETS, and QSSA models upon acute

induction at time 0 h with $\eta = 2.42$ (left) or $\eta = 200$ (right). (d) The full model-to-QSSA difference in

response time as a function of $1/\sqrt{(\eta - \eta_c)/\eta_c}$. Both the simulated and analytically-estimated

differences are presented. The analytical estimation is based on Eq. (9). For more details of (b)–(d),

refer to Text S9 and Table S8.

Related to this point, the ETS allows the analytical calculation of response time and its QSSA-

based estimate. In this calculation, we considered two different stages of protein growth—its

early and late stages (Text S9) and found that the QSSA model underestimates response time

mainly at the early stage. This calculation suggests that the exact response time would be

longer than the QSSA-based estimate by

328
$$\frac{2\pi}{r\sqrt{\frac{\eta-\eta_c}{\eta_c}}}\left(\frac{1}{D}+\frac{1}{D_{TF}}\right)+\frac{1}{r}\ln\left(1+\frac{1}{D}+\frac{1}{D_{TF}}\right)+\frac{1}{r}\left(\frac{1}{D}+\frac{1}{D_{TF}}\right)\ln\left\{1+\frac{DD_{TF}(\bar{u}-1)[DD_{TF}(\bar{u}-1)-2(D+D_{TF})]}{(D+D_{TF})^2}\right\}, (9)$$

where η and η_c denote an inducer level and its value at the transition point, respectively, r is

the sum of protein degradation and dilution rates, and D and D_{TF} are parameters inversely

proportional to the effective time delays in dimerization and dimer–promoter interaction,

respectively. The additional details and the definition of parameter $\bar{u}$ are provided in Text S9.

Notably, the above response time difference vanishes as $D^{-1} + D_{TF}^{-1} \to 0$. In other words,

the total effective time delay is responsible for this response time difference. Strikingly, this

difference indefinitely grows as η decreases towards η_c, as a linear function of

$1/\sqrt{(\eta-\eta_c)/\eta_c}$. This prediction can serve as a testbed for our theory and highlights far

excessive elongation of near-transition response time (compared to the QSSA) as an

amplified effect of the relaxation time in complex formation. This amplified effect is the

result of the near-transition ultrasensitivity that we indicated above. Consistent with our

prediction, the full model simulation always shows longer response time than the QSSA

model simulation and the difference is linearly scaled to $1/\sqrt{(\eta-\eta_c)/\eta_c}$ as exemplified by

Fig. 2(d) ($R^2 > 0.98$ in simulated conditions; see Text S12). Moreover, its predicted slope

against $1/\sqrt{(\eta-\eta_c)/\eta_c}$ [i.e., 2π (6.28 $\cdots$) multiplied by $\left(D_{TF}^{-1} + D^{-1}\right)r^{-1}$] is comparable

with the simulation results [7.3 ± 0.3 (avg. ± s.d. in simulated conditions) multiplied by

$\left(D_{TF}^{-1} + D^{-1}\right)r^{-1}$; see Text S12]. The agreement of these nontrivial predictions with the

numerical simulation results proves the theoretical value of the ETS. Again, we raise a

caution against the quasi-steady state assumption, which unexpectedly fails for very slow

dynamics with severe underestimation of response time, e.g., by a few tens of hours in the

case of Fig. 2(d).

352 Rhythmic degradation of circadian proteins

Circadian clocks in various organisms generate endogenous molecular oscillations with ~24 h

periodicity, enabling physiological adaptation to diurnal environmental changes caused by

the Earth's rotation around its axis. Circadian clocks play a pivotal role in maintaining

biological homeostasis, and the disruption of their function is associated with a wide range

of pathophysiological conditions [7,9,21,28–30]. According to previous reports, some

circadian clock proteins are not only rhythmically produced but also decompose with

rhythmic degradation rates [Figs. 3(a) and 3(b)] [42–46]. Recently, we have suggested that

the rhythmic degradation rates of proteins with circadian production can spontaneously

emerge without any explicitly time-dependent regulatory mechanism of the degradation

processes [42,47]. If the rhythmic degradation rate peaks at the descending phase of the

protein profile and stays relatively low elsewhere, it is supposed to save much of the

biosynthetic cost in maintaining a circadian rhythm. A degradation mechanism with multiple

post-translational modifications (PTMs), such as phospho-dependent ubiquitination, may

elevate the rhythmicity of this degradation rate in favor of the biosynthetic cost saving

[42,45]. Can the ETS explain this inherent rhythmicity in the degradation rates of circadian

proteins?

First, we constructed the kinetic model of circadian protein production and degradation

without the ETS or other approximations. This model attributes a circadian production rate

of the protein to a circadian mRNA expression or translation rate. Yet, a protein degradation

rate in the model is not based on any explicitly time-dependent regulatory processes, but on

constantly-maintained proteolytic mediators such as constant E3 ubiquitin ligases and

kinases. In realistic situations, the protein turnover may require multiple preceding PTMs,

like mono- or multisite phosphorylation and subsequent ubiquitination. Our model covers

these cases, as well. The model comprises the following equations:

379
$$\frac{\mathrm{d}A_0(t)}{\mathrm{d}t} = g(t) - a_0 A_0(t), \tag{10}$$

381
$$\frac{\mathrm{d}A_i(t)}{\mathrm{d}t} = a_{i-1} A_{i-1}(t) - a_i A_i(t), \tag{11}$$

where $A_0(t)$ and $A_i(t)$ represent the concentrations of the unmodified and i-th modified

proteins, respectively ($i = 1, 2, \cdots, n$ and n is the total number of the PTMs with $n \geq 1$), $g(t)$

is the protein production rate through mRNA expression and translation, a_i denotes the

protein's $(i + 1)$-th modification rate ($i = 0, 1, \cdots, n - 1$), and a_n denotes the turnover rate

of the n-th modified protein.

Next, we apply the ETS to the PTM processes in the model for the analytical estimation of

the protein degradation rate. We observed the mathematical equivalence of the PTM

processes and the above-discussed TF–DNA interactions, despite their different biological

contexts (Text S11). This observation leads to the estimate $r_\gamma(t)$ of the protein degradation

rate as

394
$$r_\gamma(t) \equiv \frac{a_v}{A(t)} \min\left[\frac{a_u}{a_u+a_v} A\left(t - \frac{1}{a_u+a_v}\right), A(t)\right] \approx \frac{a_u a_v}{a_u+a_v}\left\{1 - \frac{1}{a_u+a_v}\left[\frac{1}{A(t)}\frac{\mathrm{d}A(t)}{\mathrm{d}t}\right] + \cdots\right\}, \tag{12}$$

where $A(t)$ is the total protein concentration, a_u and a_v are the rates of the two slowest

PTM and turnover steps in the protein degradation pathway (the step of a_u precedes that of

a_v; see Text S11), and the rightmost formula is to simplify $r_\gamma(t)$ with the Taylor expansion.

The use of $r_\gamma(t)$ may not satisfactorily work for the degradation depending on many

preceding PTMs, but still helps to capture the core feature of the dynamics.

Strikingly, the quasi-steady state assumption does not predict a rhythmic degradation rate,

as the QSSA version of Eq. (12) gives rise to a constant degradation rate, $a_u a_v/(a_u + a_v)$

(Text S11). In contrast, the ETS naturally accounts for the degradation rhythmicity through

the effective time delay in the degradation pathway. The rightmost formula in Eq. (12)

indicates that the degradation rate would be an approximately increasing function of

$-A'(t)/A(t)$ and thus increase as time goes from the ascending to descending phase of the

protein profile. This predicted tendency well matches the experimental data patterns in Figs.

3(a) and 3(b). Fundamentally, this degradation rhythmicity roots in the unsynchronized

interplay between protein translation, modification, and turnover events [42]. For example,

in the case of protein ubiquitination, ubiquitin ligases with a finite binding affinity would not

always capture all newly-translated substrates, and therefore a lower proportion of the

substrates can be ubiquitinated during the ascending phase of the substrate profile than

during the descending phase. The degradation rate partially follows this ubiquitination

pattern. Additional PTMs like phosphorylation, if required for the ubiquitination, can further

retard the full substrate modification and thereby increase the degradation rhythmicity for a

given substrate profile. One may expect that these effects would be enhanced with more

limited ubiquitin ligases or kinases, under the condition when the substrate level shows a

strong oscillation. This expectation is supported by the relative amplitude of the degradation

rate estimated by Eq. (12):

$$\frac{\max_t[r_\gamma(t)] - \min_t[r_\gamma(t)]}{\langle r_\gamma(t)\rangle} \approx \frac{1}{a_u + a_v}\left\{\max_t\left[\frac{1}{A(t)}\frac{\mathrm{d}A(t)}{\mathrm{d}t}\right] - \min_t\left[\frac{1}{A(t)}\frac{\mathrm{d}A(t)}{\mathrm{d}t}\right]\right\}, \qquad (13)$$

where $\langle\cdot\rangle$ denotes a time average. Here, the relative amplitude of the degradation rate is

proportional to $1/(a_u + a_v)$ as well as to the amplitude of $A'(t)/A(t)$. Therefore, limited

ubiquitin ligases or kinases, and strong substrate oscillations increase the rhythmicity of the

degradation rate. Given a substrate profile, multiple PTMs can further enhance this

degradation rhythmicity because they invite the possibility of smaller a_u and a_v values than

expected for the case of only a single PTM. Moreover, Eq. (12) predicts that the degradation

rate would peak around the peak time of $-A'(t)/A(t)$.

In the example of Fig. 3(c) for a single PTM case, the simulated degradation rate from the

aforementioned full kinetic model exhibits the rhythmic profile in excellent agreement with

the ETS-predicted profile. Notably, the peak time of the simulated degradation rate is very

close to that of $-A'(t)/A(t)$ as predicted by the ETS. Indeed, the peaks of the degradation

rates show only < 1h time differences from the maximum $-A'(t)/A(t)$ values across most

(89–99%) of the simulated conditions of single to triple PTM cases [Fig. 3(d); Text S12 and

Table S6]. In addition, for each substrate profile, the simulated degradation rate tends to

become more rhythmic and have a larger relative amplitude as the number of the PTMs

increases [Fig. 3(e)], supporting the above argument that multiple PTMs can facilitate

degradation rhythmicity. The estimated relative amplitude in Eq. (13) also shows this

tendency for single to double PTMs, yet not clearly for triple PTMs unlike the simulated

relative amplitude [Fig. 3(e)]. This inaccuracy with the triple PTMs comes from the

accumulated errors over multiple PTMs in our estimation, as we indicated early. Still, the

estimate in Eq. (13) accounts for at least the order of magnitude of the simulated relative

amplitude, as the ratio of the simulated to estimated relative amplitude almost equals 1 for a

single PTM case and remains to be $O(1)$ for double and triple PTM cases [Fig. 3(f)].

Fig. 3. Rhythmic degradation of circadian proteins. (a) The experimental abundance levels (solid line)

and degradation rates (open circles) of the mouse PERIOD2 (PER2) protein [43]. (b) The experimental

abundance levels (dots, interpolated by a solid line) and degradation rates (open circles) of PSEUDO

RESPONSE REGULATOR 7 (PRR7) protein in *Arabidopsis thaliana* [44,45,59]. Horizontal white and black

segments correspond to light and dark intervals, respectively. (c) A simulated protein degradation rate

from the full kinetic model and its ETS- and QSSA-based estimates, when the degradation depends on

a single PTM. In addition, the protein abundance profile is presented here (gray solid line). A vertical

dashed line corresponds to the peak time of $-A'(t)/A(t)$ where $A(t)$ is a protein abundance. The

parameters are provided in Table S9. (d) The probability distribution of the peak-time difference

between a degradation rate and $-A'(t)/A(t)$ for each number of PTMs (n) required for the

degradation. The probability distribution was obtained with randomly-sampled parameter sets in

Table S6. (e) The probability distribution of the relative amplitude of a simulated degradation rate

(top) or its estimate in Eq. (13) (bottom) for each n, when the relative amplitude of a protein

abundance is 1. (f) The probability distribution of the ratio of the simulated to estimated relative

amplitude of a degradation rate for each n. For more details of (a)–(f), refer to Text S11.

464
465 Together, the ETS provides a useful theoretical framework of rhythmic degradation of

466 circadian proteins, which is hardly explained by the quasi-steady state assumption.

467
468 **Parameter estimation**

469
The use of an accurate function of variables and parameters is important for good parameter

estimation by the fitting of the parameters [13,48,49]. Parameter estimation is a crucial part

of pharmacokinetic–pharmacodynamic (PK–PD) analysis for drug development and clinical

study design [50,51]. Yet, the MM rate law is widely deployed for PK–PD models integrated

into popular simulation and statistical analysis tools. To raise a caution against the

unconditional use of the quasi-steady state assumption in parameter estimation, we here

compare the accuracies of the tQSSA- and ETS-based parameter estimates. Because the

sQSSA-based parameter estimates have already been known as less accurate than the

tQSSA-based ones [12,49] and our own analysis supports this claim (Fig. S8), we will

henceforth skip the use of the sQSSA.

Specifically, we consider a protein–protein interaction model with time-varying protein

concentrations (Text S12). To the "true" profile of the protein complex [i.e., $C(t)$ in Eq. (1)],

we fit the ETS [$C_\gamma(t)$ in Eq. (6)] or the tQSSA [$C_{tQ}(t)$ in Eq. (2)] and estimate the original

parameters of the model [52]: the ETS-based fitting can estimate both parameters K and k_δ,

and the tQSSA-based fitting can estimate only K. Likewise, we consider a TF–DNA interaction

model with time-varying TF concentration (Text S12). The ETS-based fitting can estimate

both K and k_δ, and the QSSA-based fitting can estimate only K.

In the case of protein–protein interactions, Fig. 4(a) reveals that the ETS tends to improve

the parameter estimation over the tQSSA, with more accurately estimated K: most of K

values (89.4%) estimated by the ETS show smaller relative errors than the tQSSA-based

estimates and their 69.3% even show relative errors less than half the tQSSA's. In the case of

TF–DNA interactions, the ETS still offers an improvement in the estimation of K [Fig. 4(b)]:

most of the ETS-estimated K values (90.3%) show smaller relative errors than the QSSA-

estimated ones and their 51.8% even show relative errors less than half the QSSA's.

Fig. 4. Parameter estimation for protein–protein and TF–DNA interaction models. (a) The scatter plot

of the relative errors of the tQSSA- and ETS-estimated K values for a protein–protein interaction

model. (b) The scatter plot of the relative errors of the QSSA- and ETS-estimated K values for a TF–

DNA interaction model. In (a) and (b), a diagonal line corresponds to the cases where the two

estimates have the same relative errors. (c) The probability distribution of the relative error of the

ETS-estimated k_δ for the protein–protein interaction model in (a). (d) The probability distribution of

the relative error of the ETS-estimated k_δ for the TF–DNA interaction model in (b). Regarding (a)–(d),

a subset of simulated conditions gave relative errors outside the presented ranges here, but they did

not alter the observed patterns. For more details, refer to Text S12.

Unlike K, k_δ can only be estimated through the ETS, and hence the comparison to the

tQSSA- or QSSA-based estimate is not possible. Still, k_δ is found to have the relative error <

0.1 for most of the ETS-based estimates, 81.3% and 81.0% in the cases of protein–protein

and TF–DNA interactions, respectively [Figs. 4(c) and 4(d)].

Discussion

The quasi-steady state assumption involves the approximation by time-scale separation where the "fast" components of a system undergo instantaneous equilibrium and only the "slow" components govern the relevant dynamics. The time-scale separation has been a long practice in many different areas, such as the Monod–Wyman–Changeux model of allosteric effects, the Ackers–Johnson–Shea model of gene regulation by λ phage repressor, and the Born–Oppenheimer approximation in quantum chemistry [53–55]. If some prediction from the time-scale separation deviates from empirical data, our study may provide a useful intuition about this deviation based on an overlooked time-delay effect in that system.

   We here proposed the ETS as a theoretical framework of molecular interaction kinetics with time-varying molecular concentrations. The utility of the ETS for transient or oscillatory dynamics originates in the rigorous estimation of the relaxation time in complex formation, i.e., the effective time delay. In the cases of protein–protein and TF–DNA interactions, the ETS manifests the importance of the effective time delay for the time-course molecular profiles distinct from the quasi-steady states. Accordingly, the ETS provides valuable analytical insights into the signal response time under autogenous regulation and the spontaneous establishment of the rhythmic degradation rates of circadian proteins. In addition, the ETS improves kinetic parameter estimation with a caution against the unconditional use of the quasi-steady state assumption. Our approach enhances the mathematical understanding of the time-varying behaviors of complex-complete mass-action models [38,42,56] beyond only their steady states.

   Further elaboration and physical interpretation of our framework, in concert with extensive experimental profiling of molecular complexes in regulatory or signaling pathways [18,19], are warranted for the correct explanation of the interplay of cellular components and its functional consequences. Although the simulation and empirical data presented here are supportive of the ETS, direct experimental validation is clearly warranted. This validation could involve the measurements of the time-series of molecular complex concentrations by co-immunoprecipitation assays or other techniques. High temporal resolution data are preferred for their comparison with the ETS-based profiles. Lastly, comprehensive consideration of stochastic fluctuations in molecular binding events [32,57,58] beyond the TF–DNA interactions in this study would be a fruitful endeavor for more complete development of our theory, through possible extension of the existing stochastic QSSA [33,34].

Acknowledgments

We thank Haneul Kim for useful discussions. This work was supported by Hong Kong Baptist University, Startup Grant Tier 2 (RC-SGT2/18-19/SCI/001) and Blue Sky Research Fund (RC-BSRF/21-22/09) (R.L., T.L.P.M., and P.-J.K.), the Health and Medical Research Fund (HMRF 17182691) (R.L. and P.-J.K.), and the National Research Foundation of Korea Grants (NRF-2020R1A4A1019140, NRF-2020R1F1A1075942, and NRF-2018K1A4A3A01063890) funded by the Ministry of Science and ICT (J.C., W.J.K., and C.-M.G). This work was partially conducted with the resources of the High Performance Cluster Computing Centre, Hong Kong Baptist University, which receives funding from Research Grant Council, University Grant Committee of the HKSAR and Hong Kong Baptist University. We also acknowledge the support of the UNIST Supercomputing Center for the computing resources.

References

1. V. Henri, *Lois Générales de l'Action des Diastases* (Librairie Scientifique A. Hermann, 1903).

2. L. Michaelis and M. L. Menten, *Die Kinetik der Invertinwirkung*, Biochem. Z. **49**, 333–369 (1913).

3. G. E. Briggs and J. B. S. Haldane, *A Note on the Kinetics of Enzyme Action*, Biochem. J. **19**, 338 (1925).

4. J. Gunawardena, *Time-Scale Separation–Michaelis and Menten's Old Idea, Still Bearing Fruit*, FEBS. J. **281**, 473–488 (2014).

5. C. Gérard and A. Goldbeter, *A Skeleton Model for the Network of Cyclin-Dependent Kinases Driving the Mammalian Cell Cycle*, Interface Focus **1**, 24–35 (2011).

6. K. C. Chen, A. Csikasz-Nagy, B. Gyorffy, J. Val, B. Novak, and J. J. Tyson, *Kinetic Analysis of a Molecular Model of the Budding Yeast Cell Cycle*, Mol. Biol. Cell **11**, 369–391 (2000).

7. J. C. Leloup and A. Goldbeter, *Toward a Detailed Computational Model for the Mammalian Circadian Clock*, Proc. Natl. Acad. Sci. USA **100**, 7051–7056 (2003).

8. J. Garcia-Ojalvo, *Modeling Gene Expression in Time and Space*, Annu. Rev. Biophys. **42**, 605–627 (2013).

9. M. Foo, D. E. Somers, and P.-J. Kim, *Kernel Architecture of the Genetic Circuitry of the Arabidopsis Circadian System*, PLOS Comput. Biol. **12**, e1004748 (2016).

10. T. D. Pollard, *A Guide to Simple and Informative Binding Assays*, Mol. Biol. Cell **21**, 4061–4067 (2010).

11. A. D. Attie and R. T. Raines, *Analysis of Receptor–Ligand Interactions*, J. Chem. Educ.

72, 119 (1995).

12. J. K. Kim and J. J. Tyson, *Misuse of the Michaelis–Menten Rate Law for Protein*

Interaction Networks and Its Remedy, PLOS Comput. Biol. **16**, e1008258 (2020).

13. S. Schnell and P. K. Maini, *A Century of Enzyme Kinetics: Reliability of the K_M and v_{max}*

Estimates, Comm. Theor. Biol. **8**, 169–187 (2003).

14. L. A. Segel and M. Slemrod, *The Quasi-Steady-State Assumption: a Case Study in*

Perturbation, SIAM Rev. **31**, 446–477 (1989).

15. S. Schnell, *Validity of the Michaelis–Menten Equation – Steady-State or Reactant*

Stationary Assumption: That is the Question, FEBS J. **281**, 464–472 (2014).

16. J. Eilertsen, M. R. Roussel, S. Schnell, and S. Walcher, *On the Quasi-Steady-State*

Approximation in an Open Michaelis–Menten Reaction Mechanism, AIMS Math. **6**,

6781–6814 (2021).

17. J. Eilertsen, S. Schnell, and S. Walcher, *On the Anti-Quasi-Steady-State Conditions of*

Enzyme Kinetics, Math. Biosci. **350**, 108870 (2022).

18. A. Fujioka, K. Terai, R. E. Itoh, K. Aoki, T. Nakamura, S. Kuroda *et al.*, *Dynamics of the*

Ras/ERK MAPK Cascade as Monitored by Fluorescent Probes, J. Biol. Chem. **281**, 8917–

8926 (2006).

19. N. Blüthgen, F. J. Bruggeman, S. Legewie, H. Herzel, H. V. Westerhoff, and B. N.

Kholodenko, *Effects of Sequestration on Signal Transduction Cascades*, FEBS J. **273**,

895–906 (2006).

20. S. Carmi, E. Y. Levanon, S. Havlin, and E. Eisenberg, *Connectivity and Expression in*

Protein Networks: Proteins in a Complex Are Uniformly Expressed, Phys. Rev. E. **73**,

031909 (2006).

21. J. K. Kim and D. B. Forger, *A Mechanism for Robust Circadian Timekeeping via*

Stoichiometric Balance, Mol. Syst. Biol. **8**, 630 (2012).

22. J. A. M. Borghans, R. J. de Boer, and L. A. Segel, *Extending the Quasi-Steady State*

Approximation by Changing Variables, Bull. Math. Biol. **58**, 43–63 (1996).

23. N. E. Buchler and M. Louis, *Molecular Titration and Ultrasensitivity in Regulatory*

Networks, J. Mol. Biol. **384**, 1106–1119 (2008).

24. A. R. Tzafriri, *Michaelis–Menten Kinetics at High Enzyme Concentrations*, Bull. Math.

Biol. **65**, 1111–1129 (2003).

25. H. C. Lim, *On Kinetic Behavior at High Enzyme Concentrations*, AICHE J. **19**, 659–661

(1973).

26. S. Cha, *Kinetic Behavior at High Enzyme Concentrations: Magnitude of Errors of*

Michaelis–Menten and Other Approximations, J. Biol. Chem. **245**, 4814–4818 (1970).

27. K. J. Laidler, *Theory of the Transient Phase in Kinetics, with Special Reference to*

Enzyme Systems, Can. J. Chem. **33**, 1614–1624 (1955).

28. F. Gachon, E. Nagoshi, S. Brown, J. Ripperger, and U. Schibler, *The Mammalian*

Circadian Timing System: From Gene Expression to Physiology, Chromosoma **113**, 103–

112 (2004).

29. D. H. Nagel and S. A. Kay, *Complexity in the Wiring and Regulation of Plant Circadian*

Networks, Curr. Biol*.* **22**, R648–R657 (2012).

30. T. S. Hatakeyama and K. Kaneko, *Reciprocity between Robustness of Period and*

Plasticity of Phase in Biological Clocks, Phys. Rev. Lett. **115**, 218101 (2015).

31. N. Mosheiff, B. M. C. Martins, S. Pearl-Mizrahi, A. Grünberger, S. Helfrich, I.

Mihalcescu, D. Kohlheyer, J. C. W. Locke, L. Glass, and N. Q. Balaban, *Inheritance of*

Cell-Cycle Duration in the Presence of Periodic Forcing, Phys. Rev. X **8**, 021035 (2018).

32. V. Kampen and N. Godfried, *Stochastic Processes in Physics and Chemistry* (Elsevier,

1992).

33. J. K. Kim and E. D. Sontag, *Reduction of Multiscale Stochastic Biochemical Reaction*

Networks Using Exact Moment Derivation, PLOS Comput. Biol. **13**, e1005571 (2017).

34. Y. M. Song, H. Hong, and J. K. Kim, *Universally Valid Reduction of Multiscale Stochastic*

Biochemical Systems Using Simple Non-Elementary Propensities, PLOS Comput. Biol. **17**,

e1008952 (2021).

35. E. Levine and T. Hwa, *Stochastic Fluctuations in Metabolic Pathways*, Proc. Natl. Acad.

Sci. USA **104**, 9224–9229 (2007).

36. Y. T. Maeda and M. Sano, *Regulatory Dynamics of Synthetic Gene Networks with Positive*

Feedback, J. Mol. Biol. **359**, 1107–1124 (2006).

37. N. Rosenfeld, M. B. Elowitz, and U. Alon, *Negative Autoregulation Speeds the Response*

Times of Transcription Networks, J. Mol. Biol. **323**, 785–793 (2002).

38. C. Jeynes-Smith and R. P. Araujo, *Ultrasensitivity and Bistability in Covalent-Modification*

Cycles with Positive Autoregulation, Proc. R. Soc. A **477**, 20210069 (2021).

39. F. Nazarimehr, S. Jafari, M. Perc, and J. C. Sprott, *Critical Slowing Down Indicators*,

Europhys. Lett. **132**, 18001 (2020).

40. M. I. Maturana, C. Meisel, K. Dell, P. J. Karoly, W. D'Souza, D. B. Grayden, A. N. Burkitt, P.

Jiruska, J. Kudlacek, J. Hlinka *et al.*, *Critical Slowing Down as a Biomarker for Seizure*

Susceptibility, Nat. Commun. **11**, 1–12 (2020).

41. M. Scheffer, *Foreseeing Tipping Points*, Nature **467**, 411–412 (2010).

42. R. Lim, J. Chae, D. E. Somers, C.-M. Ghim, and P.-J. Kim, *Cost-Effective Circadian*

Mechanism: Rhythmic Degradation of Circadian Proteins Spontaneously Emerges without

Rhythmic Post-Translational Regulation, iScience **24**, 102726 (2021).

43. M. Zhou, J. K. Kim, G. W. Eng, D. B. Forger, and D. M. Virshup, *A Period2 Phosphoswitch*

Regulates and Temperature Compensates Circadian Period, Mol. Cell **60**, 77–88 (2015).

44. E. M. Farré and S. A. Kay, *PRR7 Protein Levels Are Regulated by Light and the Circadian*

Clock in Arabidopsis, Plant J. **52**, 548–560 (2007).

45. H.-H. Jo, Y. J. Kim, J. K. Kim, M. Foo, D. E. Somers, and P.-J. Kim, *Waveforms of Molecular*

Oscillations Reveal Circadian Timekeeping Mechanisms, Commun. Biol. **1**, 207 (2018).

46. G. van Ooijen, L. E. Dixon, C. Troein, and A. J. Millar, *Proteasome Function Is Required for*

Biological Timing throughout the Twenty-Four Hour Cycle, Curr. Biol. **21**, 869–875 (2011).

47. J. Chae, R. Lim, C.-M. Ghim, and P.-J. Kim, *Backward Simulation for Inferring Hidden*

Biomolecular Kinetic Profiles, STAR Protoc. **2**, 100958 (2021).

48. W. Stroberg and S. Schnell, *On the Estimation Errors of K_M and V from Time-Course*

Experiments Using the Michaelis–Menten Equation, Biophys. Chem. **219**, 17 (2016).

49. B. Choi, G. A. Rempala, and J. K. Kim, *Beyond the Michaelis–Menten Equation: Accurate*

and Efficient Estimation of Enzyme Kinetic Parameters, Sci. Rep. **7**, 1–11 (2017).

50. H.-M. Back, H.-Y. Yun, S. K. Kim, and J. K. Kim, *Beyond the Michaelis–Menten: Accurate*

Prediction of In Vivo Hepatic Clearance for Drugs with Low K_Ms, Clin. Transl. Sci. **13**, 1199–

1207 (2020).

51. M. Holinstat and P. F. Bray, *Protease Receptor Antagonism to Target Blood Platelet*

Therapies, Clin. Pharmacol. Ther. **99**, 72–81 (2016).

52. M. J. D. Powell, *An Efficient Method for Finding the Minimum of a Function of Several*

Variables without Calculating Derivatives, Comput. J. **7**, 155–162 (1964).

53. J. Monod, J. Wyman, and J.-P. Changeux, *On the Nature of Allosteric Transitions: A*

Plausible Model, J. Mol. Biol. **12**, 88–118 (1965).

54. G. K. Ackers, A. D. Johnson, and M. A. Shea, *Quantitative Model for Gene Regulation by λ*

Phage Repressor, Proc. Natl. Acad. Sci. USA **79**, 1129–1133 (1982).

55. M. Born and R. Oppenheimer, *Zur Quantentheorie der Molekeln*, Ann. Phys. **389**, 457–484

(1927).

56. A. K. Manrai and J. Gunawardena, *The Geometry of Multisite Phosphorylation*, Biophys. J.

95, 5533–5543 (2008).

57. C.-M. Ghim and E. Almaas, *Two-Component Genetic Switch as a Synthetic Module with*

Tunable Stability, Phys. Rev. Lett. **103**, 028101 (2009).

58. P.-J. Kim and N. D. Price, *Macroscopic Kinetic Effect of Cell-to-Cell Variation in*

Biochemical Reactions, Phys. Rev. Lett. **104**, 148103 (2010).

59. N. Nakamichi *et al.*, *PSEUDO-RESPONSE REGULATORS 9, 7, and 5 Are Transcriptional*

Repressors in the Arabidopsis Circadian Clock, Plant Cell **22**, 594–605 (2010).