Homologous recombination between tandem paralogues drives evolution of Type VII

secretion system immunity genes in firmicute bacteria

Stephen R. Garrett[1]*, Giuseppina Mariano[1], and Tracy Palmer[1]*

[1] Microbes in Health and Disease Theme, Newcastle University Biosciences Institute,

Newcastle University, Newcastle upon Tyne, NE2 4HH, UK;

*To whom correspondence should be addressed.

e-mail: s.garrett2@newcastle.ac.uk, tracy.palmer@newcastle.ac.uk

Tel +44 191 208 3219

Running title: Expansion of T7 immunity gene families in firmicutes

ABSTRACT

The Type VII secretion system (T7SS) is found in many Gram-positive firmicutes and secretes

protein toxins that mediate bacterial antagonism. Two T7SS toxins have been identified in

Staphylococcus aureus, EsaD a nuclease toxin that is counteracted by the EsaG immunity

protein, and TspA, which has membrane depolarising activity and is neutralised by TsaI. Both

toxins are polymorphic, and strings of non-identical *esaG* and *tsaI* immunity genes are

encoded in all *S. aureus* strains. During genome sequence analysis of closely related *S.*

aureus strains we noted that there had been a deletion of six consecutive *esaG* copies in one

lineage. To investigate this further, we analysed the sequences of the tandem *esaG* genes

and their encoded proteins. We identified three blocks of high sequence homology shared by

all *esaG* genes, and identified evidence of extensive recombination events between *esaG*

paralogues facilitated through these conserved sequence blocks. Recombination between

these blocks accounts for loss of *esaG* genes from *S. aureus* genomes. TipC, an immunity

protein for the TelC lipid II phosphatase toxin secreted by the streptococcal T7SS, is also

encoded by   multiple gene paralogues. Two blocks of high sequence homology locate to the

5' and 3' end of *tipC* genes, and we found strong evidence for recombination between *tipC*

paralogues encoded by *Streptococcus mitis* BCC08. By contrast, we found only a single block

of homology across *tsaI* genes, and little evidence for intergenic recombination within this

gene family. We conclude that homologous recombination is one of the drivers for the

evolution of T7SS immunity gene clusters.

Key Words: T7SS, homologous recombination, immunity gene families, *Staphylococcus*

aureus, Streptococcus

INTRODUCTION

The type VII protein secretion system (T7SS) is found in many Gram-positive bacteria. Following its discovery in pathogenic Mycobacteria, it has since been described in a range of other actinobacteria and in firmicutes (1-5). Cryo-electron microscopy studies have shown that the ESX-5 T7SS from Mycobacteria exists as a 2.3 mDa membrane complex, with a central ATPase, EccC, forming a hexameric pore (6, 7). The firmicutes T7SS is only distantly related to the actinobacterial T7SSa system, and has been designated T7SSb. The hexameric ATPase is the only common component found across all T7SSs, and is designated EssC in the T7SSb systems (8, 9).

While the T7SSa is heavily linked with Mycobacterial virulence, there is growing evidence that the T7SSb plays an important role in bacterial antagonism (10, 11). In *Streptococcus intermedius*, three T7SSb-secreted antibacterial effectors have been identified, including TelB, an NADase, and TelC a lipid II phosphatase (12). The *Enterococcus faecalis* T7SSb also mediates contact-dependent inhibition of some firmicute bacteria, and a bioinformatic analysis in *Listeria monocytogenes* has identified over 40 potential antibacterial substrates of the T7SS (13, 14).

A role for the T7SSb in interbacterial competition was first characterised in the opportunistic pathogen *Staphylococcus aureus* (15). The T7SS-encoding locus in this organism is highly variable. Sequence divergence initiates towards the 3' end of *essC*, with *essC* sequences falling into one of four variants, termed *ess1 – ess4* (16). Downstream of each *essC* subtype is a cluster of variant-specific genes. In *essC1* variant strains one of these genes, *esaD*, encodes a secreted nuclease toxin with antibacterial activity. Protection from the toxic activity of EsaD is mediated by EsaG, which is encoded immediately adjacent to *esaD* in *essC1* strains. EsaG inactivates EsaD by forming a tight complex with the EsaD nuclease domain (15). While all *essC1* strains encode *esaD*, the toxin nuclease domain is polymorphic, and these strains also encode additional copies of *esaG* genes in a highly variable immunity gene

island located at the 3' end of the *ess*/*T7SS* locus (10, 15, 16). These additional *esaG* copies

are genetically diverse, but share some core regions of similarity within the encoded amino

acid sequences (17, 18). Strings of up to nine non-identical *esaG* genes are also found in a

similar genomic location in *essC2*, *essC3* and *essC4* strains (10).

TspA is a second antibacterial toxin secreted by the *S. aureus* T7SS. TspA has a C-terminal

membrane-depolarising domain, and immunity from intoxication is provided by the membrane-

bound TsaI protein (19). The *tspA-tsaI* locus is encoded distantly from the T7 gene cluster,

and is found across all four *essC* variant strains. Similar to EsaD, the TspA toxin domain is

polymorphic, and all strains encode clusters of TsaI variants (between two and sixteen copies)

directly downstream of *tspA* (19).

It is common for bacteria to encode repertoires of immunity proteins for protection against

polymorphic effector proteins. For example, bacteria in the gut accumulate immunity genes in

genomic islands that provide protection against type VI secretion system effectors (20). This

includes orphan immunity genes in strains that do not encode the cognate effector protein, as

we observe for *esaG* genes in *S. aureus essC2*, *essC3* and *essC4* strains that do not contain

esaD. While it is common for immunity islands to carry many predicted immunity genes, it is

less common to see so many homologues of the same immunity gene clustered together. At

present, little is known about the origin of the T7SS immunity repertoires.

The *Staphylococcus aureus* strain NCTC8325, and its derivatives, are commonly used as a

model strains for laboratory studies (21). NCTC8325 was initially isolated from a corneal ulcer

and has been used extensively for the propagation of the lysogenic bacteriophage

Staphylococcus virus 11 (Previously Phage 47) (22). Since NCTC8325 also carries two other

lysogens in its genome, *Staphylococcus* viruses 12 and 13 (23) UV-induced curing was used

to rid the strain of prophage, to give strain RN450 (24) (Fig 1a). RN450 has been used to

construct many laboratory strains over the years, including RN6390, which is one of the key

strains we have employed for T7SS secretion studies (e.g. (25, 26)).

During our analysis, we found that there were two separate genome sequences reported for

NCTC8325. When comparing the first sequence, published in 2006 (GenBank accession

number CP000253; (27)), with the later sequence (uploaded in 2018 by the Wellcome Trust

Sanger Institute to GenBank; accession number LS483365) we noted that there were a

number of differences, including in *esaG* repertoire encoded at the 3' end of the *ess* gene

cluster (Fig 1b). In this study we used whole genome sequence analysis to identify which of

these two genome sequences was most closely related to RN6390. We subsequently used

gene phylogeny and cluster analyses to identify the processes that drive the accumulation of

T7SS immunity genes. Our findings indicate that intergenic recombination is a major factor in

the evolution of *esaG* genes. We also noted that a similar process drives the evolution of *tipC*

genes, which encode Streptococcal TelC immunity proteins (12). By contrast, while there is

some evidence of intergenic recombination within *tsaI* clusters, it does not appear to be the

main mechanism of evolution for this gene family.

METHODS

Strains, genome sequencing and phylogenic analysis of prophage

Strain RN6390 was obtained from Professor Jan Maarten van Dijl (University of Groningen,

NL). All other strains were obtained from Professor José Penadés (Imperial College, UK). All

genome sequencing was carried out by MicrobesNG (Birmingham, UK) with enhanced

genome sequencing for RN6390 and standard whole genome service for RN25 and RN450;

sequences   are   available   at   NCBI   under   accession   numbers   CP090001.1,

JAJSOX000000000.1 and JAJSOY000000000.1, respectively. Whole genome alignment was

executed using progressiveMauve (28). SNP calling was carried out using Snippy v4.6.0 (29).

To determine the taxonomy of the phage identified in RN6390 strain, its nucleotide sequence

was submitted to VIPTree browser with default parameters (30). From the resulting proteomic

tree, phage genomes associated with *S. aureus* species were selected to generate the tree in

Fig 2.

Gene and protein alignment

Nucleotide sequences were obtained from NCBI and aligned with MAFFT v7.489 (31). Amino

acid sequences were obtained from NCBI and aligned using MUSCLE v3.8.1551 (32). To

construct   similarity   plots   for   both   nucleotide   and   amino   acid   sequences,   Plotcon

(https://www.bioinformatics.nl/cgi-bin/emboss/plotcon)   was   executed   using   aligned

sequences, with a window size of 5. For *esaG4,* the full pseudogene was used for the

nucleotide alignment. For the alignment of the EsaG4 amino acid sequence, the two predicted

open reading frames (ORFs) were used. For Plotcon analysis on large input sequences,

alignments were manually curated to remove partial sequences and pseudogenes.

Recombination Prediction

Aligned nucleotide sequences were opened in the RDP4 software (33) and Run All selected.

TrimAl v1.2 (34) was used to remove the unaligned 5' region of *tsaI* genes before RDP4

analysis was carried out.

Gene phylogeny

Maximum likelihood trees for nucleotide sequences were built with IQTREE v 2.1.4 (35), with

1000 ultrafast bootstraps. Trees were visualised and annotated using iTOL (36).

Comparison of genetic loci and gene cluster analysis

T7SS-encoding loci from each variant were subjected to pairwise comparisons with the

RN6390 *T7SS* locus using BLAST (37). A similar approach was used to compare

Staphylococcus phage RN6390 with *Staphylococcus* phage SAP40 genomes. The report

produced by BLAST was used as a comparison file in genoPlotR package (38) within RStudio

(39) to plot regions of similarities between the two loci. FlaGs (40) was used to assess the

variation in number of *tsaI* repeats across *S. aureus* strains. Examples were selected to

represent diversity at this locus.

Construction and analysis of a plasmid database

A database containing all available plasmid sequences was built from PLSDB (https://ccb-

microbe.cs.uni-saarland.de/plsdb/plasmids/) (41). Subsequently, Hmmsearch from the

HMMER suite (v 3.3.2) (42) was used to identify *esaG* or *tsaI* genes encoded within the

database. For each *esaG* or *tsaI* copy identified, efetch from the entrez-utilities (43) was used

to retrieve the specific assembly ID where the genes were encoded. EsaG and TsaI proteins

identified as encoded on plasmids together with their corresponding assembly ID, were then

analysed using FlaGs to define the specific gene neighbourhood of *esaG* and *tsaI* on these

assemblies.

RESULTS

Analysis of the RN6390 genome.

Two genome sequences are available for NCTC8325, here designated NCTC8325-Oklahoma

(GenBank accession number CP000253) and NCTC8325-Sanger (GenBank accession

number LS483365). The genomes of NCTC8325-Oklahoma and NCTC8325-Sanger were

aligned to identify differences between the strains (Table S1). Following SNP calling and

manual curation we identified a total of 78 SNPs and other small polymorphisms between

these two strains. We also identified a duplication of 16S and 23S ribosomal RNA genes in

NCTC8325-Oklahoma relative to NCTC8325-Sanger (Table S1). A further notable difference

is that NCTC8325-Sanger carries six additional *esaG* genes, relative to NCTC8325-

Oklahoma, at the *ess* cluster (Table S1, Fig 1b).

In order to identify which NCTC8325 strain is the likely precursor strain of RN6390, we carried

out whole genome sequencing of RN6390 (GenBank accession number CP090001.1) and

aligned it with the genomes of NCTC8325-Oklahoma and NCTC8325-Sanger, respectively.

Differences between RN6390 and each NCTC8325 strain are recorded in SNP tables (Tables

S2 and S3). As expected, a major difference found between both NCTC8325 genomes and

RN6390 is the presence of three prophages in the NCTC8325 genomes which were

successively cured out during the construction of the RN6390 progenitor, RN450 (Fig 1a). A

prophage, *Staphylococcus* phage 6390 (φ6390) has been identified previously in the genome

of RN6390, integrated in the *rpmF* gene (44, 45). This is at a different locus from the three

prophage loci found in the two NCTC8325 genome sequences. To determine whether φ6390

is related to any of the prophages cured from NCTC8325, the genome sequence was

extracted and used to determine its taxonomy. As shown in Fig 2, φ6390 is a distinct prophage

to those cured from NCTC8325. Analysis of φ6390 using blastn gave an almost identical

match to *Staphylococcus* phage SAP40 (GenBank accession number: MK801683.1, 99%

coverage and 99.98% identity). As φ6390 is not present in RN4220, this suggests that it was

introduced during genetic manipulation steps performed following generation of RN450.

Further analysis of the RN6390 genome sequence revealed that it aligned more closely to

NCTC8325-Sanger than NCTC8325-Oklahoma, and also carries the six additional *esaG*

genes found in NCTC8325-Sanger. This suggests that NCTC8325-Sanger is the likely

progenitor of the 8325 lineage. To confirm this, we sequenced the genomes of the

intermediate strains RN25 and RN450 (Genbank accession numbers JAJSOX000000000.1

and JAJSOY000000000.1, respectively), as they are directly on the lineage of RN6390. Both

of these strains share the additional copies of *esaG* and other genomic features of NCTC8325-

Sanger, confirming that this strain is likely to be the true parent of RN6390.

The origin of the NCTC8325-Oklahoma strain from GenBank accession CP000253 is

described as the 'University of Oklahoma Health Sciences Center'. This strain is clearly closely

related to NCTC8325-Sanger, but may potentially have accumulated mutations through serial

passaging within a laboratory setting. Strikingly, the biggest difference between the two strains

apart from the ribosomal RNA gene duplication is the loss of the six *esaG* homologues.

Intergenic recombination is a driving force in the evolution of *esaG* genes.

To determine how the six *esaG* copies may have been lost in NCTC8325-Oklahoma, we

examined this region of the chromosome in more detail. RN6390 encodes 12 homologues of

esaG, all of them at the *ess* gene cluster. The first copy is found directly downstream of *esaD*

and encodes the cognate immunity protein for the nuclease toxin (Fig 3a)(15). We have named

this *esaG1*. This is followed by three genes encoding hypothetical proteins (SAOUHSC_00270

_00271 and _00272), before a stretch of four further *esaG* homologues (*esaG2* - *esaG5*). This

is followed by a further homologue of *SAOUHSC_00270* (annotated as *SAOUHSC_00270b*

on Fig 3a, sharing 96.49% identity with *SAOUHSC_00270*) and then an additional seven *esaG*

homologues (*esaG6 – esaG12*; Fig 3a). Note that while *esaG4* is annotated as a pseudogene,

due to a short additional stretch of nucleotides close to the centre of the gene introducing a

premature stop and subsequent start codon, it actually encodes two smaller ORFs (Fig 3a). A

genome alignment using progressiveMauve predicts that the genomic deletion in NCTC8325-

Oklahoma spans *esaG2 – esaG7.*

To assess variation between the 12 homologues of EsaG, the amino acid sequences were

aligned (Fig 3b), and regions of homology between the proteins were analysed using Plotcon

(Fig 4a). Numerous regions of sequence homology were observed, including two major blocks

of high sequence homology at amino acids 13-31 and 84-119. To assess whether these were

general features of *S. aureus* EsaG proteins, we downloaded approximately 4,000 *S. aureus*

EsaG sequences from RefSeq and collectively analysed them using Plotcon. Fig S1a indicates

that a similar profile of homology is seen across all EsaG proteins.

We next examined the intergenic regions between the RN6390 *esaG* homologues. Strikingly,

we noted that most of the intergenic regions were of a very similar length, other than when

they directly preceded a non-*esaG* gene (for example *SAOUHSC_00270b*). They also share

a high degree of homology (Fig S2a). When we undertook Plotcon analysis on the *esaG*

genes, including the 3' intergenic regions (Fig 2a, Fig S2b), we noted the same two major

blocks of homology that we had seen from the amino acid analysis, but in addition a third block

encompassing the end of the gene and the downstream intergenic region (Fig 3a, Fig S2b,

Fig S3).

Given the substantial levels of homology between the *esaG* genes, we used the recombination

prediction software, RDP4, to determine whether there had been recombination events

between the genes. The RDP4 output, shown in Table S4 and summarised Fig 4b, predicts

with high significance that there have been extensive recombination events within most (but

not all) of these genes. As shown in Fig 4b, recombination appears to occur at the three points

within the genes that correspond to the regions of high nucleotide sequence homology (Fig

4b, Table S4). To support these findings, we constructed a maximum-likelihood tree for the

12 *esaG* genes (Fig 4c). No recombination is predicted within *esaG3*, which is genetically

distant from other *esaG* homologues. Conversely, *esaG2* and *esaG5* appear to vary only in

their central regions, and these cluster closely on the tree (Fig 4c), consistent with the RDP4

output.

Based on the RDP4 results, we built a schematic representation showing the homologous

regions of the *esaG* genes that are likely involved in the recombination events (Fig 4d).

Specific regions of the genes seem to share high homology with others, for example, the mid-

section of many of the homologues share high homology to equivalent sections of *esaG1*

(coloured teal). Conversely, genes such as *esaG3* and *esaG6* appear to be much more

diverse. Based on our observations, it is probable that a recombination event between the first

conserved regions of *esaG2* and *esaG8* was responsible for the loss of the seven genes at

this locus in NCTC8325-Oklahoma. This is corroborated by an alignment of the nucleotide

sequences of *esaG2* and *esaG8 with SAOUHSC_00274* (Fig S2c). The alignment indicates

that *SAOUHSC_00274* is mosaic comprising the 5' region of *esaG2* with the middle and 3'

sections of *esaG8*, consistent with recombination at homology block 1 (Fig S2d).

esaG recombination in an epidemic strain of _S. aureus_

USA300 is a methicillin-resistant *S. aureus essC1* strain, and a dominant cause of community-

acquired *S. aureus* infection in the USA (46). A recent study analysed the community spread

and evolution of a USA300 variant during a New York outbreak (47). Comparing the whole-

genome sequences of the epidemic lineage showed that these strains carry only seven *esaG*

genes in comparison to the ten copies in the closely related USA300 FPR3757. Using strain

BKV_2 as a representative of the outbreak lineage, we aligned the region spanning from *esaE*

to *SAUSA300_0303* with USA300 FPR3757. The alignment showed almost complete

sequence identity, other than in a region spanning from the middle of *SAUSA300_0295* to the

middle of *SAUSA300_0299* (Fig S4a) which was absent from BKV_2. When the nucleotide

sequences of *SAUSA300_0295* and *SAUSA300_0299* were aligned with *esaG4* of BKV_2,

esaG4 was again seen to be mosaic, with most of the gene being identical to

SAUSA300_0295 but the 3' end showing 100% identity to *SAUSA300_0299* (Fig S4b). This

is consistent with recombination between homology block 2 of *SAUSA300_0295* and

SAUSA300_0299, with loss of the intervening DNA.

esaG diversity across S. aureus essC2, essC3 and essC4 variants.

Although *S. aureus essC2*, *essC3* and *essC4* variants do not encode EsaD, these strains all

accumulate *esaG* genes at the 3' end of their T7SS-encoding loci. To ascertain whether the

homologues of *esaG* encoded in these strains are close relatives of those found in RN6390,

we analysed the *esaG* genes encoded in strains ST398, MRSA252 and HO 5096 0412, as

representatives of *essC2*, *essC3* and *essC4* variants, respectively. The sequence of the *T7SS*

loci for each of these strains was subjected to pairwise comparison, using BLAST, with the

RN6390 locus, and further analysed using genoPlotR (48) to produce a graphic representation

of the alignment of these regions (Fig 5). In this output, regions of homology are highlighted

by red-connecting blocks, and the colour intensity of these blocks reflects the percentage

identity found between the two compared regions.

As shown in Fig 5a, the *essC2* strain, ST398, shares higher homology with the RN6390 *T7SS*

cluster than either MRSA252 or HO 5096 0412. This strain harbours two intact copies of *esaG*

(*SAPIG_0310* and _0311), and one pseudogene (covering the two small genes *SAPIG_0314-*

0315, and herein referred to as a single *SAPIG_0314* pseudogene). We used RDP4 analysis

with *esaG1-12* from RN6390, and phylogenetic tree construction, to determine whether there

were any regions of shared homology. The analysis showed that *SAPIG_0314-0315* has the

highest homology to *esaG12* (Fig 5a) and appears to cluster in the same branch of the

phylogenetic tree (Fig 5d). No recombination events are predicted for this pseudogene,

suggesting this is most likely a copy of *esaG12* that has accrued mutations (Fig S5a).

Conversely, *SAPIG_0310* and *SAPIG_0311* have much lower homology to the *esaG*

homologues in RN6390 (Fig 5a), although recombination events are predicted with *esaG4* for

SAPIG_0310 and with *SAPIG_0314-315* for *SAPIG_0311* (Fig S5a).

The *essC3* strain, MRSA252, has four tandem *esaG* homologues at its *ess* locus, *SAR_0293*

- *SAR_0296* (Fig 5b). These are predicted by BLAST to share the highest homology with

RN6390 *esaG9-esaG12*. However, these homologues do not cluster together in the

phylogenetic tree (Fig 5d), and for example *SAR_0295*, predicted by genoPlotR to be a

homologue of *esaG11*, clusters with *esaG3*. Recombination events are also detected for all of

the *esaG* genes in MRSA252 (Fig S5b).

The *essC4* strain, HO 5096 0412, harbours two *esaG* copies, *SAEMRSA15_02570* and

SAEMRSA15_02580, which are predicted by BLAST to share highest homology with *esaG11*

and *esaG12*, respectively (Fig 5c). However, phylogenetic analysis indicates that

SAEMRSA15_02580 clusters with *esaG11* as opposed to *esaG12*, and *SAEMRSA15_02570*

does not cluster closely with any of the RN6390 *esaG* genes (Fig 5d). RDP4 analysis also

predicts recombination events for both genes in HO 5096 0412 (Fig S5c).

In summary, whilst *SAPIG_0314-0315* is a copy of *esaG12*, the remaining *esaG* homologues

found in the three representative strains of *essC* variants 2, 3 and 4 all appear to be

recombinants. Some of the recombination events are between homologues present in these

strains (Tables S5 - S7), suggestive of common ancestry. However, for other recombination

events, the parent *esaG* gene is unknown and is not present in any of these representative

strains.

Accumulation of *tsaI* genes in the RN6390 genome.

TspA is a T7SS-secreted antibacterial toxin that is highly conserved across all *essC* variant

strains (19). In all of these strains it is encoded away from the T7SS gene cluster, at a genomic

location bounded by *SAOUHSC_00583* and *iolS* (*SAOUHSC_00603*). The toxic activity of

TspA is neutralised by TsaI, a membrane protein of the DUF443 family (19). Multiple copies

of *tsaI* genes are encoded downstream of *tspA* (Fig 6, Fig S6), and in RN6390 there are 11

copies, *tsaI1 – tsaI11* (Fig 6a). A small pseudogene, encoded by *SAOUHSC_00600* shares

homology to part of the toxin region of TspA and is also found at this locus.

We wondered whether recombination events, similar to those we have observed for *esaG*

genes, also contributed to the evolution of *tsaI* genes. An alignment of the amino acid

sequences for RN6390 TsaI proteins (Fig 6b), shows that there is much greater sequence

variability between these proteins than the EsaG homologues. This is particularly apparent in

the C-terminal region of the protein, with only limited sequence identity observed between TsaI

homologues. Similar variability was observed in a representative alignment of around 3000

TsaI sequences (Fig S1b). Much greater variability was also observed in the *tsaI* intergenic

regions, in both length and DNA sequence compared with *esaG* (Fig 6c).

To identify the regions of highest homology at both the protein and DNA level, we ran Plotcon

analysis of the 11 *tsaI* genes and their encoded ORFs. Unlike EsaG*,* TsaI homologues have

only a single region of high similarly, covering approximately the first 75 amino acids, which is

also mirrored at the DNA level (Fig 7a, Fig S7). As only one block of high homology is detected

and there is a high degree of sequence variability in the *tsaI* intergenic regions, recombination

within individual genes is unlikely. To analyse this, we used RDP4 to predict recombination

events within the 11 homologues of *tsaI* genes (Fig 7b). Far fewer potential recombination

events were predicted than for *esaD* genes, and with lower probability, which could arise from

evolutionary processes other than recombination (Table S8).

tsaI genes are found on Staphylococcal plasmids

Horizontal gene transfer (HGT) is a major mechanism for the movement of genomic material

between bacteria (49). Plasmids are one of the key drivers of HGT and help to mediate the

spread of resistance genes among bacterial populations (e.g. 50, 51). To investigate whether

the *esaG* and/or *tsaI* toxin resistance genes may also be disseminated by plasmids, we

constructed a database of bacterial plasmids as described in the methods section and

interrogated this for the presence of *tsaI* and *esaG* genes. We identified a single

Staphylococcal plasmid carrying an *esaG* gene and five Staphylococcal plasmids that encode

one or more TsaI homologues (Fig 8). Two of the five plasmids encoding TsaI (pCAPBN21

and an unnamed plasmid from *Staphylococcus caprae* 26D) also encode a full-length TspA

along with the two WXG100-like proteins (DUF5344 and DUF3958 family proteins) that have

been proposed to serve as TspA-specific T7SS targeting factors (10). A further two plasmids

(pSB1-57-a and an unnamed plasmid from *Staphylococcus warneri* SWO) code for a fragment

of TspA alongside TsaI, with the SWO plasmid also encoding two further *tsaI* genes. An

unnamed plasmid from *Staphylococcus simulans* MR1 encodes an orphan *tsaI* with no

detectable *tspA* remnant. The SWO plasmid is particularly interesting as this carries further

T7-related genes including the *esaG* we identified along with a portion of *esaD*, and a fragment

of an HNH-nuclease gene along with a SM1/KNR4 protein encoding gene (a family implicated

as a nuclease immunity protein (52) and found in Staphylococcal T7SS immunity gene islands

(10)). For three of the plasmids, the immunity genes are close to recombinase genes, which

may provide a mechanism for their accumulation, and four of them carry nearby *IS* elements

that could facilitate their transfer to the chromosome without the need for homologous

recombination.

Intergenic recombination between *tipC* immunity genes in *Streptococcus*.

Three T7SS-secreted antibacterial toxins have been identified in *Streptococcus intermedius*.

TelA and TelB are both cytoplasmic-acting toxins neutralised by the TipA and TipB immunity

proteins, respectively (12). Our genome analysis indicates that strains generally encode only

a single *tipA* gene, while *tipB* is found in up to five copies. The third *S. intermedius* toxin is

TelC, a lipid II phosphatase. Protection from TelC toxicity is provided by TipC, a membrane-

bound immunity protein that faces the extracellular space (12, 53). Klein *et al*. reported that

the number of *tipC* genes encoded at the *telC* locus was highly variable between strains (53).

We used gene neighbourhood analysis across Streptococcal genomes to compare the

number of *tipC* genes present at the *telC* gene cluster, identifying 15 of them in a strain of

Streptococcus mitis BCC08 (Fig 9a).

Alignment of the *S. mitis* BCC08 TipC sequences and their encoding DNA (Fig 9b, Fig S8)

showed two regions of high sequence conservation close to the start and end, with a central

region of much higher sequence variability (Fig 10a). Using RDP4 to screen for recombination,

at least five recombination events were predicted between these genes (Fig 10b, Table S9),

in each case almost certainly through the blocks of high homology we identified. To analyse

this further we constructed a maximum likelihood tree to compare *tipC* homology. Genes

D8786_RS05910 and *D8786_RS0585* cluster closely in this tree, which corresponds to the

recombination event predicted between these two genes (Fig 10b, Table S9). Likewise

D8786_RS05920, which is predicted to be the major parent to *D8786_RS05865* (Table S9)

also clusters phylogenetically with this gene. We conclude that similar to *esaG*, intergenic

recombination drives the evolution of *tipC* immunity gene repertoires.

DISCUSSION

Through comparative genome analysis we noted that the *esaG* copy number is distinctly different in closely related strains of *S. aureus*, a finding that had also been previously described within the NCTC8235 lineage (17). To investigate how copy number variability may arise, we undertook sequence analysis of EsaG proteins and their encoding DNA, including their 3' flanking regions. We found three blocks of highly conserved nucleotide sequence, a large central one of approximately 100 nucleotides in length, and a 5' and 3' block both of around 55 nucleotides each. Homologous recombination occurs at regions of high homology within nucleotide sequences (Reviewed in 54). The minimum length requirement for efficient recombination in *S. aureus* is unclear, but stretches of 40-70 nucleotides have been reported for other bacteria (e.g. 55 - 57). Using RDP4 to predict recombination within *esaG* genes, we found strong evidence for recombination, corresponding to events within each of the three homology blocks we identified. Furthermore, our analysis revealed that the loss of six *esaG* genes in NCTC8325-Oklahoma arose from recombination across homology block 1, whereas three *esaG* genes have been lost in an outbreak strain of USA300 through recombination across homology block 2.

Previous work has reported that the *S. aureus* tandem-like lipoproteins, encoded on the νSaα island, also show extensive copy number variation across strains (58, 59). Similar analysis to that reported here showed that each *lpl* gene shares a stretch of approximately 130 nucleotides of high homology in its central region. Recombination was demonstrated to occur between the central conserved region of one gene and the same region of the neighbouring gene (57), thus spanning the 3' portion of gene 1, the intergenic region and the 5' region of gene 2.

To investigate whether intergenic recombination might represent a general mechanism for the evolution of T7SS immunity gene families, we examined the organisation of the *tipA*, *tipB* and *tipC* immunity genes in Streptococci. It has previously been noted that *tipC* copy number is

highly variable in Streptococcal genomes (53), and our analysis identified that up to 15 copies

of *tipC* could be present. Examination of recombination events within *tipC* revealed that

intergenic recombination is also a feature, and that it primarily occurs between homologous

blocks of sequence identity at the start and end of the genes. The structure of TipC reveals

that it has seven beta strands forming a concave face, with three alpha helices made up from

the N- and C-terminal regions of the protein (53, Fig 10d). Analysis of colicin DNase toxins

and their immunity proteins has shown that sequence divergence between related toxins and

immunities tends to concentrate at the binding interface (60). In agreement with this, site-

directed mutagenesis has strongly implicated the concave face as the region that binds to

TelC, and this region of TipC shows the highest level of sequence divergence (Fig 10d; 53).

We speculate that for EsaG the regions of high homology are not directly involved in toxin

binding, but may provide a structural framework on which amino acid substitutions in the

variable regions accumulate to alter the toxin binding specificity. Mechanisms to allow rapid

evolution of immunity proteins are likely to be essential to allow strains to rapidly acquire

resistance to novel toxin variants.

TspA is a second polymorphic toxin encoded by *S. aureus*. Protection from its membrane-

depolarising activity is provided by the immunity protein TsaI (19). TsaI is a polytopic

membrane protein predicted to have five transmembrane domains. As with *esaG*, multiple *tsaI*

genes are found in *S. aureus* genomes, however, our analysis has indicated that there is little

evidence of recombination between them. We found only a single block of high nucleotide

sequence homology at the 5' end of *tsaI* genes. This encodes approximately the first 75 amino

acids of TsaI, which would encompass the first two transmembrane domains. At present it is

not known whether TsaI neutralises TspA though direct interaction, or through the

sequestering of a membrane-bound partner protein with which TspA must interact to facilitate

its insertion or folding (or a combination of both of these). In this context, membrane

permeabilising peptide bacteriocins produced by some Gram-positive bacteria require a

membrane-bound receptor, such as the membrane components of the mannose

phosphotransferase system, for their activity. In the case of the *Lactococcus lactis* lactococcin

A bacteriocin, the immunity protein LciA acts through formation a complex with both the

receptor protein and the bacteriocin (61). By analogy it is possible that the first two

transmembrane domains of TsaI interact with a candidate receptor, constraining their

sequence, whereas the remainder of the immunity protein binds to the toxin and is therefore

under diversifying selection. At present it is unclear how *tsaI* gene clusters evolve, although

we did note that there was evidence for carriage of *tsaI* genes on Staphylococcal plasmids,

which may facilitate gene movement within and between strains. Further work would be

required to clarify the mechanisms that drive *tsaI* diversity.

ACKNOWLEDGEMENTS

We thank Prof Jan-Maarten van Dijl (University of Groningen, NL) for providing us with strain

RN6390, and Prof José Penadés (Imperial College, UK) for providing all of the other strains

in this study. We thank Prof Victor J. Torres and Dr Alejandro Pironti (NYU Langone, USA) for

providing the genome sequence of USA300 BKV_2. This work was supported by Wellcome

through Investigator Award 110183/A/15/Z to TP and Sir Henry Wellcome Postdoctoral

Fellowship 218622/Z/19/Z to GM. SG is funded by the Newcastle-Liverpool-Durham BBSRC

DTP2 Training Grant, project reference number BB/M011186/1.

FIGURE LEGENDS

Figure 1. Lineage of NCTC8325 daughter strains. a. A selection of daughter strains of the

NCTC8325 lineage and how they were generated. RN25 (also called 8325-3) was generated

from NCTC8325 following UV curing of prophages 11 and 12 (23). RN450 (also called 8325-

4) was generated from RN25 by a second round of UV exposure, to cure prophage 13 (23).

Methylnitronitrosoguanidine (MNNG)-mediated mutagenesis of RN450 was used to generate

RN4220, which is restriction deficient and capable of accepting foreign DNA (62). RN450 was

separately transduced with pRN3032, a Tn551 donor plasmid to generate RN1478 (63)

ISP479 is a cadmium-resistant revertant of RN1478 (64). RN6390 was generated from ISP479

by curing of pRN3032 (24). b. The genetic layout of the *ess* locus in NCTC8325- Oklahoma

and NCTC8325-Sanger. The dashed lines and bar represent the region that is missing from

NCTC8325-Oklahoma, as identified in this study.

Figure 2. Analysis of *Staphylococcal* phage 6390. A phylogenetic tree of φ6390 against a

database of *Staphylococcal* phages generated using VIPtree (29). The positions of φ6390 and

Staphylococcal phages 11, 12 and 13 (excised from the NCTC8325 parent strain during

construction of RN6390) on the tree are indicated with red stars.

Figure 3. Homologues of *esaG* encoded at the *ess* locus in RN6390. a. Illustration of the 3' of the RN6390 *ess* locus which encodes the T7SS nuclease toxin, EsaD, its cognate immunity protein, EsaG and 11 further homologues of EsaG (numbered *esaG2 – esaG12*). Genes *esaG2* to *esaG7* are indicated by a black bar, and are absent from NCTC8325-Oklahoma, while the dotted line indicates the actual region lost by recombination. Note that *esaG4* is shown in hatched shading because it is annotated as a pseudogene. However, it does encode two predicted ORFs, EsaG4i and EsaG4ii. b. Sequence alignment of EsaG homologues encoded by RN6390. The black boxes represent regions of high homology based in this alignment.

Figure 4. Recombination within the RN6390 *esaG* homologues. a. Regions of high homology across the RN6390 EsaG protein sequences (middle panel) and the corresponding nucleotide sequences (bottom panel). The positions of blocks of high homology are shown in grey shading and along with their relative positions along the gene sequence (top panel). The basepair positions that define the conserved regions are taken from the nucleotide sequences of *esaG1*. b. RDP4 was used to predict recombination events within the *esaG* homologues encoded by RN6390. The identity of each gene is given in black at the left, with regions of recombination labelled directly below, in the colour of the gene from which the recombinant section originated. c. A maximum likelihood tree was generated for RN6390 *esaG* homologues in IQTREE and visualised and annotated in iTOL. d. Illustration of regions of homology in the *esaG* homologues in RN6390. Black bars represent conserved regions of the gene and the variable regions have been assigned a colour and corresponding number. Homologous regions are coloured with the same colour. Numbers were assigned based on the first gene in the series that had the unique variable region. White hatched shading indicate a pseudogene.

Figure 5. Recombination events within the *esaG* genes encoded in representative *essC2, essC3* and *essC4* variant strains. a-c. The genes downstream of *essC* are different between each of the four *essC* variants. The regions spanning *essC* to the conserved gene *SAOUHSC_00279* were aligned between RN6390, and a. ST398 (*essC2* variant), b. MRSA252 (*essC3* variant) and c. HO 5096 0412 (*essC4* variant). Alignments were visualised using genoPlotR. d. A maximum likelihood tree constructed with IQTREE and annotated in iTOL for all *esaG* homologues found across the four representative *essC* variant strains RN6390, ST398, MRSA252 and HO 5096 0412.

Figure 6. Homologues of *tsaI* encoded at the *tspA* locus of RN6390. a. Genetic arrangement of *tsaI* genes in RN6390. *rclA* and *iolS* are a conserved gene found flanking the TspA locus in *S. aureus* strains, encoding a pyridine nucleotide-disulfide oxidoreductase and an aldo-keto reductase, respectively. b. An alignment of the encoded TsaI homologues. The black boxes represent regions of high homology based in this alignment. c. Alignment of the intergenic region downstream of each *tsaI* gene.

Figure 7. Assessing recombination events within *tsaI* homologues. a. A single region of high homology across the RN6390 TsaI protein sequences (middle panel) and the corresponding nucleotide sequences (bottom panel). The numbers which dictate the limits of the conserved region are taken form the nucleotide sequence of *tsaI1*. b. RDP4 was used to predict recombination events within the *tsaI* homologues encoded in RN6390. Each gene is labelled in black, with regions of recombination labelled directly below in the colour of the gene from which the recombinant section originated.

Figure 8. *tsaI* genes are carried on Staphylococcal plasmids. The *tsaI*-encoding regions of unnamed plasmids from *S. simulans* MR1 (accession NZ_CP015643), *S. caprae* 26D (accession NZ_CP031272), *S. warneri* SWO (accession NZ_CP033101), along with plasmids pCAPBN21 (accession NZ_CP042342) and pSB1-57-a (accession CP070965) are shown.

Figure 9. Homologues of *tipC* encoded at the *telC* locus of *Streptococcus mitis* BCC08.

a. Genetic arrangement of *tipC* genes in *S. mitis* BCC08. B. An alignment of the encoded TipC homologues. The blue boxes represent regions of high sequence homology, and the dashed line at the N-terminus of the aligned sequences indicates a probable lipoprotein signal peptide.

Figure 10. Recombination within the *S. mitis* BCC08 *tipC* homologues. a. Regions of high homology across the *S. mitis* BCC08 TipC protein sequences (middle panel) and the corresponding nucleotide sequences (bottom panel). The positions of blocks of high homology are shown in grey shading, with their relative positions along the gene sequence (top). The first 18 amino acids of TipC form a predicted lipoprotein signal sequence which is indicated by pale grey shading. The basepair positions that define the conserved regions are taken from the nucleotide sequences of *D8786_RS05940*. b. RDP4 was used to predict recombination events within the *tipC* homologues. The identity of each gene is given in black at the left, with regions of recombination labelled directly below, in the colour of the gene from which the recombinant section originated. c. A maximum likelihood tree was generated for *tipC* homologues in IQTREE and visualised and annotated in iTOL. d. The conserved (cyan) and variable (orange) regions of TipC were mapped to the crystal structure of *S. intermedius* TipC2 (pdb:6DHX; 53).

Figure S1. Similarity plots for representative EsaG and TsaI amino acid sequences. All available amino acid sequences for EsaG and TsaI were obtained from RefSeq. Sequences were aligned and a similarity plot produced using plotcon for a. EsaG and b. TsaI.

Figure S2. Homology in the intergenic regions downstream of *esaG* genes in RN6390. a. The intergenic regions found directly downstream of each *esaG* gene were aligned and visualised using boxshade. b. Plotcon analysis of RN6390 *esaG* genes and their 3' intergenic regions. Alignment of *esaG* nucleotide sequences including the downstream IGR for *esaG*

genes from RN6390. e*saG1, esaG5 and esaG12* were excluded due to having a longer

intergenic region (as seen in Fig S2a). c. Nucleotide sequence alignment

for *esaG2* (NCTC8325_00242) and *esaG8 (NCTC8325_00250)* from NCTC8325-Sanger

with *SAOUHSC_00274* from NCTC8325-Oklahoma. d. Schematic representation of the

mosaic nature of *SAOUHSC_00274*.

Figure S3. Alignment of the nucleotide sequences of *esD1-esaD12*. The blocks of high

sequence homology corresponding to Fig 4a are outlined in blue.

Figure S4. A recombination event in an epidemic lineage of USA300 results in loss of

part of an *esaG* cluster and generation of a novel *esaG* gene. a. The *esaD* locus of

USA300 FPR3757 and USA300 BKV_2. The dashed lines represent the region that is missing

from the epidemic strain, USA300 BKV_2, when compared to the USA300 FPR3757 type

strain. b. Nucleotide sequence alignment for *SAUSA300_0295* and *SAUSA300_0299* from

USA300 FPR3757 and *esaG4* from USA300 BKV_2. Coloured blocks indicated homology

between *BKV_2 esaG4* and the genes with which it is aligned.

Figure S5. Assessing recombination events in *esaG* homologues from a representative

***essC2, essC3* and *essC4* strain.** Alignments of *esaG* homologues from RN6390 with a.

ST398 (*essC2* variant), b. MRSA252 (*essC3* variant) and c. HO 5096 0412 (*essC4* variant)

were analysed using RDP4 to analyse recombination events. Each gene is labelled in black,

with regions of recombination labelled directly below in the colour of the gene from which the

recombinant section originated.

Figure S6. Representation of the variability in the numbers of *tsaI* homologues encoded

by *S. aureus* strains. Homologues of TsaI1 were obtained from RefSeq and genes

neighbouring the *tsaI* genes were identified - a selection of strains were used to visualise the

variability in number of *tsaI* genes encoded at this locus.

Figure S7. Alignment of the nucleotide sequences of *tsal1-tsal11* from RN6390. The block of high sequence homology corresponding to Fig 7a is outlined in blue.

Figure S8. Alignment of the nucleotide sequences of *tipC* homologues from *S. mitis* BCC08. The blocks of high sequence homology corresponding to Fig 9a are outlined in blue. The region encoding the predicted lipoprotein signal peptide is indicated with a dashed line.

Table S1. SNP table for NCTC_8325-Oklahoma compared to NCTC8325-Sanger. Large deletions and insertions are included below.

Table S2. SNP table for NCTC_8325-Oklahoma compared to RN6390.

Table S3. SNP table for NCTC_8325-Sanger compared to RN6390.

Table S4. RDP4 output for recombination within the *esaG* genes from RN6390. Predicted recombination events are highlighted in blue. Events that may have occurred due to an evolutionary process other than recombination are highlighted in yellow.

Table S5. RDP4 output for recombination within the *esaG* genes from RN6390 and ST398. Predicted recombination events are highlighted in blue.

Table S6. RDP4 output for recombination within the *esaG* genes from RN6390 and MRSA252. Predicted recombination events are highlighted in blue. Events that may have occurred due to an evolutionary process other than recombination are highlighted in yellow.

628 **Table S7. RDP4 output for recombination within the *esaG* genes from RN6390 and HO**

629 **5096 0412.** Predicted recombination events are highlighted in blue. Events that may have

630 occurred due to an evolutionary process other than recombination are highlighted in yellow.

632 **Table S8. RDP4 output for recombination within the *tsaI* genes from RN6390.** Events that

633 may have occurred due to an evolutionary process other than recombination are highlighted

634 in yellow.

636 **Table S9. RDP4 output for recombination within the *tipC* genes from *S. mitis* BCC08.**

637 Predicted recombination events are highlighted in blue. Events that may have occurred due

638 to an evolutionary process other than recombination are highlighted in yellow.

References

1. **Hsu T, Hingley-Wilson SM, Chen B, Chen M, Dai AZ *et al.*** The primary mechanism of attenuation of bacillus Calmette-Guerin is a loss of secreted lytic function required for invasion of lung interstitial tissue. *Proc Natl Acad Sci U S A* 2003;100:12420-12425.

2. **Pym AS, Brodin P, Majlessi L, Brosch R, Demangel C, *et al.*** Recombinant BCG exporting ESAT-6 confers enhanced protection against tuberculosis. Nat Med 2003;9:533-539.

3. **Stanley SA, Raghavan S, Hwang WW, Cox JS.** Acute infection and macrophage subversion by *Mycobacterium tuberculosis* require a specialized secretion system. *Proc Natl Acad Sci U S A* 2003;100:13001-13006.

4. **Akpe San Roman S, Facey PD, Fernandez-Martinez L, Rodriguez C, Vallin C *et al*.** A heterodimer of EsxA and EsxB is involved in sporulation and is secreted by a type VII secretion system in *Streptomyces coelicolor*. *Microbiology* 2010;156:1719-1729.

5. **Burts ML, Williams WA, DeBord K, Missiakas DM.** EsxA and EsxB are secreted by an ESAT-6-like system that is required for the pathogenesis of *Staphylococcus aureus* infections. *Proc Natl Acad Sci U S A* 2005;102:1169-1174.

6. **Bunduc CM, Fahrenkamp D, Wald J, Ummels R, Bitter W *et al*.** Structure and dynamics of a mycobacterial type VII secretion system. *Nature* 2021;593:445-448.

7. **Beckham KSH, Ritter C, Chojnowski G, Ziemianowicz DS, Mullapudi E *et al*.** Structure of the mycobacterial ESX-5 type VII secretion system pore complex. *Sci Adv* 2021;7:eabg9923.

8. **Abdallah AM, Gey van Pittius NC, Champion PA, Cox J, Luirink J, *et al.*** Type VII secretion--mycobacteria show the way. *Nat Rev Microbiol* 2007;5:883-891.

9. **Zoltner M, Ng WM, Money JJ, Fyfe PK, Kneuper H, *et al.*** EssC: domain structures inform on the elusive translocation channel in the Type VII secretion system. *Biochem J* 2016; 473:1941-1952.

10. **Bowman L, Palmer T.** The Type VII Secretion System of *Staphylococcus*. *Annu Rev Microbiol* 2021;75:471-494.

11. **Tran HR, Grebenc DW, Klein TA, Whitney JC.** Bacterial type VII secretion: An important

player in host-microbe and microbe-microbe interactions. *Mol Microbiol* 2021;115:478-489.

12. **Whitney JC, Peterson SB, Kim J, Pazos M, Verster AJ, *et al.*** A broadly distributed toxin

family mediates contact-dependent antagonism between gram-positive bacteria. *Elife*. 2017;

6:e26938.

13. **Chatterjee A, Willett JLE, Dunny GM, Duerkop BA.** Phage infection and sub-lethal

antibiotic exposure mediate *Enterococcus faecalis* type VII secretion system dependent

inhibition of bystander bacteria. *PLoS genet* 2021;17:e1009204.

14. **Bowran K, Palmer T.** Extreme genetic diversity in the type VII secretion system of Listeria

monocytogenes suggests a role in bacterial antagonism. *Microbiology* 2021;167: doi:

10.1099/mic.0.001034

15. **Cao Z, Casabona MG, Kneuper H, Chalmers JD, Palmer T.** The type VII secretion

system of *Staphylococcus aureus* secretes a nuclease toxin that targets competitor bacteria.

Nat Microbiol 2016;2:16183.

16. **Warne B, Harkins CP, Harris SR, Vatsiou A, Stanley-Wall N, *et al.*** The Ess/Type VII

secretion system of *Staphylococcus aureus* shows unexpected genetic diversity. *BMC*

Genomics 2016;17:222.

17. **Baek KT, Frees D, Renzoni A, Barras C, Rodriguez N *et al*.** Genetic variation in the

Staphylococcus aureus 8325 strain lineage revealed by whole-genome sequencing. PLoS

ONE. 2013;8:e77122.

18. **Ohr RJ, Anderson M, Shi M, Schneewind O, Missiakas D.** EssD, a Nuclease Effector

of the *Staphylococcus aureus* ESS Pathway. *J Bacteriol* 2016;199:e00528-16.

19. **Ulhuq FR, Gomes MC, Duggan GM, Guo M, Mendonca C, *et al.*** A membrane-

depolarizing toxin substrate of the *Staphylococcus aureus* Type VII secretion system mediates

intra-species competition. *Proc Natl Acad Sci U S A* 2020;117:20836-20847.

20. **Ross BD, Verster AJ, Radey MC, Schmidtke DT, Pope CE *et al*.** Human gut bacteria

contain acquired interbacterial defence systems. *Nature* 2019;575:224-228.

21. **Novick RP.** Genetic systems in staphylococci. *Methods Enzymol* 1991;204:587-636.

22. **Herbert S, Ziebandt AK, Ohlsen K, Schafer T, Hecker M *et al*.** Repair of global

regulators in *Staphylococcus aureus* 8325 and comparative analysis with other clinical

isolates. *Infect Immun* 2010;78:2877-2889.

23. **Novick R.** Properties of a cryptic high-frequency transducing phage in *Staphylococcus*

aureus. Virology 1967;33:155-166.

24. **Peng HL, Novick RP, Kreiswirth B, Kornblum J, Schlievert P.** Cloning,

characterization, and sequencing of an accessory gene regulator (*agr*) in *Staphylococcus*

aureus. *J Bacteriol* 1988;170:4365-4372.

25. **Kneuper H, Cao ZP, Twomey KB, Zoltner M, Jäger F, *et al.*** Heterogeneity in ess

transcriptional organization and variable contribution of the Ess/Type VII protein secretion

system to virulence across closely related *Staphyloccus aureus* strains. *Mol Microbiol*

2014;93:928-943.

26. **Casabona MG, Kneuper H, Alferes de Lima D, Harkins CP, Zoltner M *et al*.** Haem-iron

plays a key role in the regulation of the Ess/type VII secretion system of *Staphylococcus*

aureus RN6390. *Microbiology* 2017;163:1839-1850.

27. **Gillaspy AF, Worrell V, Orvis J, Roe BA, Dyer DW, Iandolo JJ.** *Staphylococcus aureus*

NCTC8325 genome. In: Fischetti V, Novick R, Ferretti J, Portnoy D, Rood J, editors. Gram-

positive pathogens. Washington, D.C.: ASM Press; 2006. p. 381-412.

28. **Darling AE, Mau B, Perna NT.** progressiveMauve: multiple genome alignment with gene

gain, loss and rearrangement. *PLoS ONE* 2010;5:e11147.

29. **Seemann T.** snippy: fast bacterial variant calling from NGS reads

https://github.com/tseemann/snippy. 2015.

30. **Nishimura Y, Yoshida T, Kuronishi M, Uehara H, Ogata H, Goto S.** ViPTree: the viral

proteomic tree server. *Bioinformatics* 2017;33:2379-2380.

31. **Katoh K, Misawa K, Kuma K, Miyata T.** MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 2002;30:3059-3066.
751

752 32. **Edgar RC.** MUSCLE: multiple sequence alignment with high accuracy and high
753 throughput. *Nucleic Acids Res* 2004;32:1792-1797.
754

755 33. **Martin DP, Murrell B, Golden M, Khoosal A, Muhire B.** RDP4: Detection and analysis
756 of recombination patterns in virus genomes. *Virus Evol* 2015;1:vev003.
757

758 34. **Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T.** trimAl: a tool for automated
759 alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 2009;25:1972-1973.
760

761 35. **Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD *et al*.** IQ-TREE 2:
762 New models and efficient methods for phylogenetic inference in the genomic era. Mol Biol
763 Evol 2020;37:1530-1534.
764

765 36. **Letunic I, Bork P.** Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
766 display and annotation. *Nucleic Acids Res* 2021;49:W293-W6.
767

768 37. **Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ.** Basic local alignment search
769 tool. *J Mol Biol* 1990;215:403-410.
770

771 38. **Guy L, Kultima JR, Andersson SG.** genoPlotR: comparative gene and genome
772 visualization in R. *Bioinformatics* 2010;26**:**2334-2335.
773

774 39. **RStudio Team.** RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL
775 http://www.rstudio.com/ 2020.
776

777 40. **Saha CK, Sanches Pires R, Brolin H, Delannoy M, Atkinson GC.** FlaGs and webFlaGs:
778 discovering novel biology through the analysis of gene neighbourhood conservation.
779 *Bioinformatics* 2021;37:1312-1314.
780

781 41. **Galata V, Fehlmann T, Backes C, Keller A.** PLSDB: a resource of complete bacterial
782 plasmids. *Nucleic Acids Res* 2019;47:D195-D202.
783

784 42. **Eddy SR.** Accelerated profile HMM searches. *PLoS Comput Biol* 2011;7:e1002195.
785

43. **Coordinators NR.** Database resources of the National Center for biotechnology
information. Nucleic Acids Res 2017;45:D12–17.

44. **Goerke C, Wirtz C, Fluckiger U, Wolz C.** Extensive phage dynamics in *Staphylococcus*
aureus contributes to adaptation to the human host during infection. *Mol Microbiol*
2006;61:1673-1685.

45. **Goerke C, Pantucek R, Holtfreter S, Schulte B, Zink M *et al*.** Diversity of prophages in
dominant *Staphylococcus aureus* clonal lineages. J Bacteriol 2009;191:3462-3468.

46. **Seybold U, Kourbatova EV, Johnson JG, Halvosa SJ, Wang YF *et al*.** Emergence of
community-associated methicillin-resistant *Staphylococcus aureus* USA300 genotype as a
major cause of health care-associated blood stream infections. *Clin Infect Dis* 2006;42:647-
656.

47. **Copin R, Sause WE, Fulmer Y, Balasubramanian D, Dyzenhaus S *et al*.** Sequential
evolution of virulence and resistance during clonal spread of community-acquired methicillin-
resistant Staphylococcus aureus. *Proc Natl Acad Sci U S A* 2019;116:1745-1754.

48. **Guy L, Kultima JR, Andersson SG.** genoPlotR: comparative gene and genome
visualization in R. *Bioinformatics* 2010;26:2334-2335.

49. **Arnold BJ, Huang IT, Hanage WP.** Horizontal gene transfer and adaptive evolution in
bacteria. *Nat Rev Microbiol* 2021 in press doi: 10.1038/s41579-021-00650-4.

50. **Stevenson C, Hall JP, Harrison E, Wood A, Brockhurst MA.** Gene mobility promotes
the spread of resistance in bacterial populations. *ISME J* 2017;11:1930-1932.

51. **Buckner MMC, Ciusa ML, Piddock LJV.** Strategies to combat antimicrobial resistance:
anti-plasmid and plasmid curing. *FEMS Microbiol Rev* 2018;42:781-804.

52. **Zhang D, de Souza RF, Anantharaman V, Iyer LM, Aravind L.** Polymorphic toxin
systems: Comprehensive characterization of trafficking modes, processing, mechanisms of
action, immunity and ecology using comparative genomics. *Biol Direct* 2012;7:18.
820

53. **Klein TA, Pazos M, Surette MG, Vollmer W, Whitney JC.** Molecular Basis for Immunity

Protein Recognition of a Type VII Secretion System Exported Antibacterial Toxin. *J Mol Biol*

2018;430:4344-4358.

54. **Michel B, Leach D.** Homologous Recombination-Enzymes and Pathways. *EcoSal Plus*

2012;5: doi: 10.1128/ecosalplus.7.2.7.

55. **Khasanov FK, Zvingila DJ, Zainullin AA, Prozorov AA, Bashkirov VI.** Homologous

recombination between plasmid and chromosomal DNA in *Bacillus subtilis* requires

approximately 70 bp of homology. *Mol Gen Genet* 1992;234:494-497.

56. **Shen P, Huang HV.** Homologous recombination in *Escherichia coli*: dependence on

substrate length and homology. *Genetics* 1986;112:441-457.

57. **Fujitani Y, Yamamoto K, Kobayashi I.** Dependence of frequency of homologous

recombination on the homology length. *Genetics* 1995;140:797-809.

58. **Tsuru T, Kawai M, Mizutani-Ui Y, Uchiyama I, Kobayashi I.** Evolution of paralogous

genes: Reconstruction of genome rearrangements through comparison of multiple genomes

within *Staphylococcus aureus*. *Mol Biol Evol* 2006;23:1269-1285.

59. **Tsuru T, Kobayashi I.** Multiple genome comparison within a bacterial species reveals a

unit of evolution spanning two adjacent genes in a tandem paralog cluster. *Mol Biol Evol*

2008;25:2457-2473.

60. **Ruhe ZC, Low DA, Hayes CS.** Polymorphic Toxins and Their Immunity Proteins:

Diversity, Evolution, and Mechanisms of Delivery. *Annu Rev Microbiol* 2020;74:497-520.

61. **Diep DB, Skaugen M, Salehian Z, Holo H, Nes IF.** Common mechanisms of target cell

recognition and immunity for class II bacteriocins. *Proc Natl Acad Sci U S A* 2007;104:2384-

2389.

62. **Kreiswirth BN, Lofdahl S, Betley MJ, O'Reilly M, Schlievert PM *et al.*** The toxic shock

syndrome exotoxin structural gene is not detectably transmitted by a prophage. *Nature*

1983;305:709-712.

63. **Novick RP.** Studies on plasmid replication. III. Isolation and characterization of replication-

defective mutants. *Mol Gen Genet* 1974;135:131-147.

64. **Pattee PA.** Distribution of Tn551 insertion sites responsible for auxotrophy on the

Staphylococcus aureus chromosome. *J Bacteriol* 1981;145:479-488.

a

b

a

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

a

b

Fig 7

Staphylococcus simulans MR1 unnamed plasmid

Staphylococcus caprae 26D unnamed plasmid

Staphylococcus warneri SWO unnamed plasmid

Staphylococcus capitis BN2 plasmid pCAPBN21

Staphylococcus sp. SB1-57 plasmid pSB1-57-a

■ Tsal	■ DUF4176 hypothetical protein	■ IS element
■ TspA	■ DUF3958 hypothetical porotein	■ Recombinase family protein
■ EsaG	■ DUF5344/SACOL2603-like T7SS effector	■ Replication initiator protein A
■ EsaD	■ ADP-ribosylating toxin	■ DUF536 hypothetical protein

■ Hypothetical protein	■ ParA family protein	■ Enoyl-ACP reductase, FabI
■ Hypothetical protein	■ Plasmid replication protein	■ Ketoacyl-ACP synthase III
■ SdpI family protein	■ HNH endonuclease	■ Sodium/proline symporter, PutP
■ Type I toxin-antitoxin system Fst family toxin	■ SMI1/KNR4 family protein	■ Replication protein

Fig 8

Fig 9

Fig 10

a

b

Sup Fig 1

Sup Fig 2

Sup Fig 3

Sup Fig 4

a

b

c

Sup Fig 5

Staphylococcus aureus

21305
O46
CA-347
21334
LGA251
LTT89
GHA13
NCTC6135
PS/BAC/169/17/W
M6K323
BSAC1226

Flavodoxin Family Protein
Pseudogene

Hypothetical Protein
Aldo/keto reductase

tspA
tsaI

Rrf2 Transcriptional Regulator
Hypothetical Protein

Sup Fig 6

Sup Fig 7

Sup Fig 8