Distinct sub-second dopamine signaling in dorsolateral striatum measured by a genetically-encoded fluorescent sensor

Armando G. Salinas[1,2,7,†*], Jeong O. Lee[1,†], Shana M. Augustin[1,8], Shiliang Zhang[3], Tommaso Patriarchi[4,5], Lin Tian[4], Marisela Morales[6], Yolanda Mateo[1], David M. Lovinger[1*]

1, Laboratory for Integrative Neuroscience, National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Rockville, MD, USA

2, Department of Bioengineering, George Mason University, Fairfax, VA, USA

3, Confocal and Electron Microscopy Core, National Institute on Drug Abuse, Baltimore, MD, USA

4, Department of Biochemistry and Molecular Medicine, University of California at Davis, Davis, CA, USA

5, Institute of Pharmacology and Toxicology, University of Zurich, Zurich, Switzerland

6, Neuronal Networks Section, Integrative Neuroscience Research Branch, National Institute on Drug Abuse, Baltimore, MD, USA

7, Current affiliation, Department of Pharmacology, Toxicology & Neuroscience, Louisiana State University Health Sciences Center – Shreveport, Shreveport, LA, USA

8, Current affiliation, Department of Pharmacology, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA

† equal contribution

* corresponding authors, lovindav@mail.nih.gov

armando.salinas@lsuhs.edu

Abstract

Dopamine produces neuromodulation throughout the basal ganglia, cortex and other brain regions, and is implicated in movement control, neural mechanisms of reward and actions of misused substances. The efferent projections of dopaminergic neurons with somata in the substantia nigra pars compacta and ventral tegmental area strongly innervate different striatal subregions. While much is known about the function of these neurons, there is a relative deficiency of information about *in vivo* dopamine dynamics in the nigrostriatal projections, especially those to the dorsolateral striatum (DLS). In past studies, subsecond dopamine changes were measured predominantly with fast-scan cyclic voltammetry (FSCV) both in brain slices and *in vivo*. However, traditional FSCV has limitations in discriminating among catecholamines, and cannot be used for simultaneous measurement of both slow and fast/phasic dopamine changes. In addition, FSCV has been most useful for measuring dopamine in the ventral striatum *in vivo* with less utility for measurement in dorsolateral striatum. The development of genetically encoded dopamine sensors has provided a new approach to measuring slow and fast dopamine dynamics both in brain slices and *in vivo*, raising the hope of more facile measurement of *in vivo* dopamine measurements, including in areas where measurement was previously difficult with FSCV. To this end, we first evaluated dLight photometry in brain slices with simultaneous FSCV. We found that both techniques yielded comparable findings. However, differences were noted in responses to dopamine transporter inhibitors, including cocaine. We then used *in vivo* fiber photometry with dLight to examine responses to cocaine in DLS and compared responses during Pavlovian conditioning in DLS to two other striatal subregions. These experiments show that dopamine increases are readily detectable in DLS and provide new information about dopamine transient kinetics and slowly developing signaling during conditioning. Overall, our findings indicate that dLight photometry is well suited to measuring dopamine dynamics in a striatal region of great interest where such measurements were difficult previously.

Introduction

Dopamine is a catecholamine neurotransmitter found throughout the mammalian nigrostriatal and cortical-mesolimbic circuit with critical roles in many psychiatric and neurological disorders including Parkinson's Disease, psychosis, schizophrenia, and addiction (e.g. Nestler and Carlezon, 2006; Calabresi et al., 2009; Grace, 2016; Berke, 2018; Ott and Nieder, 2019). Midbrain dopaminergic neurons provide extensively branching axons to the striatum. The dopamine released from these afferents produces neuromodulation with key roles in neuronal function, synaptic plasticity and behavior. The rodent striatum has several subregions that are part of associative (dorsomedial striatum, DMS), limbic (nucleus accumbens/ventral striatum, NAc) and sensorimotor (dorsolateral striatum, DLS) circuits that have different roles in learning, movement control and reward (Yin and Knowlton, 2004). Thus, understanding the dynamics of dopamine release in different striatal subregions is important for a full understanding of the function of these striatal regions in the context of different brain circuits.

Dopamine release has been measured in isolated cells, brain slices and *in vivo* with techniques including microdialysis sampling coupled with electrochemical detection and variants of fast-scan cyclic voltammetry (FSCV). Microdialysis allows for precise chemical identification and concurrent measurements of multiple neurotransmitters from each sample (Chefer et al., 2009). Variants of microdialysis can also be used to estimate the absolute dopamine concentration in a brain region. However, this technique has a relatively slow sampling time, typically 5-20 minutes per sample that does not allow for precise correlation of dopamine levels with discrete behavioral events. Fast-scan cyclic voltammetry is an electrochemical method that can be used to measure phasic release of catecholamines at sampling rates of 10Hz or higher (Venton and Cao, 2020). Thus, FSCV allows for real-time measurement of dopamine release in response to stimuli in brain slices, as well as *in vivo* detection of sub-second changes in neurotransmitter release in relation to behavioral events. Indeed, real-time dopamine measurements with FSCV revolutionized the study of fast dopamine changes and motivated behaviors for well over a decade. However, FSCV has limitations. Conventional FSCV discerns fast signals from a recent baseline or pre-stimulus period, and thus this technique cannot measure absolute dopamine concentrations and is not well suited to measurement of slow changes in dopamine levels. In addition, FSCV cannot distinguish between different catecholamine neurotransmitters. In practice, FSCV measurements of *in vivo* dopamine have been made mainly in the rat Nucleus accumbens, although a few studies have examined the rat DMS and DLS (Brown et al., 2011; Howe et al., 2013; Klanker et al., 2017; Willuhn et al., 2012). Unfortunately,

in vivo measurements in mouse dorsal striatum, and the DLS in particular, are lacking. This is

presumably due to difficulties in detecting small changes accompanied by contaminating signals.

Genetically encoded fluorescent biosensors for dopamine were recently developed

(Patriarchi et al., 2018; Sun et al., 2018). These membrane-targeted, G-protein-coupled receptor

(GPCR)-based sensors employ modified dopamine receptors inactive for intracellular signaling

that have had a circular permuted Green Fluorescence Protein (cpGFP) molecule in place of the

third intracellular loop. The sensors work by coupling the dopamine induced conformational

changes to changes in cpGFP emission intensity. Although fluorescent dopamine sensors cannot

measure absolute concentrations of dopamine, they can be used to simultaneously detect slow

and fast (or phasic) changes in dopamine relative to baseline levels (Mohebi et al., 2019 ;

Jørgensen et al. 2023; Van Zessen et al. 2021). These sensors also have the sensitivity to

measure dopamine in brain regions where such measurements were previously difficult (e.g.

cerebral cortex Patriarchi et al., 2018). Using genetically encoded fluorescent dopamine sensors

allows for the real-time measurement of dopamine dynamics in relation to distinct behaviors in

areas such as DLS.

In the current study we first evaluated the ability of the genetically encoded dopamine

sensor dLight to detect stimulation-induced dopamine in dorsal striatum using brain slice

photometry with simultaneous FSCV. We found many similarities in indices of regulation of

dopamine release using both methods. We also noted differences in cocaine-induced effects on

dorsal striatal dopamine release measured with dLight photometry and FSCV that challenge the

tacitly accepted, but controversial, notion that cocaine enhances dopamine release *in vitro*. Next,

we examined striatal dopamine dynamics using *in vivo* fiber photometry with dLight to assess

slow and fast/phasic dopamine changes in DLS in comparison to other striatal subregions. In

Pavlovian conditioning driven by a natural food reinforcer, we also found distinct characteristics

of phasic dopamine dynamics in DLS compared to other striatal subregions. These findings

provide new information about dopamine dynamics in dorsal striatum, and should promote further

physiological and behavioral studies in dopamine dynamics throughout the striatum.

Results

<u>Simultaneous dLight and FSCV dopamine measurement in DLS slices</u>

We first wanted to examine the characteristics of dLight under conditions where dopamine is

readily detected even in DLS. Thus, we performed simultaneous recordings using a PMT based

photometer and traditional FSCV in acutely prepared brain slices (**Figure 1a**). Electrical
stimulation evoked fluorescence transients were readily observed in DLS and were blocked by
D1 dopamine receptor antagonism (**Figure 1b and Supplementary Figure 1**). Simultaneous
dLight photometry and FSCV recordings in response to different stimulus intensities revealed
similar input—output curves, with both methods showing significant responses to 100µA stimuli
(**Figure 1c**, one sample t-test at 100µA, photometry t=5.128 p=0.0022, FSCV t=2.512, p=0.0458,
df=6).

<u>dLight fluorescence signals and FSCV originate from dopamine neurons and are dynamic</u>

To ensure that the electrically evoked dLight fluorescence transients were attributable to
dopamine release from midbrain neurons, we used a viral strategy to genetically ablate substantia
nigra dopamine neurons. We infused a Cre-dependent Caspase3-encoding virus into the
substantia nigra and dLight virus into the dorsal striatum of DAT Cre mice (**Figure 1d**). At least
five weeks later, brain slices were prepared and simultaneous dLight and FSCV recordings were
conducted. We found that genetic ablation of nigral DA neurons resulted in significantly reduced
dopamine release in DAT Cre+ relative to DAT Cre- mice, as measured with dLight (*, p<0.05,
$F_{(1,14)}=5.257$) and FSCV (*, p<0.05, $F_{(1,7)}=7.63$) across several stimulation intensities **Figure**
1e & f). Also, the electrically evoked fluorescence signal was not detected in eGFP control
experiments (**Supplementary Figure 2**)

We next assessed whether dLight and FSCV signals would show similar dynamics during
manipulations that decrease or increase dopamine release. Thus, we applied the D2 dopamine
receptor agonist, quinpirole, to slices to inhibit dopamine release (via activation of presynaptic D2
dopamine autoreceptors). We found that 30 nM quinpirole inhibited dopamine release equally
with both methods (**Figure 1h**; t=2.032; p=0.076, df=8). Having shown that dLight signals could
be decreased, we assessed whether they could be increased by increasing extracellular calcium
levels, which would enhance dopamine release. Indeed, increasing extracellular calcium
increased dLight fluorescent responses (**Figure 1i & j**; *, p<0.05, t=2.611, df=12) demonstrating
that dLight signals are truly dynamic. Further, these results demonstrate comparable performance
between dLight photometry and FSCV measurements in terms of sensitivity to presynaptic
inhibition and excitability induced changed in dopamine release.

<u>Differential regulation of dLight photometry and FSCV dopamine signals by DAT inhibitors</u>

There have been many reports of dopamine transporter (DAT) inhibitor-mediated increases in DA
release published using FSCV. However, in Patriarchi et al. (2018), it appeared that cocaine did
not increase the dLight dopamine transient peak amplitude while it increased transient duration.
This intriguing observation directly conflicted with a large body of literature and prompted us to
directly compare cocaine effects using dLight photometry and FSCV dopamine measurements.
We began by comparing the effect of increasing cocaine concentrations on simultaneously
collected dLight photometry and FSCV measurements (**Figure 2a**). We found that cocaine
increased the peak amplitude of transients measured with FSCV, but not those measured with
dLight photometry (**Figure 2b,c**). Cocaine increased the duration of both dLight photometry- and
FSCV-measured transients, suggesting that cocaine was in fact inhibiting DAT. Notably, cocaine
concentrations greater than ~10µM led to a decrease in peak transient amplitude measured with
both techniques. This is consistent with previous findings using FSCV and is due to off target
cocaine inhibition of nAChRs, which contribute significantly to striatal dopamine release. This off
target cocaine inhibition of nAChRs was first shown electrophysiologically (Francis et al., 2000;
Ma et al., 2020) and then with striatal FSCV (Acevedo-Rodriguez et al., 2014; Chen et al., 2019).

We next determined whether a more specific DAT inhibitor, nomifensine (1 µM), would have
effects similar to cocaine. Like cocaine, nomifensine increased the peak transient amplitude
measured with FSCV **(Figure 2e,h)**, but not with dLight photometry. Application of nomifensine
increased the transient decay time measured with both techniques (**Figure 2d,f,h**).

Altering FSCV adsorption altered DAT inhibitor effects on dopamine release

To further probe the factors underlying the DAT inhibitor-induced increase in peak transient
amplitude measured with FSCV, we altered the triangle voltage wave form to applied to the carbon
fiber electrode (CFE) to peak at +1.0V instead of +1.2V. (**Figure 2g,i**). This waveform change will
decrease the dopamine aborption profile and effectively decrease the sensitivity of the FSCV CFE
(Heien et al., 2003; Venton and Cao, 2020). As expected, nomifensine (1 µM) effects measured
with dLight photometry were not affected by the waveform modification (**Figure 2f,g**). Note that
the normalized amplitude of the fluorescence intensity was around 1, indicating that the dLight
photometry measurement reports no increased dopamine release in response to the DAT
inhibitor.

When we assessed evoked dopamine release before and after application of nomifensine (1 µM)
using the modified triangle waveform for FSCV measurements, we found that DAT inhibition
increased dopamine transient duration **(Figure 2h)** consistent with the dLight photometry

measurement. However, no change in peak transient amplitude was detected during nomifensine

application when the altered triangle waveform (peak at +1.0V) was used for FSCV

measurements **(Figure 2i)**. Thus, the apparent increase in dopamine release measured with

FSCV cannot be detected if CFE absorption is decreased.

<u>DAT inhibitors do not affect dopamine release in DAT KO mice</u>

FSCV has been used extensively to study the effects of cocaine and other uptake blockers on the

dynamics of dopamine clearance and evoked release. Indeed, to explain how cocaine might lead

to increased dopamine release (as measured with FSCV), it was posited that cocaine acted via a

DAT-independent mechanism (Venton et al., 2006). To evaluate this hypothesis, we again

performed simultaneous dLight photometry and FSCV recordings during which cocaine was

applied to dorsal striatum brain slices from DAT KO mouse expressing dLight (**Figure 2j**). In DAT

KO mice, the duration of stimulation-induced dopamine transients is prolonged compared to WT

mice with a decay in the 10s of seconds, compared to 1-2s in WT mice (Jones et al., 1998;

Budygin et al., 2002; Mateo et al., 2004). Accordingly, in our experiments, both dLight photometry

and FSCV evoked dopamine transients were prolonged relative to WT mice (**Figure 2j,k**).

Cocaine did not affect either dopamine transient amplitude or decay time measured with FSCV

or dLight photometry. If the putative cocaine-induced increases in dopamine release assessed

with FSCV are due to DAT-independent mechanisms, then the increase in FSCV dopamine

transient peak height should be present in DAT KO mice. Since this was not the case, we

considered other potential mechanisms for the putative DAT inhibitor-induced increases in

dopamine release observed with FSCV.

<u>DAT inhibitors do not increase dopamine release: a new interpretation of results to reconcile</u>

<u>differences with different methods</u>

To reconcile these results, we posited a new interpretation for the effects of DAT inhibitors on

dopamine transients measured with FSCV (**Figure 2l**). The detection of dopamine with FSCV

relies on the electrochemical oxidation of dopamine at the surface of the CFE. Upon electrical

stimulation of the slice, dopamine is released and diffuses across a given volume. The extent of

this diffusion is determined by several factors but most notably DAT (Venton et al., 2003; Cragg

and Rice, 2004). Thus, in a given space, the sampling volume obtained with FSCV is limited to

the distance that dopamine can diffuse to the CFE surface. Under conditions of impaired DAT

function, however, the diffusion distance of dopamine increases (**Figure 2m**). This increased

release point diffusion of dopamine also increases the effective sampling volume of the CFE,

resulting in a larger signal, but not increased release per se. When the dopamine
absorption/sensitivity of the CFE is reduced (e.g. with the altered triangle waveform, Figure 2g,i)
the effect of the increased sampling volume is reduced and thus the enhanced overflow does not
alter peak transient amplitude.

<u>dLight sensor traffics to synaptic and extrasynaptic sites</u>

We next sought to determine if dLight could traffic to synaptic sites. This would mark a major
advance in the measurement of dopamine as both microdialysis and FSCV can only measure
non-synaptic dopamine overflow. We performed immunohistochemistry for tyrosine hydroxylase
(TH), the rate limiting step in catecholamine synthesis and GFP to label dLight in mouse dorsal
striatum. We found putative dopamine axons (i.e. TH+ axons) juxtaposed to dLight/GFP (**Figure
3a**), suggesting the possibility of dLight expression in close proximity to dopamine release sites.
We followed up these experiments with electron microscopy (EM) experiments to directly assess
whether dLight would traffic to synapses and thus, at least partially, report synaptic dopamine
release (**Supplementary Table S1**). We found that dLight trafficked exclusively to plasma
membrane (**Figure 3b**) or membrane-associated regions within the cell (**Figure 3c**). We also
found that dLight could indeed traffic to synaptic sites, including putative dopamine synapses
(**Figure 3d**), and was also present at nearby extrasynaptic sites ranging from a few μm to 10s of
μm from synapses.

<u>dLight measures both slow and fast/phasic dopamine changes in DLS with *in vivo*</u>
<u>pharmacological manipulations</u>

To determine the utility of dLight for measuring dopamine dynamics in DLS *in vivo*, we expressed
the sensor in this striatal subregion and measured the fluorescence intensity profile using a
custom-built *in vivo* fiber photometry system based on TCSPC principles (**Figure 4a**). The *in vivo*
fiber photometry measurement is similar to that used in our previous work (Cui et al., 2013; Cui
et al., 2014; Kupferschmidt et al., 2017). We measured dLight fluorescence from the DLS of mice
treated with cocaine (15 mg/kg i.p.) after a 10 min observation period in their home cage. We
found that cocaine produced an increase in fluorescence that plateaued after 20 min (**Figures 4b
& c**). We also observed transient (subsecond to second duration) increases in fluorescence both
before and after cocaine administration (**Figure 4d**). To examine the profiles of these
spontaneous transients, we time-locked the transients to their peaks (**Figure 4e & f**). We then
analyzed cocaine-induced changes in transient frequency, amplitude, and decay (**Figure 4g-j**).

We found that cocaine increased dopamine transient frequency (t=4.887, p<0.001, df=16) and

amplitude (t=13.81, p<0.001, df=16). Furthermore, we found that cocaine increased the decay

time constant of spontaneous dopamine transients (t=4.730 p<0.001, df=79).

In vivo changes in fluorescence originate from dLight

To determine if the fluorescence changes observed during our recordings originated from dLight

and to ensure that the observed changes in fluorescence were due to dopamine actions on dLight

(and not altered fluorescence readout due to changes in blood flow, for example), we further

analyzed the fluorescence signals in eGFP-expressing mice (**Supplementary Figure 3**) in freely

moving conditions and after cocaine injection. In these recordings we did not observe

spontaneous, fast transients either before or after cocaine injection. There was also no slow

increase in eGFP fluorescence after cocaine injection. We also validated, through benchtop

testing, that we do not observe transient changes in fluorescence due to fiber bending or

movement artifact when recording with the TCSPC fiber photometry system (**Supplementary**

Figure 3). We also performed recordings in which dLight-expressing mice were treated with the

D1 dopamine receptor antagonist SCH23390 to block dopamine binding to the sensor. Indeed,

treatment with SCH23390 eliminated phasic dopamine transients and this did not change with

application of cocaine plus a D2 receptor antagonist (**Figure 4k**). Further, in the presence of

SCH23390, the fluorescence profile for dLight resembled that of a static eGFP control mouse

(**Figure 4l**).

dLight measures distinctive phasic dopamine dynamics in DLS in comparison to other striatal

subregions during Pavlovian conditioning

Next, we characterized the phasic DA profiles in the DLS using *in vivo* dLight photometry, and

assessed regional differences in dopamine transients during Pavlovian conditioning as this

training paradigm produces clear behavioral changes associated with consistent and reliable

phasic DA release in NAc. We trained animals in a well-established conditioning paradigm for a

food reinforcer using a tone as the conditioned stimulus (**Figure 5a-d**). During 14 sessions of daily

training, the latency between reward delivery and head-entry decreased and stabilized (**Figure**

5b and c; two-tailed paired t-test, p<0.0001). **Figure 5e** shows example traces of phasic

dopamine transients in the *in vivo* dLight photometry measurement across striatal subregions.

The average dopamine transients evoked by the CS+ were compared across the striatal
subregions (**Figure 5f-h**). We found that the average decay time of individual transients in DLS
was significantly faster than in NAc (**Figure 5f-h and i**; one-way ANOVA, $F_{(2,27)}=14.93$,
p<0.0001).

A training-dependent temporal shift in the dopamine increases is shown in **Figure 5j,k and l** (as
well as Supplementary Figure 4 showing a color plot of increases on all training trials). Similar to
previous work (Brown et al., 2011; Willuhn et al., 2012; Patriarchi et al., 2018), we observed
dopamine transients in NAc with a short latency after reinforcer delivery that gradually shifted to
a short latency after the presentation of the CS+ during the conditioning (**Figure 5j,m,n**). The
response in NAc associated with reward delivery decreased over the course or training.
Interestingly, in the DMS and DLS, the transient response to the reward delivery was maintained
late in conditioning (**Figure 5 k,l**), and thus the reward-related DA response was not significantly
different across the training period in either DMS or DLS (**Figure 5m**).

Dopamine transients tied to the CS+ showed regionally distinctive occurrence during training
(**Figure 5n**; two-way ANOVA, repeated measure, region effect: p=0.0001***, training effect:
p<0.0001**** , region x training interaction: p=0.0240 *, $F_{(2,17)}=4.682$). For example, in the NAc
and DMS there was a gradual and significant increase in the dopamine transient from the early
to late training phase (**Figure 5n**; Sidak's multiple comparison test, NAc p<0.0001, DMS
p=0.0197). Some animals exhibited a CS+-associated dopamine transient during the first few
conditioning trials (**Supplementary Figure 4**) presumably in response to the presentation of this
novel environmental event. On subsequent trials early in training this response disappeared, but
re-emerged later in training.

Figure 5o-s and Figure6 show conditioning-related dopamine transients in all three striatal
subregions. The reward-related DA transients (left top panels of **Figure 6a-c**) were reduced or
absent in unexpected reward omission tests (middle top panels of **Figure 6a-c and Figure 5s)**,
despite vigorous head entry activity **(**middle bottom panels of **Figure 6a-c,** omission rate <17%
within a session). Thus, the reward-induced dopamine increases are not simply the product of
increased reward-directed behaviors. As expected, in the CS- trials, we observed a lack of
dopamine transients accompanied by low levels of head entry activity (right top/bottom panels of
Figure 6a-c and Figure 5r). It is important to note that no responses were observed in eGFP-
expressing control animals under any of the conditioned stimulus or reward delivery conditions
despite the same training and reward-retrieval behavior (**Figure 6d**). Thus, the conditioning-
related transients observed with dLight are not due to movement of an optic patch cable or other

non-physiological changes, but rather appear to reflect changes in dopamine related to
conditioning and behavior. Interestingly, we observed sustained fluorescence increases in NAc
between 0-5 sec after CS+, but not CS-, presentation regardless of reward delivery (**Figure 5o**
and Figure 6a). These sustained responses were not observed in DMS or DLS (**Figure 5p,q,**
Figure 6b,c; Supplementary Figure 5).

Discussion

We first used dLight photometry to examine electrical stimulation-induced dopamine
increases, with simultaneous FSCV in brain slice experiments. In general, we found similar
responses with both techniques across a range of stimulus intensities. We did detect photometric
signals at slightly lower stimulus intensities, but still within the range where FSCV signals are
observed in some studies (e.g. Salinas et al., 2016). We also found that the dLight signal was
near maximal at mid and higher stimulation intensities (>400 μA) sometimes used in FSCV.
Several observations indicated that the electrical stimulation-induced changes in fluorescence we
observed were indeed due to dLight activation by dopamine released from nigrostriatal afferents.
The transients were blocked by a D1R antagonist, strongly reduced following virally expressed
Caspase3 ablation of DAT Cre+ substantia nigra neurons, and no responses were observed in
slices expressing eGFP in place of dLight. We next sought to determine if regulation of dopamine
release by the D2 autoreceptor would be similar when comparing the two methods and found that
application of quinpirole (D2 dopamine receptor agonist) resulted in a similar average inhibition
of evoked dopamine release measured with both techniques.

Interestingly, evoked photometric responses were nearly maximal at stimulation
intensities yielding a ~50% maximal voltammetric response. It is difficult to compare the sensitivity
of the two techniques, as one needs to account for differences in sampling volumes that yield a
response value per unit of measurement (e.g. 5% dF/ F_0 or 500nM DA per μm^3). Thus, the
fundamentally different sampling volumes for each method preclude statements about direct
differences in sensitivity. Nonetheless, the inherently greater sampling volumes in *in vivo* and *in*
vitro photometric methods facilitate the detection of dopamine (or any "volume" neurotransmitter)
in a way that is not possible with single electrode voltammetric methods. We must emphasize,
however, that our findings do not indicate that dLight is a more sensitive technique for detecting
dopamine release. Findings with the two techniques cannot be compared directly, given factors
including differences in the locations of dopamine detection and the carbon fiber dimensions.

However, it seems safe to say that dLight photometry has a detection capability comparable to
FSCV.

Slow and fast/phasic dopamine dynamics

Striatal (and likely extra-striatal) dopamine operates on multiple time scales and dopamine
levels, including tonic as well as slow and fast/phasic changes (Liu et al., 2021). Tonic dopamine
levels are set by the basal firing rate of midbrain dopamine neurons and possibly modulated by
local striatal mechanisms. Slowly-developing and sustained changes in dopamine occur, e.g. in
response to application of DAT blockers. In contrast, phasic dopamine changes are faster,
typically lasting on the order of seconds, and are mediated by burst firing of midbrain dopamine
neurons, local control of dopamine release (e.g. by acetylcholine or glutamate receptors on
dopamine axons), or both (Zhou et al., 2001; Cachope et al., 2012; Threlfell et al., 2012). Slow
and phasic dopamine play distinct roles in behavior and motivation (Beeler et al., 2010; Hamid et
al., 2016; Berke, 2018; Mohebi et al., 2019; Wang et al., 2021). Microdialysis sampling with
electrochemical detection techniques allows for measurement of absolute concentrations of
dopamine that reach the probe (at least when using variants such as no-net-flux). Assessment of
tonic dopamine levels within and between test sessions can also be obtained with microdialysis,
but cannot be used to assess phasic changes in dopamine levels on a behaviorally-relevant time
scale.

Like FSCV, dLight photometry cannot measure the absolute tonic extracellular dopamine
concentration. The subsecond sampling in FSCV allows for measurement of phasic dopamine
levels but due to the need to subtract a recent baseline signal, it is generally not useful for
determining slow changes in dopamine levels. A variant of FSCV using an altered waveform
(known as FSCAV; Burrell et al., 2015) can be used to measure slower changes in dopamine, but
requires slower data acquisition (10s or sec) that precludes simultaneous measurement of phasic
dopamine changes. Given these limitations, dLight (and other fluorescent biosensors) represents
a technological advance. As we demonstrate here, dLight allows for simultaneous assessment of
slow and phasic dopamine changes within an *in vivo* testing session. This was evident in our
experiments examining the effect of acute cocaine treatment on DLS dopamine levels where we
observed an increase in baseline dopamine levels that developed over several minutes following
cocaine administration. This was accompanied by increased amplitude, frequency, and duration
of phasic dopamine transients. In contrast to our *in vitro* work, we observed a cocaine-induced

increase in transient amplitude *in vivo*. This is likely due to the slower sampling rates used for our

in vivo recordings (20 Hz) which may slightly underestimate phasic responses to the highest burst

firing frequencies of dopamine neurons (up to 100Hz) which are thought to underlie phasic

dopamine transients (Hyland et al., 2002; Lohani et al., 2018). Indirect cocaine effects may also

contribute to increased transient amplitudes *in vivo*. For example, cocaine could alter midbrain

dopaminergic neuron firing *in vivo*, and this could also contribute to transient amplitudes (Covey

et al., 2014). The dopamine transients in DLS preferentially reflect locomotion in naturalistic motor

behaviors (Markowitz et al. 2023) and even after cocaine injections (Jørgensen et al. 2023).

Increased locomotion induced by cocaine injection may also contribute to increases in transient

amplitude and frequency. Such effects would not contribute to dopamine release in brain slices.

An understanding of the significance of slow and fast/phasic dopamine changes is

important. Changes in phasic dopamine release can occur in the absence of slower changes in

dopamine levels and vice versa. Also, tonic and slow, long-lasting changes in dopamine levels

may serve to influence global striatal activity or metaplasticity. For example, the chronic DA

deficiency observed in PD models is often accompanied by changes in synaptic plasticity or

altered cellular physiology (Calabresi et al., 2009; Surmeier et al., 2014; Wang and Zhang, 2016;

Zhai et al., 2018; Graves and Surmeier, 2019). Thus, the ability to simultaneously assess slow

and phasic changes in dopamine release in freely behaving animals will be of great use to the

field.

<u>Technical considerations for sampling</u>

Sampling rates can be much faster with photometry than with FSCV. FSCV sampling is

inherently limited by the electrochemical properties of the triangle wave form (~8ms) and the time

required for desorption of dopamine from the CFE surface, resulting in a maximum practical

sampling rate of up to ~50Hz (though 10Hz is typical for dopamine measurements). In contrast,

the sampling rates for photometric methods are limited by the binding and unbinding rates of the

sensor and the digitization rate of the data acquisition hardware used. Our *in vitro* photometric

responses were collected at 100Hz or 1,000Hz, although rates up to 20,000Hz are possible with

our hardware configuration. *In vivo* fiber photometry sampling rates with most available systems

can also exceed 100Hz. However, given the known time course of dopamine signaling and dLight

kinetics we chose to perform our recordings at 20Hz. It is important to note that, with photometric

methods, the fluorescent on/off rates of dLight (or other biosensors) will limit the utility of high

sampling rate data. For example, the on and off rates for dLight1.1 are ~10 and 100ms,

respectively (Patriarchi et al., 2018; Sabatini and Tian, 2020). Thus, it is possible that sampling at

200Hz with dLight1.1 will yield a large data set that would not differ practically from a data set

collected at 100 or even 20Hz. These fluorescence kinetics may also limit interpretation of

transient rise times or modeling of decay kinetics, as the fluorescent signal may not actually

represent the termination of neuromodulator signaling but rather the off rate of the sensor. Thus,

consideration of the experimental question and the type of data that can be collected should be

considered when choosing a fluorescent biosensor (Labouesse et al., 2020; Sabatini and Tian,

2020).

<u>Synaptic dopamine vs overflow</u>

Using dLight it may be possible to assess truly "synaptic" dopamine release. That is to

say, both microdialysis and FSCV are methods that measure dopamine overflow out of the

synapse collected or detected at a site distant from most release sites. Because of the size of the

probes, sampling of dopamine at synaptic release sites has not been possible with either of these

methods. In this context, dLight represents another advance for the field because it can be

genetically targeted to distinct cellular compartments. For example, dLight sensors are integral

membrane proteins so they will only be expressed in membranes. We confirmed this with our

electron microscopy work and show that indeed, dLight traffics strongly to plasma membrane,

Golgi, and ER compartments of the neuron (**Figure 3c**). Further, our results show that dLight

traffics to synaptic and nearby extrasynaptic sites (**Figure 3b,d**). Thus, dLight signals likely

represent dopamine release that acts at these proximal sites. Work from the Ford and Williams

labs (Ford et al., 2009; Marcott et al., 2014; Mamaligas et al., 2016) utilized an indirect

electrophysiological approach to measure GIRK-mediated currents activated by D2 dopamine

receptors. Like the dLight method we used, this approach likely represents a mixture of synaptic

and extrasynaptic actions of dopamine, but is not amenable to *in vivo* use. More recent work from

the Williams lab used two photon excitation with dLight to measure dopamine release at spatially

discrete sites in the midbrain (Condon et al., 2021). These authors concluded that the time course

of D2 receptor-mediated responses was dictated largely by dopamine release and not diffusion.

Given the sparsity of dopamine release sites in midbrain, it is possible that these spatially discrete

sites represent dopamine synapses. Thus, the use of dLight with high resolution imaging methods

may allow for measurement of synaptic dopamine release. Furthermore, with dual color

fluorescence imaging methods, synaptic markers could be employed to allow for labeling of pre-

or post-synaptic elements to colocalize with dLight (or other biosensor) signals, facilitating

measurement of truly synaptic neuromodulator release.

A new method settles an old controversy

Pharmacologically, the direct mechanism of action of cocaine is inhibition of DAT (and
other monoamine transporters). This should result in prolonged phasic dopamine transients and,
in regions where the DAT function is the primary mechanism of dopamine clearance, increases
in tonic dopamine levels. Indeed, with microdialysis, the expected increases in tonic dopamine
have been confirmed (Church et al., 1987; Hurd et al., 1988). Cocaine increases the duration of
evoked dopamine transients measured with FSCV. Interestingly, measurements with FSCV show
that cocaine (and other DAT blockers) also increase the observed dopamine transient peak
amplitude in brain slices. This is often interpreted to mean that DAT blockers enhance or facilitate
electrically evoked dopamine release in slice. From a pharmacological perspective, this is not
intuitive as dopamine release should not be affected by DAT inhibition. Interestingly, Patriarchi et
al. (2018) noted that there was no increase in dLight dopamine transient peak amplitude in
response to cocaine application. Following up on this, we examined cocaine effects on dopamine
transient peak amplitude and duration in our simultaneous dLight and FSCV recordings. Similar
to previous work, we observed a large cocaine-induced increase in dopamine transient peak
amplitude with FSCV but only modest or no effects with dLight (**Figure 2**). With both methods,
the dopamine transient decay was prolonged by the drug. Similar results were obtained with the
more specific DAT blocker nomifensine. To examine this discrepancy, we first considered that the
physical interaction of dopamine with the carbon fiber electrode may be the culprit. We thus
modified the triangle waveform used in typical FSCV to reduce dopamine adsorption/sensitivity
at the carbon fiber electrode. This modification resulted in a loss of the nomifensine-induced
increases in dopamine transient peak amplitude normally observed in FSCV recordings with the
traditional voltage triangle waveform (**Figure 2h-j**). We posit that if DAT inhibitors did in fact
increase dopamine release (Wu et al., 2001; Venton et al., 2006; Oleson et al., 2009; Kile et al.,
2010; Yorgason et al., 2011; Covey et al., 2014; Hoffman et al., 2016; Salinas et al., 2016), then
the change in CFE sensitivity to dopamine with the modified waveform should not adversely affect
putative DAT blocker-induced increases in dopamine release.

In a follow up experiment, we applied cocaine to dorsal striatal brain slices from DAT KO
mice. We did not observe any cocaine-induced increase in dopamine release (**Figure 2**) in DAT
KO mice. Our observation that dopamine transients were not prolonged by cocaine in DAT KO
mice is consistent with the lack of functional transporter. These findings are in line with previous
work in DAT KO mice showing no contribution of other catecholamine transporters to DA
clearance in NAc (Budygin et al., 2002; Mateo et al., 2004).

Altogether, our data do not support the idea of DAT blocker-induced increases in
dopamine release. Therefore, we posit the following alternative interpretation of the observed
DAT blocker-induced increases in dopamine transient peaks using FSCV. Rather than increasing
or facilitating dopamine release, DAT blockers by virtue of their inhibition of dopamine clearance
allow for an increase in the point diffusion of dopamine from its release site. That is to say, under
normal conditions, the diffusion of dopamine is limited by several factors, most notably DAT
(Cragg and Rice, 2004; Hoffman et al., 2016), but under conditions of impaired DAT function (e.g.
in the presence of a DAT blocker) the diffusional spread of dopamine from its release site is
increased (Venton et al., 2003). This would effectively increase the sampling volume of CFEs
allowing for greater FSCV detection of dopamine (and increased dopamine transient peak
heights) in the absence of an actual increase in release. Further, because the sampling volume
used in photometric methods is already much larger than with FSCV methods, the cocaine-
induced increase in dopamine diffusion from its release site has a more negligible effect on
dLight/photometric DA transients. Thus, we believe that dLight more faithfully reflects dopamine
release spatial dynamics than FSCV, especially under conditions where dopamine overflow or
uptake may be affected.

<u>dLight allows for insights into striatal subregion dopamine release differences *in vivo*</u>

We compared dopamine transients across the three striatal subregions: the NAc, DMS,
and DLS as animals learned a Pavlovian association between a CS+ and reward. We found that
dopamine transient decay times were faster in DLS than in NAc, with intermediate levels in DMS.
These findings are consistent with the kinetics of electrical stimulation-induced dopamine
transients measured in brain slices and *in vivo* (Garris et al., 1994; Calipari et al., 2012; although
see Brown et al, 2011). However, there is to date little information about subregional differences
in dopamine transients driven by environmental stimuli or associated with behavior and measured
in vivo.

Our findings reveal several interesting differences in dopamine dynamics over the course
of Pavlovian conditioning in the different striatal subregions. While responses to the CS+ generally
increase over the course of training, the responses to reward delivery are maintained in DMS and
DLS, but reduced over training in NAc. These responses in dorsal striatum did not track with head
entries, as observed in the unexpected omission trials (Figures 5p,q and 6b,c), indicating that they
are dependent on reward delivery per se. The CS+ and reward-related responses in NAc are

consistent with previous findings examining dopaminergic neuronal firing and FSCV (e.g. Roitman
et al., 2004; Schultz et al., 1997). The sustained elevation between CS+ delivery and reward is
not always observed, but has been seen in past studies that used FSCV or dLight fiber photometry
during performance of *in vivo* learning tasks (Hamid et al., 2016; Mohebi et al., 2019). In our
experiments this increase appeared to be tied to expectation of reward delivery (i.e. Figures 5o,
6a and Supplementary Figure 4). Sustained responses of this type were not observed in DLS or
DMS. It will be interesting to determine the neural basis of this sustained increase in dopamine,
as well as if and how it contributes to task performance.

In vivo fiber photometry technical difficulties include potential artifacts related to fiber
bending which can result in light exposure to fiber cladding, usually related to animal movement.
Thus, it is important to include crucial analyses and control procedures that can detect such
contaminating signals. We and others have used time-correlated single photon counting-based
fiber photometry to measure signals with a variety of genetically-encoded sensors in tasks
including the open field, operant lever pressing and the accelerating rotarod (Cui et al., 2013; Cui
et al., 2014; Kupferschmidt et al., 2017). In general, we find few artifacts due to light entering the
cladding unless fibers are severely kinked which generally does not happen in open field or
operant box settings. Unlike continuous-wave (CW) laser or light emitting diode (LED), the MHz
pulsed laser system used in the TCPSC measurement also minimizes the period of light exposure
to the fiber cladding and has a high-pass filtering effect that reduces the likelihood of fiber bending-
based artifacts. This system does not include the "isosbestic" excitation control used in several
fiber photometry systems (Kim et al., 2016). Instead, using eGFP alone fluorescent controls in
the current study and in previous work from the lab we observed no evidence of artifactual
changes in fluorescence (Cui et al., 2013; Kupferschmidt et al., 2017). For these reasons we are
confident that the dLight signals we measured *in vivo* with TCSPC truly indicate changes in striatal
dopamine. Likewise, several control experiments including recordings with eGFP alone, blockade
of the signal by SCH23390 and loss of signal after lesioning SNc neurons indicate that the
stimulus-induced fluorescence increases we observed in brain slices reflect dopamine increases,
and these findings are consistent with data from our previous studies indicating that photometry
signals using this approach are not contaminated by endogenous fluorescence or other non-
sensor sources (Sgobio et al., 2014; Kupferschmidt and Lovinger, 2015).

Dopamine release in DLS was measured previously using FSCV in rat conditioning and
y-maze paradigms (Brown et al. 2011, Howe et al., 2013, Klanker et al. 2017). The findings in
NAc and DMS in the Brown et al. study are similar to those that we observe with dLight fiber

photometry in well-trained mice. However, measurements with FSCV in DLS show little-to-no fast

dopamine release in response to unexpected food reward or a positive discriminative stimulus

(DS+), and a non-significant slower-developing increase that persists for a few seconds after DS+

presentation. The behavioral correlate of this late, slow component of dopamine release is

unclear, but it might be related to movement. From this study it was not clear if the lack of

dopamine changes in DLS were due to the inability of FSCV to detect dopamine in this region or

if DLS dopamine increases are not produced by reward or predictive stimuli. Our findings indicate

that both stimulus and reward-related dopamine increases can be detected in DLS in mouse using

dLight fiber photometry, consistent the former conclusion. It should also be noted that Brown and

coworkers did not provide information regarding the time course of dopamine release changes

over the course of training, as we have been able to do with dLight fiber photometry. In the Howe

et al. (2013) study, gradual "ramping" increases in dopamine were observed in DLS on less than

half the trials as rats approached the goal in a Y-maze task, but no fast/phasic changes in

dopamine were observed. The Klanker et al. (2017) study showed small increases in DLS that

appeared to be related to movement initiation in a well-learned operant task, while van Elzelingen

et al. (2022) showed dopamine signals in rat DLS during Pavlovian conditioning behavior. These

responses were not followed over the course of initial training, but were examined during the

course of reversal training. Willuhn and coworkers (2012) were able to detect fast/phasic

dopamine increases in rat DLS that developed over the course of cocaine self-administration

using FSCV. However, the prolongation of dopamine increases produced by this DAT blocker

likely facilitated detection of this increase. It should now be clear that past studies showed

dopamine detection in rat DLS, but our findings provide new information on responses to drugs

and environmental events in this important striatal subregion in mouse, as well as evidence that

we can measure both slow and fast changes in dopamine simultaneously throughout the course

of pharmacological and extended behavioral studies.

Materials and Methods

Subjects

Three-month-old, male C57BL/6J mice were obtained from the Jackson Laboratory (Strain 000664) and pair-housed in the vivarium for at least one week before any experimental use. DAT-IRES-Cre mice were obtained from the Jackson Laboratory (Strain 006660) and bred in house. DAT KO mice were developed as previously described (Giros et al., 1996) and obtained from the Sara Jones laboratory at Wake Forest University. Male and female DAT-IRES-Cre and DAT KO transgenic mice were used in all experiments. All mice were housed with 2-4 mice per cage and maintained on a 12:12 hour light cycle and ad libitum access to food and water.

All procedures performed in this work follow the guidelines of the Institutional Animal Care and Use Committee of the Division of Intramural Clinical and Biological Research, National Institute on Alcohol Abuse and Alcoholism, and the animal care and use protocol was approved by this committee.

Viruses and Stereotaxic Injections

The Caspase3-coding virus (AAV1-EF1a-FLEX-taCasp3-TEVp) was custom packaged by Vigene Biosciences and a gift from Dr. Huaibin Cai, National Institute on Aging. dLight1.1 viruses were used for *in vitro* experiments and were either generated in the Tian laboratory and sent to NIAAA (AAV5-CAG-dLight1.1 and AAV9-CAG-dLight1.1) or purchased from Addgene (pAAV5-CAG-dLight1.1; Addgene viral prep # 111067-AAV5). AAV9-CAG-dLight1.1 are only partially used in Fig2. All other in-vitro experiments used AAV5-CAG-dLight1.1. dLight1.3b virus was used for *in vivo* experiments and supplied by theTian laboratory or custom packaged by Vigene Biosciences (AAV9-CAG-dLight1.3b).

All stereotaxic injections were conducted using sterile technique on mice at least 3 months of age. Mice were anesthetized with a 5% isoflurane/oxygen mixture and placed in a Kopf stereotaxic frame. Anesthesia was maintained with 1-2% isoflurane/oxygen mixture. The skulls were leveled and an incision was made to expose the skull. Craniotomies were made over the dorsal striatum (AP +1.0, ML +/-1.8, from Bregma in mm), nucleus accumbens (AP +1.2, ML +/-0.8, from Bregma in mm) or substantia nigra (AP -3.0, ML +/-1.2, from Bregma in mm) and a 1 µL Neuros Hamilton Syringe was lowered slowly to the desired depth from the brain surface (-2.25mm, -3.75mm, and -4.1mm for dorsal striatum, nucleus accumbens, and substantia nigra, respectively). For in-vivo experiments, viruses were injected in the following coordinates. NAc (AP +1.2, ML +/-0.8, DV -3.75), DMS (AP +1.0, ML +/-1.2, DV -2.25), DLS (AP +0.8, ML +/-2.2, DV -2.25) where AP and ML in mm from Bregma, DV in mm from the brain surface. Virus infusion volumes were 300nL for dorsal striatum and 500nL for VTA and viruses were infused at 50nL/minute. After infusions were completed, the syringe was left in place for 10 minutes before withdrawal and the incisions were closed with VetBond. Mice recovered for at least three weeks before being used for in vitro experiments or fiber optic implantation for *in vivo* experiments.

Simultaneous Photometry and Voltammetric Recordings

Dual photometric and voltammetric recordings were conducted as previously described (Sgobio et al., 2014; Sgobio et al., 2019) except that the photometry recordings in the current study used dLight1.1 or dLight1.3b to compare dopamine release between voltammetric and photometric methods simultaneously in the same preparation. Brain slices containing the striatum were

prepared as previously described (Salinas et al., 2016). Briefly, mice were anesthetized with isoflurane, rapidly decapitated, brains extracted, and 300μm thick coronal sections prepared on a vibratome (Leica VT 1200S). The slices were hemisected and inspected to ensure viral expression of dLight in the region of interest using an epifluorescent Zeiss AxioZoom microscope equipped with a GFP filter set (Carl Zeiss Microscopy Filter Set Lumar # 38 BP470/40, FT495, BP525/50). Then the slices were incubated at 32°C for 30 minutes before being moved to room temperature for one hour before beginning experiments.

Brain slices with dLight expression were moved to an upright Zeiss AxioSkop2 microscope mounted on a XY translational stage and equipped with a GFP filter set. Oxygenated ACSF was perfused at 1.5-2mL/minute and warmed to 30-32°C. The recording region of interest was located under 4X magnification and fluorescent illumination to ensure dLight expression in the region of interest. Then a stainless steel twisted bipolar stimulating electrode (P1 Technologies) was placed on the tissue surface near the area of dLight expression. For simultaneous photometry-voltammetry recordings, carbon fiber electrodes (CFE) were fabricated as previously described (Salinas et al., 2016; Salinas et al., 2021) and placed in the tissue in the center of the recording region of interest under 4X magnification. Slices were then visualized with 40X objective (0.8 NA) and the field of view (~180um x 180um) was adjusted so the stimulating electrode was just outside the field of view and the CFE was in the center of the field of view. Under 40X magnification, the focus was adjusted to a focal layer beneath the slice surface where fluorescent cells could be identified. Fluorescent transients were quantified with a PMT-based system (PTI D-104 photometer) coupled with a Digidata 1322A (Molecular Devices LLC) to digitize the PMT signal (100-1,000Hz). Clampex software was used to collect photometry data and synchronize photometric and voltammetric recordings through the Digidata. A mechanical shutter (Uniblitz V25) was used to limit exposure to fluorophore-exciting light to discrete periods and minimize photobleaching of the dLight between recordings. FSCV recordings were carried out as previously described (Sgobio et al., 2014; Salinas et al., 2016; Sgobio et al., 2019; Salinas et al., 2021). Briefly, a triangle wave form voltage was applied to the CFE beginning at -0.4V to +1.2V and back to -0.4V. This scan was applied at 400V/s and repeated at 10Hz. Dopamine was identified electrochemically by the oxidation peak at +0.6V on the ascending phase of the triangle ramp. Dopamine release was evoked with electrical stimulation delivered every three minutes using a constant current stimulus isolator (DigiTimer DS3). Input-output (IO) curves were generated to examine evoked dopamine release measured with both techniques across varying electrical stimulation intensities (50-800 μA, 1 ms). For pharmacological experiments, a stimulation intensity yielding approximately 30-60% of the maximal responses with both methods was used to ensure that any subsequent treatments were not limited by floor or ceiling response effects. Baseline responses were collected for 12-20 minutes before drugs (dissolved in ACSF) were bath applied as indicated for each experiment.

In Vivo Fiber Photometry

At least three weeks after virus infusion surgeries, mice used for _in vivo_ experiments underwent a second surgery to implant the optical fiber, similar to those previously described (Kupferschmidt et al., 2017). Briefly, mice were anesthetized and mounted into the stereotaxic apparatus and an incision was made in the scalp along the skull midline as before. The skull was cleaned with a 3% H_2O_2 solution to remove any connective tissue and the skull was scored several times with a scalpel to create a better surface for the dental cement headcap at the end of the procedure. Craniotomies were made over the brain region of interest and two distal sites for anchor screw

placement. Fiber implants (Thorlabs, # CFMC 12L05) were placed in the following coordinates: NAc (AP +1.2, ML +/-0.8, DV -3.75), DMS (AP +1.0, ML +/-1.2, DV -2.25), DLS (AP +0.8, ML +/-2.2, DV -2.25) where AP and ML in mm from Bregma, DV in mm from the brain surface. The anchor screws were placed before proceeding to the fiber optic placement. To ensure optimal fiber optic placement, fluorescence intensity was monitored intraoperatively using a custom designed fiber photometry system consisting of a 473 nm picosecond pulsed laser, a HPM-100-40 hybrid detector, and a SPC-130EM time correlated single photon counting (TCSPC) module and software (Becker & Hickl). The measurement rate for the fluorescence lifetime and intensity profile was set at 20Hz during all animal experiments. The output was a custom multimode patch cord from Thor labs with a 0.22NA and 200µm fiber core diameter terminating in a 2.5mm ceramic ferrule. The patch cord ferrule was connected to the fiber optic cannula (CFMC22L05, Thor Labs) to be implanted with a ceramic mating sleeve. The implantable fiber optic cannula was lowered slowly into the craniotomy over the brain region of interest while the fluorescence intensity at the desired emission wavelengths (~500-540 nm) was monitored. The fluorescence intensity typically increased as the fiber optic approached the area of dLight expression. Once fluorescence intensity plateaued, typically ~300-500µm dorsal to the virus infusion site, the fiber optic cannula was cemented in place and the surgical site was closed around the dental cement headcap. Mice recovered for at least two weeks before further experiments.

For the Pavlovian conditioning experiment, the TCSPC system was synchronized to the operant behavior boxes (MED-PC) through a TTL channel, in which the TTL pulse generated by MED-PC controller was used to trigger the TCSPC system acquisition for 10s prior to each trial (CS+ or CS-) and maintained 60s of data acquisition. The pulsed laser was continuously turned on and sustained stable laser illumination during the entire session. *In vivo* pharmacology measurements were continuously recorded for 60min. Before the recording, the animals were tethered with an optical patch cord with laser illumination, and habituated for 20 min in the open-field arena without recording.

In Vitro Fiber Photometry Benchtop Testing

To test the possibility of patch cable vibration effects on fluorescent signals captured with TCSPC we devised a benchtop testing system (Supplementary Figure 3). The tip of patch cable connect to our *in vivo* photometry system was mounted on an optical table, and a sample tube containing a fluorescent polymer (World Precision Instruments, cat # KWIK-CAST) was placed at the tip of patch cable. Single-photon counting measures of fluorescence using the same laser and detector used for *in vivo* photometry were carried out for 5 min under conditions in which the patch cable was stable, and when the cable was vibrated by movements simulating those that occur was mice traverse the behavioral apparatuses used in our experiments.

Pavlovian Conditioning

At least three weeks following fiber optic implant surgeries, mild food restriction was carried out to gradually reduce the animals' body weight to 85-90% of their initial body weight. Thus, chow was limited to 2–3 g per day for 3 days. Concurrently, mice were handled and tethered to an optical patch cord cable within the designated behavior training time. For the first 2 days of pre-training, mice received habituation sessions in the operant training box for 1-hr while tethered to the optical patch cord. In the final day of pre-training, mice underwent a magazine training session in which 30 food pellets were delivered on a random-interval schedule (RI-60). No stimulus cue was provided during these sessions. After 3 days of food restriction and pre-training, mice

underwent Pavlovian conditioning sessions. The discrimination training entailed two conditioned
stimuli; one conditioned stimulus (CS+) was a high-frequency auditory cue followed by the
unconditioned stimulus (US, one 15mg food pellet) as a reinforcer, and a second stimulus (CS-)
was a low-frequency auditory cue which was not accompanied by a reinforcer delivery. Within a
single session, mice received 20 CS+ trials and 20 CS- trials. In the operant conditioning box
within sound and light attenuating enclosures, the auditory cue was presented for 4 seconds with
random intertrial intervals (60-sec to 120-sec). In the CS+ trials, an US reinforcer was delivered
to a pellet receptacle 1 second after the CS+ cue offset. *In vivo* photometry measurements were
conducted every other day during the training, but the mice underwent the same continuous daily
Pavlovian training sessions while tethered to an optical patch cord (Thor Labs, 200um core, 3 m
length) even if photometry measurements were not conducted. Therefore, DLS and DMS mice
received 10 weeks of consecutive daily Pavlovian training. NAc mice received four weeks of
Pavlovian training. All animals received the behavior training with 24 (+/-2) hours of interval at
their own designated time, and animals were acclimated to the behavior room for at least 1hr
before all training sessions.

<u>Fluorescence microscopy</u>

For anatomical studies, dLight virus injected mice were deeply anesthetized with chloral hydrate
(35 mg per 100 g), and perfused transcardially with 4% (wt/vol) paraformaldehyde (PFA) with
0.15% (vol/vol) glutaraldehyde and 15% (vol/vol) picric acid in 0.1 M phosphate buffer (PB, pH
7.3). Brains were left in this fixative solution for 2 h at 4 °C, solution was replaced with 2% PFA
and left overnight at 4 °C. Brains were rinsed with PB, and cut into coronal serial sections (40 μm
thick) with a vibratome (Leica). All animal procedures were approved by the National Institute on
Drug Abuse Animal Care and Use Committee.

Free floating coronal vibratome sections were incubated for 1 h in PB supplemented with 4%
bovine serum albumin (BSA) and 0.3% Triton X-100. Sections were then incubated with mouse
anti-tyrosine hydroxylase (TH) primary antibody (1:1,000 dilution, Millipore-Sigma, Cat# MAB318,
RRID: AB_2201528) and guinea pig anti GFP (1:500, Nittobo Medical, Cat# GFP-GP-Af1180,
RRID: AB_2571575) overnight at 4°C. After rinsing 3 × 10 min in PB, sections were incubated in
Alexa Fluor 594-affiniPure donkey anti-mouse (1:100 dilution, Jackson Immunoresearch
Laboratories, Cat# 715-585-151, RRID: AB_2340855) and Alexa Fluor 488-affiniPure Donkey
anti guinea pig (1:100, Jackson Immunoresearch Laboratories, Cat# 706-545-148, RRID:
AB_2340472) for 2 h at room temperature. After rinsing, sections were mounted on slides and
air-dried. Fluorescent images were collected with Zeiss LSM880 Airyscan Confocal System
(Zeiss, White Plains, NY). Images were taken with 20× objectives and z-axis stacks were
collected at 1 μm. The confocal images were collected from 3 mice.

<u>Electron microscopy</u>

Vibratome tissue sections were rinsed with PB, incubated with 1% sodium borohydride in PB for
30 min to inactivate free aldehyde groups, rinsed in PB, and then incubated with blocking solution
[1% normal goat serum (NGS), 4% BSA in PB supplemented with 0.02% saponin] for 30 min.
Sections were then incubated with guinea pig anti-GFP primary antibody (1:500 dilution, Nittobo
Medical, Cat# GFP-GP-Af1180, RRID: AB_2571575); or both guinea pig anti-GFP (1:500 dilution)
and mouse anti-TH (1:1,000 dilution) for 24 h at 4°C. Sections were rinsed and incubated
overnight at 4°C in the secondary antibody goat anti-guinea pig IgG Fab fragment coupled to 1.4
nm gold (1:100 dilution for GFP detection; Nanoprobes Inc., Cat# 2055, RRID: AB_2802149); or

743 the corresponding secondary antibodies: biotinylated goat anti-guinea pig antibody (1:100 dilution
744 for GFP detection; Vector Laboratories, Cat# PK-4007; RRID:AB_2336816) and secondary
745 antibody goat anti-mouse IgG coupled to 1.4 nm gold (1:100 dilution for TH detection; Nanoprobes
746 Inc., Cat# 2001, RRID:AB_2877644). Sections were incubated in avidin-biotinylated horseradish
747 peroxidase complex in PB for 2 h at room temperature and washed. Peroxidase activity was
748 detected with 0.025% 3,3′-diaminobenzidine (DAB) and 0.003% H_2O_2 in PB for 5-10 min. Sections
749 were rinsed in PB, and then in double-distilled water, followed by silver enhancement of the gold
750 particles with the Nanoprobe Silver Kit (2012; Nanoprobes Inc., Stony Brook, NY) for 7 min at
751 room temperature. Next, sections were rinsed with PB and fixed with 0.5% osmium tetroxide in
752 PB for 25 min, washed in PB followed by double-distilled water and then contrasted in freshly
753 prepared 1% uranyl acetate for 35 min. Sections were dehydrated through a series of graded
754 alcohols and with propylene oxide. Afterwards, they were flat embedded in Durcupan ACM epoxy
755 resin (14040; Electron Microscopy Sciences, Fort Washington, PA). Resin-embedded sections
756 were polymerized at 60°C for 2 days. Sections of 65 nm were cut from the outer surface of the
757 tissue with an ultramicrotome UC7 (Leica Microsystems Inc., Buffalo Grove, IL) using a diamond
758 knife (Diatome, fort Washington, PA). The sections were collected on formvar-coated single slot
759 grids and counterstained with Reynolds lead citrate. Sections were examined and photographed
760 using a Tecnai G2 12 transmission electron microscope (Fei Company, Hillsboro, OR) equipped
761 with a digital micrograph OneView camera (Gatan Inc., Pleasanton, CA).

763 Ultrastructural analysis

764 Serial thin sections of dorsal striatum of the mice were used in this study. Synaptic contacts were
765 classified according to their morphology and immuno-label and photographed at a magnification
766 of 6800-13000×. The morphological criteria used for identification and classification of cellular
767 components observed in these thin sections were as previously described (Zhang et al., 2015).
768 Type I synapses, here referred as asymmetric synapses, were defined by the presence of
769 contiguous synaptic vesicles within the presynaptic axon terminal and a thick postsynaptic density
770 (PSD) greater than 40 nm. Type II synapses, here referred as symmetric synapses, were defined
771 by the presence of contiguous synaptic vesicles within the presynaptic axon terminal and a thin
772 PSD. Serial sections were obtained to determine the type of synapse. In the serial sections, a
773 terminal or dendrite containing greater than 5 immunogold particles were considered as immuno-
774 positive terminal or dendrite. Pictures were adjusted to match contrast and brightness by using
775 Adobe Photoshop (Adobe Systems Incorporated, Seattle, WA). The frequency of gold particles
776 for GFP signals near to asymmetric synapses was counted from 3 mice. This experiment was
777 successfully repeated three times.

778 Photometry Data Analysis

779 *In vitro* photometric data were acquired in Clampex 9 and initially analyzed using Clampfit v9 or
780 v10 (Axon Instruments/Molecular Devices), exported to Excel for organization, and plotted and
781 analyzed in GraphPad Prism 7. The PMT readout was offset corrected to zero (with the shutter
782 closed) before basal fluorescence (F_0) and stimulation-induced increases in fluorescence (dF)
783 values were obtained. Most photometric data are presented as dF/F_0 values to compensate for
784 differences in basal fluorescence or dLight expression differences between slices. Voltammetric
785 data were acquired in DEMON Voltammetry Software suite (Yorgason et al., 2011), exported to
786 Excel for organization, and plotted and analyzed in GraphPad Prism 9.

In vivo fiber photometry data were collected using the SPCM64 9.8 Software (Becker & Hickl).
The raw photon counts were exported in ascii format for analysis with custom Python or MATLAB
scripts. Briefly, the fluorescence (F_{raw}) values were plotted as a function of time. Then, these
fluorescence time series were converted to dF/F_0 in two ways. For the segmented photometry
data (60-s per individual trial) obtained from operant conditioning sessions, F_0 was set to a moving
average of each point before the onset of CS+, using a sliding window of +/-15s, similar to
previously published methods (Patriarchi et al., 2018). Therefore, the baseline was normalized
across all measurements such that only phasic dopamine transients were quantified.
Occasionally, there was a negligible bleaching effect (<0.1% of F_0 degradation) within the 60-s of
acquisition in our measurement setup even without the moving average normalization. For the
continuous photometry measurements over 60min duration in the *in vivo* pharmacology
experiments, actual recording was performed after a 20min initial bleaching period without
recording. The F_0 was estimated by a curve fitting method. We first validated that the degradation
of fluorescence intensity was well predicted by an exponential decaying model. For this validation,
we used control mice virally transduced to express GFP in dorsal striatum and dLight expressing
mice having no drug injection to confirm the baseline prediction method. Then, for the actual
experimental animals with cocaine and saline injections, the baseline curves of F_0 were estimated
from the 20-min baseline measurement prior to the i.p. injection. Once the F_0 baseline was
estimated, dF/F_0 was calculated using the standard method, in which $dF = F_{raw}-F_0$ and F_0 is the
predicted baseline by exponential curve fitting.

Dopamine Transient Decay Analysis

For each CS+ trial, maximum peak location was identified using custom Matlab code. Then the
fluorescence profile (dF/F_0) was normalized to the maximum intensity. The data points following
the maximum intensity peak were fit using a double exponential decay model. In the fitted curve,
the time point where the normalized fluorescence profile passed under 36.8% of the maximum
intensity was selected as the transient decay time (or lifetime) of the phasic dopamine activity in
each brain regions.

Drugs and Reagents

Cocaine-HCl was obtained from the National Institute on Drug Abuse. Dihydro-β-erythroidine
hydrobromide (DHBE), nomifensine, quinpirole-HCl, and SCH23390 were purchased from Tocris
Bioscience. Dopamine-HCl was purchased from Sigma Aldrich. All drugs were dissolved in ACSF.
All other drugs and reagents, unless otherwise indicated, were obtained from Sigma Aldrich.

Statistics

GraphPad Prism 9.2 was used for all data analysis and statistics. For I-O curve comparison, two
way ANOVA (method and stimulation intensity) was used. For the analysis of the caspase
lesion curves, a mixed effects model was used (genotype and stimulation intensity factors). For
the quinpirole inhibition and high calcium experiments, unpaired two-tailed t-tests were used.
For data comparing *in vivo* transient amplitude, frequency, and decay constant changes,
unpaired two-tailed t-tests were used. For analysis of the dorsal striatum subregion changes in
decay constant, one way ANOVA was used. Unless otherwise indicated, all data represent the
mean ± SEM.

Acknowledgements

We would like to thank Dr. Huaibin Cai for the Casp3 virus, Dr. Joseph Cheer for his comments on the data and manuscript, Dr. Gabriel Loewinger for advice on data analysis/statistics, and Jacob Nadel and Lucas Voyvodic for their expert technical assistance with stereotaxic virus infusions. We also thank Rong Ye and Kevin Yu of the Confocal and Electron Microscopy Core, NIDA IRP, for collecting confocal and immune-electron microscopic images used in this study.

This work was supported by research grants from the National Institute on Alcohol Abuse and Alcoholism (K99R00 AA025991 to AGS, K99R00 AA027740 to SMA, and ZIA AA000407 & ZIA000416 to DML), the Intramural Research Program of the National Institute on Drug Abuse, the BRAIN Initiative (U01NS090604 and U01NS013522 to LT), and the National Institutes of Health (DP2MH107056 to LT).

Author Contributions

Designed Research: AGS, JOL, MM, YM, DML

Contributed Essential Reagents: TP, LT

Performed Research: AGS, YM, JOL, SMA, SZ

Analyzed Data: AGS, YM, JOL, SMA, SZ, MM

Wrote the Paper: AGS, JOL, YM, DML

Competing Interests

The authors declare no competing interests.

Bibliography

Acevedo-Rodriguez A, Zhang L, Zhou F, Gong S, Gu H, De Biasi M, Zhou FM, Dani JA (2014) Cocaine inhibition of nicotinic acetylcholine receptors influences dopamine release. Front Synaptic Neurosci 6:19.

Beeler JA, Daw N, Frazier CR, Zhuang X (2010) Tonic dopamine modulates exploitation of reward learning. Front Behav Neurosci 4:170.

Berke JD (2018) What does dopamine mean? Nat Neurosci 21:787-793.

Brown HD, McCutcheon JE, Cone JJ, Ragozzino ME, Roitman MF (2011) Primary food reward and reward-predictive stimuli evoke different patterns of phasic dopamine signaling throughout the striatum. Eur J Neurosci 34:1997-2006.

Budygin EA, John CE, Mateo Y, Jones SR (2002) Lack of cocaine effect on dopamine clearance in the core and shell of the nucleus accumbens of dopamine transporter knock-out mice. J Neurosci 22:RC222.

Burrell MH, Atcherley CW, Heien ML, Lipski J (2015) A novel electrochemical approach for prolonged measurement of absolute levels of extracellular dopamine in brain slices. ACS Chem Neurosci 6:1802-1812.

Cachope R, Mateo Y, Mathur BN, Irving J, Wang HL, Morales M, Lovinger DM, Cheer JF (2012) Selective activation of cholinergic interneurons enhances accumbal phasic dopamine release: setting the tone for reward processing. Cell Rep 2:33-41.

Calabresi P, Mercuri NB, Di Filippo M (2009) Synaptic plasticity, dopamine and Parkinson's disease: one step ahead. Brain 132:285-287.

Calipari ES, Huggins KN, Mathews TA, Jones SR (2012) Conserved dorsal-ventral gradient of dopamine release and uptake rate in mice, rats and rhesus macaques. Neurochem Int 61:986-991.

Chefer VI, Thompson AC, Zapata A, Shippenberg TS (2009) Overview of brain microdialysis. Curr Protoc Neurosci Chapter 7:Unit7 1.

Chen D, Gao F, Ma X, Eaton JB, Huang Y, Gao M, Chang Y, Ma Z, Der-Ghazarian T, Neisewander J, Whiteaker P, Wu J, Su Q (2019) Cocaine Directly Inhibits alpha6-Containing Nicotinic Acetylcholine Receptors in Human SH-EP1 Cells and Mouse VTA DA Neurons. Front Pharmacol 10:72.

Church WH, Justice JB, Jr., Byrd LD (1987) Extracellular dopamine in rat striatum following uptake inhibition by cocaine, nomifensine and benztropine. Eur J Pharmacol 139:345-348.

Condon AF, Robinson BG, Asad N, Dore TM, Tian L, Williams JT (2021) The residence of synaptically released dopamine on D2 autoreceptors. Cell Rep 36:109465.

Covey DP, Roitman MF, Garris PA (2014) Illicit dopamine transients: reconciling actions of abused drugs. Trends Neurosci 37:200-210.

Cragg SJ, Rice ME (2004) DAncing past the DAT at a DA synapse. Trends Neurosci 27:270-277.

Cui G, Jun SB, Jin X, Pham MD, Vogel SS, Lovinger DM, Costa RM (2013) Concurrent activation of striatal direct and indirect pathways during action initiation. Nature 494:238-242.

Cui G, Jun SB, Jin X, Luo G, Pham MD, Lovinger DM, Vogel SS, Costa RM (2014) Deep brain optical measurements of cell type-specific neural activity in behaving mice. Nat Protoc 9:1213-1228.

Day JJ, Roitman MF, Wightman RM, Carelli RM (2007) Associative learning mediates dynamic shifts in dopamine signaling in the nucleus accumbens. Nat Neurosci 10:1020-1028.

Ford CP, Phillips PE, Williams JT (2009) The time course of dopamine transmission in the ventral tegmental area. J Neurosci 29:13344-13352.

Francis MM, Vazquez RW, Papke RL, Oswald RE (2000) Subtype-selective inhibition of neuronal nicotinic acetylcholine receptors by cocaine is determined by the alpha4 and beta4 subunits. Mol Pharmacol 58:109-119.

Garris PA, Ciolkowski EL, Wightman RM (1994) Heterogeneity of evoked dopamine overflow within the
striatal and striatoamygdaloid regions. Neuroscience 59:417-427.
Giros B, Jaber M, Jones SR, Wightman RM, Caron MG (1996) Hyperlocomotion and indifference to
cocaine and amphetamine in mice lacking the dopamine transporter. Nature 379:606-612.
Grace AA (2016) Dysregulation of the dopamine system in the pathophysiology of schizophrenia and
depression. Nat Rev Neurosci 17:524-532.
Graves SM, Surmeier DJ (2019) Delayed Spine Pruning of Direct Pathway Spiny Projection Neurons in a
Mouse Model of Parkinson's Disease. Front Cell Neurosci 13:32.
Hamid AA, Pettibone JR, Mabrouk OS, Hetrick VL, Schmidt R, Vander Weele CM, Kennedy RT, Aragona
BJ, Berke JD (2016) Mesolimbic dopamine signals the value of work. Nat Neurosci 19:117-126.
Heien ML, Phillips PE, Stuber GD, Seipel AT, Wightman RM (2003) Overoxidation of carbon-fiber
microelectrodes enhances dopamine adsorption and increases sensitivity. Analyst 128:1413-
1419.
Hoffman AF, Spivak CE, Lupica CR (2016) Enhanced Dopamine Release by Dopamine Transport Inhibitors
Described by a Restricted Diffusion Model and Fast-Scan Cyclic Voltammetry. ACS Chem
Neurosci 7:700-709.
Howe MW, Tierney PL, Sandberg SG, Phillips PE, Graybiel AM (2013) Prolonged dopamine signalling in
striatum signals proximity and value of distant rewards. Nature 500:575-579.
Hurd YL, Kehr J, Ungerstedt U (1988) In vivo microdialysis as a technique to monitor drug transport:
correlation of extracellular cocaine levels and dopamine overflow in the rat brain. J Neurochem
51:1314-1316.
Hyland BI, Reynolds JN, Hay J, Perk CG, Miller R (2002) Firing modes of midbrain dopamine cells in the
freely moving rat. Neuroscience 114:475-492.
Jones SR, Gainetdinov RR, Jaber M, Giros B, Wightman RM, Caron MG (1998) Profound neuronal
plasticity in response to inactivation of the dopamine transporter. Proc Natl Acad Sci U S A
95:4029-4034.
Jørgensen SH., et al (2023) Behavioral encoding across timescales by region-specific dopamine dynamics.
Proceedings of the National Academy of Sciences 120.7: e2215230120.
Kile BM, Guillot TS, Venton BJ, Wetsel WC, Augustine GJ, Wightman RM (2010) Synapsins differentially
control dopamine and serotonin release. J Neurosci 30:9762-9770.
Kim CK, Yang SJ, Pichamoorthy N, Young NP, Kauvar I, Jennings JH, Lerner TN, Berndt A, Lee SY,
Ramakrishnan C, Davidson TJ, Inoue M, Bito H, Deisseroth K (2016) Simultaneous fast
measurement of circuit dynamics at multiple sites across the mammalian brain. Nat Methods
13:325-328.
Klanker M, Fellinger L, Feenstra M, Willuhn I, Denys D (2017) Regionally distinct phasic dopamine release
patterns in the striatum during reversal learning. Neuroscience 345:110-123.
Kupferschmidt DA, Lovinger DM (2015) Inhibition of presynaptic calcium transients in cortical inputs to
the dorsolateral striatum by metabotropic GABA(B) and mGlu2/3 receptors. J Physiol 593:2295-
2310.
Kupferschmidt DA, Juczewski K, Cui G, Johnson KA, Lovinger DM (2017) Parallel, but Dissociable,
Processing in Discrete Corticostriatal Inputs Encodes Skill Learning. Neuron 96:476-489 e475.
Labouesse MA, Cola RB, Patriarchi T (2020) GPCR-Based Dopamine Sensors-A Detailed Guide to Inform
Sensor Choice for In vivo Imaging. Int J Mol Sci 21.
Liu C, Goel P, Kaeser PS (2021) Spatial and temporal scales of dopamine transmission. Nat Rev Neurosci
22:345-358.
Lohani S, Martig AK, Underhill SM, DeFrancesco A, Roberts MJ, Rinaman L, Amara S, Moghaddam B
(2018) Burst activation of dopamine neurons produces prolonged post-burst availability of
actively released dopamine. Neuropsychopharmacology 43:2083-2092.

Ma ZG, Jiang N, Huang YB, Ma XK, Brek Eaton J, Gao M, Chang YC, Lukas RJ, Whiteaker P, Neisewander J,
Wu J (2020) Cocaine potently blocks neuronal alpha(3)beta(4) nicotinic acetylcholine receptors
in SH-SY5Y cells. Acta Pharmacol Sin 41:163-172.
Mamaligas AA, Cai Y, Ford CP (2016) Nicotinic and opioid receptor regulation of striatal dopamine D2-
receptor mediated transmission. Sci Rep 6:37834.
Markowitz JE, Gillis WF, Jay M et al. (2023) Spontaneous behaviour is structured by reinforcement
without explicit reward. Nature 614: 108–117.
Marcott PF, Mamaligas AA, Ford CP (2014) Phasic dopamine release drives rapid activation of striatal
D2-receptors. Neuron 84:164-176.
Mateo Y, Budygin EA, John CE, Banks ML, Jones SR (2004) Voltammetric assessment of dopamine
clearance in the absence of the dopamine transporter: no contribution of other transporters in
core or shell of nucleus accumbens. J Neurosci Methods 140:183-187.
Mohebi A, Pettibone JR, Hamid AA, Wong JT, Vinson LT, Patriarchi T, Tian L, Kennedy RT, Berke JD (2019)
Dissociable dopamine dynamics for learning and motivation. Nature 570:65-70.
Nestler EJ, Carlezon WA, Jr. (2006) The mesolimbic dopamine reward circuit in depression. Biol
Psychiatry 59:1151-1159.
Oleson EB, Salek J, Bonin KD, Jones SR, Budygin EA (2009) Real-time voltammetric detection of cocaine-
induced dopamine changes in the striatum of freely moving mice. Neurosci Lett 467:144-146.
Ott T, Nieder A (2019) Dopamine and Cognitive Control in Prefrontal Cortex. Trends Cogn Sci 23:213-
234.
Patriarchi T, Cho JR, Merten K, Howe MW, Marley A, Xiong WH, Folk RW, Broussard GJ, Liang R, Jang MJ,
Zhong H, Dombeck D, von Zastrow M, Nimmerjahn A, Gradinaru V, Williams JT, Tian L (2018)
Ultrafast neuronal imaging of dopamine dynamics with designed genetically encoded sensors.
Science 360.
Radke AK, Kocharian A, Covey DP, Lovinger DM, Cheer JF, Mateo Y, Holmes A (2019) Contributions of
nucleus accumbens dopamine to cognitive flexibility. Eur J Neurosci 50:2023-2035.
Roitman MF, Stuber GD, Phillips PE, Wightman RM, Carelli RM (2004) Dopamine operates as a
subsecond modulator of food seeking. J. Neurosci. 24, 1265–1271
Sabatini BL, Tian L (2020) Imaging Neurotransmitter and Neuromodulator Dynamics In Vivo with
Genetically Encoded Indicators. Neuron 108:17-32.
Salinas AG, Davis MI, Lovinger DM, Mateo Y (2016) Dopamine dynamics and cocaine sensitivity differ
between striosome and matrix compartments of the striatum. Neuropharmacology 108:275-
283.
Salinas AG, Mateo Y, Carlson VCC, Stinnett GS, Luo G, Seasholtz AF, Grant KA, Lovinger DM (2021) Long-
term alcohol consumption alters dorsal striatal dopamine release and regulation by D2
dopamine receptors in rhesus macaques. Neuropsychopharmacology 46:1432-1441.
Schultz W, Dayan P, Montague, PR(1997) A neural substrate of prediction and reward. Science 275,
1593–1599.
Sgobio C, Sun L, Ding J, Herms J, Lovinger DM, Cai H (2019) Unbalanced calcium channel activity
underlies selective vulnerability of nigrostriatal dopaminergic terminals in Parkinsonian mice. Sci
Rep 9:4857.
Sgobio C, Kupferschmidt DA, Cui G, Sun L, Li Z, Cai H, Lovinger DM (2014) Optogenetic measurement of
presynaptic calcium transients using conditional genetically encoded calcium indicator
expression in dopaminergic neurons. PLoS One 9:e111749.
Sun F et al. (2018) A Genetically Encoded Fluorescent Sensor Enables Rapid and Specific Detection of
Dopamine in Flies, Fish, and Mice. Cell 174:481-496 e419.
Surmeier DJ, Graves SM, Shen W (2014) Dopaminergic modulation of striatal networks in health and
Parkinson's disease. Curr Opin Neurobiol 29:109-117.

Threlfell S, Lalic T, Platt NJ, Jennings KA, Deisseroth K, Cragg SJ (2012) Striatal dopamine release is
996   triggered by synchronized activity in cholinergic interneurons. Neuron 75:58-64.

997   Van Elzelingen W, et al (2022) Striatal dopamine signals are region specific and temporally stable across
998   action-sequence habit formation. Current Biology 32.5: 1163-1174.

Van Zessen R, et al. (2021) Dynamic dichotomy of accumbal population activity underlies cocaine
sensitization. Elife 10: e66048.

Venton BJ, Cao Q (2020) Fundamentals of fast-scan cyclic voltammetry for dopamine detection. Analyst
145:1158-1168.

Venton BJ, Zhang H, Garris PA, Phillips PE, Sulzer D, Wightman RM (2003) Real-time decoding of
dopamine concentration changes in the caudate-putamen during tonic and phasic firing. J
Neurochem 87:1284-1295.

Venton BJ, Seipel AT, Phillips PE, Wetsel WC, Gitler D, Greengard P, Augustine GJ, Wightman RM (2006)
Cocaine increases dopamine release by mobilization of a synapsin-dependent reserve pool. J
Neurosci 26:3206-3209.

Wang Q, Zhang W (2016) Maladaptive Synaptic Plasticity in L-DOPA-Induced Dyskinesia. Front Neural
Circuits 10:105.

Wang Y, Toyoshima O, Kunimatsu J, Yamada H, Matsumoto M (2021) Tonic firing mode of midbrain
dopamine neurons continuously tracks reward values changing moment-by-moment. Elife 10.

Willuhn I, Burgeno LM, Everitt BJ, Phillips PE (2012) Hierarchical recruitment of phasic dopamine
signaling in the striatum during the progression of cocaine use. Proc Natl Acad Sci U S A
109:20703-20708.

Wu Q, Reith ME, Kuhar MJ, Carroll FI, Garris PA (2001) Preferential increases in nucleus accumbens
dopamine after systemic cocaine administration are caused by unique characteristics of
dopamine neurotransmission. J Neurosci 21:6338-6347.

Yin HH, Knowlton BJ (2004) Contributions of striatal subregions to place and response learning. Learn
Mem 11:459-463.

Yorgason JT, Espana RA, Jones SR (2011) Demon voltammetry and analysis software: analysis of cocaine-
induced alterations in dopamine signaling using multiple kinetic measures. J Neurosci Methods
202:158-164.

Zhai S, Tanimura A, Graves SM, Shen W, Surmeier DJ (2018) Striatal synapses, circuits, and Parkinson's
disease. Curr Opin Neurobiol 48:9-16.

Zhang S, Qi J, Li X, Wang HL, Britt JP, Hoffman AF, Bonci A, Lupica CR, Morales M (2015) Dopaminergic
and glutamatergic microdomains in a subset of rodent mesoaccumbens axons. Nat Neurosci
18:386-392.

Zhou FM, Liang Y, Dani JA (2001) Endogenous nicotinic cholinergic activity regulates dopamine release in
the striatum. Nat Neurosci 4:1224-1229.

1031

Figure Legends

Figure1. Simultaneous comparison of dLight and FSCV dopamine responses in dorsal striatal slices. **(a)** Schematic diagram of simultaneous dLight photometric and FSCV recordings. **(b)** dLight signal is completely blocked by application of the D1 dopamine receptor antagonist, SCH23390 **(c)** Input-output curves of dLight photometric and FSCV DA measurements (*p<0.05, **p<0.01; ***p<0.005, one sample t-test). **(d)** Schematic diagram illustrating the viral strategy used to ablate substantia nigra DA neurons to confirm that changes in striatal dLight fluorescence are due to dopamine. **(e & f)** Genetic ablation of nigral DA neurons results in markedly reduced dopamine release measured with dLight and FSCV. Representative traces **(g)** and summarized data **(h)** showing that application of the D2 dopamine receptor agonist, quinpirole, inhibits dopamine release equally with both methods. **(i & j)** Increasing extracellular calcium increased dLight fluorescent responses. * p<0.05, ** p<0.01. ***p<0.001 Error bars represent the SEM.

Figure 2. DAT blockers do not increase DA release. **(a)** Representative traces of simultaneously collected dLight (blue shades) and FSCV (orange shades) DA transients before and after application of cocaine. **(b)** Summary data showing that application of cocaine to dorsal striatal slices dose dependently increases peak DA transient peak amplitude measured with FSCV but not dLight photometry methods. **(c)** Summary data showing that DA transient duration is increased with both methods following cocaine application. **(d,e)** Similarly, the specific DAT inhibitor, nomifensine (1µM) increases DA transient peak amplitude with FSCV but not dLight photometry measurements. **(f)** Nomifensine experiment: Amplitude and duration of transients measured with dLight photometry normalized to pre-drug conditions with the unmodified triangle waveform (-0.4V to 1.2V) **(g)** The same normalized dlight photometry readout with nomifensine and modified triangle waveform (-0.4V to 1.0V). **(h)** The normalized FSCV amplitude and duration changes with nomifensine and unmodified triangle waveform (-0.4V to 1.2). **(i)** The normalized FSCV amplitude and duration changes with nomifensine and modified triangle waveform (-0.4V to 1.0V) **(j,k)** Cocaine does not increase DA transient peak height in DAT KO mice in photometry (j) and FSCV measurement (k). **(l,m)** Schematic diagram of the model for DAT inhibitor-induced increases in dopamine transient peak height measured using FSCV without DAT inhibitor(l) and with DAT inhbitor(m). Error bars represent the SEM.

Figure 3. Dorsal striatum expression of the dLight sensor on plasma membrane of dendrites synapsing with TH-axons. Immunofluorescence detection of dLight sensor with an antibody targeted to the GFP moiety contained in the sensor (green) and tyrosine hydroxylase (TH, red) in the dorsal striatum from a mouse injected with a viral vector encoding dLight sensor. **(a)** Apposition of a dLight-positive dendrite and TH-positive axon is seen at high magnification (right panels). **(b)** Electron micrographs of immunogold detection of dLight sensor (gold particles, green arrowheads) on the plasma membrane of a dendrite (green outline in b). Note the apposition of dendritic dLigh and a presynaptic axon (blue arrow in b). **(c)** dLight expression on the plasma membrane of a soma (green outline). Note also the dLight sensor (gold particles) in association with Golgi apparatus and endoplasmic reticulum (ER). **(d)** Detection of dLight sensor (scattered dark material, green arrowheads) in a dendrite (green outline) establishing synapses (blue arrows) with a TH-positive axon (gold particles, red arrowheads).

Figure 4. *In vivo* DA measurement in DLS following cocaine administration. **(a)** Schematic
diagram of *in vivo* fiber photometry system. **(b)** Sample fluorescence (dF/F_0, % baseline) profile
of DA activity in DLS before and after i.p. cocaine injection. **(c)** Average fluorescence (dF/ F_0,
normalized to session maximum) profile of mice (n=4) before and after i.p. cocaine injection (left
panel), and comparison of the average dF/ F_0 between 10min of baseline and 10 min of cocaine
injection (right panel, two-tailed paired t-test ** p=0.0050) **(d)** Magnified dF/ F_0 profile illustrating
DA activity before (Baseline) and after i.p. cocaine injection. **(e)** Averaged fluorescence (dF/ F_0,
normalized to session maximum) profile of baseline DA transients, time-locked to the transient
peaks (n=64). **(f)** Averaged fluorescence (dF/ F_0, normalized to session maximum) profile of DA
transients following cocaine injection, time-locked to the transient peaks (n=165). DA transient
frequency **(g)** and amplitude **(h)** were increased following cocaine injection. Each point denotes
a five minute average. **(i)** Average exponential fit of normalized dopamine transient decay. **(j)**
Fitted exponential decay constant values showing longer DA transient decay time following
cocaine injection; individual points correspond to individual transients. **(k)** dLight fluorescence is
blocked by *in vivo* administration of the D1R antagonist, SCH23390. **(l)** Representative dF/ F_0
profiles for dLight in DLS before and after SCH23390 administration as well as for an eGFP control
mouse. *** p<0.001. Shaded areas and error bars represent the SEM.

1092

Figure 5. *In vivo* DA dynamics across striatal regions during Pavlovian training. **(a)** Schematic
diagram of the experimental design and **(b,c)** behavioral learning curve during Pavlovian training
(n=20 mice, two-tailed paired t-test, p<0.0001). **(d)** Head-entry probability in the early (days 1-3)
and late training sessions (days 12-14). **(e)** Monitoring sites in the striatum and example traces of
dF/ F_0 session average profiles in NAc, DMS, and DLS. **(f,g,h)** Averaged decay constant fitting of
normalized DA transients aligned to the onset of CS+. **(i)** Summarized decay constants across
striatal regions. Each point is an average of 15 CS+ trials per animal (one-way ANOVA with
Bonferroni's correction for multiple comparisons, * p=0.01, *** p=0.0005). **(j,k,l)** DA responses to
CS+ (open circles) and reward-delivery (filled squares) during the Pavlovian training in NAc (left,
blue, n=6 mice two-way ANOVA time x agent interaction: p<0.0001 ****), DMS (mid, green, n=7
mice two-way ANOVA time x agent interaction: p<0.0006 ***), and DLS (right, red, n=7 mice, two-
way ANOVA time x agent interaction: p<0.0001 ****), respectively. **(m)** DA responses coupled to
reward delivery (two-way ANOVA main effect :   training : n.s   region : p=0.0202). **(n)** DA
responses coupled to CS+ in the early and late training sessions (two-way ANOVA region effect :
p=0.0202*, training effect : n.s. p=0.1078, region x training interaction: n.s. p=0.0911 F (2, 17) =
2.767) **(o,p,q)**   Across striatal regions, DA responses to CS- presentation (top row), CS+ (mid
row) and unexpected omission (top row) in the late training phase. **(r)** DA responses coupled to
CS+ or CS- presentation (two-way ANOVA, cue effect: p<0.0001**** region effect: p=0.0075 **
cue x region interaction: p=0.0268*). **(s)** DA responses in unexpected-omission test (two-way
ANOVA reward effect :   p<0.0001**** region : ns p=0.6399,   reward x region interaction : ns
p=0.7884). Shaded orange areas indicate CS duration.

Figure 6. Head-entry rate and DA activity patterns in: **(a)** NAc with CS+ and reward (left) CS+ and unexpected omission (mid), and CS- without reward (right); **(b)** DMS with CS+ and reward (left) CS+ and unexpected omission (mid), and CS- without reward (right); and **(c)** DLS with CS+ and reward (left) CS+ and unexpected omission (mid), and CS- without reward (right), and **(d)** Fluorescence signals in GFP control mice with CS+ and reward (left) CS+ and unexpected omission (mid), and CS- without reward (right). Shaded orange areas indicate CS duration. Note the sustained increase in fluorescence between the offset of the transient induced by the CS+ and reward delivery or unexpected omission in the NAc but not DMS or DLS.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6