TITLE

Patterns of Structural Variation Define Prostate Cancer Across Disease States

AUTHORS/AFFILIATIONS

Meng Zhou[1,2,9], Minjeong Ko[4,9], Anna C. Hoge[4], Kelsey Luu[4], Yuzhen Liu[4], Magdalena L. Russell[4], William W. Hannon[4], Zhenwei Zhang[1,5], Jian Carrot-Zhang[1,2,3], Rameen Beroukhim[1,2], Eliezer M. Van Allen[1,2,6], Atish D. Choudhury[1,3], Peter S. Nelson[4,7], Matthew L. Freedman[1,3,8], Mary-Ellen Taplin[1,3,*], Matthew Meyerson[1,2,3,*], Srinivas R. Viswanathan[1,2,3,*], Gavin Ha[4,7,10,*]

[1] Department of Medical Oncology, Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, MA 02215
[2] Broad Institute of MIT and Harvard, 415 Main Street, Cambridge, MA 02142
[3] Harvard Medical School, 25 Shattuck St, Boston, MA 02115
[4] Public Health Sciences and Human Biology Divisions, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave. N, Seattle, WA 98109
[5] Department of Pathology, UMass Memorial Medical Center, 1 Innovation Dr. #2, Worcester, MA 01605
[6] Center for Cancer Genomics, Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, MA 02215
[7] Department of Genome Sciences, University of Washington, 1959 Pacific St, Seattle, WA, 98195
[8] Center for Functional Cancer Epigenetics, Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, MA 02215
[9] These authors contributed equally
[10] Lead Contact
* Correspondence: mtaplin@partners.org (M-E.T.), matthew_meyerson@dfci.harvard.edu (M.M.), srinivas_viswanathan@dfci.harvard.edu (S.R.V.), gha@fredhutch.org (G.H.)

KEYWORDS

Prostate cancer; castration-resistant prostate cancer; androgen receptor; whole genome sequencing; structural variant; enhancer

SUMMARY

The complex genomic landscape of prostate cancer evolves across disease states under therapeutic pressure directed toward inhibiting androgen receptor (*AR*) signaling. While significantly altered genes in prostate cancer have been extensively defined, there have been fewer systematic analyses of how structural variation reflects the genomic landscape of this disease. We comprehensively characterized structural alterations across 278 localized and 143 metastatic prostate cancers profiled by whole genome and transcriptome sequencing. We observed distinct significantly recurrent breakpoints in localized and metastatic castration-resistant prostate cancers (mCRPC), with pervasive alterations in noncoding regions flanking the *AR, MYC*, *FOXA1*, and *LSAMP* genes in mCRPC. We defined nine subclasses of mCRPC based on signatures of structural variation, each associated with distinct genetic features and clinical outcomes. Our results comprehensively define patterns of structural variation in prostate cancer and identify clinically actionable subgroups based on whole genome profiling.

INTRODUCTION

Over the past decade, genomic sequencing studies have progressively sharpened our view of the genetic landscape of prostate cancer (Leinonen et al., 2011). Such studies have defined key driver genes in prostate cancer and have enabled the deployment of therapeutic agents in molecularly-defined disease subsets, including potent androgen receptor (*AR*)-targeted therapies (de Bono et al., 2011; Scher et al., 2012), poly (ADP-ribose) polymerase (PARP) inhibitors in *BRCA1/2*-altered prostate cancers, and immune checkpoint inhibitors in cancers with microsatellite instability (Abida et al., 2019, 2020; de Bono et al., 2020; Pritchard et al., 2016).

To date, most cancer genomic studies have employed whole exome sequencing (WES) and have thus been focused on mutations or copy number alterations that occur within the protein-coding regions of genes, which represent only 1-2% of the genome. More recent studies applying whole genome sequencing (WGS) to prostate and other cancers have identified previously underappreciated recurrent alterations in regulatory (non-coding) regions of the genome and have illuminated complex mechanisms of genomic alterations – driven by structural variants (SVs) – that are difficult to discern by WES (Baca et al., 2013; Campbell et al., 2020; van Dessel et al., 2019; Fraser et al., 2021; Glodzik et al., 2017; Hadi et al., 2020; Nik-Zainal et al., 2016; Quigley et al., 2018; Stephens et al., 2011; Viswanathan et al., 2018; Weinhold et al., 2014). These studies highlight the need for continued high-resolution genomic discovery efforts in prostate cancer.

In addition to efforts characterizing entire cancer genomes, recent studies have illustrated the importance of molecularly profiling prostate cancer across disease states. While many localized prostate cancers can be cured with surgery or radiotherapy, a substantial portion of higher-risk cancers recur and progress to metastatic disease, which is incurable. Recurrent prostate cancer may have a long natural history, during which time a patient may receive several lines of therapy – with androgen deprivation therapy (ADT) as a backbone – that may shape the cancer's genomic landscape (Mateo et al., 2020).

Indeed, while hormone-refractory castration-resistant prostate cancer (CRPC) has been less extensively profiled than localized prostate cancer, several studies have indicated that CRPCs display genomic landscapes distinct from treatment-naïve disease (Armenia et al., 2018; Grasso et al., 2012). A cardinal hallmark of CRPC is the reactivation of *AR* signaling in the face of maximal ADT (Chen et al., 2004; Yuan et al., 2014). This may occur via diverse mechanisms, including the production of constitutively active *AR* splice variants (*AR-V*s) and activating mutations or copy number amplifications of the *AR* gene (Brand and Dehm, 2013; Céraline et al., 2004; Henzler et al., 2016) or of regulatory elements distal to the gene body (Quigley et al., 2018; Takeda et al., 2018; Viswanathan et al., 2018). To date, the relative contribution of each of these mechanisms in driving *AR* reactivation in CRPC has not been systematically explored. Also needed is a more global map of significant hotspots of structural variation in prostate cancer genomes, drawn within a rigorous statistical framework.

In this study, we performed linked-read WGS on 36 mCRPC tumor-normal pairs. We combined these data with WGS and whole transcriptome sequencing (RNA-Seq) data from previously described localized and metastatic CRPC cohorts (Campbell et al., 2020; Li et al., 2020; Quigley et al., 2018; Viswanathan et al., 2018). We then established a harmonized workflow for the integrative genomic analysis of 278 localized and 143 metastatic CRPC samples, interrogated both hotspots and genome-wide patterns of structural variation, and evaluated their consequences.

RESULTS

WGS analysis of localized and metastatic prostate cancer cohorts

We performed linked-read whole genome sequencing on 36 biopsy specimens from 33 mCRPC patients and matched blood normal controls. After quality control, 17 samples were excluded based on insufficient tumor purity and contamination (**Methods, Figure 1A, Table S1**). We re-analyzed a linked-read WGS dataset of 23 samples published previously (Viswanathan et al., 2018), resulting in a total of 42 linked-read WGS samples from 38 patients with mean coverage of 34X (range 21X - 54X) and 33X (range 25X - 45X) for tumor and normal samples, respectively (**Table S1A**). The mean molecule length was 29 kB and 34 kB in tumor and normal samples, respectively (**Table S1A**).

We further combined these data with 101 mCRPC samples sequenced with standard short-read sequencing, published previously (Quigley et al., 2018). This resulted in the generation of a final combined cohort of 143 tumor-normal pairs (**Figure 1A**). Fifty-four samples (37.8% of 143 samples) were collected at castration resistance, prior to receiving treatment of second-generation androgen receptor signaling inhibitor (ARSi) such as abiraterone and/or enzalutamide ("pre-treatment"), while the remaining 89 samples (62.2% of 143 samples) were collected at progression ("post-treatment", **Figure 1B**, **Table S1B**). We analyzed the somatic single nucleotide variant (SNVs), insertion-deletions (indels), copy number alterations (CNAs), and SVs in the combined cohort and identified recurrent somatic alterations in each of these classes (**Figure 1A, Methods**).

A total of 2,315,452 SNVs and indels were called, with a mean tumor mutation burden (TMB) of 2.82 mutations per million bases (Mb). We confirmed that known driver genes of prostate cancer were enriched for non-synonymous mutations, including *TP53*, *RB1*, *PTEN*, *FOXA1*, *CDK12*, *AR* and *SPOP* among known COSMIC Cancer Gene Census genes (dndscv, q ≤ 0.1, **Table S1C and S1D, Methods**). We detected an average of 272 (range 96-833) SV events per sample. Based on breakpoint orientations, SV events were classified into deletions, inversions, tandem duplications, inter-chromosomal translocations, and intra-chromosomal translocations, while intra-chromosomal translocations were further divided into balanced and unbalanced events based on copy number information (**Methods**). Chromoplexy was detected in 53 samples (37.1% of 143 samples) while chromothripsis was detected in 37 samples (25.9%); these events were not mutually exclusive (Fisher's exact test, log-odds=1.417, p-value=0.612). Ten cases (7.0%) harbored a genome-wide tandem duplicator phenotype (TDP), all of which had *CDK12* inactivating alterations, as recently reported (Viswanathan et al., 2018; Wu et al., 2018). We found that TDP was mutually exclusive with ETS rearrangements (Fisher's exact test, log-odds ratio=0.133, p=0.043) and chromothripsis (log-odds ratio=0.301, p-value=0.007), as previously reported (van Dessel et al., 2019; Quigley et al., 2018; Viswanathan et al., 2018; Wu et al., 2018).

Analysis of CNA events across the genome revealed amplification and deletion peaks in the regions of known prostate cancer genes (Armenia et al., 2018; van Dessel et al., 2019; Quigley et al., 2018; Viswanathan et al., 2018). Many oncogenic drivers of mCRPC, such as *AR* and *MYC*, are within peaks of amplification across the cohort, while tumor suppressors such as *PTEN, TP53,* and *KMT2C* were found within deletion peaks (**Figure S1C**, **Table S1E and S1F**).

Recurrent somatic structural variants in prostate cancer-associated genes

Structural variants may either activate or inactivate gene function, depending on the location of the breakpoints and the specific class of SV. We analyzed the impact of SVs across our cohort, distinguishing between those with predicted inactivating ("gene transecting events") and activating ("gene flanking events") effects (**Figure 1C, Figure S1C, Table S1G and S1H**).

Frequent gene transecting alterations were observed at the *TTC28* (37.1% of 143 samples),
LSAMP (31.5%), and *PTPRD* (23.8%) loci, which have not been extensively studied in prostate
cancer. Rearrangements involving *TTC28* were predominantly inter-chromosomal translocations
between the gene body and various non-recurrent partner loci (**Figure S2B**). This likely
represents retrotransposon activity, given that the *TTC28* locus harbors an active L1
retrotransposon element (Pitkänen et al., 2014; Pradhan et al., 2017; Tubio et al., 2014).
Transecting SVs within the *LSAMP* and *PTPRD* genes were predominantly deletions. Both of
these genes are sites of deletion/rearrangement in cancer and have been reported to function as
tumor suppressors, though they have not been extensively studied within the context of prostate
cancer (Chen et al., 2003; Kresse et al., 2009; Kühn et al., 2012; Veeriah et al., 2009) (**Figure
1C**). Of note, although gene transecting events would be predicted to disrupt individual genes,
the most frequent transecting events identified via this analysis were deletion events that span
the adjacent *TMPRSS2* and *ERG* genes (observed in 37.8%), which actually produces an
activating*TMPRSS2-ERG* fusion.

Duplication events that flank an intact gene could activate oncogenes, either by resulting in copy
number gain of the gene or by duplicating non-coding regulatory regions (Quigley et al., 2018;
Viswanathan et al., 2018). Indeed, we observed recurrent tandem duplication events with
breakpoints located in the flanking gene regions of several known prostate cancer oncogenes,
including *AR* (35.7%), *FOXA1* (16.8%), *MYC* (16.8%), and *CCND1* (14.0%) (**Figure 1C**).

Certain prostate cancer driver genes were altered by multiple classes of structural alterations in
both the gene body and flanking regions (e.g., *AR*, *PTEN*), while others were predominantly
altered by a single alteration class (e.g., SNVs for *TP53*, intragenic translocations for *TTC28*, or
flanking tandem duplications for *MYC*) (**Figure 1C, Figure S1C**). Collectively, these results
demonstrate that prostate cancer is associated with diverse classes of rearrangements, both
within genes and in intergenic regions.

Significantly recurrent breakpoint regions in the mCRPC genome are enriched within enhancer regions and *AR* binding sites

Next, we sought to identify significantly recurrent breakpoint (SRB) regions across our combined
mCRPC cohort of 143 cases in a genome-wide, unbiased manner. We applied a Gamma-Poisson
regression approach to model the occurrences of SV breakpoints within 100 kB windows across
the cohort as previously described (Imielinski et al., 2017). Importantly, this model nominates
significantly recurrent breakpoint regions likely to function as cancer drivers by accounting for six
different covariates, including sequence features (e.g., GC-content and transposable elements),
fragile sites, heterochromatin regions, DNase I hypersensitivity sites (DHS), and replication timing
(**Methods**).

We identified a total of 55 significantly recurrent breakpoint regions genome-wide across our
combined mCRPC cohort (Benjamini-Hochberg corrected, q-value ≤ 0.1, **Figure 2A**, **Table S2A**).
Thirty-six (65.5%) SRB regions were located within 1 Mb of 14 known prostate cancer driver
genes, including *AR* and its enhancer, *TMPRSS2/ERG*, *TP53*, *PTEN*, *FOXA1*, and *MYC*. For
these 14 driver genes, we did not observe significant differences in SV alteration frequencies
when comparing between pre-treatment (N=54) and post-progression (N=89) samples, except in
the case of *ERG, for* which the SV frequency was enriched in pre-treatment samples (Fisher's
exact test, p = 0.0395; all other genes had p > 0.05, **Figure S3B**). We also did not identify any
major differences in the alteration frequencies of prostate cancer genes in four patients who had
paired samples collected both before treatment with and after progression on an ARSi. (**Figure
S3A**).

We then sought to compare how SVs drive prostate cancer across disease states. For the
localized disease state, we utilized genome alteration calls from 278 primary localized prostate
cancer tumors from the PCAWG study (Campbell et al., 2020; Li et al., 2020). Using Gamma-
Poisson regression, we first identified 47 SRB regions in localized prostate cancer tumors (**Figure
S2A, Table S2B**). Six prostate cancer genes (*TMPRSS2*, *ERG*, *TP53*, *PTEN*, *IL6ST*, *ELK4*) within
mCRPC SRB regions were also found within or in proximity (less than 1 Mb) to an SRB region in
localized disease. By contrast, four SRBs (three near *SEL1L3* and one near *PRKDC*) were unique
to localized disease, while 27 SRBs were unique to mCRPC with six genes nearby (*LSAMP*,
ETV1, *MYC*, *PTPRD*, *FOXA1*, *AR*). When comparing SV alteration frequencies for the 14 genes
located within SRB regions in either mCRPC or localized tumors, 12 genes were significantly
more altered in mCRPC samples, while *TMPRSS2* and *ERG* were significantly more altered in
localized disease (Fisher's exact test, $p < 0.05$ for all genes, **Figure 2B**). Thus, localized prostate
cancer and mCRPC have significantly different landscapes of recurrent SVs.

To explore the potential functional consequences of SVs in intergenic SRB regions, we
overlapped SV breakpoints with locations of H3K27ac marks specific to mCRPC (Pomerantz et
al., 2020). We observed that intergenic SVs within SRB regions in the mCRPC cohort included
gene flanking events that were enriched at putative enhancer regions for *AR*, *MYC*, and *FOXA1*,
which all had frequent focal duplication events at sites marked by mCRPC-specific H3K27ac
deposition (**Figure 2C**). Interestingly, an intragenic deletion SRB region was observed near the
transcription start site of *LSAMP*, also overlapping H3K27ac marks. *PTEN* had a high level of
both gene transecting and flanking deletions, leading to SV breakpoints that were spread more
broadly around the gene.

We also observed an enrichment of metastatic-specific *AR* binding sites (ARBS) compared to
localized primary ARBS within the 55 mCRPC SRB regions (**Figure 2D**, one-sided proportion test,
$p = 1.05 \times 10^{-8}$). This enrichment was not observed for localized primary SRB regions ($p = 0.22$).
These results highlight that SVs within mCRPC SRB regions may be capturing the genome-wide
footprint of activated *AR* signaling that occurs with castration resistance.

Refined landscape of ETS gene fusions from integrated analysis of the genome and transcriptome

We applied gene fusion analysis by integrating both genome rearrangements and fusion RNA
transcript information from 127 samples with RNA-seq data (**Figure 1A**, **Table S2C**, **Methods**).
For gene fusions involving E26 transformation-specific (ETS) transcription factor gene family
members (*ERG*, *ETV1*, *ETV4* and *ETV5*), we detected 50 events supported by both DNA and
RNA evidence, 15 supported by only DNA evidence, and 10 supported by only RNA evidence
(**Figure 2E**, **Figure S2D**). Overall, 74 samples (51.7% of 143 samples) harbored a fusion event
of the *ETS* gene family, consistent with previous reports (Tomlins et al., 2005, 2007) (**Figure 1B,
Table S2C**).

Among the ETS fusions, *ERG* was most commonly involved with *TMPRSS2* as the fusion partner
(54 out of 57 cases, **Figure 2G**). Other common ETS fusion partners were *SLC45A3* (7 cases)
and lncRNA RP11-356O9.1 downstream of *FOXA1* (3 cases). *ETV1* had eight distinct fusion
partners, which is consistent with previous reports that *ETV1* is a promiscuous ETS fusion
member (Kumar-Sinha et al., 2015) (**Figure 2F**).

We observed that fusions of the ETS family members *ERG*, *ETV1*, *ETV4* and *ETV5* were mutually
exclusive, except for one sample which harbored fusions of both *ERG* and *ETV1* (**Figure S2D**).
In addition, gene fusion events were correlated with higher expression of the corresponding ETS
genes they involved (Wilcoxon rank-sum tests, $p < 0.05$ for all genes, **Figure 2E**). In the 38 cases
which did not show any evidence for an ETS fusion, we noted that presence of high-level

expression (z-score > 1) of ETS genes *ERG*, *ETV1*, *ETV4*, and *ETV5* were also mutually
exclusive (Fisher's exact test, p = 0.480 for *ETV4*, p = 0.363 for *ETV5*, **Figure S2D**). These may
represent cases of missed fusion calls, or cases in which ETS family members are
transcriptionally activated through non-genetic mechanisms.

Interestingly, we also observed 20 cases (14.0% of 143 cases) involving fusions between the ETS
family member *ELK4* and its upstream gene *SLC45A3*. While the *ELK4* locus was an SRB in our
analysis (**Figure 2A** and **Figure S2B**), manual inspection of individual samples revealed evidence
for a genomic event capable of producing an *ELK4* fusion in only 1 out of 20 cases (**Figure S2D**
and data not shown). In contrast, 19 other cases showed *ELK4* fusions on RNA-sequencing
alone, consistent with a mechanism of cis-splicing or transcriptional read-through events that may
perhaps be induced by local genomic alterations (Qin et al., 2017; Rickman et al., 2009; Zhang
et al., 2012) (**Table S2C**). Importantly, although *ELK4* fusions were significantly correlated with
higher expression of *ELK4* (Wilcoxon rank-sum test, p = 7.91×10^{-5}, **Figure S2D**), these events
were not mutually exclusive with fusions of other ETS family members (Fisher's exact test, p =
0.472). Thus, the functional consequences of these *ELK4* fusions and whether they contribute to
prostate cancer pathogenesis in a manner similar to other ETS fusions remains to be determined.

Diverse and complex rearrangements driving *AR* signaling in mCRPC

Genomic alterations involving the *AR* locus play an important role in sustaining *AR* signaling in
mCPRC (Chen et al., 2004; Quigley et al., 2018; Visakorpi et al., 1995; Viswanathan et al., 2018).
However, the complete spectrum of diverse structural mechanisms that underlie *AR* activation in
mCRPC has not been fully characterized. To understand the relationship between different modes
of somatic *AR* activation, we determined copy number at the *AR* gene body and its upstream
enhancer and categorized samples into distinct groups of: **(1)** co-amplification (N = 99, 69.2% of
143 cases); **(2)** selective *AR* gene body amplification (N = 4, 2.8% of 143 cases); **(3)** selective *AR*
enhancer gains (N = 17, 11.9% of 143 cases), and **(4)** lack of amplification for both (N = 23, 16.0%
of 143 cases) (**Figure 3A-C, Table S3**). For the 122 samples with expression data available, we
observed that *AR* gene expression was higher in the co-amplification and selective enhancer
categories compared to samples with no amplification, after accounting for tumor purity and ploidy
(ANCOVA/TukeyHSD p-values 5.6×10^{-11} and 4.5×10^{-4}, respectively), but not for selective *AR*
status (ANCOVA p = 0.098) (**Figure 3B, Methods**). Interestingly, we observed that samples with
selective enhancer duplication exhibited similar *AR* expression levels to samples with co-
amplification (ANCOVA, p = 0.31), even though enhancer duplications involved lower-copy gains
(mean 2.73, range 1.97 - 5.02) compared to co-amplified samples (mean 12.87, range 1.55 -
150.57) (**Figure 3A**). This is consistent with previous results (Viswanathan et al., 2018) and
suggests a mechanism whereby *AR* expression levels are increased through even modest
genomic expansion of enhancer elements.

We then systematically and manually curated the diverse mechanisms of rearrangements
activating *AR* signaling by analyzing patterns of SVs at the *AR* locus (**Figure 3C, Table S3,**
Methods). We observed a total of 62 samples (43.4% of 143 samples) with tandem duplication
SV events that spanned the enhancer with breakpoints located within 1 Mb, including 16 cases
(11.2% of 143 samples) with selective enhancer copy number amplification status (**Figure 3D**).
Thirty-two samples (22.4% of 143 samples) harbored intragenic rearrangements within *AR*, which
may have implications for the production of truncated, constitutively-active *AR* splice variants
(Henzler et al., 2016). For example, in case DTB-124-BL, we observed a focal intragenic deletion
spanning exons 4-8 of *AR*, which includes the ligand binding domain, resulting in the expression
of truncated *AR* variants (Kanayama et al., 2021) (**Figure 3E**, **Figure S3C**). Interestingly, in the
21 samples with selective *AR* enhancer or selective *AR* gene body copy number gain, none
harbored intragenic SV events in *AR*.

We also examined the landscape of complex rearrangement mechanisms involving *AR*; these
mechanisms involve multiple SV events and copy number patterns, including chromothripsis,
extrachromosomal DNA (ecDNA), chromoplexy, and breakage-fusion-bridge cycle (BFB)
(**Methods**). Chromothripsis of a region or the entire X chromosome involving the *AR* locus was
detected in 5 samples, all of which had co-amplification of *AR* and enhancer, suggesting that
following repair after catastrophic DNA shattering the *AR* locus was retained or further amplified
(**Figure 3F, Figure 3G**). Thirteen samples (9.1% of 143 samples) showed very high levels of *AR*
and enhancer copy number, suggesting the possibility of their presence on extrachromosomal
elements (ecDNA, **Figure 3H**). In 40 samples (28.0% of 143 samples), the most frequent complex
rearrangement mechanism, BFB, led to *AR* locus amplification, including instances following
chromothripsis (Stephens et al., 2011; Umbreit et al., 2020) (**Figure 3G**). Overall, we noted that
complex rearrangement events, which frequently co-occurred, were significantly enriched in
samples with co-amplification of *AR* and enhancer compared to those with selective enhancer
copy number gain status (Fisher's exact test, p = 1.52×10^{-4}).

Distinct signatures of structural rearrangement patterns in mCRPC

To systematically characterize genome-wide structural rearrangement patterns in mCRPC, we
performed rearrangement signature analysis using SV breakpoint features, non-negative matrix
factorization, and known reference signatures (Degasperi et al., 2020; Nik-Zainal et al., 2016)
(**Methods**). First, we derived signatures *de novo*, which identified eight signatures: six that
matched reference signatures (RefSigs) also observed in localized prostate cancer (> 0.91 cosine
similarity), one that matched an ovarian cancer RefSig.R14 associated with large segment (100
kB-10 Mb) TDP (0.96 cosine similarity), and one that was likely an artifact specific to linked-read
sequencing (**Figure S4A-C, Table S4A and 4B**). Therefore, we excluded the linked-read data
and focused on standard WGS data from 101 mCRPC cases for further SV signature analysis.
We fit these samples to the nine known RefSigs from localized prostate cancer (R1-4, R6a-b, R8-
9, R15) and the one (R14) from ovarian cancer (**Figure S4A**, **Table S4C**). Overall, eight of the
RefSigs were detected across our cohort (R1-2, R4, R6a-b, R9, R14-15). Notably absent in
mCRPC were RefSig.R8 (short, 1-10 kB inversions) and RefSig.R3, which is associated with
germline *BRCA1* mutations and short (1-100 kB) tandem duplications (Degasperi et al., 2020;
Glodzik et al., 2017; Nik-Zainal et al., 2016; Willis et al., 2017) (**Figure S4D**). By contrast, we
observed increased prevalence of some signatures in mCRPC compared to localized disease,
including RefSig.R2 (large SV classes, abundant translocations; 97% vs. 60%), RefSig.R4
(clustered translocation events; 37% vs. 27%), and RefSig.R15 (large deletions and inversions,
48% vs. 37%) (**Figure S4D**).

To investigate whether molecular subtypes in mCRPC can be grouped based on SV patterns, we
applied hierarchical clustering on the exposure of the eight fitted signatures and identified nine
distinct SV clusters (**Figure 4, Table S4C**). We observed that samples in SV Cluster 1 were
composed of non-clustered translocation events and were significantly enriched for the presence
of chromoplexy (χ^2 test, FDR corrected, q = 0.12). SV Cluster 3 was characterized by many short
deletions and was significantly enriched for *BRCA2* mutations (q = 5.01×10^{-4}). SV Cluster 5 was
significantly enriched for *SPOP* mutations (q = 0.02), with no instances of ETS gene family fusion
(q=0.06), consistent with previous reports (Barbieri et al., 2012). SV Cluster 6 had the highest
prevalence of *TP53* mutation (q = 0.02), while SV Cluster 7 samples harbored the TDP associated
with *CDK12* inactivation (q = 3.52×10^{-11}) as well as enrichment for *CCND1* gains (q = 0.02),
consistent with previous reports (Nguyen et al., 2020; Wu et al., 2018). The remaining clusters
did not have enrichment for any alterations in known driver genes; however, distinct SV patterns
were still evident in SV Cluster 4 (non-clustered tandem duplications), 8, and 9 (increased
clustered SV events of various classes).

While SV Clusters 3, 5 and 6 had significant enrichment of mutations in *BRCA2*, *SPOP,* and *TP53*,
respectively, not all samples within each cluster harbored these mutations. Intriguingly, we further
noted that clinical outcomes showed significantly better stratification when using SV Clusters 3,
5, and 6 for outcome stratification compared to using the associated mutation status itself (**Figure
S4D-E**). Specifically, SV Cluster 5 had significantly better overall survival than SV Clusters 3 and
6 (log-rank test, p=0.01), while the sample group with *SPOP* mutations did not have significantly
greater survival compared to the sample groups with *BRCA2* and *TP53* mutations (log-rank test,
p=0.45) in this cohort. Together, these results indicate the analysis of genome-wide patterns of
rearrangements may provide a way to further refine molecular subtypes in mCRPC.

DISCUSSION

We present a large-scale and comprehensive integrative genomic analysis of both localized
prostate cancer and mCRPC, with a focus on how structural variation drives each of these
clinically distinct disease states. The size of our cohort as well as our harmonized analysis pipeline
enable a sharper view of the genetic alterations that drive prostate cancer across its natural history
as compared with prior studies, which have involved either smaller cohorts or been limited to a
single disease state (Campbell et al., 2020; Cancer Genome Atlas Research Network, 2015;
Quigley et al., 2018; Viswanathan et al., 2018).

In contrast to somatic SNVs/indels and CNAs that occur within coding regions, the functional and
clinical significance of alterations within noncoding regions has often been more challenging to
interpret, as localized variations in mutability may result in the nomination of certain recurrently
mutated sites that do not necessarily drive cancer (Glodzik et al., 2017; Imielinski et al., 2017;
Nik-Zainal et al., 2016). This issue is even more complex for SVs, in which different classes of
SVs spanning the same loci would be predicted to have distinct functional consequences. Our
study addresses the former issue by identifying genomic hotspots of structural variation with
rigorous correction for covariates including nucleotide composition, replication timing, sensitivity
to DNA breaks, repetitive elements, and chromatin state. We address the latter issue by careful
curation of SV classes to distinguish those that are likely to be activating versus inactivating
(**Figures 1B and 3; Methods**).

Our approach has produced several insights into the recurrent rearrangements that drive prostate
cancer. First, several top hotpots of rearrangement genome-wide lie in noncoding regions outside
the boundaries of known prostate cancer genes. In many cases, such as for *AR*, *MYC*, and
FOXA1, these hotspots overlap with active chromatin marks and likely represent distal regulatory
regions for neighboring prostate cancer genes (**Figure 2**). These data are intriguing in light of the
observation that a majority of prostate cancer germline susceptibility loci are in noncoding regions
(Giambartolomei et al., 2021). Second, the loci altered by rearrangements differ across prostate
cancer disease states (**Figure 2B**). For example, *TMPRSS2-ERG* rearrangements are enriched
in localized prostate cancer versus mCRPC, while alterations in *AR, FOXA1, MYC*, and *LSAMP*
are more frequent in mCRPC than in localized disease. Third, certain driver genes are enriched
for alteration by SVs as compared to other mutagenic processes. For example, *PTEN* inactivation
frequently occurs via gene transecting SV events, while *TP53* inactivation is primarily caused by
SNVs (**Figure 1C and Figure S1**).

Our systematic genomic discovery efforts again highlight the primacy of *AR* as a target of somatic
alteration in hormone-refractory mCRPC. We have precisely catalogued the diverse genomic
mechanisms leading to *AR* activation across our large cohort and find that different alteration
mechanisms are associated with differing levels of *AR* amplification. Whether the precise
mechanism by which *AR* is altered in a given patient is associated with differences in response
to *AR* pathway inhibition warrants further investigation in clinically annotated cohorts. High levels

of *AR* signaling in mCRPC may also underlie the patterns of structural variation seen in this disease state. Strikingly, we found that *AR* binding sites overlapped several of the top SV hotspots in mCRPC (**Figure 2D**), consistent with the notion that androgen signaling may induce DNA double-strand breaks that resolve as rearrangements (Haffner et al., 2010).

In addition to alterations in highly validated prostate cancer genes, we identified highly recurrent rearrangements near or involving genes that have not been extensively studied in prostate cancer, such as *LSAMP, PTPRD*, and *TTC28*. *LSAMP* encodes a cell-surface glycoprotein and has a possible tumor suppressor role in several cancers (Chen et al., 2003; Kresse et al., 2009; Kühn et al., 2012); notably, deletions near the *LSAMP* locus have been shown in one report to be enriched in African American men with prostate cancer (Petrovics et al., 2015). *PTPRD*, a receptor protein tyrosine kinase, has been previously identified as a target of inactivating alteration in glioblastoma (Veeriah et al., 2009). We observed frequent SVs near the *TTC28* locus, which encodes an L1 retrotransposon element, specifically in mCRPC (**Figure 1C**). L1 retrotranspositions originating from *TTC28* have been reported previously in colorectal cancer (Pitkänen et al., 2014; Pradhan et al., 2017; Tubio et al., 2014); our results raise the intriguing possibility that they may also be frequent in prostate cancer, and may be activated by the pressure of hormonal therapy. Interestingly, we also observed SRBs near *ELK4* along with a relatively high frequency of *SLC45A3-ELK4* chimeric transcripts, although it was not clear how the rearrangements at this locus produced the chimeric transcripts in most cases. Whether this fusion functions similarly to or in a distinct mode from other ETS fusions is an exciting area for future study.

Our study also extends beyond the analysis of SVs at individual loci to molecularly subclassify prostate cancers based on their genome-wide signatures of structural variation. Sample clustering based on SV signature exposure defines distinct molecular subtypes of prostate cancer and may find utility alongside signatures of single base substitution and copy number to more precisely define tumor subtypes (Alexandrov et al., 2013, 2020; Degasperi et al., 2020; Macintyre et al., 2018; Wang et al., 2021). In the mCRPC cohort, we identified 9 molecular subtypes based on SV signature, and several clusters had clear associated genomic alterations including chromoplexy (cluster 1), *BRCA2* alterations (cluster 3), *SPOP* alterations (cluster 5), *TP53* alterations (cluster 6) and *CDK12/CCND1* alterations (cluster 7). Future studies with larger WGS cohorts may identify associated alterations in the remaining clusters. Notably, unsupervised clustering identified samples with clear SV signatures but without detectable associated mutations in genes or pathways that plausibly contribute to the genomic alterations (**Figure 4**). Moreover, clinical outcomes were more separated by SV signature cluster than by alterations of the mutations associated with those clusters (**Figure S4D-E**).

In sum, these results highlight the dynamic complexity of rearrangements in prostate cancer across disease states and provide insights into new mechanisms of oncogenesis that can be functionally prioritized in future studies. More broadly, our work underscores the key role of large-scale WGS studies in the derivation of a comprehensive molecular taxonomy of prostate cancer.

ACKNOWLEDGEMENTS

We thank the many patients and their families for their generosity in contributing to this study. We also thank the Prostate Cancer Foundation (PCF) and Stand Up 2 Cancer (SU2C) International Prostate Cancer Dream Team for contributions to specimen acquisition.

This work was supported by the National Institutes of Health (K22 CA237746 to G.H.; P01 CA163227 and R01 CA234715 to P.S.N.; R01 GM107427, R01 CA193910, and R01 CA251555 to M.L.F.; R35 CA197568 to M.M.), Department of Defense Prostate Cancer Research Program (Physician Research Award W81XWH-17-1-0358 to S.R.V.; W81XWH-19-1-0565 and W81XWH-

21-1-0234 to M.L.F.; PC200262 to P.S.N.), PCF Young Investigator Awards (G.H. and S.R.V.),
PCF-Movember Challenge Award (to E.M.V.), Brotman Baty Institute for Precision Medicine (to
G.H.), the Fund for Innovation in Cancer Informatics Major Grant (to G.H.), the V Foundation
Scholar Grant (to G.H.), Wong Family Award in Translational Oncology and Dana-Farber Cancer
Institute Medical Oncology grant (to A.D.C.), H.L. Snyder Medical Research Foundation and the
Cutler Family Fund for Prevention and Early Detection (to M.L.F.), and the Pan-Mass Challenge
team IMAGINE (to M-E.T.), American Cancer Society Research Professor (M.M.).

This research was also supported in part by the NIH/NCI Cancer Center Support Grant P30
CA015704, Pacific Northwest Prostate Cancer SPORE (P50 CA097186), and Scientific
Computing Infrastructure (ORIP Grant S10OD028685).

AUTHOR CONTRIBUTIONS

Conceptualization: M-E.T, M.M., S.R.V., G.H.

Methodology: M.Z., M.K., S.R.V., G.H.

Software: M.Z., M.K., A.C.H., G.H.

Formal Analysis: M.Z., M.K., A.C.H., K.L., Y.L., M.R., W.H., J.C-Z, S.R.V., G.H.

Data Curation: M.Z., M.K., A.C.H., Z.Z., S.R.V., G.H.

Writing – Original Draft: M.Z., M.M., G.H., S.R.V.

Writing – Review & Editing: M.Z., R.B., E.M.V., A.D.C. P.S.N., M.L.F., M-E.T., M.M., G.H., S.R.V.

Visualization: M.Z., M.K., A.C.H., S.R.V., G.H.

Supervision: M-E.T., M.M., S.R.V., G.H.

Funding Acquisition: S.R.V., G.H., M.M.

DECLARATION OF INTERESTS

A.D.C.: Honoraria: OncLive, Bayer, Targeted Oncology, Aptitude Health, Journal of Clinical Pathways, Cancer Network; Consulting: Blackstone; Advisory Board: Clovis, Dendreon, Bayer, Eli Lilly, AstraZeneca, Astellas, Blue Earth; Research Funding: Bayer

E.M.V.: Advisory/Consulting: Tango Therapeutics, Genome Medical, Invitae, Enara Bio, Janssen, Manifold Bio, Monte Rosa; Research support: Novartis, BMS; Equity: Tango Therapeutics, Genome Medical, Syapse, Enara Bio, Manifold Bio, Microsoft, Monte Rosa; Travel reimbursement: Roche/Genentech; Patents: Institutional patents filed on chromatin mutations and immunotherapy response, and methods for clinical interpretation; intermittent legal consulting on patents for Foaley & Hoag

M-E.T.: Advisory boards: Janssen, Pfizer, Astra Zeneca, Bayer

M.L.F.: Served as a consultant to and has equity in Nuscan Diagnostics. This activity is outside of the scope of this manuscript.

M.M.: Consultant for Bayer, Interline and Isobl; an inventor of patents licensed to LabCorp and Bayer; and receives research funding from Bayer, Janssen, and Ono Pharmaceuticals.

P.S.N.: Served as a consultant to Bristol Myers Squibb, Janssen, and Pfizer in work unrelated to the present study.

451   S.R.V.: Consulting (current or previous 3 years), MPM Capital and Vida Ventures; spouse is an
452   employee of and holds equity in Kojin Therapeutics.

453   All other authors declare no competing interests.

 MAIN FIGURE LEGENDS

Figure 1. Study design and genomic landscape of mCRPC.

(A) Workflow of study and data analysis. Tumor specimens (grey) from both primary prostate
cancer and mCRPC were included in this study. Linked-read and short-read whole-genome
sequencing (WGS) and RNA-sequencing datasets were either generated for this study or
reanalyzed from prior studies (Quigley et al., 2018; Viswanathan et al., 2018). A pooled dataset
of 143 mCRPC samples with WGS data was used in this study after curation (**Methods**). Genomic
alteration call-sets for 278 primary localized prostate cancer samples were obtained from
ICGC/TCGA Pan-Cancer Analysis of Whole Genomes (PCAWG) (Campbell et al., 2020; Li et al.,
2020). For 125 mCRPC samples, RNA-seq was used. The overview of the genomic alteration
and characterization analysis is shown.

(B) Clinical annotations and somatic alterations for 143 patient samples in the pooled mCRPC
cohort. Samples are ordered by treatment type; the four patients with pre-treatment and post-
progression pairs are placed at the right. (Top) Clinical and sample information and genomic
pattern classifications. (Middle) Distribution of genomic rearrangement types in individual samples.
(Bottom) Mutational burden for SNVs and indels computed as number of mutations per mega-
base pair (Mb). Y-axis shown in logarithmic scale. Threshold lines indicates mutational burden at
2.5 and 5 mutations per Mb.

(C) Genomic rearrangement alteration profiles of key mCRPC genes. (Top) Events were
categorized into gene transecting and gene flanking events (**Methods**). Gene transecting: if any
of its breakpoints was located within the gene body region. Gene flanking: rearrangements which
were not gene transecting and had breakpoints located within 1 Mb of either transcription start
site or termination site of the gene. Only 159 genes reported and known to be involved in prostate
cancer were considered in this analysis (**Table S1G and S1H**). (Middle) Frequency and
distribution of rearrangement types for gene transecting events; genes with ≥ 10% frequency are
shown. Gene transecting events were prioritized over flanking events during annotation. The
category "Multiple" represents gene-sample pairs carrying more than one type of rearrangement
event. (Bottom) Frequency of gene flanking events by tandem duplication; genes with ≥ 10% are
shown.

Figure 2. Genome-wide analysis of genomic rearrangements in mCRPC.

(A) Analysis of significantly recurrent breakpoint (SRB) identified regions of rearrangement
hotspots, genome-wide, using a Gamma-Poisson regression model. Each dot corresponds to a
100 kB bin (n=26,663 total bins). Statistically significant SRB bins with FDR (Benjamini-Hochberg)
q-value ≤ 0.1 (n=55) are colored based on the distance to the nearest known prostate cancer
driver gene, within 1 Mb. The driver genes within 1 Mb of the SRB bins are labeled. A square
bracket is used for genes spanning multiple bins. Bins with q-value > 0.1 were not significant
(grey).

(B) Comparison of SV alteration frequency in mCRPC versus primary localized prostate cancer.
The union set of genes (n=14) within 1 Mb of SRB hotspot regions in mCRPC and localized
prostate cancer cohorts was included in the comparison. The frequencies represent total gene
transecting and flanking SV events. All labeled genes were significantly enriched in either mCRPC
or primary localized tumors (Fisher's test, p-value < 0.05).

(C) Patterns of rearrangements at the loci of driver genes identified at SRB regions in mCRPC
cohort of 143 tumors. Cumulative counts of intra-chromosomal SV events (tandem duplications
"TandemDup", deletions, and inversions) were computed based on the breakpoints and span of
the events. Histone H3 lysine 27 acetylation (H3K27ac) and *AR* binding sites (ARBS) specific to

mCRPC were obtained from a previous study (Pomerantz et al., 2020). Inter-chromosomal
translocations are not shown. Genome coordinates based on hg38 build.

(D) Overlap of *AR* binding sites (ARBS) within SRB hotspots of mCRPC (55 regions) and primary
localized prostate (47 regions) cohorts. Metastatic-specific and primary localized-specific ARBS
were obtained from previous studies (Pomerantz et al., 2015, 2020). χ^2 test of independence p-
values are shown.

(E) Fusion status and expression of selected genes in ETS transcription factor gene family in the
mCRPC cohort with WGS and RNA-seq data. Fusion type was defined as the data evidence that
supported the event: DNA-only, corresponds to WGS; RNA-only, corresponds to RNA-seq;
DNA+RNA, corresponds to support from both WGS and RNA-seq. Each dot represents a tumor
sample and is colored based on fusion type of each sample; grey indicates no evidence of fusion
event. Data shown for samples with available expression data for the specific ETS gene. Gene
expression values of full-length transcripts are z-score normalized.

(F) Fusion profile of *ETV1*. DNA rearrangement breakpoints supporting the fusion (purple bars)
are indicated with the corresponding fusion partners. Exons of the ETS domain (red) are indicated.
Genome coordinates based on hg38 build.

(G) Summary of fusion partners for selected genes in ETS transcription factor gene family in
mCRPC cohort. Fusion events and partners are indicated by flow connections. Total counts of
individual fusion events and partners across the cohort are shown.

Figure 3. Modes of *AR* activation in mCRPC.

(A) Copy number of *AR* gene and its enhancer (~624 kB upstream) for mCRPC cohort samples
after adjustment by tumor purity and sample ploidy normalization. Data shown for samples with
available *AR* gene expression data. (Left) Copy number of *AR* and its enhancer are shown in $\log_2$
scale, colored based on *AR* gene expression level (transcripts per million, TPM). (Right) Excerpt
of figure highlighting *AR* expression for samples with lower copy number values.

(B) *AR* expression for *AR* locus copy number status for 122 samples with available *AR* gene
expression data. ANCOVA test was performed to account for tumor purity and ploidy as
covariates. TukeyHSD p-values for pair-wise comparisons between groups with *AR* locus
amplification status and groups with no amplification.

(C) Patterns of rearrangements involving the *AR* locus in 143 mCRPC samples. Presence of
specific alteration events and complex rearrangements (black) were predicted automatically and
manually curated. *AR* gene expression shown (blue shades) for same samples in (B); samples
with no available expression data are indicated in grey. Representative examples of each
category are presented in (D) to (H).

(D-H) Complex and simple rearrangement patterns involving the *AR* locus, including focal
duplication events on *AR* enhancer **(D)**, intragenic deletion event leading to loss of ligand binding
domain of *AR* **(E)**, chromosomal level chromothripsis events involving *AR* and enhancer **(F)**, arm-
level chromothripsis coinciding with *AR* amplification by break-fusion-break cycle **(G)**, extra-
chromosomal DNA amplicon including *AR* and enhancer **(H)**. *AR* gene boundary (green) and its
enhancer (yellow) are shown; concave arcs, intra-chromosomal SV events; convex arcs, inter-
chromosomal SV events. Copy number values represent 10 kB bins and have been tumor purity
corrected.

Figure 4. Clustering of mCRPC SV signatures

SV signature analysis and hierarchical clustering identifies nine distinct molecular groups. (Top)
Dendrogram of the clustering of SV signature exposure. The prevalence of each signature was
computed based on having ≥ 0.05 exposure (proportion of SVs). (Middle) Enrichment of altered
prostate cancer drivers. Enriched alterations in Cluster 1, 3, 5, 6, and 7 are shown based on
statistical significance by χ^2 test. (Bottom) Composition of SV types and sizes for each SV cluster,
separated by non-clustered (nc) and clustered (c) SV events.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Gavin Ha (gha@fredhutch.org).

Materials availability

This study did not generate new unique reagents.

Data and code availability

- Whole genome sequencing data have been deposited at dbGaP under accession number phs001577 and access is available upon request.
- All original code has been deposited at GitHub and is publicly available as of the date of publication. Links are provided in the key resources table.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Human subjects

For tumor biopsies profiled via linked-read sequencing, samples were collected from individuals with mCRPC who provided informed consent on institutional IRB-reviewed protocols, as previously described (Viswanathan et al., 2018). Uniformly reanalyzed data were generated as described in the respective studies (Campbell et al., 2020; Quigley et al., 2018).

METHOD DETAILS

Sequence data processing for linked-read genome sequencing data

Data processing of the linked-read genome sequencing data include high molecular weight DNA preparation and sequencing library construction followed protocols as previously described (Viswanathan et al., 2018). DNA was extracted from tumor samples using the MagAttract HMW DNA Kit (QIAGEN), and then quantified using Quant-it Picogreen assay kit (Thermo Fisher) on a Varioskan Flash Microplate Reader (Thermo Fisher). For germline samples, pre-extracted DNA was size-selected on the PippinHT platform (Sage Science) and then quantified using the Quant-it Picogreen assay kit (Thermo Fisher) on a Varioskan Flash Microplate Reader (Thermo Fisher). Libraries were constructed using the 10X Chromium protocol (10X Genomics), with the fragment sizes determined using the DNA 1000 Kit and 2100 BioAnalyzer (Agilent Technologies) and quantified using qPCR (KAPA Library Quantification Kit, Kapa Biosystems). WGS libraries were sequenced using the Illumina HiSeqX platform. The Long Ranger v2.2.2 pipeline (10X Genomics) was used for aligning sequence reads to the human genome hg38 (GRCh38).

Samples were excluded from the analysis based on having tumor purity less than 15% estimated by TitanCNA or based on cross-individual contamination indicated by SNP fingerprinting. A total of 17 samples with linked-read data was excluded (**Table S1J**).

QUANTIFICATION AND STATISTICAL ANALYSIS

List of known prostate cancer driver genes

For analyses limited to established prostate cancer driver genes, a curated list of 159 known prostate cancer driver genes was assembled from several prior studies (Armenia et al., 2018; van Dessel et al., 2019; Quigley et al., 2018; Viswanathan et al., 2018). The list of genes are provided in **Table S1**.

Somatic mutation analysis

Somatic mutation detection

Somatic mutation calls for samples based on linked-read sequencing were generated by Mutect2 from the Genome Analysis Toolkit (GATK) (Van der Auwera and O'Connor, 2020). Default parameters were used on individual pairs of tumor and normal samples following the standard GATK pipeline. A panel of normals based on all normal samples was used to filter out germline variants. The SNV calls were further processed using the modified version of LoLoPicker (Carrot-Zhang and Majewski, 2017) as described previously (Viswanathan et al., 2018). The panel of normals for LoLoPicker was generated from 52 normal samples based on linked-read sequencing. The final SNV call set was composed of the common variants called by both Mutect2 and LoLoPicker. Somatic indels for linked-read samples were called by Strelka (Saunders et al., 2012). All parameters were default except the following modifications: sindelNoise = 0.000001, minTier1Mapq = 20. Somatic mutation calls for the 101 WGS samples based on short-read sequencing including SNV and indels based on Strelka were obtained from a prior study (Quigley et al., 2018). All variants were further annotated using annovar with "table_annovar.pl" to functionally annotate genetic variants. The paramter -neargene was set to 5000 to define the promoter region as 5 kB upstream of the transcription start site of a protein coding gene.

Analysis of significantly mutated genes

R package dndscv (Martincorena et al., 2017) was used to identify significantly mutated genes. For driver discovery on GRCh38, a precomputed database corresponding to human genome GRCh38.p12 was downloaded and used as the reference database. A global q-value ≤ 0.1 was applied to identify statistically significant (novel) driver genes. To reduce false positives and increase the signal to noise ratio, we only considered mutations in Cancer Gene Census genes (v81) (Tate et al., 2019).

Copy-number analysis of linked-read WGS and short-read WGS data

Copy-number calls

The ploidy and purity corrected copy-number of all mCRPC samples in this study was analyzed by TitanCNA (Ha et al., 2014) and ichorCNA (Adalsteinsson et al., 2017), with different pipeline settings. For WCDT samples, the snakemake workflow for Illumina sequencing was applied with the following parameters modified: ichorCNA_normal: c(0.25, 0.5, 0.75); ichorCNA_ploidy: c(2,3,4); ichorCNA_includeHOMD: TRUE; ichorCNA_minMapScore: 0.75; ichorCNA_maxFracGenomeSubclone: 0.5; ichorCNA_maxFracCNASubclone: 0.7; TitanCNA_maxNumClonalClusters: 3; TitanCNA_maxPloidy: 4. The workflow is available at https://github.com/GavinHaLab/TitanCNA_SV_WGS.

For linked-read data samples, a Snakemake workflow for 10X Genomics whole genome sequencing data was used with the following parameters modified: TitanCNA_maxNumClonalClusters: 3; TitanCNA_maxPloidy: 4. TitanCNA solutions were generated for number of clonal clusters from 1 to 3 and ploidy initializations from 2 to 4. Optimal

630 solutions were selected as described, with manual inspection to confirm tumor ploidy and clonal
631 cluster selection (Viswanathan et al., 2018); solutions are provided in **Table S1J**. The workflow
632 can be accessed at https://github.com/GavinHaLab/TitanCNA_10X_snakemake. The final copy-
633 number call-set is included in **Table S1I**.

Recurrent somatic copy-number alteration

GISTIC 2.0 was used to detect regions with recurrent CNA in mCRPC samples. For input, all copy
numbers (logR_Copy_Number from TITAN output) were converted to log2 copy ratio using the
median logR copy number from genome-wide (separately for autosomes and X chromosome) as
denominator. We set corrected logR copy number to -1.5 for segments where corrected log R
copy number below -1.5 and set values to 0 if copy neutral. GISTIC2.0 was run with the following
parameters: td 0.5; ta 0.1; genegistic 0; maxseg 5000; js 4; cap 1.5; broad 1; brlen 0.75; conf 0.99;
qvt 0.25; armpeel 1; rx 0; gcm mean; do_gene_gistic 1; savegene 1; scent median. Wide peaks
detected by GISTIC2 were re-annotated based on overlapping genomic coordinates, using
prostate cancer driver genes.

Structural variant analysis

Structural variant detection in linked-read and short-read whole genome sequencing data

For each tumor-normal pair of samples with linked-read genome sequencing data, three variant
callers were used to detect structural variants: SvABA (Wala et al., 2018), GROC-SVS (Spies et
al., 2017), Long Ranger version 2.2.2 (https://support.10xgenomics.com/genome-
exome/software/pipelines/latest/using/wgs).

The SvABA analysis was performed using default tumor-normal paired settings. Re-analysis of
low confidence (based on evidence from discordant and split reads) events filtered by SvABA was
performed to 'rescue' SVs using linked-read barcode overlap between pairs of breakpoints within
a given SV event, as previously described (Viswanathan et al., 2018). Only SV events having
span of 1.5 times the mean molecule length in the library were considered for rescue. We further
rescued low confidence intra-chromosomal SV events with span > 50 kB filtered by SvABA if at
least one of the breakpoint pair was within 100 kB of a CNA boundary or (2) if both breakpoints
were each within 1 Mb of the boundaries for the overlapping CNA event and the length of the SV
overlaps this CNA event by > 75%. Inter-chromosomal translocation SV events filtered by SvABA
are rescued if both breakpoints were within 100 kB of CNA boundaries.

GROC-SVS analysis was performed using two-sample (tumor-normal paired) mode or three-
sample (pre-treatment, post-progression, normal) mode when applicable. SV events were
retained if all following conditions were satisfied: (1) $p < 1 \times 10^{-10}$, (2) minimum barcode overlap $\geq$
2 on the same haplotype, (3) no more than 1 barcode overlap between different haplotypes, (4)
FILTER value reported by the software was within this set {"PASS", "NOLONGFRAGS",
"NEARBYSNVS", or "NEARBYSNVS; NOLONGFRAGS"}, and (5) classified as somatic.

Long Ranger analysis generated SV calls for tumor and normal samples, independently. For each
tumor-normal pair, both large SVs ("large_sv_calls.bedpe") and deletions ("dels.vcf") were
combined for individual samples. Somatic tumor SVs were determined as events that were not
found in the matched normal sample based on the left breakpoints in tumor and normal being
within 1 kB and the right breakpoints in tumor and normal samples being within 1 kB. Only SV
events with FILTER values within this set {"PASS", "LOCAL_ASM", "SV", "CNV, SV"} and intra-
chromosomal events with span $\geq$ 100 kB were considered. SV events were only retained if both
breakpoints of an SV event were within 500 kB the boundaries of an overlapping CNA event and
the length of SV overlaps this CNA event by > 75%.

SV events from these three callers were then combined by taking the union of the filtered events
from. Intersecting events between 2 or more call-sets were determined if both breakpoints of one
event were located within 5 kB from both breakpoints of the event detected by the other tool. Then
the details of this event were retained based the priority ordered by SvABA, GROC-SVS, Long
Ranger. Long Ranger SV events were further filtered out if they were not intersecting events
detected by at least one other tool. SV events with span less than 1 kB were excluded from
downstream analyses.

An SV panel of normals (PoN) was generated using germline events from SvABA and Long
Ranger calls. There are two components to this panel: (1) frequency of germline events at exact
breakpoint locations (SVpon.bkpt) and (2) frequency of germline event breakpoint overlapping
within tiled windows of 1 kB (SVpon.blackListBins). The PoN was used to filter events in the
combined SV call-set when an SV has at least one breakpoint with SVpon.bkpt ≥ 2 and
overlapping bin with SVpon.blackListBins ≥ 100.

The workflow for SV analysis from linked-read sequencing data can be accessed at
https://github.com/GavinHaLab/SV_10X_analysis. Manual curation of filtered SV events in the *AR*
locus was performed and rescued events were labeled "Manual". The final SV call-set is included
in **Table S1K**.

For samples based on short-read WGS, SvABA was used in tumor-normal paired mode for SV
detection with default parameters. Intra-chromosomal SV events with span > 1 kB were retained.
The SvABA workflow can be accessed at https://github.com/GavinHaLab/TitanCNA_SV_WGS

Classification of structural variants in mCRPC

SV types were annotated based on orientations of breakpoints and bin-level copy-number around
breakpoints. The orientation of one breakpoint was defined based on the fragment of DNA
molecule being connected to the altered molecule. If the connected fragment was to the 5'-end of
the breakpoint, *i.e.*, "upstream" or "left" to the breakpoint, then the orientation was annotated as
forward or "+"; on the contrary, if the connected fragment was located to the 3'-end of the
breakpoint, the orientation was annotated as reverse or "-". The copy-number near each
breakpoint was evaluated using 10 kB bins. For one SV event, copy-number values of the bins
located to the upstream and downstream of breakpoint 1 were denoted as c_1^{up} and c_1^{down},
respectively; similarly, the copy-number values for breakpoint 2 were denoted as c_2^{up} and c_2^{down}.
In addition, then mean copy-number c^{mean} of the 10 kB bins between the two breakpoints of one
SV event and the number of bins s were also considered during SV classification. Intra-
chromosomal SV events, *i.e.*, both breakpoints were located on the same chromosome, were
classified to the list of SV types below following the corresponding classification criteria.

• Deletion. Events having the orientation combination (reverse, forward) and length
between 10 kB and 1 Mb were classified as deletions. The copy-number values of
breakpoints should satisfy $c_1^{up} > c_1^{down}$ or $c_2^{up} < c_2^{down}$, and $c_1^{up} > c^{mean}$ or $c_2^{down} > c^{mean}$, and
$s \leq 5$. In addition, events overlapping copy-number deletion or LOH segments were also
considered as deletions.
• Tandem duplication. Events having the orientation combination (forward, reverse) and
length between 10 kB and 1 Mb were classified as tandem duplications. The copy-number
values of breakpoints should satisfy $c_1^{up} < c_1^{down}$ or $c_2^{up} > c_2^{down}$, and $c_1^{up} < c^{mean}$ or $c_2^{down} <$
c^{mean}, and $s \leq 5$. In addition, events overlapping copy-number gain or copy neutral LOH
segments were also considered as tandem duplications.
• Inversion. Events having the orientation combination (forward, forward) or (reverse,
reverse) and length between 10 kB and 5 Mb were classified as inversions. Furthermore,
inversion events shorter than 30 kB with unequal copy-numbers around either breakpoint
were classified as fold-back inversions.

- Balanced rearrangement (balanced). Events having the orientation combination same to inversion (forward, forward) or (reverse, reverse), but length larger than 5 Mb were classified as balanced events. The copy-number values of breakpoints should satisfy $c_1^{up} = c_1^{down}$ and $c_2^{up} = c_2^{down}$, or $c_1^{up} = c^{mean}$ and $c_2^{down} = c^{mean}$.
- Unbalanced rearrangement (unbalanced). Intra-chromosomal events which did not fulfill any of the above criteria and having length larger than 10 kB were classified as unbalanced events.
- All SV events with two breakpoints located on different chromosomes were classified as translocations.

ICGC/TCGA PCAWG localized prostate cancer structural variants

We obtained localized prostate cancer structural variation calls from ICGC Data Portal release 28 (https://dcc.icgc.org/releases/PCAWG/consensus_sv). In this consensus SV file, each SV event was predicted by at least two variant callers. Samples that were classified as prostate adenocarcinoma (PRAD) and early onset prostate cancer (EOPC) were selected. A total of 278 samples successfully lifted over to genome build GRCh38. To maximize consistency with mCRPC datasets, we used only the PCAWG consensus SVs that included "SNOWMAN" as one of the tools. Note that "SNOWMAN" was the previous name for SvABA. Intrachromosomal SV events shorter than 10 kB were excluded.

Tandem duplicator phenotype

For all samples in the combined cohort, the TDP status was predicted using copy-number and SV by counting the number of copy-number segments overlapping with tandem duplication SV events, *i.e.,* gain segments. A sample was considered as TDP if it has more than 300, or 90 gain segments for samples based on linked-read sequencing and short-read sequencing, respectively. The number of segments with gain and median length SV are reported in **Table S1L**.

Chromothripsis analysis

Chromothripsis events were detected by ShatterSeek R package (Cortés-Ciriano et al., 2020). Structural variants calls by SvABA and copy-number calls by TitanCNA were used as input data (excluding Y chromosome). In the input, consecutive segments were joined as one if they had the same copy-number value and centromere regions were filtered out.

Manual inspection was performed for reported chromothripsis-like events after adapting criteria thresholds. For samples based on short-read sequencing, confidence classification criteria were refined from the ShatterSeek documentation. Following criteria were used for high confidence calls: total number of intra-chromosomal structural variants events involved in the event ≥ 10; max number of oscillating CN segments (two states) ≥ 10; satisfying either the chromosomal enrichment or the exponential distribution of breakpoints test ($p \leq 0.05$). For samples based on linked-read sequencing, we filtered these calls based on a weighted score that is primarily determined by the number of SVs in a cluster, with less weight given to CN oscillations. In this analysis, events with a score over 0.8 were considered as high confidence and all other events were excluded. The score is defined based on the following terms (ranges from 0 to 1).

- Weight 0.6 if total number of intra-chromosomal structural variants events involved in the event ≥ 10.
- Weight 0.2 for max number of oscillating CN segments (two states) ≥ 7 or max number of oscillating CN segments (three states) ≥ 14.
- Weight 0.1 for passing chromosomal enrichment test by ShatterSeek.
- Weight 0.1 for passing exponential distribution of breakpoints test.

Chromoplexy analysis

ChainFinder was used to detect chromoplexy events (Baca et al., 2013). Ten samples that were considered as TDP (01115374-TA2, 01115202-TC2, 01115248-TA3, 01115503-TC2, 01115257-TA4, 01115284-TA9, 01115414-TA1, DTB-063-BL, DTB-183-BL, DTB-214-BL) were excluded from this analysis. In addition, four samples that were found to cause numeric instabilities of ChainFinder were also excluded (DTB-023-BL, DTB-102-PRO, DTB-111-PRO, DTB-151-BL). The SV calls of remaining samples were further filtered to exclude those that were located within 5 Mb from chromosomal ends or overlapping chromothripsis regions. For copy-number input, segments that were determined as copy neutral by TitanCNA were set to have log copy-ratio of 0. Copy-ratio of the other segments were computed from copy-number values generated by TitanCNA divided by 2 for autosomes or 1 for X chromosome. Log copy-ratio values less than -1.5 were set to -1.5. The output of ChainFinder was used for determining chromoplexy status of individual samples. A chromoplexy event was defined as a chain including at least 5 rearrangement events and involving more than 2 different chromosomes. Samples having at least 2 such events were considered positive for chromoplexy status.

ChIP-seq data analysis

ChIP-seq data used in this study were downloaded from Gene Expression Omnibus (GEO) (Barrett et al., 2013) and the Sequence Read Archive (SRA) (Leinonen et al., 2011). Short reads were mapped to the human genome GRCh38 (hg38) using bwa (Li and Durbin, 2009). Because read lengths were less than 50bp, the bwa aln command with default parameters was used for mapping. MACS2 (Zhang et al., 2008) was used to identify peaks from mapped ChIP-seq data. For histone modification marks, MACS2 callpeak command was applied with --nomodel --broad –extsize 146. For CTCF data, MACS2 callpeak command was used with --nomodel --extsize 200. Below is the list of ChIP-seq datasets involved in this analysis.

- H3K4me3, H3K27me3 and CTCF (GSE38685) (Bert et al., 2013).
- H3K36me3 and H3K9me3 (GSE98732) (Du et al., 2019).
- H3K4me1 and H3K27ac (GSE73785) (Taberlay et al., 2016).

For *AR* binding site (ARBS), the peak files were downloaded from two different datasets and converted to hg38 coordinates. For primary prostate cancer, ARBS data were downloaded from GSE70079 (Pomerantz et al., 2015). The union of all tumor sample peaks was used. For mCRPC, met-specfic ARBS data were obtained from a previous study (Pomerantz et al., 2020).

Identification of SRB regions

Masking the human genome based on mappability

The human genome was divided into 100 kB non-overlapping bins for detection of significantly recurrent breakpoint regions (SRB). A low-mappability mask was generated for the hg38 genome to screen out out regions that are difficult for variant calling based on short-read sequencing. We adopted procedures from a previous study (Mallick et al., 2016) to construct a mask corresponding to regions with low mappability in the human genome. The unmasked regions were defined as the eligible territories for SRB detection. The 100 kB bins with less than 75% overlap with eligible territories were excluded from the analysis. Below is a list of masked regions included in the low-mappability mask.

- Composition mask. This set of masked regions includes regions with low sequence complexity detected by mdust, regions with long homopolymers detected by seqtk, satellite regions annotated by RepeatMasker (Smit, AFA, Hubley, R & Green, P, 2013), and low complexity regions annotated by RepeatMasker.

• Mappability mask. This mask was based on mappability of *k*-mers in the human genome
hg38. The value *k* was set to 75 which is half of the read length of WGS data in this study.
Each base in the genome was assigned a mappability level, based on the mapping
ambiguity of all 75-mers overlapping this specific base. See below for the list of mappability
levels.
o Level 0: all 75-mers overlapping this base could not be mapped to the genome
uniquely.
o Level 1: more than 50% of overlapping 75-mers are not uniquely mapped.
o Level 2: more than 50% of overlapping 75-mers are uniquely mapped with 1-
mismatch hits.
o Level 3: more than 50% of overlapping 75-mers are uniquely mapped without 1-
mismatch hits.

Regions with mappability level 0 and 1 were included in the low-mappability mask.

Generating covariates for regression analysis

To accurately model the genomic features of mCRPC, we incorporated the following covariates.

• Nucleotide composition, including GC content, CpG fraction and TpC fraction per 10 kB
non-overlapping bin in the genome.
• Replication timing of LNCaP (data obtained from ENCODE under accession
ENCFF995YGM, lifted over from hg19 to hg38) (Davis et al., 2018; ENCODE Project
Consortium, 2012).
• DNase I hypersensitive sites (data obtained from ENCODE under accession
ENCFF434GSJ, lifted over to hg38).
• Repeats annotated by RepeatMasker, including LINE, SINE, LTR, DNA transposon and
simple repeats.
• Heterochromatin regions inferred by ChromHMM (Ernst and Kellis, 2012) with the 18-state
model parameters from the Roadmap Epigenomics Project (Roadmap Epigenomics
Consortium et al., 2015), based LNCaP ChIP-seq data of H3K4me1, H3K4me3, H3K4ac
H3K27me3, H3K36me3 and H3K9me3.
• Common fragile sites downloaded from HGNC biomart (Tweedie et al., 2021).

SRB detection

Structural variants from the final call set were used for statistical enrichment of recurrent
breakpoints within 100 kB bins using a Gamma-Poisson regression implemented in the package,
fish.hook (Imielinski et al., 2017). Breakpoints of SVs were treated independently. The Benjamini-
Hochberg procedure was used for multiple testing correction and bins with q-value ≤ 0.1 were
determined to be significant. The distances of individual known driver genes to those significant
bins were evaluated based on the shortest genomic distance between the gene and bin
boundaries, regardless of gene orientations.

Annotation of gene alteration status

Gene alteration by copy-number

Copy-number segments were excluded if their cellular fraction was lower than 0.8, except for
those which were determined as copy neutral or copy-number greater than 4. The gene
annotation was based on known protein coding genes from GenCode release 30 (GRCh38.p12)
(Frankish et al., 2019). For each gene, its copy-number was assigned to the copy-number value
and LOH status of the segment that has the largest overlap with it. The gene-level copy-number
was normalized based on ploidy of the corresponding sample, with autosomal genes normalized

by the inferred ploidy rounded to nearest integer, and X-linked genes normalized by half such
value. Then the copy-number status of each gene was categorized based on the following criteria.

- Amplification. Normalized gene-level copy-number is greater than or equal to 2.5.
- Gain. Normalized gene-level copy-number is between 2 and 2.5.
- Homozygous deletion. Normalized gene-level copy-number is 0.
- Deletion with LOH. Normalized gene-level copy-number is between 0 and 1, and LOH
status was found.
- Copy neutral LOH. Normalized gene-level copy-number is 1 and LOH status was found.

Gene alteration by structural variant

Gene coordinates were based on ENSEMBL v33 of hg38 (Howe et al., 2021). Gene body region
of one gene was defined as the widest region of all known isoforms collapsed. Gene flanking
region was defined as the corresponding two 1 Mb regions next to the gene body region on 5'-
end and 3'-end, respectively.

Gene alteration status by genome rearrangements was defined based on the breakpoints and
directions of involving structural variant events. A gene in one WGS sample (gene-sample pair)
was considered having gene transecting events if any breakpoints of SV events were located
within the gene body region. If the gene transecting status did not apply, then this gene-sample
pair was examined for gene flanking status if the breakpoints of any intra-chromosomal SV events,
including tandem duplications, deletions, and inversions, were located within the gene flanking
regions. Additionally, translocation events including intra-chromosomal balanced and unbalanced
events which spanned over 10 Mb, and inter-chromosomal translocation events were considered
altering the gene flanking regions if any of their breakpoints was in the gene flanking region, and
the direction of the SV was going towards the gene body region. The alteration status of
rearrangements for each gene-sample pair was exclusive between gene transecting and gene
flanking, with the former being prioritized in report.

AR alteration analysis

Copy-number of the *AR* gene (chrX:67,544,623-67,730,619) and the *AR* enhancer region
(chrX:66,895,000-66,910,000) were each computed as the mean corrected total copy-number
across the 10 kB bins overlapping each region. The copy-number was further normalized by
sample ploidy as previously described. Amplification status of *AR* was determined by comparing
the log2 fold-change *FC* of enhancer-level over gene-level copy-number. Four distinct groups
were defined based on copy-number and *FC* as below.

- Co-amplification. Ploidy normalized copy-number values of both *AR* gene body and
enhancer are greater than 1.5.
- Selective *AR* amplification. $FC < -\log2(1.5)$ and enhancer copy-number is less than 1.5.
- Selective enhancer copy gain. $FC > \log2(1.5)$ and *AR* gene body copy-number is less than
1.5.
- Lack of amplification for both. All other cases were considered as no amplification for both
regions.

ANCOVA test was used to test if different patterns of *AR* amplification have an impact on *AR*
expression. Batch corrected log10(TPM+1) values using ComBat from sva R package (v3.34.0)
were used for *AR* expression level. We fit the ANCOVA model using *AR* expression as the
response variable, *AR* amplification status as the predictor variable, and ploidy, purity as
covariates. The function Anova in the car package (v3.0-5) was used with Type III sum of squares
for the model. Post hoc analysis was performed to determine the specific differences among four

different *AR* amplification status. The function glht was used within the multcomp package (v1.4-
11) in R to perform Tukey's Test for multiple comparisons.

Gene expression

906 TPM values for a subset of the samples based on linked-read sequencing were obtained from
907 cBioportal (Cerami et al., 2012; Gao et al., 2013). For samples based on short-read sequencing
908 the TPM values were obtained from a previous study (Quigley et al., 2018).

Gene fusion analysis

Fusion status of the main members of the ETS family, including *ERG*, *ETV1*, *ETV4*, *ETV5* and
ELK4 was analyzed. Determination of gene fusion status was based on both DNA and RNA levels.
For DNA, structural variants transecting gene body regions were used. SV events were
considered supporting gene fusion only if they satisfy the following criteria: (1) the breakpoints of
this event must be located within the ETS gene and another protein coding gene, respectively; (2)
the orientation of the breakpoint located within the ETS gene must be pointing towards the coding
sequence of ETS domain. For RNA, arriba was used to detect fusion transcripts from RNA-seq
data (Uhrig et al., 2021). The fusion status was only confirmed if all following conditions were
satisfied: (1) the complete ETS domain was included in the fusion product; (2) detection
confidence reported by arriba is "high"; (3) coding sequence in the fusion transcript was in sense
orientation and no out-of-frame shifts.

SV signature analysis

Signature extraction and clustering

De novo signature extraction was performed on all SV events called by SvABA of the combined
cohort using signature.tools.lib (Degasperi et al., 2020) with the recommended settings of 20
bootstraps, 200 repeats, the clustering with matching algorithm, the KLD objective function, and
RTOL = 0.001. The exposure of one signature in one sample is defined as the median activity of
the signature within the sample across all bootstraps. For clustering, the reference signature
exposure values for each sample based on short-read sequencing were normalized such that the
sum of exposure values per sample is 1, and the normalized exposure values for each signature
were mean-centered across all samples. A Euclidean distance matrix was computed and then
samples were clustered with the Ward.D2 algorithm using R's hclust function. We chose the
number of clusters to be k = 9 based on dendrogram using cutree function in R.

Enrichment of alterations in SV clusters

All 9 identified SV clusters were analyzed for enrichment of alterations. To make the analysis
unbiased by SV signature, we limited our search to alteration types that were orthogonal to
rearrangements, which include SNV, copy-number gain and copy-number loss. We performed
hypothesis testing on each driver-alteration pair, and also on chromoplexy and chromothripsis.
For each SV cluster, a χ^2 test was performed for each driver gene alteration status, with samples
within group being tested against samples belonging to all 8 other SV clusters. Multiple testing
adjustment based on Benjamini-Hochberg FDR was performed to compute q-values. Alteration
categories with q-values less than 0.25 were determined as enriched in the corresponding SV
cluster.

Survival analysis

944 Survival data was obtained from (Chen et al., 2019). Survival analyses were conducted using the
945 Kaplan-Meier method with log-rank testing for significance. The function survfit from survival R
946 package was used to perform the analysis.

 SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Recurrent CNA and alteration profiles of most frequently altered genes, related to Figure 1.

(A) Recurrent copy number gain events in the genome. The frequencies of copy number gain are plotted in red according to their genomic coordinates. Regions with significantly recurring CNA are colored in black. Known driver genes that are within those regions are labeled.

(B) Recurrent copy number loss events in the genome. The frequencies of copy number loss are plotted in blue with y-axis inverted.

(C) Alteration profiles of known prostate cancer driver genes. Alterations are categorized into CNA, SNV and SV, with SV being further divided into gene transecting (SV tr.) and gene flanking (SV fl.). The percentages of samples carrying corresponding alterations are shown as stacked bars. All known prostate cancer driver genes were considered and the top 16 genes with overall alteration frequencies above 30% are shown.

Figure S2. Recurrent SV in localized prostate cancer and landscape of ETS fusion in mCRPC, related to Figure 2.

(A) SRBs detected in the cohort of localized prostate cancers. The criteria for coloring and labeling are the same as Figure 2.

(B) Translocation events originating from *TTC28*. In the circos plot, *TTC28* is labeled with a vertical bar at the 22q12.1 locus. Translocation events which have breakpoints located within 5 kB to the 3'-end of the L1 retrotransposon are visualized as blue arcs.

(C) Schematics of cumulative counts from intra-chromosomal SV events. Individual SV events are indicated by a grey arc, and colored crosses correspond to breakpoints of each event.

(D) Expression and fusion status for main genes of the ETS family. The expression values were normalized from TPM to z-score within each gene. Grey boxes indicate expression data are not available. For fusion status, color indicates the data type which was used to call fusion.

Figure S3. Comparison of genomic alterations in disease states, related to Figure 3.

(A) Alteration status of paired samples from the same patients before and after treatment. Known prostate driver genes with alteration status in any of the included samples are shown.

(B) Comparison of rearrangement frequency in different disease states of mCRPC. The known prostate cancer driver genes that were located within 1 Mb to any SRB region are included.

(C) *AR* splice variants in sample DTB-124-BL. The expression values of all known *AR* exons, including both canonical and cryptic ones, are shown in the top panel. In the bottom panel, the number of reads covering the junction sites of two exons are indicated by weighted arcs.

Figure S4. Signature analysis of SV events, related to Figure 4.

(A) Workflow of SV signature analysis. Samples involved in this analysis are described in green boxes. Details of relevant signatures are shown in blue boxes. The steps for obtaining the final 9 SV clusters are indicated by numbers.

(B) Signature exposure based on de novo SV signatures. The exposure values of each sample were normalized such that the sample-wise sum is 1. Samples are ordered alphabetically based on names. The sequencing technology used for each sample is labeled at the bottom.

(C) Similarity between de novo SV signatures and reference signatures. Pairwise cosine similarity
between de novo and reference SV signatures is shown.

(D) Comparison of reference signature (RefSig) prevalence between mCRPC and localized
prostate cancer. The prevalence value for a signature in mCRPC was computed based on
samples harboring at least 5% signature exposure. Localized prostate cancer prevalence values
were obtained from signal.mutationalsignatures.com (Degasperi et al., 2020) computed from 199
PCAWG samples.

(E) Kaplan Meier curve of prediction using mutation class of key marker genes. Samples were
grouped based on the mutation status of the corresponding marker gene.

(F) Kaplan Meier curve of prediction using SV cluster information. Samples were grouped based
on their assignments of the corresponding SV cluster.

SUPPLEMENTAL TABLE LEGENDS

Table S1. Sequencing, clinical and alteration information of all samples involved in this
study. Related to Figures 1, 2, 4, S1-3.

(A) Sequencing metrics of all samples based on linked-read sequencing.

(B) Clinical properties and key genomics metrics of the cohort.

(C) Somatic mutation status of 159 prostate cancer drivers in the cohort. Sample and genes were
sorted alphabetically. Genes with no detected mutations were left blank.

(D) Significantly mutated genes (q ≤ 0.1) detected by dN/dS algorithm.

(E) Somatic copy-number alteration status of 159 prostate cancer drivers in the cohort.

(F) Recurrent copy-number alteration peaks detected by GISTIC.

(G) Gene transecting rearrangements of 159 prostate cancer drivers in the cohort. Types of
rearrangement events were included.

(H) Gene flanking rearrangements of prostate cancer drivers in the cohort.

(I) TITAN copy number segments for all samples. Columns with "Corrected_*" were used for
analysis in this study.

(J) TITAN optimal solutions selected for all samples.

(K) Structural variant calls for all samples. For samples with linked-read data ("CRPC10X"), union
set of detected calls from SvABA, GROC-SVS, and Long Ranger are indicated. `SV.Filter`
indicate SV events after filtering. `support` contain evidence from various callers; manual curation
of events is indicated here. `CN_overlap_type` contain the final SV classification after annotation
with copy number information.

(L) TDP status, copy number gain event counts, and median tandem duplication lengths for all
samples.

Table S2. Significantly recurrent breakpoint regions and ETS fusion. Related to Figure 2,
S2 and S3.

(A) Significantly recurrent breakpoints (SRB) regions (q ≤ 0.1) in the mCRPC cohort of 143
samples.

(B) Significantly recurrent breakpoints (SRB) regions (q ≤ 0.1) in the localized prostate cancer
cohort of 278 samples.

(C) Fusion status of the ETS family genes. Gene expression was normalized to z-score for each
gene. Genes with no detected fusion events or available expression data were left blank.

Table S3. *AR* alteration patterns in the mCRPC cohort. Related to Figure 3 and S3.

Table S4. SV signature in the mCRPC cohort. Related to Figure 4 and S4.

(A) Matrix of cosine similarity with rows representing reference signatures and columns
representing *de novo* signatures.

(B) Exposure of all 8 *de novo* signatures in the cohort. Values were not normalized.

(C) Exposure of 8 chosen reference signatures in the cohort. Values were not normalized.

1036

REFERENCES

Abida, W., Cheng, M.L., Armenia, J., Middha, S., Autio, K.A., Vargas, H.A., Rathkopf, D., Morris, M.J., Danila, D.C., Slovin, S.F., et al. (2019). Analysis of the Prevalence of Microsatellite Instability in Prostate Cancer and Response to Immune Checkpoint Blockade. JAMA Oncol *5*, 471–478.

Abida, W., Patnaik, A., Campbell, D., Shapiro, J., Bryce, A.H., McDermott, R., Sautois, B., Vogelzang, N.J., Bambury, R.M., Voog, E., et al. (2020). Rucaparib in Men With Metastatic Castration-Resistant Prostate Cancer Harboring a BRCA1 or BRCA2 Gene Alteration. J Clin Oncol *38*, 3763–3772.

Adalsteinsson, V.A., Ha, G., Freeman, S.S., Choudhury, A.D., Stover, D.G., Parsons, H.A., Gydush, G., Reed, S.C., Rotem, D., Rhoades, J., et al. (2017). Scalable whole-exome sequencing of cell-free DNA reveals high concordance with metastatic tumors. Nat. Commun. *8*, 1324.

Alexandrov, L.B., Nik-Zainal, S., Wedge, D.C., Aparicio, S.A.J.R., Behjati, S., Biankin, A.V., Bignell, G.R., Bolli, N., Borg, A., Børresen-Dale, A.-L., et al. (2013). Signatures of mutational processes in human cancer. Nature *500*, 415–421.

Alexandrov, L.B., Kim, J., Haradhvala, N.J., Huang, M.N., Tian Ng, A.W., Wu, Y., Boot, A., Covington, K.R., Gordenin, D.A., Bergstrom, E.N., et al. (2020). The repertoire of mutational signatures in human cancer. Nature *578*, 94–101.

Armenia, J., Wankowicz, S.A.M., Liu, D., Gao, J., Kundra, R., Reznik, E., Chatila, W.K., Chakravarty, D., Han, G.C., Coleman, I., et al. (2018). The long tail of oncogenic drivers in prostate cancer. Nature Genetics *50*, 645–651.

Baca, S.C., Prandi, D., Lawrence, M.S., Mosquera, J.M., Romanel, A., Drier, Y., Park, K., Kitabayashi, N., MacDonald, T.Y., Ghandi, M., et al. (2013). Punctuated evolution of prostate cancer genomes. Cell *153*, 666–677.

Barbieri, C.E., Baca, S.C., Lawrence, M.S., Demichelis, F., Blattner, M., Theurillat, J.-P., White, T.A., Stojanov, P., Van Allen, E., Stransky, N., et al. (2012). Exome sequencing identifies recurrent SPOP, FOXA1 and MED12 mutations in prostate cancer. Nat Genet *44*, 685–689.

Barrett, T., Wilhite, S.E., Ledoux, P., Evangelista, C., Kim, I.F., Tomashevsky, M., Marshall, K.A., Phillippy, K.H., Sherman, P.M., Holko, M., et al. (2013). NCBI GEO: archive for functional genomics data sets--update. Nucleic Acids Res *41*, D991-995.

Bert, S.A., Robinson, M.D., Strbenac, D., Statham, A.L., Song, J.Z., Hulf, T., Sutherland, R.L., Coolen, M.W., Stirzaker, C., and Clark, S.J. (2013). Regional activation of the cancer genome by long-range epigenetic remodeling. Cancer Cell *23*, 9–22.

de Bono, J., Mateo, J., Fizazi, K., Saad, F., Shore, N., Sandhu, S., Chi, K.N., Sartor, O., Agarwal, N., Olmos, D., et al. (2020). Olaparib for Metastatic Castration-Resistant Prostate Cancer. N Engl J Med *382*, 2091–2102.

de Bono, J.S., Logothetis, C.J., Molina, A., Fizazi, K., North, S., Chu, L., Chi, K.N., Jones, R.J., Goodman, O.B., Saad, F., et al. (2011). Abiraterone and Increased Survival in Metastatic Prostate Cancer. N Engl J Med *364*, 1995–2005.

Brand, L.J., and Dehm, S.M. (2013). Androgen Receptor Gene Rearrangements: New
Perspectives on Prostate Cancer Progression. Curr Drug Targets *14*, 441–449.

Campbell, P.J., Getz, G., Korbel, J.O., Stuart, J.M., Jennings, J.L., Stein, L.D., Perry, M.D., Nahal-
Bose, H.K., Ouellette, B.F.F., Li, C.H., et al. (2020). Pan-cancer analysis of whole genomes.
Nature *578*, 82–93.

Cancer Genome Atlas Research Network (2015). The Molecular Taxonomy of Primary Prostate
Cancer. Cell *163*, 1011–1025.

Carrot-Zhang, J., and Majewski, J. (2017). LoLoPicker: detecting low allelic-fraction variants from
low-quality cancer samples. Oncotarget *8*, 37032.

Céraline, J., Cruchant, M.D., Erdmann, E., Erbs, P., Kurtz, J.-E., Duclos, B., Jacqmin, D., Chopin,
D., and Bergerat, J.-P. (2004). Constitutive activation of the androgen receptor by a point mutation
in the hinge region: a new mechanism for androgen-independent growth in prostate cancer. Int.
J. Cancer *108*, 152–157.

Cerami, E., Gao, J., Dogrusoz, U., Gross, B.E., Sumer, S.O., Aksoy, B.A., Jacobsen, A., Byrne,
C.J., Heuer, M.L., Larsson, E., et al. (2012). The cBio Cancer Genomics Portal: An Open Platform
for Exploring Multidimensional Cancer Genomics Data: Figure 1. Cancer Discovery *2*, 401–404.

Chen, C.D., Welsbie, D.S., Tran, C., Baek, S.H., Chen, R., Vessella, R., Rosenfeld, M.G., and
Sawyers, C.L. (2004). Molecular determinants of resistance to antiandrogen therapy. Nature
Medicine *10*, 33–39.

Chen, J., Lui, W.-O., Vos, M.D., Clark, G.J., Takahashi, M., Schoumans, J., Khoo, S.K., Petillo,
D., Lavery, T., Sugimura, J., et al. (2003). The t(1;3) breakpoint-spanning genes LSAMP and
NORE1 are involved in clear cell renal cell carcinomas. Cancer Cell *4*, 405–413.

Chen, W.S., Aggarwal, R., Zhang, L., Zhao, S.G., Thomas, G.V., Beer, T.M., Quigley, D.A., Foye,
A., Playdle, D., Huang, J., et al. (2019). Genomic Drivers of Poor Prognosis and Enzalutamide
Resistance in Metastatic Castration-resistant Prostate Cancer. Eur Urol *76*, 562–571.

Cortés-Ciriano, I., Lee, J.J.-K., Xi, R., Jain, D., Jung, Y.L., Yang, L., Gordenin, D., Klimczak, L.J.,
Zhang, C.-Z., Pellman, D.S., et al. (2020). Comprehensive analysis of chromothripsis in 2,658
human cancers using whole-genome sequencing. Nat Genet *52*, 331–341.

Davis, C.A., Hitz, B.C., Sloan, C.A., Chan, E.T., Davidson, J.M., Gabdank, I., Hilton, J.A., Jain,
K., Baymuradov, U.K., Narayanan, A.K., et al. (2018). The Encyclopedia of DNA elements
(ENCODE): data portal update. Nucleic Acids Res *46*, D794–D801.

Degasperi, A., Amarante, T.D., Czarnecki, J., Shooter, S., Zou, X., Glodzik, D., Morganella, S.,
Nanda, A.S., Badja, C., Koh, G., et al. (2020). A practical framework and online tool for mutational
signature analyses show inter-tissue variation and driver dependencies. Nat Cancer *1*, 249–263.

van Dessel, L.F., van Riet, J., Smits, M., Zhu, Y., Hamberg, P., van der Heijden, M.S., Bergman,
A.M., van Oort, I.M., de Wit, R., Voest, E.E., et al. (2019). The genomic landscape of metastatic
castration-resistant prostate cancers reveals multiple distinct genotypes with potential clinical
impact. Nat Commun *10*, 5251.

Du, Q., Bert, S.A., Armstrong, N.J., Caldon, C.E., Song, J.Z., Nair, S.S., Gould, C.M., Luu, P.-L.,
Peters, T., Khoury, A., et al. (2019). Replication timing and epigenome remodelling are associated
with the nature of chromosomal rearrangements in cancer. Nat Commun *10*, 416.

ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in the human
genome. Nature *489*, 57–74.

Ernst, J., and Kellis, M. (2012). ChromHMM: automating chromatin-state discovery and
characterization. Nat Methods *9*, 215–216.

Frankish, A., Diekhans, M., Ferreira, A.-M., Johnson, R., Jungreis, I., Loveland, J., Mudge, J.M.,
Sisu, C., Wright, J., Armstrong, J., et al. (2019). GENCODE reference annotation for the human
and mouse genomes. Nucleic Acids Res *47*, D766–D773.

Fraser, M., Livingstone, J., Wrana, J.L., Finelli, A., He, H.H., van der Kwast, T., Zlotta, A.R.,
Bristow, R.G., and Boutros, P.C. (2021). Somatic driver mutation prevalence in 1844 prostate
cancers identifies ZNRF3 loss as a predictor of metastatic relapse. Nat Commun *12*, 6248.

Gao, J., Aksoy, B.A., Dogrusoz, U., Dresdner, G., Gross, B., Sumer, S.O., Sun, Y., Jacobsen, A.,
Sinha, R., Larsson, E., et al. (2013). Integrative analysis of complex cancer genomics and clinical
profiles using the cBioPortal. Sci Signal *6*, pl1.

Giambartolomei, C., Seo, J.-H., Schwarz, T., Freund, M.K., Johnson, R.D., Spisak, S., Baca, S.C.,
Gusev, A., Mancuso, N., Pasaniuc, B., et al. (2021). H3K27ac HiChIP in prostate cell lines
identifies risk genes for prostate cancer susceptibility. The American Journal of Human Genetics
108, 2284–2300.

Glodzik, D., Morganella, S., Davies, H., Simpson, P.T., Li, Y., Zou, X., Diez-Perez, J., Staaf, J.,
Alexandrov, L.B., Smid, M., et al. (2017). A somatic-mutational process recurrently duplicates
germline susceptibility loci and tissue-specific super-enhancers in breast cancers. Nat Genet *49*,
341–348.

Grasso, C.S., Wu, Y.-M., Robinson, D.R., Cao, X., Dhanasekaran, S.M., Khan, A.P., Quist, M.J.,
Jing, X., Lonigro, R.J., Brenner, J.C., et al. (2012). The mutational landscape of lethal castration-
resistant prostate cancer. Nature *487*, 239–243.

Ha, G., Roth, A., Khattra, J., Ho, J., Yap, D., Prentice, L.M., Melnyk, N., McPherson, A.,
Bashashati, A., Laks, E., et al. (2014). TITAN: inference of copy number architectures in clonal
cell populations from tumor whole-genome sequence data. Genome Res *24*, 1881–1893.

Hadi, K., Yao, X., Behr, J.M., Deshpande, A., Xanthopoulakis, C., Tian, H., Kudman, S., Rosiene,
J., Darmofal, M., DeRose, J., et al. (2020). Distinct Classes of Complex Structural Variation
Uncovered across Thousands of Cancer Genome Graphs. Cell *183*, 197-210.e32.

Haffner, M.C., Aryee, M.J., Toubaji, A., Esopi, D.M., Albadine, R., Gurel, B., Isaacs, W.B., Bova,
G.S., Liu, W., Xu, J., et al. (2010). Androgen-induced TOP2B-mediated double-strand breaks and
prostate cancer gene rearrangements. Nat Genet *42*, 668–675.

Henzler, C., Li, Y., Yang, R., McBride, T., Ho, Y., Sprenger, C., Liu, G., Coleman, I., Lakely, B.,
Li, R., et al. (2016). Truncation and constitutive activation of the androgen receptor by diverse
genomic rearrangements in prostate cancer. Nat Commun *7*, 13668.

Howe, K.L., Achuthan, P., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M.R., Armean, I.M.,
Azov, A.G., Bennett, R., Bhai, J., et al. (2021). Ensembl 2021. Nucleic Acids Res *49*, D884–D891.

Imielinski, M., Guo, G., and Meyerson, M. (2017). Insertions and Deletions Target Lineage-
Defining Genes in Human Cancers. Cell *168*, 460-472.e14.

Kanayama, M., Lu, C., Luo, J., and Antonarakis, E.S. (2021). AR Splicing Variants and Resistance
to AR Targeting Agents. Cancers (Basel) *13*, 2563.

Kresse, S.H., Ohnstad, H.O., Paulsen, E.B., Bjerkehagen, B., Szuhai, K., Serra, M., Schaefer, K.-
L., Myklebost, O., and Meza-Zepeda, L.A. (2009). LSAMP, a novel candidate tumor suppressor
gene in human osteosarcomas, identified by array comparative genomic hybridization. Genes
Chromosomes Cancer *48*, 679–693.

Kühn, M.W.M., Radtke, I., Bullinger, L., Goorha, S., Cheng, J., Edelmann, J., Gohlke, J., Su, X.,
Paschka, P., Pounds, S., et al. (2012). High-resolution genomic profiling of adult and pediatric
core-binding factor acute myeloid leukemia reveals new recurrent genomic alterations. Blood *119*,
e67.

Kumar-Sinha, C., Kalyana-Sundaram, S., and Chinnaiyan, A.M. (2015). Landscape of gene
fusions in epithelial cancers: seq and ye shall find. Genome Medicine *7*, 129.

Leinonen, R., Sugawara, H., Shumway, M., and International Nucleotide Sequence Database
Collaboration (2011). The sequence read archive. Nucleic Acids Res *39*, D19-21.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics *25*, 1754–1760.

Li, Y., Roberts, N.D., Wala, J.A., Shapira, O., Schumacher, S.E., Kumar, K., Khurana, E., Waszak,
S., Korbel, J.O., Haber, J.E., et al. (2020). Patterns of somatic structural variation in human cancer
genomes. Nature *578*, 112–121.

Macintyre, G., Goranova, T.E., De Silva, D., Ennis, D., Piskorz, A.M., Eldridge, M., Sie, D.,
Lewsley, L.-A., Hanif, A., Wilson, C., et al. (2018). Copy number signatures and mutational
processes in ovarian carcinoma. Nat Genet *50*, 1262–1270.

Mallick, S., Li, H., Lipson, M., Mathieson, I., Gymrek, M., Racimo, F., Zhao, M., Chennagiri, N.,
Nordenfelt, S., Tandon, A., et al. (2016). The Simons Genome Diversity Project: 300 genomes
from 142 diverse populations. Nature *538*, 201–206.

Martincorena, I., Raine, K.M., Gerstung, M., Dawson, K.J., Haase, K., Van Loo, P., Davies, H.,
Stratton, M.R., and Campbell, P.J. (2017). Universal Patterns of Selection in Cancer and Somatic
Tissues. Cell *171*, 1029-1041.e21.

Mateo, J., McKay, R., Abida, W., Aggarwal, R., Alumkal, J., Alva, A., Feng, F., Gao, X., Graff, J.,
Hussain, M., et al. (2020). Accelerating precision medicine in metastatic prostate cancer. Nat
Cancer *1*, 1041–1053.

Nguyen, B., Mota, J.M., Nandakumar, S., Stopsack, K.H., Weg, E., Rathkopf, D., Morris, M.J.,
Scher, H.I., Kantoff, P.W., Gopalan, A., et al. (2020). Pan-cancer Analysis of CDK12 Alterations

Identifies a Subset of Prostate Cancers with Distinct Genomic and Clinical Characteristics.
European Urology *78*, 671–679.

Nik-Zainal, S., Davies, H., Staaf, J., Ramakrishna, M., Glodzik, D., Zou, X., Martincorena, I.,
Alexandrov, L.B., Martin, S., Wedge, D.C., et al. (2016). Landscape of somatic mutations in 560
breast cancer whole-genome sequences. Nature *534*, 47–54.

Petrovics, G., Li, H., Stümpel, T., Tan, S.-H., Young, D., Katta, S., Li, Q., Ying, K., Klocke, B.,
Ravindranath, L., et al. (2015). A novel genomic alteration of LSAMP associates with aggressive
prostate cancer in African American men. EBioMedicine *2*, 1957.

Pitkänen, E., Cajuso, T., Katainen, R., Kaasinen, E., Välimäki, N., Palin, K., Taipale, J., Aaltonen,
L.A., and Kilpivaara, O. (2014). Frequent L1 retrotranspositions originating from TTC28 in
colorectal cancer. Oncotarget *5*, 853–859.

Pomerantz, M.M., Li, F., Takeda, D.Y., Lenci, R., Chonkar, A., Chabot, M., Cejas, P., Vazquez,
F., Cook, J., Shivdasani, R.A., et al. (2015). The androgen receptor cistrome is extensively
reprogrammed in human prostate tumorigenesis. Nat Genet *47*, 1346–1351.

Pomerantz, M.M., Qiu, X., Zhu, Y., Takeda, D.Y., Pan, W., Baca, S.C., Gusev, A., Korthauer, K.D.,
Severson, T.M., Ha, G., et al. (2020). Prostate cancer reactivates developmental epigenomic
programs during metastatic progression. Nat. Genet.

Pradhan, B., Cajuso, T., Katainen, R., Sulo, P., Tanskanen, T., Kilpivaara, O., Pitkänen, E.,
Aaltonen, L.A., Kauppi, L., and Palin, K. (2017). Detection of subclonal L1 transductions in
colorectal cancer by long-distance inverse-PCR and Nanopore sequencing. Sci Rep *7*, 14521.

Pritchard, C.C., Mateo, J., Walsh, M.F., De Sarkar, N., Abida, W., Beltran, H., Garofalo, A., Gulati,
R., Carreira, S., Eeles, R., et al. (2016). Inherited DNA-Repair Gene Mutations in Men with
Metastatic Prostate Cancer. New England Journal of Medicine *375*, 443–453.

Qin, F., Zhang, Y., Liu, J., and Li, H. (2017). SLC45A3-ELK4 functions as a long non-coding
chimeric RNA. Cancer Lett *404*, 53–61.

Quigley, D.A., Dang, H.X., Zhao, S.G., Lloyd, P., Aggarwal, R., Alumkal, J.J., Foye, A., Kothari,
V., Perry, M.D., Bailey, A.M., et al. (2018). Genomic Hallmarks and Structural Variation in
Metastatic Prostate Cancer. Cell *174*, 758-769.e9.

Rickman, D.S., Pflueger, D., Moss, B., VanDoren, V.E., Chen, C.X., de la Taille, A., Kuefer, R.,
Tewari, A.K., Setlur, S.R., Demichelis, F., et al. (2009). SLC45A3-ELK4 is a novel and frequent
erythroblast transformation-specific fusion transcript in prostate cancer. Cancer Res *69*, 2734–
2738.

Roadmap Epigenomics Consortium, Kundaje, A., Meuleman, W., Ernst, J., Bilenky, M., Yen, A.,
Heravi-Moussavi, A., Kheradpour, P., Zhang, Z., Wang, J., et al. (2015). Integrative analysis of
111 reference human epigenomes. Nature *518*, 317–330.

Saunders, C.T., Wong, W.S.W., Swamy, S., Becq, J., Murray, L.J., and Cheetham, R.K. (2012).
Strelka: accurate somatic small-variant calling from sequenced tumor-normal sample pairs.
Bioinformatics *28*, 1811–1817.

Scher, H.I., Fizazi, K., Saad, F., Taplin, M.-E., Sternberg, C.N., Miller, K., de Wit, R., Mulders, P.,
Chi, K.N., Shore, N.D., et al. (2012). Increased Survival with Enzalutamide in Prostate Cancer
after Chemotherapy. New England Journal of Medicine *367*, 1187–1197.

Smit, AFA, Hubley, R & Green, P (2013). RepeatMasker Open-4.0.

Spies, N., Weng, Z., Bishara, A., McDaniel, J., Catoe, D., Zook, J.M., Salit, M., West, R.B.,
Batzoglou, S., and Sidow, A. (2017). Genome-wide reconstruction of complex structural variants
using read clouds. Nature Methods.

Stephens, P.J., Greenman, C.D., Fu, B., Yang, F., Bignell, G.R., Mudie, L.J., Pleasance, E.D.,
Lau, K.W., Beare, D., Stebbings, L.A., et al. (2011). Massive genomic rearrangement acquired in
a single catastrophic event during cancer development. Cell *144*, 27–40.

Taberlay, P.C., Achinger-Kawecka, J., Lun, A.T.L., Buske, F.A., Sabir, K., Gould, C.M., Zotenko,
E., Bert, S.A., Giles, K.A., Bauer, D.C., et al. (2016). Three-dimensional disorganization of the
cancer genome occurs coincident with long-range genetic and epigenetic alterations. Genome
Res *26*, 719–731.

Takeda, D.Y., Spisák, S., Seo, J.-H., Bell, C., O'Connor, E., Korthauer, K., Ribli, D., Csabai, I.,
Solymosi, N., Szállási, Z., et al. (2018). A Somatically Acquired Enhancer of the Androgen
Receptor Is a Noncoding Driver in Advanced Prostate Cancer. Cell *174*, 422-432.e13.

Tate, J.G., Bamford, S., Jubb, H.C., Sondka, Z., Beare, D.M., Bindal, N., Boutselakis, H., Cole,
C.G., Creatore, C., Dawson, E., et al. (2019). COSMIC: the Catalogue Of Somatic Mutations In
Cancer. Nucleic Acids Res *47*, D941–D947.

Tomlins, S.A., Rhodes, D.R., Perner, S., Dhanasekaran, S.M., Mehra, R., Sun, X.-W., Varambally,
S., Cao, X., Tchinda, J., Kuefer, R., et al. (2005). Recurrent fusion of TMPRSS2 and ETS
transcription factor genes in prostate cancer. Science *310*, 644–648.

Tomlins, S.A., Laxman, B., Dhanasekaran, S.M., Helgeson, B.E., Cao, X., Morris, D.S., Menon,
A., Jing, X., Cao, Q., Han, B., et al. (2007). Distinct classes of chromosomal rearrangements
create oncogenic ETS gene fusions in prostate cancer. Nature *448*, 595–599.

Tubio, J.M.C., Li, Y., Ju, Y.S., Martincorena, I., Cooke, S.L., Tojo, M., Gundem, G., Pipinikas,
C.P., Zamora, J., Raine, K., et al. (2014). Mobile DNA in cancer. Extensive transduction of
nonrepetitive DNA mediated by L1 retrotransposition in cancer genomes. Science *345*, 1251343.

Tweedie, S., Braschi, B., Gray, K., Jones, T.E.M., Seal, R.L., Yates, B., and Bruford, E.A. (2021).
Genenames.org: the HGNC and VGNC resources in 2021. Nucleic Acids Res *49*, D939–D946.

Uhrig, S., Ellermann, J., Walther, T., Burkhardt, P., Fröhlich, M., Hutter, B., Toprak, U.H.,
Neumann, O., Stenzinger, A., Scholl, C., et al. (2021). Accurate and efficient detection of gene
fusions from RNA sequencing data. Genome Res *31*, 448–460.

Umbreit, N.T., Zhang, C.-Z., Lynch, L.D., Blaine, L.J., Cheng, A.M., Tourdot, R., Sun, L.,
Almubarak, H.F., Judge, K., Mitchell, T.J., et al. (2020). Mechanisms generating cancer genome
complexity from a single cell division error. Science *368*, eaba0712.

Van der Auwera, G.A., and O'Connor, B. (2020). Genomics in the Cloud.

Veeriah, S., Brennan, C., Meng, S., Singh, B., Fagin, J.A., Solit, D.B., Paty, P.B., Rohle, D.,
Vivanco, I., Chmielecki, J., et al. (2009). The tyrosine phosphatase PTPRD is a tumor suppressor
that is frequently inactivated and mutated in glioblastoma and other human cancers. PNAS *106*,
9435–9440.

Visakorpi, T., Hyytinen, E., Koivisto, P., Tanner, M., Keinänen, R., Palmberg, C., Palotie, A.,
Tammela, T., Isola, J., and Kallioniemi, O.P. (1995). In vivo amplification of the androgen receptor
gene and progression of human prostate cancer. Nat. Genet. *9*, 401–406.

Viswanathan, S.R., Ha, G., Hoff, A.M., Wala, J.A., Carrot-Zhang, J., Whelan, C.W., Haradhvala,
N.J., Freeman, S.S., Reed, S.C., Rhoades, J., et al. (2018). Structural Alterations Driving
Castration-Resistant Prostate Cancer Revealed by Linked-Read Genome Sequencing. Cell *174*,
433-447.e19.

Wala, J.A., Bandopadhayay, P., Greenwald, N.F., O'Rourke, R., Sharpe, T., Stewart, C.,
Schumacher, S., Li, Y., Weischenfeldt, J., Yao, X., et al. (2018). SvABA: genome-wide detection
of structural variants and indels by local assembly. Genome Res *28*, 581–591.

Wang, S., Li, H., Song, M., Tao, Z., Wu, T., He, Z., Zhao, X., Wu, K., and Liu, X.-S. (2021). Copy
number signature analysis tool and its application in prostate cancer reveals distinct mutational
processes and clinical outcomes. PLoS Genet *17*, e1009557.

Weinhold, N., Jacobsen, A., Schultz, N., Sander, C., and Lee, W. (2014). Genome-wide analysis
of noncoding regulatory mutations in cancer. Nat Genet *46*, 1160–1165.

Willis, N.A., Frock, R.L., Menghi, F., Duffey, E.E., Panday, A., Camacho, V., Hasty, E.P., Liu, E.T.,
Alt, F.W., and Scully, R. (2017). Mechanism of tandem duplication formation in BRCA1-mutant
cells. Nature.

Wu, Y.-M., Cieślik, M., Lonigro, R.J., Vats, P., Reimers, M.A., Cao, X., Ning, Y., Wang, L., Kunju,
L.P., de Sarkar, N., et al. (2018). Inactivation of CDK12 Delineates a Distinct Immunogenic Class
of Advanced Prostate Cancer. Cell *173*, 1770-1782.e14.

Yuan, X., Cai, C., Chen, S., Chen, S., Yu, Z., and Balk, S.P. (2014). Androgen receptor functions
in castration-resistant prostate cancer and mechanisms of resistance to new agents targeting the
androgen axis. Oncogene *33*, 2815–2825.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C.,
Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS).
Genome Biol *9*, R137.

Zhang, Y., Gong, M., Yuan, H., Park, H.G., Frierson, H.F., and Li, H. (2012). Chimeric transcript
generated by cis-splicing of adjacent genes regulates prostate cancer cell proliferation. Cancer
Discov *2*, 598–607.

Gene Expression Omnibus: NCBI gene expression and hybridization array data repository -
PubMed.

1301

Figure 1

Figure 2

Figure 3

Figure 4